Distribution Category: UC-601

ANL-91/40

ANL--91/40

DE92 006763

ARGONNE NATIONAL LABORATORY
9700 South Cass Avenue
Argonne, Illinois 60439

RADIATION CHEMISTRY OF SYNTHETIC WASTE

by

D. Meisel, H. Diamond, E. P. Horwitz, C. D. Jonah, M. S. Matheson,
M. C. Sauer, Jr., and J. C. Sullivan

Chemistry Division

November 1991

Work performed for Westinghouse Hanford Company in accordance with P. O. M652408.

MASTER

1 OF 1 OF THIS DOCUMENT IS UNCLASSIFIED

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency Thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

TABLE OF CONTENTS

	<u>Page</u>
ABSTRACT	1
SUMMARY	2
1. INTRODUCTION.....	3
A. Statement of Problem and Purpose	3
B. Material to be Covered	3
2. RADIATION CHEMISTRY OF DILUTE AQUEOUS SOLUTIONS	4
3. EFFECTS OF HIGH-SOLUTE CONCENTRATIONS ON YIELDS.....	6
A. General	6
B. Effect of Hydroxide	8
C. Effect of Nitrate	11
D. Effect of Nitrite	13
E. Other Inorganic Solutes	13
4. MECHANISMS AND YIELDS OF NO _x RADIOLYSIS.....	13
5. REACTIONS OF ORGANIC CONSTITUENTS.....	15
6. EFFECTS OF HIGH-SOLUTE CONCENTRATIONS ON RATES.....	20
A. Viscosity Effects	20
B. Ionic Strength Effects	22
C. Time-Dependent Rate Constants	23
D. Quantum Mechanical Effects	24
7. EFFECTS OF TEMPERATURE AND PRESSURE ON YIELDS AND RATES	25
8. CONCLUSIONS.....	27
9. REFERENCES	29

FIGURE

<u>No.</u>	<u>Page</u>
1. The ratio of H ₂ yield, G(H ₂), in the presence of the solute concentration [S] to the yield of H ₂ in the absence of solute, G(H ₂) ⁰	10

LIST OF TABLES

1. Yields and Rate Constants for Primary Species Produced in the Radiolysis of Water.....	7
2. Measured and Corrected Values of C ₃₇ , and Rate Constants, k_{dil} , with e_{aq}^-	9
3. Acid-Base Equilibria and Rate Constants for Primary Products of Water Radiolysis	11
4. Standard Redox Potentials of Nitrogen Species in Water	14
5. Representative Rate Constants for Reactions of Organic Additives with the Primary Radicals from Water Radiolysis (numbers in parentheses are pHs)	16
6. Rate Constants for the Reaction of OH Radicals with EDTA and NTA, the Decay (first and second order) of the Radicals, and Their Reaction with Oxygen	17
7. Temperature Effects on Rate Constants for Reactions of the Primary Radical.....	26

RADIATION CHEMISTRY OF SYNTHETIC WASTE

by

D. Meisel, H. Diamond, E. P. Horwitz, C. D. Jonah, M. S. Matheson,
M. C. Sauer, Jr., and J. C. Sullivan

ABSTRACT

The yield of H_2 from radiolysis of aqueous solutions is substantially reduced by the presence of nitrate and nitrite in the waste solutions. Nitrate is more efficient in scavenging the precursors to H_2 than is nitrite, therefore, the latter should be maintained at higher levels if minimization of radiolytic gas production is required. Nitrate is the major scavenger for e_{aq}^- and nitrite is the major scavenger for H atoms. At the concentration levels of the waste solutions some fraction of the radiation energy will be absorbed directly by the solutes, primarily the nitrate/nitrite components. Organic additive will increase the generation of H_2 and mechanistic information is available to allow predictive modeling of trends in the rate of the generation. Physical parameters such as temperature, viscosity, and pressure will not significantly affect the gas generation relative to its generation under normal conditions. Radiolytic generation of N_2O is very inefficient in the absence of organic solutes. No mechanistic information is available on its generation in the presence of organic additives. At the concentration levels of the inorganic salts in the waste solutions, it will be very difficult to find a chemical additive that could efficiently reduce the yield of the generated H_2 , except, perhaps, increasing the concentration of the nitrite/nitrate components.

SUMMARY

An in-depth understanding of the radiation chemistry of aqueous solutions, based on extensive data on yields of products and reaction rates, has been achieved over the last four decades. These two elements, mechanistic understanding and a broad data base, are essential for a meaningful analysis of the radiolytic processes that occur in mixed-waste solutions. This report summarizes the basic principles, along with some quantitative data, of the radiation chemistry of highly concentrated solutions that allow such an analysis. It specifically addresses systems and issues similar to those encountered in the tank safety program (high-base content, high-nitrate and -nitrite contents, and moderate concentrations of organic chelators). It is intended to provide some background information for the nonspecialist along with citations for further exploration. It is also meant to furnish quantitative guidelines for the daily activities of the task force that has been assembled in the Chemistry Division of Argonne National Laboratory.

The high concentrations, primarily of nitrate and nitrite, dictate analysis of the radiolytic processes that occur on very fast time scales (psec). Both nitrate and nitrite are essential in the minimization of dihydrogen generation. Of the two, nitrite is the more important ingredient, and steps must be taken to ensure that known high concentrations of nitrite are maintained in the waste solutions. Any hydrogen-containing organic compound will add to the production of H_2 , in addition to a small H_2 component that results directly from the radiolysis of water. Minimization of the organic components, for example, via carefully planned degradation or separation, should be considered as a remediation strategy.

Although detailed information is available on the radiolytic degradation of the nitrate/nitrite system, its interaction with organic systems has not been adequately studied. The generation of N_2O in the waste solution most probably involves interaction of the NO_x system with the organic components. However, there is no reason to believe that the nitrogen in the N_2O necessarily originates from the organic nitrogenous content.

Except for nitrate, nitrite, and the organic chelators, none of the other components that exist in the waste solution can react with the primary radicals (e_{aq}^- , H, OH) of water radiolysis to any significant extent. The high hydroxide ion concentration ensures complete transformation of most radicals to their base form within some short time. Secondary radicals such as NO_3 , NO_2 , NO, organic radicals, etc., are often longer lived and may react with minor components. This radiation-induced chemistry of the secondary radicals is highly specific, but could be minimized if patterns of the secondary chemistry are established in future studies. The effects of the phase heterogeneity (particulates) of the waste solutions have not been addressed in the survey, but they will probably intervene only in reactions of secondary radicals. Other unconventional physical conditions in the tank (viscosity, temperature, high inert salt concentrations, etc.) will affect rates of individual processes but will not alter the overall product yields. Exceptional in this regard are hydrogen abstraction reactions by H atoms and reactions of the NO_x radicals, which may have significant activation energies and thus may be strongly temperature dependent.

1. INTRODUCTION

A. Statement of Problem and Purpose

The episodic venting of gases in waste-storage tanks may pose significant risk because relatively high levels of H_2 and N_2O gases are detected during these events. A possible source for these gases are radiolytic processes that are initiated by the absorption of ionizing radiation by the waste solution. Preliminary calculations based on the information available in the Waste Tank Science Report⁽¹⁾ indicate that about 25% of the H_2 generated in the tank results directly from the radiolysis of water (assumptions made: rate of H_2 generated in the tank is 37 moles/d; $G(H_2) = 0.06$ molecules/100 eV). In the presence of organic compounds, the yield may be higher and, therefore, a larger fraction of the dihydrogen may be accounted for by radiolysis. The goal of the Argonne task force is to examine the role of radiolytic sources in the generation of gases and to propose remediation strategies. As a first step within this broad mission, we have surveyed the literature in order to inform members of the task force, as well as others, of the available information on yields and mechanisms of dihydrogen and nitrous oxide production in solutions of similar composition to that of the tank. This information will be critically evaluated, and its reliability will be assessed.

B. Material to be Covered

This literature search will focus on the radiolytic reactions that occur in the highly concentrated basic aqueous solutions of sodium nitrate/nitrite that will also be experimentally studied by the task force. Inasmuch as the effects of heterogeneity (slurries, particulates, etc.) are not now under study by the task force, this review will not cover heterogeneous systems. Nevertheless, we note in passing that even in cemented waste (0.2 water/cement ratio, 15% wt sodium nitrate, and ~0.5% organic constituents) the yield of dihydrogen from radiolysis is comparable to that in simulated waste.⁽²⁾ It is obvious that water and/or organic material is required for the generation of H_2 by radiolytic processes, and it is reasonable to infer that aqueous slurries will follow similar trends to those observed in the concentrated solutions. Material specific to the modeling computations performed by our task force (mathematical treatment and computer algorithms) is considered too technical and not directly applicable to the waste problem and thus will not be covered here.

Three data bases were searched for this survey. These were (a) ACS-Chemical Abstracts Data Base (for the period 1967-1990); (b) The Radiation Chemistry Data Center (RCDC) data base of the Radiation Laboratory at the University of Notre Dame; (c) Nuclear Science Abstracts (for the period 1948-1990). These data bases include only open literature; classified relevant information does exist, but is excluded from this report. We observed that much scientific activity on the radiation chemistry of solutions relevant to our task occurred during the late 1950s to early 1970s. Very little has been done since then on this aspect of radiation chemistry despite other important advances in the field. Much more information than could be summarized here on the kinetics of radiolytically produced species is available in several compilations from the RCDC.^(3,4) Similarly, compilations of redox potentials of organic⁽⁵⁾ and inorganic⁽⁶⁾ radicals are available.

The absorption of ionizing radiation by water initiates a sequence of reactions that ultimately leads to the formation of the final products. As the reaction sequence is analyzed and the species involved become removed from the initial act of radiolysis, it becomes increasingly difficult to label the process as "radiolytic". Some judgment had to be made, therefore, as to the extent that this survey would branch into other fields of chemistry beyond what is commonly labeled as radiation chemistry. In general, the dividing line in this summary was determined by the nature of the data bases surveyed. This is particularly felt in the discussion of the NO_x system where many of the

radiolytically generated radicals are stable, though highly reactive, species. Hence, more information on that system probably exists in sources not covered by this survey.

2. RADIATION CHEMISTRY OF DILUTE AQUEOUS SOLUTIONS

Several comprehensive reviews of the radiation chemistry of aqueous solutions provide background information on the field.⁽⁷⁻¹⁰⁾ It is apparent from the high-solute concentrations in the waste solutions that the fate of the primary products of water radiolysis is determined at very early stages. Therefore, we will address these early events in our survey. The following paragraphs summarize the relevant information.

Upon absorption of the energy of ionizing radiation (β or γ) by water, a water molecule ionizes to give thermally unrelaxed electrons and holes (H_2O^+). Although most of the absorbed energy leads directly to ionization of water molecules, some excitation events may also occur. The role of excited water molecules, however, is still not well established. A highly excited water molecule may ionize to give similar products to those obtained by direct ionization,⁽¹¹⁾ or it may dissociate to give either $\text{H} + \text{OH}$ radicals or $\text{H}_2 + \text{O}$. These excited water molecules may play a significant role in highly concentrated solutions and may account for some of the observed H_2 . The sequence of events following the absorption of the radiation energy is illustrated in Scheme I. The primary species are produced within localized domains, labeled "spurs," approximately 20 Å in diameter and separated by several thousand Å from one another along the electron track. Each spur contains the dissociation products of a few water molecules. The processes occurring within these spurs are relevant to our task because some of them produce H_2 and could be intercepted and thus

Scheme I

prevented from producing H₂. In addition, the presence of some solutes (in particular, organic compounds) may enhance H₂ formation by reacting with radicals (e.g., H atoms or e_{aq}⁻) within the spurs.

The number of water molecules dissociated in each spur as well as other spur parameters, have been estimated by Schwarz.⁽¹²⁾ About 50% of the spurs contain the products of one or two water molecules. Approximately 13% of the energy is deposited in short tracks ("blobs") that contain a larger number of dissociation products and that are created as the energy of the original energetic particle degrades. Within <10⁻¹² s following the ionization act, the electrons thermalize, localize, and solvate. The times required for these early processes have been determined by Gauduel et al.^(13,14) with femtosecond laser techniques and are given in reactions 1-3.

The local concentration of the radicals, H, OH, etc., is initially quite high (in the range of molar concentrations) due to the small volume of the spur. A competition between radical recombination, radical scavenging by solutes, and diffusion of the radicals out of the spur is thus initiated. In spite of the short times involved, the precursor to the solvated electrons, e_t⁻, can be scavenged at high-solute concentrations (see below). In the following discussion we will not analyze in detail the processes of the oxidizing species. These reactions are directly relevant to the reactions of the NO_x systems (and thus may be important in the production of N₂O and will be discussed in that context). The organic solutes will often produce reducing radicals as a consequence of their reaction with oxidizing radicals. These reducing organic radicals can generate H₂ from water only under unusual conditions (significantly, such "unusual" conditions include catalysis by particulates). The reactions of the oxidizing radicals, however, are important as competing reactions within the spurs and will be taken into account in our calculations.

Of particular interest are radical recombination reactions that produce dihydrogen, reactions 4-6 (numbers in parentheses indicate estimated yields in units of molecules per 100 eV for the particular reaction in dilute solutions).⁽¹²⁾

The large value for formation of dihydrogen from hydrated electrons (reactions 4 and 5) emphasizes the importance of this species as a source of H₂. Some hydrogen atoms are also formed by the reaction of solvated electrons with protons within the spurs (reaction 7):

In addition, a yield of G(H₂)⁰ = 0.15 (where the superscript 0 indicates direct, nonscavengeable formation of dihydrogen, e.g., from an excited water molecule) has been estimated by Schwarz.⁽¹²⁾ The observation that at high-nitrate concentrations the yield of H₂ is reduced to G(H₂) = 0.06,⁽¹⁾ along with our own experimental results, indicates that direct formation of H₂

must be smaller than the estimate in reference 12. Mahlman's estimate of $G(\text{H}_2)^0 = 0.1$, based on the dependence of the yield of H_2 on nitrate concentration,⁽¹⁵⁾ also seems too high. The yield of directly produced dihydrogen is of primary concern in our considerations. At present, it seems reasonable to assume that $G(\text{H}_2)^0 \leq 0.06$.

It is commonly accepted that the product distribution and the yield of H_2 at "long" times (when spur reactions are complete, at ~ 100 ns) in dilute solutions can be represented by equation 8,⁽¹⁶⁾ where the numbers in parentheses represent the corresponding G values:

At 1 ps, prior to the completion of spur reactions, the yields can be represented by equation 9:

Obviously, after 100 ns some of the reducing radicals have reacted to produce dihydrogen within the spur. The earliest species detectable by existing technology (best suited for such studies is the fast pulse radiolysis system at the Low Energy Linac Facility at CHM/ANL) are e_{aq}^- and OH radicals. The yields quoted in equation 9 were experimentally observed,^(17,18) and their time dependence has been recorded in dilute solutions and concentrated solutions. *It is clear from this discussion that the term "yield" is a time- and concentration-dependent quantity.* The yield of H_2 , and the yields of its precursors, will depend on the time at which they are measured after the ionization act. High-solute concentrations may reduce the yield of H_2 if the solute reacts with a precursor of H_2 before that precursor can produce dihydrogen. Table 1 summarizes the known yields of the primary products from the radiolysis of water at short and long times and the rate constants for the reactions among these radicals. These data are important in our modeling task.

3. EFFECTS OF HIGH-SOLUTE CONCENTRATIONS ON YIELDS

A. General

As the concentrations of solutes increase, several effects may be expected. The most obvious effect is the competitive reaction of the solutes for the primary radicals. To the extent that the composition of the solution is known (questionable in the tank), this effect can be calculated easily and accurately from the known rate constants summarized in the tables or elsewhere.⁽³⁾ However, the rate constants listed in the tables were measured in dilute solutions. These rate constants will be modified at the high-solute concentrations by the reasonably well-understood and predictable effects discussed in section 6. At high concentrations the solute will penetrate the spur, and its reactions may efficiently compete with intraspur reactions. The effect of competition with spur reactions can be reasonably estimated by any of several variants of the continuum diffusion model.^(12,19-21) The computational effort of our task force will utilize a form of this model that has been modified to include the effects of high-solute concentrations.

When solute competition for spur species is significant (as we expect it to be in the tank solutions), the solute may compete for the precursors of the well-defined primary products. These precursors (e.g., the localized electrons of equation 1 or H_2O^+ of equation 2) cannot be directly observed, even with psec pulse radiolysis, due to their short lifetimes, but the reactivity of the solute with the precursor can be estimated from the reduction in yield of the primary radical at very short times.^(22,23) It was shown that the fraction, ϕ , of electrons that escape reaction with the solute, S, prior to their solvation is given by

TABLE 1. Yields and Rate Constants for Primary Species Produced in the Radiolysis of Water

Yields

Species	Yield at 1 ps	Yield at Long Times ^a
e_{aq}^-	4.78	2.70
H	0.62	0.61
OH	5.7	2.87
H ⁺	4.78	2.70
HO ₂	-	0.026
H ₂	0.15 ^b	0.43
H ₂ O ₂	-	0.61

^aLong times refer to the time when spurs have homogenized, $\sim 10^{-7}$ s. ^bCalculated value.

Rate Constants

Reactants		Products	k (M ⁻¹ s ⁻¹) ^c
$e_{aq}^- + e_{aq}^-$	→	H ₂ + 2OH ⁻	$2k = 1.1 \times 10^{10}$
$e_{aq}^- + H^+$	→	H	2.3×10^{10}
$e_{aq}^- + H$	→	H ₂ + OH ⁻	2.5×10^{10}
$e_{aq}^- + OH$	→	OH ⁻	3.0×10^{10}
$e_{aq}^- + O^-$	→	2OH ⁻	2.2×10^{10}
$e_{aq}^- + H_2O_2$	→	OH ⁻ + OH	1.1×10^{10}
$e_{aq}^- + HO_2^-$	→	2OH ⁻ + OH	3.5×10^9
H + H	→	H ₂	$2k = 1.55 \times 10^{10}$
H + OH	→	H ₂ O	7.0×10^9
H + H ₂ O ₂	→	OH + H ₂ O	9.0×10^7
OH + OH	→	H ₂ O ₂	$2k = 1.1 \times 10^{10}$
OH + H ₂	→	H + H ₂ O	4.2×10^7
OH + H ₂ O ₂	→	H ₂ O + HO ₂	2.7×10^7
OH + HO ₂ ⁻	→	H ₂ O + O ₂ ⁻	7.5×10^9
OH + O ⁻	→	HO ₂ ⁻	$\leq 2.0 \times 10^{10}$
O ⁻ + O ⁻	→	O ₂ ²⁻ ⇌ HO ₂ ⁻	unknown
O ⁻ + H ₂	→	H + OH ⁻	1.1×10^8 ^d
O ⁻ + H ₂ O ₂	→	O ₂ ⁻ + H ₂ O	$\leq 5.0 \times 10^8$
O ⁻ + HO ₂ ⁻	→	O ₂ ⁻ + OH ⁻	4×10^8

^cAll values are critically evaluated from reference 3. ^dFrom reference 93.

$$\phi = \exp\left(\frac{-\{S\}}{C_{37}}\right),$$

where C_{37} is a constant typical for the solvent and solute and $\{S\}$ is the solute concentration. When $\{S\} = C_{37}$, the yield of solvated electrons will drop to $\frac{1}{e}$ of its value in dilute solutions. To emphasize the significance of this parameter, for nitrate $C_{37} = 0.47$ M;⁽²⁴⁾ hence, at 2 M nitrate concentration, less than 1.5% of the electrons will become solvated. The remainder will react with nitrate prior to solvation. Since the reaction with nitrate does not produce H_2 , it is clear that nitrate is an extremely efficient scavenger in reducing the yield of dihydrogen from reactions 4, 5, and 7. Table 2 summarizes the presently known values of C_{37} . Significant to our task, nitrite (with $C_{37} = 1.6$ M)⁽²⁵⁾ is considerably less effective than nitrate in scavenging the hydrated electron precursors. Other solutes with small C_{37} (Table 2), which are also present in the tank, may also have some effect. With the exception of H^+ , C_{37} is generally smaller (i.e., smaller concentrations are required to scavenge a given percentage of electrons) the higher the reaction rate of the solute with solvated electrons. The products of these reactions are not well established, but we can surmise that they are the same as those obtained in the reactions of the hydrated electrons with the same solutes.

At yet higher solute concentrations, an appreciable fraction of the energy is deposited in the solute rather than in the water. This is commonly labeled as the "direct effect". The energy absorbed by the solute is commonly assumed to be proportional to its electron density and thus can be readily calculated. It was recently suggested that the valence-electron density is a more appropriate parameter than the total electron density.⁽²⁶⁾ The products of the direct effect are often unknown. In the absence of detailed information, one has to assume that the products of the direct effect will be a hot electron and a hole of the absorbing solute. Detailed information on this effect in concentrated nitrate solutions is available and will be discussed below. More difficult to predict is the fraction of energy that is absorbed by water at such high-salt concentrations and that still produces the same primary products as in dilute solution. This complication apparently is due to the inefficiency of water of solvation in producing the "normal" products of water radiolysis. Thus, even inert ions such as alkali metal ions will affect the yields of molecular products (e.g., H_2). The higher the solvation energy of the ion, the smaller the yield of free water radiolysis products. The effect of inert cations on the yield of H_2 in concentrated nitrate solutions has been determined experimentally.⁽²⁷⁾

The yield of dihydrogen from radiolysis of water in the presence of several solutes has been measured over a wide concentration range. It was noted by Hayon and Moreau⁽²⁸⁾ and earlier by Schwarz⁽²⁹⁾ that the dependence of these yields on concentration for the various solutes could be superimposed on a unified curve by multiplying the solute concentration by a correction factor, f , specific to the solute. Because such a universal curve may find practical utility, we present it in Figure 1. It contains information on the yield of H_2 in both nitrate and nitrite solutions but not on their mixtures. The theoretical rationalization given for this empirical curve is that it represents the efficiency of scavenging of solvated electrons by the solute at low concentrations and of H atoms at higher concentrations. The factor f thus reflects the reactivity of the solute towards these two H_2 precursors.

B. Effect of Hydroxide

The main effect of high-hydroxide concentrations on the yield of molecular products results from the shift in acid-base equilibria of OH , e_{aq}^- , H^+ , and H .

TABLE 2. Measured and Corrected Values of C_{37} , and Rate Constants, k_{dil} , with e_{aq}^-

Compound	C_{37}^a M	$C_{37(\text{corr})}^b$ M	k_{dil}^c ($\times 10^{-10}$) $\text{M}^{-1} \text{s}^{-1}$
SeO_4^{2-}	0.42	0.42	0.11
MoO_4^{2-}	0.63	0.65	0.2
OCl^-	0.48	0.52	0.7
acetone	1.4	1.6	0.87
NO_3^-	0.42	0.47	1.1
Te(OH)_6	0.075	0.09	3.2
salicylate	0.8	1.0	1.0
Cu^{2+}	0.9	1.3	3.3
HgCl_2	0.08	0.09	3.4
Ag^+	0.14	0.17	4.0
Cr^{3+}	0.30	0.84	6.0
CH_3NO_2	0.30	0.37	3.0
Cd^{2+}	0.38	0.55	5.0
Fe^{3+}	1.3	>3.5	6.0
Hg^{2+}	0.062	0.09	8.0
CrO_4^{2-}	0.2	0.26	2.0
$\text{CCl}_3\text{CO}_2^-$	0.5		0.85
$\text{ClCH}_2\text{CO}_2^-$	6.0		0.12
cystamine	1.1		4.0
NO_2^-	1.6		0.4
IO_4^-	0.14	0.33	3.7
Zn^{2+}	8.0		0.15
H^+	>10		2.5

^aMeasured C_{37} values from references 24 and 25. ^bCorrected for time dependence of the rate constant k_{dil} . ^cRate constant with e_{aq}^- in dilute solutions.

Universal Curve
Fraction of Hydrogen Yield vs Corrected Solute Conc.

Figure 1: The ratio of H₂ yield, G(H₂), in the presence of the solute concentration [S] to the yield of H₂ in the absence of solute, G(H₂)°. From reference 28. The concentration correction factor, *f*, for the solutes tested are: KNO₂, 1.0; NaNO₃, 2.4; H₂O₂, 2.5; acrylamide, 4.0; Na₂Cr₂O₇, 8.2; Cu(NO₃)₂, 10.

However, it should be recognized that equilibrium takes some time to evolve. Therefore, the initial distribution of the primary products is far from the distribution at equilibrium; as time progresses, these acid-base reactions will compete with the other processes mentioned in the previous section. Furthermore, the different acid-base forms of each species will have different reactivities and different sets of rate constants. For that reason, the rate constants for OH as well as for O⁻ have to be considered and are included in Table 1. Relevant acid-base rate and equilibrium constants of the primary radicals are summarized in Table 3. According to equation 13, e_{aq}⁻ is the base form of the H atom.

Many organic radicals that may be produced by the reaction of the primary radicals with the organic additives will also have pK_a 's that are widely different from those of the parent molecules and may ionize in the basic solution in the tank. For example, hydroxylic moieties often will retain the proton in the molecule (ROH) but will convert to the base form (RO⁻) in the radical (e.g., $pK_a(\text{CH}_3\text{CH}_2\text{OH}) = 18$; $pK_a(\text{CH}_3\text{CH}\cdot\text{OH}) = 11.5$; ⁽³⁰⁾ for the α -hydroxy radical of glycolate, $pK_a = 8.8$). ⁽³¹⁾ In the base form, these radicals may be expected to be stronger reductants than in the acid form. For alcohols, the pK_a of the radical is 4-5 pH units below that of the molecule.

TABLE 3. Acid-Base Equilibria and Rate Constants for Primary Products of Water Radiolysis

Acid form		Base form	k_f^a (M ⁻¹ s ⁻¹)	k_b^a	pK
OH	$\rightleftharpoons$	O ⁻ + H ⁺	$2.5 \times 10^{-2} \text{ s}^{-1}$	2×10^{10}	11.9
OH + OH ⁻	$\rightleftharpoons$	O ⁻ + H ₂ O	1.3×10^{10}	1.8×10^6	-3.9
H + OH ⁻	$\rightleftharpoons$	H ₂ O + e _{aq} ⁻	2.2×10^7	19	-6.1
H ₂ O	$\rightleftharpoons$	H ⁺ + OH ⁻	$3.6 \times 10^{-5} \text{ s}^{-1}$	2.0×10^{11}	15.7
H	$\rightleftharpoons$	e _{aq} ⁻ + H ⁺	4.8 s^{-1}	2.3×10^{10}	9.6

^a k_f and k_b are the rate constants for the forward and backward reactions as written.

C. Effect of Nitrate

Much effort has been expended in the study of the radiation chemistry of nitrate solutions over a very broad concentration range. Nitrate ions do not react with OH radicals, even at 5 M concentration, ⁽³²⁾ and react only slowly with H atoms. They are, however, extremely efficient scavengers of dry and solvated electrons.

Nitrate is the only species of the NO_x group for which the direct effect has been studied in detail. ^(27,33,34) This effect is the major source for the production of dioxygen in highly concentrated nitrate solutions. The results of the direct effect on nitrate are initiated by reaction 14.

The ejected electron will either solvate or react with NO₃⁻; the NO₂ species may enter any of the pathways available to it in Scheme II; the oxygen atom may react either with NO₂ to give the NO₃ radical, with a scavenger, or with another NO₃⁻ ion to give O₂ (and NO₂⁻); and all of these reactions will occur in competition with diffusion out of the spur and with the various intraspur recombination reactions. In practice, the experimental observation is an increase in G(O₂) and G(NO₂⁻) that parallels the decrease in G(H₂) upon increasing nitrate concentrations in this high-concentration regime. Thus, it is well established that increasing nitrate concentrations in the high-concentration regime leads to a decrease in dihydrogen yields but to an increase in dioxygen yields.

Reports on the yield of N₂O production in nitrate solutions are scarce. Some may be found in still-classified documents. The generation of N₂O in high-level liquid waste from fast breeder reprocessed fuel has been previously observed but mechanistic understanding from this observation is minimal. ^(35a) Hart and Bibler, ^(35b) independently, reported recently that the addition of organic solutes to nitrate solutions leads to the radiolytic production of N₂O. A significant experiment in which both nitrite and nitrate are present in the irradiated solutions has not been performed yet in their studies, but our results show that high yields of N₂O are produced in the radiolysis of nitrite/nitrate mixtures when organic substances are also present.

D. Effect of Nitrite

The radiation chemistry of nitrite will be considered in section 4 and in Scheme II. In the waste solutions it will be the major OH and H-atom scavenger. The direct effect on NO_2^- has not been studied, but we may reasonably speculate that it will follow an analogous course to that of reaction 14. Due to its efficient scavenging of hydrogen atoms, it is one of the most efficient additives in the reduction of $\text{G}(\text{H}_2)$ (see Figure 1). It is clear from these considerations and those discussed in section 4 that maintaining high-concentration levels of nitrite is a vital component in efforts to reduce radiolytic production of H_2 . That nitrate alone, even at high concentrations, is insufficient to maintain the low levels of dihydrogen yields necessary for stabilization of the tank is amply shown in the work of Sutton and co-workers.^(36,37) These researchers have shown that even in 2 M nitrate, competitive reactions with organic substrates (e.g., alcohols, formate) will generate significant yields of H_2 .

E. Other Inorganic Solutes

Among the other inorganic solutes present in the tank, aluminate is a major component. The rate constant for the reaction of Al^{3+} with e_{aq}^- has been determined ($2 \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$). The rate of reaction of aluminate ions with the solvated electron, however, is much slower ($5.5 \times 10^6 \text{ M}^{-1} \text{ s}^{-1}$) and could not significantly compete with the other components in the waste solutions. The product of the reduction, Al^{2+} , can react with H atoms (as well as with reducing organic radicals) at low pH to produce H_2 via the hydride mechanism;⁽³⁸⁾ at high pH, production of e_{aq}^- from metallic aluminum has been reported.⁽³⁹⁾ It is difficult to see how aluminate could directly interfere with any of the radiolytic processes because the rates of its reactions are rather slow. However, the existence of alumina particulates in the heterogeneous mixture may alter many reaction pathways, either catalytically or simply by altering the microenvironment at which the reaction would otherwise occur.

Another component of some significance is carbonate. It is a poor electron scavenger but a reasonably efficient OH scavenger ($k = 4.2 \times 10^8 \text{ M}^{-1} \text{ s}^{-1}$). At the high pH prevailing in the tank, the O^- radical will react much more slowly than the OH radical due to the electrostatic repulsion. Considering its relatively low concentration ($<0.5 \text{ M}$), it is unlikely to compete with the NO_x system for any of the primary radicals. Similar conclusions may be reached regarding other components of the waste solutions. None of them are likely to interfere directly with the reactions of the primary radicals. However, many of them could react with secondary radicals, primarily the longer lived ones from the NO_x system.

4. MECHANISMS AND YIELDS OF NO_x RADIOLYSIS

The generation of N_2O in the tank solutions may result from the radiolytic reduction of nitrate, nitrite, or an adduct of any of these compounds with organic substrates. We, therefore, consider the state of knowledge on the degradation of NO_x systems under the field of radiation. A comprehensive description of all the known reactions of the primary radiolytically produced radicals with the nitrate and nitrite systems is given in Scheme II. This scheme includes rate constants and equilibrium constants for the corresponding reactions. Much of the scheme has been compiled from the work of Henglein and coworkers,⁽⁴⁰⁻⁴³⁾ but contributions from other groups are also included.⁽⁴⁴⁻⁴⁶⁾ Some standard redox potentials (1 M H^+) of various nitrogen-containing intermediates are collected in Table 4.⁽⁶⁾ In estimating relative reactivities at a given pH from standard redox potentials, it should be remembered that the pK_a 's of both the reduced and the oxidized species of a redox couple must be known in order to perform the conversion. In the following text, we emphasize only the major features of the NO_x system. As stated in section 1B, the

literature on reactivity of the NO_x system from sources other than the radiation chemistry literature is only marginally covered in this survey.

TABLE 4. Standard Redox Potentials of Nitrogen Species in Water (taken from reference 6)

Oxidation / Reduction	E ⁰ , V(NHE)
NO ₃ / NO ₃ ⁻	2.49
NH ₃ ⁺ / NH ₃	2.13
NO ₂ ⁺ / NO ₂	1.51
N ₃ / N ₃ ⁻	1.33
NH ₂ OH ⁺ / NH ₂ OH	≤1.26
NO ⁺ / NO	1.21
NO ₂ / NO ₂ ⁻	1.04
NH ₂ / NH ₂ ⁻	-0.75
N ₂ H ₄ ⁺ / N ₂ H ₄	≤0.73
NH ₂ OH / (NH ₂ , OH ⁻)	-0.12
NO / NO ⁻ (singlet)	-0.35
NO / NO ⁻ (triplet)	-0.39
NO ₃ ²⁻ / NO ₃ ⁻	≤-0.4
NH ₃ OH ⁺ / (NH ₃ , OH)	-0.58

The NO₃ radical in the waste solution will be produced primarily by the direct effect on nitrate (top left, Scheme II). The redox potential of the NO₃/NO₃⁻ couple has been estimated to be 2.3-2.6 V.⁽⁴⁷⁾ This radical is, thus, a very strong oxidant. Its reaction pathways parallel those of OH radicals, leading to similar products in many cases, albeit at a reduced rate. The reactions of NO₃ radicals with many inorganic substrates are electron transfer processes, whereas the reactions with saturated organic compounds are hydrogen abstraction reactions; with unsaturated organic substrates they are an addition reaction. Rate constants for some of these reactions have been measured and are summarized in reference 4. For primary alcohols, the rate constant is linear with the number of carbon atoms ($k = 3.2 \times 10^5 \text{ M}^{-1} \text{ s}^{-1}$ per carbon atom).⁽⁴⁸⁾ From the redox potentials, rate constants, and concentrations, it is evident that most of the NO₃ radicals produced in the waste solution will be converted to NO₂ radicals by their reaction with nitrite (reaction 15).

Nitrite ions are the primary scavenger for OH radicals producing NO₂. The NO₂ radical will also be produced in the tank by reaction of e_{aq}⁻ with NO₃⁻.

Due to the high rate constants of reactions 16 and 17 and the large concentrations of nitrite and nitrate, both reactions are expected to be the dominant reactions of the two corresponding primary

radicals; hence, the NO_2 radical is expected to be an important radical in the tank solutions. Furthermore, in the absence of any reductants (e.g., organic chelators), this is a well-known stable species. It is a mild oxidant with an estimated redox potential (for $\text{NO}_2/\text{NO}_2^-$) of 1.03 V.⁽⁴⁹⁾ The rate constants for its reaction with several phenoxides and their temperature dependence have been determined.^(50,51) All reported reactions of NO_2 with stable scavengers are one-electron transfer reactions, with the notable exception of SO_3^{2-} , which proceeds via an oxygen-atom transfer. Most importantly, nitrite ions are the primary scavengers for hydrogen atoms (reaction 18). Reaction 18

competes with the reaction of H atoms with organic substrates, which often leads directly to the production of H_2 . It is already clear from this discussion, as well as from our experimental and modeling results, that it is essential to determine the concentration of nitrite in the tank solution and to maintain it at high levels of concentration if minimization of radiolytic dihydrogen production is desired.

Another stable radical, produced during the radiolytic reduction of nitrite (by either e_{aq}^- or H atoms or organic radicals), is NO. Its chemistry is well known and will not be surveyed here. If NO and NO_2 (or their dimers) accumulate in the solution, the cross recombination product, N_2O_3 , will also form. The high concentration of hydroxide in the tank solution will shift the equilibria so that N_2O_3 will be mostly transformed to NO_2^- , and NO_2 will dimerize, disproportionate, and hydrolyze to give NO_3^- and NO_2^- . The steady state concentration of each of these species (NO_3^- , NO_2 , NO, etc.) will depend on the exact concentrations of base, nitrite and nitrate, dose rate, total dose, and reactions with organic compounds. Except for the reactions with organic compounds, the information given in Scheme II should allow a rather accurate prediction of the concentrations of these intermediates. Production of N_2O from NO in basic solutions has long been known⁽⁵²⁾ and has been verified in our experiments. NO is also expected to react efficiently with many organic radicals but specific information on the rates of such reactions is not available in the data bases surveyed for this report. One-electron reduction of NO will lead directly to nitrogen in the +1 oxidation state, the same as in N_2O . Hence, a ready pathway for the radiolytically induced generation of N_2O can be envisaged.

5. REACTIONS OF ORGANIC CONSTITUENTS

Organic constituents are potentially a source for the production of dihydrogen, primarily via hydrogen abstraction reactions by hydrogen atoms. Some rate constants for scavenging of the radiolytic primary radicals by representative organic substrates are given in Table 5. Many more are available in the compilation by Buxton et al.⁽³⁾ Note, however, the paucity of data on the rate of reaction of hydrogen atoms with organic substrate at high pH. This may be significant for the organic ligating agents of interest here, because the rates of these reactions will depend on their acid-base form. Also missing in the literature are some rate constants for organic components that are believed to be present in the tank.

Chelating agents are of major concern as sources of dihydrogen and N_2O gaseous products in the tank solutions; we shall concentrate on the radiation chemistry of these compounds in this section. We start with the common chelating substances and proceed stepwise through their likely degradation products. EDTA is one of the principal organic compounds originally put in the tank. The pK_a for the ionization of the last $-\text{COOH}$ group is 9.95 (at 1.0 M ionic strength), so this compound is completely ionized to EDTA^{4-} . This will also be true for all other chelating agents discussed here; they will exist in the completely ionized form in the tank solution. The reactivity of e_{aq}^- toward EDTA^{4-} is rather low ($k \cong 4 \times 10^5 \text{ M}^{-1} \text{ s}^{-1}$ at pH 11.5 and ionic strength 1.6).⁽⁵³⁾ Because there is a correlation between the rate constant for the e_{aq}^- reaction and the reactivity

TABLE 5. Representative Rate Constants for Reactions of Organic Additives with the Primary Radicals from Water Radiolysis (numbers in parentheses are pHs). Rate constants for NO_3^- and NO_2^- are shown for comparison.

Compound	$k(e_{aq}^-)$	$k(H)^c$	$k(O^-)^d$	$k(OH)^d$
Ethanol		1.3×10^7 ^a (1)	12.2×10^8	17.9×10^8
Acetic acid	2×10^8 ^a (5.4)	8×10^4		
Acetate	1.1×10^6 ^a (~10)	3.5×10^5 ^a (7)	0.5×10^8	0.7×10^8
Oxalic acid		4.1×10^5		
Oxalate	3.1×10^7 ^a (7.6)	$< 4 \times 10^4$ ^a (7)		7.7×10^6 ^a (6)
Glycolic acid	4.3×10^8 (4)	1.8×10^7		6.6×10^8 (1)
Glycolate	8.2×10^6 ^a (7)	3.7×10^7 ^a		5.6×10^8 ^a (5.5)
Glyoxalic acid	1.3×10^9 ^a (4)			
Citric acid		4.3×10^5		5.0×10^7 ^a
Citrate	$< 10^5$ ^a		0.42×10^8	
Glycine	2.8×10^8 ^a (11.8)	8×10^4	5.6×10^8	29.5×10^8
IDA	1.2×10^7 ^a (10.5)	4.0×10^5	7.9×10^8 ^e (13)	8.6×10^7 ^e (6)
NTA		7.5×10^6		2.5×10^9 ^a (9)
EDTA	4.5×10^5 ^b (11.5)	6.5×10^7	1×10^8 ^f	2.8×10^9 ^g (9)
NO_2^- ^a	4.1×10^9	7.1×10^8	3.1×10^8	1.0×10^{10}
NO_3^- ^a	9.7×10^9	1.4×10^6	no reaction	no reaction

^aReference 3; ^bReference 53; ^cReference 66 at pH 1 unless otherwise indicated; ^dReference 65 unless otherwise indicated; ^eReference 58; ^fReference 55; ^gReference 54.

towards its precursor, most probably EDTA^{4-} will also be unreactive towards the precursor of the hydrated electron. The reactive oxidizing species in the highly alkaline waste solutions is the ionized form of OH, i.e., O^- radical. It may be safely assumed that at the concentration level of the organic molecules, ~0.1 M, acid-base equilibria of the primary radicals are established prior to the reaction with the organic scavenger. Bhattacharyya and Kundu⁽⁵⁴⁾ find in neutral solution $k(\text{OH} + \text{EDTA}) = 2.8 \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$. As the pH increases, the rate decreases; Mulazzani and co-workers⁽⁵⁵⁾ quote $k(\text{O}^- + \text{EDTA}^{4-}) \approx 1 \times 10^8 \text{ M}^{-1} \text{ s}^{-1}$. The high ionic strength of the waste solution would likely minimize this largely electrostatic effect. The main reaction pathway is an abstraction of an H atom from the position α to the $-\text{CO}_2^-$ group. Other possibilities are electron transfer from the CO_2^- or the amino groups, and an H abstraction from the ethylene groups. However, it was postulated that the carboxy radical ($-\text{CO}_2^\cdot$) is produced only in negligible amounts, whereas the other two radicals are converted to the α -carboxy radical.⁽⁵⁵⁾ Electron spin resonance results⁽⁵⁶⁾ (at pH ~ 11) also reveal only the α -carboxy radicals. Above about 10^{-3} M , $G(-\text{EDTA}) = 1.6 \pm 0.3$ is independent of concentration at pH 13 in deaerated solutions;⁽⁵⁴⁾ no carbonyl-containing products (e.g., formaldehyde or glyoxalic acid) were found at this pH in deaerated

solution. However, this last observation is probably due to the efficient reaction of e^-_{aq} with the carbonyl products, because the yield of glyoxalic acid can be restored on addition of e^-_{aq} scavengers (e.g., N_2O).^(57,58)

The radicals obtained from the OH reaction with EDTA and NTA decay via a first-order reaction followed by a second-order decay (studied, though, only at $pH \leq 12$).⁽⁵⁹⁾ Some relevant results are given in Table 6. The fast reaction rate with dioxygen (Table 6) indicates that these are, indeed, aliphatic radicals. The first-order reaction was attributed to the transformation of the amino or alkyl radicals to the α -carboxy radical, the second-order reaction to the disproportionation reaction, followed by hydrolysis of the latter to give the final products (glyoxalic acid, formaldehyde, CO_2 , and the ligand with one less acetate group). The sequence of reactions that leads to the degradation of EDTA to NTA is shown in Scheme III. The reaction of EDTA with H atoms is expected to follow the abstraction from $-CH_2-$ pathways.

Bhattacharyya and Srisankar⁽⁵⁷⁾ irradiated aqueous solutions of 0.01 M NTA and found $G(-NTA) = G(IDA)$ in neutral and pH 11.5 solutions. In deaerated alkaline solution, $G(-NTA) = 1.6 \pm 0.2$ and $G(IDA) = 1.4 \pm 0.1$, whereas $G(\text{carbonyl}) = 0.5$ (measured as formaldehyde). The G values increase linearly with increasing NTA concentration up to $\sim 8 \times 10^{-3}$ M and then become independent of concentration. The mechanism proposed to explain the observed products includes the disproportionation reaction between two radicals, reaction 19,

followed by the hydrolytic reactions of the hydroxy acid and the keto acid (reactions 20a, 20b).

TABLE 6. Rate Constants for the Reaction of OH Radicals with EDTA and NTA, the Decay (first and second order) of the Radicals, and Their Reaction with Oxygen^a

	pH	$10^{-8} \times k (\text{OH} + \text{Ligand})$ $M^{-1} s^{-1}$	$10^{-2} \times k_{D1}$ $M^{-1} s^{-1}$	$2k_{D2}$ $M^{-1} s^{-1}$	$10^{-8} \times k (R\cdot + O_2)$
EDTA	4.5	4.0	30	2.0×10^7	2.0
	9.0	20	2.6	1.0×10^6	4.5
NTA	4.0	7.5	30	5.0×10^7	1.7
	9.0	25		1.0×10^5	3.0
	11.3			5.0×10^6	
IDA	6.0	0.86			
	13.0 ^b	7.9			
Glycine	6.0	$0.082(e^-_{aq})^c$			
	7.8			1.0×10^9 ^d	

^aValues for EDTA and NTA are from reference 59; for IDA from reference 58.

^bAt this pH OH is in the O^- form.

^cRate constant for reaction with e^-_{aq} from reference 62.

^dDecay rate constant for $\dot{C}H_2CO_2^-$ radicals from reference 62.

It should be recognized here that the disproportionation reaction 19 requires a reaction between two radicals that might be at very low concentrations in a complex solution such as the waste system and, therefore, very slow. However, an oxidation reaction (e.g., by the long-lived NO_2 radicals) may lead to identical degradation products that were obtained in the oxidation side of the disproportionation, ultimately leading to ammonia when $\text{R} = \text{H}$. Bhattacharyya and Saha⁽⁵⁸⁾ also studied the γ radiolysis of IDA. For glycine and carbonyl compounds, G values again level off, at $\sim 4.0 \times 10^{-2}$ M IDA; $\text{G}(-\text{IDA}) = \text{G}(\text{glycine}) = 1.9$ at pH 13 in deaerated solution, whereas $\text{G}(\text{CH}_2\text{O}) = 0.5$ and $\text{G}(\text{CHOCOOH}) = 0.1$. Again, the carbonyl-containing products efficiently compete for e_{aq}^- , but when the latter is scavenged (by N_2O), $\text{G}(\text{glycine}) = 3.3$, $\text{G}(\text{CH}_2\text{O}) = 1.2$, and $\text{G}(\text{glyoxalic}) = 2.0$.

The hydrated electron reacts with glycine, $\text{H}_3^+\text{NCH}_2\text{COO}^-$, in neutral solution⁽⁶⁰⁾ via the two pathways shown below:

The $\dot{\text{C}}\text{H}_2\text{CO}_2^-$ radicals that are obtained in the β -scission reaction 21a were also observed for NTA and IDA. Irradiation of solutions of glycine at pH > 13 (in the presence of OH, $\text{O}^\bullet$, and H scavengers) gave an appreciable yield of $\dot{\text{C}}\text{H}_2\text{CO}_2^-$, although less efficiently than in neutral solution (probably because the reaction with e_{aq}^- is slower when the glycine molecule is negatively charged; the fraction that undergoes deamination, however, is probably larger). Thus, some of the deamination reaction with e_{aq}^- occurs at high pH as well:

Markovic et al.⁽⁶¹⁾ find $k_{21a}/k_{21b} = 1.6$, hence, ca. 60% of the reaction will proceed via the deamination pathway in neutral solutions. Simic and Hayon⁽⁶²⁾ find only 40% deamination of glycine (reasonably good agreement). The C_{37} at pH 6 for glycine is >10, so $\text{H}_3^+\text{NCH}_2\text{CO}_2^-$ is quite unreactive toward the e_{aq}^- precursor.⁽⁶³⁾ Reaction of glycine with OH radicals again yields the α -carboxy radical (identical to the α amino in this case). This radical, $\text{H}_2\text{N}\dot{\text{C}}\text{HCO}_2^-$, was identified by ESR.⁽⁵⁶⁾ Neta et al.⁽⁶⁴⁾ determined the following acid-base equilibria ($\text{pK}_{a1} < 1$, $\text{pK}_{a2} = 6.6$):

Thus, the glycy radical (and probably the analogous radicals considered above) is completely deprotonated at pH 14. Rate constants for the reaction of $\text{O}^\bullet$ radicals with glycine were also measured in 1 M NaOH.⁽⁶⁵⁾ The Coulombic repulsion reduces reaction rates for singly, doubly, and triply charged anions by factors of 0.36, 0.10, and 0.03, respectively, at low ionic strength.⁽⁶⁵⁾ However, the high ionic strength in the tank would substantially reduce this Coulombic effect. Hydrogen atoms will react similarly to the reaction of OH with glycine and similar amino acids; hydrogen abstraction by H atoms will, of course, directly produce dihydrogen. Ionization of $-\text{COOH}$ to $-\text{COO}^-$ would change the rate constant of the H atom reaction by only a factor of ≤ 2 , whereas deprotonation of the amino group to NH_2 might increase it by a factor of 10^2 to 10^3 .⁽⁶⁶⁾ Several rate constants for the reactions of e_{aq}^- , H, OH, and $\text{O}^\bullet$ with these organic constituents are listed in Table 5.

HEDTA differs from EDTA in that it has a C-H bond α to the hydroxyl group. At this position, the hydrogen atom is activated relative to other positions and is easier to abstract. However, no experimental verification of this expectation is available in the literature. Furthermore, some degradation products may appear as hydroxy-carboxylic acids (e.g., glycolic acid). For these organic substances, OH and H will abstract the H atom α to the hydroxyl group, and the resulting radicals will dimerize or disproportionate.⁽³¹⁾

In aqueous solutions of glycolic acid, the main radiolysis products are H_2 ($G = 4.2$)⁽⁶⁷⁾ and tartaric acid ($G = 2.1$).⁽⁶⁸⁾ The pK_a of $\text{HO}\dot{\text{C}}\text{HCOOH}$ is 4.6, and the pK_a of $\text{HO}\dot{\text{C}}\text{HCOO}^-$ is 8.8. Thus, even the hydroxyl group of the radical will ionize in the tank solution. This ionization will increase the lifetime of the radical significantly. For the recombination of two $-\text{O}\dot{\text{C}}\text{HCOO}^-$ radicals, $2k = 1.5 \times 10^7 \text{ M}^{-1} \text{ s}^{-1}$ at pH 12; $2k = 3.6 \times 10^6 \text{ M}^{-1} \text{ s}^{-1}$ for the citrate radicals ($-\text{OOCCH}_2\text{C}(\text{OH})(\text{COO}^-)\dot{\text{C}}\text{HCOO}^-$) at pH 13. In comparison, many of the fully protonated radicals will recombine with $2k = 2 \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$. Garrison⁽⁶⁹⁾ gives G values for initial yields from 1 M glycine at pH = 6.4 [$G(\text{NH}_3) = 4.3$; $G(\text{ketoacid}) = 2.1$; $G(\text{fatty acid}) = 1.2$; $G(\text{aldehyde}) = 0.5$; and $G(\text{H}_2) = 2.0$], and concludes that deamination by both e_{aq}^- and OH occur. Maxwell et al.⁽⁷⁰⁻⁷²⁾ note that formaldehyde and, especially, glyoxalic acid yields decrease with increasing dose.

To transpose our discussion on the effects of chelating agents and their radiolytic degradation products to the situation in the tank: Little reaction with the solvated electrons or their precursors is expected to occur for EDTA, NTA, IDA, or glycine. Some e_{aq}^- reactions with these chelators may

lead to deamination; thus, ammonia is a reasonable product to expect. Nonetheless, because NO_3^- and NO_2^- will efficiently compete for solvated electrons, the yield will be low. A small fraction of the H atoms (and OH radicals) will probably attack these species (and other organic substrates) to produce H_2 (and water). Two of the α -carboxy radicals thus produced will disproportionate and yield carbonyl-containing compounds. However, in the tank, many of the α radicals may react with NO_3 , NO_2 , NO , or O_2 ; little information on such reactions is available.

6. EFFECTS OF HIGH-SOLUTE CONCENTRATIONS ON RATES

The high-solute concentration in the tank will have several effects on rates of reactions and, therefore, may indirectly affect yields of products. Three of these effects might be significant and will, therefore, be explored here. These include viscosity, ionic strength, and the effect of time from the event of absorption of the radiation energy on the rate constants. A fourth effect, quantum mechanical tunneling, will be mentioned only briefly. In ideal (relatively dilute) solutions, these effects are well understood (see, e.g., reference 73) and their influence can be accurately estimated, but the high-salt concentrations in the tank generally complicate the assessment of this influence.

A. Viscosity Effects

In general, it can be shown that a rate constant, k , for any reaction can be expressed as the sum of reciprocals of the diffusion-controlled rate constant, k_D , and the activation-controlled rate constant, k_A .

$$\frac{1}{k} = \frac{1}{k_D} + \frac{1}{k_A} \quad (24)$$

Obviously, from this equation, no rate constant can exceed k_D no matter how large k_A may become. Although reasonable predictions of the diffusion-controlled rate constant may be made, no such estimates can be made *a priori* for activation-controlled rate constants.

Many of the reactions of the primary radicals with solutes or among themselves are diffusion controlled. Any parameter that will affect the rate of diffusion may, therefore, modify the rate of such reactions as well. The diffusion-controlled rate constant may be expressed (in $\text{M}^{-1} \text{s}^{-1}$) as

$$k_D = \frac{4\pi D_{ab} \beta N_0}{1000} \quad (25)$$

where $D_{ab} = D_a + D_b$ is the sum of the diffusion coefficients (in $\text{cm}^2 \text{s}^{-1}$) of the reacting species, N_0 is Avogadro's number (6.02×10^{23} molecules/mole), k is Boltzmann's constant (1.38×10^{-16} erg deg $^{-1}$) and

$$\beta^{-1} = \int_R^\infty \left(\exp \frac{V(r)}{kT} \right) r^2 dr \quad (26)$$

$V(r)$ and R are the potential energy between the two reactants and the radius of reaction, respectively. If at least one of the reactants is uncharged, then $\beta^{-1} = R^{-1}$ from equation 26. Equation 25 then simplifies to

$$k_D = \frac{4\pi D_{ab} R N_0}{1000} \quad (27)$$

Typical diffusion coefficients are in the range of $2 \times 10^{-5} \text{ cm}^2 \text{s}^{-1}$, whereas for the solvated electron $D = 4.7 \times 10^{-5} \text{ cm}^2 \text{s}^{-1}$. Reaction radii are often estimated as the sum of the ionic radii of the reactants or estimated from bond lengths ($R = R_a + R_b$; $r_{e\text{aq}} = 2.5 \text{ \AA}$; often $r_b = 2 \text{ \AA}$). Thus, in

agreement with experiment, $k_D = 2.3 \times 10^{10} \text{ M}^{-1} \text{ s}^{-1}$ is a typical rate constant for diffusion-controlled reactions of the solvated electron with uncharged scavengers.

Equation 27 shows that the diffusion-controlled rate constant depends on both the diffusion coefficients and on the reaction radius. Because both are related to the viscosity, η , via the Stokes-Einstein relation ($D_a = kT/6\pi\eta R_a$), equation 27 may be further simplified (assuming $R_a = R_b = R/2$) to give

$$k_D = \frac{8kN_0T}{3\eta 1000} = \frac{8RT}{3\eta 1000} \quad (28)$$

With these assumptions, the rate constant is independent of the radii and, thus, of the species involved in the reaction. At 25 °C with $\eta = 1 \text{ cP} = 10^{-2} \text{ Poise}$, and $R = 8.31 \times 10^7 \text{ erg mol}^{-1} \text{ deg}^{-1}$, one obtains $k_D = 6.6 \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$. This estimate of a "universal" rate constant for a diffusion-controlled reaction may be compared with the more accurate one calculated from equation 27. Because the viscosity of the tank solution may approach large values,⁽⁷⁴⁾ the rate constants of many of the diffusion-controlled reactions will be smaller than the dilute-solution values quoted in the tables. However, because of the lack of specificity in the dependence of rate on diffusion, the relative rates of many of these reactions are not expected to change significantly. Thus, the viscosity may influence individual rate constants, but it should exert only a small effect on the final yields of products. Because, to a first approximation, the viscosity decreases exponentially with increasing temperature, k_D exhibits an Arrhenius-like temperature dependence, equation 29a.

$$\eta \approx A \exp\left(\frac{E_a}{RT}\right) \quad (29a)$$

$$k_D = \left(\frac{8RT}{3000A}\right) \exp\left(\frac{-E_a}{RT}\right) \quad (29b)$$

Typical activation energies for diffusion are $\sim 14 \text{ kJ mol}^{-1}$. At the prevailing temperature in the tank ($\sim 60 \text{ °C}$), individual diffusion-controlled reaction rate constants will therefore increase by a factor of ~ 3 relative to their rate constants at 25 °C. Again, the relative rates will not be significantly altered.

Equation 27 is valid only for reactions in which at least one reactant is uncharged. For reactions between ions, the term β in equations 25 and 26 should be evaluated. This term can be expressed as

$$\beta = -\frac{z_a z_b r_0}{\left\{1 - \exp\left(\frac{z_a z_b r_0}{R}\right)\right\}} \quad (30)$$

where z_i are the charges of the ions, $r_0 = e^2/\epsilon kT = 7.1 \times 10^{-8} \text{ cm}$ for water at 25 °C, e is the charge on the electron, $4.80 \times 10^{-10} \text{ esu}$, and ϵ is the macroscopic dielectric constant of the medium (78.6 for water at 25 °C). Equation 25, then, will assume the form

$$k_D = -\frac{4\pi D_{ab} z_a z_b r_0 N_0}{\left\{1000 \left[1 - \exp\left(\frac{z_a z_b r_0}{R}\right)\right]\right\}} \quad (31)$$

Many of the reactions of the primary products of water radiolysis have been evaluated according to this equation, in particular those of the hydrated electron.⁽⁷⁵⁾ Good agreement between the predicted and measured values of k_D is observed when the reaction is between two oppositely charged ions. For ions of the same charge, estimates according to equation 31 are very sensitive to the choice of the value of R .⁽⁷³⁾

B. Ionic Strength Effects

High-electrolyte (inert or otherwise) concentrations may affect the rates of reactions between ions, whether they are diffusion or activation controlled. The well-known formalism developed by Debye and Huckel will suffice here, although more sophisticated models have also been developed (e.g., Harned and Owens⁽⁷⁶⁾). In principle, the effect arises from the fact that counterions accumulate in the vicinity of the reactant ions. Thus, an ionic atmosphere is created around the reactant ions that will screen some of the electrostatic interaction between them. For a diffusion-controlled reaction, this is equivalent to modifying $V(r)$ in equation 26. For an activation-controlled reaction, a similar Boltzmann term appears in k_A because the reactants themselves, both of which are charged ions, will assume a distribution around one another that is different from the bulk homogeneous distribution. Their concentration in the vicinity of one another is, therefore, different from their bulk concentration. As a result of the formation of the ionic atmosphere around the reactant ions, reactions between similarly charged ions will increase in rate, because the repulsive interaction between them will be partially screened at high ionic strength. Conversely, for oppositely charged ions, the rate of reaction will decrease at high ionic strength.

Quantitatively, the Debye Huckel treatment leads to

$$k = k^0 \exp(z_a z_b r_0 \kappa) , \quad (32)$$

where k^0 is the rate constant (diffusion or activation controlled) at zero ionic strength, and z_a, z_b, r_0 were defined in the previous section. The parameter κ^{-1} , the so-called Debye radius, is often regarded as the radius of the ionic atmosphere around the ion, and is given by

$$\kappa^2 = \frac{8\pi e^2 N_0 \rho I}{(1000 \epsilon kT)} , \quad (33)$$

where ρ is the solution density and I is the ionic strength (defined as $I = (1/2) \sum c_i z_i^2$; c_i is the molal concentration of the i -th ion, and z_i is its charge). For water at 25 °C (where $\kappa^2 = 1.08 \times 10^{15} \cdot I \text{ cm}^{-2}$) equation 32 translates to

$$\log k = \log k^0 + 1.02 z_a z_b I^{1/2} \quad (34)$$

This equation is obeyed up to concentrations of approximately 10^{-2} M. A somewhat more precise description, which is obtained by taking into account the finite size of the ions, is

$$\log k = \log k^0 + \frac{1.02 z_a z_b I^{1/2}}{(1 + I^{1/2})} . \quad (35)$$

The latter often holds up to 0.1 M concentrations. Above this concentration, the effect of ionic strength is weaker than that predicted from equation 35. For all practical purposes, for the ionic concentrations that exist in the tank, any electrostatic interactions will be completely screened by the high ionic strength, and the reactions between charged ions will probably proceed with rate constants that are typical for uncharged species (excluding specific interactions, complex formation, etc.).

As emphasized in preceding sections, many of the reactions of the primary radiolytic radicals occur very rapidly. It is relevant, therefore, to question whether the ionic atmosphere around an instantaneously created ion will develop prior to its reaction with a scavenger. The time required to develop the ionic atmosphere, τ , is approximately equal to the time required for an ion to diffuse a distance equal to the radius of the ionic atmosphere, i.e., κ^{-1} . To calculate τ , the Einstein equation of diffusion,

$$\tau = \frac{1}{(4Dk^2)} \quad (36)$$

can be used. For $I = 1 \text{ M}$ and a typical D of $2 \times 10^{-5} \text{ cm}^2 \text{ s}^{-1}$, one obtains $\tau \approx 1 \times 10^{-11} \text{ s}$ for the ionic atmosphere to develop. In comparison, the half-lifetime for the reaction of e_{aq}^- with NO_3^- at 1 M will be $\tau_{1/2} = 0.69/(1.8 \times 10^{10} \times 1) \approx 3.8 \times 10^{-11} \text{ s}$ (at this high concentration $k = 1.8 \times 10^{10} \text{ M}^{-1} \text{ s}^{-1}$ and not the value in Table 2). It seems, therefore, that the ionic atmosphere will relax around the solvated electron at a rate that is only slightly faster than the reaction rate. Somewhat slower relaxation times of the ionic atmosphere are predicted by the Falkenhagen equation:⁽⁷⁷⁾

$$\tau = \frac{6.92 \times 10^{-9}}{(c\Lambda_{\infty})} \quad (37)$$

where c is the molar concentration of the counterions, and Λ_{∞} is their molar conductance at infinite dilution. Since relaxation of the ionic atmosphere always occurs under attractive electrostatic interaction, it will usually develop before any reaction occurs between ions of similar charges. Thus, the ionic relaxation around e_{aq}^- is more rapid than the rate of its reaction with NO_3^- . However, for the reaction between oppositely charged ions (e.g., the hydrated electron and metal ions), the reactions at high-salt concentrations often occur before the full ionic atmosphere around e_{aq}^- has developed.⁽⁷⁸⁾ How would the rate constant for an ion, for which the ionic atmosphere has not yet attained its equilibrium configuration, change? This has been only marginally investigated,⁽⁷⁹⁾ but it seems reasonable to assume a geometric average of the rate constant at the high ionic strength and in dilute solutions for the unrelaxed ion:

$$k' = (k_0 k)^{1/2} \quad (38)$$

where k' is the rate constant for the unrelaxed ion and k is the rate constant calculated from equation 35.

To summarize the implications of high ionic strength of the waste solutions in yields: the only charged primary radicals of concern in this respect are e_{aq}^- and $\text{O}^\cdot$. For the former, because the overwhelming scavengers for e_{aq}^- are nitrate and nitrite, a similar increase in rate may be expected for both. Thus, the fate of e_{aq}^- will hardly change in the concentrated solution. For $\text{O}^\cdot$, the major scavenger is NO_2^- , and the rate of this reaction will increase relative to the rate in dilute solutions. Secondary scavengers for $\text{O}^\cdot$ are the organic chelators, and their rate will also increase relative to dilute solutions because these chelators exist in the waste solution as negatively charged ions. In general, it may be concluded that the reactivity of $\text{O}^\cdot$ with either NO_2^- or the organic chelators will approach that of its acid form, the OH radical.

C. Time-Dependent Rate Constants

The fact that many of the reactions of interest occur on a very short time scale introduces considerations that are commonly neglected in classical kinetics. One such effect is the so-called time dependence of the rate constant. When reactants (e.g., solvated electrons) are instantaneously generated by radiolysis in a solution, their original distribution within the solution will be completely random. Hence, some of the reactants will be created very close to a scavenger molecule whereas others will be generated far from any reactant. Those reactants that are generated near a scavenger will react faster, and the population of those created close to scavengers will decrease more rapidly than those of the more distant species. Eventually, a steady-state situation will be achieved around the various reactants in which the concentration gradient of the scavenger around the reactant is identical for all reactant molecules throughout the solution. At earlier times, the rate constants are

time dependent. The considerations discussed in the two previous sections apply. This problem has been discussed in detail by Noyes.⁽⁸⁰⁾ His expression for the rate constant at time t , k_t , can be recast as

$$k_t = k_{\infty} \left(1 + \frac{k_{\infty}}{\left(4\pi R D_{ab} \frac{N}{1000} \right)} \times \frac{R}{(\pi D_{ab} t)^{1/2}} \right), \quad (39)$$

where k_{∞} is the rate constant at long times (those given in the tables), and the other symbols have their customary meanings. Experimentally, these effects have been observed for many scavengers of the hydrated electron.⁽⁷⁸⁾ The faster the reaction (higher concentrations and larger k_{∞}), the more significant the effect. For uncharged species (measured for nitromethane, $D_{ab} = 6.5 \times 10^{-5} \text{ cm}^2 \text{ s}^{-1}$; $R = 6.0 \times 10^{-8} \text{ cm}$; $k_{\infty} = 2.9 \times 10^{10} \text{ M}^{-1} \text{ s}^{-1}$) at 1 M, the rate constant at the half-lifetime for the reaction ($t = 2.4 \times 10^{-11} \text{ s}$) is nearly twice k_{∞} . For slower reaction rates, the effect is smaller (e.g., for acetone $D_{ab} = 6.5 \times 10^{-5} \text{ cm}^2 \text{ s}^{-1}$; $R = 5.0 \times 10^{-8} \text{ cm}$; $k_{\infty} = 0.85 \times 10^{10} \text{ M}^{-1} \text{ s}^{-1}$; at 1 M $k_t = k_{\infty}$ within a few percent at the half-time). When both reactants are charged, the electrostatic effects discussed in the two preceding sections have to be taken into account in estimating k_{∞} , but otherwise the considerations are identical. For nitrate at 1 M concentration, k_t at the half-life is twice k_{∞} .⁽⁷⁸⁾

D. Quantum Mechanical Effects

The effects discussed in sections 6A-6C are all classical effects in which quantum mechanics play no role. However, the electron is a quantum particle, and quantum effects will appear when the reaction of interest is an electron transfer reaction. It is now well established that the electron can be transferred over long distances, much longer than the sum of the radii of the two reaction species. Consequently, situations may arise in which the electron is transferred with no physical collision between the reactants. When the diffusion of the two reactants towards one another is relatively slow (e.g., two ions of opposing charges), and when a strong driving force for the electron to be transferred exists, the electron may be transferred, via a tunneling mechanism, even though the reactants are quite far apart. This may occur regardless of the solute concentration. Long-range electron transfer by such a mechanism is currently under active investigation in many laboratories. Our present assessment has little implication on the processes in the tank solutions. Nevertheless, we evaluate this aspect of the chemistry because it implies that large radii, much larger than could be predicted from the sum of the radii of the reactants, may be necessary to predict rate constants by using the phenomenological description of equation 25. Indeed, very large reaction radii are required in order to explain the rates of reactions of the hydrated electron with several negatively charged ions.⁽⁷⁵⁾

When a reaction between two radicals is evaluated, it should be recognized that only encounters in which the two unpaired spins produce a singlet state could be productive. Statistically, only 25% of all the encounters could lead to a singlet radical-pair. This means that the diffusion-controlled rate constant, k_D (equation 24), has to be multiplied by this factor (0.25) when the rate constant for such a reaction is estimated. However, if the spin-lattice relaxation time of the radicals is much shorter than the encounter time (10^{-8} - 10^{-10} s) the spins will reorient during the encounter and each encounter will yield products. The latter is the situation for OH radicals (relaxation time $< 10^{-9} \text{ s}$). Hence, most reactions of OH radicals will not be hindered by the spin requirement. The majority of radicals, however, have spin-lattice relaxation times in the μs time regime and the maximum reaction rate constants achievable by these radicals will be only a quarter of the diffusion-controlled rate constant.

7. EFFECTS OF TEMPERATURE AND PRESSURE ON YIELDS AND RATES

The chemistry in the tank solutions occurs at approximately 60 °C; hence, the effects of temperature on yields and reaction rates of the primary radicals were also surveyed. Extensive studies have been performed on these effects over a broad temperature range,⁽¹⁶⁾ as they are important in pressurized-water reactor systems. However, these studies were limited to relatively low solute concentrations. Because many of the reactions relevant to this survey are close to the diffusion-controlled limit, and because the physical properties of water do not significantly change in the limited temperature range that exists in the tank, the effect of temperature is expected to be minimal. However, the secondary reactions that occur below the diffusion-control limit may exhibit pronounced temperature effects.

The effect of temperature on the sum of yields $\Sigma G(e_{aq}^-) + G(H) + G(OH)$ has been measured over the temperature range 20-200 °C.⁽⁸¹⁾ These studies establish that the spur size extends to larger distances at higher temperatures. This is mostly a result of the lower density of water at higher temperatures, which, in turn, increases the thermalization length of the ejected electrons. Consequently, yields of primary radicals, at times when spur processes are complete, will be more dependent upon temperature than the yields at early times. In other words, the concentration of radicals in the spur is lower at higher temperature, and, therefore, intraspur reactions (e.g., reactions 4-6 for dihydrogen production) are somewhat slower at higher temperature. At the temperature range of the waste solution, the sum of yields of primary radicals that escape the spur increases by 10% upon increasing the temperature from 20 to 60 °C.⁽⁸¹⁾ It was also concluded that $G(e_{aq}^-) + G(H) \approx G(OH)$, and thus the increase in the yield of the reducing radicals is also ~10%. More importantly, because the waste solutions are highly concentrated, most of the radicals will react with the scavengers within the spur, and their escape yield will change little over the same temperature range. In summary, we expect no effect of the elevated temperature on the yield of primary radicals.

Activation energies for some of the rate constants of the primary radical reactions have been measured and are summarized in Table 7. A more comprehensive compilation of activation energies for reactions of radiolytic processes can be found in reference 16. When the reaction is near the diffusion-control limit (but somewhat slower than the limit), the effect of temperature on both k_D and k_A has to be taken into account. This can be done using equations 24 and 31. The dependence of k_D on temperature enters, then, through the dependence of the diffusion coefficients, D_{ab} , and the parameter r_0 (equation 31), on temperature. For reactions between ions, the Debye radius, κ^{-1} (equations 32 and 33), is also temperature dependent. The best agreement with experimental data was obtained when the temperature dependence of the diffusion coefficients of the primary radicals was assumed to be identical to that of the self diffusion of water.⁽⁸⁷⁾ A simplified treatment that assumes an activation energy for diffusion of $E_{aD} = 12.6 \text{ kJ mol}^{-1}$ often yields reasonable agreement with experiments when the reaction is truly diffusion controlled.⁽⁸⁷⁾ In equation 24, the dependence of k_A on temperature is the usual Arrhenius dependence. For the cases where this analysis was employed, k_A and its energy of activation are given in Table 7.

The value of the activation energy for the important reaction 11, which converts H atom to e_{aq}^- , is still controversial.^(84,85) We quote in Table 7 the most recent value. Activation energies for the important scavenging reaction of hydrated electrons by nitrate and nitrite (and also for $e_{aq}^- + e_{aq}^-$)⁽⁸⁶⁾ are difficult to quantify because they do not follow an Arrhenius dependence on temperature.⁽⁸⁷⁾ The rate constant for the reaction of NO_3^- with e_{aq}^- increases by a factor of two (1×10^{10} to $2 \times 10^{10} \text{ M}^{-1} \text{ s}^{-1}$), whereas that for the NO_2 increases even less (3.7×10^9 to $5.2 \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$) upon increasing the temperature from 20 to 60 °C.⁽⁸⁷⁾ The activation parameters for the recombination of e_{aq}^- to produce dihydrogen, given in Table 7, are valid only in the temperature range of 20 to 150 °C.

TABLE 7. Temperature Effects on Rate Constants for Reactions of the Primary Radical

Reaction	E_{a}^{a} kJ mol ⁻¹	E_{aA}^{a} kJ mol ⁻¹	$k_{\text{A}}(298 \text{ K})$ M ⁻¹ s ⁻¹	$k_{\text{obs}}(298 \text{ K})$ M ⁻¹ s ⁻¹	Reference
$\text{e}_{\text{aq}}^- + \text{e}_{\text{aq}}^-$	23.0			$2k = 1.1 \times 10^{10}$	86
$\text{H} + \text{H}$	14.6			$2k = 1.5 \times 10^{10}$	92
$\text{OH} + \text{OH}$		3.7	$2k = 1.0 \times 10^{10}$		87
$\text{HO}_2 + \text{HO}_2$		22.8	$2k = 1.6 \times 10^6$	<i>b</i>	87
$\text{OH} + \text{H}_2$		18.0	4.0×10^7	<i>b</i>	87
$\text{OH} + \text{H}_2\text{O}_2$		13.0	3.0×10^7	<i>b</i>	87
$\text{OH} + \text{CO}_3^{2-}$		23.6	4.2×10^8	<i>b</i>	87
$\text{OH} + \text{HCO}_3^-$		21.2	8.5×10^6	<i>b</i>	87
$\text{OH} + \text{FeCN}_6^{4-}$		7.0	4.0×10^{10}		87
$\text{OH} + \text{Cu}^{2+}$		13.3	3.4×10^8	<i>b</i>	87
$\text{OH} + \text{Fe}^{2+}$		9.2	4.6×10^8	<i>b</i>	87
$\text{OH} + \text{HCO}_2^-$		4.0	6.5×10^9		87
$\text{OH} + \text{MeOH}^{\text{c}}$	4.8			1.0×10^9	83
$\text{OH} + \text{i-PrOH}^{\text{c}}$	3.5			2.9×10^9	87
$\text{OH} + \text{t-BuOH}^{\text{c}}$	11.0			7.0×10^8	87
$\text{O}^- + \text{MeOH}^{\text{c}}$	2.8			8.6×10^8	83
$\text{O}^- + \text{i-PrOH}^{\text{c}}$	2.4			1.5×10^9	83
$\text{e}_{\text{aq}}^- + \text{O}_2$		11.5	6.5×10^{10}		87
$\text{e}_{\text{aq}}^- + \text{H}^+$		14.5	3.0×10^{10}		87
$\text{e}_{\text{aq}}^- + \text{Cd}^{2+}$		12.0	5.0×10^{11}		87
$\text{H} + \text{OH}^-$	38.5			2.4×10^7	84
$\text{H} + \text{O}_2$		6.2	3.2×10^{10}		87
$\text{H} + \text{H}_2\text{O}_2$		16.6	5.0×10^7	<i>b</i>	87

^a k_{A} is the activation-controlled rate constant and E_{aA} is the activation energy for k_{A} to be used in equation 31. $E_{\text{aD}} = 12.6 \text{ kJ mol}^{-1}$ is often used for the diffusion-controlled rate constant, see text.

^bThe reaction is slow enough to justify $k_{\text{obs}} = k_{\text{A}}$.

^cMeOH, i-PrOH, t-BuOH are methanol, isopropanol, and t-butanol, respectively.

Few other thermodynamic parameters for other relevant reactions have been measured. For reaction 40, ΔH is temperature dependent.⁽⁸²⁾ Experimentally, the pK_a of this reaction changes

from 11.85 at 20 °C to 10.81 at 80 °C, but most of this change could be accounted for by the change in K_w with temperature.⁽⁸³⁾ For the pK_a of the HO_2 radical, $\Delta H = 0$ has been estimated and a value of $\Delta H = -15.4$ kJ/mole has been determined for reaction 41.⁽⁸²⁾

Activation energies for hydrogen-atom abstraction reactions by OH or O^- from organic substrates (Table 7) are even smaller than the activation energy for self diffusion of the radical in water. This was taken to indicate that these reactions are slower than the diffusion-controlled limit and probably proceed via a short-lived intermediate.⁽⁸⁷⁾ Significantly, no information is available on the effect of temperature on the rate of dihydrogen production via hydrogen abstraction from organic compounds (reaction 42). Because the rate constants for this reaction are often smaller than

the diffusion-controlled limit, they may be expected to have relatively high-activation energies. The increase in temperature may, therefore, increase the rate of reaction 42 relative to the other reactions that H atoms undergo and thus increase the yield of H_2 . The activation parameters for the oxidation of a few phenolates by NO_2 radicals have been recently measured ($E_a = 31 \pm 4$ kJ mol⁻¹; $\log A = 13.4 \pm 0.4$).⁽⁵⁰⁾ No information is available on the effect of temperature on rates of other NO_x radicals.

It seems clear from this discussion that the effect of temperature on the rates of reactions of the primary radicals with the substrates available in the tank solution will not be large. However, the effect of temperature on hydrogen abstraction from organic substrates by H atom may be significant.

The pressure in various regions within the tank will be somewhat elevated due to hydrostatic forces. The density of the waste solution implies that the pressure at the bottom of the tank may be 2 atm. The question therefore arises as to the possible effects of the pressure on the yields and rates of the radiolytic processes. Several studies have been conducted to measure these effects.⁽⁸⁸⁻⁹¹⁾ In the range of $1-8 \times 10^3$ atm, no effect of pressure on $G(H_2)$, $G(H)$, or $G(e^-_{aq})$ could be observed in neutral solutions. Secondary reactions, including hydrogen abstraction by hydrogen atoms from organic substrates, do exhibit an increase in $G(H_2)$ upon increasing the pressure in this range. However, within the pressure ranges relevant to the tank solution, this increase is completely negligible. In summary, pressure effects should be of no concern in any consideration of dihydrogen yields. No documentation of the effects of pressure on NO_x yields was found in the literature.

8. CONCLUSIONS

Several points of importance that emanate from this survey will be summarized below in the same order in which they were discussed in the text.

a. Nonhomogeneous kinetics should be considered in analyzing the primary radiolytic reactions. Considerations based on homogeneous models are only marginally relevant to the concentration regime prevailing in the waste solutions.

b. The vast majority of the species produced radiolytically will convert to their basic forms. However, some scavenging reactions will efficiently compete with the neutralization reactions at early times. The major scavenger for e_{aq}^- and its precursors is NO_3^- , and for H and OH (O^-) it is NO_2^- .

c. Dioxygen will be produced by the direct effect of the radiation on nitrate and nitrite. The organic content cannot prevent O_2 formation by this pathway, but the presence of organic molecules may lead to the destruction of oxygen via organic radical reactions with O_2 in later steps.

d. Much information is available on the yields of H_2 , O_2 , and other products in nitrate solutions in the more conventional pH ranges. Less work has been done on nitrite systems and very little on mixtures of nitrate and nitrite. The general observations are clear:

High-nitrate and -nitrite concentrations lead to lower H_2 yields and higher O_2 yields; higher organic content always leads to higher H_2 yields. There is an empirical universal dependence of dihydrogen yields on scavenger reactivity, and this relationship may be used to estimate the yield. Aside from the direct production of H_2 by water radiolysis (i.e., from dissociation of water molecules or from recombination of the reducing radicals, all of which amounts to $G(H_2) < 0.06$), the only radiolytic pathway for the production of H_2 is via hydrogen abstraction by hydrogen atoms from organic substrates. Nitrate is so extremely efficient in scavenging e_{aq}^- and its precursors that at a concentration of ~ 1 M it will completely prevent H_2 production from e_{aq}^- sources. Nitrite, even though it is an efficient H-atom scavenger, is not so dominating. Other H-atom scavengers (organic compounds) may efficiently compete with nitrite.

e. While an extensive data base is available on the reactivity of the primary products of water radiolysis, very little is known about the reactivity of secondary radicals, in particular of the NO_x system, with organic substrates. This should be a major effort in future experiments. We hope the parallel literature search on thermal reactions of this system will provide more information on this aspect of the waste chemistry. Although there does not appear to be a convenient pathway to the conversion of these substances to dihydrogen, these substances may be converted to N_2O .

f. The effects of temperature, pressure, viscosity, and high ionic strength on yields of products and reactivity of the primary radicals are reasonably well understood. None of these factors are expected to appreciably affect the reactivity and yields. Most importantly, though, the effects of temperature on the reactivity of hydrogen atoms and of the secondary radicals (again, NO_x are the focal point) are not known, and they are expected to be strongly temperature dependent.

Some considerations of chemical (as distinct from physical or mechanical) remediation strategies, internal to the tank (as distinct from external), are appropriate at this point. In order to minimize hydrogen production, the intuitive approach would be to utilize a hydrogen-atom scavenger that will not lead to H_2 . Among such candidates, nitrite is undoubtedly the most attractive scavenger (assuming that nitrate is at the ~ 1 M concentration level). We, therefore, recommend that high priority be assigned to the analytical determination of nitrite concentration in the tank. Any depletion below the ~ 2 M level could be corrected with little additional deleterious effects. Other inorganic candidates pose a dilemma because they will necessarily be relatively strong oxidants that may lead to the undesirable generation of potentially explosive mixtures. Nonetheless, they may be explored on a laboratory scale prior to specific recommendation. Sulfide (polysulfide at high pH) was considered, but, from the purely radiolytic mechanistic point of view, it is believed that HS^- will be the scavenger for hydrogen atoms and its reaction will produce dihydrogen. Organic scavengers are considered less desirable because of less favorable economics and the chemical complexities that they may introduce. Hydrogen atoms will react with organic additives either via hydrogen abstraction (the reaction to avoid) or via addition to unsaturated bonds, olefinic or aromatic. Perchlorinated organic substrates, and perhaps aromatic derivatives, will eliminate the former route but their price, stability, solubility, and the chemical consequences (production of HCl is not a major concern) may be prohibitive.

REFERENCES

- (1) W. S. P. Tank, PNL-7595 (1991) "Chemical and Physical Processes in Tank 241-SY-101: A Preliminary Report".
- (2) H. J. Mockel and R. H. Koster, Nucl. Tech. (1982) 59, 494-7; "Gas Formation During the γ -Radiolysis of Cemented Low- and Intermediate-Level Waste Products".
- (3) G. V. Buxton, C. L. Greenstock, W. P. Helman and A. B. Ross, J. Phys. Chem. Ref. Data (1988) 17, 513-886; "Critical Review of Rate Constants of Hydrated Electron, Hydrogen Atom and Hydroxyl Radicals in Aqueous Solutions".
- (4) P. Neta, R. E. Huie and A. B. Ross, J. Phys. Chem. Ref. Data (1988) 17, 1027-284; "Rate Constants for Reactions of Inorganic Radicals in Aqueous Solution".
- (5) P. Wardman, J. Phys. Chem. Ref. Data (1989) 18, 1637-755; "Reduction Potentials of One Electron Couples Involving Free Radicals in Aqueous Solution".
- (6) D. M. Stanbury, Adv. Inorg. Chem. (1989) 33, 69-138; "Reduction Potentials Involving Inorganic Free Radicals in Aqueous Solution".
- (7) Farhataziz and M. A. J. Rodgers, (1987) VCH Publishers; "Radiation Chemistry: Principles and Applications".
- (8) J. W. Spinks and R. J. Woods, (1976) J. Wiley & Sons; "An Introduction to Radiation Chemistry".
- (9) I. G. Draganic and Z. D. Draganic, (1971) Academic Press; "The Radiation Chemistry of Water".
- (10) A. O. Allen, (1961) Van Nostrand Co.; "The Radiation Chemistry of Water and Aqueous Solutions".
- (11) F. S. Dainton and S. R. Logan, Trans. Faraday Soc. (1965) 61, 715-22; "Radiolysis of Aqueous Solutions Containing Nitrite Ions and Nitrous Oxide".
- (12) H. A. Schwarz, J. Phys. Chem. (1969) 73, 1928-37; "Applications of the Spur Diffusion Model to the Radiation Chemistry of Aqueous Solutions".
- (13) Y. Gauduel, S. Pommeret, A. Migus and A. Antonetti, Chem. Phys. (1990) 149, 1-10; "Some Evidence of Ultrafast H_2O^+ -Water Molecule Reaction in Femtosecond Photoionization of Pure Water: Influence on Geminate Pair Recombination".
- (14) Y. Gauduel, S. Pommeret, A. Migus and A. Antonetti, J. Phys. Chem. (1991) 95, 533-9; "H/D Isotope Effects on Femtosecond Electron Reactivity in Aqueous Media".
- (15) H. A. Mahlman, J. Chem. Phys. (1960) 32, 601-3; "Hydrogen Formation in the Radiation Chemistry of Water".
- (16) W. G. Burns and W. R. Marsh, J. Chem. Soc. Faraday Trans. 1 (1981) 77, 197-215; "Radiation Chemistry of High Temperature (300-410 °C) Water".
- (17) C. D. Jonah, M. S. Matheson, J. R. Miller and E. J. Hart, J. Phys. Chem. (1976) 80, 1267-70; "Yields and Decay of Hydrated Electron from 100 ps to 3 ns".
- (18) C. D. Jonah and J. R. Miller, J. Phys. Chem. (1977) 81, 1974-6; "Yield and Decay of OH Radical from 200 ps to 3 ns".
- (19) H. Y. Yamaguchi, Rad. Phys. Chem. (1989) 34, 801-7; "A Spur Diffusion Model Applied to Estimate Yields of Species in Water Irradiated by Nonhomogeneous Photons of 50 eV to 2 MeV".
- (20) G. Girija and C. Gopinathan, Radiat. Phys. Chem. (1982) 19, 107-12; "Comprehensive Calculations on the "Variable Size" Spur Diffusion Model of Aqueous Radiolysis".
- (21) C. N. Trumbore, D. R. Short, J. E. Fanning Jr. and J. H. Olson, J. Phys. Chem. (1978) 82, 2762-7; "Effects of Pulse Dose on Hydrated Electron Decay Kinetics in the Pulse Radiolysis of Water. A Computer Modeling Study".
- (22) R. K. Wolff, M. J. Bronskill and J. W. Hunt, J. Chem. Phys. (1970) 53, 4211-5; "Picosecond Pulse Radiolysis Studies. II. Reactions of Electrons with Concentrated Scavengers".
- (23) K. Y. Lam and J. W. Hunt, J. Phys. Chem. (1974) 78, 2414-6; "Yield of Solvated Electrons in the Aliphatic Alcohols at Picosecond Times".

- (24) C. D. Jonah, J. R. Miller and M. S. Matheson, *J. Phys. Chem.* (1977) 81, 1618-22; "The Reaction of the Precursor of the e_{aq}^- with Electron Scavengers".
- (25) K. Y. Lam and J. W. Hunt, *Int. J. Radiat. Phys. Chem.* (1975) 7, 317-38; "Picosecond Pulse Radiolysis Studies. VI. Fast Electron Reactions in Concentrated Solutions of Scavengers in Water and Alcohols".
- (26) A. J. Swallow and M. Inokuti, *Rad. Phys. Chem.* (1988) 32, 185-9; "Radiation Energy Partition Among Mixture Components: Current Ideas on an Old Question".
- (27) J. T. Kiwi and M. Daniels, *J. Inorg. Nucl. Chem.* (1978) 40, 576-9; "On the Radiolysis of Concentrated Alkaline and Calcium-Nitrate Solutions".
- (28) E. Hayon and M. Moreau, *J. Phys. Chem.* (1965) 69, 4058-62; "Reaction Mechanism Leading to the Formation of Molecular Hydrogen in the Radiation Chemistry of Water".
- (29) H. A. Schwarz, *J. Am. Chem. Soc.* (1955) 77, 4960-4; "The Effect of Solutes on the Molecular Yields in the Radiolysis of Aqueous Solutions".
- (30) A. J. Swallow, *Prog. Reaction Kinetics* (1978) 9, 195-366; "Reactions of Free Radicals Produced from Organic Compounds in Aqueous Solution by Means of Radiation".
- (31) M. Simic, P. Neta and E. Hayon, *J. Phys. Chem.* (1969) 4214-9; "Pulse Radiolysis of Aliphatic Acids in Aqueous Solutions. II. Hydroxy and Polycarboxylic Acids".
- (32) H. A. Mahlman, *J. Phys. Chem.* (1960) 64, 1598; "The G(OH) in the ^{60}Co Radiolysis of Aqueous NaNO_3 Solutions".
- (33) M. Daniels and E. E. Wigg, *J. Phys. Chem.* (1969) 73, 3703-9; "Radiation Chemistry of the Aqueous Nitrate System. II. Scavenging and pH Effects in Co-60 γ Radiolysis of Concentrated Sodium Nitrate Solutions".
- (34) S. P. Kulkarni and A. N. Garg, *J. Radioanal. Nucl. Chem. Lett.* (1991) 153, 211-9; "An Evaluation of Rate Constants for Radiation Induced Decomposition of Alkali Metal Nitrates".
- (35) a. B. G. Brodda, S. Dix, and E. R. Merz, *Analyst* (1989) 114, 335-7; "Radiolytic Gas Formation in High-level Liquid Waste Solutions".
b. E. J. Hart and N. E. Bibler, (1991) Independent unpublished results on the effect of organic solutes on the radiolytic production of N_2O in concentrated nitrate solutions, presented at the "Tank Waste Science Panel" meeting; Denver, CO; February 7, 1991.
- (36) M. Chouraqui and J. Sutton, *Trans. Faraday Soc.* (1966) 62, 2111-21; "Origin of Primary Hydrogen Atom Yield in Radiolysis of Aqueous Solutions".
- (37) C. Baquey, J. C. Roux and J. Sutton, *J. Phys. Chem.* (1970) 74, 4210-6; "Radiolysis of 1 M Aqueous Ethanol Solutions in Potassium Nitrate".
- (38) M. Fiti and R. Morar, *Radiochem. Radioanal. Lett.* (1980) 43, 137-46; "Radiolytic Oxidation of Al^{2+} Intermediate".
- (39) J. Kalecinski, *Bull. Acad. Pol. Sci., Ser. Sci. Chim.* (1970) 18, 263-9; "The Hydrated Electron in Reaction of Aluminum with Aqueous Solution of Alkali Hydroxides".
- (40) M. Gratzel, A. Henglein, J. Lilie and G. Beck, *Ber. Bunsenges. Phys. Chem.* (1969) 73, 646-53; "Pulse Radiolytic Study of the Elementary Processes of Oxidation and Reduction of Nitrite Ions".
- (41) M. Gratzel, A. Henglein and S. Taniguchi, *Ber. Bunsenges. Phys. Chem.* (1970) 74, 292-6; "Pulse Radiolytic Investigation of the Reduction of NO_3^- Ions and the Buildup and Decay of Pernitrous Acid in Aqueous Solutions".
- (42) M. Gratzel, S. Taniguchi and A. Henglein, *Ber. Bunsenges. Phys. Chem.* (1970) 74, 488-92; "Pulse Radiolytic Investigation of the Oxidation of NO and the Equilibrium $\text{N}_2\text{O}_3 \rightleftharpoons \text{NO} + \text{NO}_2$ in Aqueous Solutions".
- (43) M. Gratzel, S. Taniguchi and A. Henglein, *Ber. Bunsenges. Phys. Chem.* (1970) 74, 1003-10; "Pulse Radiolytic Investigation of the Short-lived Intermediates of the Reduction of NO in Aqueous Solutions".
- (44) W. A. Seddon and M. J. Joung, *Can. J. Chem.* (1970) 48, 393-4; "Pulse Radiolysis of Nitric Oxide in Aqueous Solution".
- (45) W. A. Seddon and H. C. Sutton, *Trans. Faraday Soc.* (1963) 59, 2323-33; "Radiation Chemistry of Nitric Oxide Solutions".

- (46) R. J. Knight and H. C. Sutton, *Trans. Faraday Soc.* (1967) 63, 2628-39; "Radiolysis of Aqueous Solutions of Nitric Oxide".
- (47) P. Neta and R. E. Huie, *J. Phys. Chem.* (1986) 90, 4644-8; "Rate Constants for the Reactions of NO₃ Radicals in Aqueous Solutions".
- (48) L. V. Shastri and R. E. Huie, *Int. J. Chem. Kinet.* (1990) 22, 505-12; "Rate Constants for Hydrogen Abstraction Reactions of NO₃ in Aqueous Solution".
- (49) W. K. Wilmarth, D. M. Stanbury, J. E. Byrd, N. H. Po and C. P. Chuba, *Coord. Chem. Rev.* (1983) 51, 155; "Electron Transfer Reactions Involving Simple Free Radicals".
- (50) Z. B. Alfassi, R. E. Huie, P. Neta and L. C. T. Shoute, *J. Phys. Chem.* (1990) 94, 8800-5; "Temperature Dependence of the Rate Constants for Reactions of Inorganic Radicals with Organic Reductants".
- (51) Z. B. Alfassi, R. E. Huie and P. Neta, *J. Phys. Chem.* (1986) 90, 4156-8; "Substituent Effects on Rates of One Electron Oxidation of Phenols by the Radicals ClO₂, NO₂, and SO₃".
- (52) E. Barnes, *J. Chem. Soc.* (1931) 2605-20; "The Action of NO on Alkaline Hydroxide".
- (53) R. Buitenhuis, C. M. N. Bakker, F. R. Stock and P. W. F. Louwrier, *Radiochim. Acta* (1977) 24, 189-92; "Rate Constants for the Reaction of e_(aq)⁻ with EDTA and Some Metal EDTA-Complexes".
- (54) S. N. Bhattacharyya and K. P. Kundu, *Int. J. Radiat. Phys. Chem.* (1972) 4, 31-41; "X-Irradiation of Aqueous Solutions of Ethylenediamine Tetraacetic Acid (EDTA)".
- (55) Q. G. Mulazzani, M. Venturi and M. Z. Hoffman, *J. Phys. Chem.* (1985) 89, 722-8; "Radiolytically Induced One-Electron Reduction of Methylviologen in Aqueous Solution. Reactivity of EDTA Radicals Toward Methylviologen".
- (56) R. Poupko, B. L. Silver and A. Lowenstein, *Chem. Comm.* (1968) 453; "Study of Radicals Derived from Glycine and Alanine in Solution".
- (57) S. N. Bhattacharyya and E. V. Srisankar, *Int. J. Radiat. Phys. Chem.* (1976) 8, 667-71; "Radiolysis of Aqueous Solutions of Nitrilotriacetic Acid".
- (58) S. N. Bhattacharyya and N. C. Saha, *Radiat. Res.* (1976) 68, 234-43; "Gamma Radiolysis of Aqueous Solutions of Iminodiacetic Acid".
- (59) J. Lati and D. Meyerstein, *J. Chem. Soc., Dalton Trans.* (1978) 1105-18; "Oxidation of First-Row Bivalent Transition-Metal Complexes Containing Ethylenediaminetetra-acetate and Nitrilotriacetate Ligands by Free Radicals: A Pulse-Radiolysis Study".
- (60) P. Neta and R. W. Fessenden, *J. Phys. Chem.* (1970) 74, 2263-6; "Electron Spin Resonance Study of Deamination of Amino Acids by Hydrated Electrons".
- (61) V. Markovic, D. Nikolic and O. I. Micic, *Int. J. Radiat. Phys. Chem.* (1974) 6, 227-32; "Pulse-Radiolysis of Glycine and Alanine at pH 0-7".
- (62) M. Simic and E. Hayon, *Radiat. Res.* (1971) 48, 244-55; "Reductive Deamination of Oligopeptides by Solvated Electrons in Aqueous Solution".
- (63) J. E. Aldrich, K. Y. Lam, P. C. Shragge and J. W. Hunt, *Radiat. Res.* (1975) 63, 42-52; "Fast Electron Reactions in Concentrated Solutions of Amino Acids and Nucleotides".
- (64) P. Neta, M. Simic and E. Hayon, *J. Phys. Chem.* (1972) 76, 3507-8; "On the pK_a of the ⁺H₂NCHCO₂H Radical".
- (65) P. Neta and R. H. Schuler, *J. Phys. Chem.* (1975) 79, 1-6; "Rate Constants for the Reaction of O⁻ Radicals with Organic Substrates in Aqueous Solution".
- (66) P. Neta and R. H. Schuler, *Radiat. Res.* (1971) 47, 612-27; "Rate Constants for Reaction of Hydrogen Atoms with Compounds of Biochemical Interest".
- (67) E. Hayon and J. Weiss, *J. Chem. Soc.* (1960) 5091-7; "The Chemical Action of Ionising Radiations in Solution. Part XXIV. Action of X-Rays (200 kV) on Aqueous Solutions of Acetic Acid and Glycolic Acid".
- (68) P. M. Grant and R. B. Ward, *J. Chem. Soc.* (1959) 2654-9; "Effects of γ-Radiation, Part II. Characterisation of the Product from Glycolic Acid".
- (69) W. M. Garrison, in "Current Topics in Radiation Research", Ed. M. Ebert and A. Howard, (1968) 43-94; "Radiation Chemistry of Organo-Nitrogen Compounds".

- (70) C. R. Maxwell, Radiat. Res. (1964) 4, 175-8; "Addendum: Radiolysis of Aqueous Glycine".
- (71) C. R. Maxwell, D. C. Peterson and N. E. Sharpless, Radiat. Res. (1954) 1, 530-45; "The Effect of Ionizing Radiation on Amino Acids I. The Effect of X-rays on Aqueous Solutions of Glycine".
- (72) C. R. Maxwell, D. C. Peterson and W. C. White, Radiat. Res. (1955) 2, 431-8; "The Effect of Ionizing Radiation on Amino Acids III. The Effect of Electron Irradiation on Aqueous Solutions of Glycine".
- (73) R. E. Weston and H. A. Schwarz, (1972) Prentice-Hall, Inc.; "Chemical Kinetics".
- (74) B. M. Mauss, Rockwell International Internal Letter No. 65453-86-079 (1986) "101-SY Samples: Laboratory Analysis and Results".
- (75) M. Anbar and E. Hart, Adv. Chem. Ser. (1968) 81, 79-94; "On the Reactivity of Hydrated Electrons towards Inorganic Compounds".
- (76) H. S. Harned and B. B. Owens, (1958) Van Nostrand; "Chemistry of Electrolyte Solutions".
- (77) H. Falkenhagen, (1971) Verlag; "Theory of Electrolytes".
- (78) C. D. Jonah, J. R. Miller, E. J. Hart and M. S. Matheson, J. Phys. Chem. (1975) 79, 2705-11; "Picosecond Pulse Radiolysis. I. Time or Concentration Dependent Rate Constants".
- (79) P. J. Coyle, F. S. Dainton and S. R. Logan, Proc. Chem. Soc. (1964) 219; "The Probable Relaxation Time of the Ionic Atmosphere of the Hydrated Electron".
- (80) R. N. Noyes, Progress in Reaction Kinetics (1961) 1, 129-60; "Effects of Diffusion Rates on Chemical Kinetics".
- (81) G. V. Buxton and N. D. Wood, Rad. Phys. Chem. (1989) 34, 699-704; "Effect of Temperature and Scavenging Power on the Sum of $G(e_{aq}^-) + G(H) + G(OH)$ in the Range 20-200 °C. A Pulse Radiolysis Study".
- (82) G. V. Buxton, N. D. Wood and S. Dyster, J. Chem. Soc., Faraday Trans. 1 (1988) 84, 1113-21; "Ionization Constants of OH and HO₂ in Aqueous Solution up to 200 °C".
- (83) A. J. Elliot and D. R. McCracken, Rad. Phys. Chem. (1989) 33, 69-74; "Effect of Temperature on O⁻ Reactions and Equilibria: A Pulse Radiolysis Study".
- (84) P. Han and D. M. Bartels, J. Phys. Chem. (1990) 94, 7294-9; "Reevaluation of Arrhenius Parameters for $H + OH^- \rightarrow e_{aq}^- + H_2O$ and the Enthalpy and Entropy of Hydrated Electrons".
- (85) B. Hickel and K. Sehested, J. Phys. Chem. (1985) 89, 5271-4; "Activation Energy for the Reaction of $H + OH^- \rightarrow e_{aq}^-$. Kinetic Determination of the Enthalpy and Entropy of Solvation of e_{aq}^- ".
- (86) H. Christensen and K. Sehested, J. Phys. Chem. (1986) 90, 186-90; "The e_{aq}^- and Its Reactions at High Temperature".
- (87) A. J. Elliot, D. R. McCracken, G. V. Buxton and N. D. Wood, J. Chem. Soc., Faraday Trans. (1990) 86, 1539-47; "Estimation of Rate Constants for Near-diffusion-controlled Reactions in Water at High Temperatures".
- (88) R. R. Hentz and D. W. Brazier, J. Chem. Phys. (1971) 54, 2777-80; "γ Radiolysis of Liquids at High Pressures. X. The Reaction $H + OH^-$ and the Partial Molal Volume of the Hydrated Electron".
- (89) R. R. Hentz and R. J. Knight, J. Chem. Phys. (1970) 52, 2456-9; "γ Radiolysis of Liquids at High Pressures. VIII. Primary Yields at 8.7 kbar and Reactions of the Hydrated Electron with H₂O and H₂O⁺".
- (90) R. R. Hentz, Farhatziz and D. J. Milner, J. Chem. Phys. (1967) 47, 5381-4; "γ Radiolysis of Liquids at High Pressures. V. Reaction of the Hydrated Electron with Water".
- (91) R. R. Hentz, Farhatziz and D. J. Milner, J. Chem. Phys. (1967) 47, 4865-7; "γ Radiolysis of Liquids at High Pressures. IV. Primary Yields in Neutral Aqueous Solutions".
- (92) K. Sehested and H. Christensen, Rad. Phys. Chem. (1990) 36, 499-500; "The Rate Constant of the Bimolecular Reaction of H Atoms at Elevated Temperatures".
- (93) B. Hickel and K. Sehested, J. Phys. Chem. (1991) 95, 744-7; "Activation Energies for the Reactions $O^- + H_2$ and $O^- + D_2$ in Aqueous Solution".

Distribution for ANL-91/40

Internal:

F. Barnabas
F. A. Cafasso
E. Cerny
Y.-D. Cheng
H. Diamond
E. P. Horwitz
C. D. Jonah
M. S. Matheson

D. Meisel (29)
M. C. Sauer, Jr.
L. M. Stock
J. C. Sullivan
Y. Vojta
ANL Patent Dept.
ANL Contract File
TIS Files (3)

External:

DOE-OSTI for distribution per (UC-601) (12)
ANL Library
Manager, Chicago Operations Office, DOE
G. D. Johnson, Westinghouse Hanford Company (40)