Old and New Physics in Nucleon Spin Structure
DOE/ER/40561--30
DE92 004385

Harry J. Lipkin

*Department of Physics, Weizmann Institute of Science
Rehovot, Israel*

and

*Institute for Nuclear Theory, University of Washington
HN-12, Seattle, Washington 98195, USA*

PREPARED FOR THE U.S. DEPARTMENT OF ENERGY
UNDER GRANT DE-FG06-90ER40561

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, mark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

OLD AND NEW PHYSICS IN NUCLEON SPIN STRUCTURE

Harry J. Lipkin

*Department of Physics, Weizmann Institute of Science, Rehovot,
Israel*

and

*Institute for Nuclear Theory, University of Washington,
Seattle, WA 98195*

ABSTRACT

EMC implies quarks carry very little of the proton spin. $(G_A/G_V)_{n \rightarrow p}$ implies that valence quarks carry 70% of the spin, but disagrees with SU(6). $(G_A/G_V)_{\Sigma^- \rightarrow n}$ agrees strikingly with SU(6). All semileptonic decay data can be fit by SU(3), but no model fits the data by breaking SU(6) without also breaking SU(3). Considerable data on hadron masses and magnetic moments are fit by a simple constituent quark model with only constituent quarks. A toy model for the proton with valence quarks and sea fits nearly everything.

TABLE 1. Theoretical Predictions and Experimental Values of G_A/G_V

DECAY	Simple SU(6)	Constituent SU(6)	SU(3)	Experiment
$n \rightarrow p$	5/3	<i>input</i>	<i>input</i>	1.261 ± 0.004
$\Lambda \rightarrow p$	1	0.756 ± 0.003	0.727 ± 0.007	0.718 ± 0.015
$\Xi^- \rightarrow \Lambda$	1/3	0.252 ± 0.001	0.193 ± 0.012	0.25 ± 0.05
$\Sigma^- \rightarrow n$	-1/3	-0.252 ± 0.001	<i>input</i>	-0.340 ± 0.017

The experimental values of G_A/G_V for semileptonic nucleon and Λ decays disagree strongly with simple SU(6), but are fit by constituent SU(6), which fixes G_A/G_V for the constituent quark to fit the nucleon decay data and reduces simple SU(6) predictions by the same factor. The Σ data agree perfectly with simple SU(6) and disagree with constituent SU(6). SU(3) fits Λ data within two standard deviations using the SU(3) "equal spacing rule"¹ with nucleon and Σ as input. All three predictions fit Ξ data within two standard deviations. The contrast between the SU(6) agreement with the Σ and disagreement with N and Λ is a puzzle for model builders as is also any success of SU(3) since no reasonable quark model breaks SU(6) without breaking SU(3).

The large value $(G_A/G_V)_{n \rightarrow p} > 1$ shows that the overlap with a proton of a state produced by a $d \rightarrow u$ transition in a neutron is greater for spin-flip than non-flip, while the EMC result^{2,3} implies that the proton spin is carried mainly by non-quark-spin degrees of freedom which are spectators in beta decay. This inconsistency can be resolved by a toy-model⁴ wave function

$$|B^\dagger\rangle = \cos\theta \cos\phi |b^\dagger\rangle + \cos\theta \sin\phi |[b\epsilon]^\dagger\rangle + \sin\theta |[bD]^\dagger\rangle \quad (1)$$

where $|b\rangle$ is the traditional three-quark SU(6) wave function, ϵ and D denote additional isoscalar constituents respectively with vacuum quantum numbers $J^{PC} = 0^{++}$ and quantum numbers $J^{PC} = 1^{++}$ and which can represent combinations of quark-antiquark pairs, gluons and orbital angular momentum.

From this wavefunction and the experimental result $(G_A/G_V)_{n \rightarrow p} \approx (5/4)$

$$(G_A/G_V)_{n \rightarrow p} = (5/3) \cos^2 \theta - (5/9) \sin^2 \theta \approx (5/4); \quad \sin \theta \approx \pm \sqrt{3/16}. \quad (2)$$

$$\langle \Sigma_V \rangle = \cos^2 \theta - \frac{1}{3} \sin^2 \theta \approx \frac{3}{4} \quad (3)$$

$$\langle \Sigma_S \rangle = \frac{1}{8} \langle D^\dagger | \Sigma_S | D^\dagger \rangle \pm \frac{\sqrt{13}}{8} \sin \phi \langle D^0 | \Sigma_S | \epsilon \rangle \quad (4)$$

where Σ_V and Σ_S denote respectively the contributions of the valence quark spins and the spin of a sea of quark-antiquark pairs respectively to the spin of the proton.

The ϵ component with vacuum quantum numbers which does not contribute to magnetic moments, weak decays nor proton spin was overlooked in previous analyses.⁵ However its excitation to $J = 1$ in EMC can produce a relatively large contribution via the off-diagonal element $\langle D^0 | \Sigma_S | \epsilon \rangle$. The direct effect on EMC of the D component is proportional to $\sin^2 \theta$ and constrained to be small to fit G_A/G_V . The off-diagonal contribution is *linear* in $\sin \theta$ and can be much larger as shown by the factor $\sqrt{13}$ in eq. (4). The three-quark component preserves traditional quark model results for magnetic moments, weak charges, *etc.*⁶ except for the depolarization (3) which reduces all spin-dependent observables by 3/4.

The model-dependent matrix elements in eq. (4) have been evaluated⁴ for a model with a single quark-antiquark pair in a state of spin one coupled nonrelativistically to $J=0$ or 1 with an additional angular momentum of unity, denoted by Λ_S , arising from any combination of orbital angular momentum, gluons or other “inert” degrees of freedom which do not contribute to weak decays nor EMC.

$$\langle \Sigma_S \rangle = -\frac{1}{8} \left(6\sqrt{\frac{26}{27}} \sin \phi - 1 \right) \quad (5a)$$

$$\langle \Lambda_S \rangle = 1 - \langle \Sigma_V \rangle - \langle \Sigma_S \rangle = -\frac{1}{8} \left(-6\sqrt{\frac{26}{27}} \sin \phi - 1 \right) \quad (5b)$$

Substituting into Eq. (4) the values of Σ_S required to fit EMC data gives values of $\sin \phi = 1 \pm 0.3$, nearly independent of the flavor content of the sea.⁶ This gives a baryon wave function $|B\rangle$ with only a small amplitude to be a bare three-quark baryon, $|b\rangle$. The largest term in the wave function, $[[b\epsilon]^1]$, contains the “inert” (0^{++}) quark pair. But the strangeness content of the sea is completely undetermined.

A constituent quark model with the same physics has the usual SU(6) wave function for three and only three quarks having an internal structure directly analogous

the baryon wave function Eq. [1], with v denoting a valence current quark and parameters determined by fitting data.

$$|c^\uparrow\rangle = \cos\theta_c \cos\phi_c |v^\uparrow\rangle + \cos\theta_c \sin\phi_c |v\epsilon]^\uparrow\rangle + \sin\theta_c |[vD]^\uparrow\rangle \quad (6)$$

$$\sin\theta_c \approx \pm\sqrt{3/16}; \quad \sin\phi_c \approx \frac{1}{6}\sqrt{\frac{27}{26}}(1 - 8\langle\sigma_s\rangle) \quad (7)$$

where σ_s denotes the contribution of the sea to the constituent quark spin and gives the same values for $\sin\phi_c$ as obtained in the previous case for $\sin\phi$ for both models.

Many analyses of EMC suggested drastic modifications of the standard quark model^{5,7} which might destroy its successes in fitting other data. These had large values of angular momentum carried by other degrees of freedom like a polarized sea of quark-antiquark pairs, orbital angular momentum and gluons. The toy model (1) shows that data can be fit without these drastic modifications, but the puzzles of Table I remain.

The new puzzles must be balanced against the many successes of a constituent quark model with baryons made of three quarks, mesons of a quark-antiquark pair and no additional degrees of freedom.⁸⁻¹¹ All flavor dependence is in the constituent quark effective masses denoted by m_i which appear with the same values in additive mass terms in the mass operator¹¹ and in a hyperfine interaction inversely proportional to the product of these masses,¹⁰

$$M = \sum_i m_i + \sum_{i>j} \frac{\vec{\sigma}_i \cdot \vec{\sigma}_j}{m_i m_j} \bar{v}_{ij}. \quad (8)$$

where $\bar{v}_{ij}$ is flavor independent and the effective mass includes not only the full single particle energy with the kinetic energy but also the spin and flavor independent part of the two-body interaction.

A very striking prediction for the sum of the proton and neutron magnetic moments^{12,13} in impressive agreement with experiment and with a scale determined entirely by masses and no free parameters follows from simply assuming isospin-symmetric nucleon wave functions consisting only of u and d quarks having Dirac moments with effective masses related to hadron masses by eq. (8) and neglecting orbital angular momentum contributions.

$$\begin{aligned} \mu_p + \mu_n = 0.88 \text{ n.m.} &= \mu_u \langle S_{zu} \rangle_p + \mu_d \langle S_{zd} \rangle_p + \mu_u \langle S_{zu} \rangle_n + \mu_d \langle S_{zd} \rangle_n = \\ &= (\mu_u + \mu_d) \langle S_{zu} + S_{zd} \rangle_p = \frac{M_p}{3m_u} = \frac{2M_p}{M_N + N_\Delta} = 0.865 \text{ n.m.} \end{aligned} \quad (9)$$

Two more magnetic moment predictions with no free parameters, and hadron mass predictions for the "elementary excitations of the nucleon"; i.e. the excitations produced by changing the flavor and/or the spin orientation of a single quark are

$$-1.46 = \frac{\mu_p}{\mu_n} = -\frac{3}{2} \quad (10a)$$

$$\mu_\Lambda = -0.61 \text{ n.m.} = -\frac{\mu_p \mu_s^{EM}}{3 \mu_d^{EM}} = -\frac{\mu_p \mu_s^{col}}{3 \mu_d^{col}} = -\frac{\mu_p}{3} \frac{M_{\Sigma^*} - M_\Sigma}{M_\Delta - M_N} = -0.61 \text{ n.m.} \quad (10b)$$

$$\langle m_s - m_u \rangle_{Bar} = M_\Lambda - M_N = 177 \text{ MeV} = \frac{M_N + M_\Delta}{6} \cdot \left(\frac{M_\Delta - M_N}{M_{\Sigma^*} - M_\Sigma} - 1 \right) = 190 \text{ MeV.} \quad (11a)$$

$$\begin{aligned} \langle m_s - m_u \rangle_{mes} &= \frac{3(M_{K^*} - M_\rho) + M_K - M_\pi}{4} = 180 \text{ MeV} \\ &= \frac{3M_\rho + M_\pi}{8} \cdot \left(\frac{M_\rho - M_\pi}{M_{K^*} - M_K} - 1 \right) = 178 \text{ MeV.} \end{aligned} \quad (11b)$$

where μ_f^{col} denotes the color magnetic moment of a quark of flavor f .¹⁰ Assuming the same additive mass parameter m_i for both mesons and baryons gives

$$\langle m_u \rangle_{Bar} = \frac{M_N + M_\Delta}{6} = 362 \text{ MeV} = \langle m_u \rangle_{Mes} = \frac{3}{8} \cdot M_\rho + \frac{1}{8} \cdot M_\pi = 306 \text{ MeV.} \quad (11c)$$

The mass formula (8) also gives

$$M_{\Omega^-} - M_\Delta = 440 \text{ MeV} = (M_\Lambda - M_N) \cdot \left(3 - \frac{(M_\Delta - M_N)(m_s + m_u)}{2m_s^2} \right) = 451 \text{ MeV.} \quad (12)$$

The basic physics tested by these mass relations is the change in the hadron mass when the spin of a quark is flipped or when the flavor of a nonstrange quark in a nonstrange hadron is changed, as well as the relation between meson and baryon masses. The success of these relations is based upon: (1) The validity of the mass operator (8) with all the flavor dependence in the quark masses m_i , and the *same* value for m_i in *both* terms. (2) The validity of first-order perturbation theory using the same single albeit unknown unperturbed wave function for all states considered.

References

1. Z. Dziembowski and J. Franklin, J. Phys. G17 (1991) 213
2. J. Ashman *et al.*, Nucl. Phys. B328 (1989) 1.
3. Stanley J. Brodsky, John Ellis and Marek Karliner, Phys. Lett. 206B, 309 (1988).
4. R. L. Jaffe and H. J. Lipkin, Phys. Lett. to be published.
5. H. J. Lipkin, Phys. Lett. B237 (1990) 130.
6. H. J. Lipkin, Phys. Lett. B256 (1991) 284.
7. See, for example, G. Altarelli and G. G. Ross, Phys. Lett. B212 (1988) 391.
8. Ya. B. Zeldovich and A.D. Sakharov, Yad. Fiz 4 (1966) 395; Sov. J. Nucl. Phys. 4 (1967) 283.
9. P. Federman, H.R. Rubinstein and I. Talmi, Phys. Lett. 22 (1966) 203; H.R. Rubinstein, Phys. Lett. 22 (1966) 210.
10. A. De Rujula, H. Georgi and S.L. Glashow, Phys. Rev. D12 (1975) 147.
11. I. Cohen and H. J. Lipkin, Phys. Lett. 93B, (1980) 56.

12. Harry J. Lipkin, In: Proceedings of the 7th International Symposium on High Energy Spin Physics, Protvino (USSR), 22-27 September, (1986) Edited and Published by The Institute of High Energy Physics, Serpukhov (USSR), (1987) Vol. I, p.98.
13. Harry J. Lipkin, Nucl. Phys. A478, (1988) 307c.

END

**DATE
FILMED
01/30/92**

I