VOLUME II**LITERATURE REVIEW ON THE USE OF BIOACCUMULATION FOR
HEAVY METAL REMOVAL AND RECOVERY**

by

John R. Benemann**Consultant
1212 Kelly Court
Pinole, CA 94564**

and

Edward W. Wilde**Westinghouse Savannah River Company
Savannah River Site
Aiken, SC 29808****DISCLAIMER**

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

M8909021

MASTER**DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED** *So*

ABSTRACT

Bioaccumulation of metals by microbes - "bioremoval" - is a powerful new technology for the concentration, recovery, and removal of toxic heavy metals and radionuclides from waste streams and contaminated environments. Algae are particularly well suited for metal bioremoval. A recent commercial application of bioremoval utilizes inert (dead) immobilized microalgae biomass as ion exchange materials for the removal of heavy metals from industrial waste waters. Also, living microalgal cultures have been used to remove metals from mine effluents. Other types of microbial biomass can also be used in similar processes. Commercialization of this technology is under active development.

Microbial cells and biomass can bioaccumulate metals and radionuclides by a large variety of mechanisms, both dependent and independent of cell metabolism. Microbial cell walls can act as ion exchange and metal complexation agents. Heavy metals can precipitate and even crystallize on cell surfaces. Metabolically produced hydrogen sulfide or other metabolic products can bio-precipitate heavy metals. Many microbes produce both intra- and extracellular metal complexing agents which could be considered in practical metal removal processes. Microorganisms have also evolved highly efficient and often selective transport systems that allow them to accumulate essential trace metals, often from extremely low (<ppb) concentrations. These transport systems can be harnessed for the selective removal of trace amounts of toxic metals and radionuclides. Several distinct mechanisms can operate in sequence or simultaneously. Bioremoval processes are greatly affected by the microbial species and even strain used, pH, redox potential, temperature, and other conditions under which the microbes are grown. Development of practical applications of bioremoval requires applied research using the particular waste solutions to be treated, or close simulations thereof.

From a practical perspective, the selection of the microbial biomass and the process for contacting the microbial biomass with the metal containing solutions are the key issues. Much of the recent commercial R&D has emphasized commercially available, inert, microbial biomass sources (e.g., microalgae powders, bakers yeast, etc.), as these can be acquired in sufficient quantities at affordable costs. However, for improved effectiveness, the screening for, and eventual production of, microbial strains under growth conditions which maximize bioaccumulation will be required. Also, the use of living, metabolically active, microorganisms in bioremoval processes has significant potential. Contacting of the microbial biomass with the heavy metal containing solutions can be accomplished by immobilizing the cells or using microbial biomass sources easily concentrated by sedimentation or filtration.

The fundamental research and practical applications of bioaccumulation by microalgae suggests these organisms warrant a high priority in the development of advanced bioremoval processes. The further development of these processes could have significant applications at the SRS and other DOE facilities, as well as environmental protection generally.

TABLE OF CONTENTS

Abstract	i
Table of Contents	ii
1.0 INTRODUCTION	1
2.0 HEAVY METALS AND MICROORGANISMS	6
2.1 Trace Metal Uptake	6
2.2 Metal Toxicity	8
2.3 Heavy Metal Resistance and Bioaccumulation	9
3.0 MICROBIAL METAL BIOSORPTION	13
3.1 Introduction	13
3.2 Microbial Cell Walls	14
3.3 Mechanisms of Biosorption	16
3.4 Metal Desorption and Biomass Regeneration	18
4.0. OTHER MICROBIAL BIOACCUMULATION PROCESSES	20
4.1 Biodeposition Reactions	20
4.2 Anaerobic Biodeposition	21
4.3 Intracellular Metal Bioaccumulation	22
5.0. MICROBIAL METAL BINDING AGENTS	25
5.1 Metallothioneins, Other Metal Binding Proteins, and Phytochelatins	25
5.2 Siderophores and Other Complexing Agents	27
5.3 Extracellular Polysaccharides and Other Exudates	28

TABLE OF CONTENTS (Contd)

6.0 SOURCES OF MICROBIAL BIOMASS FOR BIOACCUMULATION	30
6.1 Microbial Biomass Sources	30
6.2 Screening for Bioaccumulating Microbes	31
6.3 Genetic Selection in Continuous Cultures	32
6.4 Screening for Bioaccumulation Mutations	34
7.0 PROCESS DESIGN FOR BIOREMOVAL SYSTEMS	36
7.1 Introduction	36
7.2 Immobilization of Biomass for Bioremoval	36
7.3 Use of Biofilms in Bioremoval	38
7.4 Conclusions	39
8.0 APPLICATIONS OF MICROALGAE IN BIOREMOVAL	40
8.1 Introduction	40
8.2 Microalgae Bioremoval from Mine Effluents	40
8.3 Microalgae Bioremoval from Treatment Ponds	41
8.4 Microalgae Bioremoval from Industrial Wastewaters	43
9.0 CONCLUSIONS	45
10.0. REFERENCES	46

1.0 INTRODUCTION

The practical use of bioaccumulation of metals by microbes is a powerful new technology termed "bioremoval" for the concentration, recovery and removal of toxic heavy metals and radionuclides from waste streams and contaminated environments. Some practical applications of bioaccumulation have already been commercialized in the treatment of industrial wastewater and mining effluents. Based on available data, current processes can be greatly improved and new bioremoval technologies could be developed in the near term which could solve many of the environmental problems associated with the release of heavy metals and radionuclides.

There are several mechanisms of bioaccumulation. A major distinction between mechanisms is that some require active metabolism by living microbes (active processes), while others can occur with dead biomass (passive processes), or even specific components thereof (e.g. cell walls, polysaccharides, chelating agents, etc.). Within these broad categories, several distinct bioaccumulation mechanisms can be distinguished:

- the action of cell walls as ion exchange materials;
- the precipitation and crystallization of heavy metals on cell surfaces;
- the anaerobic bioprecipitation of metals by metabolically generated H_2S or other precipitating agents;
- the active uptake and intracellular storage of metals;
- the metabolism dependent oxidation or reduction of metals and their precipitation (intra - or extracellularly); and
- the complexing of metals by specific binding agents produced by the cells.

Often two or more of these mechanisms can operate simultaneously with the same microbe. Thus, it is generally observed that metal accumulation by living microbes occurs in two phases - one fast (complete in less than 10 minutes) and one slow, often requiring cell metabolism and even growth. In the green alga *Chlorella vulgaris*, these two phases were exemplified by rapid extracellular adsorption followed by relatively slow intracellular accumulation (Khummonghol et al., 1982). More recent studies with the green alga *Chlamydomonas reinhardtii* demonstrated a similar phenomena (Xue et al., 1988). However, this is not always the case. For example, U bioaccumulation by the bacterium *Pseudomonas aeruginosa* occurred intracellularly very rapidly (<10sec) while under similar environmental conditions with the yeast *Saccharomyces cerevisiae* it took place extracellularly with a slower reaction (3-5 min) (Strandberg et al., 1981). Metabolism was not required in either case.

The actual mechanism(s) involved in bioaccumulation processes are unknown in many cases and several mechanisms may be involved simultaneously or sequentially. Therefore, the experimental data is generally interpreted phenomenologically in terms of basic kinetic parameters: equilibrium concentrations, removal ions, etc. In many cases, only a few of these are reported and available data is often insufficient to allow extrapolation from laboratory to larger scale systems.

Nevertheless, existing data and experience clearly demonstrate that bioremoval processes can be harnessed in practical applications for treatment of wastewaters and restoration of contaminated environments.

Depending on the metal, microbial biomass, and experimental variables, it is possible to concentrate metals from the environment into the microbial biomass by factors of many thousands, or in some cases, over one million fold; from concentrations ranging from hundreds of ppm to less than one ppb; over time periods of a few seconds to several hours, and even days (if culture growth is required); and with the final metal content in the biomass up to and even exceeding the organic biomass dry weight.

The concentration factors, metal contents, equilibrium concentrations, and kinetics of metal accumulation by the microbial biomass are dependent on a large number of variables during both the metal accumulation process (particularly pH and concentrations of other ions), and during the growth and preparation of the microbial biomass. Added to this are the fundamental differences in metal binding exhibited by different microbial species, even strains of the same species, and the multiplicity of different mechanism by which metals are accumulated by microbes. The difficulties in dealing with so many variables are exacerbated by the problems of comparing results from different laboratories, which follow different approaches and protocols, collecting different types of data, often expressed in different units, which are usually impossible to compare directly.

Most of the literature data on removal and concentration of heavy metals from solutions deals with laboratory solutions, not actual waste streams where multiple metals and high, often variable, concentrations of other ions can interfere with, or at least significantly affect, the bioaccumulation process. Nevertheless, it is evident from the literature that bioaccumulation is a potentially very powerful technique, suitable for the removal of specific metal ions from process and waste solutions, and even from contaminated environments, particularly in the low ppm to low ppb concentration ranges that are difficult for conventional technologies (i.e., ion exchange, or activated carbon) to handle.

Potential applications of bioremoval processes in environmental clean-up range from the use of immobilized microalgae for the recovery of gold in mining effluents to the use of living algae cultures in "meander" systems for the removal of heavy metals from industrial effluents (see Section 8). A major area of current development that has already reached commercial application, is the use of microbial biomass, specifically microalgae biomass, immobilized on various supports to remove key metals from industrial wastes.

The ion exchange materials from microalgae and other microbes have some properties, such as lower interference by divalent ions and greater specificities, that favor their applications in specific situations where conventional ion exchange materials do not perform well. If the cost and performance of these materials can be improved, they could offer broad economic advantages over conventional ion exchange resins.

Most of the research in bioaccumulation has been carried out using dead biomass, capitalizing on the ion exchange properties of microbial biomass. However, despite comments in the literature that "bioaccumulation of metals by living microbes is not feasible because the metal toxicity often inhibits microbial growth" (Volesky, 1986), and similar sentiments expressed by many other authors, living biomass has significant promise in practical applications (Maeda et al., 1985; Olson and Brinkman, 1987; Bender and Ibeanusi, 1987).

In some cases, only a metabolism dependent mode of bioaccumulation has been observed. For example, Maeda et al. (1985) reported that arsenic bioaccumulation by the green alga Chlorella vulgaris occurred only when living cells were involved. Living cells bioaccumulated up to 50 mg As/g dry weight of the algae but no arsenic was bioaccumulated when cells had been pretreated with a respiratory inhibitor or heat killed. Similarly, Bender and Ibeanusi (1987) reported that passive adsorption on cell surfaces was not a primary mechanism in the long-term uptake of lead by the filamentous blue-green alga Anabaena. Lead uptake was increased by or the addition of carbon dioxide and other nutrients.

Metabolism-driven active uptake systems may be more effective in reducing specific metal ion concentrations below that which could be achieved with simpler adsorption processes, and can result in higher metal concentrations in the biomass. Living biomass can also change the speciation of heavy metals by oxidation-reduction reactions, allowing much greater flexibility in the overall process. "Once the limitations of working within the constraints of a living system have been overcome there is a far greater potential for a living continuous process" than for the use of dead biomass (Macaskie and Dean, 1989).

One of these limitations is that heavy metals inhibit microbial growth. However, growth and metal accumulation need not be coincidental and would likely be de-coupled in most practical process, as growth is more sensitive to the metals than energy metabolism, required for active processes (e.g., Macaskie and Dean, 1984). The issue is not living vs. dead biomass in bioaccumulation application; both processes have distinct and complementary capabilities in specific situations and both should be considered in bioaccumulation R&D.

From a practical perspective, a major issue is the method used to contact the microbial biomass with the metal containing solution. In the laboratory, it is possible to contact metal and microbes in shake flasks. However, laboratory methods for separating the microbial biomass from the metal solution after contacting, such as filtration and centrifugation, are generally not applicable when dealing with large volumes of metal containing solution. More cost-effective methods must be used.

The preferred option is the immobilization of the microbial biomass onto some type of inert support matrix that allows passing large volumes of solution through a relatively small bed of active material. Alternatively, the biomass can be formed into pellets that can be used in mixed tank or fluidized bed reactors and easily separated from the process liquor by sedimentation. Next to the development of the metal biosorbing microbial biomass itself, the method for biomass immobilization or separation is the key to the success of this technology. Many fungi and microalgae grow in large colonies, flocs, or filaments that can be easily separated by gravity sedimentation or filtration. Therefore, these organisms could have significant advantages in bioremoval processes.

The objective of any R&D project in bioaccumulation is to develop a microbial biomass that, for a particular application, exhibits high specificity, high capacity, lack of interference by other components, low residual metal ion concentrations, ready recovery of the metal ions in a concentrated form and reusability of the biomass. Meeting these criteria within allowable economic boundaries for a specific application requires applied R&D, as well as a fundamental understanding of the mechanisms involved. It also suggests that rather than a few universally applicable bioaccumulation systems, specialized bioremoval processes are likely to be developed for different types of applications.

The commercial potential of bioaccumulation is reflected in part by the ever increasing number of researchers in this field. The market for the biological treatment of metal-containing wastes is potentially enormous. The U.S. market for treatment of metal-containing waste streams projected for the year 1998 is estimated at over \$200 million dollars for current discharges and \$1.5 billion for groundwater and leachate treatment systems. About one third of this, or over \$0.5 billion, is projected by one market forecast to be captured by biological treatment processes (Biotechnology Processing, 1989). This estimate does not include the potential applications of bioaccumulation in the cleanup of existing sites, such as the DOE facilities.

From a commercial perspective this field is still in its infancy, as reflected by the fact that only one company, Bio-Recovery Systems, Inc. of Las Cruces, New Mexico, is currently offering commercial products based on microalgae biomass. Although several other companies are becoming active in this field (Anonymous, 1989), none appears to actually have significant sales. Obviously, this is an area ready for additional commercial activity.

There are a number of technologies competing with bioremoval: Plant derived polymers, such as modified cellulose or starch derivatives (Chaufer et al., 1986), lignins, tannins (Sakaguchi and Nakajima, 1986) and even peat (Bencheikh-Lehovine, 1989), to mention only some examples, can bind metals. Other biological sources of metal binding agents are chitin and chitosan (Muzarelli et al., 1985, 1986; McKey et al., 1986, de la Noue and Proulx, 1988) obtained from the shrimp exoskeleton, and seaweed products such as alginate (Wilkinson, et. al., 1989, Kuyucak and Volesky, 1988, 1989a, b, c).

Furthermore, the ion exchange technologies and other conventional technologies already in use in heavy metals recovery are being improved (Streat and Naden, 1987), and may provide suitable, and better understood, alternatives in many specific situations. Also, even though a large potential market exists, market penetration is difficult due to the diverse nature of the applications and widely dispersed customer base, resulting in high service and technical support costs. Despite these potential competitors, bioremoval is likely to become a major technology in the future - based on the present experience with already commercially available microalgae based products and the research data on microbial bioaccumulation.

Microbial cells have the ability to bioaccumulate metal ions "...in a manner which should be the envy of the industrial chemist pursuing metal recovery . . ." (Hunt, 1986). The rewards that will be derived from practical applications of this technology are potentially enormous: microbes could help us to recover and remove toxic metals and radionuclides from waste waters and even from the environment, preventing the potentially disastrous environmental and public health consequences resulting from such releases.

Bioaccumulation is covered in many of the reviews and books dealing with biotechnology in the mining industry (e.g., Norris and Kelly, 1987; Ehrlich and Holmes, 1986; Lawrence et al., 1986; Eccles and Hunt, 1986; Karavaiko and Groudev, 1985). Several more books are in press or preparation, but were not available (except for selected chapters) for review during preparation of this report (Beveridge, 1989; Volensky, 1989). In addition, several general reviews specifically dealing with bioaccumulation have recently appeared (Hancock, 1986; Ross, 1987a, 1987b; Beveridge, 1986; Volesky, 1987; Gadd et al., 1988; Gadd, 1989; Olson et al., 1989; Macaskie and Dean, 1989).

This report is designed primarily as a guide to the literature and a discussion of the current state-of-the-art and required R&D, specifically as it may apply to problems identified at the SRS and other DOE facilities. First, microbial interactions with heavy metals are reviewed, starting with the uptake of metals and ions required for microbial growth. Then, the use of both dead biomass in ion-exchange type of bioaccumulation processes is reviewed, followed by other mechanisms of biosorption, involving both living and dead biomass. Some specific examples of processes using microalgae are discussed in the concluding sections.

2.0 HEAVY METALS AND MICROORGANISMS:

2.1 Trace Metal Uptake

Microorganisms need a variety of heavy metals for growth: Fe, Mo, Ni, Co, V, Se, Mn, Mg, Zn, Cu, Cr, etc., for a total of over twenty different metals and ions. Not all microbes use all these elements; however, a few are universally required and many are needed by a majority of microorganisms. Although most metals are required in only small amounts (usually less than 0.01% of cell dry weight), without them, the organisms can not survive and reproduce. These trace elements, or micronutrients, bind to various enzymes where they perform specific catalytic functions essential to cell survival, at least under certain environmental conditions.

The need for such essential trace elements has resulted in the evolution of highly efficient, and often highly specific, mechanisms for their uptake, particularly under conditions where these trace elements become limiting factors in growth and reproduction. Under such circumstances of growth limitation, many trace elements can be sequestered from concentrations at or below the parts per billion range, and stored intracellularly, resulting in relatively high concentration factors. (Concentration factors are the ratio of extracellular to intracellular concentrations, the latter generally given in units of dry weight of the biomass).

Microbial cells normally store trace elements in amounts that greatly exceed their immediate metabolic or survival needs. Upon transfer of a microbial culture to a new growth solution devoid of one of the trace elements it often grows at an undiminished rate for several generations, increasing its biomass several-fold, before intracellular storage is exhausted and growth slows down. When intracellular storage reserves are exhausted, many microbes can adapt their metabolic machinery and reduce their requirements for a particular metal to a minimum, by efficiently recycling the few atoms present in each cell, substituting other elements (even though they may work at greatly reduced efficiency), or, in some extreme cases, even apparently doing without.

One example of doing without is described in recent reports showing that some nitrogen-fixing soil bacteria can, in the absence of molybdenum, induce a nitrogenase enzyme that is totally devoid of this metal previously thought to be essential. (Robson, 1986). Earlier work had already established that under Mo limiting conditions these bacteria will take up and replace Mo with V or W, in the presence of those metals (Benemann et al., 1972, 1973). However, the metal substituted enzymes were not catalytically active. In conclusion, even though growth is greatly reduced under conditions where a trace element is limiting, it may not always be completely inhibited.

One major microbial adaptation to conditions of low nutrient availability (and this is a general phenomenon for both macro - and micronutrients) is the induction of a highly efficient uptake system for that nutrient, capable of concentrating the metals from much lower concentrations than is the case with cultures grown with an excess of that nutrient.

Thus, it is commonly observed that there are two (sometimes even more) distinct assimilation mechanisms for a particular nutrient: a "low affinity" system normally present constitutively under conditions of ample nutrient supply and a "high affinity" uptake system induced under conditions of nutrient limitation (Silver, 1985). However, regulation of metal uptake systems can be complex. For example, in the yeast *Candida utilis*, high Mn^{2+} represses the Mn^{2+} transport systems, but the same was not the case for Cu^{2+} transport (Perkin and Ross, 1986).

The high affinity systems are characterized by both a high specificity and a high affinity for the element. In some cases, less than 100 parts per trillion (0.1 ppb) can result in measurable stimulation of growth. For example, the Ni uptake system of the filamentous blue-green alga *Anabaena cylindrica*, does not transport the very similar Mg^{2+} and Co^{2+} and Ni uptake can be measured at levels as low as 0.05 ppb. The ability of microbes to utilize such low levels of trace elements suggests that, in at least some cases, the growth of microbes observed in the absence of specific nutrients could have been due to the limitations of the analytical techniques used and the difficulty in removing the last trace contamination from the growth media.

The difficulties of working with such trace levels of metals makes this a field of research where new discoveries of essential trace elements for microbial metabolism are still being made, and where technical difficulties can lead to erroneous conclusions. For example, returning to the example of nitrogen fixation, early reports of catalytic activities by V-substituted nitrogenase enzymes (McKenna et al., 1970) overlooked the small contamination by molybdenum, a problem that cannot yet be eliminated from the more recent reports on molybdenum-free nitrogenases, mentioned above.

The high affinity uptake systems often exhibit a relatively high rate of metal influx when suddenly exposed to saturating levels of the trace element. Metal uptake by microbes is under feed-back control; once intracellular storage requirements are filled, further uptake is much diminished. Normally, high affinity systems only result in the accumulation of the small amounts of trace elements required for growth. During conditions of trace element limitation, when there is a maximal induction of transporters, an "overshoot" of metal uptake can be observed. Thus the yeast *Candida utilis* grown under low Zn^{2+} conditions will hyper-accumulate Zn when exposed to this metal, resulting in a biomass containing almost 1% Zn by dry weight, over 10-fold the amount contained in normal cells (Failla and Weinberg, 1977; Failla 1977).

The control of such "overshoot" phenomena, coupled to the high uptake rates from very low concentrations observed, suggest that it could result in greatly improved bioaccumulation systems, compared to those observed with normally grown cells. One major challenge for future research is to overcome the feedback inhibition to metal uptake, and the limitations on intracellular storage, to maximize the longer term accumulation rates and amounts sequestered in the biomass. Increasing metal storage can be achieved by both physiological and genetic methods. This is an area where basic research could readily point the way to practical applications.

Indeed, the study of trace element storage in microbial systems is an area which has been relatively neglected and which deserves much further study. Heavy metal storage is important in applications of such processes in practical bioaccumulation systems. Thus, although algae are known to accumulate metals up to 20% of their dry weight (Ahlf et al., 1980), most studies on heavy metal uptake by living microbes report intracellular metal contents of less than one percent, often less than 0.1%, while several percent, preferably 10% or more, would be desirable in many cases to make practical systems a commercial reality. Of course, the required metal accumulation by the biomass depends on the parameters of the specific process being investigated: the concentration in the feed solution, the re-utilization of the biomass, the cost of the biomass, and the economics of competitive effluent treatment processes. Thus, even a relatively low metal content would be adequate in some cases.

Another characteristic of the high affinity uptake systems is that their specificity for different metal ions is quite distinct from that exhibited by the "low affinity" uptake systems. A few examples will be discussed later (Section 4.3); however, this field is too undeveloped to allow a more detailed discussion.

This is an area which requires fundamental research. A major distinction between the high and low uptake systems, is the higher energy requirements of the former, as would be expected from the need to transport a nutrient against a high concentration gradient. This explains the generally inducible nature of the high affinity uptake system; when ample supplies of the trace element are present the energy consuming high affinity system would be an unnecessary drain on cell metabolism. One consequence of this inducibility is that the culture conditions of the microbial biomass would have a significant effect on the uptake capabilities of the cells. This could explain some of the variability found between different literature reports, although, in general, it is not apparent from the literature that there has been much attention paid to the potential of using trace element limited cultures in bioaccumulation studies. This approach is experimentally straight forward and could lead to significant advances in the short-term.

The potential to manipulate these high affinity uptake systems, such that they exhibit desirable specificities for particular metal ions, or combinations thereof, is alluring. Genetic engineering methods might be used in the future to manipulate the "transporter" proteins to affect their specificities. However, before this is possible, a better understanding of the underlying biochemistry and genetics is required. The one broad generalization that can be made is that microbes are complex, highly adaptable molecular machines that have stored in their two-thousand-odd gene library a significant repertoire of metabolic capabilities and alternatives on how to obtain from the environment the elements they require for growth. In the context of bioaccumulation, the challenge is to use and tailor these metabolic systems to result in the accumulation.

2.2 Metal Toxicity

Many of the essential trace elements are toxic given sufficient concentrations - which in some cases are not much higher than those required for optimal growth. A good example is Se, required in trace amounts by some microbes (and most higher organisms, including humans) but a highly toxic material, to higher organisms, at only about ten-fold higher concentration than required for survival. In other cases, for example Cu and Zn, toxic concentrations are considerably higher than those required for growth, but the elements often are found in the environment at concentrations higher than desirable for microbial, or human, health.

Thus, the microbial concentration mechanisms could be used to not only recover essential trace elements of economic importance (e.g., Co, V, Mo, etc.) from low concentrations in process or waste streams but also to remove these and other trace metals (e.g., Se, Cu, Zn) from environments in which they are present at potentially toxic levels. However, application of the microbial trace metal uptake systems to such situations requires the selection of strains that are relatively resistant to such elements without diminishing their uptake capabilities.

In some cases, the microbial concentrating mechanisms will mistake an otherwise unessential, or even toxic, metal for a required one - resulting in its uptake and cell death. Indeed, this is the major cause for toxicity by heavy metals such as Hg, Cd, Pb, etc., which have no biological function but sometimes substitute for essential trace metals during the uptake process and, in some

cases, are even incorporated into specific enzymes, resulting in inhibition of particular metabolic functions. The reason for this confusion on the part of the microbes is that these elements are only rarely found in appreciable concentrations in the natural environment, and therefore the uptake systems have failed to evolve a clear specificity against them. Their chemical similarities (ionic radius, shape, ligand affinities, etc.) to the required elements for which they are mistaken account for their ready acceptance by the microbes.

Toxic metals, once introduced into the cell can attach to and alter the function of any number of enzymes, particularly to free -SH groups (which are usually critical to enzyme action). Indeed, the relative toxicities of Hg, Zn, Pb, Cd, and Mn correlate with the solubility products of the metal sulfides, suggesting that -SH (thiol group) binding is the common mechanism of toxicity (Fisher and Jones, 1981). The most sensitive of enzymes are those that have a dithiol configuration at their active site (Kagi and Hapke, 1984). Thus, it is not surprising that intracellular protection against heavy metals poisoning involves molecules with thiol groups, such as glutathione and metallothioneins (see below).

Heavy metal toxicity can be both specific, affecting a single enzyme and therefore metabolic pathway, or general, affecting many different enzymes. The exact mechanisms in metal inhibition are sometimes difficult to ascertain, as the observable effects (cessation or decreased growth, altered cell morphology, and eventually, cell death by lysis and dissolution) are nonspecific.

In conclusion, practical applications of the natural microbial uptake mechanisms need not be limited to essential trace elements. They could also be used in the removal from solution of other toxic elements. However, these active uptake systems are coupled to the energy generating metabolic machinery of the cell, and thus require living organisms. The conditions optimal for, or even permissible of, microbial growth are usually quite different from those required in a heavy metal concentration process. Although heavy metals can inhibit both replication and metabolism, these two processes are usually not affected to the same degree. Metabolism by living cells is typically much less sensitive than growth and reproduction. Thus, by decoupling growth from heavy metal concentration, it should be possible to make use of the active transport processes for toxic heavy metal removal. This is an area that warrants additional investigation.

ire O

2.3. Heavy Metal Resistance and Bioaccumulation

Microbes can become resistant to various heavy metals by increasing exposure to the metal over a period of many generations, resulting in selection of resistant cultures adapted to living with such metals. Microbes have developed a number of different strategies to become metal tolerant. These strategies include (Misra et al., 1984; Silver, 1985):

- making the cell impermeable to the heavy metal;
- reduction/oxidation of heavy metals into nontoxic forms;
- excretion of extracellular metal binding agents;
- synthesis of intracellular binding agents;
- intracellular storage of metals;

- metabolic conversion to volatile forms;
- production of cell surface metal adsorbing polymers; and
- increasing the efflux rates of heavy metal ions.

The field of microbial heavy metal inhibition and resistance is vast, and would require a review in its own right.

Among the best studied systems is the reduction of Hg^{2+} to elemental mercury (Summers, 1985), which often volatilizes and is removed from the cell. In most bacteria, the gene for mercury volatilization is carried on plasmids. Plasmids are small pieces of DNA which can replicate independently of the chromosome and are often exchanged between different bacteria. Plasmids often carry useful genetic information such as antibiotics and heavy metals resistance, which allows the cells to survive in novel environments. The genes coding for resistance to metals other than mercury are also often, though by no means universally, found on plasmids. Plasmid mediated heavy metal protection mechanisms include examples of most of the above listed mechanisms: reduction of metals and intracellular storage, exclusion of the heavy metal from the cell, energy-dependent efflux systems, and specific heavy metal binding and storage proteins and other molecules.

Often heavy metal resistance mechanisms are general, that is, multiple heavy metal resistance is observed. A blue-green algae isolate (*Synechococcus* sp.) obtained from an industrial effluent was simultaneously resistant to Hg, Cd, and Pb (Bisen et al., 1987). Acquisition of resistance to one metal can confer resistance to several others. For example, the fungi *Rhizopus stolonifera* and *Cunninghamella blakesleana* adapted to tolerate elevated concentrations of Cu also became tolerant to elevated levels of Cd, Co, Ni, and Pb (Garcia-Toledo et al., 1985). This would be expected from the nonspecific nature of some of the resistance mechanisms. Also, multiple heavy metal resistance (along with antibiotic resistance) can be conferred by plasmids found in bacteria isolated from a natural environments (Schuett, 1989).

Heavy metal resistance need not always reflect genetic mutations, because inducible resistance mechanisms are also observed. However, this type of protection generally appears only after a long period of exposure to the heavy metal and is lost when the metal is removed. For example, in the case of the fungus *Neurospora crassa*, Cu toxicity resulted in a blue-colored biomass (due to the adsorption of Cu) which contained 26% more amino groups and a higher content of acidic amino acids (Subramanyam et al., 1983).

Such inducible mechanisms can be of significant value to an organism living in an environment where heavy metal contamination is sporadic. This is because, in general, the heavy metal protection mechanisms require significant expenditures of metabolic energy, which results in a reduction in growth potential. Under conditions of low heavy metal concentrations, repression of the heavy metal resistance mechanisms would thus help maximize growth. Indeed, one of the best known metal protection mechanisms, also found in higher organisms including humans, the metallothionein system, is such an inducible protection mechanism (see Section 5.1).

Heavy metal resistant strains may or may not be good candidates for heavy metal bioaccumulation. As heavy metal resistance often involves mechanisms which either exclude the metal to start with (due to a greatly reduced affinity for the metal), or eject it (e.g., by pumping out the metal that penetrates into the cell), such situations would result in a decreased uptake, and thus little net removal of the heavy metals (see, for example, Baldry and Dean, 1980). In the screening for Cd uptake, bacterial strains derived from a Cd contaminated area did not adsorb Cd well, although one strain of Citrobacter derived from a Pb contaminated area exhibited good Cd uptake (and resistance) (Macaskie and Dean, 1982). In another example, Cu tolerant strains of the green alga Scenedesmus did not transport Cu into the cell, compared to a Cu sensitive strain. However, initial cell surface adsorption was similar (Mierle and Stokes, 1976).

In some cases, it appears to make little difference whether strains are resistant or sensitive to a heavy metal: uptake of Zn by Zn resistant strains of the blue-green alga Anacystis was similar to that of Zn sensitive strains (Shehata and Whitton, 1982). Also, two strains of the bacterium Bacillus subtilis differing in sensitivity to Cd^{2+} exhibited similar degrees of Cd^{2+} bioaccumulation (Surowitz et al., 1984). In other cases, such as those involving heavy metal adsorption to the cell surface or intracellular storage in an inert form, resistant strains would be clearly superior to the wild type strain in metal removal. In one of the first studies of metal binding by yeast cells, a Co resistant strain of baker's yeast Saccharomyces cerevisiae accumulated much more Co, up to 10% by dry weight, than a sensitive strain (Parker and O'Brien, 1954). The mechanism of this process, e.g., active bioaccumulation or surface biosorption, was not clear.

In a more recent example, a more or less specific, metabolism linked, metal uptake into the cell was demonstrated in resistant microbes: in plasmid encoded Hg resistance, a metal reductase results in the detoxification of the mercury ion, which is brought into the cell to the reductase by an active uptake system. Evidence for this was the fact that some reversion mutants (i.e., mutants that lost mercury resistance) were super-sensitive to this metal: they still had the uptake system but lost the reductase enzyme (Foster et al., 1979). In the same vein, one Cd resistant strain of the bacterium Pseudomonas putida accumulated Cd which is stored in both polyphosphate and bound to low molecular weight metallothionein-like proteins (Higham et al., 1985). Similarly, in a Cd resistant strain of Saccharomyces cerevisiae, almost all of the Cd was bound to cytosol protein, indicating uptake into the cell (Joho et al., 1985). Another example is that of Ni resistant strains of a blue-green alga, Synechococcus, which effectively removed Ni from solution via an intracellular uptake and storage mechanism, resulting in dense intracellular granules (Wood and Wang, 1983).

In many cases, increased metal bioaccumulation by metal resistant microbes is due to increased bioaccumulation to cell wall and surface polysaccharides, as first demonstrated for two Zoogloea ramigera strains (Friedman and Dugan, 1967). Similarly, Scott et al. (1986) found that a strain of Arthrobacter viscosus (as the name implies, it has a noticeable capsular coat) removed both Cd and Pb at greater efficiency than A. globiformis, a noncapsular species (although for the case of Pb at the lowest concentration tested, 1ppm, the results were similar). Further work by this group (Scott and Palmer, 1988), demonstrated that it was the exopolysaccharide of A. viscosus which accounted for most of the metal adsorption.

The mycelium of the fungus Penicillium ochro-chloron, isolated from a copper plating solution was shown to accumulate higher amounts of copper than common in other fungi (Stokes and Lindsay, 1979). In another study, the mycelium of fungi adapted to tolerate Cu removed about twice as much copper from solutions as those of nontrained cultures of the same species, although no extracellular polysaccharides were apparent. In these examples, the higher metal uptake was most likely due primarily to adsorption to the outer cell wall and capsular materials, a subject which is discussed in detail next.

3.0. MICROBIAL METAL BIOSORPTION

3.1 Introduction

Ion exchange resins find extensive use in waste treatment and the mining industry for heavy metal removal (Streat and Naden, 1987; Streat, 1987). The function of microbial cell walls, their constituents and other biopolymers as ion exchange materials and similar reversible metal binding agents ("biosorption") has been recognized for several decades and is the subject of extensive current research and even commercial ventures. The use of microbial biomass to replace or substitute for conventional ion exchange materials in environmental applications does not involve major process modifications or unfamiliar technology, it is therefore a logical and near-term application of bioremoval technology.

Because the biosorption properties and materials of the microbes do not involve growth or metabolism, inert (dead) biomass can be used in microbial ion exchange, and is often found to be superior to living cells. The biosorption properties of the microbial biomass are primarily associated with the cell wall and cell surface polymers. Therefore, fractionation of the microbial biomass can be used to enrich for these active agents, in some cases significantly increasing heavy metal binding capabilities.

It is often difficult to ascertain from the literature whether metabolism-independent metal adsorption is taking place exclusively. In some cases, particularly when the biomass is clearly nonliving (e.g., heat killed), this is not an issue. Even with living cells, in many cases, bioaccumulation of metals is clearly due to passive adsorption to the cell wall (e.g., biosorption). Thus, Cd uptake by two strains of Bacillus subtilis differing in sensitivity to cadmium, was almost entirely (about 90%) due to adsorption on the cell wall, with only a small amount found in the membrane and soluble cell fractions (possibly contaminants) (Surowitz et al., 1984). The blue-green alga Chroococcus parvis, accumulated Cd, Cu, and Zn in a rapid (<1 min) process that was reversed by metal chelators, clearly indicating surface adsorption (biosorption) (Les and Walker, 1984).

In some cases, it is reported that "non-living" biomass was used, either because it was dried or because no energy source was present. However, neither condition eliminates metabolic activity based on endogenous substrates (such as polybetahydroxy-butyrate in bacteria). However, in most cases, this does not appear to have been a problem, particularly where the reversible nature of the bioaccumulation process has been demonstrated. It is the reversibility of the process which, for the purposes of this discussion, distinguishes biosorption from other bioaccumulation processes.

Initial metal cation binding by biopolymers is to a large extent due to ionizable groups: phosphates, carboxyl, and sulfate being the main ones. Both polysaccharides and proteinaceous materials participate in surface binding of metal ions, with the proteins often providing specific coordination sites for particular metals. Thus, binding of metals is not limited to ion exchange mechanisms, as chelation mechanisms also occurs on the cell surface. The formation of coordination compounds and nucleation and outgrowth of metal aggregates may also take place. Indeed, several mechanisms are usually involved simultaneously, and the kinetics of metal accumulation are likely to be complex (Hunt, 1986). The microbial cell wall, and the surface biopolymers, are the major focus of interest in this context and are discussed next.

3.2 Microbial Cell Walls

The microbial cell wall is the outer most portion of the cell that separates the environment from the protoplasm. By this definition, the cell wall includes the superficial layers exterior to the cell wall proper (e.g., the slime layers, sheaths, capsules, S-layers, etc.); the surface structures such as pili and flagella; and even the underlying cell membrane and the periplasmic space between the cell membrane and the cell wall. Operationally, this is an acceptable definition as these structures are usually difficult to separate into clean and distinct fractions. It should be noted that not all microbes have a conventional cell wall. Some, such as the green alga *Dunaliella*, are equipped to survive with only a thin cell membrane, while in others, the cell membrane is not distinct from the cell wall proper.

Microbial cell walls are not inert or passive structures. They actively participate in cell metabolism in many ways, from the control and regulation of cell permeability to cell division. They are not uniform structures, but are composed of many different biopolymers - carbohydrates, proteins, lipids, and combinations thereof - each with particular properties and functions, and metal binding capabilities. Thus, the cell surface of the green alga *Chlorella* has been characterized as "literally a mosaic of metal ion binding sites which differ in affinity and specificity" (Darnall et al., 1986a). This is also the case with procaryotic cells, which possess "unique negative surface charges related to the various groups within the cell walls" that can "exhibit a certain degree of selectivity..." for binding specific ions (Ferris et al., 1986; for specific examples see Doyle et al., 1980; Beveridge et al., 1982).

In many cases, however, a particular component, usually a polysaccharide, dominates the cell surface, particularly in those strains that exhibit a thick sheath, mucilage or capsule (the distinctions between these being mainly morphological). In such cases, this is the main site where metal adsorption takes place. Indeed, the frequently observed elaboration of such mucilaginous materials in response to metal toxicity indicates that surface polysaccharides may be produced as metal binding agents in some microbes. However, mucilage is also produced in response to other adverse environmental conditions, from chlorination to desiccation, and thus may not represent a specific protection mechanism against heavy metals. Whatever the rationale, it is clear that such polymers do function in metal adsorption and ion exchange and are of particular interest as they may be able to be isolated and provide a more defined ion exchange material than whole microbial biomass.

The cell wall is usually rigid and resists the high internal (several atmospheres) pressure inside the cell. (Beveridge, 1988). It is negatively charged and adsorbs Mg^{2+} and Ca^{2+} ions, which are essential for the maintenance and properties of the wall (Ferris et al., 1986). The important role of cations in maintaining cell wall integrity can be demonstrated by washing microbes with EDTA, a metal chelator, which removes both Mg^{2+} and other ions and liberates large amounts of surface polymers (Beveridge, 1989). Cell walls also adsorb, to differing extents, other metals including toxic ones (Hg, Cd, Cu, etc.), which exchange for the normal cations in cell walls.

There is, of course, great variability among the cell walls of different microbes, with major chemical divisions between the procaryotes and eucaryotes, and further divisions among these. Thus, the cell wall structures among various procaryotes differ fundamentally, with a major division being the gram positive, gram negative and archaebacterial type of cell walls.

The gram positive bacteria, such as *Bacillus subtilis*, and the cyanobacteria (blue-green algae) have cell walls composed of a rigid peptidoglycan (polysaccharide backbone with peptide side-chains) to which secondary polymers (teichoic and teichuronic acid) are attached. Extensive covalent cross-linking make this cell wall essentially a single molecular structure and accounts for its extreme toughness. Gram negative bacteria, such as *Escherichia coli*, have a more complex cell wall, also consisting of a peptidoglycan material, but much thinner than that of gram positive bacteria and with a lipopolysaccharide outer surface. A large variety of proteins are embedded in the cell wall and underlying cell membrane. These proteins function in various capacities, including metal binding and transport.

In general, gram positive bacteria walls bind more metal than gram negative cell walls, due to the higher anionic charge capacity of the former (Beveridge and Koval, 1981). However, the actual binding agents, on the cell walls (e.g., teichoic or teichuronic acids) can vary, depending on the actual organism studied and their growth conditions. Removal of teichoic acids from the cell wall did not result in loss of most of the binding capacity, which remained with the peptidoglycan (Beveridge and Murray, 1980a). Some gram positive bacterial cell walls can interact with heavy metals as open, low density ion exchange resins (Marquis et al., 1976), while others exhibit at least some selectivity, with an increased partitioning ability (Beveridge and Murray, 1980a,b; Beveridge et al., 1982; Doyle et al., 1980).

Gram negative bacteria, with their thin peptidoglycan cell wall and an outer lipoprotein membrane are quite different from the gram positive bacteria, and generally do not bind metal as avidly (Hoyle and Beveridge, 1983). However, in some cases gram negative bacteria can also mediate a strong metal deposition processes (Beveridge and Koval, 1981). Any generalizations are tenuous, as there is great variability in terms of the actual polymers (and, in this context, metal binding agents) presented by these microbes to the surrounding environment. Although the archaeobacteria have not been as well studied as other procaryotes, they are also involved in metal binding (Beveridge, 1989).

Not only does the microbial type and species, even specific strain used, affect metal binding, but the exact experimental conditions (nutrients, temperature, pH and other ions) are also critical. Thus, Baldry and Dean (1981) found that Cu uptake varied greatly depending on which nutrient (e.g., C, Mg, N, K, etc.) limited cell growth in continuous cultures of the bacterium *Pseudomonas fluorescens* and *Bacillus subtilis*. Even in batch cultures (nonlimiting growth), several studies showed that the stage of the batch culture (lag, log, linear, stationary) affected metal binding (e.g., Townsley and Ross, 1985).

The physicochemical characteristics of the solution being treated, particularly pH and redox potential, are major factors influencing algal bioaccumulation. Generally, algal cell walls act as polyfunctional cation exchangers with negatively charged ligands depending on the pH and metal species. Adsorption on cell surfaces typically decreases with pH until it is close to zero at pH 4 (Ahlf, 1988). Another type of bioaccumulation involves the precipitation of metal ions at cell wall surfaces via metabolic activity. This latter type of bioaccumulation was recently shown to be very significantly influenced by redox potential (Ahlf, 1988). Studies with the microalga *Cyanidium caldarium* showed that bioaccumulation of Cd, Cu, Cr, and Pb were negatively correlated with redox potential while bioaccumulation of iron was positively correlated with redox potential (Ahlf, 1988).

In addition to bacteria and algae, the literature on heavy metal binding by fungi and yeasts is also extensive (see Gadd, 1986a,b and 1989 for reviews). Differences in cell wall chemistry between the different groups, and genetic variability within a single species, affect their metal binding also. Examples of microbial ion exchange in fungi and yeast include radium adsorption to Penicillium chrysogenum (Tsezos and Keller, 1983) which exceed the capacity of conventional sorbents by almost 14 fold and the use of Rhizopus arrhizus for recovery of Uranium (Tsezos, 1984). As with bacteria, fungi also bind metals differently depending on the batch culture growth phase and the composition of the growth media, such as the presence of Mg or Ca, due to their influence on the cell wall component biosynthesis (Treens-Sears et al., 1984a,b and 1986).

The fungus Rhizopus arrhizus was found to absorb a variety of different metal cations and anions, but did not absorb alkali metals (Tobin et al., 1984). The amount of uptake of the cations was directly related to their ionic radii, which accounts for the stronger binding of the toxic heavy metals. The uptake of all cations was consistent with absorption of the metals by phosphate, carboxylate, and other negatively charged groups. However, anions such as vanadate and molybdate, are also adsorbed to the surface of these microbes, and the pH dependence of this adsorption indicates electrostatic attraction to positively charged functional groups (Tobin et al., 1984). Thus, microbial cell walls can function as both anionic and cationic ion exchangers.

Dead (generally heat killed or dried) microbial biomass exhibits different metal binding properties than living biomass, with the method of treatment also affecting metal binding. In some cases metal binding rates increase after heat treatment of the biomass, without increasing total metal bound (Siegel, et al., 1987). Other activators included alkali and dimethylsulfoxide pretreatment (Galun et al., 1987a,b). Heat killed cell of the alga Chlorella adsorbed more metal than living cells (Nakajima et al., 1981). Similarly, Kuyucak and Volesky (1988b) recently reported that nonliving yeast cells exhibited greater U, Zn, and Cd binding than living cells. This was not the case with heat killed gram positive bacteria like Bacillus subtilis and, indeed, is not likely to be a general phenomenon.

There is now considerable information about the chemical nature of microbial cell walls and a good understanding of the basic mechanisms of metal adsorption and binding to the surfaces of microbial cell walls and polymers. However, the study of metal binding to cell walls is still to a large degree, empirical. The challenge of basic research in this area is to develop screening protocols that predict the biosorption capabilities of a particular biomass from a select number of measurements.

3.3 Mechanisms of Biosorption

The mechanisms and kinetics of metal binding to microbial biomass, living or dead, can be complex. In general, a two step process of biosorption is observed: simple ion-exchange followed by the formation of complexes (coordination compounds) which may be chelates (Hunt, 1986). The first, ion-exchange, phase is relatively rapid, normally being complete in a few minutes (typically 5 to 10), and exhibits reversible saturation equilibria (e.g., the amount bound depends on the concentration in solution). Both cation (carboxyl, phosphate, sulfate) and anion (N-compounds) ion exchange binding sites are present on microbial cell walls.

In most cases, this phase is followed by a slower reaction resulting in either the uptake of the metal intracellularly or the deposition of the metal on the surface of the cell, in much larger amounts than could be accounted for by the ion exchange groups available (Beveridge and Murray, 1980a,b). In a few cases, an even slower phase, resulting in actual crystallization of the metals (again either intra- or extracellularly) is observed. These mechanisms can be observed both with living and dead biomass, although intracellular uptake is usually (though not always) limited to living organisms.

The intracellular absorption and surface crystallization of metals will be addressed in the next section. Here we only deal with the surface adsorption of metals, "biosorption", which is defined in this context as the relatively easily reversible binding of metals to microbial cell walls, involving ion exchange and chelation processes.

Such binding can be very strong if it involves chelation to proteins, which would make the process irreversible from a practical perspective. For example, Murray and Kidby (1975) found that Hg was adsorbed to yeast cell walls in large amounts (equivalent to the weight of the cell wall) and the binding was strong - the Hg^{+} ion was not eluted by NaOH or EDTA. Evidence indicated that the metal ion specifically bound to a major cell wall protein.

The biosorption of heavy metals onto cells by ion exchange mechanisms is due primarily to ion pair formation, e.g., the negatively charged phosphate, carboxyl or sulfate groups attach to the positively charged metal ions, displacing in the process more weakly bound cations (sodium, magnesium, etc.). Indeed, it is possible to establish "affinity sequences" for the relative binding of various metals for a particular biomass, condition, etc. Numerous such affinity sequences have been published (see Gadd, 1989).

The general pattern of such affinity sequences is that the divalent ions are much more strongly bound than the monovalent alkaline earth ions and binding depends on ion radius, accounting for the stronger binding of the heavy metals with larger ionic radii. However, other mechanisms and forces are also involved, such as Van der Waals forces and formation of metal coordination complexes. These secondary mechanisms affect the kinetics of metal binding, as well as binding affinities. Thus, although biosorption processes are similar to conventional ion exchange systems, there are fundamental differences between them, which allow biosorption systems to have a greater specificities, higher affinities, and larger capacities than is possible with available ion exchange resins.

Metal binding to microbial biomass exhibits typical adsorption-desorption equilibria. Adsorption isotherms are a plot of metal adsorbed by the biomass (X , usually expressed as mg metal/g dry weight biomass) as a function of residual metal in solution after adsorption (C = mg/L). Adsorption isotherms are a most important measurement, as they determine both capacity and affinity. The data are often plotted as Freundlich ($\ln X = [1/n]\ln C + \ln K$, where n and K are constants) and/or Langmuir ($X/C = C/b + 1/Kb$, where b and K are constants) isotherms. Usually the data fit either equation reasonably well, often better the former than the latter, even though the latter is derived from mechanistic adsorption-desorption equilibria and the former is empirical (its constants have no physical reference).

Regardless of equation used, such isotherms are of predictive value, as they can be used to determine systems capacities and for process design. Of course, the actual biomass must be used in such experiments). Most important, they allow a direct comparison over a wide range of metal concentrations of interest. Finally, they allow a measure of the concentration factor ($F = [X/C] \times 1,000$ in the above units) for the biomass in question, at various metal concentrations.

The adsorption isotherms are dependent on many experimental factors, of which pH and competing cations are the most important, as evidenced from a large literature. One parameter that should not, in theory, be a factor, but in practice is found to significantly affect binding isotherms, is biomass concentrations. This could be explained by the fact that at increasing biomass concentrations the interaction between the biomass particles results in a decrease in the available binding sites.

The binding isotherms do not consider the kinetics of such a process, as they are carried out under equilibrium conditions. (The biomass is contacted with the metal solution for sufficient time to achieve equilibrium). As mentioned, biosorption is usually a relatively fast process, with equilibrium being attained in a few minutes. However, if there are significant diffusion limitations, such as with immobilized or pelletized biomass, much longer time periods, even several hours, are required for the process to come to equilibrium. In some cases, where secondary reactions result in an irreversible precipitation, binding kinetics can be complex. A typical example of this is for U binding (see discussion below).

Selectivity, the differential binding of metal ions, is one of the major issues in this field. There are distinct preferences for metals by various microbial biomass sources, with divalent toxic heavy metals (e.g., Cd, Pb, etc.) generally adsorbing much more strongly than Ca^{2+} or Mg^{2+} (see Gadd, 1989 for a review). Also, biosorption can be affected by other cations or protons (Sakaguchi et al., 1979; Nakajima et al., 1981). However, in general, there are relatively few studies where several metals were presented simultaneously or biomass was exposed to solutions of similar compositions as would be found in process waters (for examples of exceptions see Dugan and Pickrum, 1972; Lakshmanan, 1986; McCready and Lakshmanan, 1986).

3.4 Metals Desorption and Biomass Regeneration

By the above definition, biosorption refers to bioaccumulation processes in which the metals can be desorbed from the biomass relatively readily and the biomass reused in further adsorption cycles. Although biomass reuse may, in theory, be feasible with intracellular bioaccumulation processes, in practice, it has only been demonstrated with surface bound metals.

The process by which the metals are desorbed is a major parameter in the design of a bioaccumulation process: the milder the conditions the lower the cost of the process is likely to be, both in terms of the requirement for chemicals and the degradation of the biomass, which can be severe in many cases. Besides cost and biomass recovery (reuse), the other major optimization factor in desorption is the concentration of the metal in the eluate: there is little point to the exercise unless the metal concentration in the final waste or recovery flow exhibits a large increase over the original waste or process stream.

Thus, the parameters of interest in desorption are: the conditions required for elution, the residual metal on the biomass (e.g., recovery), the volume required for elution, and the number of regenerations feasible (e.g., the amount of degradation per cycle). Of course, if the biomass is cheap enough, and binds sufficient amounts of metal, no recycling would be required, as new biomass could be added on a continuous basis (to a fluidized bed, for example) or the columns combusted or otherwise discarded after use. This has been proposed (Shumate et al., 1980; Voiesky, 1987) in the case of U removal with biomass material obtained from a waste treatment process. However, in general, no such ready source of biomass is available. Thus, desorption is usually a critical parameter in the evaluation of bioaccumulation processes.

There have been many studies carried out regarding metal desorption and biomass reuse. A few representative ones are mentioned here. Norberg and Ryding (1984) and Norberg and Persson (1984), used Zoogloea ramigera, to sorb Cu and Cd to nearly 0.3 and 0.1 g/g dry weight. Acid treatment rapidly desorbed the material in several cycles with some selectivity in elution provided by pH. Acid treatment can also be effective with fungal and algal biomass (McCready and Lakshmanan, 1986; Treen-Sears et al., 1984a; Tsezos, 1984). Zimnik and Sneddon (1988) were able to desorb Al from microalgae biomass with acid and reuse the algal biomass about half a dozen times. However, eventually acid treatment can destroy the ion exchange properties of the biomass (although there are, as usual, exceptions (Scott et al., 1986).

A less destructive treatment is the use of bicarbonate solutions. This treatment was very effective in recovering U from fungal biomass (Galun et al., 1983). Similarly, Tsezos (1984) tested six different eluates and also concluded that bicarbonate was the best choice for the elution of U from Rhizopus arrizus. That treatment allowed biomass reuse for several cycles, with 90% of the original U uptake preserved. However, bicarbonate solutions do not work in all cases: for the desorption of radium, Tsezos et al. (1987) were forced to use HCl, which resulted in degradation of their immobilized biomass after only two to three cycles.

Where applicable, bicarbonate has been a widely used choice in biomass elution, both because of its gentleness to the biomass and effectiveness in elution (and relatively high concentration factors achieved), in most cases. Thus, it has been applied in heavy metal removal using microalgal biomass (e.g., Nakajima et al., 1982; Nakajima and Sakaguchi, 1986). In another example, involving surface deposition rather than biosorption (see next section), Macaskie and Dean (1984), using immobilized cells of Citrobacter, found that adsorbed Cd could be readily recovered using a bicarbonate solution.

Many different elution processes have been tested for their use in desorbing heavy metals from biomass with varying degrees of success (Gadd, 1989). One of the most promising areas of future research is the potential for differential elution of the metals sorbed to biomass. Darnall et al. (1986b) found that polyacrylamide immobilized Chlorella, to which a mixture of Cu, Zn, Hg, and Au had been sorbed, would differentially elute these metals by first washing the column with acetate buffer at pH 5 (Zn) and then pH 3.5 (Cu) followed by washing with mercaptoethanol at pH 5 (Au) and pH 2 (Hg). Such selective elution is of obvious interest in applications where normally a large variety of metals are present in the feed streams and their selective recovery and separation is of interest.

4.0 OTHER MICROBIAL BIOACCUMULATION PROCESSES

4.1 Biodeposition Reactions

In some cases, the cell wall acts as a nucleation site for metal deposition, resulting in much larger metal accumulation than would be possible with normal ion exchange materials (Beveridge and Murry, 1980b). Such cell surface precipitation reactions can be metabolism independent (they can occur with dead cells) but, in many cases, they are affected by cell metabolism, either through oxidation-reduction reactions, or excretion of metabolites. Oxidative and reductive processes are involved in the deposition of Fe and Mn on cell surfaces of iron and manganese bacteria, resulting in extensive precipitation of these metals by bacteria such as Metallogenium and Sphaerotilus (see Ghiorse, 1984 for a review). However, these aspects are not discussed further in this report, which focuses on toxic metals.

Cell surface precipitation reactions occur at a much slower rate than biosorption (mainly ion exchange) reactions, which often are a prerequisite. Metal deposition on the surface of microbial cells accounts for the very high metal bioaccumulation observed by many authors with a variety of metals and microbes: About 50% by dry weight lead compounds were associated with the bacterium Micrococcus luteus, essentially all of which was present on the cell surface (less than 1% being cytoplasmic) and it had no apparent effect on viability (Tornabene and Edwards, 1972). Similarly, aggregation of hydrated iron oxides and crystallization of elemental gold can occur on Bacillus subtilis cell walls (Beveridge and Murray, 1980a,b; Hoyle and Beveridge, 1983). Wood and Wang (1983) described the precipitation of NiS dendritic crystals at algal cell surfaces, a reaction which may be useful in precipitation of toxic metals.

A classic example of cell precipitation is the accumulation of U on cell surfaces, often in very large amounts. The fungus Rhizopus arrhizus accumulates U and Th at over 180 mg/g, double the capacity of IRA-400 ion exchange resins (Tsezos and Volesky 1981, 1982). This large amount of metal accumulation appears to involve three sequential mechanisms. First, there is coordination of the metal to the amine nitrogen (with participation of a free radical of a OH group) of the chitin component of the cell wall. This is a classical ion exchange adsorption process; however, it only accounts for 6 mg/g adsorption (Tsezos, 1983). After this initial adsorption, the complexed U acts as a nucleation site for accumulation of additional uranium ions, fitting a Freundlich adsorption isotherm model and accounting for 66% of the U bound, to 120 mg/g. Finally, a slower process, requiring about 30 minutes to completion, involves hydrolysis and precipitation of uranyl hydroxide on the cell wall, adding another 60 mg U/g of biomass (see Tsezos, 1986 for a review).

Galun et al., (1982) also studied adsorption of U to fungal biomass cell walls and recently reported that U bioaccumulation by Penicillium digitatum resulted in deposits of UO_2Cl_2 on the surface of the hyphae, as revealed by X-ray dispersive analysis (Galun et al., 1987a). In another study, Friis and Myers-Keith (1986), working with lyophilized cells of Streptomyces longwoodensis, which had a high capacity for bioaccumulation of uranyl ions (0.44 g/g dry biomass), bound the U through both the cell surface and intracellular components, and bioaccumulation correlated with phosphate content of biomass components.

Workers at the Oak Ridge National Laboratory (Shumate, et al., 1978, 1980; Strandberg, et al., 1981) pioneered the study of U accumulation by microorganisms. They observed that there was a distinct difference between the accumulation of U by yeast (*Saccharomyces cerevisiae*) and bacteria (*Pseudomonas aeruginosa*), with the latter accumulating U primarily intracellularly in an extremely rapid (<10 sec) metabolism independent reaction, while the former adsorbed U via a surface mediated reaction, from which the U could be removed and the biomass reused. Interestingly, electron microscopic observations of the cells revealed that less than half of the cells of either organisms actually accumulated any metal. Thus, some individual cells contained two to three fold more metal per gram than the biomass as a whole. The reason for this is not clear. (See also Shumate and Strandberg, 1985). These observations are similar to reports of biodeposition of U and other metals in studies of microalgae (Fisher, 1985; Horikoshi et al., 1979).

There are a number of mechanisms by which microbes can induce the precipitation of metals at the cell surface. One involves hydrogen peroxide (a significant metabolic by-product in many algae and some aerobic bacteria), which can result in surface non-enzymatic oxidation of the metals and their subsequent precipitation (Wood and Wang, 1983). Another is due to the secretion of metal binding agents, such as oxalic acids excreted by some fungi, resulting in metal (typically Cu) precipitation at the cell surface (Murphy and Ley, 1983).

Cell surfaces can also accumulate particulate materials. Cu, Pb, and Zn sulfides were adsorbed by fungal mycelia (Wainwright et al., 1988). Heavy metal removal from polluted waters can also occur through metal accumulation and trapping of colloidal metals in algae (Jennett et al., 1982). Algal cells were found to bind large amounts of Au through a reductive process by which ionic gold is biodeposited as atomic gold (Hosea et al., 1986).

An interesting biodeposition reaction, described by Dean and colleagues in England, involves cell surface phosphatases. These enzymes allow cells to hydrolyze organic phosphate molecules and adsorb the phosphate moiety. In the presence of a suitable substrate (e.g., glycerol-2-phosphate) and a metal, the hydrolysis of the phosphate results in the precipitation of a metal phosphate complex on the cell surface (e.g., Baldry and Dean, 1980, Aikin and Dean, 1979a,b). In experiments using living (but non-growing) cells of the bacterium *Citrobacter*, a total of 100 mgPb/g dry weight biomass was accumulated, but on addition of glycerol-2-phosphate, the adsorption of the lead increased by over three-fold (Aikin and Dean, 1979a,b). This process was extensively investigated in recent years by this group and the metals bioaccumulated by this process include U and Cd (Macaskie and Dean, 1989). For Cd adsorption, the surface precipitate was identified as CdHPO_4 (Macaskie and Dean, 1987). However, there are significant difficulties in practical applications, since not only must living cells be used, but a relatively complex substrate (organic phosphate) is required.

4.2 Anaerobic Bioprecipitation

In sediments and under many anaerobic conditions sulfate is reduced to hydrogen sulfide by sulfate reducers. Sulfide, of course, has a high affinity for many metals. The use of sulfate reducers for the precipitation of metals from industrial and other effluents has been proposed (Dockins et al., 1980; Illyaletinov, 1985). Brierley et al. (1980) suggested heavy metal removal by sulfide precipitation in uranium wastewaters, however nutrient limitations in these systems resulted in only partial removal of U, Se, and Mo.

Heavy metal removal by anaerobic sulfide production requires an organic substrate to serve as the electron donor in sulfate reduction. Microalgae growing in polluted waters can provide such organic electron donors when they settle and decompose in the sediments. Jackson (1978) found that heavy metals were removed in a series of lakes and streams that received sewage and effluents from mining activities. These systems contained algae and sulfate reducing bacteria.

More direct reactions have also been reported. Wood and Wang (1983) found that the acidophilic alga Cyanidium concentrated 20% by weight heavy metals when grown on acid mine drainage waters, and that these metals were in the form of surface precipitated metal sulfides, implicating a sulfate reduction process. In another study, production of H₂S by fungi resulted in heavy metal precipitation and accumulation on the cell wall (Minney and Quirk, 1985).

Thus, the concept of using anaerobic bioprecipitation reactions for heavy metals removal is both straightforward and reasonably well documented. There are, however, some caveats. One is that the presence of extracellular microbial heavy metal complexing agents could greatly affect the equilibrium constants for the metal sulfides, increasing metal solubilities.

Using natural ecosystems, such as ponds, swamps, wetlands, or their engineered analogues, to remove metals through anaerobic bioprecipitation could result in long term problems: if the anaerobic conditions are ever removed a rapid oxidation of the metal sulfides would result (bioleaching) which would convert the sulfide to sulfuric acid and release the soluble metals. Thus, "natural" systems, e.g., existing wetlands, should not be used for such purposes and engineered systems must include a decommissioning phase in which the accumulated sediments are removed to a long term repository where exposure to aerobic conditions would be prevented indefinitely.

In this context, the use of anaerobic digesters and anaerobic biomass for heavy metal removal should be considered. The role of toxicity, and fate of heavy metals in anaerobic digestion of municipal and industrial wastes has been reviewed by Rudd (1987). Morper (1986) reported on an upflow sludge-blanket reactor (USBR) which was fed a high Cu containing soluble feed. Elimination of 90% of the incoming copper was observed. The copper ended up in the biomass sludge due to Cu precipitation. A laboratory model revealed a clearly visible cuprophilic zone in the reactor which could not "be explained by mere adsorption and precipitation, but must be associated with biological action" (Morper, 1986). Ag and Au were also accumulated from waste waters with high efficiencies and at relatively high concentrations from rather dilute waste waters. Thus, anaerobic digestion processes, and anaerobic biomass warrants further consideration in the development of bioaccumulation processes.

4.3 Intracellular Metal Bioaccumulation

Active transport, also termed energy dependent uptake, requires metabolic energy input. The mechanisms involved are not yet fully understood: metal influx is probably driven by the membrane potential and more or less specific membrane proteins (the "transporters"). In addition, the metals are often bound to intracellular proteins and/or other polymers. Specific uptake systems for particular metals have been demonstrated in several systems, such as those involving Fe, Ni, Co, Mo, etc. Some general properties of such systems were discussed earlier, including the fact that

such uptake systems can be either specific for a single metal or capable of actively transporting several distinct metals.

Considerable complexity is encountered. Thus, at low concentrations (μM), a combined Zn and Mn uptake system can be demonstrated, but at higher concentrations (mM) Mn is taken up by the Mg transporter system (Perkin and Ross, 1986). However, this picture does not always hold: in Brevibacterium ammonigenes, Mn and Zn transporters are distinct, even at low metal concentrations (Schmid and Auling, 1981).

Another interesting mechanism of heavy metal uptake is "facilitated transport" in which the heavy metals are transported into the cell when complexed with an organic molecule. Chmielowski and Kalpiska (1986) reported that germanium (found at 20 to 280 ppm in ash from coal and coke) uptake by a strain of Pseudomonas putida was facilitated in the presence of catchetol, but not acetate, through a non-specific uptake of the Ge-catchetol complex. A more specific type of facilitated transport is the siderochrome system for iron uptake, discussed in detail in the next subsection.

Some nonessential heavy metals, are also taken up by energy dependent metabolism, such as Cd by some fungi (Norris and Kelly, 1977) and algae. In algae, Cd transport depends on the specific species used: in Chlorella fusca, Cd was taken up by the Zn transporter (Mand and Trombella, 1978), while in Chlorella pyrenoidosa it entered via the Mn transporter (Hart et al., 1979), as evidenced by the relative competitive effects of the two metals on Cd uptake in these algae. Similar differences are observed among the bacteria: in Staphylococcus aureus, Cd uptake is through the Mn transporter system (and is even more efficient than Mn uptake) (Tynecke et al., 1981). In the case of the intestinal bacterium Escherichia coli, Cd uptake competes with Zn transport.

These active transport systems result in the poisoning of the cells in the presence of such toxic metals. In the natural environment, such conditions will result in strong selection pressures, and resistance mechanisms will arise, which can involve loss of these transporters or specific toxic ion efflux systems, usually plasmid encoded (Tyneka et al., 1981). Despite these examples of active transport, in most cases, toxic metals appear to be transported by diffusion followed by binding to intracellular polymers, resulting in cell poisoning.

Not only cations but also metal anions are concentrated by such transporter systems. Thus, a specific molybdate uptake system is found in bacteria, which is competitively inhibited by sulfate (which has very similar structure and ionic radius) (Cole et al., 1986) and, similarly, vanadate enters cells via the phosphate II (high affinity) system (Bowman, 1983). Arsenate, because of its chemical similarity to phosphate is also transported via the phosphate uptake system (Mobley and Summers, 1987). Indeed, its toxicity is due to substitution for phosphate in biochemical molecules, where it hydrolyzes spontaneously. Chromate, another toxic metal anion, can be taken up by microbial cells via the sulfate transport system (Ohtake et al., 1987).

Active transport systems generally exhibit slower uptake kinetics than surface binding (ion exchange) mechanisms, although in at least some cases much greater amounts of metal may be accumulated (Gadd, 1986a,b). One problem is that intracellular uptake would make recovery of the metal more difficult (and reuse of the biomass essentially impossible) than with reversible binding mechanisms. However, this would not be an insurmountable problem in situations where relatively small mass amounts (e.g., in the ppb range) of heavy metals are being removed from

solution. It does suggest that a relatively inexpensive biomass source must be available, as it cannot be reused. It also suggests that greater effort should be placed on developing systems with increased intracellular storage.

The intracellular storage systems exhibit different specificities and discrimination mechanisms than the transporter systems. Thus a metal transported into the cell may not fit into the storage process and accumulate in rather high concentrations. For example, tungstate is a competitive inhibitor of molybdate metabolism, and is taken up by bacterial cells when it exceeds molybdate concentrations. However, uptake is not feedback regulated and intracellular tungstate concentrations can exceed 10% of the cell dry weight. Surprisingly, such high tungstate concentration did not result in any apparent ill effect on cell growth or metabolism (Benemann et al., 1973).

A major disadvantage of such active processes is the need for living, or at least metabolically active, microbial biomass under the conditions of heavy metal removal. The complications introduced by having to deal with living organisms include the need to supply an energy source, and the maintenance of favorable environmental conditions over relatively long periods of time. These constraints could be severe and may limit the practical applications of these phenomena.

However, living microbes in waste treatment has been successfully used in very large plants, treating over 100 million gallons per day, and for a variety of applications, from the oxidation of wastes to nitrification/denitrification and even biodegradation of toxic organics. Thus, the arguments that living microbes can not be used in metal bioaccumulation and removal processes do not have merit.

Indeed, the current biological waste treatment processes such as activated sludge and anaerobic digestion, already play an important part in the removal of significant proportions of the heavy metals found in municipal and industrial waste waters (Lester, 1987). The large variability in metal removal from activated sludge systems is a reflection of the generally uncontrolled conditions in such processes, at least from the perspective of optimization for metals removal.

Since trace metal transport systems typically function with very low concentrations of metals, they are considered by some to be unlikely candidates for metal recovery from metal process wastes (e.g., Olson, 1986, Volesky, 1986). This may be true for concentrated wastes, but would not be the case for more dilute wastes, or for the removal of traces of the most toxic materials, such as Cd, Pb, or Hg, which must be reduced to as low a concentration as possible to meet acceptable drinking water standards (e.g., Sterrit and Lester, 1986). In many situations, sufficient dilution waters are not available. In others, public consciousness of the ultimate consequences of poisoning the global environment with heavy metals discourages dilution as a treatment substitute. Thus, the powerful concentrating mechanisms evolved by microbial systems could be adapted to the needs of environmental protection.

5.0 MICROBIAL METAL BINDING AGENTS

In addition to bioaccumulation processes associated with conventional cell surface properties and the metabolic pathways described above, microbial metal bioaccumulation can also be effectuated by specific intracellular and extracellular metal binding agents, such as metallothioneins, phytochelatin complexes, siderophores, extracellular polysaccharides, and other metal chelating agents. These microbial products suggest the possibility that specific metal binding agents could be mass-produced to develop highly effective and selective processes for metals extraction and removal.

5.1 Metallothioneins, Other Metal Binding Proteins, and Phytochelatins

Metallothioneins are small (generally <10,000 molecular weight) proteins with high metal binding capacities (50-150 mg metal/g protein). Metallothioneins are characterized by a high cysteine content, and it is the cysteine groups which bind the metal (Kagi and Hapke, 1984). They are produced by most higher organisms in response to the administration of some heavy metals (e.g., they are "inducible"), and are also reported from many microbes, including bacteria (Butt and Ecker, 1987; Leeper and Wolfram, 1985).

Considerable work has been carried out on the molecular biology and genetic and protein engineering of these agents, as they have proven useful tools in studying and exploiting gene control in higher organisms. Indeed, metal-regulated DNA sequences are among the most efficient elements for trans-species gene transfer and expression discovered to date. Thus, they have been extensively used in genetic engineering studies and applications (Butt and Ecker, 1987). The famous giant mice were produced by splicing the human growth hormone gene to a metallothionein gene and then transferring this construct into the germline of the mice.

Metallothioneins are of great interest in heavy metal binding as they have a high affinity for heavy metals and exhibit some specificity. However, natural metallothioneins are rather susceptible to air oxidation. Although minor structural changes reportedly render the molecule more resistant to oxygen, without reducing its capability to bind heavy metals such as Hg^{2+} (Holmes, 1988), it is unclear whether sufficient resistance can be achieved in practice. This is an area where protein engineering could produce some results, although that is a technology which is not very predictable in terms of specific project objectives. The problems of lability (not only to oxygen, but to pH, foaming, etc.) is a major one, and, in general, proteins are difficult reagents to use in large scale industrial applications.

Of course, one of the major industrial biotechnology processes, the conversion of corn starch to high fructose sugars, uses isolated enzymes. And, indeed, a recent report on the use of rabbit metallothionein immobilized onto cyanogen-bromide activated agarose beads demonstrated a very high affinity, capacity, and regenerability during the removal of Cd from a feed stream (Furlong et al., 1988). These authors also suggested the use of smaller peptides, related to glutathione, which are produced by plants and are effective binders of Cd. These peptides are now known to be phytochelatin complexes, ubiquitous to all algae and higher plants (Gekeler et al., 1988). Finally, the possibility of using purified specific metal binding proteins derived from the trace metal membrane transport mechanism, can be considered, as these can have a very high affinity for specific metal ions.

It will require significant breakthroughs in biochemical process engineering to allow an economical production of metallothioneins for relatively large scale applications, even if the problems of immobilization, stability, and regeneration are solved. Also, it is by no means clear that sufficiently selective metallothioneins can indeed be produced, to allow the absorption of specific metals in the presence of other metal ions. Although it is possible to consider the possibility of "smart proteins" designed with hydrophobic ends for froth flotation or with other characteristics suitable for matrix attachment, which could be reused (Holmes, 1988), this must be considered a rather far in the future concept. Nevertheless, the potential of using metallothionein proteins is significant and this subject warrants continued attention.

Indeed, metallothioneins have been studied by several groups with the objective of developing practical processes for metal recovery. In one example, Leeper et al. (1985) and Leeper and Wolfram (1985) investigated metallothioneins for recovery of Co from dump leach waters that also contained Ni, Fe, Al, Zn, etc. However, after isolation and characterization of a couple of metallothioneins from Co resistant *Saccharomyces cerevisiae* and from *Bacillus subtilis* this approach was abandoned as being too difficult. Butt and Ecker (1987) in their detailed review of the molecular biology and biotechnological applications of yeast metallothioneins, report on unpublished observations that suggest that gold resistance and uptake are linearly related, and that this is due to metallothionein binding of gold. As gold does not induce metallothioneins (the metallothionein coded by this gene locus is specifically and exclusively induced by Cu) they suggest that a strain in which the metallothionein gene is always expressed would be useful in such an application.

The Cu inducible yeast metallothionein was recently genetically cloned into the bacterium *Escherichia coli*. It was fused to a larger protein (Romeyer et al., 1988) to prevent its breakdown (the fate of small homologous proteins such as metallothioneins). The bacteria produced the fusion protein in high amounts (8% of cell dry weight) and accumulated Cd^{2+} and Cu^{+} intracellularly, even though no enhanced resistance to these metals was noted. These results suggest that it may be possible to augment microbial uptake mechanisms by introduction and expression of metallothionein genes into microbes.

It should be noted that although most attention has been paid to metallothioneins as metal binding proteins, many enzymes contain metals and bind these with both very high affinity and specificity. In more and more cases the three dimensional structure of these proteins, and, in particular, the active site which binds the metals, is being elucidated by X-ray crystallography and by other advanced technologies. The uptake of a particular metal by a specific site is a function of both thermodynamic and kinetic parameters, which, although not yet predictable, are becoming understood for the proteins whose structures we have now available. Using these structures as guides, and with the techniques of genetic and protein engineering, it should be possible to design new and improved metal binding proteins for particular applications. Although such products will require long term R&D, much of the basic understanding on which such a product development would need to be based has already been achieved.

The use of living microbes with a specifically selected, or even purposefully designed, metallothionein and other metal binding protein(s), preferably coupled to an active uptake system, would be an attractive goal for the future. This approach is certainly preferable over the alternative concept of using purified proteins in metal removal applications.

Gekeler et al. (1988) demonstrated the ability of algae to synthesize phytochelatins in response to heavy metal exposure. These peptides have been isolated from higher plants and characterized to be $(\gamma\text{-Glu-Cys})_n\text{-Gly}$, where $n = 2$ to 11 (Grill et al., 1985, 1987). A three-step isolation procedure yielded a virtually pure Cd-binding complex from algae as demonstrated by amino acid analysis. HPLC, amino acid analysis, and chemical sequencing followed by 2-dimensional TLC revealed phytochelatin production. Several phylogenetically distant algae were induced to form phytochelatins by a wide range of heavy metals. Algal cells exposed to a variety of toxic metal ions Cd^{2+} , Pb^{2+} , Zn^{2+} , Ag^+ , Cu^{2+} , and Hg^{2+} clearly revealed phytochelatin induction (Gekeler et al. (1988)). The data showed that the ability to synthesize phytochelatins is ubiquitous in the algae. Taking into account the induction of phytochelatins by Cd^{2+} and Cu^{2+} , and the preferential binding of heavy-metals by these peptides (Grill et al., 1987), Gekeler et al. (1988) concluded that the previously published Cu- and Cd-binding 'proteins' occurring in algae (Stokes et al., 1977; Hart and Bertram, 1980; Nagano et al., 1984; Gingrich et al., 1986) were actually phytochelatin complexes that were erroneously reported as proteins.

It appears that all types of algae are able to inactivate heavy metal ions by a cytoplasmatic detoxification mechanism, the phytochelatins. Thus, an evolutionary divergence in heavy metal sequestration has occurred between plants and animals. Animals respond to heavy metal stress by induction of metallothionein gene expression, plants by induction of phytochelatin formation via enzymic polymerisation of peptidic precursors.

5.2 Siderophores and Other Complexing Agents

Microbes excrete a large number of organic compounds, often in significant amounts. Among these are low molecular weight (generally $<1,000$) Fe and Cu (and possibly other metal) complexing agents, although these compounds also complex other heavy metals. The Fe^{3+} complexing agents are the best studied - they are collectively known as siderophores.

Chemically, siderophores are characterized as hydroxamates and phenolate-catecholate derivatives, although several other types of structures are also found. Their major characteristic is their extremely high affinity for Fe^{3+} , with dissociation constants in the range of 10^{-30} to 10^{-50} , compared with 10^{-12} for ferric hydroxide. Thus, siderophores can solubilize iron from insoluble particulates. Siderophores are excreted by microbes under conditions of Fe limitation, and their main function is to help solubilize Fe^{3+} from particulate precipitates usually present in the environment. The extremely low concentrations of iron in oxic environments at neutral or alkaline pH, makes iron availability one of the major nutrient supply problems for any microbe. The very high affinity constants of siderophores for Fe^{3+} , as well as the very specific and high affinity ligands present on the cell walls which recognize and internalize (take up) siderophore-Fe complexes, results in iron accumulation by the cells. Once in the cell the ferric iron is reduced and thereby released from the siderophore, which can be again excreted. (Braun, 1985). As far as is known iron is the only metal which is taken up by this type of transport system.

Sometimes more than one siderophore is produced by one organism, as in the case of Azotobacter, which produces possibly four iron binding exudates (azotochelin, azotobactin, 2,3-dihydroxybenzoic acid and aminochelin). At least the first two are coordinately synthesized: over thirty

siderophore lacking mutants lack both of these (the others not having been studied) (Glick et al., 1988). The function of such a variety of different siderophores is unknown. Blue-green algae also produce siderophores (McKnight and Morel, 1979; Clarke et al., 1987) as well as other metal binding extracellular materials (see below).

Although iron is the key metabolite involved in siderophore biosynthesis and activity, other metals also play a not well understood role, as evidenced by the fact that Zn can result in increased siderophore production in at least one bacterium (Hyer and Page, 1988). Siderophores are also known to complex other metals, such as Cu, however with lower affinity constants. Olson et al. (1989), following-up on earlier observations, found that iron deficient Pseudomonas fluorescens cultures excreted siderophores which would result in the taking up of Gallium (up to about 0.23% of dry weight), which was toxic to cell metabolism.

Sometimes, Cu complexing agents are excreted by Fe deficient algal cells, suggesting that they are siderophores (McKnight and Morel, 1979). However, it is unlikely that all extracellular Cu and other heavy metal complexing agents excreted by microorganisms are siderophores. This is indicated from the fact that Cu and other metal complexing activities are induced by the presence of heavy metals and sometimes correlate with heavy metal resistance.

Siderophores and similar metal complexing compounds could be considered in the removal of heavy metals from process and waste streams. This has been demonstrated by DeVoe and Holbein (1986), who covalently linked siderophores to a matrix to remove ferric iron from growth media (to be able to obtain purified media that does not support growth in the absence of added iron). These same immobilized siderophores were tested in the removal of plutonium from high salt containing nuclear fuel reprocessing waste liquors, which resulted in undetectable levels of Pu in the column effluents. These authors also patented siderophore analogues which were demonstrated to be highly effective in removing both radioactive and nonradioactive toxic metals from plating effluent, cooling water blow-down, photographic wastes, and other waste waters. Removals in all cases were essentially complete, with no detectable residues in the column effluents. An important aspect of this work is that it was carried out with actual wastewaters, containing high salt concentrations and interfering metals. Practical applications are listed, specifically that of commercial units that remove Ag to below 0.06 ppm, recovering one ounce of Ag per kilogram of absorbent. This material has promise and should be further evaluated.

5.3 Extracellular Polysaccharides and Other Exudates

Microbes release a significant amount of organic matter into their environment, including polysaccharides, proteins, enzymes, amino acids, other nitrogenous organic compounds, organic acids, vitamins, etc. Although a fraction of this material could be considered leakage from the cytoplasm and periplasmic space, unwanted end products of metabolism and the remains of cell lysis, part is excreted for some specific function important to the survival of the microbe. Even anaerobic fermentative bacteria, which must excrete organic molecules to balance their redox reactions, excrete compounds which are not related to the requirements of intermediate metabolism.

Of course, in the laboratory, such activities may be baffling, unless a specific function (e.g., antibiotics) can be ascribed to such products. However, in nature, in competition with a myriad of other organisms for limiting nutrients and energy sources, such extracellular materials

could aid in the survival of the cell. One function of excreted organic compounds would be in managing the speciation, availability, and toxicity of the metal ions present in the environment. This was suggested, for example, for polysaccharides excreted by microalgae (Crist et al., 1981).

Among these microbial excreted compounds, the polysaccharides represent a major fraction and have been the subject of intensive investigations regarding their metal binding properties, particularly in relation to their role in metal removal from waste treatment systems (Lester, 1987; Sterrit and Lester, 1986). Algae often excrete polysaccharides, some of which have been shown to bind Cu (Strong et al., 1982; Kaplan et al., 1987a,b). The role of copper complexing agents in making blue-green algae resistant to this potent algicide is well established (Jardim and Pearson, 1985). Extracellular products of microalgae have also been demonstrated to bind other heavy metals, including Zn^{2+} and Cd^{2+} (Kaplan et al., 1987a,b; Imber et al., 1985). Large variations in both polysaccharide excretion and specific metal binding were noted between different species and growth conditions. It should be noted that there is not much, if any, difference between the exopolysaccharides elaborated by the cells as mucilaginous material exterior to the cell wall and the excreted polysaccharides, which are the fraction of these materials that have sloughed off from the microbial cell wall and are found freely dissolved in the culture media. The amount of such materials produced, and the fraction which actually is sloughed off, or which can be readily removed from the microbial biomass (by a treatment such as washing with a saline solution or warm water) depends to a large degree on the growth conditions of the microbes. Thus, in this context, metal adsorption by extracellular polysaccharides and loosely associated polysaccharide is equivalent.

A number of workers have reported on the binding of various ions and metals by purified extracellular polysaccharides from *Zooglea ramigera* (Dugan and Pickrum, 1972), *Pseudomonas* sp. (Corpe, 1974), and *Klebsiella aerogenes* (Brown and Lester, 1982). The latter authors found that after gently removing extracellular polysaccharides from the cells of a capsulated strain of *K. aerogenes*, metal adsorption by the cells decreased about 90%, while the soluble polysaccharide adsorbed most, but not all, of the metal complexed by the intact microbes.

Norberg and colleagues (Norberg and Persson, 1984; Norberg and Rydin, 1984) showed that the activated sludge bacterium *Zooglea ramigera*, was an active bioaccumulator of various heavy metals, due primarily to the acidic polysaccharide produced by this organism. The high uptake of metals by this microbe in activated sludge was ascribed by Sterrit and Lester (1986) to mutually exclusive roles of the extracellular polysaccharides in metal binding and flocculation.

Considering the central role of extracellular, or cell wall, polysaccharides in metal binding, the question arises of whether microbial polysaccharides could be used as metal binding agents, thus avoiding the complications of using the entire biomass, of which only a fraction is actually useful in metal binding. Indeed, Muzarelli and co-workers have for several years studied the use of both microbial and other biological materials, as well as modified and synthetic, polymers in metal binding (Muzarelli et al., 1985, 1986). Several of the polysaccharides and derivatives studied by this group, as well as others, exhibit binding isotherms comparable to those of microbial systems.

In comparing metal binding by microbial polysaccharides with microbial biomass, the advantage at present is definitely with the latter. The cost of isolating and purifying the polysaccharides does not make-up for the value added. More importantly, the biomass generally has a higher metal binding capacity, as found with chitin from *Aspergillus niger* (Muzarelli et al., 1986).

6.0 SOURCES OF MICROBIAL BIOMASS FOR BIOACCUMULATION

6.1 Microbial Biomass Sources.

To be useful as ion exchange materials, or for metal bioaccumulation generally, the microbial biomass, must meet a number of utility criteria: capacity (g metal/g biomass); specificity; reproducibility; minimal interference by other ions; reusability; and, of course, economics (Lakshmanan et al., 1986). Although each of these issues has been addressed in various studies, few studies have considered all these requirements as they may apply to a specific application and there is still a considerable amount of R&D needed to develop microbial ion exchange materials which deliver high utility at a reasonable cost.

The issue of economics explains the interest in using commercially available biomass materials, such as bakers yeast (Saccharomyces cerevisiae), microbes used in antibiotics production (Penicillium sp., Streptomyces sp.), activated sludge (including constituent bacteria such as Zooglea ramigera), anaerobic sludge, denitrification biofilms, citric acid fermentations (Aspergillus niger), food fermentations by-products (Bacillus subtilis), and microalgae that are produced commercially (e.g., Chlorella and Spirulina).

Each of these has been used by various groups for bioaccumulation processes and/or bioaccumulation studies. It should be noted, however, that many studies that used the above listed microbes, actually carried out their research with laboratory grown biomass cultures, rather than obtaining these from commercial sources. Thus, the data in the literature may not always extrapolate to the use of commercially available materials.

Commercially available fungal and bacterial biomass has been used in the development of commercial bioaccumulation products. The best known of these are the AMT-BIOCLAIM pellets of Advanced Mineral Technologies, Inc., based on a proprietary immobilization process for Bacillus subtilis (although the nature of the microbial biomass is not identified in the publications) (Brierley et al., 1985, 1986). Other commercial products are reportedly being developed: B.V. Sorbex of Montreal, Canada, founded by Volesky, is using proprietary immobilization techniques for the fungus Rhizopus arrhizus and algae (and possibly other fermentation byproducts, see Kuyucak and Volesky, 1988a,b and 1989a,b). Tsezos, also in Canada, has developed proprietary immobilization technology for activated sludge and other biomass sources (e.g., Tsezos et al., 1988) which could become a basis for future commercial products. However, none of the fungal or bacterial based products is currently sold commercially.

Presently, the only commercially available bioaccumulation material, AlgaSORB™, utilizes microalgae biomass, specifically dried cells of Chlorella and Spirulina. AlgaSORB™ technology was developed by Bio-Recovery Systems, Inc. of New Mexico. The development was, at least in part, due to the extensive research carried out by a number of groups on algal biosorption in recent years (for examples: Sakaguchi et al., 1979, 1981; Horikoshi et al., 1981; Nakajima et al., 1981; Crist et al., 1981; Fisher et al., 1983; Ferguson et al., 1989), and most directly due to the work by Darnall and colleagues in New Mexico (Darnall et al., 1986a,b, 1988a,b; Darnall and Bedell, 1989; Greene et al., 1986a,b, 1987; Bedell and Darnall, 1989; Watkins et al., 1987; Greene and Darnall, 1988; Greene and Bedell, 1989). The proprietary AlgaSORB™ materials are currently used in the metals plating, electronics, and mining industries. The use of AlgaSORB™ has reportedly

increased dramatically during the past several months despite the relatively high cost of the algal biomass (over \$10/kg, even for non-food grade material).

By using biomass specifically produced for bioaccumulation applications, it should be possible to significantly improve the metal binding capabilities of these materials. The trade-off is between economics and utility: a biomass specifically produced for metal adsorption would be relatively expensive and the increased utility may not warrant the additional cost, compared to the biomass sources available as by-products from other processes. Although the use of available biomass sources are presently favored from an economic perspective, in the future, as markets expand, the production of specific biomass materials for such applications is likely to become practical, and even necessary.

The question thus arises of how superior microbial strains are to be developed: through screening of existing cultures and new isolates or by genetic improvements. These subjects are addressed next.

6.2 Screening for Bioaccumulating Microbes

Considering the many different microbial species and strains, growth conditions, environmental conditions, interferences by metals and other ions, and biomass pretreatments, the effort for screening for an optimal biomass biosorbent could be quite large. The screening protocols are also generally laborious, compounding this problem. Several researchers have carried out screening studies comparing different microbial species, and in some cases strains, for bioaccumulation for one or a few metals. However, in general, these surveys were very limited in both the number of strains tested (generally less than ten) and the conditions under which the tests were carried out.

There have been a few studies where relatively large numbers of different microbial species were screened. For example, a Japanese group (Horikoshi et al., 1981; Nakajima and Sakaguchi, 1986) screened over 100 strains of bacteria and fungi for uranium bioaccumulation, including (in the latter study) the effect of a mixture of different metals. Differences of up to ten fold in U adsorption between different strains were noted, although total metal adsorption differed typically by a smaller factor (about three fold). Macaskie and Dean (1982) screened 174 bacteria and 29 moulds, grown in the presence of 10 mg/L of Cd, for Cd bioaccumulation. Several strains accumulated over 1% by dry weight of this toxic cation. The same group (Aikin and Dean, 1979b) screened 47 bacteria and 9 fungi for Pb uptake, finding that bioaccumulation ranged from 0.1 to 36% of dry weight. Thus, the need for screening a wide variety of microbes is apparent.

The large amount of work required for such extensive screening efforts accounts for the fact that most reports in the literature have dealt with far fewer strains, and most studies deal with the adsorption of a single metal by one microbe. This compares to the literally hundreds of thousands of strains that have been screened for the production of antibiotics by the pharmaceutical industry. Although such extensive screening programs may not be necessary for bioaccumulation strain selection, there are significant differences in the ability of various types of microbes (e.g., algae, yeast, gram positive and negative bacteria, etc.) to bind metals and large differences are also observed between different strains of the same species. Thus, a screening program which allows at least a preliminary assessment of a large number of different strains under standardized conditions is required.

A screening protocol, using a radioactive tracer microplate technique, was developed by Hassett et al. (1981) for microalgae. This method involves using the standard microplate wells. Single strains are inoculated into each well and, after a period of growth, a radioactive metal solution is added. After contacting for a given time, the supernatant solution is drawn off and counted in a scintillation counter. Using this technique, Hassett et al. (1981) tested over ten different algal species, with sufficient replication for statistical analysis. There were very large differences in concentration coefficient between the various parameters tested (pH, culture age, metals, strains), and few generalizations could be made. (One was that bioaccumulation by older cultures was less than 10% as effective, in terms of concentration factors, as by younger cultures).

The results of this research, as well as other less exhaustive studies, suggests that the current protocols for the searching for superior biosorbing microbes are inadequate. However, it is also not clear whether a wider search for improved bioaccumulating strains will be worth the effort. Nevertheless, it is clear that improved screening protocols, along the lines of those of Hassett et al. (1981) are desirable. The question arises of whether only empirical research is possible to provide the best biomass material and operation conditions for bioaccumulation or whether a more theoretical basis can be established to guide such screening efforts. This is an additional area for future research.

In conclusion, the issue is whether to restrict attention to existing biomass sources which can be obtained in relatively large quantities at reasonable prices or to find new and novel microbial strains that exhibit superior characteristics to the available biomass types. This will, to a large degree, depend on the particular application under consideration: its technical requirements and economic constraints. If available biomass sources cannot provide sufficient capacity, removal efficiencies, and recovery (metal or biomass), and if the higher price (several dollars per kg) which would be entailed in a custom produced microbial biomass could be justified, then search for novel microbial metal bioaccumulators would be justified. Another approach is to develop such improved bioaccumulators through strain improvement and genetic techniques, as discussed next.

6.3 Genetic Selection With Continuous Cultures

Genetic techniques could improve existing microbial bioaccumulation systems. Applying genetic engineering methods, except as a tool of basic research, is premature, as there is only very limited understanding of the basic mechanisms involved in the biosynthesis of cell walls and the metal binding functional groups, or bioaccumulation processes generally. However, conventional genetic techniques would appear to be applicable in the near term, although no work appears to have been done in this area thus far.

Continuous (chemostat) cultures could be considered in such research. Indeed, there is some data in the literature: continuous cultures were used to establish a mixed culture of three bacteria which were able to remove 76% of the silver (appx. 100 mg/L) after an almost three month run (Charley and Bull, 1979) and up to 92% of Cu was removed by a more complex mixed culture which exhibited wall attachment (Dunn and Bull, 1983). However, in these cases, the selection pressure in the continuous cultures was not directly related to metal adsorption.

To develop a more effective process for selecting a superior microbial genotype, continuous cultures should be subjected to some type of selection which would result in preferential growth of the best metal adsorbing microbial population. How to achieve such selection pressures for metal adsorption in continuous or batch cultures is the challenge of this research.

One possible approach is to use a recycle system in which the effluent microbial population would be separated based on density, through differential centrifugation. The heaviest cells (presumably those which adsorb the most metal during the process) would be selectively recycled and thereby given a large growth rate advantage over the lighter cells, even if their growth rate is actually slower (as it is likely to be). Such selective recycle systems are a powerful method for controlling and selecting microbial species and strains in continuous cultures (Weissman and Benemann, 1979). The key question in this case is whether, from a practical perspective, differential centrifugation can be carried out with sufficient precision to differentiate between metal concentrating and normal cells. Also, density differences could result from mechanisms other than a genetic improvement in metal adsorption.

If essential metal absorption mechanisms are to be used, then selection pressures could be used which are based on the storage capabilities of the microbial species: short periods of exposure to the metal followed by relatively long periods of metal limitation. This would select for strains which have a high storage capacity for that specific metal. Such selection processes would also select for strains with a propensity to bioaccumulate other, nonessential, even toxic, heavy metals which mimic the behavior of the essential trace elements.

Use of ion specific electrodes or similar online or rapid analytical techniques could perhaps be used to establish a feedback mechanism between the free heavy metal in the culture and the feed supply and dilution rate of the culture. As the free metal concentration drops the dilution rate would be increased - thereby selecting for cultures which would grow relatively fast under conditions which result in metal removal from solution.

Variations on this theme could be to maintain the growth rate, cell concentration, or even productivity (dilution x concentration) constant while manipulating metal concentration and supply. Modern techniques of computer control make such approaches feasible at relatively low cost.

One problem with the latter approach is that the selection is at the culture level, not at the individual microbe level as in the prior examples. If metal adsorption is otherwise counter-selective (e.g., decreases growth rate), as it is likely to be, there would be a balance established between metal adsorbing and non-adsorbing individuals, which would reduce the overall effectiveness of the selection process and, most importantly, would result in the rapid loss of the desired genotypes as soon as the selection pressure is removed. For such cases, the initial culture would be screened for desirable genotypes which are then isolated in pure culture.

One other approach, particularly when dealing with toxic metals, is the use of two stage processes, one for biomass growth and the second for metal adsorption. This could be considered for the last two selection examples discussed. Such a separation between growth and adsorption phases is of significant interest in the removal of heavy metals present at concentrations which normally inhibit microbial growth.

6.4 Screening for Bioaccumulation Mutations.

One approach to developing microbial strains with increased metal bioaccumulation capacities, would be to isolate for metal resistant strains, and screen among these for strains which exhibit higher metal uptake. As discussed above, increased binding of heavy metals by cell surfaces is only one of many resistance mechanisms. Nevertheless, heavy metal resistance is a relatively easy mutation to screen or select for (e.g., by colony formation on plates or by growth in liquid cultures) and should be one of the mainstays in the search for improved bioaccumulation strains. A variation of this approach would be to search for strains highly sensitive for a particular metal. The rationale is that such strains would have an efficient adsorption/uptake system which results in that metal becoming more toxic than normal.

One advantage in screening for metal resistance (or sensitivity) is that this involves identification of individual mutations, rather than selection of whole populations, as in a chemostat. The difficulties inherent in continuous culture selection processes makes such an approach appealing. The major issue is the development of highly efficient (low labor) screening protocols which would allow the rapid searching for a desirable mutation among, literally, tens and even hundreds of thousands of other cells.

Indeed, to speed up the process, a batch growth step is desirable during which the desired mutants would be favored to grow (or at least survive) compared to the wild type. This "enrichment" would be feasible with metal resistance by exposing the culture to the metal for a few generations. Alternatively, sensitivity can be selected for by exposing the culture to a level of the metal which is non-lethal to the wild type in the presence of an antibiotic which kills only growing cells (e.g., penicillin). The survivors of such a process would then be grown in liquid culture for a few generations and replica plated on permissive and nonpermissive media, following standard microbiological practices.

Alternative screening and enrichment techniques, not depending on heavy metal resistance, would be desirable. Again, density differences could be used through the technique of density gradient centrifugation, which is able to separate cells based on density differences. This technique would appear to be applicable to such a genetics program, to seek out a few mutations among a very large background of wild type cells. Cells binding significant quantities of metals (extra- or intracellularly) would be expected to be denser than wild type cells, and even small density differences would result in clean separations. Thus, rather than requiring a single mutational event, this system would allow, as is the case for continuous culture selections, the step-wise increase in metal binding capacity by a particular microbial line, similar to the step-wise improvement in conventional fermentation processes.

If the metal content of the cells can be visualized or detected by some type of rapid spectroscopy, it should be possible to use cell sorting techniques to select for desirable individual phenotypes in a microbial culture. Cell sorting techniques are very powerful and can also be used to establish the variability in a particular culture. It has, for example, been reported that during U accumulation by yeast and bacteria only a fraction of cells actually accumulated this metal, in high amounts, with the remainder of the cells being devoid of the metal (Strandberg et al., 1981). Whether this has a genetic basis is not known.

This type of research is not even at its initial stages. The above discussion only identifies some of the potential approaches that could be followed. Once the techniques for selection and screening have been worked out, they could be applied to a variety of problems and applications. Also, they would provide an entry to the application of more powerful genetic techniques, which could be initially used to analyze the genetics and biochemistry of the mutations which result in improved bioaccumulation and ultimately would result in more advanced bioaccumulation processes based on enhanced understanding of the mechanisms involved.

7.0 PROCESS DESIGN FOR BIOREMOVAL SYSTEMS

7.1 Introduction

In principle, the use of microbial biomass in bioremoval processes is relatively straightforward: the metal containing solution is contacted with the microbial biomass for a sufficient period of time to allow the system to achieve equilibrium or reach a desirable reduction in metal concentrations, and then the microbial biomass is separated from the solution. If feasible, the microbial biomass is regenerated for further use by some selective desorbing solution, with the concentrated metal solution subjected to further treatment (recovery or disposal). Alternatively, the metals are extracted from the biomass by some irreversible process or the biomass is discarded appropriately. In practice, this simple scheme is seldom feasible outside the laboratory.

From a practical perspective, native microbial biomass, whether alive or dead, has several highly undesirable properties: small particle size, low mechanical strength, and relatively low density (Tsezos, 1986). Before microbial biomass can be used as a replacement for ion exchange resin, or in other bioaccumulation applications, these properties must be altered. Otherwise, the costs of separating the contacted biomass from the process streams become an overwhelming economic factor.

Thus, Lakshmanan et al. (1986) and McCready and Lakshmanan (1986) who carried out an economic analysis of the use of several different microbial biosorbents in U recovery, concluded that costs were several fold higher in both capital and operating than current technology (ion exchange). Although the biosorption capacities of the various biomass sources used, 20-60 mg U/g of material, compared favorably with available ion exchange resins, they based their analysis on a process involving a mixed tank reactor followed by a centrifuge for biomass recovery, these being the two major cost factors in the overall process. This demonstrated the need for a low cost immobilization or similar stabilization process for the biomass, which would allow ready contacting and separation of the biomass and metals containing solution.

Thus, the use of biomass materials in bioaccumulation is just as much a problem in process design as is finding suitable biomass materials. The key aspect of process design is in the contacting of the biomass and the metal solution to result in good mass transfers while facilitating easy separation of the two phases after contacting is completed. The major approaches to this problem have been artificial immobilization and natural biofilm or floc formation. These are discussed next.

7.2 Immobilization of Biomass for Bioremoval

There has been extensive work on immobilization of biomass into or on some type of solid support, pellet, bead, granule, or similar structure. The factors that require attention during the immobilization of microbial biomass, whether living or dead, in addition to its bioaccumulation characteristics (capacity, affinity), are good flow rates (low pressure drop), little compression, no plugging or channeling, high surface area to immobilized biomass, good mechanical strength, long service life (e.g., reutilization), and, of course, low cost. If metabolically active cells are to be used, metabolism should not be adversely affected by the immobilization procedure.

One major issue is that the kinetics and capacity factors observed with dispersed microbial cells are significantly degraded during any immobilization process, principally because of diffusion limitations near the surface of the immobilization matrix and due to the decrease in effective surface area of the biomass. From a kinetic and capacity perspective, it would be preferable to use dispersed microbial cells. However, any practical system must separate, at low cost, the biomass and process flow and this often favors some type of immobilization.

The technology for immobilizing microbial cells has been a major focus of biotechnology R&D for over two decades, with literally thousands of publications, scores of reviews, and even a few practical applications. There are several different approaches to such artificial immobilization: chemical immobilization by covalent or non-covalent (adsorption), bonding of the biomass to a (more or less) inert support (including cross-linking the cells to each other), and physical entrapment into a porous structure. The resulting materials are of two types: solid materials, suitable for passing liquid over or through (e.g. columns or other high surface area:volume contactors), and pellets or flocs which can be suspended in the liquid with relatively little mixing energy but which will settle and compact readily for separation purposes. The latter can be used in mixed tanks or fluidized bed reactors. Fluidized bed reactors are particularly applicable to metals removal because as the bead or pellets fill up with metals they become heavier and can be selectively removed from the bottom of the reactor (Brierley et al., 1986; Yakubu and Dudeney, 1986).

Many different immobilization systems, matrixes and chemicals have been used in metal bioaccumulation studies: coal particles for yeasts (Shumate et al., 1980) polyacrylamide gels (Nakajima et al., 1982 ; Greene et al., 1986a,b, Macaskie and Dean, 1984, 1985), silicate gels and, , 1986b; de la Noue and Proulx, 1988), polyvinyl chloride granules, and polysulfonate beads (Ferguson et al., 1989). In several cases, the method of immobilization is deemed proprietary, such as in the work of Brierley et al. (1986) mentioned above and that of Tsezos et al., (1988), who report metal binding results but not the method of immobilization. Tsezos et al. (1988), working with the fungus *Rhizopus arrhizus*, determined that the immobilized particles produced, 0.5 to 1.2 mm in size, had good resistance to compression, high porosity, good wetting ability, low content of inactive material, good resistance to chemicals and favorable equilibrium kinetics, all properties desirable in immobilized biomass for bioaccumulation applications.

Although for some applications these immobilization techniques are adequate, none yet optimize low cost, high surface to volume areas, physical strength (e.g., lack of compressibility, which allows large column operations). This, however, is probably due primarily to the relative undeveloped nature of the field (in many cases questions of costs are still to be addressed), rather than to any inherent limitations. It is expected that the development of better immobilization techniques will remain a major area of applied R&D in this field, as it is a critical aspect to the commercialization of this technology.

7.3 Use of Biofilms in Bioremoval

A more direct method of immobilizing microorganisms than the artificial processes described above, is the use of the natural tendencies of microbes to aggregate as wall growth, flocs and other naturally immobilized biomass assemblages. The formation of biofilms and flocs, depends on the growth properties of the microbial biomass. This suggests that such systems would be most readily utilized in applications where the bioaccumulation is based on living or at least metabolically active biomass, as non-living biomass would tend to slough off or disperse from a biofilm.

Indeed, various types of biofilm reactors, such as rotating biological contactors (RBC's) and attached film reactors, are used in a number of wastewater treatment processes, including nitrification-denitrification and trickling filters. Even the most widely used process in waste treatment, activated sludge, is a type of immobilized microbial reactor, because the activated sludge flocs are aggregations of microbes.

Microbes attach to solid surfaces and each other via extracellular polysaccharides. These polysaccharides coat exposed surfaces as the first step in a biofouling process and then direct the attachment of microbial cells in a reversible binding process that is made irreversible by the elaboration of additional attached exopolymer materials. Although other biofilm forming mechanisms have also been reported (e.g., direct attachment without any apparent prior exopolysaccharide production), this description suffices for this discussion.

A major question in the development of bioaccumulation technology is not the how of microbial attachment but the why. Although a large number of publications belabor this issue, particularly in the sanitary engineering literature, the simple answer is that microbial attachment, either to walls or each other, is the result of strong selection pressures, similar to those which result in wingless insects on ocean islands.

Thus, for example, the activated sludge bacteria form flocs because the flocs settle relatively rapidly in the settling basins and are then recycled back to the aeration tank. The action of recycling of the flocs allows them to out-compete the dispersed bacteria (which are not recycled as they do not settle) even though the microbes in the flocs actually exhibit lower growth rates than the dispersed bacteria. This is a general phenomenon, which is applicable not only to activated sludge bacteria separated by settling but also with other microbes selectively separated by other means, such as filtration, and can be demonstrated mathematically (Weissman and Benemann, 1979). Thus, in principle, and practice, it is possible to establish selective conditions which would result in the establishment of biofilm or biofloc processes.

Although biofilms actively adsorb heavy metals, both in natural systems (e.g., benthic algae, see next section) and in engineered reactors (activated sludge, rotating biological contactors, etc.), there has been relatively little study of such systems specifically designed for heavy metals removal (See Dunn and Bull, 1983, for an example). A recent comprehensive review of the literature on metals interactions and adsorption by biofilms (Lion et al., 1988) yields only a meager amount of information on how to use biofilms in heavy metals removal. The issue is also skirted in a recent book devoted to heavy metals in waste water treatment (Lester, 1987), in which much data is reviewed on metal adsorption by exopolymers, the nature and formation of such exopolymers, and their role in

the formation of biofilms and the removal of heavy metals during the waste treatment processes. However, little unification between these aspects is apparent. Thus, the use of controlled biofilms as a method of heavy metals removal or recovery is one that still needs to be developed.

7.4 Conclusions

Despite the emphasis on immobilization processes, whether natural or artificial, there may be a few situations where dispersed organisms may be more useful. Dispersed organisms would exhibit better kinetics of metal adsorption, as they would not be diffusion limited. Also, when using living organisms, particularly in those situations where growth is required, these may not be readily amenable to immobilization or biofilm formation. In those cases low cost separation technology must be applied. This has apparently not been specifically investigated for metals removal, except for the work by Smith (1988) and Smith et al. (1989) on the flotation of microalgae. Another alternative to immobilization could be to rely on the self-flocculating characteristics of many microbial cells, allowing maximal exposure to metals during a dispersed growth phase followed by an induction of flocculation, using a two stage reactor (e.g., Norberg and Ryding, 1984).

The difficulties of dealing with immobilized processes and with living microbes are multiplied when dealing with a combination of both in a practical system. However, dispersed growth, flocculant growth and biofilm reactors comprising living microbes are already widely in use in the waste treatment industry, at scales which would dwarf anything contemplated in bioaccumulation. Thus, applications of such technologies in this area does not require any breakthroughs in technology but can be extrapolated from existing systems.

In conclusion, the engineering problems inherent in such processes must, in many cases, still be resolved. For example, U is bound 14 times more efficiently by Asperigillus niger than by IRA-400 (a standard ion exchange resin), however the bioexchange material occupied 20 times the volume and had 1/5th the flow rate of the competing ion exchange material (Yakubu and Dudeney, 1986). Such disadvantages must be overcome by the applied R&D in the areas of biomass immobilization. It is likely that applied R&D in this area will continue to accelerate, given an impetus by the promising results obtained over the past decade, the application of bioexchangers in some commercial applications, and the burgeoning number of companies entering this field (anonymous, 1989).

8.0 APPLICATIONS OF MICROALGAE IN BIOREMOVAL

8.1 Introduction

Algae are known to accumulate metals up to 20% of their dry weight (Ahlf et al., 1980). They can accumulate heavy metals to concentrations several orders of magnitude higher than the surrounding medium (Becker, 1986) and are thus extremely well suited for the purification of heavy-metal contaminated waters. However, despite the extensive research in bioaccumulation processes, the actual applications in this area are presently limited. One area which has received a significant amount of attention from a practical perspective is the use of natural populations of microalgae, in "meander" or similar semi-engineered open systems for the removal of heavy metals from mine operation effluents.

Another already practical use of microalgae is the use of biosorbents for the recovery of gold from mining operations. Reportedly one operation using microalgae for gold recovery has been operating in South Africa at a Gencor Co. mine for several years, although no details have been reported (Grobellar, personal communication 1988). Gold recovery by microalgae has also been developed by Bio-Recovery Systems, Inc., of Las Cruces, New Mexico (Darnall et al., 1988b; Bedell, personal communication, 1989). Indeed, Bio-Recovery Systems, Inc. has commercialized microalgae based biosorbents for the removal of heavy metals from various types of industrial wastewaters.

8.2 Microalgae Bioremoval from Mine Effluents

The fact that microalgae accumulate metals has been known for many decades, and more recently, algal bioaccumulation has been studied in algae growing in, or as part of, the treatment systems of mine waste waters. For example, Wong (1979) found that marine algae growing in tailings of an iron ore mine concentrated Fe, Mn, Pb, and Zn on the cell wall. Jackson (1978) used a mixed culture of algae and bacteria for the anaerobic bioprecipitation of metal sulfides from a mine waste water. Trollope and Evans (1976) in a study of algae growing in a pond polluted by Zn smelting wastes found that green algae accumulate Cu, Fe, Zn, and Pb, with cyanobacteria also accumulating Ni. More recently, Kalin and Smith (1985) and Mann and Fyfe (1985) published brief abstracts on the use of Chara and Euglena, respectively, for the bioaccumulation of heavy metals from uranium mine waste waters, although detailed reports are lacking.

Probably the best known work in this category is that of Gale and Wixson (1979, see also Gale, 1986) who studied an artificial meander system discharging from a lead mine tailing lagoon. The meander stream was dominated by benthic attached filamentous green algae (Cladophora, Spirogyra, Hydrodictyon, etc.). These algae could grow even in areas of very high metal contamination, with one algal sample found growing in a zinc-rich leachate reported to contain this metal at 25% by weight (dry). Cladophora collected from the meander, when packed (11 g wet weight) into a column could efficiently remove lead (as the acetate) at a concentration of 100 ppm, with other cations (K, Mg, Ca, Na) being displaced by the metal - clearly indicating the ion exchange nature of the process. Field specimens contained much lower amounts of lead (about 1/10th those observed in the laboratory), but also contained Zn and some Cu. The ion exchange nature of that binding was also evidenced by the release of the metals by EDTA washing.

Although ion exchange was demonstrated to take place, this was only considered to be the third stage in the process of metal removal, with the first stage being a physical entrapment of suspended mineral particles, containing most of the heavy metals contained in the waste stream. This was followed by reactions at or near the algal cell surface resulting in the solubilization of the minerals and finally reassociation and redistribution of the available cations in a typical cation exchange process. The authors also quote earlier work which demonstrated that metal adsorption by peat was a cation exchange process and that sphagnum moss together with humic acids are responsible for the fixation of elements by peat bogs overlying mineral deposits.

Another field study of microalgae for heavy metals removal from a mining waste water was carried out by Brierley and Brierley (1981) and Brierley et al. (1980) who monitored a treatment process for U and Mo removal (about 0.8 ppm each) from a U mine water involving a series of three consecutive algae ponds, each 1 to 2 acres in size, for final settling and contaminant removal. These ponds contained mostly filamentous algae, including the macroscopic green alga Chara, the microscopic green algae Spirogyra and Rhizolconium, the blue-green alga Oscillatoria, and several minor microflora components. A major process for metals removal was the decomposition of the algae in the sediments, resulting in metal precipitation (accounting for the high levels of metals in the sediments.). However, the chemistry of U and Mo ions precipitation under anaerobic conditions is more complex than the Zn, Cd, and Cu sulfides which were precipitated from zinc mine waste water by Jackson (1978) using a similar algae-sediment-bacterial system. Although the sedimentation and decay of the algal biomass led to metals precipitation, there also was ongoing metals solubilization taking place.

These conclusions were supported with laboratory studies involving both long term algae decomposition studies and short term adsorption studies. The metal adsorption studies indicated that the algae could accumulate 2 g/kg dry weight of U. The major limitation to metals removal into the sediments was lack of sufficient algal biomass production to maintain a constantly anaerobic sediment environment. If all the U in the sediments were to come from settled algal biomass, they concluded that (based on 0.8 ppm and a 4.4 MGD flow) they would need to produce 2,500 tons of algae in these ponds, which is too high by a factor of at least 20 (for about 5 acres) even with optimal algae production, which was far from the case. However, as direct removal in the sediments could probably result in removing all of the metal, the role of the algae would have been simply to supply reductant, which would not have required that much productivity. Other investigators have reported on U binding by microalgae under better defined conditions, reporting both higher concentration factors and contents in the algae (e.g., Greene et al, 1986b). Those reports are reviewed after consideration of large scale algae culture systems in heavy metals removal.

8.3 Microalgae Bioremoval from Treatment Ponds

Microalgae are used extensively in municipal and industrial waste treatment ponds. These ponds, generally called oxidation ponds, are relatively large (usually a few acres), deep (4 to 6 feet) ponds that typically contain a mixed population of microalgae that develops naturally and provides the oxygen required for the aerobic breakdown of wastes.

Tarifeno-Silva et al. (1982) studied the concentrations of 15 heavy metals in successive stages of artificial food chains composed of algae, herbivorous crustaceans and carnivorous fish. These food chains were grown in secondarily treated chlorinated/dechlorinated domestic wastewaters. Bioaccumulation of heavy metals occurred mainly in the algae (CF's of 500-30,000). Later stages in the food chains showed successively lower accumulations (CF's of 30-6,000 for cladocerans and 4-1600 for fish).

One of the first studies of removal of heavy metals from wastewaters using microalgae was that of Filip et al. (1979) who studied the use of oxidation pond algae in the removal of Cd, Cu, and Cr from waste waters. They exposed the mixed cultures of algae obtained from such ponds to various levels and combinations of these metals and studied, under laboratory conditions, the growth and metal adsorption in semicontinuous cultures, extending over several weeks. Finally, they used intermittent sand filtration to remove the algae and demonstrate the effectiveness of the process in removing metals. They demonstrated relatively high degrees of Cd and Cu adsorption (from sub ppm and low ppm levels, respectively) and nearly total removal by intermittent sand filtration for these two metals (Cr was not removed well).

The positive evaluation of the prospects of such a process by Filip et al. (1979) were not shared by Becker (1983) who investigated concentration factors and growth rates by several algae exposed to actual municipal waste waters using a dialysis system. The wastewater was, however, very low in metals, maximally 13 ppb Pb and 2 ppb Cd - making extrapolations to a more contaminated situation difficult. Also, not enough information about conditions (algae concentrations, time of exposure, temperatures, pH, etc.) were provided. Concentration factors (ppb metal in algae dry weight/ppb metal in wastewater) were almost 8,000 for Pb and about 1,000 for Cd, in the better cases, in broad agreement with a few prior literature data points quoted by Becker (1983) (although much higher concentration factors can also be found, typically over 100,000 and even over 1 million). He then evaluated the requirement for a practical process and concluded that it would not be feasible for a number of reasons, including the long retention times required (several weeks) and the difficulties of separating the algae. The basis for the requirement for a long retention time was not apparent from the data, particularly as he showed that metal adsorption itself was essentially complete in less than 6 hours. Also, the mass balance seemed to be wrong: using his assumptions about concentration factors (5,000) and heavy metal concentration to be removed (1 ppm), sufficient algal biomass is produced in oxidation ponds (over 200 mg/l) to effectuate complete metal removal.

However, there are certainly significant problems with the concept of using oxidation ponds, or similar algal waste treatment systems in any heavy metal removal process. A major problem is the uncontrolled nature of the existing production system, where no control over algal species growing in the ponds is practiced (or possible) and where even algal growth fluctuates widely. A second, perhaps even more crucial, problem is the harvesting of the algal biomass - no cost effective process has been demonstrated.

Both of these limitations, could, however, be overcome by the use of "high rate ponds", which are shallow, raceway type, hydraulically mixed (often with paddle wheels) ponds. With such systems algal species are more controllable, suggesting that metal binding behavior, if not optimal is at least predictable. Also, by controlling algal species, very low cost algae harvesting processes, such as "bioflocculation" can be applied (Benemann et al., 1980, 1987). Alternative microalgae harvesting processes could also be considered. For example, in recent work, Smith, (1988), and colleagues (Smith et al., 1989), investigated floatation harvesting of microalgae specifically for heavy metals

removal. Other suitable processes include microstraining or chemical flocculation of the algae. Algal biomass harvested from such waste treatment systems could provide a cheap and low cost source of algal biomass for the removal of heavy metals from industrial wastewaters by bioaccumulation processes, which is discussed next.

8.4 Microalgae Bioremoval from Industrial Wastewaters

As with other microbes, microalgae exhibit both active and passive bioaccumulation processes, although few active transport systems have been studied. One recent exception is the work of Campbell and Smith (1986), who found that Ni^{2+} was accumulated by the blue-green alga *Anabaena cylindrica* via an energy (membrane potential) dependent process with high affinity (binding constant of less than one ppb) although the uptake rate was relatively slow. The slow uptake could be accounted for by the low requirements for this metal, used by only one or two enzymes not required for growth in the laboratory. Most importantly, the uptake was highly specific, being unaffected by other metals present, including the ionically similar Mg^{2+} and Co^{2+} . The first step in the process was a rapid (EDTA inhibited) surface binding of Ni, followed by a slower, sustained, energy dependent and specific intracellular uptake of the metal. However, the concentration factors observed were only about 2,500, independent of external concentration (within the limits tested) - which, based on thermodynamic arguments, would suggest that transport did not involve Ni-binding carriers but, rather an electrochemical equilibrium. An active transport process would be expected to accumulate Ni^{2+} until a certain internal concentration is reached, regardless of the external levels. This work clearly demonstrates the power of active uptake systems, both in terms of very high affinity and specificity for particular metals, as well as their potential limitations, in terms of rate of uptake and accumulation factors. However, these would be expected to be much higher with metals that are required in higher amounts or are more essential for the growth of the organism. They also would be more easily manipulated by genetic selection.

Probably the most extensive comparative study of concentration factors was carried out by Hassett et al. (1981). Using a microtiter plate assay with radioactive metals, eleven different algae were exposed to three different metals (Cd, Pb, Hg) at seven pH values (from 4 to 10) and concentration factors determined for 11 and 44 day old cultures after a 3 hr incubation in the dark. Uptake measured in a few lighted control samples were not statistically different. As may be expected, there were large variations in concentration factors for the different metals, algal species, pH levels, and metals, with few clear trends appearing. Mercury exhibited higher concentration factors (to a maximum of almost 100,000 in one case and over 10,000 in most cases) by more algae than cadmium or lead, with the latter only being concentrated by one species over the whole pH range tested. However, the most spectacular difference was between the 11 and 44 day old cultures, with the latter exhibiting concentration factors ten to over one hundred fold lower than the former. This was attributed to both the higher cell concentrations in the older cultures, and probably more importantly, the differences in cell wall composition of aged cultures, for which supposition there is considerable evidence in the literature. The reasons for this are unknown.

Microalgae have been shown to accumulate a large number of different metals, with concentration factors from negligible to over 1 million fold (Fisher, 1985; Horikoshi et al, 1979). An extensive literature has developed over the past decade on this phenomena (Sakaguchi et al., 1979, 1981; Crist et al, 1981; Nakjima et al., 1981; Khummongkol et al., 1982; Fisher et al., 1983, 1985; Trevors et al., 1985; Ferguson et al., 1989; Darnall et al., 1986a,b; Darnall et al., 1988a,b; Greene et al., 1986a,b; Greene and Bedell, 1989).

9.0 CONCLUSIONS

There appears to be great potential in the application of microbial bioaccumulators in the commercial treatment of industrial wastewaters. Microbial bioaccumulators have the potential of recovering metals while simultaneously improving water quality in other ways. They are competitive with established ion exchange and other treatment technologies (and the expected future improvements in these technologies), and thus the economics of the overall process are of importance.

The potential advantages of microalgal bioremoval processes are their greater specificity for particular metals, higher binding capacities, lower equilibrium metal concentrations, insensitivity to interfering ions, and lower cost. The challenge for the future is to combine these advantages into practical bioremoval processes that can be used to improve the environment.

10.0 REFERENCES

- Ahlf, W. "Recovery of Metals from Acid Waste Water by Cyanidium caldarium", Appl. Microbiol. Biotechnol. 28:512-513 (1988).
- Ahlf, W., U. Irmer, and A. Weber. "Accumulation of Lead by Fresh Water Algae Under Various Conditions", Z. Pflanzenphysiol 100:197-207 (1980).
- Aikin, R.M., and A.C.R. Dean, "Lead Accumulation by Pseudomonas fluorescens and by a Citrobacter sp.", Microbios Letters 9:55 (1979a).
- Aikin, R.M., A.C.R. Dean, "Lead Accumulation by Microorganisms", Microbios Letters 5:129-133 (1979b).
- Anonymous, "Metal-containing Wastes: Biological Treatment Can Recover Metals. 1998 U.S. Market to Reach \$510 Million" Biotech. Processing 11(1):4-5 (1989).
- Baldry, M.G.C., and A.C.R. Dean, "Environmental Change and Copper Uptake by Bacillus subtilis supsp. Niger and by Pseudomonas fluorescens", Biotech. Letters 3:142-147 (1981).
- Baldry, M.G.C. and A.C.R. Dean, "Copper Accumulation by Bacteria, Moulds, and Yeasts", Microbios 22:7-14 (1980).
- Becker, E.W., "Limitations of Heavy Metal Removal from Waste Water by Means of Algae", Water Res. 17:459-466 (1983).
- Bedell, G.W., and D.W. Darnall, "Immobilization of Non-Viable, Biosorbent, Algal Biomass for the Recovery of Metal Ions", in Voleksy, B. (ed.), Biosorbents and Biosorption Recovery of Heavy Metal Ions CRC Press, (1989 in press).
- Bencheikh-Lehovine, "Zinc Removal Using Peat Adsorption", Environmental Tech. Letters 10:101-107 (1989).
- Bender, J. and V. Ibeanusi, "Effects of Supplements on the Bioaccumulation of Lead in Anabaena spp.", Bull. Environ. Contam. Toxicol. 39:209-213 (1987).
- Benemann, J.R., D.M. Tillett, and J.C. Weissman, "Microalgae Biotechnology", Trends in Biotechnology 5:47-53 (1987).
- Benemann, J.R., B.L. Koopman, J.C. Weissman, D.M. Eisenberg, and R. P. Goebel. "Development of Microalgae Harvesting and High Rate Pond Technologies in California." In G. Shelef and C.J. Soeder (editors) Algae Biomass Production and Use. El Sevier/North Holland Biomedical Press, Amsterdam (1980).
- Benemann, J.R., "The Future of Microalgae Biotechnology" in Algal Biotechnology (R.C. Cresswell, T.A.V. Rees, and N. Shah, eds.) Longman, London (1989 in press).

- Benemann, J.R., G. Smith, P.J. Kostel, and C.E. McKenna, "Tungsten Incorporation into Azotobacter vinelandii Nitrogenase." FEBS Letter 29:219-221 (1973).
- Benemann, J.R., C.E. McKenna, R. F. Lie, T.G. Taylor and M.D. Kamen "The Vanadium Effect in Nitrogen Fixation by Azotobacter vinelandii" Biochem. Biophys. Acta, 264 25-30 (1972).
- Beveridge, T.J., "The Immobilization of Soluble Metals by Bacterial Walls", in Ehrlich, H.L., and D.S. Holmes (eds.), Workshop on Biotechnology for the Mining, Metal Refining Industries, Biotech Bioeng Symp.16 (1986).
- Beveridge, T.J., "Metal Ions and Bacteria", in Beveridge, T., and R. Doyle (eds.), Metal Ions and Bacteria, John Wiley & Sons, in press (1989).
- Beveridge, T.J., C.W. Forsberg, and R.J. Doyle, "Major Sites of Metal Binding in Bacillus licheniformis Walls", J. Bacteriol. 150:1438-1448 (1982).
- Beveridge, T.J., and R.G.E. Murray, "Uptake and Retention of Metals by Cell Walls of Bacillus subtilis", J. Bacteriol. 127: 1502-1518 (1980a).
- Beveridge, T.J., and R.G.E. Murray, "Sites of Metal Deposition in the Cell Wall of Bacillus subtilis", J. Bacteriol. 141:876-887 (1980b).
- Beveridge, T.J., and S.F. Koval, "Binding of Metals to Cell Envelopes of Escherichia coli", Appl. Env. Microbiol. 42:325-335 (1981).
- Beveridge, T.J., "The Bacterial Surface: General Considerations Towards Design and Function", Can. J. Microbiol. 34:363-372 (1988).
- Bisen, P.S., H.D. Shukla, A. Gupta, and S.N. Bagchi, "preliminary Characterization of an Novel Synechococcus Isolate Showing Mercury, Cadmium, and Lead Tolerance", Environmental Tech. Letters 8:427-432 (1987).
- Bowman, B.J., "Vanadate Uptake in Neurospora crassa Occurs via Phosphate Transport System II", Appl. Env. Microbiol. 153:286-291 (1983).
- Braun, V., "The Unusual Features of the Iron Transport Systems of Escherichia coli", Trends Biochem. (1985).
- Brierley, J.A., G.M. Goyak, and C.L. Brierley, "Considerations for Commercial Use of Natural Products for Metals Recovery", in Eccles, H., and S. Hunt, (eds.), Immobilization of Ions by Bio-Sorption, Ellis Horwood, Ltd. (1986).
- Brierley, J.A., C.L. Brierley, and G.M. Goyak, "AMT-Bioclaime Process for Treatment of Metalliferous Wastewater from Electroplating and other Industries", in McCready, R.G.L. (ed.), Proc. Second Ann. General Meet. of BIOMINET (1985).
- Brierley, J.A., C.L. Brierley, and K.T. Dreher, "Removal of Selected Inorganic Pollutants from Uranium Mine Waste Water by Biological Methods", in Brawner, C.O., (ed.), Uranium Mine Waste Disposal, (First Int. Conf., May 19-21 (1980).

- Brierley, C.L., and J.A. Brierley, Biological Processes for Concentrating Trace Elements from Uranium Mine Waters, New Mexico Water Resources Research Institute December (1981).
- Brown, M.J., and J.N. Lester, "Role of Bacterial Extracellular Polymers in Metal Uptake in Pure Bacterial Culture and Activated Sludge - VII. Effects of Mean Cell Retention Time", Water Res. 16:1549-1560 (1982).
- Butt, T.R., and D.J. Ecker, "Yeast Metallothionein and Applications in Biotechnology", Microbiol. Rev. 51:351-364 (1987).
- Campbell, P.M., and G.D. Smith, "Transport and Accumulation of Nickel Ions in the Cyanobacterium Anabaena cylindrica", Arch. Biochem. Biophys. 244:470-477 (1986).
- Charley, R.C., and A.T. Bull, "Bioaccumulation of Silver by a Multispecies Community of Bacteria", Arch. Microbiol. 123:239-244 (1979).
- Chaufer, B., S.A. Ouiminga, A. Deratani, and B. Seville, "Metal Ion Extraction by an Ultrafiltration Process with Modified Starch Derivatives", Eccles, H., and S. Hunt (eds.), Immobilization of Ions by Bio-Sorption, Horwood (1986).
- Chmielowski, J. and B. Klapcinska, "Bioaccumulation of Germanium by Pseudomonas putida in the Presence of two Selected Substrates", Appl. Env. Microbiol. 51:1099-1103 (1986).
- Cole, J.J., R.W. Howarth, S.S. Nolan, and R. Marino, "Sulfate Inhibition of Molybdate Assimilation by Planktonic Algae and Bacteria: Some Implications for the Aquatic Nitrogen Cycle", Biogeochemistry 2:179-196 (1986).
- Corpe, W.A., "Metal-Binding Properties of Surface Materials from Marine Bacteria", Dev. Ind. Microbiol. 26:249-255 (1974).
- Crist, R.H., and K. Oberholser, N. Shank, and M. Nguyen, "Nature of Bonding between Metallic Ions and Algal Cell Walls", Environ. Sci. Tech. 15:1212-1217 (1981).
- Darnall, D.W., and G.W. Bedell, "Heavy Metal Ion Recovery Using Biological Biosorbents", Presented at the Symposium Biotechnology in Minerals and Metal Processing, held in Las Vegas, Nevada, Society of Mining Engineers (1989).
- Darnall, D.W., B. Green, M. Hosea, R.A. McPherson, M. Henzl and M.D. Alexander, "Recovery of Heavy Metals by Immobilized Algae", in Kelly, D.P., and Norris, P.R. (eds.), Biohydrometallurgy 1987, Warwick, Science and Technology Letters, London (1988a).
- Darnall, D.W., B. Greene, and J. Gardea-Torresdey, "Gold Binding to Algae" in Kelly, D.P., and Norris, P.R. (eds.), Biohydrometallurgy 1987, Warwick, Science and Technology Letters, London pp. 487-498 (1988b).
- Darnall, D.M., B. Greene, M. Hosea, R.A. McPherson, M. Henzl, and M.D. Alexander, "Recovery of Heavy Metals by Immobilized Algae", in Thompson, R. (ed.), Trace Metal Removal from Aqueous Solution, The Industrial Div Royal Soc. (1986a).

- Darnall, D.W., B.Green, M.T. Henzl, J.M. Hosea, R.A. McPherson, J. Sneddon, and M.D. Alexander, "Selective Recovery of Gold and Other Metals Ions from an Algal Biomass", Env. Sci. Technol. 20:206-208 (1986b).
- de la Noue, J. and D. Proulx. "Tertiary Treatment of Urban Wastewaters by Chitosan-Immobilized Phormidium sp." In: Algal Biotechnology, T. Stadler, J. Mollion, M.-C. Verdus, Y. Karamanos, H. Morvan et d. Christiaen (Eds), Elsevier Applied Science Publ., Barking, Engl., 159-168 (1988).
- DeVoe, I.W., and B.E. Holbein, "Metal Absorption by Modified Chitins", in Thompson, R., (ed.), Trace Metal Removal from Aqueous Solution, The Industrial Division of the Royal Society of Chemistry, IV Series, Whitstable Kent, (1986).
- Dockins, W.S., G.J. Olson, G.A. McFeters, and S.C. Turbak, "Dissimilatory Bacterial Sulfate Reduction in Montana Groundwaters", Geomicrobiol. J. 2:83-89 (1980).
- Doyle, R.J., T.H. Matthews, and U.N. Streips, "Chemical Basis for Selectivity of Metal Ions by the Bacillus subtilis Cell Wall", J. Bacteriol. 143:471-480 (1980).
- Dugan, P.R., and H.M. Pickrum, "Removal of Mineral Ions from Water by Microbially Produced Polymers", Proc. 27th Purdue Ind. Waste Conf., pp.1019-1038 (1972).
- Dunn, G.M., and A.T. Bull, "Bioaccumulation of Copper by a Defined Community of Activated Sludge Bacteria", Europ. J. Appl. Microbiol. Biotechnol. 17:30-34 (1983).
- Eccles, H., and S. Hunt, (eds.), Immobilization of Ions by Bio-Sorption, Ellis Horwood, Ltd., Chichester (1986).
- Ehrlich, H.L., and D.S. Holmes, (eds.) Workshop on Biotechnology for the Mining, Metal Refining and Fossil Fuel Processing Industries, Biotech. Bioeng. Symp. No. 16 Rensselaer Polytechnic Inst., Troy NY (1986).
- Failla, M.L., "Zinc: Functions and Transport in Microorganisms", in Weinber, E.D., ed., Microorganisms and Minerals, Marcel Dekker, Inc., New York, pp 151-213 (1977).
- Failla, M.L., and E.D. Weinberg, "Cyclic Accumulation of Zinc by Candida utilis during Growth in Batch Culture", J. Gen. Microbiol. 99:85-97 (1977).
- Ferguson, C.R., M.R. Peterson, T.H. Jeffers, "Removal of Metal Contaminants from Waste Waters Using Biomass Immobilized in Polysulfone Beads", Preprint to be presented at Las Vegas, February 1989.
- Ferris, F.G., W.S. Fyfe, and T.J. Beveridge, "Metal Sorption by Bacterial Cell Walls", in McCready, R.G.L. (ed.), Proceeding of the Second Annual General Meeting of BIOMINET, October 24-25 (1985) Mississauga, Canada, CANMET (1986).
- Filip, D.S., T. Peters, V. D. Adams and E.J. Middlebrooks, "Residual Heavy Metal Removal by an Algae-Intermittent Sand Filtration System", Water Res. 13: 305-313 (1979).
- Fisher, N.S., "Accumulation of Metals by Marine Picoplankton", Marine Biology, 87: 137-142 (1985).

- Fisher, N.S., P. Bjerregaard, and S.W. Fowler, "Interactions of Marine Plankton with Transuranic Elements. 1. Biokinetics of Neptunium, Plutonium, Americium, and Californium in Phytoplankton", Limnol. Oceanogr., 28: 432-447 (1983).
- Fisher, N.S., and G.J. Jones, "Heavy Metals and Marine Phytoplankton: Correlation of Toxicity and Sulfhydryl- binding", J. Phycol. 17:108-111 (1981).
- Foster, T.J., H. Nakahara, A.A. Weiss, and S. Silver, "Transposon A-Generated Mutations in the Mercuric Resistance Genes of Plasmid R100-1", J. Bacteriol. 140:167-181 (1979).
- Friedman, B.A., and P. R. Dugan, "Concentration and Accumulation of Metallic Ions by the Bacterium Zoogloea", Dev. Ind. Microbiol. 2:381-388 (1967).
- Fris N. Myers-Keigh P., Biosorption of uranium and lead by Streptomyces longwoodensis. Biotechnol. Bioeng. 28:21 (1986).
- Furlong, C.E., J.A. Sundstrom, E.B. Weiler, P.W. Chung, J. Yin, and H.W. Blanch., "Exploitation of the High Affinity and Specificity of Proteins ...", in Ommen, G.S., (ed.), Environmental Biotech., Plenum, pp 281-293 (1988).
- Gadd, G.M., "Fungal Responses towards Heavy Metals", in Herbert, R.A., and G.A. Codd (eds.), Microbes in Extreme Environments, Academic Press for the Soc. General Microbiol., pp 83-110 (1986a).
- Gadd, G.M., "The Uptake of Heavy Metals by Fungi and Yeasts: The Chemistry and Physiology of the Process and Applications for Biotechnology", in Eccles, H., and S. Hunt (eds.), Immobilization of Ions by Bio-Sorption, Ellis (1986b).
- Gadd, G.M., C. White, and L. deRome, "Heavy Metal and Radionuclide Uptake by Fungi and Yeasts" in Norris P.R., and D.P. Kelly, Biohydrometallurgy, Proc. of the Inter. Symp. Warwick. Science and Tech. Lett., Kew, pp 421-435 (1988).
- Gadd, G.M., "Accumulation of Metals by Microorganisms and Algae", in K.-J. Rhem (ed.), Biotechnology Handbook, Vol. 6B. Special Microbial Processes, VCH Verlagsgesellschaft: Weinheim, in press (1989).
- Gale, N.L., "The Role of Algae and Other Microorganisms in Metal Detoxification and Environmental Clean-Up", in Ehrlich, H.L., and D.S. Holmes (eds.), Workshop on Biotechnology for the Mining, Metal Refining and Fossil Fuel Processing (1986).
- Gale, N.L., and B.G. Wixson, "Removal of Heavy Metals from Industrial Effluents by Algae" Dev. Ind. Microbiol. 20:259-273 (1979).
- Galun, M., P. Keller, H. Feldstein, E. Galun, S. Siegel, and B. Siegel, "Recovery of Uranium (VI) from Solution using Fungi. II. Release from Uranium-loaded Penicillium biomass", Water, Air and Soil Pollution 20:227-285 (1983).
- Galun, M., S.M. Sielgel, M.I. Cannon, B.Z. Siegel, and E. Galun, "Ultrastructural Localization of Uranium Biosorption in Penicillium digitatum by Stem X-Ray Microanalysis", Environmental Pollution 43:209-218 (1987a).

- Galun, M., E. Galun, B.Z. Siegel, P. Keller, H. Lehr, and S.M. Siegel, "Removal of Metal Ions from Aqueous Solutions by Penicillium Biomass: Kinetic and Uptake Parameters", Water, Air, Soil Pollution 33:359-371 (1987b).
- Galun, M., P. Keller, D. Malki, H. Feldstein, E. Galun, S.M. Siegel, and B.Z. Siegel, "Removal of Uranium(VI) from Solution by Fungal Biomass and Fungal Wall-related Biopolymers", Science 219:285-287 (1982).
- Garcia-Toledo, A., H. Babich, and G. Stotzky, "Training of Rhizopus stolonifer and Cunninghamella blakesleeana to Copper: Cotolerance to Cadmium, Cobalt, Nickel and Lead", Can. J. Microbiol. 31:485-492 (1985).
- Gekeler, W., E. Grill, E. L. Winnacker, and M. H. Zenk. "Algae Sequester Heavy Metals via Synthesis of Phytochelatin Complexes", Arch. Microbiol. 150:197-202 (1988).
- Ghiorse, W.C., "Biology of Iron- and Manganese-depositing Bacteria", Ann. Rev. Microbiol. 38: 515-550 (1984).
- Gingrich, D. J., D. N. Weber, C. F. Shaw, J. S. Garvey, and D. H. Petering, "Characterization of a highly negative and labile binding protein induced in Euglena gracilis by cadmium". Environ. Health Perspect. 65:77-85 (1986).
- Glick, B.R., N. Menhart, N.W. Soong, W.H. Farmer, T. Viswanatha, and J.J. Pasternak, "Isolation and Partial Characterization of Siderophore Mutants of Azotobacter vinelandii" 17:343-346 (1988).
- Greene, B., M. Hosea, R. McPherson, M. Henzl, M.D. Alexander, and D.W. Darnall, "Interaction of Gold(I) and Gold (III) Complexes with Algal Biomass", Environ. Sci. Tech. 20:627-632 (1986a).
- Greene, B., M.T. Henzl, J. M. Hosea, and D.W. Darnall, "Elimination of Bicarbonate Interference in the Binding of U(VI) in Mill-Waters to Freeze dried Chlorella vulgaris", Biotech. Bioeng. 28:764-767 (1986b).
- Greene, B., R. McPherson, and D. Darnall, "Algal Sorbents for Selective Metal Ion Recovery", in J.W. Patterson, and R. Passino, (eds.). Metals Speciation, Separation, and Recovery Lewis Publishers, Inc., (Proc. Int. Symp.) (1987).
- Greene, B., and D.W. Darnall, "Temperature Dependence of Metal Ion Sorption by Spirulina", Biorecovery 1:27-41 (1988).
- Greene, B., and G. W. Bedell, "Algal Gels or Immobilized Algae for Metal Recovery", An Introduction to Applied Phycology., I. Akatsuta, ed., SPB Academic Publishing ((1989) In press.
- Grill, E., E-L Winnacker, and M. H. Zenk. "Phytochelatin: The Principal Heavy Metal-Complexing Peptides of Higher Plants". Science 230:574-576 (1985).

- Grill, E., Winnacker E-L, and M. H. Zenk. "Phytochelatin, a class of heavy metal-binding peptides from plants, are functionally analogous to metallothioneins". Proc Natl Acad Sci USA, 84:439-443 (1987).
- Hancock, I.C., "The Use of Gram-positive Bacteria for the Removal of Metals from Aqueous Solution", in Thompson, R. (ed.), Trace Metal Removal from Aqueous Solution, The Industrial Div. the Royal Society of Chemistry, IV Series (1986).
- Hart, B. A., P. E. Bertram. "A cadmium-binding protein in a cadmium tolerant strain of Chlorella pyrenoidosa", Envir. Exp. Bot. 20:175-180 (1980).
- Hart, B.A., P.E. Bertram, and B.D. Scaife, "Cadmium Transport by Chlorella pyrenoidosa", Environ. Res. 18:327 (1979).
- Hassett, J.M, J.C. Jennett, and J.E. Smith, "Microplate Technique for Determining Accumulation of Metals of Algae", Appl. Environm. Microbiol. 41:1097-1106 (1981).
- Higham, D.P., P.J. Sadler, and M.D. Scawen, "Cadmium Resistance in Pseudomonas putida: Growth and Uptake of Cadmium", J. Gen. Microbiol. 131:2539-2544 (1985).
- Holmes, D.S., "Biotechnology in the Mining and Metal Processing Industries: Challenges and Opportunities", Mineral Metallurgical Processing 5:49-56 (1988).
- Horikoshi, T., A. Nakajima, and T. Sakaguchi, "Uptake and Uranium by Various Cell Fractions of Chlorella regularis", Radioisotopes 28:9-12 (1979).
- Horikoshi, T., A. Nakajima, and T. Sakaguchi, "Studies on the Accumulation of Heavy Metal Elements in Biological Systems. XIX. Accumulation of Uranium by Microorganisms", European J. App. Microbiol Biotech. 12:90-96 (1981).
- Hosea, M., B. Greene, R. McPherson, M. Henzl, M. D. Alexander, and D.W. Darnall, "Accumulation of Elemental Gold on the Alga Chlorella vulgaris", Inorganica Chimica Acta, 123:161-165 (1986).
- Hoyle, B., and T.J. Beveridge, "Binding of Metallic Ions to the Outer Membrane of Escherichia coli", App Env. Microbiol. 46:749-752 (1983).
- Hunt, S., "Diversity of Biopolymer Structure and its Potential for Ion Binding Applications" in Eccles, H., and S. Hunt (eds.), Immobilization of Ions by Bio-Sorption, Ellis Horwood, Ltd., Chichester (1986).
- Hyer, M. and W.J. Page, " Zn^{2+} Increases Siderophore Production in Azotobacter vinelandii", App Environ. Microbiol. 54:2625-2631 (1988).
- Ilyaletinov, A.N., "Environmental Protection and Use of Microorganisms for Industrial Waste Treatment", in Karavaiko, G.I., and S.N. Groudev (eds.), Biotechnology of Metals, Moscow, pp 344-351 (1985).
- Imber, B.E., M.G. Robinson, A.M. Ortega, and J.D. Burton, "Complexation of Zinc exudates from Skeletonema costatum grown in Culture", Marine Chemistry 16:131 (1985).

- Jackson, T.A., "The Biogeochemistry of Heavy Metals in Polluted Lakes and Astreams at Flin Flon Canada and a Proposed Method for Limiting Heavy-Metal Pollution of Natural Waters" Environmental Geology 2:173-189 (1978).
- Jardim, W.F. and H.W. Pearson, "Copper Toxicity to Cyanobacteria and its Dependence on Extracellular Ligand Concentration and Degradation", Microb. Ecol. 11:139-148 (1985).
- Jennett, J.C., J.E. Smith, and J.M. Hassett, "Factors Influencing Metal Accumulation by Algae", NITS PB83149377 (1982).
- Joho, M., M. Imai, and T. Murayama, "Different Distribution of Cd+2 between Cd-sensitive and Cd-resistant Strains of Saccharomyces Cervisiae", J. Gen. Microbiol. 131:53-56 (1985).
- Kagi, J.H.R., and H. J. Hapke, "Biochemical Interactions of Mercury Cadmium, and Lead", in Nriagu, J.O. (ed.), Changing Metal Cycles and Human Health, Dahlem Konferenzen 1984. Springer Verlag, Berlin, pp 237-250 (1984).
- Kalin, M. and M.P. Smith, "Biological Polishing Agents for Mill Wastewater An Example", in McCready, R.G.L. (ed.), Proceeding of the Second Annual General Meeting of BIOMINET, October 24-25 (1985).
- Kaplan, D., D.Christiaen, and S. Arad, "Binding of Heavy Metals by Algal Polysaccharides", in Stadler, T., J. Mollion, M. C. Verdus, Y. Karamanos, H. Morvan, and D. Christiaen (eds.) Algal Biotechnology, Proc.4th Int.Meet. SAA,e (1987a).
- Kaplan, D., D. Christianen, and S.(M.) Arad, "Chelating Properties of Extracellular Polysaccharides from Chlorella spp.", Appl. Environm. Microbiol. 53:2953-2956 (1987b).
- Karavaiko, G.I. and S.N. Groudev, Eds. Biogeotechnology of Metals, United Nations Environment Program (1985).
- Khummonghol, D., G.S. Canterford, and C. Fryer, "Accumulation of Heavy Metals in Unicellular Algae", Biotech. Bioeng. 24:2643-2660 (1982).
- Kuyucak, n., and B. Volesky, "Recovery of Cobalt by a New Biosorbent", CIM Bulletin, pp 95-99 (1988a).
- Kuyucak, N., and B. Volesky, "Biosorbents for Recovery of Metals from Industrial Solutions", Biotech. Lett. 10:137-142 (1988b).
- Kuyucak, N., and B. Volesky, "New Algal Biosorbent for Gold Recovery Process", in Norris, P.R., and D.P. Kelly (eds.), BioHydroMetallurgy, Warwick, pp 453-464 (1988c).
- Kuyucak, N., and B. Volesky, "Desorption of Cobalt-Laden Algal Biosorbent", Biotech. Bioeng. 33: 815-822 (1989a).
- Kuyucak, N., and B. Volesky, "The Mechanism of Cobalt Biosorption", Biotech. Bioeng. 33: 823-831 (1989b).

- Kuyucak, N., and B. Volesky, "Accumulation of Cobalt by Marine Alga", Biotech. Bioeng. 33: 809-814 (1989c).
- Lakshmanan, V.I., J. Christison, R. A. Knapp, J.M. Scharer, and V. Sanmugasunderam, "A Review of Bioadsorption Techniques to Recover Heavy Metals from Mineral-Processing Streams", in McCready, R.G.L. (ed.), Proc. Second Annual Meeting Biominet (1986).
- Lawrence, R.W., R.M.R. Brannion, and H.G. Ebner, Eds. Fundamental and Applied Biohydrometallurgy, Elsevier Pub. Co. (1986).
- Leeper, S.A., J.H. Wolfram, and T.E. Ward, "Low Molecular Weight, Cobalt-binding Proteins from Saccharomyces cerevisiae and Bacillus subtilis", in McCready, R.G.L. (ed.), Third Meeting of BIOMINET (1985).
- Leeper, S.A., and J.H. Wolfram, "Biologically Assisted Minerals Processing: Metallothionein Status Report, February 1985", Internal Technical Report SE-S-85-001, EE&G (Idaho National Energy Laboratory) (1985).
- Les, A., and R.W. Walker, "Toxicity and Binding of Copper, Zinc, and Cadmium by the Bluegreen Alga Chroococcus parvulus", Water, Air, Soil Poll. 23:129-139 (1984).
- Lester, J.N., "Biological Treatment", in J.N. Lester, ed. Heavy Metals in Wastewater and Sludge Treatment Processes. Vol. II, CRC Press, Boca Raton, Florida, pp, 15-40 (1987).
- Lion, L.W., M.L. Shuler, K.M. Hsieh, and W.C. Ghiorse, "Trace Metal Interactions with Microbial Biofilms in Natural and Engineered Systems", CRC Critical Reviews in Environmental Control 17:273-306 (1988).
- Macaskie, L.E., and A.C.R. Dean, "Microbial Metabolism, Desolubilization, and Deposition of Heavy Metals: Metal Uptake by Immobilized Cells and Application". in Mizrahi, A. (ed.), Adv. Biotech Processes 12:, in press (1989).
- Macaskie, L.E., and A.C.R. Dean, "Cadmium Accumulation by Micro-organisms", Environmental Tech. Let. 3:49-56 (1982).
- Macaskie, L.E., and A.C.R. Dean, "Cadmium Accumulation by Citrobacter sp.", J. Gen. Microbiol. 130:53-62 (1984).
- Macaskie, L. E., and A.C.R. Dean, "Uranium Accumulation by Immobilized Cells of a Citrobacter sp.", Biotech. Letters 7:457-460 (1985).
- Macaskie, L. E., A.C.R. Dean, "Use of Immobilized Biofilm of (Citrobacter sp. for the Removal of lead from aqueous flows". Enzyme Microbiol. Technol. 9:2 (1987).
- Maeda, S., S. Nakashima, and T. Takeshita. "Bioaccumulation of Arsenic by Freshwater Algae and the Application to the Removal of Inorganic Arsenic from an Aqueous Phase. Part 11. Chlorella vulgaris Isolated from Arsenic-Polluted Environment". Separation Sci. Technol. 20:153-161 (1985).
- Mang, S. and H. W. Trombala, "Uptake of Cadmium by Chlorella fusca Zeitschrift Pflanzenphysiol. 190:293 (1978).

- Mann, H. and W.S. Fyfe, "Retardation of Heavy Metal Dispersion in Tailings Seepage Waters from Elliot Lake, Via Metal Uptake by Algae", in McCready, R.G.L. (ed.), Proc. of the 2nd Ann. General Meeting of BIOMINET, October 24-25 (1985).
- McCready, R.G.L., and V.I. Lakshmanan, "Review of Bioadsorption Research to Recover Uranium from Leach Solutions in Canada", in Eccles, H., and S. Hunt (eds.), Immobilization of Ions by Bio-Sorption, Ellis Horwood, Ltd., (1986).
- McKay, G., H.S. Blair, and A. Findon, "Sorption of Metal Ions by Chitosan" in Eccles, H., and S. Hunt, (eds.), Immobilization of Ions by Bio-Sorption, Ellis Horwood, Ltd., Chichester, pp 59-70 (1986).
- McKenna, C.E., J.R. Benemann and T.G. Traylor. "A Vanadium-Containing Nitrogenase Preparation - Implications for the Role of Molybdenum in Nitrogen Fixation." Biochem. Biophys. Res. Commun. 41:1505-1508 (1970).
- McKnight, D.M., and F.M. Morel, "Release of Weak and Strong Copper Complexing Agents by Algae", Limnol. Oceanogr. 25:62-71 (1979).
- Mierle, G., and P.M. Stokes, "Heavy Metal Tolerance and Metal Accumulation by Planktonic Algae", pp 113-122 (1976).
- Minney, S.F., and A.V. Quirk, "Growth and Adaptation of Saccharomyces cerevisiae at Different Cadmium Concentrations", Microbios 42:37-44 (1985).
- Misra, T.K., S. Silver, H.L.T. Mobley, and B.P. Rosen, "Molecular Genetics and Biochemistry of Heavy Metal Resistance in Bacteria", in Tashjian, A.H.Jr. (ed.), Molecular and Cellular Approaches to Mechanisms of Toxicity, (1984).
- Mobley, H.L., and A.O. Summers, "Plasmid-encoded Ion Transport Systems", in Rosen, B.P., and S. Silver (eds.), Ion Transport in Prokaryotes, Academic Press, San Diego, pp 305-326 (1987).
- Morper, M., "Anaerobic Sludge - A Powerful and Low Cost Sorbent for Heavy Metals" in Eccles, H., and S. Hunt (eds.), Immobilization of Ions by Bio-Sorption, Ellis Horwood, Ltd., Chichester, pp 91-104 (1986).
- Murphy, R.J., and J.F. Levy, "Copper Precipitation on Fungal Cell Surfaces", Trans. Br. Mycol. Soc. 81:165-168 (1983).
- Murray, A.D. and D.K. Kidby, "Sub-Cellular Location of Mercury in Yeast Grown in the Presence of Mercuric Chloride", J. Gen. Microbiol. 86:66-74 (1975).
- Muzarelli, R.A.A., F. Tanfani, M. Emanuelli, and L. Bolognini, "Aspartate, Glucan, Glycine Glucan, and Serine Glucan for the Removal of Cobalt and Copper from Solutions and Brines", Biotech. Bioeng. 27:1115-1121 (1985).
- Muzarelli, R.A.A., F. Bregani, and F. Sigon, "Chelating Capacities of Amino Acid Glucans and Sugar Acid Glucans Derived from Chitosan", in Eccles, H., and S. Hunt (eds.), Immobilization of Ions by Bio-Sorption, Ellis Horwood (1986).

- Nagano, T., M. Miwa, S. Suketa, S. Okada. "Isolation, physicochemical properties, and amino acid composition of a cadmium-binding protein from cadmium-treated Chlorella ellipsoidea". J. Inorg. Biochem. 21:61-71 (1984).
- Nakajima, A., T. Horikoshi, and T. Sakaguchi, "Studies on the Accumulation of Heavy Metal Elements in Biological Systems. XVII. Selective Accumulation of Heavy Metal Ions by Chlorella vulgaris", Eur. J. App. Microb. Biotech. 12:76 (1981).
- Nakajima, A., and T. Sakaguchi, "Selective Accumulation of Heavy Metals by Microorganisms", Appl Microbiol. Biotech. 24:59-64 (1986).
- Nakajima, A., T. Horikoshi, and T. Sakaguchi, "Recovery of Uranium by Immobilized Microorganism", Eur. J. App Microbiol. Biotech. 16:88-91 (1982).
- Norberg, A.B., and S. Ryding, "Development of a Continuous Process for Metal Accumulation by Zoogloea ramigera", Biotech. Bioeng. 26:265-268 (1984).
- Norberg, A.B., and H. Persson, "Accumulation of Heavy Metal Ions by Zoogloea ramigera", Biotech. Bioeng. 26:239 (1984).
- Norris, P.R., and D.P. Kelly, "Accumulation of Cadmium and Cobalt by Saccharomyces cerevisiae", J. Gen. Microbiol. 99:317-324 (1977).
- Norris, P.R., and D.P. Kelly, eds. Biohydrometallurgy. Proc. Internatl. Symp. Warwick, 1987 Science and Technology Letters (1987).
- Ohtake, H., C. Cervantes, and S. Silver, "Decreased Chromate uptake in Pseudomonas fluorescens Carrying a Chromate Resistance Plasmid", J. Bacteriol. 169:3853-3856 (1987).
- Olson, G.J., F.E. Brinckman, T.K. Trout, and D. Johnsonbaugh, "Bioaccumulation of Metals from Solution: New Technology for Recovery, Recycling and Processing", Int. Symp. Metallurgical Processes for the Year 2000... Las Vegas (1989).
- Olson, G.J., "Microbial Intervention in Trace Element-containing Industrial Process Streams and Waste Products", in Bernhard, M., F.E. Brinckman, and P.J. Sadler (eds.), The Importance of Chemical "Speciation" in Environment (1986).
- Olson, G.J., and F.E. Brinckman, "Review and Discussion -Algal Sorbents for Selective Metal Ion Recovery", in J.W. Patterson, and R. Passino, Metals Speciation, Separation, and Recovery Lewis Publishers, Inc. (1987).
- Parker, W. and M. O'Brien, J. Bacteriol. "Cobalt Binding by Saccharomyces cerevisiae" 68:167 (1954).
- Perkin, M.J., and I.S. Ross, "Mn and Mg Transport", Microbios Lett. 29:115 (1986).
- Robson, R. L., "Nitrogen Fixation in Strains of Azotobacter chroococcum Bearing Deletions or a Cluster of Genes Coding for Nitrogenase", Arch. Microbiol. 146:74-79 (1986).

- Romeyer, F.M., F.A. Jacobs, L. Masson, Z. Hanna, and R. Brousseau, "Bioaccumulation of Heavy Metals in Escherichia coli Expressing an Inducible Synthetic Human Metallothionein Gene", J. Biotech. 8: 207-220 (1988).
- Ross, I.S., "Uptake of Heavy Metals by Microorganisms", International Industrial Biotechnology, October/November, 184-188 (1987a).
- Ross, I.S., "The Use of Microorganisms for the Removal and Recovery of Heavy Metals from Aqueous Effluents", International Industrial Biotechnology, June/July, 330-333 (1987b).
- Rudd, T., "Sludge Treatment", in J.N. Lester, ed. Heavy Metals in Wastewater and Sludge Treatment Processes. Vol. II, CRC Press, Boca Raton, Florida, pp 41-68 (1987).
- Sakaguchi, H., T. Tsuji, A. Nakajima, and T. Horikoshi, "Accumulation of Cadmium by Green Microalgae", Eur. J. Appl. Microbiol. Biotech. 8:207-215 (1979).
- Sakaguchi, T., and A. Nakajima, "Recovery of Uranium from Seawater by Immobilized Tannin", Separation Science and Technology, 22: 1609-1623 (1986).
- Sakaguchi, T., Nakajima, A., T. Horikoshi, "Studies on the Accumulation of Heavy Metal Elements in Biological Systems. XVIII. Accumulation of Molybdenum by Green Microalgae", Eur. J. Appl. Microbiol. Biotech. 12:84-89 (1981).
- Schmid, J. and G. Auling, "Manganese Transport in Brevibacterium ammoniagenes", J. Bacteriol. 169:3385-3387 (1987).
- Schuett, C., "Plasmids in the Bacterial Assemblage of a Dystrophic Lake: Evidence for Plasmid-encoded Nickel Resistance", Microbial Ecol. 17:49-62 (1989).
- Scott, J.A., S.J. Palmer, and J. Ingham, "Microbial Metal Adsorption Enhancement by Naturally Excreted Polysaccharide Coatings", in Eccles, H., and S. Hunt (eds.), Immobilization of Ions by Bio-Sorption, Ellis Horwood, Ltd.(1986).
- Scott, J.A., and S. J. Palmer, "Cadmium Bio-Sorption by Bacterial Exopolysaccharide" Biotechn. Lett. 10:21-24 (1988).
- Shehata, F.H.A., and B.A. Whitton, "Zinc Tolerance in Strains of the Blue-green Alga Anacystis nidulans", Br. Phycol. 17:5-12 (1982).
- Shumate, S.E., G.W. Strandberg, and J.R. Parrott, "Biological Removal of Metal Ions from Aqueous Process Streams", Biotech. Bioeng. Symp. No. 8, pp 13-20 (1978).
- Shumate, S.E., G.W. Strandberg, D.A. McWhirter, J.R. Parrott, G.M. Bogacki, and B.R. Locke, "Separation of Heavy Metals from Aqueous Solutions Using 'Biosorbents' - Development of Contacting Devices for U", Biot. Bioeng. Symp. 10, 27 (1980).
- Shumate, S.E., and G.W. Strandberg, "Accumulation of Metals by Microbial Cells" in Moo-Young, M. (ed.), Comprehensive Biotechnology. Vol. 4 pp. 235-248, Pergamon Press, Oxford (1985).

- Siegel, S.M., B.Z. Siegel, C. Barghigiani, K. Aratani, P. Penny, and D. Penny, "A Contribution to the Environmental Biology of Mercury Accumulation in Plants", Water, Air, Soil Pollution 33:65-72 (1987).
- Silver, S., "Bacterial Transformations of and Resistance to Heavy Metals", in Igrolic, K.J., and A.E. Martell (eds.), Environmental Inorganic Chemistry, VCH Publishers Inc., Deerfield Beach, Florida, pp 513-539 (1985).
- Smith, R.W., Z. Yang, and R.A. Wharton Jr., "Flotation of Microorganisms - Implications in the Removal of Metal Ions from Aquous Streams", in Metallurgical Processes for the Year 2000 and Beyond, Symp. in press (1989).
- Smith, R.W., "Flotation of Algae, Bacteria and other Microorganims", Mineral Processing and Technology Review, 1: in press (1988).
- Sterrit, R.M., and J.N. Lester, "Heavy Metal Immobilization by Bacterial Extracellular Polymers", in Eccles, H., and S. Hunt (eds.), Immobilization of Ions by Bio-Sorption, Ellis Horwood, Ltd., Chichester (1986).
- Strandberg, G.W., S.E. Shumate, and J.R. Parrott, Jr., "Microbial Cells as Biosorbents for Heavy Metals: Accumulation of Uranium by Saccharomyces cerevisiae and Pseudomonas aeruginosa", Appl. Env. Microbiol. 41:237-245 (1981).
- Streat, M., and D. Naden, Ion Exchange and Sorption Processes in Hydrometallurgy, Soc. Chem. Ind., J. Wiley & Sons (1987).
- Streat, M., "Ion Exchange in Uranium Extraction", in Streat, M., and D. Naden (eds.), Ion Exchange and Sorption Processes in Hydrometallurgy, J. Wiley & Sons pp 2-53 (1987).
- Strong, J.R.P., J.C. Madgwick, and B.J. Ralph, "Metal Binding Polysaccharide from the Alga, Klebsormidium Fluitans", Biotech. Letters 4:239-242 (1982).
- Subramanyam, C., G. Venkateswerlu, and S.L.N. Rao, "Cell Wall Composition of Nuerospora crass Under Condition of Copper Toxicity" Appl. Env. Microbiol. 46:585-590 (1983).
- Summers, A.O., "Bacterial Resistance to Toxic Elements", Trends in Biotech. 3:122-125 (1985).
- Surowitz, K.G., J.A. Titus, and R. M. Pfister, "Effects of Cadmium Accumulation on growth and Respiration of a Cadmium Sensitive Strain of Bacillus subtilis and a Selected Cadmium Resistant Mutant", Arch. Micro. 140:107-112 (1984).
- Tarifeno-Silva, E., L. Y. Kawasaki, D. P. Yu, M. S. Gordon, and D. J. Chapman. "Aquacultural Approaches to Recycling of Dissolved Nutrients in Secondary Treater Domestic Wastewaters - III. Uptake of Dissolved Heavy Metals by Artificial Food Chains", Wat. Res. 16:59-65 (1982).
- Tobin, J.M., D.G. Cooper, and R.J. Neufeld, "Uptake of Metal Ions by Rhizopus arrhizus Biomass", Appl. Env. Micro. 47:819-822 (1984).
- Tornabene, T.G., and H.W. Edwards, "Microbial Uptake of Lead", Science 176:1334-1335 (1972).

- Townsley, C.C., and I.S. Ross, "Copper Uptake by Penicillium spinulosum", Microbios, 44: 125-132 (1985).
- Treen-Sears, M.E., B. Volesky, and R.J. Neufeld, "Ion Exchange/Complexation of the Uranyl Ion by Rhizopus Biosorbent, Biotechn. Bioeng. 26:1323-1329 (1984a).
- Treen-Sears, M.E., S.M. Martin, and B. Volesky, "Propagation of Rhizopus javanicus Biosorbent", Appl. Environ. Microbiol. 48:137-141 (1984b).
- Treen-Sears, M.E., S.M. Martin, and B. Volesky, "Control of Rhizopus Biosorbents' Quality during Propagation", in Lawrence, R.W., R.M.R. Branion, and H.G. Ebner (eds.), Fundamental and Applied Biohydrometallurgy, Elsevier, (1986).
- Trevors, J.T., K.M. Oddie, and B.H. Belliveau, "Metal Resistance in Bacteria", FEMS Microbiol. Rev. 32: 39-54 (1985).
- Trollope, D. R. and B. Evans Environ. Pollut. "Concentration of Copper, Iron, Nickel, and Zinc in Freshwater Algal Blooms." 11: 109-116 (1976).
- Tsezos, M., S.H. Noh, and M.H.I. Baird, "A Batch Reactor Mass Transfer Kinetic Model for Immobilized Biomass Biosorption", Biotech. Bioeng. 32:545-533 (1988).
- Tsezos, M., M.H.I. Baird and L.W. Shemilt, "The Use of Immobilized Biomass to Remove and Recover Radium from Elliot Lake Uranium Tailing Streams", Hydrometallurgy 17:357-368 (1987).
- Tsezos, M., and D.M. Keller, "Adsorption of Radium-226 by Biological Origin Absorbents", Biotech. Bioeng. 25:201-215 (1983).
- Tsezos, M., "The Role of Chitin in Uranium Adsorption by R. arrhizus", Biotech. Bioeng. 25: 2025-2040 (1983).
- Tsezos, M. and B. Volesky, "The Mechanism of Thorium Biosorption by Rhizopus arrhizus", Biotech. Bioeng. 24:955-969 (1982).
- Tsezos, M. and B. Volesky, "Biosorption of Uranium and Thorium" Biotech. Bioeng., 23: 583-604 (1981).
- Tsezos, M., "Recovery of Uranium from Biological Adsorbents -Desorption Equilibrium", Biotech. Bioeng., 26: 973-981 (1984).
- Tsezos, M., "Adsorption by Microbial Biomass as a Process for Removal of Ions from Process or Waste Solutions", in Eccles, H., and S. Hunt (eds.), Immobilization of Ions by Biosorption, Ellis Horwood, Ltd., Chichester (1986), pp.
- Tyneka, Z., Z. Gos, and J. Zajac, "Reduced Cadmium Transport Determined By a Resistant Plasmid in Staphylococcus aureus" J. Bacteriol., 147:305-312 (1981).
- Volesky, B., Editor, Biosorption of Heavy Metals, CRC Press Inc., Boca Raton, Florida (1990)

- Volesky, B., "Biosorbents for Metal Recovery" Trends Biotechnol. 5:96-101 (1987).
- Volesky, B. "Biosorbent Materials" in Ehrlich, H.L., and D.S. Holmes (eds.), Workshop on Biotechnology for the Mining, Metal Refining and Fossil Fuel Processing Industries, Biotechnology and Bioengineering Symposium No. 16, (1986).
- Wainwright, M., I. Singleton, and R.G.J. Edyvean, "Use of Fungal Mycelium to Adsorb Particulates from Solution", in Norris, P.R., and D.P. Kelly (eds.), BioHydroMetallurgy, Warwick, pp 499-501 (1988).
- Watkins, J.W., R.C. Elder, B.Green, and D.W. Darnall, "Determination of Gold Binding in al Algal Biomass Using EXAFS and XANES Spectroscopies", Inorganic Chem. 26:1147 1151 (1987).
- Weissman, J.C. and J.R. Benemann. "Biomass Recycling and Species Competition in Continuous Cultures." Bioeng. and Biotech. 21:627-648 (1979).
- Wilkinson, S. C., K. H. Goulding and P. K. Robinson, "Mercury Accumulation and Volatilization in Immobilized Algal Cell Systems". Biotech. Letters, 11:861-864 (1989).
- Wong, M.H., Chan, K.Y., S.H. Kwan, and C.F. Ng, "Metal Contents of the Two Marine Algae Found on Iron Tailings", Marine Pollution Bulletin, 10:56-59 (1979).
- Wood, J.M., and H.-K. Wang, "Microbial Resistance to Heavy Metals", Env. Sci. Tech. 17: 582-590 (1983).
- Xue, H., W. Stumm and L. Sigg. "The Binding of Heavy Metals to Algal Surfaces", Wat. Res. 22:917-926 (1988).
- Yakubu, N.A., and A.W.L. Dudeney, "Adsorption by Microbial Biomass as a Process for Removal of Ions from Process or Waste Solutions", in Eccles, H., and S. Hunt, (eds.), Immobilization of Ions by Bio-Sorption, Ellis Horwood, Ltd.(1986).
- Zimnik, P.R., and J. Sneddon, "Binding and Removal of Aluminum Ions in Waters by an Algal Biomass", Analytical Letters 21:1383-1396 (1988).

**DATE
FILMED**

5/01/92