10
224/9288(1)

ORNL/NERP-5

**OAK RIDGE
NATIONAL
LABORATORY**

MARTIN MARIETTA

**RESOURCE MANAGEMENT PLAN
FOR THE OAK RIDGE
RESERVATION**

**VOLUME 28: WETLANDS ON THE
OAK RIDGE RESERVATION**

Maureen Cunningham
Larry Pounds

Environmental Sciences Division
Publication No. 3765

MANAGED BY
MARTIN MARIETTA ENERGY SYSTEMS, INC.
FOR THE UNITED STATES
DEPARTMENT OF ENERGY

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

This report has been reproduced directly from the best available copy.

Available to DOE and DOE contractors from the Office of Scientific and Technical Information, P.O. Box 62, Oak Ridge, TN 37831; prices available from (615) 576-8401, FTS 626-8401.

Available to the public from the National Technical Information Service, U.S. Department of Commerce, 5285 Port Royal Rd., Springfield, VA 22161.
NTIS price codes—Printed Copy: A07 Microfiche A01

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

ENVIRONMENTAL SCIENCES DIVISION

RESOURCE MANAGEMENT PLAN FOR THE OAK RIDGE RESERVATION

VOLUME 28: WETLANDS ON THE OAK RIDGE RESERVATION

Maureen Cunningham¹ and Larry Pounds²

¹Science Applications International Corp., P.O. Box 2501, MS-3057,
Oak Ridge, Tennessee 37831.

²The University of Tennessee, Knoxville.

Date Published -- December 1991

Prepared for the
Oak Ridge National Environmental Research Park
Office of Health and Environmental Research
(Activity No. KP 02 04 00 0)

Environmental Sciences Division Publication No. 3763

Prepared by the
OAK RIDGE NATIONAL LABORATORY
Oak Ridge, Tennessee 37831
managed by
MARTIN MARIETTA ENERGY SYSTEMS, INC.
for the
U.S. DEPARTMENT OF ENERGY
under contract DE-AC05-84OR21400

MASTER

CONTENTS

	<u>Page</u>
ABSTRACT	1
OBJECTIVES	1
INTRODUCTION	1
METHODS	2
RESULTS	5
LOVELL QUADRANGLE	13
Sites with Wetland Vegetation	13
Sites without Wetland Vegetation	14
BETHEL VALLEY QUADRANGLE	15
Sites with Wetland Vegetation	15
Sites without Wetland Vegetation	19
ELVERTON QUADRANGLE	20
Sites with Wetland Vegetation	20
Sites without Wetland Vegetation	22
NOTEWORTHY PLANT SPECIES IN WETLANDS ON ORR	23
DISCUSSION	26
RECOMMENDATIONS	28
ACKNOWLEDGMENTS	28
LITERATURE CITED	29
APPENDIX	31

LIST OF TABLES

<u>Table</u>		<u>Page</u>
1	Participants at a planning meeting held January 1990 to define the scope of a wetlands survey of ORR	3
2	Presence or absence of hydrophytic vegetation on sites in the USGS Lovell quadrangle that were visited during a wetlands survey of ORR	7
3	Presence or absence of hydrophytic vegetation on sites in the USGS Bethel Valley quadrangle that were visited during a wetlands survey of ORR	8
4	Presence or absence of hydrophytic vegetation on sites in the USGS Elverton quadrangle that were visited during a wetlands survey of ORR	9
5	Common wetland genera and species on ORR	10
6	Sites on the NWI maps of ORR that were not visited during the wetlands survey	12
7	Additions to the checklist of vascular plants on ORR	24

LIST OF FIGURES

<u>Figure</u>		<u>Page</u>
1	The hierarchical system of codes assigned by NWI to wetlands on ORR	6
A1	Index map, giving the coverage of the ORR by ten other maps, shows possible wetland sites	35

ABSTRACT

A survey of wetlands on the Oak Ridge Reservation (ORR) was conducted in 1990. Wetlands occurring on ORR were identified using National Wetlands Inventory (NWI) maps and field surveys. More than 120 sites were visited and 90 wetlands were identified. Wetland types on ORR included emergent communities in shallow embayments on reservoirs, emergent and aquatic communities in ponds, forested wetland on low ground along major creeks, and wet meadows and marshes associated with streams and seeps. Vascular plant species occurring on sites visited were inventoried, and 57 species were added to the checklist of vascular plants on ORR. Three species listed as rare in Tennessee were discovered on ORR during the wetlands survey. The survey provided an intensive ground truth of the wetlands identified by NWI and offered an indication of wetlands that the NWI remote sensing techniques did not detect.

OBJECTIVES

The goals of the survey were as follows:

1. to identify the wetlands resource on ORR,
2. to describe the types of wetlands present,
3. to map the locations of wetlands on the ORR,
4. to inventory the vascular plant species on wetlands on ORR, and
5. to consider the potential threats to wetlands on ORR.

INTRODUCTION

Wetlands have received considerable attention in the past several years as a natural resource in need of protection. Wetlands perform a variety of important functions in the ecosystem (Sharitz and Gibbons 1989). They play a major role in the quantity and quality of water in local and regional hydrologic regimes. Floodwaters can be absorbed by wetlands, and water quality can dramatically improve after passing through a wetland. The rate of productivity in wetlands is impressively high, and wetlands provide a necessary habitat for both resident and migratory wildlife.

Wetlands vary in size from large areas such as the Everglades National Park to small riparian zones along streams. Regulations protecting wetlands have traditionally focused on relatively large wetlands and relatively large-scale disturbances. Disturbances of small wetlands and cumulative impacts from small-scale disturbances of large wetlands,

however, may also be significant and warrant attention in wetland protection.

The rate of loss of wetlands in the United States is staggering. Wetlands represent a small percentage of the land base of the United States, and in the lower forty-eight states wetland acreage is estimated to have been reduced by half over the past 200 years (Dahl 1990). Wetlands continue to annually disappear by hundreds of thousands of acres (U.S. Congress 1986). These factors have shaped a national commitment to reduce loss of wetlands in the United States.

Previous investigators on the Oak Ridge Reservation (ORR) have studied selected aquatic and riparian habitats. Loar (1984) described the aquatic habitats on ORR, DeSelm and Shanks (1967) studied vegetation development on the drained White Oak Lake bed, and Ryon and Smith (1988, 1989) and Ryon et al. (1990) have documented riparian vegetation on several areas of ORR. Prior to 1990 a systematic survey of wetlands on the reservation had not been conducted. The valuable wetland resource across ORR was not documented, and the potential impact of Department of Energy (DOE) project development on wetlands could not be assessed. Potential impacts to wetlands must be considered for compliance with floodplain/wetlands environmental review requirements issued by DOE (1979). To address this deficiency the Resource Management Organization, an ad hoc group of resource scientists and managers from the three DOE facilities on ORR, requested that a survey be conducted to identify wetlands across the reservation.

METHODS

A planning meeting to define the scope of the survey was held in January 1990 and was attended by representatives from the DOE Oak Ridge facilities (ORNL, K-25 Site, and Y-12 Plant) as well as the U.S. Army Corps of Engineers and the U.S. Fish and Wildlife Service, two agencies that have responsibilities for jurisdictional wetlands. The meeting attendees and their respective organizations are listed in Table 1. At the meeting, a decision was made to use the definition of wetland contained in the Federal Manual for Identifying and Delineating Jurisdictional Wetlands (Federal Interagency Committee for Wetland Delineation 1989). Because of the large area to be surveyed and the time frame allotted for completion, only a general survey was deemed possible. Thus the locations and descriptions of wetlands would be provided, but the lengthy task of delimiting individual wetland boundaries and other detailed descriptions would be delegated to site-specific surveys to be requested by environmental compliance staff during review of proposed projects.

Once the scope of the survey was defined, the first step was to identify potential wetland sites across ORR. National Wetlands Inventory (NWI) maps were acquired; these are prepared by the U.S. Fish and Wildlife Service using interpretation of aerial photography and overlaid on

Table 1. Participants at a planning meeting held January 1990 to define the scope of a wetlands survey of ORR

U.S. Army Corps of Engineers	Michael Bell, Brad Bishop
U.S. Fish and Wildlife Service	Robert Willis
U.S. DOE OR	David Page
Y-12 Environmental Management	Patty Phillips, Mick Wiest
K-25 Environmental Management	Bill Scheib
Martin Marietta Energy Systems, Inc., Environmental and Safety	Jim Rogers
Oak Ridge National Laboratory (ORNL) Environmental Compliance	James Hall
ORNL Environmental Sciences	Maureen Cunningham, Jim Loar, Webb Van Winkle, Pat Parr, Mike Ryon, Glenn Cada, Roger Kroodsma

U.S. Geological Survey (USGS) quadrangle maps. Aerial photographs of ORR used in the construction of the maps were taken in 1981. The NWI maps were used as base maps of wetlands on ORR, with the realization that all wetlands present would not be shown on the maps. Unique identifiers consisting of a USGS quadrangle initial and a number were added to wetland codes assigned by NWI to each mapped wetland. A wetland on the Bethel Valley quadrangle, for example, would be designated "B-NWI code-ORR number."

Additional areas not shown on the NWI maps but considered by the field investigators to be possible wetland sites were also identified and coded with a quadrangle initial and a number. These were all visited either on foot or by canoe, and data sheets were prepared for each visited site. Data collected for each site included a general description of vegetation and abundant plant species, notes on any abnormal conditions (e.g., unusual weather or a recent anthropogenic disturbance) prevailing at the time of the visit, and other comments by the field surveyors. It is important to note that the survey reports on visited sites but does not suggest that all wetlands on ORR are identified. For example, the location of wetlands on upland areas is difficult to identify from aerial photographs or to predict from topographic maps.

Three criteria must be met for an area to be considered wetland (Federal Interagency Committee for Delineating Jurisdictional Wetlands 1989). Hydric soils must be present, wetland hydrology must exist (the ground saturated or inundated for at least one week during the growing season), and hydrophytic vegetation must be dominant. If hydrophytic vegetation is dominant, it is assumed that the other criteria are met. Determining whether hydrophytic vegetation is dominant requires two time-consuming techniques, quantitative sampling and definition of the wetland boundary. Because of time constraints in this survey, dominance of hydrophytic vegetation was not determined; abundance of hydrophytic plants was used as a field indicator of wetlands. The U.S. Fish and Wildlife Service (1988) manual National List of Plant Species That Occur in Wetlands: Tennessee was used to verify hydrophytic vegetation. Thus the survey provides information on sites that have an abundance of wetland species and have a high probability of meeting the definition of wetland, at least in some portion of the site. Hydrophytic vegetation may be confined to a narrow zone or may be in discontinuous locations at the sites.

Ponds on parcel A at the ORR's extreme eastern end (Clinton quadrangle), soon to be sold to the City of Oak Ridge, were not surveyed. An environmental assessment for sale of the site has been conducted, and the status of the ponds as wetlands is described therein (DOE, June 1990).

RESULTS

Field visits to 122 possible wetland sites were completed. Of the sites visited 90 supported hydrophytic vegetation; 61 were shown on the NWI maps, and 29 sites were not. Twenty-five sites mapped by NWI as wetlands lacked hydrophytic vegetation; these were mostly farm ponds that no longer held water. Wetlands identified on NWI maps of ORR fell into one of three major categories: palustrine, lacustrine, or riverine. The vast majority of wetlands identified by NWI on ORR were palustrine; however, their combined area may not be greater than lacustrine habitat on ORR. Explanation of the hierarchical system of codes assigned by NWI to ORR wetlands is given in Fig. 1.

Definitions of the various system codes used by NWI are provided in The Classification of Wetland and Deepwater Habitats of the United States (U.S. Fish and Wildlife Service 1989) and are presented here in an abridged form (e.g., references to wetlands with ocean-derived salts are omitted).

Riverine wetlands and deepwater habitats are contained within a channel, with the exception of those dominated by trees, shrubs, or persistent emergents. A channel is "an open conduit either naturally or artificially created which periodically or continuously contains moving water, or which forms a connecting link between two bodies of standing water" (Langbein and Iseri 1960).

Lacustrine wetlands and deepwater habitats exhibit the following characteristics: (1) location in a topographic depression or a dammed river channel; (2) lack of trees, shrubs, and persistent emergents with greater than 30% areal coverage; and (3) total area in excess of 8 ha (20 acres).

Palustrine wetlands include all nontidal wetlands dominated by trees, shrubs, or persistent emergents. Also included are wetlands lacking such characteristics but with all of the following: (1) area less than 8 ha; (2) lack of active wave-formed or bedrock shoreline features; and (3) water depth in the deepest part of the basin less than 2 m at low water.

The following section provides a brief description of the sites visited. Sites are arranged by USGS quadrangle and NWI code. Sites without an NWI code are not shown on the NWI maps for ORR. The first section under each quadrangle name describes sites that supported hydrophytic vegetation; the second lists sites that lacked wetland vegetation.¹ Tables 2, 3, and 4 list the sites visited for the Lovell, Bethel Valley, and Elverton quadrangles, respectively, and indicate whether wetland vegetation was present. Plant species commonly found on ORR wetlands are in Table 5.

¹Each site visited is referenced to a map in the Appendix. An index map of ORR in the Appendix shows coverage of the individual maps.

Fig. 1. The hierarchical system of codes assigned by NWI to wetlands on ORR.

ORNL-DWG 91-17713

R - Riverine

P - Palustrine

L - Lacustrine

Modifiers

Table 2. Presence or absence of hydrophytic vegetation on sites in the USGS Lovell quadrangle that were visited during a wetlands survey of ORR

<u>Hydrophytes present</u>	<u>Hydrophytes not present</u>
L-L1OWHh	L-PF01C-1
L-PF01A-2	L-POWHh-1
L-POWHh-8	L-POWHh-2
L-POWHh-9	L-POWHh-3
L-POWHh-15	L-POWHh-10
L-POWHh-21	L-POWHh-11
L-POWHh-22	L-POWHh-16
L-POWHh-23	L-POWHx-1
L-POWHh-24	L-POWHx-3
L-POWHh-25	L-POWHx-4
L-POWHh-26	L-POWHx-7
L-POWHh-28	L-3
L-POWHh-29	
L-POWHh-31	
L-POWHh-32	
L-POWHh-33	
L-POWHh-35	
L-POWHh-36	
L-1	
L-2	

Table 3. Presence or absence of hydrophytic vegetation on sites in the USGS Bethel Valley quadrangle that were visited during a wetlands survey of ORR

<u>Hydrophytes present</u>		
B-L1OWHh	B-POWHh-13	B-19
B-L2OWHh-1	B-POWHx-1	B-20
B-L2OWHh-2	B-R4SB3A-1	B-21
B-PFO1Ah-5	B-R4SB3U-4	B-22
B-PFO1A-7	B-1	B-25
B-PFO1A-11	B-2	
B-PFO1A-17	B-3	
B-PFO1A-18	B-4	
B-PFO1A-19	B-5	
B-PFO1A-21	B-6	
B-PFO1A-22	B-7	
B-PFO1A-24	B-8	
B-PFO1A-27	B-9	
B-PFO1A-32	B-10	
B-PFO1A-99	B-11	
B-POWHh-1	B-12	
B-POWHh-2	B-13	
B-POWHh-3	B-14	
B-POWHh-7	B-15	
B-POWHh-8	B-16	
B-POWHh-11	B-17	
B-POWHh-12	B-18	
B-PFO1A-6		
B-PFO1A-25		
B-PFO1A-32		
B-POWHr-1		
B-POWH-2		
B-POWHh-9		
B-POWHh-10		
B-7		
B-23		
B-24		

Table 4. Presence or absence of hydrophytic vegetation on sites in the USGS Elverton quadrangle that were visited during a wetlands survey of ORR

Hydrophytes present

E-L1OWHh-1
 E-L1OWHh-2
 E-L2OWHh-1
 E-L2OWHh-5
 E-L2OWHh-6
 E-PAB1H-1
 E-PF01A-1
 E-PF01A-2
 E-PF01A-96
 E-PF0SF/SS1F
 E-POWHh-1
 E-POWHh-2
 E-POWHh-3
 E-POWHh-6
 E-POWHh-97
 E-POWHx-2
 E-POWHx-3
 E-R2UB3H-99
 E-1
 E-2
 E-3
 E-4
 E-5

Hydrophytes not present

E-POWHr-1
 E-POWHr-2
 E-POWHh-4
 E-POWHh-5
 E-POWHx-1
 E-R2UB3H-1
 E-6
 E-7
 E-8
 E-9

Table 5. Common wetland genera and species on ORR

<u>Trees</u>	<u>Herbaceous emergent</u>
<u>Acer negundo</u> , boxelder	<u>Acorus calamus</u> , sweet flag
<u>Fraxinus pennsylvanica</u> , green ash	<u>Carex</u> sp., sedge
<u>Platanus occidentalis</u> , sycamore	<u>Juncus</u> sp., rush
<u>Salix nigra</u> , black willow	<u>Justicia americana</u> , water willow
	<u>Eleocharis</u> sp.
	<u>Hibiscus</u> sp., mallow
	<u>Samolus parviflorus</u>
<u>Shrubs</u>	<u>Saururus cernuus</u> , lizard's tail
<u>Alnus serrulata</u> , alder	<u>Sagittaria</u> sp.
<u>Amorpha fruticosa</u> , indigobush	<u>Typha</u> sp., cattail
<u>Cephalanthus occidentalis</u> , buttonbush	
<u>Cornus amomum</u> , silky dogwood	
<u>Hypericum densiflorum</u>	
<u>Floating aquatics</u>	<u>Submerged aquatics</u>
<u>Wolffia</u> sp.	<u>Elodea</u> sp.
<u>Lemna</u> sp.	<u>Myriophyllum</u> sp.
	<u>Potamogeton</u> sp.

Some sites on ORR shown on the NWI maps were not visited during the survey (Table 6); the majority are identified as palustrine forested or riverine and are small tributaries through upland forest, where major wetland development was not considered likely because of the steep gradient. Stream drainages and ponds with significant amounts of radioactive contaminants were not visited. The White Oak Creek floodplain and White Oak Lake have extensive wetland communities and should be surveyed by staff certified to work on contaminated sites.

Table 6. Sites on the NWI maps of ORR that were not visited during the wetlands survey

B-PF01A-3	B-R4SB3U-1
B-PF01A-4	B-R4SB3U-2
B-PF01A-5	B-R4SB3U-3
B-PF01A-9	B-R4SB3U-5
B-PF01A-26	B-R4SB3U-6
B-PF01A-28	B-R4SB3U-7
B-PF01A-29	B-R4SB3U-8
B-PF01A-30	B-R4SB3U-9
B-PF01A-31	B-R4SB3U-10
B-PF01A-33	B-R2UB3H-1
B-PF01A-34	B-L2OWHh-1
B-PF01A-35	L-R4SB5A-2
B-PF01A-36	
B-PF01A-37	
B-PF01A-97	
B-PF01A-98	
B-PF01A-199	
B-POWH-3	
B-POWH-1	
B-POWHx-2	

LOVELL QUADRANGLE

I. Wetland vegetation is present at these sites. (Letters in parentheses refer to maps in Appendix.)

L-L1OWHh - Lowland along the Clinch River (Melton Hill Lake). Wetland shrubs and herbaceous emergent vegetation are abundant in some areas, primarily in embayments and small coves with very shallow water along the dissected shoreline. (I and J)

L-PFO1A-2 - A relatively open, flat, wet area adjoining McCoy Branch near Rogers Quarry. Black willow and cattail dominate. (I)

L-POWHh-8 - A large pond with cattail along the edges; floating and rooted aquatic plants present. (I)

L-POWHh-9 - A shallow pond covered with duckweed; emergent herbaceous species along the edge. (I)

L-POWHh-15 - Pond with a narrow zone of emergent herbaceous species at the edge. Pond is in a bowl-shaped depression; planted pine trees are on three sides. (I)

L-POWHh-21 - An embayment of Melton Hill Lake that is separated from the reservoir by Edgemoor Road except for a culvert. Emergent herbaceous vegetation dominates in some areas along the bank. (I)

L-POWHh-22 - A section of Melton Hill Lake that is isolated by Edgemoor Road. The water level follows that of the reservoir, so it is perhaps joined to it by a culvert that is not visible. Many emergent herbaceous species are along the banks of this shallow pool, and numerous wet meadow species occur back from the shoreline. (I)

L-POWHh-23 - An embayment of Melton Hill Lake that is separated from the reservoir by Edgemoor Road except for a culvert. Herbaceous emergent vegetation is along two sides of the pool. (I)

L-POWHh-24 - Two adjacent ponds near the shore of Melton Hill Lake. Aquatic plant species are present. (I)

L-POWHh-25 - An embayment of Melton Hill Lake that is impounded by a road. A culvert connects the embayment to the reservoir. A perennial stream flows into the embayment. Wetland shrubs and an abundance of emergent herbaceous species are present at the narrow end of the embayment. (I)

L-POWHh-26 - An embayment of Melton Hill Lake that is impounded by a road. A culvert connects the embayment to the reservoir. A small perennial stream flows into the site. A small wetland community is at the western end of the embayment. (I)

L-POWHh-28 - An embayment of Melton Hill Lake that is impounded by a road; a culvert connects it to the reservoir. Green ash, black alder, and black willow dominate the wetland community at the narrow end of the embayment. (J)

L-POWHh-29 - A shallow pond dominated by a hydrophytic grass. Another small pond is nearby. (J)

L-POWHh-31 - A steep-sided pond surrounded by black willow and silky dogwood. (J)

L-POWHh-32 - A small pond with emergent herbaceous species at the edges. (J)

L-POWHh-33 - A pond near Melton Hill Lake with numerous aquatic species and emergent herbaceous species along the shallow shoreline. (J)

L-POWHh-35 - A small pond with some emergent herbaceous vegetation and wetland shrubs. (I)

L-POWHh-36 - A broad shallow pond with an abundance of emergent herbaceous species. Some wetland shrubs are also present. (I)

L-POWHx-4 - A small pond with some wetland vegetation that dries out during the summer.

L-1 - A pond that has almost filled in with emergent herbaceous species; very little open water is left in the center. (I)

L-2 - East Fork Poplar Creek from Lake Reality to ORR boundary. The banks and floodplain of the creek are forested wetland dominated by sycamore, boxelder, and green ash. (I)

II. Sites that support little to no hydrophytic plants. (Letters in parentheses refer to maps in Appendix.)

L-PFO1C-1, 2, and 3 - Forested islands with a steep bank and very few emergent species at shoreline. (I)

L-POWHh-1 - An earthen dam is intact, but no water is present. ((J)

L-POWHh-2 - An earthen dam is intact, but no water is present. (J)

L-POWHh-3 - No wetland present. Perhaps a shallow pond that has dried up. (J)

L-POWHh-10 - Could not locate this pond; it must be dry. It does not appear on the most recent topographic maps for ORR. (I)

L-POWHh-11 - New Hope Pond is now drained and filled. (I)

L-POWHh-16 - This pond no longer holds water; no hydrophytic species are present. (I)

L-POWHx-1 - Rogers Quarry. A deep steep-sided rock quarry now filled with water. No hydrophytic vegetation at the edges. We did not take a boat out to look for aquatics. (I)

L-POWHx-3 - Steep sides to this now dry, and apparently consistently dry, pond. (I)

L-POWHx-7 - Two very steep-sided ponds side by side; no hydrophytic vegetation. (J)

L-3 - Headwaters of East Fork Poplar Creek to the now filled New Hope Pond. Headwaters within the Y-12 fence are mixed with a large volume of Y-12 process water and flow through a channel with rip-rap banks. Very little natural vegetation or creek bank remains. As the water flows around the old New Hope Pond area, it is contained in a cement channel and remains in the channel until it flows into Lake Reality. (I)

BETHEL VALLEY QUADRANGLE

I. Wetland vegetation is present at these sites. (Letters in parentheses refer to maps in Appendix.)

B-L1OWHh - Clinch River (Melton Hill Lake). Much of the river edge is a relatively steep bank with little wetland development, but hydrophytic vegetation occurs in some lowland areas. In these areas, particularly embayments, wetland trees, shrubs, and herbaceous emergent vegetation occur. (E, G, and H)

B-L2OWHh-1 - White Oak Creek Embayment. Only a small spur of this site was visited. The main body of the embayment is contaminated and so was not visited by canoe. The upper end of the embayment spur near the 0800 field site supported emergent wetland vegetation. (E)

B-L2OWHh-2 - White Oak Lake. From a distance, abundant emergent wetland vegetation was observed along the edges of White Oak Lake, but because of contamination we did not get a close look. (E)

B-PF01Ah-5 - Two small impounded ponds flow into a stream with a wet floodplain. The stream flows into a small embayment of Melton Hill Lake. Black willow and sedges are abundant. (G)

B-PF01A-7 - Wetland community development in areas where the stream crosses through a cleared power-line right-of-way. Emergent herbaceous species dominate. (D)

B-PF01A-8 - A sedge meadow is present where the stream crosses through a cleared power-line right-of-way. (D)

B-PF01A-11 - Floodplain of Bear Creek. Forested wetland vegetation occurs in varying width bordering the stream. A small tributary to Bear Creek flows into this area; the upstream section of the tributary flows through a power line right-of-way. In the wet opening, Platanthera peramoena, an orchid listed as threatened in Tennessee, was found. (D)

B-PF01A-17 - Relatively little wetland species except where the stream crosses the power-line right-of-way. In this area herbaceous emergent species occur. Platanthera peramoena, a species listed as threatened in Tennessee, grows in the wet opening. (D)

B-PF01A-18 - Forested wetland along floodplain of Bear Creek. The width of the wetland vegetation varies and is not always continuous. (F)

B-PF01A-19 - A thicket with small areas of wetland mostly along Bear Creek. (D)

B-PF01A-21 - Forested wetland along an unnamed tributary to Bear Creek. An interesting wetland not shown on the NWI map adjoins the tributary. It is partly forested but grades into a shrub area and then into an opening dominated by sweet flag, Acorus calamus. The open area has standing and slowly moving water fed by seeps. The site is part of a registered Tennessee Natural Area. During the wetlands survey a twayblade orchid tentatively identified as Liparis loeselii, a state endangered species, was found in the forested part of the wetland. Confirmation of identification was made in 1991. (D)

B-PF01A-22 - Forested wetland composed of seeps, channels, swamps, low wet woods, and peaty/marshy areas. Large patches of Carex prosina with distinct boundaries. Large areas dominated by Solidago patula and Impatiens capensis. Swamp areas with standing open water and with bare soil (void of vegetation) or blackened litter; water marks on the trees. Water in channels was flowing in some places. Sparse pine throughout the maple- and ash-dominated woodland contributed a larger than expected share of litter on the forest floor. A state endangered species, Platanthera flava var. herbiola, grows in this wetland. (D)

B-PF01A-24 - Forested wetland along East Fork Poplar Creek. Sycamore, boxelder and green ash dominate. (C and D)

B-PF01A-27 - Forested wetland along an unnamed tributary to East Fork Poplar Creek. As stream flows through a power-line opening, a wet meadow/marshy area is developed, partly fed by seeps away from the stream channel. (C)

B-PF01A-32 - Visited only the headwater of the stream. Seeps and springs are in the area. The stream has a gravel bottom. Exotic species dominate on one side of the stream but a wetland orchid, Platanthera clavelata, was present on the other side. Hydrophytic vegetation is confined to certain areas. (G)

B-PFO1A-99 - Area visited was relatively flat bottomland along an unnamed tributary to East Fork Poplar Creek. Herbaceous emergent vegetation is present. (F)

B-POWHh-1 - A narrow embayment separated from the reservoir by a roadbed and connected with a culvert. Strong zonation in the emergent wetland; Scirpus and Juncus at water's edge are bordered by a shrub Hypericum, and farthest from the shore were wetland dogwood, willow, and alder. (E)

B-POWHh-2 - Raccoon Creek embayment. A narrow embayment separated from the reservoir by a road. A culvert connects it to the main channel. Herbaceous and shrubby emergent/wet meadow vegetation is present on the long shoreline. (E)

B-POWHh-3 - A small embayment cut off from the reservoir by a road and connected by a culvert. There is a large area of wetland dominated by a single grass species. (E)

B-POWHh-7 - A very small pond created by Bethel Valley Road isolating a part of the Walker Branch embayment. Cattail and Myriophyllum are present. (G)

B-POWHh-8 - A pond with some emergent vegetation and Wolffia, a floating aquatic species. (F)

B-POWHh-11 - A large pond with zones of emergent vegetation around the perimeter. No aquatics observed. Slopes to the pond are not steep, but pond gets deep quickly. (F)

B-POWHh-12 - A shallow pond with many emergent species and large beds of aquatic species. (F)

B-POWHh-13 - A stream drainage feeding Bearden Creek embayment is cut off from the reservoir by a gravel road with a culvert underneath. Standing water next to the road supports herbaceous emergent vegetation. There is a dense alder thicket in the shady stream drainage. (G)

B-POWHx-1 - A group of ponds constructed and used for research. Two aquatic species, Myriophyllum and Elodea nuttallii, a state-listed aquatic species, are present. (D)

B-R4SB3A-1 - Headwaters and upper reaches of Bear Creek. Wetland vegetation varies through this highly disturbed section of the creek. It ranges from narrow strips of boxelder, sycamore, and willow along the bank to more open sections with low banks or broader channel where emergent and wet meadow vegetation are present. (F)

B-R4SB3U-4 - A wet stream bottom with hydrophytic species present, including a wetland orchid. (F)

B-1 - Low, wet drainage on south-facing slope of Pine Ridge near Y-12. This is a recently disturbed area with lots of rock fill and rip-rap on the sides. Water was standing and flowing through a culvert under Bear Creek Road. Some herbaceous emergent species are established. (F)

B-2 - A shallow pond at the edge of a woodland; standing dead trees, some with apparent nest holes. Water flows through a culvert to Bear Creek. Black willow and herbaceous emergents are present. (F)

B-3 - A pond in a disturbed area with a few emergent species present. Looks recently formed or reworked. (F)

B-4 - A broad, wet area bordering Bear Creek Road connected by a culvert to another wet area upslope. Water was flowing at the upslope location. A rich collection of mostly herbaceous emergent vegetation and wet meadow species. (F)

B-5 - A low, wet area with flowing water on the south-facing slope of Pine Ridge near Bear Creek Road. Many emergent and wet meadow species present. (F)

B-6 - A small pond in a low area along Bear Creek Road. Cattails abundant and aquatic species present also. (F)

B-8 - The Y-12 fly ash pond. This large area is a mosaic of bare ash, grasses and forbs, and black willow. In some sections hydrophytic species are clearly dominant. The ash sluice is no longer deposited in the pond, so the hydrology is altered. Judging from the downstream flow, however, there may still be enough flow from the springs feeding McCoy Branch to continue the wetland hydrology. (F)

B-9 - A series of small wetlands along the power-line right-of-way that runs parallel to Bear Creek Road. In 1989 a section of the right-of-way was turned into a haul road for trucks carrying clay to cap burial grounds at Y-12. The widening and grading of the right-of-way have reduced the extent of the wet areas at this site. (F)

B-10 - A series of marshy areas under the Bear Creek power-line right-of-way, east of the section that is used as a haul road (see B-9). Some emergent species present, along with wet meadow species and an alder thicket. (F)

B-11 - A pool that either is a natural oxbow of East Fork Poplar Creek or was cut off from the creek during construction of Highway 95. A culvert at one end may connect it to the creek; the pipe is bent, however, and may not allow flow. Area is forested. Rock outcrops are present along one side of the curved pool, and the other side is marshy. Hydrophytic species include herbaceous forest floor species and duckweed, a floating aquatic. (C)

B-12 - A forested boggy wetland near New Zion Cemetery. A unique community on ORR; cool water emerges slowly and steadily from the ground in several places. Sphagnum moss dominates in seep areas, and ferns cover the boggy forest floor. (E)

B-13 - A series of wetlands under the Bear Creek power-line right-of-way west of Highway 95. A great diversity of wet meadow species occur in these sites, including Platanthera peramoena. (D)

B-14 - Spring-fed streams with alder, spicebush, and fern nearby. (G)

B-15 - An excavated pond with a synthetic liner. Cattails are present at one end. (G)

B-16 - Wet meadows along Bethel Valley Road. Emergent and wet meadow species present. (G)

B-17 - A series of wet sites under the right-of-way corridor that runs parallel to Melton Valley Drive. Emergent and wet meadow species dominate in these areas. (G)

B-18 - Small pond at the Y-12 landfill. Eroded sides. The edges of the pond support a few emergent and wet meadow species. (F)

B-19 - An apparently perennial stream flows through forest near George Jones Cemetery. As it crosses a gas pipeline right-of-way, hydrophytic herbs and shrubs are present. (D)

B-20 - A section of the Bearden Creek embayment is separated from the reservoir by Melton Valley Drive, except for a culvert under the road. A small stream flows into the isolated section. Many emergent species are present along the shore. Water milfoil is abundant. (G)

B-21 - A spring-fed pond near ORNL. The water is clear, and Elodea nuttallii (an aquatic species listed by Tennessee as threatened) dominates the pond. (D)

B-22 - A small stream that leads to B-PSS1A-1. Several hydrophytic ferns are present in the area. (D)

II. These sites do not support hydrophytic vegetation. (Letters in parentheses refer to maps in Appendix.)

B-PF01A-6 - Island in the Melton Hill Reservoir that does not support wetland vegetation. (G)

B-PF01A-25 - Stream in a forested area. No wetland vegetation, even in the power-line right-of-way. (D)

B-PF01A-32 - Stream in a forested area near Walker Branch Watershed Laboratory. Banks of the stream were steep-sided; no wetland community development. (G)

B-POWHr-1 - The settling basins have been closed and filled. No hydrophytic vegetation. (F)

B-POWH-2 - Pond no longer exists. No wetland vegetation. (F)

B-POWHh-9 - A dam was noted, but no water or hydrophytic vegetation was present. (G)

B-POWHh-10 - A dam was noted, but no water was present. (G)

B-7 - A disturbed area in the Y-12 fenced enclosure. A few cattails and a willow, but most of the vegetation is not hydrophytic. (F)

B-23 - Several sites along the power-line corridor that runs parallel to Bethel Valley Road near ORNL. Wetlands have not developed in these topographic low places along the right-of-way as they have in Bear Creek Valley. (D)

B-24 - A small unnamed tributary to East Fork Poplar Creek that is not a perennial stream. Both the forested section and the opening under the power line did not support wetland vegetation. (D)

ELVERTON QUADRANGLE

I. Wetland vegetation is present at these sites. (Letters in parentheses refer to maps in Appendix.)

E-L10WHh-1 - Poplar Creek. Wetland vegetation exists in discontinuous bands along the creek on low ground and small islands and in shallow embayments. A wide array of forested wetland species and of herbaceous emergent species occurs primarily along the west bank of the creek. Limestone outcrops are abundant on the east bank. Aureolaria patula and Cimicifuga rubifolia, two rare plant species that are always associated with rivers and creeks, occur on the rock outcrops and adjacent forest. Two great blue heron rookeries are in the Poplar Creek drainage in ORR. (A and B)

E-L10WHh-2 - Deepwater channel of Watts Bar Reservoir. Much of this area is upland habitat at the river bank, but sizeable areas of well-developed wetland communities are present in several shallow embayments and on lowland. Forested wetland dominates on lowland near Gallahar Bridge, and one of the largest areas of emergent vegetation on ORR is in a narrow embayment near the bridge. Emergent vegetation is also abundant just upstream and downstream of Brashear Island. Aureolaria patula, a rare plant species that is a candidate for federal listing, occurs at shoreline just downstream of K-25. (B)

E-L2OWHh-1 - A silt bar in the sharp bend of Poplar Creek before its junction with the Clinch River. Some herbaceous emergent vegetation present but far less than in the protected shallow areas along the creek. (B)

E-L2OWHh-5 - This narrow channel supports emergent species. Wet meadow species, including an uncommon orchid, Platanthera lacera, are in the power-line corridor near the channel. (B)

E-L2OWHh-6 - Roberts Branch embayment. A community of emergent wetland species occurs in the narrow end of the embayment. (B)

E-PAB1H-1 - Two small, adjacent ponds surrounded by wetland shrubs. (B)

E-PF01A-1 AND 2 - Floodplain of East Fork Poplar Creek supports a forested wetland community. The wetland extends farther back from the creek in some areas than the NWI map shows. (A)

E-PF01A-96 - Forested wetland on lowland along the Watts Bar Reservoir edge. (B)

E-PF0SF/SS1-1F - A pond below a water filtration plant. Effluent from the plant empties into the pond and then flows into the reservoir. Pond is surrounded by wetland shrubs and herbaceous emergent vegetation. Water level fluctuates rapidly. (B)

E-POWHh-1 - Embayment on Poplar Creek that is cut off by a road and connects to the impoundment by a culvert. Forested up to the edges. Herbaceous emergent species present. (A)

E-POWHh-2 - A large pond previously used for waste retention and disposal; a spillway allows outflow to Watts Bar Reservoir. A beaver dam on the spillway has raised the water level in the pond. Dead trees are at the edge of pond. Cattail dominates the emergent community at the northwest end where a small stream enters the pond. The wet stream drainage is dominated by black willow. Some aquatic species are present in the pond. (A and B)

E-POWHh-3 - A large pond separated from Poplar Creek by a narrow strip of land; there is apparently no culvert connecting it to the creek. Much of the pond bank is steep. A sedge meadow is developed on one edge, however, and a wetland shrub community is well-developed in a strip on one shore. (B)

E-POWHh-6 - A pond alongside Highway 58. The pond is mostly steep-sided. A stream enters the back end of the pond, and in this area there is an alder thicket. Seeps and wet areas occur along the stream drainage. (B)

E-POWHh-97 - An embayment of the Watts Bar Reservoir cut off by an old roadbed. A culvert connects it to the reservoir. An emergent wetland community occurs where a small stream enters the embayment. (B)

E-POWHx-2 - A pond that has apparently either varied in size or been altered; the small pond in this vicinity doesn't seem to be on the NWI map. Sedge meadows are near the pond. (B)

E-POWHx-3 - A pond and associated canals have wetland shrub areas on the edges. (B)

E-R2UB3H-99 - Brashear Creek drainage into the Watts Bar Reservoir. An emergent wetland community occurs where the creek enters the embayment. (B)

E-1 - Wet area on a power-line right-of-way that supports emergent wetland species, including Platanthera peramoena. (B)

E-2 - A wetland between Harrell Road and East Fork Poplar Creek. Several springs are in this area. Abundant algae in the streams from the springs. This low wet area is dominated by wetland shrubs growing in fairly dense thickets. (A)

E-3 - A pond on a power-line right-of-way near K-25. An interesting site with a diverse array of emergent species and aquatic pondweed. (B)

E-4 - Mitchell Branch inside the fence at K-25. Area bordering the stream is in wetland trees and shrubs but the extent varies. In low areas or in sections with braided channel, wetland species extend back from the banks. (A)

E-5 - Mitchell Branch outside the K-25 fence. Near Portal 6 the small stream flows through a forest with some wetland shrubs near the stream, then into an open area near a parking lot. In the opening the stream area is dominated by wetland shrubs and herbaceous emergent species. Between Portal 5 and Poplar Creek, the stream area supports emergent and wet meadow species and wetland trees and shrubs. (A)

II. These sites do not support hydrophytic vegetation. (Letters in parentheses refer to maps in Appendix.)

E-POWHr-1 AND 2 - Former waste settling ponds that have been drained and sediments removed during environmental restoration. No hydrophytic vegetation. (A)

E-POWHh-4 AND 5 - Small artificial ponds with no wetland vegetation. (B)

E-POWHx-1 - A long, narrow, steep-sided pond that is cemented in some areas. No wetland development. (B)

E-R2UB3H-1 - The channel of East Fork Poplar Creek supports virtually no hydrophytic plants. Wetlands that do exist are out of the stream channel; see E-PF01A-1 AND 2 for a description of forested wetland on the floodplain. (A)

E-6 - A small stream flows through upland forest; there is no wetland development. (A)

E-7 - A pool of water is formed by the roadbed of Highway 58 near K-25, blocking the flow of a small stream. A very scenic area in the forest with extensive limestone outcrops. The terrain is steep, and the pond has no shallow areas that support wetland species. (B)

E-8 - Cleared areas along dirt roadways just west of K-25. No wetlands have developed in the clearings. (B)

E-9 - Openings on power-line rights-of-way west of K-25. No hydrophytic vegetation was observed. (A)

NOTEWORTHY PLANT SPECIES IN WETLANDS ON ORR

Vascular plant species occurring on sites visited were inventoried during the wetlands survey, and 57 species were added to the checklist of vascular plants on ORR (Table 7). Not all of the added species are wetland species. Plant specimens were collected for each new species. Several rare plant species not previously seen on ORR were observed in wetland habitats. Purple fringeless orchid, Platanthera peramoena, was observed in flower and positively identified. A floating aquatic species, identified as Elodea nuttallii, was found in two ponds; the species can be difficult to distinguish from the more common and variable Elodea canadensis, and the identification should be confirmed by an expert on this genus. A twayblade orchid, tentatively identified as Liparis loeselii, was observed in fruit, and on this basis the determination was made. The identification was verified in spring 1991, when the species was in flower.

Table 7. Additions to the checklist of vascular plants on ORR (not all of these are wetland species)

<u>Genus</u>	<u>Species</u>	<u>Variety</u>
Agalinis	fasiculata	
Ammannia	coccinea	
Aronia	arbutifolia	
Carex	blanda	
Carex	debilis	
Carex	festucacea	
Carex	gravida	
Carex	laevivaginata	
Carex	meadii	
Carex	oligocarpa	
Ceratophyllum	demersum	
Chasmanthium	laxum	
Chasmanthium	ssiliflorum	
Cyperus	brevifolioides	
Cyperus	retrofractus	
Cyperus	tenuifolius	
Elaeagnus	umbellata	
Eleocharis	palustris	
Eleocharis	quadrangulata	
Elodea	nuttallii	
Erianthus	giganteus	
Eulalia	vimineum	
Euthamia	graminifolia	
Galium	obtusum	
Galium	pedemontanum	
Gratiola	neglecta	
Hibiscus	moschuetos	palustris
Hypericum	drummondii	
Hypericum	prolificum	
Juncus	brachycarpus	
Juncus	scirpoides	
Leucospora	multifida	
Liatris	spicata	
Lotus	corniculatus	
Mosla	dianthera	

Table 7 (continued). Additions to the checklist of vascular plants on ORR (not all of these are wetland species)

<u>Genus</u>	<u>Species</u>	<u>Variety</u>
Najas	guadalupensis	
Platanthera	lacera	
Platanthera	peramoena	
Polygala	incarnata	
Polygonum	cespitosum	
Potamogeton	diversifolius	
Potamogeton	pusillus	
Prenanthes	trifoliata	
Ranunculus	acris	
Rapistrum	rugosum	
Rhyncospora	globularis	
Rorippa	palustris	
Rudbeckia	fulgida	sullivantii
Sagittaria	engelmanniana	
Sagittaria	engelmanniana	longirostra
Scirpus	americanus	
Scirpus	purshianus	
Scrophularia	marilandica	
Silphium	compositum	
Solidago	rugosa	
Taenidia	integerrima	
Wolffia	sp.	

DISCUSSION

The extensive survey of ORR for the presence of wetlands provides an information base for use in natural resource management and resource protection. The survey provides information on the presence of hydrophytic vegetation at the visited sites, but not every wetland on ORR was identified. The scope was to field-check sites shown on the NWI maps and other likely locations of wetlands. Upland sites may have wetlands that neither the NWI maps nor survey staff identified, and some sites on ORR were not surveyed because radioactive contaminants are present (e.g., the White Oak Creek watershed). For those concerned with environmental protection and compliance at the DOE Oak Ridge facilities, the survey may be most useful as an indication of the areas visited that do not support wetland vegetation. In addition, general knowledge of the wetland resources across ORR is essential for efficient project planning, particularly given the federal and state environmental regulations that must be addressed.

The ORR wetlands can be characterized either as (1) embayments of the Melton Hill and Watts Bar Reservoirs that border the reservation; (2) forested wetland along East Fork Poplar Creek, Poplar Creek, Bear Creek, and their tributaries; (3) ponds that are primarily remnants of previous farms; or (4) wet openings associated with streams and seeps. Most of the wetland sites visited are relatively small - less than 1 acre in size. Exceptions to this are large areas of forested wetland along East Fork Poplar Creek and the Clinch River near Gallahar Bridge. Several well-developed emergent communities in shallow embayments of the reservoirs are also greater than 1 acre in size. The emergent communities typically grade into a marsh adjoining forested wetland. Ponds on ORR vary in size and degree of wetland community development; several are large and support diverse flora and fauna. A number of wetland sites visited during the survey may be considered significant habitats because they either support rare taxa or are a unique example of a community on ORR.

The survey provided a good opportunity for field validation of the remotely sensed data used by NWI in preparing its maps and gives an indication of areas that the NWI mapping techniques missed. About one-third of the ORR wetlands identified during the survey were not shown on the NWI maps. The maps were helpful in locating ponds and pools with open water (POW code) in nonforested areas. Wet meadows and some wetlands in forests, however, were not identified on the NWI maps. The NWI maps for ORR did not include all the perennial stream channels as wetland habitat and it is not clear why. Lacustrine wetlands identified along the Melton Hill and Watts Bar Reservoirs and Poplar Creek did not show the adjoining palustrine forested wetland, which occurs extensively, though not continuously, along the shores.

Fifteen of the small ORR ponds identified by NWI no longer hold water. Eleven of these are presumably former livestock ponds that have dried up. The remaining four are now drained or filled artificial ponds that

served as settling basins for DOE activities. Many other ponds appear to be stable and have considerable wetland community development. One pond close to the reservoir's main channel was rich in aquatic plant species, perhaps because migrating birds may carry propagules on their feet that are deposited in the ponds.

Embayments of the Melton Hill and Watts Bar reservoirs are classified by NWI as palustrine if they are cut off from the reservoir by a road, or they are included in the lacustrine system if they are not separated. Embayments that support well-developed wetland communities are either very shallow or cut off by a road. In both cases protection is provided from motorboats and their wave action, and recreational-user traffic on the shore is reduced. Small streams commonly flow into the embayments and flow through a wetland community. It was evident during the survey that much of the debris and surface pollutants on the reservoirs accumulate in the narrow corners of embayments. The numerous wetland communities of varying size in embayments may be playing a role in water quality regulation through decomposition or adsorption of undesirable elements. That role may increase as the size of the wetlands grows over time; much of the existing wetland habitat associated with the reservoirs is only 30 to 40 years old. Abundance of wildlife associated with wetland habitat (e.g., herons and osprey) will likely continue to increase also.

The most surprising group of wetlands discovered on ORR are those on utility rights-of-way. These were predominantly in Bear Creek Valley and Melton Valley and occurred where small streams or moist drainages crossed the right-of-way. Although the upstream or upslope forested areas did not appear to be dominated by wetland species, wet meadow conditions prevailed in the openings. A rich variety of herbaceous wetland species were present, including an orchid listed as threatened in the state. Possible explanations for this observation are (1) the herbaceous species that develop so lushly in the openings may impede surface water and shallow groundwater flow and result in broad zones of wet soil; (2) there may be a differential in the degree of substrate wetness required to support a forested wetland community versus one dominated by herbaceous species; (3) the forest canopy may be a greater evapotransportative surface than the dense herbaceous community and thus deplete soil moisture to a much greater extent; or (4) the subsurface soil may be structurally different in the right-of-way openings from forest soil because of slope location or compaction from mowing equipment.

Potential threats to the ORR wetlands include habitat loss or degradation from project development activities, extreme and sustained water level fluctuation in the reservoirs, use of herbicides on the utility rights-of-way, and use of the rights-of-way for heavy truck traffic.

RECOMMENDATIONS

Although the purpose of this survey was to identify ORR wetlands, it is appropriate to make recommendations for responsible stewardship of significant ORR wetland habitat. Several wetland sites provide habitat for state-listed rare plant species previously unknown on ORR. Platanthera peramoena grows in wet meadows in power-line rights-of-way (B-PF01A-11, B-PF01A-17, B-13, and E-1). Elodea nuttallii is an aquatic species that grows in ponds (B-POWHx-1 and B-21). Liparis loeselii grows in the forested edge of the Hembree Cemetery marsh (B-PF01A-21). Sites supporting listed rare species are recognized as protected on ORR by designation as National Environmental Research Park Natural Areas (Parr and Pounds 1987). The newly discovered rare-plant sites should be thus designated.

Sites for either unique or representative plant communities on ORR warrant recognition as areas worthy of protection. A number of areas fitting this description are designated National Environmental Research Park Reference Areas (Parr and Pounds 1987). Several of the wetlands visited during the survey are unique communities either in terms of the elements present or the size or quality of the community. New Zion Cemetery wetland (B-12) is a forested, boggy area that has a unique assemblage of species. A large wetland on a shallow embayment (a section of E-L1OWHh-2) near Gallahar Bridge on State Highway 58 has extensive stands of emergent herbaceous species, shrubs, and forested wetland. Large wetlands provide habitat for some animals that cannot live in smaller patches. A well-developed pond community, including emergent and aquatic species, is present at L-POWHh-8. A smaller pond (L-POWHh-33) near the Clinch River on Freels Bend is rich in aquatic and emergent species. These wetlands should be considered candidates for Research Park Reference Areas.

ACKNOWLEDGMENTS

The authors want to thank Debra J. Awl for her assistance and helpful discussions during the course of this project and Vernon McNeilus and Hal De Selm for identification of graminoid specimens.

The authors would also like to thank Tom Chaffins and John Wagoner, Site Facilities Planning, Martin Marietta Energy Systems Engineering Division, for their help in preparing the maps.

This report was prepared by the Oak Ridge National Environmental Research Park, Environmental Sciences Division, Oak Ridge National Laboratory, Oak Ridge, Tennessee, 37831, managed by Martin Marietta Energy Systems, Inc., under contract DE-AC05-84OR21400 with the U.S. Department of Energy. Publication No. 3765, Environmental Sciences Division, ORNL.

LITERATURE CITED

- Dahl, T. E. 1990. Wetlands loss since the Revolution. National Wetlands Newsletter 12(6):16-17.
- DeSelm, H. H., and R. E. Shanks. 1967. Vegetation and floristic changes on a portion of White Oak lake bed. Ecology 48(3):410-425.
- Federal Interagency Committee for Wetland Delineation. 1989. Federal Manual for Identifying and Delineating Jurisdictional Wetlands. U.S. Army Corps of Engineers, U.S. Environmental Protection Agency, U.S. Fish and Wildlife Service, and U.S.D.A. Soil Conservation Service, Washington D.C.
- Langbein, W. B., and K. T. Iseri. 1960. General introduction and hydrologic definitions manual of hydrology. Part 1. General surface-water techniques. U.S. Geological Survey Water-Supply Paper 1541-A.
- Loar, J. M. 1984. Resource management plan for the U.S. Department of Energy Oak Ridge Reservation. Vol. 2. Appendix A: Aquatic habitats. ORNL-6026/V2. Oak Ridge National Laboratory.
- Parr, P. D., and L. R. Pounds. 1987. Resource management plan for the Oak Ridge Reservation. Vol. 23: Oak Ridge National Environmental Research Park, Research Sites, and State Natural Areas. ORNL/ESH-1/V23. Oak Ridge National Laboratory.
- Reed, P. B. 1988. National list of plant species that occur in wetlands: Tennessee. NERC-88/18.42. National Wetlands Inventory, U.S. Fish and Wildlife Service.
- Ryon, M. G., J. G. Smith, and M. J. Peterson. 1990. Substrate and cover. In J. M. Loar (ed.), Second Annual Report on the Y-12 Plant Biological Monitoring and Abatement Program (draft).
- Ryon M. G., and J. G. Smith. 1989. Substrate and cover. pp. 2-32 to 2-49. In J. M. Loar (ed.), Third Annual Report on the ORNL Biological Monitoring and Abatement Program. Oak Ridge National Laboratory, Oak Ridge, Tennessee (draft).
- Ryon, M. G., and J. G. Smith. 1988. Substrate and cover. pp 49-64. In G. R. Southworth (ed.), Ecological Evaluation of Bear Creek, 1984-1987. Oak Ridge National Laboratory, Oak Ridge, Tennessee (draft).
- U.S. Congress. 1986. Emergency Wetland Resources Act of 1986, S.740, 99th Congress, 2nd Session, Jan. 21, 1986.
- U.S. Department of Energy. 1990. Environmental Assessment: Proposed Sale of Parcel A2 of the Oak Ridge Reservation to the City of Oak Ridge, Tennessee (draft).

U.S. Department of Energy. 1979. Compliance with floodplain/wetlands environmental review requirements. pp. 12594-12599. Federal Register, Vol. 44, No. 46.

U.S. Fish and Wildlife Service. 1979. Classification of wetlands and deepwater habitats of the United States. Biological Services Program, FWS/OBS 79/31.

APPENDIX

APPENDIX

Maps for the sites on Oak Ridge Reservation (ORR) that were considered as possible wetland sites are included in the Appendix. Figure A1 indicates which portion of the Reservation each of the 10 maps represents. All sites shown on the National Wetland Inventory (NWI) maps for the ORR (which is located in Elverton, Bethel, and Lovell Quadrangles) are included on these maps. Additional sites on ORR visited by the wetland survey team are also shown. Codes are used to identify whether the sites were originally identified on the NWI maps or were additional sites identified by the ORR wetland survey team.

NWI IDENTIFIED SITES

On the NWI maps, a code is assigned to each site which gives a general description of the site. These are described in Figure 1 in the text. The same codes are used for ORR map sites. The code is followed by a hyphen and a number or, in the case of multiple sites, one code with a line to numbers located at the specific sites. References to these sites in the text and tables use a prefix of E (Elverton), B (Bethel), or L (Lovell) to indicate the Quadrangle map in which the site occurs. This prefix is not used on the map due to space restrictions.

NON-NWI IDENTIFIED SITES

Sites visited by the wetlands survey team that had not been identified on the NWI maps are identified with a prefix indicating the Quadrangle map in which they occur, and a number.

CODE OUTLINES

The codes are outlined by a shape (in some cases only the final number is outlined). These outline shapes and their descriptions are below.

Wetland vegetation is not present.

Wetland vegetation is present.

The site was not visited.

EXAMPLES OF MAP CODES

B-9

This is a site with code B-9 in the text. The outline indicates that the site was visited by the survey team and wetland vegetation was present. The lack of a NWI code shows that this site was not shown on the NWI maps, but identified as an additional site by the ORR wetland survey team.

POWHh-9

The NWI code POWHh indicates that this site was on the NWI maps. The outline shows that the site was visited by the wetland survey team and no significant wetland vegetation was found. References to this site in the text would be preceded with the letter indicating in which Quadrangle map the site occurs (B, E, or L).

Note that although both of these sites end with the number "9," they are different sites.

ORNL-DWG 91-17714

Fig. A1. Index to maps showing the location of some possible wetlands on the ORR.

S16A

Map A. Elverton Quad.

S16A

Map B. Elverton Quad.

0 1/4 1/2 1 = 1 mile

Map C. Bethel Valley Quad.

S16A

Map D. Bethel Valley Quad.

S16A

Map E. Bethel Valley Quad.

SIGA

0 1/4 1/2 3/4 1 = 1 mile

Map G. Bethel Valley Quad.

S16A

0 1/4 1/2 1 = 1 mile

Map H. Bethel Valley Quad.

A horizontal number line representing a distance of 1 mile. The line starts at 0 on the left and ends at 1 mile on the right. There are three major tick marks labeled $\frac{1}{4}$, $\frac{1}{2}$, and $\frac{3}{4}$ from left to right, dividing the line into four equal segments.

Map I. Lovell Quad.

S16A

Map J. Lovell Quad.

INTERNAL DISTRIBUTION

1. B. D. Barkenbus	80. D. E. Reichle
2. L. D. Bates	81. J. G. Rogers
3. W. M. Belvin	82. S. Sandberg
4. H. M. Braunstein	83. D. S. Shriner
5. J. H. Cushman	84. G. Snyder
6. K. M. Downer	85. C. L. Stair
7. C. H. Estes, III	86. S. H. Stow
8. M. P. Farrell	87. D. W. Swindle
9. D. E. Fowler	88. R. I. Van Hook
10. C. W. Gehrs	89. Central Research Library
11-30. C. A. Hardy	90-104. ESD Library
31. S. G. Hildebrand	105-106. Laboratory Records Dept.
32. V. W. Jones	107. Laboratory Records, ORNL-RC
33. M. E. Mitchell	108. ORNL Patent Section
34-78. P. D. Parr	109. ORNL Y-12 Technical Library
79. S. E. Poligone	

EXTERNAL DISTRIBUTION

- 110. M. Bell, U.S. Army Corps of Engineers, P.O. Box 1070, Nashville, TN 37202
- 111. B. Bishop, U.S. Army Corps of Engineers, P.O. Box 1070, Nashville, TN 37202
- 112-131. M. Cunningham, Science Applications International Corp., P.O. Box 2501, MS-3057, Oak Ridge, TN 37831-2501
- 132. K. L. Daniels, Science Applications International Corp., P.O. Box 2501, Oak Ridge, TN 37831
- 133. J. F. Franklin, Bloedel Professor of Ecosystem Analysis, College of Forest Resources, University of Washington, Anderson Hall (AR-10), Seattle, WA 98195
- 134. R. E. Frounfelker, DOE Field Office, Oak Ridge, P.O. Box 2001, Oak Ridge, TN 37831
- 135. C. S. Gist, DOE Field Office, Oak Ridge, P.O. Box 2001, Oak Ridge, TN 37831
- 136. G. M. Hornberger, Professor, Department of Environmental Sciences, University of Virginia, Charlottesville, VA 22903
- 137. G. Y. Jordy, Director, Office of Program Analysis, Office of Energy Research, ER-30, G-226, U.S. Department of Energy, Washington, DC 20545
- 138. L. Marz, DOE Field Office, Oak Ridge, P.O. Box 2001, Oak Ridge, TN 37831
- 139. W. G. McMillan, DOE Field Office, Oak Ridge, P.O. Box 2001, Oak Ridge, TN 37831
- 140. R. H. Olsen, Vice President for Research, University of Michigan, Medical Science Building II, #5605, 1301 East Catherine Street, Ann Arbor, MI 48109-0620

141. A. Patrinos, Acting Director, Environmental Sciences Division,
Office of Health and Environmental Research, ER-74,
U.S. Department of Energy, Washington, DC 20585
142. L. Pendergraft, Information Resource Center, 105 Broadway
Avenue, Oak Ridge, TN 37830
143. P. W. Phillips, DOE Field Office, Oak Ridge, P.O. Box 2001,
Oak Ridge, TN 37831
- 144-163. L. R. Pounds, 124 Arkansas Avenue, Oak Ridge, TN 37830
164. R. A. Wardrop, DOE Field Office, Oak Ridge, Radium at Y-12,
P.O. Box 2001, Oak Ridge, TN 37831
165. R. Willis, U.S. Fish and Wildlife Service, P.O. Box 845,
Cookeville, TN 38503
166. F. J. Wobber, Environmental Sciences Division, Office of
Health and Environmental Research, ER-74, U.S. Department of
Energy, Washington, DC 20585
167. Office of Assistant Manager for Energy Research and
Development, Oak Ridge Operations, P.O. Box 2001,
U.S. Department of Energy, Oak Ridge, TN 37831-8600
- 168-177. Office of Scientific and Technical Information, P.O. Box 62,
Oak Ridge, TN 37831

END

**DATE
FILMED**

3 / 18 / 92