Technical Progress Report

Contract #DE-FG02-87ER40342

between

U.S. Department of Energy

and

Colorado School of Mines

"Studies of nuclear reactions at very low energies"

Principal investigator, F.E. Cecil

Professor of Physics

Colorado School of Mines

January 15, 1992

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

EB 14 1992
ved OSTI

ds

Summary

This report summarizes the progress made in research contract DE-FG02-87ER40342 between the US Department of Energy and the Colorado School of Mines entitled "Studies in Nuclear Reactions at Very Low Energies" for the calendar year 1991. The report of our study of proton radiative capture reactions on ${}^6\text{Li}$, ${}^7\text{Li}$, ${}^9\text{Be}$ and ${}^{11}\text{B}$ between 40 and 180 keV has been accepted for publication in "Nuclear Physics A", bringing to a proper conclusion this multi-year effort. The deuteron radiative capture reactions on ${}^2\text{H}$, ${}^6\text{Li}$ and ${}^{10}\text{B}$ have been measured between center of mass energies of 20 and 140 keV. Of note is the observation that the gamma ray-to-charged particle branching ratio for the DD reaction appears independent of energy down to a center of mass energy of 20 keV, consistent with some and contrary to other theoretical models. We have investigated the ratio of the reactions $\text{D(d,p)}\text{T}$ and $\text{D(d,n)}{}^3\text{He}$ down to c.m. energies of 3 keV and the ratio of the reactions ${}^6\text{Li(d,p)}{}^7\text{Li}$ and ${}^6\text{Li(d},\alpha){}^4\text{He}$ down to a c.m. energy of 19 keV. The DD reaction ratio is independent of energy while the (d,p) branch of the $\text{D-}{}^6\text{Li}$ evinces a significant enhancement at the lowest measured energies. We have continued our investigation of charged particle production from deuterium-metal systems at a modest level of activity. Noteworthy in this investigation is the observation of 3 MeV protons from deuteron beam loaded Ti and LiD targets subject to extreme thermal disequilibria.

Significant facility improvements were realized during the most recent contract period. Specifically the downstream magnetic analysis system proposed to eliminate beam induced contaminants has been installed and thoroughly tested. This improvement should allow the $\text{D(a},\gamma){}^6\text{Li}$ reaction to be measured in the coming contract period. A scattering chamber required for the measurement of the ${}^7\text{Li}({}^3\text{He,p}){}^9\text{Be}$ reaction has been designed, fabricated and installed on the accelerator. A CAMAC based charged particle identification system has been assembled also for use in our proposed measurement of the ${}^7\text{Li}({}^3\text{He,p}){}^9\text{Be}$.

Table of Contents

I. Nuclear reactions at very low energies

A. Radiative capture reactions on light nuclei.

1. Protons
2. Deuterons
3. Alpha particles

B. The Oppenheimer-Phillips effect.

1. Reactions $D(d,p)T$ and $D(d,n)^3He$,
2. Reactions $^6Li(d,p)^7Li$ and $^6Li(d,\alpha)^4He$
3. Reactions $^{10}B(d,p)^{11}B$ and $^{10}B(d,\alpha)^8Be$

C. Charged particle emission from deuterium-metal systems

II. Material sciences

III. Operations

A. Facility improvements

1. Magnetic analysis system
2. Scattering chamber
3. CAMAC based charged particle identification system

B. Equipment failure

1. Isolation transformer
2. Nuclear Data ND66B multichannel analyzer
3. Ion source turbomolecular pump

IV. Productivity

1. Personnel
2. Publications
3. Related theses

Appendix. Reprints and preprints of recent publications

I. Nuclear reactions at very low energies

A. Radiative capture reactions on light nuclei

1. Protons

We have completed a multi-year experimental and theoretical effort devoted to the proton radiative capture reaction on ${}^6\text{Li}$, ${}^7\text{Li}$, ${}^9\text{Be}$ and ${}^{11}\text{B}$ at proton bombarding energies between 40 and 180 keV. A report on the results of this effort has been accepted for publication in Nuclear Physics A. A copy of the galley proofs of this paper is included in the Appendix.

2. Deuterons

One of the major goals of the present research grant is a extension of our earlier investigation of the $\text{D}(\text{d},\gamma){}^4\text{He}$ reaction, to energies below the lowest energy at which the reaction has been observed. The motivation for this extension is the theoretical interest which has been devoted to this reaction as a measure of the quadrupole moment of the alpha particle. Many models of this reaction process suggest that the gamma ray to charged particle branching ratio of the reaction is independent of energy below about 200 keV while others predict a significant energy dependence as the energy approaches zero. In the present study this reaction was measured at two c.m. energies, 20 and 40 keV. The spectrum measured at a c.m. energy of 40 keV (a bombarding energy of 80 keV) is shown in Figure 1. The 40 keV c.m. energy measurement overlapped with our earlier measurements and the measured branching ratio agrees with the earlier measurement. The present measurement at 20 keV c.m. energy yielded a branching ratio of $(1.4 \pm 0.3) \times 10^{-7}$ providing strong evidence that the branching ratio is constant as a function of energy. Any effort to extend this measurement to still lower

Energy spectrum for reaction $2\text{H}(\text{d},\gamma)4\text{He}$; 80 keV

energies, say 10 keV c.m. energy, with the present facility would involve several months of continuous bombardment and was deemed prohibitive. With the same techniques used in the measurement of the DD reaction, we have measured the gamma ray to charged particle branching ratios of the D- ^6Li and D- ^{10}B reactions. Typical spectra are shown in Figs 2 and 3. The gamma rays from these reactions, with energies of 23.8, 22.3 and 25.2 MeV for the target nuclei of ^2H , ^6Li and ^{10}B respectively, represent some of the most exothermic among light nuclei since the product nuclei are the tightly bound ^4He , ^8Be and ^{12}C . The energy calibration for the measurements of these extraordinarily energetic gamma rays was provided by the 4.44 MeV and 11.67 MeV gamma rays from the $^{11}\text{B}(p,\gamma)^{12}\text{C}$ reaction and the 17.25 MeV gamma ray from the $^7\text{Li}(p,\gamma)^8\text{Be}$ reaction as well as known low energy room contaminant gamma rays. This calibration curve is shown in Fig. 4. The branching ratios for the three reactions are indicated in Fig. 5. Because the yield of the reaction drops off precipitously at low energies, it will be difficult to extend these measurements to significantly lower energies. On the basis of our measurements it does appear that the ^{10}B branching ratio is independent of energy in the narrow range of bombarding energies between 150 and 120 keV while the branching ratio for the ^6Li reaction appears to be dropping with energy, again over a relatively narrow region of energies. It is noteworthy that both the ^2H and ^6Li branching ratios are at the level of $1\text{E}-7$. This identity should not be too surprising if one were to adopt a naive view of a ^6Li target nucleus as deuteron-alpha particle cluster. We are pursuing more realistic R-matrix calculations of this reaction⁴.

Energy spectrum for reaction $^{10}\text{B}(\text{d},\gamma)^{12}\text{C}$; 150 keV

Energy spectrum for reaction ${}^6\text{Li}(d,\gamma){}^8\text{Be}$; 140 keV

Gamma ray to charged particle branching ratios

3. Alpha particles

Two of the stated objectives of the present grant are measurements of the reactions ${}^2\text{H}(\alpha, \gamma){}^6\text{Li}$ and ${}^6\text{Li}(\alpha, \gamma){}^{10}\text{B}$, both at the astrophysically interesting energies around 150 keV. The D- α reaction, yielding a gamma ray of 1.54 MeV must be measured with a high resolution HpGe detector due to proximate room background gamma rays while an NaI(Tl) should suffice for the higher energy gamma ray from the α - ${}^6\text{Li}$ reaction. Our previous efforts at measuring the D- α reaction were thwarted by neutron background from contaminant deuterons resulting from target heating. Our proposed solution to this problem was the installation of a downstream bending magnet to separate the alpha beam from the contaminant deuteron beam. This system was installed (see Sec III.A.1 below) and our subsequent effort at measuring this reaction at an alpha bombarding energy of 170 keV was interrupted by the failure of the accelerator's isolation transformer (see section III.B.1 below) after only six hours of running corresponding to an accumulated charge of 0.29 C. The spectrum accumulated after this brief run (Fig 6) shows no evidence of the D- α capture gamma ray when compared to a long background run (Fig. 7). Having replaced the failed isolation transformer and completed the installation and calibration of the beam-line magnetic analysis system, we expect to resume our measurements of the D- α reaction. Our study of the α - ${}^6\text{Li}$ reaction was likewise aborted by the isolation transformer failure. As with the D- α reaction, a preliminary run showed no evidence for the 4.6 MeV gamma ray noted in our project proposal (compare Figs 8 and 9). The reaction will likewise be revisited.

D(α,γ)⁶Li; 170 keV; 0.29 C; 6 hrs @20 μ A 7/29/90

HpGe room background; passive Pb and steel shielding; 66 hrs

${}^6\text{Li}(\alpha, \gamma){}^{10}\text{B}$; 150 keV; 3.4 C; NaI detector

NaI(Tl) room background spectrum; with veto

B. The Oppenheimer-Phillips Effect.

An investigation of the Oppenheimer-Phillips effect, in which the ratios of outgoing channels in deuteron induced reactions at very low energies are affected by the distortion of the projectile deuteron in the Coulomb field of the target nucleus, is one of the major objectives of the present research grant. This effect has been investigated by comparing the yields of the various charged particle branches of specific deuteron induced reactions as a function of the deuteron bombarding energy. The branches which we have compared are of the reactions $D(d,p)T$ and $D(d,n)^3He$, the reactions $^6Li(d,p)^7Li$ and $^6Li(d,\alpha)^4He$ and the reactions $^{10}B(d,p)^{11}B$ and $^{10}B(d,\alpha)^8Be$.

1. The reactions $D(d,p)T$ and $D(d,n)^3He$

The ratio of these two reactions have been measured down to c.m. energies of 10 keV by Brown and Jarmie and 6.7 MeV by Krauss et al. We have extended the knowledge of this ratio to 3 keV c.m. energy. Using windowless ORTEC ruggedized Silicon surface barrier (SSB) detectors we were able to measure all charged products of these reactions; specifically the ratio of the yield of the tritons and 3He ions were used to measure the reaction ratio. Typical spectra measured at bombarding energies of 15 and 6 keV are shown in Figs 10 and 11. Particle groups corresponding to the 3.09 MeV proton, the 1.05 MeV triton and the 0.89 MeV 3He ion are well defined at the even the lowest energy. The ratio of the yields of p-T and n- 3He groups as a function of energy is given in Fig. 12 where it is compared to R-matrix calculations of Hale and DWBA calculations of Koonin and Mukerjee. Excellent agreement with Hale's calculations is noted. The results of this measurement have profound implications for cold nuclear fusion.

Specifically, if the production of heat is possible from room temperature nuclear reactions in the absence of significant external escaping radiation, then either the branching ratio of the two reactions given in Fig. 12 must change dramatically (many orders of magnitude!) between 3 keV and room temperature or the heat is being produced by reactions other than the D-D reaction. We presented these data and this important conclusion at the Second Annual Conference on Cold Fusion recently. A preprint of our contribution to the proceedings of this conference is attached to this report.

2. The reactions ${}^6\text{Li}(d,p){}^7\text{Li}$ and ${}^6\text{Li}(d,\alpha){}^4\text{He}$

The analogous investigation of the Oppenheimer-Phillips effect in the D- ${}^6\text{Li}$ system would involve the ratio of the yields of the very low energy ${}^7\text{Li}$ and ${}^7\text{Be}$ reaction products from the reactions ${}^6\text{Li}(d,p){}^7\text{Li}$ and ${}^6\text{Li}(d,n){}^7\text{Be}$. These reaction products are too low in energy for the windowless SSB detector system used in the D-D measurements and consequently we measured the ratio of the reactions ${}^6\text{Li}(d,p){}^7\text{Li}$ and ${}^6\text{Li}(d,\alpha){}^4\text{He}$ by measuring the yields of the 5 MeV protons and 8 MeV alphas. This ratio was measured on the assumption that an enhancement of the (d,p) reaction would dominate any Coulomb distortion induced change in the charge symmetric (d, α) reaction. The spectrum showing these reaction products measured at 80 keV is given in Fig. 13. The yield of the 3 MeV proton from the D(d,p)T reaction as deuterium from the beam builds up in the target is evident in this spectrum. This contaminant reaction is significant since the pile-up continuum from these protons constitutes the major limitation to the measurement of this ratio at very low energies. The region of the spectrum containing the 5 MeV protons from the ${}^6\text{Li}(d,p){}^7\text{Li}$ reaction at bombarding

d6Li; 80 keV

d6Li; 80 keV

d6Li;25 keV

Comparison of ${}^6\text{Li}(\text{d},\text{p}){}^7\text{Li}$ and ${}^6\text{Li}(\text{d},\alpha){}^4\text{He}$: D6LIAP14.GRA

energies of 80 and 25 keV is shown in Figs 14 and 15 respectively. The proton to alpha particle ratio measured between bombarding energies of 25 and 170 keV is shown in Fig. 16. Our measurements agree with the ratio derived from the published cross sections of Elwyn et al. at energies above 110 keV. Even neglecting the two lowest energy point (at 25 and 30 keV) by virtue of their relatively large error bars, our data indicates a significant enhancement of the (d,p) reaction at low energies. There are no published calculations of the (d,p) to (d, α) reaction with which to compare our results.

3. Reactions $^{10}\text{B}(\text{d,p})^{11}\text{B}$ and $^{10}\text{B}(\text{d},\alpha)^8\text{Be}$

As with ^6Li , our investigation of the Oppenheimer-Phillips effect in D- ^{10}B reactions involves a comparison of the reactions $^{10}\text{B}(\text{d,p})^{11}\text{B}$ and $^{10}\text{B}(\text{d},\alpha)^8\text{Be}$. A typical spectrum is Fig. 17 where the ^8Be ground state may be compared to the yield states in ^{10}B , the ground state of which is superimposed on the very broad first excited state of ^8Be . The ratio of the yields (d,p0) to (d, α 0) is shown in Fig. 18. As with ^6Li , there is an indication of an enhancement of the (d,p) reaction at low energies although the statistical accuracy of this measurement must be greatly improved before definitive conclusion may be drawn.

C. Charged particle emission from deuterium metal systems

We have continued our investigation of the anomalous production of energetic charged particles from deuterium metal systems. Although this work is supported in part by a separate research grant from Electric Power Research Institute, we include a discussion of this effort in the present report by virtue of the original interest expressed by DOE in our work and by virtue

B10(dp),(da); 170 keV

D8LIAP14.GRA

of the overlap in experimental techniques.

Our observations of energetic charged particles emitted in infrequent bursts from deuterium gas loaded Ti-662 was reported at the Provo Conference on anomalous nuclear effects in deuterium metal systems (1990). A copy of our contribution to the proceedings of this conference is included in the Appendix of the present report.

Based in part on our earlier studies of charged particle emission from deuterium beam loaded metal targets and in part on the work of Scaramuzzi et al. and Menlove et al., we have cooled some of the deuteron beam loaded targets used in the studies described in I.B.1 above and continued to monitor the targets under vacuum with the SSB detectors after the beam was turned off. Spectra were measured both with the windowless SSB detectors used in the D-D reaction studies discussed in Sec I.B.1 and with 6 micron Al foil covered SSB detectors. The spectra measured with the beams on are compared to the spectra with the beams off in Figs. 19 - 22. The beam-off spectra were measured with the target temperatures at about -80° C. The beam off spectra indicate the production of the 3 MeV protons from the D-D reaction. The maximum production rate corresponds to about $1\text{E-}18$ /(dd pair)/ (sec). We feel this is the clearest evidence to date of the D-D reaction occurring in deuterium-metal systems. This work will continue as time and equipment availability permit.

ssb detector w Al cover
70 keV

ssb detector w Al cover
5 min count
27 mins after gate valve closed
temp about -80°C

beam on; Ti662 tgt; 30 keV

beam on/ beam off cycles
12 cycles; 5 mins @ 80 μ A; 20 keV
2 mins gate valve closed
target temp about -80C
300 mm² det @ 2.75"; $A/(4\pi R^2)=.005$
Nov 18, 1991

II. Material Sciences

A comprehensive investigation of effect of a variety of ion beams of various doses on the high-temperature superconductor YBCO was completed at the beginning of the most recent contract period. This investigation utilized H, He, C, N, O and Ne beams from the laboratory's low energy accelerator at doses in the 10^{13} to 10^{16} cm⁻² range. XRD studies subsequent to the bombardments indicated that the samples were gradually amorphized and the critical temperature T_c likewise decreased as a function of dosage. On the other hand SEM and EDX measurements indicated that the surface morphology and stoichiometry remained largely unchanged following bombardment. The results of this work again emphasize the flexibility and versatility of the laboratory's accelerator, particularly the adaptability of the ion source to a wide variety of beam species with very little degradation in operating characteristics. A paper describing this work has been submitted to Nuclear Physics B for publication. A preprint of that paper is attached to this report.

III. Operations

A. Facility improvements

1. Magnetic analysis system

A beam-line magnetic analysis was proposed two years ago as a solution to the problem of deuteron beam contaminants in non-deuteron beams resulting from beam heating of the CD2 targets used in our proposed study of the $D(\alpha, \gamma)^6\text{Li}$ reaction. This system was installed in the most recent contract period and is indicated in Fig. 23, located immediately down-stream from the gate valve separating the target chamber from the main accelerator drift-tube and labelled "30 degree Bending magnet" in Fig. 23. A field map of the magnet (Fig. 24) was used as input to a ray-tracing code which specified the current settings required for specific beam species and energies. This code was checked by running various beams from the accelerator through the system. The optics of the system were checked by comparing the beam on the target and on the electrically separated magnet channel as functions of the voltage on the magnet power supply and as functions of the voltage on the beam line gridded einzel lens. From Figs 25 and 26 it is seen that with proper setting of the beam magnet and the gridded einzel lens, all of the beam can be delivered to the target. The yield of the D-D reaction as measured with a SSB detector while a CD2 target was bombarded with 100 microAmps of 120 keV using the beam-line magnetic analysis system was reduced by about four orders of magnitude from the yield which had been observed with the original accelerator configuration. On the basis of this improvement, we feel that we have achieved a successful solution to the beam contaminant problem and we look forward to a renewed effort at measuring the astrophysically important reaction $D(\alpha, \gamma)^6\text{Li}$.

Figure 2
CSM Low Energy Particle Accelerator

Figure 3
MAGNETIC FIELD MAP

Target and Chamber Current vs. Lens Voltage

Target and Chamber Current vs. Magnet Setting

2. Scattering chamber

A scattering chamber for use in measurements of angular distributions and particle identification has been designed in-house, fabricated in the CSM Physics Department machine shop, assembled, vacuum tested and installed on the accelerator at the target position of the 30 degree bending magnet. Drawings of the chamber are given in Fig. 27 and 28. This chamber will accommodate a two detector particle telescope which will operate at scattering angles from 0° to 150° from the forward beam direction and a monitor detector fixed at 90° . The chamber pumps down to the ultimate beam-line vacuum of $1\text{E-}7$ Torr. Initial tests with bringing beam into the chamber have succeeded in concentrating the beam spot to about a 4 mm square, with no umbral beam striking the target holder. The first application of this scattering chamber has been to measure angular distribution of the 3 MeV protons from the D(d,p)T reaction. Spectra measured in the variable angle detector and the monitor detector are shown in Figs 29 and 30. The target consisted of a 23 micron Al foil target which was deuterated by the beam. This foil is thick enough to allow stop the incident beam deuterons but thin enough to allow the 3 MeV protons from the D-D reaction to pass with an energy loss of about 500 keV. Consequently we were able to measure the yield of the reaction at zero degrees to the forward beam direction. The angular distribution measured between 0 and 152.5° is shown in Fig. 31 for a 70 keV deuteron beam and in Fig. 32 for a 20 keV beam. We believe that this is the first measurement of the charged particle branch of the D-D reaction at zero degrees. The angular distributions presented in Figs. 31 and 32 agrees with the distributions of Brown and Jarmies and with those of Krauss et al. Later

Angular dist. of 3 MeV proton from D(d,p)T Reaction
70 keV lab energy
Jan. 7, 1992

Angular dist. of 3 MeV proton from D(d,p)T Reaction
20 keV lab energy
Jan. 8, 1992

applications of the chamber will include a study of the reaction ${}^7\text{Li}({}^3\text{He},\text{p}){}^9\text{Be}$.

3. CAMAC based particle identification system

A particle identification capability is required for our proposed measurement of the two nucleon transfer reaction ${}^7\text{Li}({}^3\text{He},\text{p}){}^9\text{Be}$ since the Q-value for this reaction is comparable to that of the presumably more prolific single nucleon reaction ${}^7\text{Li}({}^3\text{He},\text{d}){}^8\text{Be}$. This capability will be provided by a versatile CAMAC system including an ORTEC 8411 octal ADC interfaced to an IBM PS/2 55Z computer. This system was acquired under a separate grant. The system has been tested with a thin (20 micron) DE detector using alpha sources and low energy triton from the reaction ${}^6\text{Li}(\text{n},\text{t})\alpha$ where the neutrons are supplied by the laboratory's PuBe source. A scatter plot of the DE and E values for these sources as processed by the particle identification system is shown in Fig. 33.

B. Equipment failure

The calendar year 1991 witnessed the failure of the elements crucial to the operation of the facility. In spite of the fact that replacement elements were ordered as soon as repair was deemed untenable, much time was lost that otherwise would have been devoted to data acquisition.

1. Isolation transformer

The main ion source rack supply isolation transformer failed by virtue of a short circuit in the internal transformer coils on July 29, 1991 during a 150 keV alpha particle experiment. Following several days of communication with the manufacturer, Universal Voltronics, we concluded that it would be

TID 65

Calibration of E- ΔE particle identifier
9/18/91

significantly more time and cost effective to order a replacement unit than to return the original for repair. Following a request to and approval by DOE for a reallocation of contract funds, the replacement was ordered for \$6100 on August 1 and delivered October 10. It was immediately installed and accelerator operation resumed. Discussions with the manufacturer and with the accelerator supplier (originally General Ionex) failed to shed any insight into the cause of the failure.

2. Nuclear Data ND66B Multichannel Analyzer

The laboratory's ND66B 2-ADC multichannel analyzer, which has been our workhorse since 1983, failed on June 10, 1991 due to a short circuit in one of the main circuit boards. Although only one board was damaged, the manufacturer, now Canberra/ND, assured us, in what might be cynically viewed as a tribute to American industry's commitment to planned obsolescence, that the replacement of the individual board was impossible. Consequently we acquired a Canberra/ND Accuspec System, transferred the two ND575 ADCs from the original ND66B and plugged the system into one of the laboratory's IBM PS/2 computers.

3. Ion source turbomolecular pump.

The bearings on the ion source turbomolecular pump seized up on October 13. This was not totally unexpected given the nominal 15,000 hour lifetime of these pumps. The excellent trade-in policy which Balzer's maintains supplied us with a replacement rebuilt pump with a 6 month warranty at a cost of \$1050 and a delivery time of a few days. The new pump was installed immediately and accelerator operation was resumed.

References

Ronald E. Brown and Nelson Jarmie, "Differential cross sections at low energies for $D(d,p)T$ and $D(d,n)^3He$ ", Physical Review C 41 (1990) 1391.

A. Krauss, H.W. Becker, H.P. Trautvetter, C. Rolfs and K. Brand, "Low energy fusion cross sections of D-D and D- 3He reactions", Nuclear Physics A465 (1987) 150.

S.E. Koonin and M. Mukerjee, "Branching ratios in low energy deuteron induced reaction", Physical Review C, 42 (1990) 1639. IV. Productivity

Alex Elwyn et al., "Measurement of D- 6Li reactions from 0.1 to 2 MeV", Physical Review C16 (1977) 1744

1. Personnel

Ph. D.:

F.E. Cecil, Professor, Principal investigator

J.A. McNeil, Associate Professor (NSF support)

C.S. Galovich, Assistant Professor, University of Northern Colorado

Graduate students:

D. Beddingfield

H. Liu

J.C. Scorby

J.S. Yan

Undergraduate students:

Todd Kuhlmann

Damon Christina

Joe Caccavilianna

Erica Nelson

John Hopkins

2. Publications

1. F.E. Cecil, D. Ferg, H. Liu, J.C. Scorby, J.A. McNeil and P.D. Kunz, "Radiative capture of protons by light nuclei at low energies". Nuclear Physics A, accepted for publication 1991. and Bulletin of the American Physical Society 36, 1242 (1991).

2. F.E. Cecil, D. Beddingfield, H. Liu and C.S. Galovich, "Observation of charged particle bursts from deuterium loaded thin titanium foils" in "Proceedings of workshop on anomalous nuclear effects in deuterium metal systems", Provo (1990). Eds. Steven E. Jones, Franco Scaramuzzi, and David Worledge. American Institute of Physics Conference Proceeding #228 (1991) pp. 375-382 and Bulletin of the American Physical Society 36, 1335 (1991).

3. F.E. Cecil and G.M. Hale, "D-D and D-⁶Li reactions at very low energies" in "Proceedings of Second Annual Cold Fusion Conference", Como (1991). Eds T. Bressani and C. Preparata. To be published (1992) by European Physical Society.

4. L.A. Greco, F.E. Cecil, J.U. Trefny and B. Yarar, "Ion irradiation studies of RF sputtered YBCO thin film superconductors" Nucl. Instrum and Methods B, submitted (1991).

5. D.H. Beddingfield, F.E. Cecil, H. Liu, C.S. Galovich and S. Asher, "Characterization of deuterium gas loaded thin Titanium foils" in "Proceedings of Second Annual Cold Fusion Conference", Como (1991). Eds T. Bressani and G. Preparata. To be published (1992) by European Physical Society.

3. Related theses

1. "Preparation and characterization of Ti-662 hydrides and deuterides",
David H. Beddingfield. Master of Science, May 1991.

2. "Magnetic analysis system for the CSM low energy particle
accelerator", Damon Christina, Bachelor of Science, December 1991.

preprints removed.
ds

END

**DATE
FILMED**

3 / 27 / 92