The Regents of the University of California
submit the proposal for FY 1992 to the

DE92 007024

Department of Energy

For Renewal of Grant No. ~~DOE DE-FG03-90ER40546~~
under the Outstanding Junior Investigator Program

*Progress Report
Task J*

1990 - 1991

STUDIES IN THEORETICAL PARTICLE PHYSICS

University of California at San Diego

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

Principal Investigator: David B. Kaplan

July 1, 1991

MASTER

ok
DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

TABLE OF CONTENTS

1. Proposal Overview	2
2. Background and Recent Work	3
A. Properties of Matter	3
1. <i>Strange Matrix Elements</i>	3
2. <i>The Structure of Valence Quarks</i>	4
B. The Origins of Matter	6
1. <i>Weak Scale Baryogenesis</i>	6
2. <i>Explicit Models</i>	7
C. The Origins of Mass	8
1. <i>A New Mechanism for Mass Generation</i>	8
2. <i>Semi-Compositeness and Family Structure</i>	10
3. Proposed Research	11
A. Constituent Quarks as Collective Excitations	11
1. <i>Constituent Quark Structure</i>	11
2. <i>B mesons</i>	12
3. <i>Dense Matter</i>	12
4. <i>Excited Quark States</i>	12
B. Weak Scale Baryogenesis	13
1. <i>Charge Bias and Transport</i>	13
2. <i>Higgs CP Violation</i>	14
3. <i>Supersymmetric Models</i>	14
C. Flavor Structure	15
1. <i>Phenomenology of Semi-Compositeness</i>	15
2. <i>Masses by Rank</i>	15
4. Publications	16
5. Conference Talks	18
6. Biographical Sketch	19
7. FY 1991/92 Budget	20
8. Current Support	33
9. Attachments: Selected Papers	34
1. <i>Qualitons</i>	35
2. <i>Spontaneous Weak Scale Baryogenesis</i>	55
3. <i>Flavor at SSC Energies...</i>	66

[1] PROPOSAL OVERVIEW

This research proposal is being submitted to the Department of Energy for renewal of grant no. DOE DE-FG03-90ER40546 under the Outstanding Junior Investigator Program. Support is requested for myself (David B. Kaplan), a graduate student (Martin Schmaltz) for the purpose of carrying out fundamental research in theoretical particle physics.

This proposal focuses on research on three distinct areas of particle physics:

1. **Nonperturbative QCD.** I intend to continue work on analytic modelling of nonperturbative effects in the strong interactions. I have been investigating the theoretical connection between the nonrelativistic quark model and QCD. The primary motivation has been to understand the experimental observation of nonzero matrix elements involving current strange quarks in ordinary matter—which in the quark model has no strange quark component. This has led to my present work on understanding constituent (quark model) quarks as collective excitations of QCD degrees of freedom.
2. **Weak Scale Baryogenesis.** A continuation of work on baryogenesis in the early universe from weak interactions. In particular, an investigation of baryogenesis occurring during the weak phase transition through anomalous baryon violating processes in the standard model of weak interactions.
3. **Flavor and Compositeness.** Further investigation of a new mechanism that I recently discovered for dynamical mass generation for fermions, which naturally leads to a family hierarchy structure.

A discussion of recent past work is found in the next section, followed by an outline of the proposed research. A recent publication from each of these three areas is attached to this proposal.

[2] BACKGROUND AND RECENT WORK

A. Properties of Matter

1. STRANGE MATRIX ELEMENTS

QCD is generally thought to be the correct fundamental theory of the strong interactions. Since the QCD coupling gets strong at long distances, trying to understand how the observed properties of hadrons can follow from QCD requires techniques beyond perturbation theory. What is particularly difficult to understand is how the successes of the nonrelativistic quark model might be understood in terms of QCD. This is not at all obvious: the u and d quarks of QCD are light and strongly interacting at hadronic scales; the same quarks in the quark model are heavy, nonrelativistic, and confined but rather weakly interacting.

A useful tool in understanding the nonperturbative properties of QCD is the experimental measurement of matrix elements in the proton of operators involving only strange quarks. Such matrix elements, if nonzero, would seem to contradict the quark model, and give direct experimental evidence about non-perturbative QCD. There are several pieces of intriguing experimental evidence. First of all, there is the so-called "K-N sigma term", the indirect measurement of $m_s \langle p | \bar{s}s | p \rangle \simeq 350$ MeV which suggests that approximately a third of the mass of the proton comes from the strange quark. Another piece of evidence comes from a recent experiment by the European Muon Collaboration (EMC) which finds a large value for the matrix element $\langle p | \bar{s} \gamma^\mu \gamma_5 s | p \rangle$ from deep inelastic electro-production. In these experiments data from high energy yields the above *low* energy matrix element by means of the Bjorken sum rule. As pointed out by Manohar and me an independent measurement of the same matrix element may be made in elastic $\nu - P$ scattering, since the Z^0 boson has an axial vector coupling to strange quarks*. With present data, the matrix elements determined from these completely different experiments are mutually consistent.

The existence of nonzero strange matrix elements suggest that the valence quarks of the non-relativistic quark model are complex conglomerates of the QCD

* D.B. Kaplan and A. Manohar, *Nucl. Phys.* B310 (1988) 527.

degrees of freedom. In particular, the valence u and d quarks appear to have a significant cloud of strange "QCD quarks" about them. This should not be too surprising since the s quark's current mass ($\simeq 150$ MeV) is well within the regime where QCD is strongly interacting. Nonetheless, physicists have usually assumed the absence of strange quarks in normal matter. The strange component of the proton wavefunction, and in particular the dependence of the proton mass on the strange quark mass, was investigated by Klebanov and myself in the context of the Skyrme model*.

Finally, earlier work by A. E. Nelson (UCSD) and myself on kaon condensation suggests that, as subtle as the strangeness content of normal matter may be, the consequences of the affinity of strange quarks for dense matter, such as found in "neutron" stars and possibly heavy ion collisions, may be quite dramatic. We showed in a series of papers that by analyzing the phenomenological chiral lagrangian for meson-baryon interactions that at densities of roughly three times nuclear density, a Bose-Einstein condensate of kaons is favored**.

2. THE STRUCTURE OF CONSTITUENT QUARKS

Rethinking the nature of valence quark structure was the subject of two recent papers*** and of proposed research discussed below. I have suggested that constituent quarks are solitons (called "qualitons") that consist of a local deformation of the color alignment $\bar{q}q$ pairs in the QCD vacuum. In particular, I considered the configuration

$$\langle (\bar{q}_L)_a (q_R)_b \rangle \propto \begin{pmatrix} e^{iF(r)\hat{r}\cdot\vec{\sigma}} & \\ & 1 \end{pmatrix}_{ab}$$

where a, b are color indices, $\vec{\sigma}$ are the Pauli matrices, and $F(r)$ is a function which equals π at $r = 0$ and zero at $r = \infty$. The gluon configuration is then found by solving the classical Yang-Mills equations for the gluon field in the presence of the mean fields for $\langle (\bar{q}_L)_a (q_R)_b \rangle$; one finds that at long distance it corresponds to a Coulomb color electric field in the T_8 component of color, as well as color magnetic

* D.B. Kaplan and I. Klebanov, *Nucl. Phys.* B335 (1990) 45.

** D. B. Kaplan and A. E. Nelson, *Phys. Lett.* B175 (1986) 57; *Phys. Lett.* B192 (1987) 193; *Nucl. Phys.* A479 (1988) 273c, 285c.

*** D.B. Kaplan, *Phys. Lett.* B235 (1990), 163; *Nucl. Phys.* B351 (1991) 137. A copy of the latter paper is attached to this proposal.

dipole fields in the $T_{1,2,3}$ components:

$$B_a^i \rightarrow \frac{r_B}{r^3} (\delta_{ia} - 3\hat{r}_i \hat{r}_a)$$

where $i = 1, 2, 3$ is a spatial index and $a = 1, 2, 3$ is an adjoint color index, and r_B is the color magnetic dipole strength of the qualitons, which can be calculated in this model.

Semiclassical quantization of these classical field configurations is a little delicate, since they possess an infinite number of zero modes corresponding to color gauge transformations. I have shown how the quantization may be accomplished, and that the quantum states one finds include ones with the same quantum numbers as the true quarks of QCD, due to the soliton topology. In particular, the states have baryon number $B = 1/3$, spin $J = 1/2$, and transform as color triplets. This is remarkable—and due to the topology of the above ansatz—considering that the $\bar{q}q$ bilinear out of which the soliton is constructed has $B = 0$, $J = 0$ and must transform under color as either a singlet or octet. Of interest is also the existence of excited quark states in the qualiton model, the lightest being a $B = 1/3$ state which has $J = 1/2$ but transforms as a $\bar{6}$ under color, thus behaving like a quark-gluon bound state.

Unlike the fundamental quarks, qualitons have a mass, comparable to the constituent quark mass, and are spatially extended with nontrivial internal quark and gluon structure. In this model, a B meson, for example, looks like a heavy, pointlike b quark sitting inside a cloud of $\bar{u}u$ pairs which carries the quantum numbers of a single $\bar{u}$ quark. One may regard this cloud as a valence $\bar{u}$ quark, or alternatively as a topological polarization of the QCD vacuum which screens the b quark's color.

While at first the qualiton model seems quite different than the bag model, I am beginning to believe that it is simply a more sophisticated version of it. For example, if one were interested in the spatial dependence of $\langle \bar{q}q \rangle$ in a B meson in the qualiton model, it is given by the trace of the above ansatz for $\langle (\bar{q}_L)_a (q_R)_b \rangle$ over the indices a, b , yielding something that varies from 3 at $r = \infty$ to -1 at $r = 0$. Taking into account the motion of the qualiton about the static b quark will smear out this result, and one will end up with a spatial dependence for $\langle \bar{q}q \rangle$ similar to that proposed in the bag model: nonzero outside the B meson, close to zero inside

the B meson. A big difference is that no quarks need to be put into the "qualiton bag", since the "bag" carries all of the quantum numbers, including color. The advantage of course, is that the qualiton model is a relativistic field theory, while the bag model is not.

The qualiton model for hadronic structure is something I intend to pursue further, as discussed below in the section on proposed research.

B. The Origins of Matter

1. WEAK SCALE BARYOGENESIS

The photon to baryon ratio observed in the universe today is about 10^{-9} . What is remarkable is that there is *any* asymmetry between baryons and anti-baryons, since we see none on the microscopic level. In the late sixties Sakharov pointed out that baryogenesis would require (1) Baryon number violation; (2) C and CP violation; (3) Nonthermal processes. In the mid-seventies it was realized that the CP violating late decay of particles from Grand Unified Theories (GUTS) could lead to a baryon asymmetry. This is the "standard" picture. One of the great drawbacks of the standard baryogenesis scenario is that it is not testable--all the interesting physics occurs at such a high temperature that the only relic around today is the baryon asymmetry itself.

In the mid-seventies 'tHooft realized that baryon number was violated in the $SU(2) \times U(1)$ theory of weak interactions, now a greatly tested and successful theory. This so-called "anomalous" baryon violation was merely a theoretical curiosity though, since any such process was hugely suppressed at zero temperature. However, starting with speculations by Dimopoulos and Susskind in the late 1970's, and by Kuzmin, Rubakov and Shaposhnikov in the mid- 1980's, it has since been generally agreed upon that these baryon violating processes might have occurred in thermal equilibrium when the temperature of the universe was greater than the $SU(2) \times U(1)$ symmetry breaking scale (about 250 GeV).

Baryogenesis requires more than just baryon violation. The ingredients that are needed, aside from anomalous baryon violation, are CP violation and

a departure from thermal equilibrium. The latter is possible, as Kuzmin *et al.* pointed out, if the weak phase transition is first order. If such is the case (and many viable models appear to undergo a first order weak phase transition), then there will always be some supercooling and departure from equilibrium.

As for CP violation, it is easy to show that Kobayashi-Maskawa (KM) CP violation is always too small to account for the observed baryon asymmetry, due to suppression from small Yukawa couplings. (Basically, the reason is that a dimensionless measure of CP violation in the KM model yields a number of order 10^{-20} , which is much smaller than the observed baryon to entropy ratio of 10^{-10} .) As a result, a successful theory of weak scale baryogenesis necessarily involves an extension of the standard model to incorporate extra CP violation.

This explains why weak scale baryogenesis is such an interesting topic to pursue: from the observed baryon asymmetry one is led to constraints on the symmetry breaking sector of the weak interactions (in order to obtain a first order phase transition) as well as predictions for new particles and new CP violation at the weak scale—all with experimental consequences.

2. EXPLICIT MODELS

Having all of the necessary ingredients in an acceptable theory does not necessarily lead to baryogenesis and it is important to construct and analyze explicit models. I have been involved in this endeavour with coauthors A. E. Nelson (UCSD) and A. G. Cohen (BU) for the last 14 months. In 1990 we developed the first explicit and phenomenologically acceptable model for weak scale baryogenesis based on an extension of the standard model involving right handed neutrinos and a CP violating neutrino mass matrix*. During the weak phase transition the phase boundary sweeps through space. In this model, CP violation in the neutrino mass matrix leads to a net lepton number flux being reflected in front of the moving phase boundary, even though there is no net lepton number impinging on the domain wall. This excess lepton number is then converted into baryon number in the unbroken phase by anomalous weak processes. The phase boundary then passes over the baryons, which saves them from decay since anomalous baryon

* A.G. Cohen, D.B. Kaplan and A.E. Nelson, *Phys. Lett.* B245 (1990) 561; *Nucl. Phys.* B349 (1991) 727.

violating processes are highly suppressed in the broken phase. As a consequence of this model, we find that in order to get a sufficiently large baryon asymmetry, the tau neutrino mass is constrained to lie in the experimentally accessible range

$$5 \text{ MeV} \lesssim m_{\nu_\tau} \lesssim 35 \text{ MeV} .$$

More recently we have been investigating models in which the requisite CP violation is in the Higgs sector* By making use of the mechanism of "spontaneous baryogenesis" developed in an earlier work by Cohen and myself** we showed how a first order weak phase transition with Higgs CP violation can temporarily generate a chemical potential for axial top quark number (only the top quark is involved, due to its large Yukawa coupling). Creating an asymmetry in left handed top quark number (balanced by an asymmetry in right handed top number, so that there is no baryon asymmetry) seems an accomplishment of dubious value! What is remarkable about this paper though is that we show by means of simple thermodynamic arguments that this asymmetry in axial top quark number is translated into a baryon number asymmetry by anomalous weak interactions.

C. The Origins of Mass

1. A NEW MECHANISM FOR MASS GENERATION

In a recent preprint I proposed a new mechanism for generating fermion masses from spontaneous $SU(2) \times U(1)$ violation ***. In the standard model, fermion masses arise from Yukawa couplings of the fermions with the Higgs field, which develops an expectation value. A similar mechanism is used in extended technicolor theories (ETC). In these models the $SU(2) \times U(1)$ violating parameter that gives the W and Z their masses is a fermion bilinear expectation value $\langle TT^c \rangle$, where T is a left handed technifermion, and T^c is a left handed anti-technifermion.

* A. G. Cohen, D. B. Kaplan and A. E. Nelson, UCSD preprint PTH 91-11, to appear in *Phys. Lett. B*.

** A. G. Cohen and D. B. Kaplan, *Phys. Lett. B* 199 (1987) 251; *Nucl. Phys. B* 308 (1988) 913.

*** D. B. Kaplan, UCSD preprint PTH 91-04, submitted to *Nucl. Phys. B*.

T and T^c carry weak quantum numbers as well as a new strong interaction – “technicolor”. In ETC models, this bilinear is coupled to ordinary fermions by means of a four fermion operator of the form

$$\frac{1}{\Lambda^2} f f^c (T T^c)^\dagger ,$$

where the Weyl fields f and f^c represent the ordinary technicolor-neutral fermions, and Λ the scale of new physics. When the technifermions condense, a mass of size $\langle T T^c \rangle / \Lambda^2$ is given to the fermion f , analogous to the scalar Higgs mechanism.

The new mechanism I have proposed also relies on technicolor to give the W and Z bosons their masses, but involves a different mechanism than ETC for giving fermions their masses. Instead of couplings of the above form, I considered the four fermion operators

$$\frac{a}{\Lambda^2} f T T T + \frac{b}{\Lambda^2} f^c T^c T^c T^c$$

These operators lead to mixing between ordinary fermions and technibaryons of strength

$$\mu \simeq \frac{|\psi(0)|^2}{\Lambda^2}$$

where $\psi(0)$ is the technibaryon wave function at the origin. The operator TTT produces a left handed technibaryon, while $T^c T^c T^c$ produces a right handed technibaryon, and the two are wedged by the technibaryon mass M at the weak scale. This leads to a mass matrix for the ordinary fermion/technibaryon system of the form

$$\mathcal{M} = \begin{pmatrix} 0 & \mu \\ \mu & M \end{pmatrix} ,$$

leading to a heavy fermion of mass M (the “technibaryon”) and a light state of mass μ^2/M , the ordinary fermion.

Alternatively, mass generation for the f and f^c fields may be viewed as resulting from a three loop diagram joining the f and f^c fields via virtual T and T^c lines which are joined by the $SU(2) \times U(1)$ violating condensates. This picture shows how the mechanism differs from the Higgs and ETC mechanisms: fermion masses are proportional to *three* powers of the $SU(2) \times U(1)$ symmetry violating parameter, rather than one.

This mechanism also realizes in a new fashion part of the old and previously unsuccessful program for constructing light fermions as composites of strongly interacting constituents. This always required non-QCD like strong interactions, which are hard or impossible to analyze. In the mechanism I have described, ordinary fermions are light and *semi*-composite—since they mix with technibaryons—but the constituents are assumed to have entirely QCD like interactions. The fermions are light not due to bizarre strong interactions, but because the mixing with composite states is small.

2. SEMI-COMPOSITENESS AND FAMILY STRUCTURE

As interesting as a new mass generating mechanism for fermions may be, the true value of the new mechanism I have described lies in its relation to flavor structure. In all previous models, involving Higgs fields or ETC interactions, the mass operator for fermions has been bilinear in the fermion fields. Thus the interactions have always transformed as $(3, \bar{3})$'s under the $SU(3) \times SU(3)$ chiral flavor symmetries of the fermions (the "3" comes from the 3 families). In contrast, an operator of the form $fTTT$ transforms as a simple triplet under the $SU(3)$ symmetry that rotates the f field through the three families. As a result, the 3×3 mass matrices for fermions of a given charge (charged leptons, for example) will necessarily be made up of products of 3-vectors, rather than 3×3 matrices. This has important consequences, since a 3×3 matrix of the form $M_{ij} = a_i b_j$ is only of rank one. If the mass matrix for the fermions turns out to have the form

$$M_{ij} = (a_i b_j) + \epsilon (c_i d_j) + \epsilon^2 (e_i f_j)$$

with $\epsilon \ll 1$, then a family hierarchy of masses results.

In the above cited preprint I have shown that a natural model can be constructed that results in such a hierarchical form, where the ϵ 's are nothing but the factors of $1/16\pi^2$ that arise from loops. In other words, a high energy theory with no family symmetry structure at all can lead to a family hierarchy for ordinary fermions, using this "semicompositeness" mechanism. This is what I consider the most interesting aspect of this mechanism.

[3] PROPOSED RESEARCH

In this section I outline the directions I expect my research to take in the coming year. These investigations fall into three categories: constituent quarks as collective excitations, weak scale baryogenesis, and flavor structure of fermion masses—all continuations of past work described in the previous section.

A. Constituent Quarks as Collective Excitations

1. CONSTITUENT QUARK STRUCTURE

The qualiton model of constituent quarks as solitons made out of current quarks and glue provides a framework for computing form factors for constituent quarks. The form factors of constituent quarks which I intend to calculate are:

- The magnetic dipole moment;
- The color-magnetic dipole moment;
- The axial form factor g_A for pion-quark couplings;
- The $\langle \bar{3}_3 \rangle$ matrix element in constituent U and D quarks;
- Gluon matrix elements (i.e. glue contribution to the spin, etc.).

All of these calculations involve numerical work, and I am presently communicating with Marek Karliner on the subject.

It is important to understand these numbers, since it might shed light on why the constituent quark model works so well—in particular, why a theory of strongly interacting, nearly massless fermions (QCD), seems to behave like a theory of weakly interacting massive particles. I have shown by means of my qualiton model how the kinematics work (i.e, quantum numbers); it remains to be seen whether dynamical properties such as those listed above work equally well.

2. B MESONS

The next step is to go beyond quark properties and examine color singlet states. Obviously, an explanation of observed hadron spectroscopy is the ultimate goal. The first states I intend to focus on are the B mesons, since they are relatively simple, containing a heavy quark serving as a static color source, and only one light quark. I believe that such a calculation is feasible, using the coherent state path integral formalism to deal with the spin and color orientation of the b quark. The goal of this work is to calculate the $B - B^*$ mass splitting, as well as the decay constant f_B .

3. DENSE MATTER

With two or three flavors of quarks, the qualiton model contains both quark-like solitons (qualitons) as well as baryon-like ones (skyrmions). As such it offers a unique framework for investigating the properties of quark dissociation in dense matter. This could shed light on the EMC effect, as well as theoretical suggestions that matter at neutron star densities might prefer to comprise of quark matter instead of baryons. As a first step I intend to use the technique devised by others to look at dense matter in the skyrme model, by confining unit baryon number to a hypersphere, and investigating the spatial distribution as a function of the volume V of this hypersphere. Previous work looked for chiral symmetry restoration in the Skyrme model; a similar calculation in the qualiton model can also look for a dissociated quark phase, as well as investigate gluon behaviour in dense matter.

4. EXCITED QUARK STATES

A final subject of inquiry in the qualiton model is to look at the excited quark states. As mentioned above, the model predicts a possibly metastable state with the quantum numbers of a $\bar{q}qq$ or qa state: baryon number $B = 1/3$, spin $J = 1/2$ transforming as a color $\bar{6}$. Such constituents could play a role in meson spectroscopy*, and would be interesting to investigate.

* T.H. Burnett and S.R. Sharpe, Ann. Rev. Nucl. Part. Sci. 40 (1990) 327.

B. Weak Scale Baryogenesis

As mentioned in the Background section of this proposal, "weak scale baryogenesis" refers to making use of the anomalous baryon violation already present in the standard model to produce the observed baryon asymmetry of the universe.

1. CHARGE BIAS AND TRANSPORT

As discussed in the on part B of the previous section of this proposal (pp. 6-8), my previous work on weak scale baryogenesis focused on two models: the first involving CP violating neutrino masses, the second involving top quarks interacting with Higgs fields with CP violating self interactions. In these models one can recognize two basic mechanisms needed for weak scale baryogenesis:

- **Charge Bias.** This refers to the discovery in our top quark model that CP violating interactions between particles and the phase boundary during a first order weak phase transition that is biased toward production of *any* $U(1)$ charge which is not orthogonal to baryon number, will cause anomalous weak processes to create a baryon number asymmetry. In our top quark model, the biased charge was axial top quark number.
- **Transport.** The other mechanism needed is charge transport, developed in our neutrino model. The idea is that the biased charge discussed above needs to be transported away from the broken phase and back into the unbroken phase of the universe, since it is only there that anomalous interactions are in thermal equilibrium. This is possible when the charge biasing mechanism arises particles being struck by the moving phase boundary. The particles are reflected back into the unbroken phase, and one can calculate the asymmetric reflection coefficients given a model for the reflected particle and its interactions with the Higgs sector.

The next step is to synthesize these two mechanisms and show how they work together in various models to produce sufficient baryon number. At the moment the promising models to look at in greater depth are those with CP violation in the Higgs sector, and the special case of supersymmetry. The ultimate goal is not to just explain the observed baryon asymmetry, but to make phenomenological

predictions for *LEP2*, *LHC*, and *SSC* experiments. Already from the neutrino model we have a prediction for the tau neutrino mass, which would be measurable in tau-charm factory being proposed in Spain.

2. HIGGS CP VIOLATION

In models with *CP* violation in the Higgs sector, heavy particles, such as the top quark, can be reflected from the $SU(2) \times U(1)$ phase boundary in a charge biased manner. The effect is a tree level one, and calculable directly from the Dirac equation with a space dependent mass term for the fermion. What happens is that reflections from various points in the bubble wall interfere with each other in a *CP* violating manner, leading to reflection coefficients that differ for components of different charges. I intend to analyze this effect in the two Higgs model with reflected top quarks to estimate the size of the *CP* violation in the Higgs sector. If the *CP* violating angle ξ is of order unity, then the electric dipole moments of both the neutron and electron will be predicted to be only slightly above the current experimental bounds.

3. SUPERSYMMETRIC MODELS

Finding a charge bias mechanism in supersymmetric models is somewhat more challenging, since *CP* violation (in the minimal supersymmetric model) may be rotated out of the Higgs sector entirely. Thus loop effects are required it seems. On the other hand, charge transport may prove to be extremely easy, as there are many heavy exotic particles in supersymmetric models which could reflect from the phase boundary. As supersymmetric models are interesting and viable candidates for extensions of the standard model, investigating whether or not weak scale baryogenesis is possible in supersymmetric theories is of great interest, especially since it appears somewhat constrained.

C. Flavor Structure

1. PHENOMENOLOGY OF SEMI-COMPOSITENESS

As described in section 2 of this proposal, I have recently devised a new mechanism for generating fermion masses*. With this "semi-compositeness" mechanism, fermion fields which are forbidden by $SU(2) \times U(1)$ symmetry from acquiring masses, mix with technibaryons at the weak symmetry breaking scale, resulting in a sort of "see-saw" mass matrix. The light states are partial mixtures of technibaryons—the lighter the fermion, the less the admixture of technibaryon. Since neither baryon number nor technibaryon number is individually conserved in these models, they may be expected to lead to interesting SSC phenomenology. In particular, I intend to investigate whether production of technibaryons at the SSC may be significantly enhanced.

Another possibly interesting phenomenological consequence of this mechanism is that left handed b quark is forced to have a large mixing with technibaryons, since it is mated to the left handed top quark which is very heavy. Thus the b may have large form factors in its couplings to the Z that would deviate from standard model predictions.

2. MASSES BY RANK

Another feature of the model I devised is that a natural family hierarchy arose without being put in by hand, due to the fact that at tree level, one loop and two loops, contributions to the fermion mass matrices are each strictly of rank one. I intend to investigate whether there are some general mechanisms at work here that are independent of the particular model I have proposed.

* "Flavor at SSC Energies...", UCSD PTH 91-04, attached to this proposal.

[4] PUBLICATIONS

David B. Kaplan

- [1] Quantitative Evaluation of Mouse Peritoneal Macrophage Spreading by Image Analyses... (with D. Schovaert-Brossault et al.), *Acta Stereol* 1983; 2/1: 159.
- [2] $SU(2) \times U(1)$ Breaking by Vacuum Misalignment (with H. Georgi), *Phys. Lett.* 136B (1984) 183.
- [3] Dynamical Generation of Supersymmetry, *Phys. Lett.* 136B (1984) 162.
- [4] Composite Higgs Scalars (with H. Georgi, S. Dimopoulos), *Phys. Lett.* 136B (1984) 187.
- [5] Calculation of the Composite Higgs Mass (with H. Georgi, P. Galison), *Phys. Lett* 143B (1984) 152.
- [6] Composite Higgs and Custodial $SU(2)$ (with H. Georgi), *Phys. Lett.* 145B (1984) 216.
- [7] Anatomy of a Composite Higgs Model (with M. Dugan, H. Georgi), *Nucl. Phys.* B254 (1985) 299.
- [8] Opening the Axion Window, *Nucl. Phys* B260 (1985) 215.
- [9] Manifesting the Invisible Axion at Low Energies (with H. Georgi, L. Randall), *Phys. Lett.* 169B (1986) 73.
- [10] Current Mass Ratios of the Light Quarks (with A. Manohar), *Phys. Rev. Lett.* 56 (1986) 2004.
- [11] Strange Goings On In Dense Nucleonic Matter (with A. Nelson), *Phys. Lett.* 175B (1986) 57.
- [12] Strange Condensate Realignment In Relativistic Heavy Ion Collisions (with A. Nelson), *Phys. Lett.* 192B (1987) 193.
- [13] Anomalous Vortices and Electromagnetism (with A. Manohar), *Nucl. Phys.* B302 (1988) 280.

- [14] Thermodynamic Generation of the Baryon Asymmetry (with A. Cohen), Phys. Lett. 199B (1988) 251.
- [15] Spontaneous Baryogenesis (with A. Cohen), Nucl. Phys. B308 (1988) 913.
- [16] Strange Matrix Elements in the Proton From Neutral Current Experiments (with A. Manohar), Nucl. Phys. B310 (1988) 527.
- [17] Isocurvature Baryon Number Fluctuations in the Inflationary Universe (with A. Cohen), Phys. Lett. 216B (1989) 20.
- [18] The Exact Metric About Global Cosmic Strings (with A. Cohen), Phys. Lett. 215B (1988) 67.
- [19] The Role of the Strange Quark in the Large-N Skyrme Model (with I. Klebanov), Nucl. Phys. B335 (1990) 45.
- [20] Constituent Quarks as Collective Excitations of QCD, Phys. Lett. B235, (1990) 163.
- [21] Weak Scale Baryogenesis (with A.G. Cohen and A.E. Nelson), Phys. Lett. B245 (1990) 561.
- [22] Baryogenesis at the Weak Phase Transition (with A.G. Cohen and A.E. Nelson), Nucl. Phys. B349 (1991) 727.
- * [23] Qualitons, Nucl. Phys. B351 (1991) 137.
- * [24] Flavor at SSC Energies: A New Mechanism for Dynamically Generated Fermion Masses, preprint UCSD PTH 91-04.
- * [25] Spontaneous Baryogenesis at the Weak Phase Transition (with A. G. Cohen and A. E. Nelson), preprint UCSD PTH91-11.

Papers marked by a * have been attached to this proposal.

[5] CONFERENCE TALKS

David B. Kaplan

- [1] The Composite Higgs Mechanism, *Proceedings of the Santa Fe Meeting (Oct. 31 - Nov. 3, 1984)* (1984), ed. T. Goldman, M. Nieto.
- [2] The Composite Higgs Mechanism, *Proceedings of the 23rd International Conference on High Energy Physics (July 16 - 23, 1985)* (1985), ed. S. Loken.
- [3] Kaon Condensation in Dense Matter (with A. Nelson), talk given at the Bad Honnef conference on *Strangeness in Hadronic Matter* (June 1987), published in: Nucl. Phys. A479 (1988) 273c.
- [4] Kaon Condensation in Heavy Ion Collisions (with A. Nelson), talk given at the Bad Honnef conference on *Strangeness in Hadronic Matter* (June 1987), published in: Nucl. Phys. A479 (1988) 285c.
- [5] Flavor Mixing in Composite Models, talk presented at *The 12th Johns Hopkins Workshop (1988)*, published in the proceedings.
- [6] Qualitons, talk given at *The Santa Fe QCD Workshop*, Aug. 1990 (unpublished).
- [7] Spontaneous Baryogenesis at the Weak Phase Transition, talk presented at *The Workshop on the Weak Phase Transition (May 16-18, 1991)*, The Institute for Advanced Studies (unpublished).

END

**DATE
FILMED**

3 / 4 / 92