100-11510

ONE REGIONAL ARM GUIDE FOR CLIMATIC EVALUATION

ROBERT M. BROWN

APRIL 1990

DEPARTMENT OF APPLIED SCIENCE

BARCLAY CENTER, NATIONAL BUREAU OF STANDARDS

100-11510-1

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

DISCLAIMER

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Department of Energy, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.

BNL--46836

DE92 007188

ONE REGIONAL ARM GUIDE FOR CLIMATIC EVALUATION

Robert M. Brown
Department of Applied Science
Brookhaven National Laboratory
Upton, NY 11973

April 1990

MASTER

UNDER CONTRACT NO. DE-AC02-76CH00016 WITH
THE UNITED STATES DEPARTMENT OF ENERGY

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

A Regional ARM Guide

Table of Content

	Page
Introduction	1-5
1. Continentality.	6-7
2. Temperature.	
a. Average temperatures. (Jan. and Jul.) Maximum, Mean and Minimum.	8-13
b. Monthly average temperatures for Wichita, Dodge City, Topeka, Concordia and Goodland.	14-18
c. First and last killing frost.	19
3. Insolation.	
a. Annual duration for Kansas.	20
b. Annual duration for the U.S.	21
4. Cloudiness.	
Monthly days of clear, partly cloudy and cloudy for Wichita, Dodge City, Topeka, Concordia and Goodland.	22-24
5. Precipitation.	
a. Annual amounts for Kansas.	25
b. Monthly amounts for Wichita, Dodge City, Topeka, Concordia and Goodland.	26
6. Thunderstorms and Hail.	
a. Annual number of thunderstorms for Kansas.	27
b. Annual number of thunderstorms for the U.S.	28
c. Hour when thunderstorms occur in the U.S.	28
d. Hail risk vs. elevation.	29
e. Hail loss ratio vs. elevation.	30
f. Average number of days of hail. Kansas and the U.S.	31
c. Incidence of large hail. Kansas and the U.S.	32
d. Tracks of Tornadoes in the U.S. including Kansas.	33
7. Wind.	
a. Table of monthly averages.	34
b. Annual wind speeds.	35
c. Wind Roses. (Jan. and Jul.)	36-37
d. Low-level jet.	38
8. Droughts.	
Palmer's drought index for Kansas.	39
9. Fog.	
a. Annual number of fog days in Kansas.	40
b. Map of Kansas rivers.	41
10. Concordia Climate.	
Table of climate conditions for major station near Manhattan, Kansas.	42
11. References.	43

One Regional ARM Guide for Climatic Evaluation

Introduction

One of the early tasks of the Atmospheric Radiation Measurements (ARM) Program is to provide climatic guides for site selection purposes including possible continental, regional, local and on-site locations. The first guide "A Preliminary ARM Guide for Climatic Evaluations" provided some climate data on a continental scale; this one is an attempt to show the variability that exists over a region.

Kansas was chosen for this particular guide because it satisfies most of the requirements given in the ARM Program Plan, i.e., climatic significance, potential for synergism with other programs and scientific and logistical viability.

Kansas has extreme climatic variations, is centrally located, is compatible with other large scale programs (Fife), has good airfields and accommodations to minimize time and effort in planning and operating an ARM site for continuous use and special campaigns.

1. General Features.

Kansas is located near the center of the the contiguous United States and has a distinctly continental climate with wide variations in temperature and precipitation. The continentality of Kansas as compared to the rest of the states is shown on pages 6 and 7. As can be seen, it ranges from the mid-40's in the south to 50 in the northwest section of the state.

The main geological features affecting the climate in Kansas are the Rocky Mountains in the west and the Gulf of Mexico to the south. The mountains prevent the importation of moisture from the Pacific Ocean while the Gulf is the primary source for most of the precipitation.

Another factor that influences the climate in Kansas is the difference in elevation which changes from about 750 ft. (mean sea level) in the eastern sections to 4000 ft. in some areas in the western sections. The terrain is gently rolling over the north-central sections, is quite level in the central and southern areas and is somewhat hilly along the streams and rivers of eastern Kansas.

The climate is described as humid in the east, moist subhumid in the central section, dry subhumid in the west and semiarid in the southwest.

The two main rivers are the Kansas and the Arkansas which provide drainage and are responsible for much of the fog that exists.

2. Temperature.

The annual mean temperature ranges from about 58 F along the southeastern boarder to 52 F in the extreme northwest. Monthly mean temperatures in the northwest range from 28 F in January to near 78 F in July while in the southeast, the range is from 34 F in January to 81 F in July. As can be seen from the temperature maps in this guide, there are near linear temperature gradients during most of the year across Kansas although there is a change in isopleth directionality depending on the season. Extreme variations are given later in this guide.

3. Insolation.

In direct contrast to the near linear decrease in precipitation across the State from east to west, a near linear increase of insolation intensity and duration exists from east to west in Kansas. The mean annual duration of sunshine is shown on pages 20 and 21. The combination of little precipitation and abundant sunshine indicates why the southwestern section is climatically classified as semiarid.

4. Cloudiness.

Statistics show that there are many more cloudy days in Kansas during the winter and spring months as compared to clear days in the summer and fall. There is an average of 140 clear days in the northern half of the State and 130 clear days in the south. Topeka has 154 cloudy days as compared to 104 in Goodland so there are more cloudy days in the eastern than western half. The plots on pages 22 to 24 show the monthly cloudiness at the five major locations. The type of clouds and their frequencies were given in the first ARM guide on continental measurements and will not be repeated in this guide. A copy of that guide can be obtained from Brookhaven if desired.

5. Precipitation.

The thirty year average precipitation measurements given in this guide show a close to linear decrease in amounts from a high of 40 inches a year in the east to 16 inches in the west. The monthly averages of the five major meteorological stations in the State are shown on pages 25 and 26 and indicate similar results as that just stated above. In the winter most of the rain is light but in the summer 24-hour falls of 5 to 10 inches have been recorded in all sections of the State. However, heavier falls are more frequently reported in the eastern than western part of the State.

The average annual snowfall ranges from about 12 inches in the southeast to 25 inches in the northwest. January and February are the months of heaviest snow in eastern Kansas and February and March in the west. The snow cover does not usually remain for long periods of time, however, there have been winters when the ground was covered for several weeks consecutively.

6. Thunderstorms and Hail.

There is an annual average of between 45 to 60 thunderstorms in Kansas. On the average, more occur in the east and south than in the north and west. The map on page 27 shows the distribution of thunderstorms in Kansas while the map on page 28 can be used to compare those results with the rest of the United States and to give the hours when thunderstorms occur.

Hail usually occurs with thunderstorms but is often reported in the vicinity of well defined frontal systems as well. Hail normally falls in narrow bands and varies in size from its usual size of peas to as big as baseballs. Falling hail can do considerable damage as evidenced by the June 23, 1957 hailstorm in Wichita and vicinity where 14 million dollars in damage occurred. It can destroy wheat and corn fields in a short period of time. As a matter of fact, hail causes more than 10 times the damage of tornadoes. Hail damage in Kansas exceeds that of any other state. The map on page 31 shows the mean annual number of days with hail. The map and plot on pages 29 and 30 show the risk ratio of hail as a function of elevation.

7. Wind.

The average monthly wind direction and speed at five locations in Kansas is shown in the Table on page 34. As can be seen, the prevailing wind direction is from the south except in winter, when northerly winds are more frequent. The wind speed is usually higher in the western part of Kansas than in the east. This can be seen in the map on page 35. The annual average wind directions are shown by the wind roses on pages 36 and 37 for January and July.

An interesting wind feature has been observed in western Texas, central Oklahoma, central Kansas and Iowa. It is a "low-level jet", a flow of air at 11-14 m/sec from the SSW to NNE 700 to 900 meters above the surface, with much slower speeds above and to the sides of the main stream. It may be the catalyst for severe hailstorms and tornadoes which occur in the summer and for icestorms that occur in the winter in that area. It is shown on page 38.

8. Drought.

Drought is more than an atmospheric condition. Plant and soil relationships are combined with atmospheric moisture properties to determine drought measurements. W. C. Palmer has devised a drought index which compares the actual difference between observed monthly rainfall and calculated evapotranspiration with the medium value for the particular month using a soil storage value of six inches of available water. This soil moisture can be depleted or replenished.

When the index is plotted against time, as shown on page 39, the slope of the curve indicates the severity of a drought. The plot shows three drought periods in western Kansas.

9. Fog.

Kansas has a relatively large number of fog days. The map on page 40 shows the annual distribution of fog days in the State. As can be seen, there are more fog days in the western part than in the eastern section.

The river system is shown on page 41. Most of the water travels from the higher elevations in the west to the lower elevations in the east. Fog normally forms in the lower valleys but that does not seem to apply to Kansas.

10. Extremes.

Temperatures of 100 F or higher occur on an average of 10 days per year in the east and west section of the State and 15 days in the central part. In some of the hotter summers the number of days increases to over 50 days in the central section. The number of days with 0 F or lower average 2 to 4 days per year in the southeast to 8 to 10 days in the northwest.

Extremes for the State are -40 F on February 13, 1905 in Lebanon to 121 F on July 18, 1936 at Fredonia.

Freezing temperatures (32 F) have been recorded somewhere in Kansas all months of the year. The July freezing temperature was reported at Tribune on July 18, 1888 during a severe hailstorm. White frost and ice build up on windows occurred during an invasion of cold air on August 26, 1910.

Thunderstorms can be violent at times spawning tornadoes. 39 tornadoes were reported on June 4, 1955 in various locations in the western two-thirds of the State. There are normally 8 to 16 tornadoes reported in Kansas each year. They have been observed in all months except January and December. May and June each have a greater number than any other month, while about twice as many occur in the east than in the west.

Strong winds are responsible for raising sand and dust into the troposphere. Soil conservation practices have stabilized much of the Great Plains, which, decimated by decreased precipitation during the mid-30's formed the notorious dust bowl. In spite of the stabilization effort, strong winds still create dust storms on occasion.

The following extremes pertain to the five major weather stations in Kansas:

Location	Temp. (F)				Monthly Precip. (in.)			
	High	Date	Low	Date	Max.	Date	Min	Date
Goodland	111	7/40	-26	1/59	10.1	7/85	0	1/33
Wichita	113	7/54	-21	2/82	10.5	5/57	T	1/86
Concordia	109	7/80	-17	1/85	14.1	6/67	0	1/86
Dodge City	109	7/86	-15	2/75	9.13	7/62	0	1/67
Topeka	110	8/84	-23	2/79	15.2	6/67	T	12/54

Continentality

KANSAS

Iselines of equal continentality---Interpolated from Frewarth

Isolines of equal continentality. (From TREWARTHA, 1961.)

[illegible]

Isolines are drawn through points of approximately equal value. Caution should be used in interpolating on these maps.

Mean Temperature (°F.), January

KANSAS

Isotherms are drawn using some interpolation.

[illegible]

Isolines are drawn through points of approximately equal value. Caution should be used in interpolating on these maps.

[illegible]

Iso-lines are drawn through points of approximately equal value. Caution should be used in interpolating on these maps.

Mean Temperature (°F.), July

KANSAS

Isotherms are drawn using some interpolation.

[illegible]

Isolines are drawn through points of approximately equal value. Caution should be used in interpolating on these maps.

Wichita Kansas

30 Year Averages

Dodge City Kansas

30 Year Averages

Topeka Kansas

30 Year Averages

Concordia Kansas Temperatures

30 Year Averages

Goodland Kansas

30 Year Averages

KANSAS

Mean annual duration of sunshine, hundreds of hours, 1931-1961
Some interpolation used to determine this map.

Mean annual duration of sunshine, in hundreds of hours, based on black-bulb sunshine recorders at 125 stations, 1931-1960, and 50 for fewer years. (After U.S. Weather Bureau National Atlas map.)

Kansas Cloudiness

Goodland - 68 years

Kansas Cloudiness

Dodge City - 46 years

Kansas Cloudiness

Wichita—35 years

Kansas Cloudiness

Topeka—42 years

Kansas Cloudiness

Concordia--26 years

[illegible]

Based on period 1931-55

isobars are drawn through points of approximately equal value. Caution should be used in interpolating on these maps.

Kansas Precipitation

30 Year Averages

KANSAS

Mean annual number of Thunderstorms. (1955-1981)
Isopleths are interpolated.

Mean annual number of days with thunderstorms, based on data from 261 stations, 1953-1962.
(Prepared by Robert Maples.)

Hour (local standard time) of maximum frequency of thunderstorms, June-August. (After RASMUSSEN, 1971.)

—*The Increase of Hail Risk with Elevation in Kansas. Dots show proportional increase of loss ratio with respect to the rise in elevation from eastern to western Kansas. As the loss ratio increases, rates for hail insurance must of necessity increase accordingly.*

(Diagram by the Crop-Hail Insurance Actuarial Association, Chicago, Illinois.)

Mean annual incidence of large hail (>19 mm) per 26,000 km², 1955-1967. (Prepared by M.L. Swift from data of PAUTZ, 1969.)

Mean annual incidence of tornadoes per 26,000 km², 1955-1967. (Prepared by M.L. Swift from data of PAUTZ, 1969.)

Tracks of violent tornadoes, 1930-1974 [after Fujita, 1976b].

Tracks of strong tornadoes, 1930-1974 [after Fujita, 1976b].

Monthly Average Wind Direction and Speed in Kansas

	Concordia 26		Dodge City 46		Goodland 40		Topeka 39		Wichita 35	
	Dir.	Sp.	Dir.	Sp.	Dir.	Sp.	Dir.	Sp.	Dir.	Sp.
Jan	N	12.2	S	13.5	WSW	12.4	NNW	10.1	S	12.2
Feb	N	12.5	N	14.0	NW	12.5	N	10.4	N	12.7
Mar	S	13.9	N	15.7	NW	14.2	NW	12.2	S	14.1
Apr	S	14.1	N	15.6	SSE	14.6	S	12.1	S	14.1
May	S	12.3	S	14.6	SSE	13.5	S	10.6	S	12.4
Jun	S	11.9	S	14.3	S	12.7	S	9.9	S	12.1
Jul	S	11.6	S	13.1	SSE	11.9	S	8.6	S	11.2
Aug	S	11.4	S	12.8	SSE	11.6	S	8.5	S	11.1
Sep	S	11.6	S	13.7	S	12.1	S	8.8	S	11.6
Oct	S	11.8	S	13.5	SSE	11.8	S	9.2	S	11.9
Nov	S	11.9	N	13.7	NNW	11.9	S	10.0	S	12.2
Dec	S	12.0	S	13.5	WSW	11.9	S	10.0	NNW	12.1

wind speed in miles/hour

number below each city=years of data

KANSAS

Average annual wind speed (mph) From Comparative Climate Data-NOAA-1988
Interpolated results.

January Wind Rose-Ave. percent frequency-1951-1960 (From Slusser, 1960.)

July Wind Rose-Ave. percent frequency=1951-1960 (From Slusser, 1961.)

Location of axis of low-level jet in 26 of 28 cases, 1959-1960. Star indicates median latitude and longitude of all 28 cases, centers of which are shown by dots. Land elevations above sea level shown by solid lines. (After BONNER, 1968.)

Droughts in Western Kansas. (After Palmer)

Annual average heavy fog days. (From *Climates of the States-Vol. 2*)
Interpolated results.

KANSAS

CLIMATIC TABLE FOR CONCORDIA, KANS.

Latitude 39°33'N, longitude 97°39'W, elevation 448 m (424 m)

Month	Mean sta. press. (mbar)	Temperature (°C)				Mean vap. press. (mbar)	Precipitation (mm)		Mean snowfall (cm)
		daily mean	mean daily range	extremes			mean	max. in 24 h	
				max.	min.				
Jan.	968.7	-2.2	10.6	26	-32	4	15	44	11
Feb.	968.0	0.5	11.1	27	-32	4	22	53	13
Mar.	965.4	5.3	11.8	36	-24	5	34	52	14
Apr.	964.6	12.2	12.1	38	-10	8	57	78	3
May	964.0	17.3	11.6	39	-4	13	92	164	tr.
June	962.9	23.3	11.8	43	6	19	107	126	0
July	965.1	26.7	12.5	46	8	21	86	145	0
Aug.	965.2	25.8	12.4	47	5	20	81	121	0
Sept.	966.4	20.8	12.6	44	-2	14	62	117	0
Oct.	967.3	14.3	12.8	39	-9	10	44	111	1
Nov.	968.1	5.6	11.5	28	-26	6	29	52	4
Dec.	968.3	-0.1	10.2	24	-26	4	15	53	8
Annual	966.2	12.5	11.7	47	-32	11	644	164	54
Rec.(yrs.)	30	30	30	76	76	10	30	76	76

Month	Number of days with			Mean cloud- iness (tenths)	Mean sun- shine (h)	Wind	
	precip. (>0.25 mm)	thunder- storm	fog			most freq. direct.	mean speed (m/sec)
Jan.	5	•	2	5.5	60	N	3
Feb.	6	•	2	5.8	60	N	4
Mar.	7	1	1	6.0	62	S	4
Apr.	9	4	1	5.9	63	S	4
May	11	7	•	5.9	66	S	4
June	10	9	•	5.0	73	S	4
July	8	8	•	4.5	78	S	3
Aug.	9	8	•	4.2	77	S	3
Sept.	7	5	•	3.9	72	S	3
Oct.	6	2	1	4.0	70	S	3
Nov.	4	1	1	4.7	64	S	3
Dec.	5	•	2	5.2	58	S	3
Annual	87	45	10	5.1	68	S	4
Rec.(yrs.)	76	76	18	17	53	75	76

References

1. *The World Book Encyclopedia*. 1964 edition.
2. *Physical Climatology*, H. Landsberg. Gray Printing Co., DuBois, PA. 1958.
3. *Comparative Climatic Data for the United States*. NOAA publication. (through 1988).
4. *Climates of the States*. Vol. 2. NOAA 1974.
5. Palmer, W.C. (1956): Drought in Western Kansas; *Weekly Weather and Crop*, Bull. 43, 7-8.
6. Bonner, W.D., 1968. Climatology of the low-level jet. *Mon. Weather Rev.*, 99:833-850.
7. *World WeatherDisc*. Version 1.0, Sept. 1989.
8. *World Survey of Climatology*. H. Landsberg, Vol. 11 Elsevier Scientific Publishing Co., 1974.
9. *Hailstorms of the United States*. S. D. Flora, University of Oklahoma Press: Norman, Oklahoma 1956.

**DATE
FILMED
314192**

I