



# Fermi National Accelerator Laboratory

FERMILAB-Pub.-84/110-A  
October, 1984

## COSMOLOGICAL CONSTRAINTS ON THE PROPERTIES OF WEAKLY INTERACTING MASSIVE PARTICLES

Gary Steigman  
Bartol Research Foundation  
University of Delaware, Newark, DE 19716

and

Michael S. Turner  
Enrico Fermi Institute, The University of Chicago  
Chicago, IL 60637  
and  
Theoretical Astrophysics, Fermi National Accelerator Laboratory  
Batavia, IL 60510

### Abstract

Considerations of the age and density of, as well as the evolution of structure in, the Universe lead to constraints on the masses and lifetimes of weakly interacting massive particles (WIMPs).

## 1. Introduction

The Universe provides a valuable laboratory for the study of elementary particle physics. During its early evolution particles, some of whose existence is yet to be confirmed or ruled out by experiments at terrestrial accelerators, were copiously abundant. Such particles may have survived in sufficient numbers to influence the subsequent evolution of the Universe and, may have played a significant role in the development of structure in it. Data on the present age, density and large scale structure of the Universe can be marshalled to provide interesting constraints on the abundances, masses and lifetimes of such relics. The purpose of this paper is to derive constraints based on these considerations.

To date, much progress has been made in constraining the masses and abundances (related to the interaction strengths) of stable or long-lived particles through the requirement that their present mass density not exceed that observed (see, e.g., ref. [1]). Unstable relics whose decay products include photons and/or electrically charged particles, have their lifetimes, as well as their masses and abundances, constrained directly by observations of the background radiation fields (see, e.g., ref. [2]) and, indirectly by considerations of stellar structure and evolution, (see, e.g., ref. [3]) and by primordial nucleosynthesis (see, e.g., ref. [4]).

It might seem that unstable particles whose decay products interact only feebly (i.e., participate in interactions which are as weak, or weaker than, the weak interaction) might have disappeared without a trace. However, such particles and their invisible decay products may have at one time dominated the energy density of the Universe and, therefore, controlled the evolution of the Universe during a crucial epoch. In a natural extension of previous

analyses we use the requirements that the present Universe be neither too young (or, equivalently too dense) and that the observed large scale structure should have been able to evolve in it to derive constraints on the masses, abundances and lifetimes of unstable, weakly interacting massive particles (WIMPs) whose decay products are "invisible" (i.e. interactionless) and sufficiently light so that for all epochs of interest they are ultrarelativistic. In the recent literature there are numerous examples of models in which particles (usually neutrinos) are relatively long-lived ( $\sim 10^2 - 10^{10}$  yr) and decay into invisible, relativistic daughter particles, for example:  $\nu\nu^* \rightarrow \nu\nu^*$  family,  $\nu\nu^* \rightarrow \nu\nu^*$  or  $\nu\nu^* \rightarrow \text{Majoron}$  (Dicus *et al* [2],[5]).

In the next section we consider the evolution of the density of a massive WIMP,  $X$ , and of the density of its relativistic decay products,  $R$ . When these densities are compared with those of the relic photons and neutrinos,  $\nu$ , and of the other stable non-relativistic particles (i.e., baryons, etc.),  $NR$ , interesting epochs during the evolution of the Universe are identified. With this as background, we proceed in the following section to derive constraints on the mass, abundance and lifetime of the  $X$ . In a concluding section we summarize our results.

### 1. The Evolution of WIMPs and Their Decay Products

Consider a particle  $X$  whose mass is  $M_X$  and whose abundance (relative to relic photons) is  $n_X$ . The lifetime of this particle is  $t_0$  ( $\equiv \Gamma^{-1}$ ;  $\Gamma$  = decay width). [Note,  $t_0/10^9$  yr =  $2.08 \times 10^{-41}$  GeV/  $\Gamma$ ]. Prior to decay the energy density in  $X$ s is,

$$\rho_X = M_X n_X n_Y = M_X n_X n_Y R^{-3} \quad (1a)$$

In equation (1a),  $n_Y$  is the number density of relic photons and  $R$  ( $\geq 1$ ) is the cosmic scale factor normalized so that, at present,  $R_0=1$ . Throughout, the present epoch is denoted by the subscript zero. If  $\theta$  is the present

microwave temperature in units of 2.7K, then  $n_{\gamma 0} = 399 \text{ e}^3 \text{ cm}^{-3}$ . For  $M_X$  measured in eV,

$$\rho_X = 399(M_X n_X \text{ e}^3) R^{-3} \text{ (eVcm}^{-3}\text{)}. \quad (1b)$$

If  $X$  were one of the "usual" (e,  $\mu$  or  $\tau$ ) neutrinos then,

$n_X = n_\nu = 3/4(T_\nu/T_\gamma)^3 = 3/11$ . It is useful to introduce  $\tilde{M}_X \equiv M_X(n_X/n_\nu)\text{e}^3$ , the mass of an equivalent neutrino in a Universe with  $T_{\gamma 0} = 2.7\text{K}$ . With this definition,

$$\rho_X = 109\tilde{M}_X R^{-3} \text{ (eVcm}^{-3}\text{)}. \quad (1c)$$

It is convenient to compare all densities today with the present critical (Einstein-deSitter) density  $\rho_c$ :

$$\rho_c = \frac{3H_0^2}{8\pi G} = 1.05 \times 10^4 h_0^2 \text{ (eVcm}^{-3}\text{)}, \quad (2)$$

where the present value of the Hubble parameter is  $H_0 = 100 h_0$  ( $\text{kms}^{-1} \text{Mpc}^{-1}$ ). From (1c) and (2) it follows that,

$$\tilde{M}_X = 96.8 (\Omega_X h_0^2) \text{ eV}. \quad (3)$$

At the epoch when  $t = t_0 = r^{-1}$  and  $R = R_0$ , we assume that all the  $X$ s decay simultaneously. After decay, the energy density of their relativistic decay products ( $R$ ) is,

$$\rho_R = 1.05 \times 10^4 (\Omega_R h_0^2) R^{-4} \text{ (eVcm}^{-3}\text{)}, \quad (4)$$

and since  $\rho_X = \rho_R$  at  $R=R_0$ ,  $\Omega_R = \Omega_X R_0$  or,

$$\tilde{M}_X R_0 = 96.8 (\Omega_R h_0^2) \text{ (eV)}. \quad (5)$$

[Note that the errors made by assuming that all the decays occur simultaneously are 10-20%; see ref. [6]].

In addition to the products of  $X$ -decay, there is of course also the relativistic background of relic photons and (light) neutrinos:

$$\rho_{\gamma\nu} = \rho_\gamma + \rho_\nu = A_{\gamma\nu} \rho_\gamma = 0.25 A_{\gamma\nu} \text{ e}^4 R^{-4} \text{ (eVcm}^{-3}\text{)}, \quad (6a)$$

where

$$A_{\gamma\nu} = 1 + \frac{7}{8} (T_\nu/T_\gamma)^4/3 N_\nu = 1.68 + 0.227(N_\nu - 3). \quad (6b)$$

In equation (6b),  $N_\nu$  is the number of relativistic 2-component neutrinos. Comparing  $\rho_{\gamma\nu}$  with  $\rho_c$  we find,

$$\Omega_{\gamma\nu} h_0^2 = 2.4 \times 10^{-5} A_{\gamma\nu} \text{ e}^4. \quad (7)$$

Finally, there will be a background of non-relativistic (NR) particles such as baryons and, possibly, other stable massive relics (and/or those massive products of  $X$  decay which have become NR):

$$\rho_{NR} = 1.05 \times 10^4 (\Omega_{NR} h_0^2) R^{-3} \text{ (eVcm}^{-3}\text{)}. \quad (8)$$

In the absence of WIMPs and their decay products, the Universe evolves from early "radiation-domination" to "matter-domination" at the epoch  $R=R_{eq}$  where,

$$R_{eq} = \frac{\rho_{\gamma\nu}}{\rho_{NR}} = 2.4 \times 10^{-5} A_{\gamma\nu} \theta^4 (\Omega_{NR} h_0^2)^{-1}. \quad (9)$$

In the presence of WIMPs and their decay products, there are several other epochs of interest. First, compare  $\rho_X$  with  $\rho_{NR}$  and call the ratio  $x$ ,

$$x = \frac{\rho_X}{\rho_{NR}} = \frac{\bar{M}_X}{96.3 \Omega_{NR} h_0^2} = \left(\frac{R}{R_D}\right) \frac{1}{R_D}. \quad (10)$$

If  $x < 1$ , neither the  $X$ s nor the  $R$ s ever dominate the dynamical evolution of the Universe. In this case the  $X$  is "cosmologically safe" (with respect to the issues considered in this paper), and we will not be able to place any constraints on its properties. In the more interesting case that  $x > 1$ , the Universe will become  $X$ -dominated at  $R=R_X$  where,

$$R_X = x^{-1} R_{eq}. \quad (11)$$

provided that the  $X$ s do not decay first (i.e.,  $R_X < R_D$ ); see Figures 1 and 2. If, however,  $R_D < R_X$ , the  $X$ s will decay prior to  $X$ -domination and, once again, neither the WIMPs nor their decay products are ever dynamically significant; see Figure 3. This case too is "cosmologically safe". Returning to the more interesting case where  $x > 1$  and  $R_X > R_D$ , it is seen (Figures 1 and 2) that the period of  $X$ -domination lasts until the  $X$ s decay at  $R=R_D$ .

Thereafter, the Universe will be  $R$ -dominated until  $R=R_{NR}$  where,

$$R_{NR} = x R_D = \frac{R}{\Omega_{NR}}. \quad (12)$$

Notice that if  $\Omega_{NR} > \Omega_{NR}$ , the present Universe is radiation-dominated(!); this is the case considered by Turner, Steigman and Krauss ([7], hereafter TSK) in an attempt to reconcile an Einstein-deSitter Universe ( $\Omega_0=1$ ) with the observational constraint that the matter associated with the visible galaxies contributes less than a third of that required. This case is illustrated in Figure 1.

For  $\Omega_{NR} < \Omega_{NR}$ , the present Universe is matter-dominated; see Figure 2. Although this is the "standard" assumption, it must be emphasized that the baryon density is small:  $\Omega_B \approx 0.14-0.19$  (ref. [4]), so that if  $\Omega_{NR} = 1$ , the non-relativistic matter is dominated by something other than baryons which, in addition, must be more smoothly distributed than the luminous matter already observed.

We have, in the above, expressed the epoch of  $X$ -decay by the scale factor  $R_D$ . We are, however, interested in constraining the lifetime  $t_D$ . In relating  $t_D$  to  $R_D$ , there are three cases to be considered. If the Universe is  $X$ -dominated at decay ( $x > 1$ ,  $R_X < R_D$ ), then  $6\pi G_0 \rho_{X0} t_D^2 = 1$  and,

$$t_D = \frac{6.5 \times 10^9 R_D^2 \text{ yr.}}{(\Omega_{NR} h_0^2)^{1/2}} = \frac{6.5 \times 10^9 R_D^{3/2} \text{ yr.}}{(\bar{M}_X/96.3 \text{ eV})^{1/2}} \quad (13)$$

If, the Universe is  $\gamma\nu$ -dominated at decay ( $x > 1$ ,  $R_D < R_X$ ), then  $(32\pi/3)G_0 \rho_{\gamma\nu 0} t_D^2 = 1$  and,

$$t_D = \frac{4.9 \times 10^9 R_D^2 \text{ yr.}}{(\Omega_{\gamma\nu} h_0^2)^{1/2}} = \frac{10^{12} R_D^2 \text{ yr.}}{(A_{\gamma\nu} \theta^4)^{1/2}} \quad (14)$$

Finally, in the less interesting case ( $x < 1$ ) where the Xs decay when the Universe is NR-dominated,  $6 \times G_0 \rho_{NR} t_D^2 = 1$  and,

$$t_D = \frac{6.5 \times 10^9 \rho_{NR}^{3/2}}{(\rho_{NR} h_0^2)^{1/2}} \text{ yr.} \quad (15)$$

Using the relations derived in this section we now proceed, through various cosmological considerations, to find constraints on  $\bar{M}_X$  and  $t_D$ .

### III. Constraints On X Masses and Lifetimes

Although relic WIMPs may have decayed long ago and their decay products may be "invisible", nonetheless the Xs and/or the Rs could have played a dynamically important role in the evolution of the Universe. If so, data on the age, density and observed structure of the present Universe can be used to constrain the masses ( $M_X$ ), abundances ( $n_X$ ) and lifetimes ( $t_D$ ) of such particles. The results to be obtained in this section are summarized in Figure 4 which displays the  $t_D$  vs.  $\bar{M}_X$  plane; the regions to the left and below the various lines are not excluded by cosmological considerations.

(A)  $t_D \geq t_0$ : In this case the stable or long-lived Xs are still present today. Such relics must not contribute excessively to the present mass density ( $\rho_X \leq \rho_0$ ). From equation (3) this constraint leads to an upper limit to  $\bar{M}_X$ .

$$\bar{M}_X(A) \leq 96.8 [\text{Max}(\rho_0 h_0^2)]^{1/2} (\text{eV}). \quad (16)$$

Current data suggests  $h_0 \leq 1$  (refs. [8-10]) and does not exclude a value as small as  $h_0 = 0.4$  (ref. [10]). The density associated with luminous matter is small:  $\rho = 0.1-0.3$  (refs. [11-14]). However, "theoretical prejudice" (i.e.: the "naturalness" of the Einstein-deSitter model and the very attractive "inflationary" Universe scenarios) strongly suggest that  $\rho_0 = 1$ . Measurements of the deceleration parameter  $q_0$ , imply that  $\rho_0 \leq \text{few}$  [15]. In setting an upper limit to the combination  $\rho_0 h_0^2$  it is, it must be emphasized, not permissible to choose  $\rho_0 \geq 1$  and  $h_0 = 1$ . The reason is that such a combination corresponds to an exceedingly youthful Universe:  $t_0 \leq 6.5 \times 10^9$  yr. The age of a matter-dominated Universe  $t_0 = 10^{10} h_0^{-1} f(\rho_0)$  yr., where  $f(\rho)$  is monotonically decreasing, from  $f(0) = 1$  to  $f(1) = 2/3$ . Asymptotically,  $f^2(\rho)$  approaches  $(\pi/2)^2$  from below, which implies that, independent of  $h_0$  and  $\rho_0$ ,  $\rho_0 h_0^2 \leq (\pi/2)^2 (t_0 / 10^{10} \text{ yrs})^{-2}$ . Various determinations of the age of the Universe suggest a lower bound [16-17] on the age:  $t_0 \geq (10-13) \times 10^9$  yr., which then restricts  $\rho_0 h_0^2$ :

$$\rho_0 h_0^2 \leq 1.5-2.5. \quad (17a)$$

Restricting  $h_0$  to be  $\geq 1/2$  and again taking  $t_0 \geq (10-13) \times 10^9$  yr results in the tighter constraint:

$$\rho_0 h_0^2 \leq 0.25 - 0.75 \quad (17b)$$

In preparing Figure 4 and, in subsequent numerical comparisons, we have used

$\text{Max}(\rho_0 h_0^2) = 1/4$  so that the limit in equation (16) becomes:

$$\bar{M}_X(A) \leq 24 \text{ eV.}$$

(B)  $\omega_R > \omega_{NR}$ : This is the case, considered by ISK, in which the present Universe is R-dominated. If the energy density in the Rs is too large, again the Universe will be too young. From equation (5) we see that,

$$\tilde{M}_\chi(B)R_D \leq 96.8 [\text{Max}(\omega_0 h_0^2)](eV). \quad (18)$$

In this case,  $\chi$ -decay has occurred when the Universe was  $\chi$ -dominated (see Fig. 1) so that the  $t_D$  vs.  $R_D$  relation is given by equation (13). It follows then that,

$$\tilde{M}_\chi(B) \leq 1/8 [\text{Max}(\omega_0 h_0^2)]^{3/4} (10^{10} \text{ yr}/t_D)^{1/2}. \quad (19)$$

For the same values of  $\omega_0$  and  $h_0$ , a radiation-dominated Universe is younger than a matter-dominated Universe,

$$RD: t_D = \frac{0.98 \times 10^{10} \text{ yr.}}{h_0 + (\omega_0 h_0^2)^{1/2}} \quad (20)$$

For  $t_D \geq 10 \times 10^9$  yr. and  $h_0 \geq 0.4-0.5$ ,  $\omega_0 h_0^2 \leq 0.23-0.34$ , whereas for  $t_D \geq 13 \times 10^9$  yr. and  $h_0 \geq 0.4-0.5$ ,  $\omega_0 h_0^2 \leq 0.06-0.12$ .

Roughly speaking then,

$$\tilde{M}_\chi(B) \approx \tilde{M}_\chi(A)(t_D/t_0)^{1/2}, \quad (21)$$

where in Fig. 4 and throughout the paper  $t_0$  is taken to be  $\sim 10^{10}$  yr. This mass-lifetime constraint, for a neutrino which decays into relativistic particles, was first discussed by Dicus et al [2].

(C)  $x \leq 1$ : If  $\nu_\chi \leq \rho_{NR}$  then neither the  $\chi$ s nor their decay products ever significantly affect the dynamical evolution of the Universe (since  $\omega_\chi \ll \omega_R$ ). This will be the case for  $\chi$ s whose mass is less than,

$$\tilde{M}_\chi(C) = 96.8 [\text{Max}(\omega_{NR} h_0^2)] \cdot M_\chi(A). \quad (22)$$

This is a "cosmologically safe" case in the sense that although it may be possible for  $M_\chi$  to exceed  $\tilde{M}_\chi(C)$  without violating any observational data,  $M_\chi \leq \tilde{M}_\chi(C)$  is not forbidden by any consideration in this paper.

(D)  $R_D < R_\chi$ : Even if  $x > 1$  so that early on  $\omega_\chi > \omega_R$ , the  $\chi$ s may decay while the Universe is  $\nu$ -dominated (see Fig. 3). Once again, neither the  $\chi$ s nor their decay products ever significantly influence the dynamical evolution of the Universe. This, too, is a "cosmologically safe" case, defined in the  $M_\chi$ - $t_D$  plane by:

$$M_\chi(D) \approx 0.023 (\omega_{\nu 0}^4)^{3/4} (10^{10} \text{ yr.}/t_D)^{1/2}, \quad (23a)$$

$$\text{or } M_\chi(D) = 10^{-3} M_\chi(A)(t_D/t_0)^{1/2} = 10^{-3} \tilde{M}_\chi(B). \quad (23b)$$

Of the cases considered to this point, (A), (C) and (D) have been "cosmologically safe" in the sense that if  $M_\chi \leq M_\chi(A)$  for  $t_D \geq t_0$  or, if  $M_\chi \leq \tilde{M}_\chi(C)$  for  $10^{-6} \leq t_D/t_0 \leq 1$  or, if  $\tilde{M}_\chi \leq M_\chi(D)$  for  $t_D \leq 10^{-6} t_0$ , neither the  $\chi$ s nor the Rs ever dominate the energy density of the Universe. In contrast, case (B) is potentially dangerous in the sense that (for  $t_D \geq t_0$ ) even for masses somewhat smaller than  $M_\chi(B)$ , there will be an epoch during which the Universe is R-dominated (see Fig. 1). It is such




FIGURE 1




FIGURE 2




FIGURE 3




FIGURE 4



12.

recent epochs during which the observed large scale structure (galaxies, clusters, etc.) of the Universe is generally believed to have formed. By requiring that such structure be able to evolve from small density perturbations in an otherwise homogeneous Universe, we will find limits to  $\bar{M}_X$  which are more constraining than those found in case B above. Although these new constraints will be weaker than those of cases C and D, our analysis will permit us to deal with a region of the mass-lifetime plane which cannot easily be excluded by "cosmologically safe" considerations alone.

Since we will be using "the formation of structure in the Universe" to constrain  $\bar{M}_X$  and  $t_D$ , we should first briefly outline the general picture of how structure is believed to have evolved. [For a more detailed discussion we refer the reader to ref. [18]]. The basic idea is that the structure we see today grew up from small density inhomogeneities via the Jeans or gravitational instability. The initial "spectrum" of density inhomogeneities is specified by the amplitudes of the various Fourier components of the density contrast:  $\delta\rho/\rho = \sum_k \delta_k e^{ikx}$ . A particular component is labeled by its comoving wavenumber  $k$  or comoving wavelength  $\lambda = 2\pi/k$ , which are related to the "proper" (or physically measureable) wavenumber and wavelength by:  $\lambda_{phys} = R\lambda$  or  $k_{phys} = k/R$ . Note at the present epoch  $\lambda_{phys} = \lambda$  and  $k_{phys} = k$ . The density contrast on a given scale is related to the Fourier component by:  $(\delta\rho/\rho)^2 = k^3 |\delta_k|^2$ . A convenient way to specify the initial spectrum is to specify the amplitude,  $(\delta\rho/\rho)_H$ , on a given scale when the scale "entered the horizon", i.e., when  $\lambda_{phys} = ct$ . For example, the scale-invariant Zeldovich spectrum predicted in inflationary models is characterized by  $(\delta\rho/\rho)_H = \text{constant}$  [19-20]. Once inside the horizon, linear density perturbations grow as:  $\delta\rho/\rho = R^n$  ( $n = 1$  - matter domination;  $n = 0$  - radiation domination) -- thus the total amount by which a linear perturbation grows is determined by the factor by which the cosmic scale factor has grown during the epoch(s) of matter-domination. For reference,  $\lambda = 1$  Mpc corresponds

to a galactic size ( $\approx 10^{12} M_\odot$ ) perturbation. Studies of the galaxy-galaxy correlation function [21] indicate that the scale which is just going non-linear today is:  $\lambda_c = 5h_0^{-1}$  Mpc. Density perturbations also result in anisotropies in the cosmic  $\mu$ -wave background radiation [22-25], and thus measurements of the isotropy of the  $\mu$ -wave background provide valuable and stringent constraints on the initial spectrum of density perturbations. While the precise constraints depend upon the scale dependence of the initial spectrum, for our purposes, we will use the constraint that  $(\delta\rho/\rho)_H$  must be  $\leq 0(10^{-3}-10^{-4})$  to be consistent with the measured smoothness of the  $\mu$ -wave background [22-25]. We believe this to be a very conservative bound -- in all likelihood the constraint is much more restrictive. Finally, we note that we are restricting ourselves to adiabatic (as opposed to isothermal) initial perturbations as they are strongly favored in light of the fact that  $\rho_0$  appears to be  $\gg \rho_B$ , the fact that isothermal perturbations are all but inconsistent with scenarios of baryogenesis, and the fact that the perturbations predicted in inflationary scenarios are adiabatic. Allowing for the possibility of isothermal perturbations would lessen only slightly the constraints which we will now discuss.

(E)  $R_{NR} \leq 10^{-3}$ : If  $R_X < R_D$ , there will be an epoch of X-domination followed at  $R=R_D$  by an epoch of R-domination (see Figure 2). Then, at  $R=R_{NR}=xR_D$ , the Universe becomes NR-dominated. If  $R_{NR} \leq 10^{-3}$ , then perturbations in the linear regime will grow by  $\geq 10^3$  from  $R=R_{NR}$  up to the present epoch. By the standards we outlined above this may be just barely enough growth to account for the observed large scale structure. For this case to apply, we first must have  $R_X = x^{-1}R_{eq} < R_D$  so that  $R_{eq} < xR_D = R_{NR} \leq 10^{-3}$ . This condition,  $R_{eq} \leq 10^{-3}$ , requires that  $\Omega_{NR} h^2 \geq 0.024 (A_{\nu\nu} e^4)$ . From equations (10) and (13),

14.

$$\tilde{M}_X(E) = 78 [(xR_D)(\Omega_{NR} h_0^2)]^{3/4} (10^{10} \text{ yr.}/t_0)^{1/2}. \quad (24)$$

For  $\Omega_{NR} h_0^2 \leq 1/4$  and  $R_{NR} = xR_D \leq 10^{-3}$  we have,

$$\tilde{M}_X(E) = 6.4 \times 10^{-3} \tilde{M}_X(A) (t_0/t_D)^{1/2}. \quad (24a)$$

or

$$\tilde{M}_X(E) = 6.4 \tilde{M}_X(D). \quad (24b)$$

(F)  $R_X < R_D < 1$ : In this case there are two epochs during which the Universe is matter-dominated and during which perturbations may grow. Perturbations which enter the horizon before or at  $R=R_X$  will grow by a factor  $R_D/R_X$  until the Universe becomes R-dominated at  $R=R_D$ . From  $R=R_D$  until  $R=R_{NR}$  the Universe is radiation-dominated and perturbations grow very little. When  $R=R_{NR}$  the Universe becomes matter-dominated again and perturbations resume their growth up to the present epoch. It is clear (see Fig. 2) that the total growth factor is

$$\gamma = \frac{R_D}{R_X} \left( \frac{1}{R_{NR}} \right) = \frac{1}{R_{eq}} = \frac{4.2 \times 10^4 \Omega_{NR} h_0^2}{(A_{\gamma\nu} \theta^4)}. \quad (25)$$

Notice that the growth factor  $\gamma$  is independent of  $R_D$ , in fact independent of whether or not there were relic Xs. Now, as TSK have emphasized, we must make sure that the "proper" scales achieve sufficient growth. Perturbations which entered the Universe before X-domination have a present size  $\lambda \leq \lambda_X$  where

$$\lambda_X = 53 (A_{\gamma\nu} \theta^4)^{1/2} (24 \text{ eV}/M_X) \text{ Mpc}. \quad (26)$$

[In TSK,  $N_\nu = 2$ , corresponding to  $A_{\gamma\nu} = 1.45$  was chosen.] Larger scales ( $\lambda > \lambda_X$ ) enter the horizon at a later epoch ( $R > R_X$ ) and undergo less growth, by a factor  $(\lambda_X/\lambda)^2$ . As discussed earlier, the scale  $\lambda_c = 5h_0^{-1}$  Mpc is just entering the non-linear regime today. If  $\tilde{M}_X \geq 600h_0 \text{ eV}$ , then  $\lambda_X \leq \lambda_c/2$  and  $\gamma(\lambda_c) \leq \gamma/4$ . Therefore, to insure that initial perturbations of amplitude  $\leq 10^{-3} - 10^{-4}$  (on a scale of  $5h_0^{-1}$  Mpc) undergo sufficient growth, we require that  $\tilde{M}_X \leq \tilde{M}_X(F)$  where,

$$\tilde{M}_X(F) = 25h_0^{-1} \tilde{M}_X(A). \quad (27)$$

#### IV. Summary

The results of our considerations are illustrated in Figure 4. The regions in the  $t_0$  vs.  $\tilde{M}_X$  plane to the right of and above lines A and B are excluded by considerations of the present universal mass density. Here we have adopted  $\Omega_D h_0^2 < 1/4$  so that,

- (i)  $\tilde{M}_X < \tilde{M}_X(A) = 24 \text{ eV}$  for  $t_0 > t_0$  and,
- (ii)  $\tilde{M}_X < \tilde{M}_X(B) = \tilde{M}_X(A) (t_0/t_D)^{1/2}$  for  $t_0 < t_0$ .

Note that for  $\tilde{M}_X = \tilde{M}_X(B)$  the present Universe is radiation-dominated and  $\Omega_D = 1$ . For  $t_0 \geq 10^{-2} t_0$ , this case is not inconsistent with the observed large scale structure (TSK). For  $t_0 \leq 10^{-2} t_0$ , the observed large scale structure excludes case (B); consistency requires that,

(iii)  $\bar{M}_X < \bar{M}_X(F) = 25h_0^{-1}M_X(A)$  for  $t_D \leq 10^{-2}t_0$ . If  $\bar{M}_X = \bar{M}_X(F)$ , then as  $t_D$  decreases, so does  $R_{NR}$ ; that is, the epoch of NR-domination occurs earlier. If  $R_{NR} \leq 10^{-3}$  then perturbations may achieve enough growth to account for the observed large scale structure. Therefore, for  $t_D < 4 \times 10^{-7} t_0$ , we only have the weaker requirement that,

$$(iv) \quad \bar{M}_X < \bar{M}_X(E) = 6.4 \times 10^{-3} \bar{M}_X(A) (t_0/t_D)^{1/2}.$$

The regions below and to the left of curves E and F in Figure 4 may be permitted. However, it is quite clear that if for  $10^{-6}t_0 \leq t_D \leq t_0$ ,

$$(v) \quad \bar{M}_X < \bar{M}_X(C) = \bar{M}_X(A), \text{ or for } t_D < 10^{-6} t_0,$$

$$(vi) \quad \bar{M}_X < \bar{M}_X(D) = 10^{-3} \bar{M}_X(A) (t_0/t_D)^{1/2},$$

then neither the Xs nor their decay products ever dominate the universal mass density. No cosmological considerations (discussed here) exclude the regions to the left and below curves C and D – these regions are cosmologically safe. The analysis in this paper allows the region bounded by curves CDEF. However, more careful consideration of the formation of structure in the Universe (for example, detailed analysis of the predicted microwave fluctuations, and numerical simulations of these scenarios) are likely to exclude this region, leaving curves C and D as the relevant constraints..

[In a recent paper which primarily concerned itself with the viability of a neutrino-dominated Universe, Hut and White [26] have indirectly discussed one of the mass-lifetime constraints (case E) we have derived here.]

#### Acknowledgements

This work was supported at Bartol by DOE grant DE-AC02-78ER-05007 and by the DOE at the University of Chicago and at Fermilab and by NASA at Fermilab. Some of this work was done while GS was a guest of the Theoretical Astrophysics Group at Fermilab and while both of us were at the Aspen Center for Physics.

#### REFERENCES

- [1] R. Cowsik and J. McClelland, *Phys. Rev. Lett.* **29**(1972)669; G. Marx and A Szalay, *Proceedings Neutrino '72*, **1**(1972)123; S. Gershtein and Ya. B. Zeldovich, *JETP Lett.* **4**(1966)174.
- [2] D. Dicus, E. Kolb, and V. Teplitz, *Astrophys. J.* **221**(1978)327; J. Gunn, B. Lee, i. Lerche, O. Schramm, and G. Steigman, *Astrophys. J.* **223**(1978)1015; M. S. Turner, *Proceedings Neutrino '81*, **1**(1981)95.
- [3] K. Sato and H. Sato, *Prog. Theor. Phys.* **54**(1975)1564; D. Dicus, E. Kolb, V. Teplitz, and R. Wagoner, *Phys. Rev.* **D18**(1978)1829 and **D22**(1980)839.
- [4] J. Yang, M. Turner, G. Steigman, O. Schramm, and K. Olive, *Astrophys. J.* **281**(1984)493.
- [5] F. Wilczek, *Phys. Rev. Lett.* **49** (1982) 1549; D. Reiss, *Phys. Lett.* **115B** (1982) 217; Y. Chikashide, R. Mohapatra and R. Peccei, *Phys. Rev. Lett.* **45** (1980) 1926; G. Gelmini and J. Valle, *Phys. Lett. B*, in press (1984).
- [6] M. S. Turner, "Cosmology with Decaying Particles", submitted to *Phys. Rev. D.*(1984).
- [7] M. Turner, G. Steigman, and L. Krauss, *Phys. Rev. Lett.* **52**(1984)2090.
- [8] M. Aaronson and J. Mould, *Astrophys. J.* **265**(1983)1; R. Buta and G. DeVaucouleurs, *Astrophys. J.* **266**(1983)1.
- [9] A. Sandage and G. Tammann, *Astrophys. J.* **256**(1982)339, and *Nature* **307** (1984) 326.
- [10] O. Branch *et al.*, *Astrophys. J.* **270**(1983)123.

- [11] S. Faber and J. Gallagher, *Ann. Rev. Astron. Astrophys.* 17(1979)135.
- [12] W. Press and M. Davis, *Astrophys. J.* 259(1982)449.
- [13] M. Davis and P.J.E. Peebles, *Ann. Rev. Astron. Astrophys.* 21(1983)109.
- [14] *Physical Cosmology*, eds. R. Balian, J. Audouze, and D. Schramm (North-Holland, Amsterdam, 1980).
- [15] J. Gunn and B. Oke, *Astrophys. J.* 195(1975)255;  
J. Kristian, A. Sandage and J. Westphal, *Astrophys. J.* 221(1978)333.
- [16] D. N. Schramm, in *Highlights of Astronomy* as presented to the 18th General Assembly of the IAU, vol. 6, ed. R. West (Reidel, Dordrecht, 1983).
- [17] I. Iben, Jr., *Ann. Rev. Astron. Astrophys.* 12(1974)215; A. Sandage, *Astrophys. J.* 252(1982)553.
- [18] P.J.E. Peebles, *The Large-Scale Structure of the Universe* (Princeton Univ. Press, Princeton, 1980).
- [19] S. W. Hawking, *Phys. Lett.* 115B(1982)295;  
A. A. Starobinsky, *Phys. Lett.* 117B(1982)175.
- [20] A. Guth and S. Y. Pi, *Phys. Rev. Lett.* 49(1982)1110; J. Bardeen, P. Steinhardt and M. S. Turner, *Phys. Rev.* 028(1983)679.
- [21] M. Davis and P.J.E. Peebles, *Astrophys. J.* 267(1983)465.
- [22] J. Silk and M. Wilson, *Astrophys. J.* 244(1981)L37.
- [23] P.J.E. Peebles, *Astrophys. J.* 263(1982)L1.
- [24] J. Bond and G. Efsthathiou, *Astrophys. J.* (1984) in press.
- [25] N. Vittorio and J. Silk, *Astrophys. J.* (1984) in press.
- [26] P. Hut and S. White, *Nature* 310(1984) 637.

## Figure Captions

Figure 1. The evolution of the densities of the several constituents of the Universe: X particles (X), their relativistic decay products (R), photons and neutrinos ( $\gamma\nu$ ), and stable, nonrelativistic particles (NR). The Universe becomes X-dominated at  $R_X$ ; the Xs decay at  $R_D$ ; at the present ( $R=1$ ), the Universe is R-dominated ( $R > R_{NR}$ ).

Figure 2. The evolution of the densities in the case that the present epoch of NR-domination has been preceded by epochs of R- and X-domination. The current epoch of NR-domination begins at  $R_{NR}$ . Linear density perturbations can grow significantly only when the Universe is matter dominated, here for  $R_X < R < R_D$  and for  $R > R_{NR}$ .

Figure 3. The evolution of the densities in the case that the Xs decay before they can dominate the Universe ( $R_D < R_X$ ).

Figure 4. The lifetime ( $t_D$ ) versus mass plane;  $\tilde{M}_X = M_X(n_X/n_\nu)\theta^3$  where  $n_X = n_X/n_\nu$  and  $n_\nu = n_\nu/n_\nu = 3/11$ . The areas to the left of and below the various curves (see the text) are permitted. The heavily shaded part of curve B is the  $R_D = 1$ ,  $R_D$  model proposed by TSK. To the left of and below lines C and D are cosmologically safe, while the region bounded by lines C, D, E, F is, at present, permitted (more careful considerations are likely to exclude this region as well).