Institute for Fundamental Theory Preprint UFIFT

DOE/ER/40272--130

DE92 008773

IFT HEP-91-11

1992

May 1991

Construction of Lax operators from weight diagrams†

Steven L. Carbon and Eric J. Piard

*Institute for Fundamental Theory
Department of Physics
University of Florida, Gainesville, FL 32611*

Abstract: We show that cyclic weight diagrams corresponding to representations of affine Lie algebras allow one to read-off the associated Lax operator. The resultant Lax pair generates the modified KdV equations, and have been shown in some cases to produce acceptable solutions of the string equation of matrix models.

MASTER

† Supported in part by U.S. Department of Energy under contract No. DE-FG05-86ER-40272.

UNIVERSITY OF FLORIDA

Gainesville, Florida 32611

Recently, matrix models have received a great deal of attention as nonperturbative descriptions of string theory^{1,2,3}. Since initial advancements, progress has proceeded in many different directions. In particular, Douglas⁴ has shown that the limited number of known matrix model solutions can be derived from the Lax pair formalism usually associated with the KdV equations. This identification with integrable systems greatly increases the number of classifiable matrix models since it was shown a long time ago⁵ that Lax operators are associated with affine Lie algebras. For example, the models discussed by Douglas are related to the canonical representations of $A_n^{(1)}$. More recently, Di Francesco and Kutasov^{6,7} have discussed $D_n^{(1)}$ based matrix models which the standard matrix techniques^{8,9,10} have yet to solve. Thus, it may be worthwhile to focus on the integrable systems approach.

The matrix construction of Lax operators, reviewed in depth by Drinfeld and Sokolov⁵, first defines the matrix operator

$$\mathcal{L} = \frac{\partial}{\partial x} + \Lambda + q(x). \quad (1)$$

The second term is constructed from the negative simple roots, E_i , of an embedding affine Lie algebra $\hat{\mathcal{G}}$, i.e.,

$$\Lambda = \sum_{i=0}^r c_i E_i, \quad (2)$$

where E_0 is the affine simple root. The coefficients are then assigned values $c_i = 1$ and $c_m = \lambda$, which corresponds to deleting the m' th vertex from the Dynkin diagram. The residual Lie algebra(s) corresponding to the remaining (disconnected) Dynkin diagram(s) is denoted $(\hat{\mathcal{G}}, c_m)$. For use below, let us define the vacuum projection operator Δ^- by

$$\Lambda = \sum_{i=0}^r E_{i \neq m} + \lambda E_m \equiv \Delta^- + \lambda E_m. \quad (2)$$

The procedure is then to reduce the system of linear equations given by the kernel matrix equation

$$\mathcal{L}\vec{\psi}(x) = 0, \quad (3)$$

where $\vec{\psi} = (\psi_1, \psi_2, \dots, \psi_N)$, to the scalar Lax eigenvalue equation

$$L(x)\phi(x) = \mu\phi(x), \quad (4)$$

where μ is a constant function of the spectral parameter λ . The vacuum solution ϕ is a linear combination of the components of the eigenfunction $\vec{\psi}$ which satisfies

$$\Delta^- \vec{\psi}(x) = 0. \quad (5)$$

Drinfeld and Sokolov found that the fact the embedding algebra is affine guarantees the solution generates an integrable system.

For the number of independent degrees of freedom in (3) to equal the rank of the affine Lie algebra $\hat{\mathcal{G}}$, or equivalently the residual basic Lie algebra $(\hat{\mathcal{G}}, c_m)$, one must fix the gauge invariance present in the coordinate dependent term $q(x)$. To derive an appropriate gauge fixed matrix operator one must find a matrix $S(x)$ such that

$$\begin{aligned} \mathcal{L}^{G.F.} &= e^{ad} S \mathcal{L} \\ &= \frac{\partial}{\partial x} + \Lambda + q^{G.F.}(x), \end{aligned} \quad (6)$$

where ad denotes the adjoint mapping. Drinfeld and Sokolov find the sufficient condition that $S \in C^\infty\{\mathbb{R}, \eta\}$, where η is the set of positive simple roots F_i with the m 'th member removed.

Here, we choose the “diagonal” gauge which is defined in terms of the standard gradation of the embedding Lie algebra regardless of which root is removed. For the present discussion we take $(\hat{\mathcal{G}}, c_0)$. The diagonal gauge is convenient since, deleting the zero'th vertex, gives the form

$$q^{diag} = \sum_{i=1}^r v_i(q_1, q_2, \dots, q_N) H_i, \quad (7)$$

where H_i are elements of the Cartan subalgebra. In this gauge, solutions to the kernel equation (3) will create scalar Lax operators which generate the modified KdV equations. On the other hand, if one expresses the coordinate dependent term in the “canonical” gauge of Drinfeld and Sokolov, the Lax operator will generate the regular KdV equations, which are related to the diagonal case *via* the Miura transformation.

Since the diagonal gauge is written directly in terms of the elements of the Cartan subalgebra, there exists a simple diagrammatic technique for constructing these (pseudo)differential operators. Here, we present the algorithm which allows one to build Lax operators by reading off information from cyclic weight diagrams corresponding to representations of the embedding affine Lie algebras. In a longer

paper we prove this ascertion and extend our work, without modification, to affine supersymmetric Lie algebras¹¹.

To motivate the algorithm we first give examples of the Drinfeld and Sokolov construction of $L(x)$ by solving the matrix system $\mathcal{L}\vec{\psi} = 0$. To illustrate, consider the embedding affine Lie alegbra $(A_2^{(1)}, c_0)$. Clearly, the residual Lie algebra is A_2 . Fig. 1 presents the cyclic weight diagram of the canonical rep of $A_2^{(1)}$. Note, eliminating the presence of the affine root results in the highest weight diagram of the fundamental rep of A_2 , with highest weight vector (10) . *i.e.*,

$$\begin{array}{c} (10) \\ \downarrow \\ (-11) \\ \downarrow \\ (0 -1). \end{array} \quad (8)$$

Reading directly from this weight diagram, we immediately write down the sum of the negative simple roots,

$$\Delta^- = \begin{pmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}, \quad (9)$$

and the elements of the Cartan subalgebra

$$H_1 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad H_2 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{pmatrix}. \quad (10)$$

Similarly, by reading off directly from the cyclic weight diagram in Fig. 1, we find the matrix Λ .

Substituting the appropriate matrices into the kernel equation (3) produces the system of equations:

$$\begin{aligned} [\partial + v_1]\psi_1 &= -\lambda\psi_3, \\ [\partial - v_1 + v_2]\psi_2 &= -\psi_1, \\ [\partial - v_2]\psi_3 &= -\psi_2. \end{aligned} \quad (11)$$

Here, on the right-hand-side, we have placed the terms due to Λ . The vacuum condition (5) determines the scalar function to be $\phi = \psi_3$. Thus, we must solve by starting with the last equation. First, we multiply this equation through by $[\partial - v_1 + v_2]$, and then eliminate ψ_2 using the second equation. This gives

$$[\partial - v_1 + v_2][\partial - v_2]\psi_3 = \psi_1. \quad (12)$$

Next, we multiply through by $[\partial + v_1]$ and use the top equation in (11). The final result is the scalar Lax eigenvalue equation

$$\begin{aligned} L^{(A_2^{(1)}, c_0)} \psi_3 &= [\partial + v_1][\partial - v_1 + v_2][\partial - v_3] \psi_3 \\ &= -\lambda \psi_3, \end{aligned} \quad (13)$$

where the spectral parameter is given by $\mu = -\lambda$.

This example exhibits a common feature relevant to our scheme below. When, the vacuum condition (5) requires the scalar eigenfunction to be given by a single component of the eigenfunction, say $\phi = \psi_i$, then the resulting characteristic equation satisfies

$$L\psi_i = \mu\psi_i. \quad (14)$$

Consequently, the system reduction must start with the i 'th equation in the matrix system, and proceed upward till the top equation is reached. If $i \leq N$ the process continues starting with the bottom equation and moves upward until the i 'th equation is reached. We shall refer to this case as trivial since the corresponding cyclic affine weight diagram is linear, containing no branch points. A second feature brought out in this example, is that the number of factors in the resultant Lax operator (13) is equal to the number of weights in the weight diagram. Unfortunately, this is valid only for trivial cases. Nevertheless, this last observation is key to our scheme.

To highlight one more property of the general procedure we turn to a non-trivial example. For this, we require a representation of an affine Lie algebra whose cyclic weight diagram has at least one branching point. Thus, consider the canonical rep of the affine algebra $(D_4^{(1)}, c_0)$ whose cyclic weight diagram, shown in Fig. 2, has two branch points. Reading off from the affine diagram gives

$$\Lambda = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & \lambda & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & \lambda \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \end{pmatrix}. \quad (15)$$

The branch points have manifested themselves by placing more than one non-zero integer in the second and sixth rows. Now, further reading off the elements of the

Cartan matrix gives the system of equations

$$\begin{aligned}
[\partial + v_1]\psi_1 &= -\lambda\psi_7, \\
[\partial - v_1 + v_2]\psi_2 &= -\psi_1 - \lambda\psi_8, \\
[\partial - v_2 + v_3 + v_4]\psi_3 &= -\psi_2, \\
[\partial - v_3 + v_4]\psi_4 &= -\psi_3, \\
[\partial + v_3 - v_4]\psi_5 &= -\psi_3, \\
[\partial + v_2 - v_3 - v_4]\psi_6 &= -\psi_4 - \psi_5, \\
[\partial + v_1 - v_2]\psi_7 &= -\psi_6, \\
[\partial - v_1]\psi_8 &= -\psi_7.
\end{aligned} \tag{16}$$

The vacuum condition (5) produces two distinct solutions, ψ_8 and the linear combination $\psi_4 - \psi_5$. Here, we consider the first case. Proceeding as before, we encounter a well-known technical problem not found in the trivial case. The components ψ_4 and ψ_5 can not both be simultaneously eliminated since the expressions $[\partial - v_3 + v_4]$ and $[\partial + v_3 - v_4]$ do not commute. This dilemma is directly linked to the fact, the corresponding cyclic weight diagram has a branch point connecting the fourth and fifth weights to a single weight located below them.

To overcome this obstacle the pseudo-differential operator ∂^{-1} must be introduced. Its operation on any function $f(x)$ is given by the expansion

$$\partial_x^{-1} f(x) = \sum_{i=0}^{\infty} (-1)^i f^{(i)}(x) \partial_x^{-1-i}. \tag{17}$$

Utilizing the pseudo-differential operator we rewrite the fourth equation in (16) as

$$\psi_4 = -[\partial - v_3 + v_4]^{-1} \psi_3, \tag{18}$$

and a similar expression for $\psi_5(x)$. Thus

$$\psi_4 + \psi_5 = -\{[\partial - v_3 + v_4]^{-1} + [\partial + v_3 - v_4]^{-1}\} \psi_3. \tag{19}$$

A helpful identity for differential operators A and B is

$$\{A^{-1} + B^{-1}\}^{-1} = \{A^{-1}[A + B]B^{-1}\}^{-1} = B[A + B]^{-1}A. \tag{20}$$

When applied to Eq.(19), a cancellation occurs among the v_i 's appearing in the curly brackets, giving

$$\psi_3 = -[\partial - v_3 + v_4]\partial^{-1}[\partial + v_3 - v_4](\psi_4 + \psi_5). \tag{21}$$

Finally, the Lax operator of $(D_4^{(1)}, c_0)$ with vacuum ψ_8 is given by

$$L' = \frac{1}{4} \partial^{-1} [\partial + v_1] [\partial - v_1 + v_2] [\partial - v_2 + v_3 + v_4] [\partial - v_3 + v_4] \times \partial^{-1} [\partial + v_3 - v_4] [\partial + v_2 - v_3 - v_4] [\partial + v_1 - v_2] [\partial - v_1], \quad (22)$$

which is proportional to the standard result.

We have chosen these two examples because they introduce the techniques needed to generate Lax operators associated with even the most complicated algebraic systems. Furthermore, they show how closely the structure of cyclic affine weight diagrams is linked with the construction of general Lax operators. As a result, we propose a set of four steps which allows one to construct Lax operators associated with cyclic representations of affine Lie algebra.

First, we propose that to every weight vector of an affine cyclic weight diagram one can associate an operator as follows:

$$\text{Step 1: } (ab \dots) \rightarrow [\partial_x + av_1(x) + bv_2(x) + \dots]. \quad (23)$$

Even before we specify how to build $L(x)$ from these factors, we can draw a significant conclusion. Namely, due to the fact we are dealing with scale invariant theories, *e.g.*, KdV equations or matrix models, two Lax operators $L(x)$ and $L'(x)$ will be equivalent if their respective cyclic weight diagrams have the same shape and number of weight vectors, and if there exists a mapping between the coefficients of the weight factors.

Second, we introduce a step which is designed to facilitate the construction of Lax operators when branch points exist in the corresponding weight diagram. Essentially, this step reduces more complicated non-trivial cases to a sum of manageable trivial cases by reducing the branched weight diagram to a sum of linear subdiagrams.

$$\begin{aligned} \text{Step 2: } & \text{Replace branching weight diagrams by the sum} \\ & \text{of linear subdiagrams, each representing a vertical route} \\ & \text{beginning with, and ending on, the vacuum weight}(s). \end{aligned} \quad (24)$$

For example, Fig. 3 presents the four linear subgraphs associated with the canonical rep of $D_4^{(1)}$.

In drawing cyclic weight diagrams, it is important that the arrows generated by the deleted vertex are distinguished from the others. Furthermore, the directions

of the arrows must also be noted. The Lax operator associated with a subdiagram is then constructed as follows:

Step 3: Circulate around the loop beginning with the vacuum solution, in the direction opposite to that indicated by the deleted vertex's arrow. If a weight vector is approached by an arrow's

(25)

a) tail, append its weight factor to the operator's left side,

b) head, append the inverse of its weight factor to the operator's left side.

For the arrow of the deleted vertex, do nothing.

For trivial cases, this completes the computation of $L(x)$. However, for non-trivial cases with branching weight diagrams we can not naively build the final Lax operator from a sum of its constituent linear subgraphs. Instead, as we prove in Ref.[11], they are added together analogously to how one computes total resistance of resistors in parallel.

Step 4: The Lax operator is given by the inverse of the sum of the inverses, of the operators resulting from step three.

(26)

Example: For the non-trivial case $(D_4^{(1)}, c_0)$, with $\phi = \psi_8$ as before, the operator corresponding to the first linear subdiagram in Fig. 3 is

$$L_1 = [\partial - v_1]^{-1} [\partial + v_1] [\partial - v_1 + v_2] [\partial - v_2 + v_3 + v_4] [\partial - v_3 + v_4] \times [\partial + v_2 - v_3 - v_4] [\partial + v_1 - v_2] [\partial - v_1], \quad (27)$$

with a similar expression for L_2 . The operators for the remaining subdiagrams are easily found along the lines of $A_2^{(1)}$. Factoring out common terms, and simplifying, the reciprocal of the sum

$$(L')^{-1} = \sum_{i=1}^4 (L_i)^{-1} \quad (28)$$

reproduces the previous expression given in Eq.(22).

Finally, we address the question of integrability of Lax pair equations generated by non-canonical reps of affine Lie algebras. Drinfeld and Sokolov found that the infinite set of conserved currents, required for integrability of the affine algebra generated Lax operator, are given by the coefficients of the Laurent expansion of $\mathcal{L}$ in λ . Thus, every rep of an affine Lie algebra leads to an integrable system since the defining properties of affine Lie algebras are representation independent.

ACKNOWLEDGMENTS

We wish to thank Zongan Qiu and Moustafa Awada for discussions, and Pierre Ramond for reading over the manuscript.

REFERENCES

1. M. Douglas and S. Shenker, Nucl. Phys. **B335**, 635 (1990).
2. E. Brezin and V.A. Kazakov, Phys. Lett. **B236**, 144 (1990).
3. D. Gross and A.A. Migdal, Nucl. Phys. **B340**, 333 (1990).
4. M. Douglas, Phys. Lett. **B238**, 176 (1990).
5. V. Drinfeld and V. Sokolov, J. Sov. Math. 30 (1984) 1975.
6. P. Di Francesco and D. Kutasov, Nucl. Phys. **B342**, 589 (1990).
7. P. Di Francesco and D. Kutasov, Princeton preprint PUPT-1206 (1990).
8. D. Bessis, C. Itzykson and J.-B. Zuber, Adv. Appl. Math. **1**, 109 (1980).
9. M. L. Mehta, Comm. Math. Phys. **79**, 327 (1981).
10. S. Chadha, G. Mahoux and M. L. Mehta, J. Phys **A14**, 579 (1981).
11. S. L. Carbon and E. J. Piard, UF preprint in preparation.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

FIGURE CAPTIONS

Figure 1. Cyclic weight diagram for canonical rep of $(A_2^{(1)}, c_0)$. Dashed line indicates arrow due to deleted vertex c_0 .

Figure 2. Cyclic weight diagram for canonical rep of $(D_3^{(1)}, c_0)$. Dashed line indicates arrow due to deleted vertex c_0 .

Figure 3. Linear subdiagrams from the cyclic weight diagram of the canonical rep of $(D_3^{(1)}, c_0)$. Dashed lines indicate arrows due to deleted vertex c_0 .

Figure 1

Figure 2

Figure 3

END

**DATE
FILMED**

4/9/92

1