55
1/2/92 JS(2)
PREPARED FOR THE U.S. DEPARTMENT OF ENERGY,
UNDER CONTRACT DE-AC02-76-CHO-3073

PPPL-2816
UC-420,426

PPPL-2816

INVESTIGATION OF GLOBAL ALFVÉN INSTABILITIES
IN TFTR

BY

K.L. WONG, R. DURST, R.J. FONCK, AND S.F. PAUL

January 1992

PPPL

PRINCETON
PLASMA PHYSICS
LABORATORY

NOTICE

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial produce, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

NOTICE

This report has been reproduced directly from the best available copy.

Available to DOE and DOE contractors from the:

Office of Scientific and Technical Information
P.O. Box 62
Oak Ridge, TN 37831;
Prices available from (615) 576-8401.

Available to the public from the:

National Technical Information Service
U.S. Department of Commerce
5285 Port Royal Road
Springfield, Virginia 22161
703-487-4650

INVESTIGATION OF GLOBAL ALFVEN INSTABILITIES IN TFTR

by

K. L. Wong, R. Durst*, R. J. Fonck*, S. F. Paul, D. R. Roberts+, E. D. Fredrickson, R. Nazikian, H. K. Park, M. Bell, N. L. Bretz, R. Budny, C. Z. Cheng, S. Cohen, G. W. Hammett, F. C. Jobes, L. Johnson, D. M. Meade, S. S. Medley, D. Mueller, Y. Nagayama, D. K. Owens, S. Sabbagh#, and E. J. Synakowski

Princeton PLasma Physics Laboratory,
Princeton University, Princeton, New Jersey 08543.

Abstract

Toroidal Alfvén Eigenmodes (TAE) were excited by the energetic neutral beam ions tangentially injected into TFTR plasmas at low magnetic field such that the injection velocities were comparable to the Alfvén speed. The modes were identified by measurements from Mirnov coils and beam emission spectroscopy (BES). TAE modes appear in bursts whose repetition rate increases with beam power. The neutron emission rate exhibits sawtooth-like behavior and the crashes always coincide with TAE bursts. This indicates ejection of fast ions from the plasma until these modes are stabilized. The dynamics of growth and stabilization was investigated at various plasma current and magnetic field. The results indicate that the instability can effectively clamp the number of energetic ions in the plasma. The observed instability threshold is discussed in the light of recent theories. In addition to these TAE modes, intermittent oscillations at three times the fundamental TAE frequency were observed by Mirnov coils, but no corresponding signal was found in BES. It appears that these high frequency oscillations do not have direct effect on the plasma neutron source strength.

MASTER
EP

1. INTRODUCTION

In an ignited d-t tokamak fusion reactor, the 3.5 MeV alpha particles produced from the fusion reactions are expected to be born with velocities faster than the Alfvén speed, so they can interact strongly with shear Alfvén waves¹. In tokamaks, toroidal Alfvén eigenmodes² (TAE) can go unstable easily because their frequencies lie within the frequency gap between two Alfvén continuum bands thus avoiding continuum damping. This instability can eject the fast alpha particles from the tokamak before they thermalize with the bulk plasma³. It is also important to reactors which rely on energetic neutral beam current drive, like the International Thermonuclear Experimental Reactor (ITER), which plans to employ 1.3 MeV neutral beams to drive the plasma current. Substantial reduction in current drive efficiency can occur if the beam particles are expelled by excited TAE modes.

In typical tokamak parameters, the Alfvén velocity $V_A = B/(\mu_0 n_i m_i)^{1/2}$ is very high; its magnitude is comparable to the velocity of MeV ions, e.g., in a 5 tesla magnetic field, deuterium density of 10^{20} m^{-3} gives $V_A = 7.7 \times 10^6 \text{ m/s}$ which corresponds to the velocity of 1.2 MeV alpha particles. Therefore, it was generally perceived that one could only investigate this instability in an ignited tokamak. Careful consideration of tokamak operation characteristics revealed that it might be possible to simulate this type of alpha particle physics⁴ with existing neutral beams. In order to excite TAE modes with neutral beam injection, it is necessary to have the beam injection velocity V_b comparable to V_A . The Alfvén velocity can be reduced by lowering the magnetic field and raising the plasma density. The maximum density achievable in a tokamak is approximately

proportional to the magnetic field. The Murakami parameter⁵, defined as $n_e R/B_T$, has an upper bound which is limited by disruption. Its value is machine dependent, and it varies with methods of plasma fueling and auxiliary heating. In TFTR with neutral beam heating and gas fueling, we can have $n_e R/B_T$ up to $6 \times 10^{19} \text{ m}^{-2} \text{ tesla}^{-1}$. The tangentially injected beam ions are passing particles in the tokamak. We prefer the orbit shift to be less than half the minor radius so that they are reasonably well confined, i.e., $\Delta \sim 2q\rho < a/2$. Some of the beam particles may be scattered into trapped particles before they are slowed down, and we prefer to have their banana width $\Delta_b \sim 2q\rho\epsilon^{-1/2} < a/2$ in order to avoid strong localized heating on the limiter. This is more restrictive than the previous condition on passing particles.

With all these constraints taken into account, Fig.1 shows a small operating region for TAE excitation in TFTR. We have assumed 100keV deuterium beam ions ($V_b = 3.1 \times 10^6 \text{ m/s}$) in a deuterium plasma with $q(a)=2.3$. Within the small area with $B \sim 1 \text{ tesla}$, $n_i \sim 2.5 \times 10^{19} \text{ m}^{-3}$, we have $V_b \geq V_A$, $n_e R/B_T \leq 6 \times 10^{19} \text{ m}^{-2} \text{ tesla}^{-1}$ and $\Delta_b \leq a/2$. We believe that this is a feasible operating point to excite TAE modes. It should be pointed out that the boundaries of this region are "soft" boundaries. TAE modes can be excited with V_b slightly below V_A , and sometimes we can have plasmas with $n_e R/B_T$ slightly above $6 \times 10^{19} \text{ m}^{-2} \text{ tesla}^{-1}$. In our experiment, we found that the plasma disrupted too often at $q(a)=2.3$, and we operate mostly at $q(a)>2.7$. Nevertheless, Fig.1 shows that we can expect to see TAE modes only in a very small operating parameter space, and there is no doubt about where to look for them.

2. MODE IDENTIFICATION

TAE modes have the following characteristics:

- a. Excitation requires $V_b \sim V_A$.
- b. The mode frequency is approximately $\omega \approx V_A/2qR$ where V_A and q are taken at the location at which the mode has maximum amplitude.
- c. Each eigenmode consists of many poloidal harmonics⁶ and the dominant m -th harmonic changes with q approximately following the relationship $m \sim nq$.

In our experiment, up to 14 MW of deuterium neutral beams at energies below 110 keV were injected into plasmas with the following parameters: $B \geq 1$ tesla, $q(a) \geq 2.7$, $R_0 = 2.4\text{m}$, $a = 0.75\text{m}$, $n_e \sim 3 \times 10^{19} \text{ m}^{-3}$, $T_e(0) \sim 2\text{keV}$, $\langle \beta_p \rangle \sim 1$. Bursts of coherent oscillations were observed by Mirnov coils and beam emission spectroscopy⁷ (BES). These oscillations have the characteristics of TAE modes as summarized in the following:

- a. With $q(a)$ fixed at 2.8 and $\langle \beta_p \rangle$ kept around 1, we raised the toroidal field from 1 tesla to 2 tesla. V_b/V_A dropped from 1 to 0.6 and those oscillations disappeared.
- b. By varying the plasma density and magnetic field, we managed to change V_A over a factor of 2 and the oscillation frequency followed the theoretical prediction $\omega \approx V_A/2qR$ over the entire range with $q = 1.3$. Doppler shift in frequency was minimized by nearly balanced neutral beam injection.
- c. The mode structure was measured by BES. Cross-phase spectra showed a radial standing wave structure with reasonably good phase coherence between signals from various locations. The coherence is less than

perfect, and its variation corresponds to the mode amplitude profile.

This indicates that a coherent global mode structure is superimposed on some incoherent fluctuations. Poloidal mode number measurements indicate that the dominant poloidal harmonic number m increases with minor radius while $m \sim nq$ holds only approximately. As expected, the direction of poloidal propagation is the same as the ion diamagnetic drift velocity.

In our experiment, $\langle \beta \rangle \sim 1\%$, about half the Troyon limit. One shot has been analysed for ideal ballooning instability. It was found that the plasma is everywhere stable against ideal ballooning modes, except near the very edge of the plasma. Based on the above results, we conclude that these are TAE modes. Some of these experimental data have been published recently⁸. Concurrent experiments⁹ on the DIII-D tokamak also produced evidence of TAE modes under similar conditions.

3. EJECTION OF ENERGETIC PARTICLES

The most significant consequence of TAE instability is the ejection of energetic particles by these modes. Evidence of this can be seen from Fig.2. The neutron emission rate S_N has sawtooth-like behavior and there is very good correlation between the TAE bursts and the drop of neutron emission from the plasma. Since beam-target reaction dominates in our plasma, a decrease in neutron emission rate is a strong indication of energetic beam ion loss. It appears that when the beam ion population reaches a certain level, TAE modes become unstable. They grow to a certain amplitude, eject

some fast ions, probably reduce the energetic ion density gradient at the same time, until they become stable again. This suggests that the beam ion pressure is clamped near the instability threshold. When the neutral beam injection power P_b is raised, the neutron source strength rises faster during the quiescent period. It takes less time to reach the instability threshold, and the TAE bursts appear more frequently with somewhat larger amplitudes. There appears to be a nonlinear relationship between the amplitude of the TAE burst and the change in S_N . Low amplitude TAE bursts seem to have no noticeable effect on S_N . This appears to support the existence of a threshold for orbit stochasticity which can cause significant fast ion loss³. More work is needed to quantify this for comparison with theory. Fig.3 shows the variation of neutron emission rate and central electron density with neutral beam power. They increase faster at higher magnetic field. However, the ratio $S_N/n_e(0)$ is not sensitive to beam power as depicted in Fig.4. When we repeat this with $I_p=530\text{kA}$ at $B_T=1.2$ tesla, $S_N/n_e(0)$ remains insensitive to P_b , but its numerical value is notably higher (by approximately 40%) than that for $I_p=420\text{kA}$.

The above observations can be explained as follows. Let us approximate the deuterium ion distribution function as the sum of a cold Maxwellian distribution (target plasma) and an energetic slowing-down distribution due to neutral beam injection. The neutron production rate from d-d reaction is the sum of beam-target, beam-beam and target-target reactions:

$$S_N = \int dV \{ n_b n_t \langle \sigma v \rangle_{bt} + 1/2 n_b^2 \langle \sigma v \rangle_{bb} + 1/2 n_t^2 \langle \sigma v \rangle_{tt} \} \quad (1)$$

In our experiment, $T_i < 2\text{keV}$ and the thermal reactivity $\langle \sigma v \rangle_{tt}$ is practically zero. The beam-beam neutron rate is estimated to be 20-30% of the total. Therefore, S_N is mainly due to beam-target reactions:

$$S_N \approx \int dV n_b n_t \langle \sigma v \rangle_{bt} \quad (2)$$

We assume that the deuterium density profile is similar to the electron density profile which is flat across the plasma minor radius, and it is insensitive to beam power. Furthermore, we assume that the slowing-down distribution $f_b(v)$ for the beam ions is also insensitive to beam power. This is substantiated by the fact that the electron temperature and the effective charge number are not sensitive to beam power, and it leads to the result that $\langle \sigma v \rangle_{bt}$ is independent of beam power. Therefore,

$$S_N/n_e(0) \propto S_N/n_t(0) \propto \int dV n_b = N_b \quad (3)$$

where N_b is the total number of beam ions in the plasma. The fact that $S_N/n_e(0)$ is independent of beam power indicates that N_b is clamped by the TAE instability. It is interesting to note that with $I_p=420\text{kA}$, $S_N/n_e(0)$ has similar values at $B_T=1$ tesla and 1.2 tesla. The value becomes 10% higher when I_p was raised to 530kA, possibly due to better fast ion confinement at higher current. We should point out that the uncertainty on $S_N/n_e(0)$ is about +/-15%, and a 40% difference is barely outside the error bars. More careful analysis is needed before we can come to a definite conclusion.

4. INSTABILITY THRESHOLD

As noted in our previous paper, the observed instability threshold is an order of magnitude higher than that predicted by Fu and Van Dam¹⁰ in which electron Landau damping due to the passing electrons is assumed to be the dominant damping mechanism. If we include the effect of the anisotropic beam-ion velocity distribution function¹¹, the disagreement is even larger. It was pointed out recently¹² that trapped electrons near the trapped/passing boundary can play an important role. The collisional damping rate due to trapped electrons is given as¹²:

$$\gamma/\omega = -2.1 \{ \ln[8(2\varepsilon\omega/\nu_e)^{1/2}] \}^{1/2} \beta_e (\nu_e/\omega)^{1/2} \quad (4)$$

where ε is the inverse aspect ratio and ν_e is the electron collision frequency. γ/ω is proportional to $(\nu_e/\omega)^{1/2}$ rather than ν_e/ω due to the electrons in the trapped/passing boundary layer¹³. For our experimental parameters, this damping rate is an order of magnitude larger than the electron Landau damping calculated by Fu and Van Dam¹⁰. Continuum damping¹⁴⁻¹⁶ would also raise the threshold substantially. It depends on the details of $q(r)$ which we cannot quantify at this moment. Ion Landau damping¹⁷ due to the sideband resonance at $v_{||} = V_A/3$ is also considered. It is not important for this experiment because of the low β plasma.

In our experiment, the $n=2$ mode dominates at $B=1$ tesla while the $n=3$ mode becomes dominant at $B=1.2$ tesla. This feature was qualitatively reproduced by the NOVA-K code⁶. We use the plasma pressure and q profiles obtained from transport analysis, and find that the $n=2$ mode has a lower

threshold at $B=1$ tesla, while the $n=3$ mode has a lower threshold at $B=1.2$ tesla. However, the absolute magnitude of the instability threshold is still an order of magnitude below the experimental value. Incorporation of collisional damping and continuum damping into NOVA-K code will be carried out soon.

5. HIGH FREQUENCY OSCILLATIONS

The TAE bursts appear very frequently at $\omega = \omega_A$ during neutral beam injection into plasmas at low magnetic field. They can be detected by Mirnov coils and BES. Intermittent oscillations at a higher frequency $\omega \approx 3\omega_A$ are also observed by Mirnov coils, but there is no corresponding signal in BES although its frequency bandwidth is wide enough for these oscillations. Fig.5 shows the amplitude modulation of these high frequency oscillations in the Mirnov coil signal. The period of modulation approximately equals the sawtooth period, but there appears to be a time lag between the sawtooth crash and the appearance of these oscillations which, unlike the TAE modes, have no instantaneous effect on the neutron source strength. As we know, there is a third order frequency gap in the Alfvén continuum at the frequency $3\omega_A$ and it is tempting to relate that to these high frequency oscillations. The width of this gap is very narrow (proportional to ϵ^3), but the temporal variation of $q(r)$ due to sawteeth activity may cause the gap to appear periodically which permits these intermittent oscillations. However, we have to point out that oscillations with similar appearance have been observed by Mirnov coils in plasmas with high magnetic field. More analysis and measurements of mode

numbers are necessary for their identification.

6. SUMMARY AND DISCUSSIONS

We have demonstrated that TAE modes can be excited by fast ions with velocities comparable to the Alfvén speed. The TAE modes can effectively clamp the number of fast particles in the plasma, and therefore present an obvious danger to fusion reactors. Fortunately, the instability threshold is not as low as originally predicted, and it may be possible to operate a fusion reactor without exciting them. It is very important to find out the instability threshold in terms of plasma parameters. In the TFTR experiment, TAE modes can be excited only in a very small operating region, and the plasma is not well diagnosed because of the low magnetic field. Many of the plasma parameters are outside the design range of the existing diagnostics. Further experimental and theoretical studies are necessary to provide better understanding of this instability. If one can drive these modes with an external antenna and study their damping rates, it can provide valuable information on the instability threshold. In case the instability cannot be avoided in a reactor, we may have to find a way to stabilize it. Feedback stabilization may work for low- n modes, but probably will not be effective for high- n modes.

ACKNOWLEDGEMENTS

The authors would like to thank the neutral beam group and the TFTR staff for their technical support which makes this experiment possible.

Discussions with many members of the theory group are also very helpful. This project is supported by USDOE under contract number DE-AC02-76-CHO-3073.

*Permanent address: Department of Nuclear Engineering, University of Wisconsin, Madison, Wisconsin 53706.

+Permanent address: The University of Texas, Austin, Texas 78712.

#Permanent address: Columbia University, New York, New York 10027.

REFERENCES

1. M. N. Rosenbluth and P. H. Rutherford, Phys. Rev. Lett. **34** (1975), 1428.
2. C. Z. Cheng, Liu Chen, and M. S. Chance, Ann. Phys. (N.Y.) **161** (1985), 21;
also C. Z. Cheng and M. S. Chance, Phys. Fluids **29** (1986), 3695.
3. D. J. Sigmar, Collection of Papers Presented at the IAEA Technical
Committee Meeting on Alpha Particles in Fusion Research at Aspenas,
Sweden (June 10-14, 1991), Vol. II, p.624. Submitted to Phys. Fluids B.
4. K. L. Wong, Bull. Am. Phys. Soc. **33** (1988), 2098.
5. M. Murakami, J. D. Callen, L. A. Berry, Nucl. Fusion **16** (1976), 347.
6. C. Z. Cheng, Phys. Fluids B **3** (1991), 2463.
7. R. J. Fonck, P. A. Dupperex, and S. F. Paul, Rev. Sci. Instrum. **61** (1990),
3487.
8. K. L. Wong, R. J. Fonck, S. F. Paul, D. R. Roberts, E. D. Fredrickson, R.
Nazikian, H. K. Park, M. Bell, N. L. Bretz, R. Budny, S. Cohen, G. W. Hammett,
F. C. Jobes, D. M. Meade, S. S. Medley, D. Mueller, Y. Nagayama, D. K. Owens,
and E. J. Synakowski, Phys. Rev. Lett. **66** (1991), 1874.
9. W. Heidbrink, E. J. Strait, E. Doyle, G. Sager, R. T. Snider, Nucl. Fusion
31 (1991), 1635.
10. G. Y. Fu and J. W. Van Dam, Phys. Fluids B **1** (1989), 1949.
11. H. L. Berk, B. N. Breizman, and Huanchun Ye, IFSR#507(July 1991),
Institute for Fusion Studies, The University of Texas at Austin.
12. N. N. Gorelenkov and S. E. Sharapov, Collection of Papers Presented at
the IAEA Technical Committee Meeting on Alpha Particles in Fusion
Research at Aspenas, Sweden (June 10-14, 1991), Vol. II, p738.
13. M. N. Rosenbluth, D. W. Ross, and D. P. Kostomarov, Nucl. Fusion **12**

(1972), 3; also V. A. Mazur, A. B. Mikhajlovskij, Nucl. Fusion **17** (1977), 193.

14. M. N. Rosenbluth, H. L. Berk, D. M. Lindberg, and J. W. Van Dam, IFSR#518 (August 1991), Institute for Fusion Studies, The University of Texas at Austin. Submitted to Phys. Rev. Lett.
15. F. Zonca and L. Chen, Bull. Am. Phys. Soc. **35** (1990), 2069. Submitted to Phys. Rev. Lett.
16. H. L. Berk, J. W. Van Dam, Z. Guo, D. M. Lindberg, IFSR#499, Institute for Fusion Studies, The University of Texas at Austin. Submitted to Phys. Fluids B.
17. R. Betti and J. P. Friedberg, Collection of Papers Presented at the IAEA Technical Committee Meeting on Alpha Particles in Fusion Research at Aspenas, Sweden (June 10-14,1991), Vol. II, p.667.

Figure Captions

Figure 1. Determination of operating point for the TAE experiment.

Figure 2. Correlation between the drop in neutron source strength and TAE bursts.

Figure 3. Variation of neutron source strength and central electron density with neutral beam power.

Figure 4. The neutron emission rate normalized by the central electron density is not sensitive to beam power and magnetic field.

Figure 5. Intermittent oscillations observed by Mirnov coils at frequency $3\omega_A$.

Fig. 1

Fig. 2

▲ Neutron Source Strength at $B_T = 1$ Tesla

△ Neutron Source Strength at $B_T = 1.2$ Tesla

● $n_e(0)$ at $B_T = 1$ Tesla

○ $n_e(0)$ at $B_T = 1.2$ Tesla

PPPL #91X0595

Fig. 3

▲ $S_N / n_e(0)$ at $B_T = 1$ Telsa

△ $S_N / n_e(0)$ at $B_T = 1.2$ Telsa

$I_p = 420\text{kA}$

PPPL #91x0597

Fig. 4

Shot 54741, $B_T = 1$ Tesla, $I_p = 420$ kA, $P_b = 14$ MW

PPPL#91X0503

Fig. 5

EXTERNAL DISTRIBUTION IN ADDITION TO UC-420

Dr. F. Paoloni, Univ. of Wollongong, AUSTRALIA	Prof. I. Kawakami, Hiroshima Univ., JAPAN
Prof. M.H. Brennan, Univ. of Sydney, AUSTRALIA	Prof. K. Nishikawa, Hiroshima Univ., JAPAN
Plasma Research Lab., Australian Nat. Univ., AUSTRALIA	Director, Japan Atomic Energy Research Inst., JAPAN
Prof. I.R. Jones, Flinders Univ, AUSTRALIA	Prof. S. Itoh, Kyushu Univ., JAPAN
Prof. F. Cap, Inst. for Theoretical Physics, AUSTRIA	Research Info. Ctr., National Instit. for Fusion Science, JAPAN
Prof. M. Heindler, Institut für Theoretische Physik, AUSTRIA	Prof. S. Tanaka, Kyoto Univ., JAPAN
Prof. M. Goossens, Astronomisch Instituut, BELGIUM	Library, Kyoto Univ., JAPAN
Ecole Royale Militaire, Lab. de Phy. Plasmas, BELGIUM	Prof. N. Inoue, Univ. of Tokyo, JAPAN
Commission-European, DG. XII-Fusion Prog., BELGIUM	Secretary, Plasma Section, Electrotechnical Lab., JAPAN
Prof. R. Bouciqué, Rijksuniversiteit Gent, BELGIUM	S. Mori, Technical Advisor, JAERI, JAPAN
Dr. P.H. Sakanaka, Instituto Fisica, BRAZIL	Dr. O. Mitarai, Kumamoto Inst. of Technology, JAPAN
Instituto Nacional De Pesquisas Especiais-INPE, BRAZIL	J. Hyeon-Sook, Korea Atomic Energy Research Inst., KOREA
Documents Office, Atomic Energy of Canada Ltd., CANADA	D.I. Choi, The Korea Adv. Inst. of Sci. & Tech., KOREA
Dr. M.P. Bachynski, MPB Technologies, Inc., CANADA	Prof. B.S. Liley, Univ. of Waikato, NEW ZEALAND
Dr. H.M. Skarsgard, Univ. of Saskatchewan, CANADA	Inst of Physics, Chinese Acad Sci PEOPLE'S REP. OF CHINA
Prof. J. Teichmann, Univ. of Montreal, CANADA	Library, Inst. of Plasma Physics, PEOPLE'S REP. OF CHINA
Prof. S.R. Sreenivasan, Univ. of Calgary, CANADA	Tsinghua Univ. Library, PEOPLE'S REPUBLIC OF CHINA
Prof. T.W. Johnston, INRS-Energie, CANADA	Z. Li, S.W. Inst Physics, PEOPLE'S REPUBLIC OF CHINA
Dr. R. Bolton, Centre canadien de fusion magnétique, CANADA	Prof. J.A.C. Cabral, Instituto Superior Tecnico, PORTUGAL
Dr. C.R. James,, Univ. of Alberta, CANADA	Dr. O. Petrus, AL I CUZA Univ., ROMANIA
Dr. P. Lukác, Komenského Univerzita, CZECHO-SLOVAKIA	Dr. J. de Villiers, Fusion Studies, AEC, S. AFRICA
The Librarian, Culham Laboratory, ENGLAND	Prof. M.A. Hellberg, Univ. of Natal, S. AFRICA
Library, R61, Rutherford Appleton Laboratory, ENGLAND	Prof. D.E. Kim, Pohang Inst. of Sci. & Tech., SO. KOREA
Mrs. S.A. Hutchinson, JET Library, ENGLAND	Prof. C.I.E.M.A.T, Fusion Division Library, SPAIN
Dr. S.C. Sharma, Univ. of South Pacific, FIJI ISLANDS	Dr. L. Stenflo, Univ. of UMEA, SWEDEN
P. Mähönen, Univ. of Helsinki, FINLAND	Library, Royal Inst. of Technology, SWEDEN
Prof. M.N. Bussac, Ecole Polytechnique,, FRANCE	Prof. H. Wilhelmson, Chalmers Univ. of Tech., SWEDEN
C. Mouttet, Lab. de Physique des Milieux Ionisés, FRANCE	Centre Phys. Des Plasmas, Ecole Polytech, SWITZERLAND
J. Radet, CEN/CADARACHE - Bat 506, FRANCE	Bibliotheek, Inst. Voor Plasma-Fysica, THE NETHERLANDS
Prof. E. Economou, Univ. of Crete, GREECE	Asst. Prof. Dr. S. Cakir, Middle East Tech. Univ., TURKEY
Ms. C. Rinni, Univ. of Ioannina, GREECE	Dr. V.A. Glukhikh, Sci. Res. Inst. Electrophys. Apparatus, USSR
Dr. T. Mui, Academy Bibliographic Ser., HONG KONG	Dr. D.D. Ryutov, Siberian Branch of Academy of Sci., USSR
Preprint Library, Hungarian Academy of Sci., HUNGARY	Dr. G.A. Eliseev, I.V. Kurchatov Inst., USSR
Dr. B. DasGupta, Saha Inst. of Nuclear Physics, INDIA	Librarian, The Ukr.SSR Academy of Sciences, USSR
Dr. P. Kaw, Inst. for Plasma Research, INDIA	Dr. L.M. Kovrizhnykh, Inst. of General Physics, USSR
Dr. P. Rosenau, Israel Inst. of Technology, ISRAEL	Kernforschungsanlage GmbH, Zentralbibliothek, W. GERMANY
Librarian, International Center for Theo Physics, ITALY	Bibliothek, Inst. Für Plasmaforschung, W. GERMANY
Miss C. De Palo, Associazione EURATOM-ENEA, ITALY	Prof. K. Schindler, Ruhr-Universität Bochum, W. GERMANY
Dr. G. Grosso, Istituto di Fisica del Plasma, ITALY	Dr. F. Wagner, (ASDEX), Max-Planck-Institut, W. GERMANY
Prof. G. Rostangni, Istituto Gas Ionizzati Del Cnr, ITALY	Librarian, Max-Planck-Institut, W. GERMANY
Dr. H. Yamato, Toshiba Res & Devel Center, JAPAN	Prof. R.K. Janev, Inst. of Physics, YUGOSLAVIA

**DATE
FILMED
314192**

I