DOE/PC/90176--T55

DOE/PC/90176--T55

[DE92 004191

FINAL TECHNICAL REPORT

September 1, 1990 through August 31, 1991

Project Title: CHARACTERIZATION OF ORGANIC NITROGEN IN IBCSP COALS

DE-FG22-90PC90176

Principal Investigator: Michael A. Kruge, Dept. of Geology, Southern Illinois University, Carbondale, IL 62901

Co-Investigators: Stephen R. Palmer, Dept. of Geology; Edwin J. Hippo, Dept. of Mechanical Engineering and Energy Processes; John C. Crelling, Dept. of Geology, Southern Illinois University, Carbondale, IL 62901

Project Monitor: Ken Ho, CRSC

ABSTRACT

When nitrogen and sulfur contents are expressed as N/C and S/C atomic ratios, it is apparent that, on a molar basis, nitrogen contents are just as high as sulfur in IBCSP coals. Removal of nitrogen from power plant effluents must be given serious consideration in any attempt to solve the acid rain problem. Unlike sulfur, there does not appear to be a relationship between maceral type and bulk nitrogen content. Nitrogen is concentrated in $\text{CH}_3\text{COOH}/\text{H}_2\text{O}_2$ -soluble oxidation products of the large majority of samples. The ether-soluble fraction of the methylated $\text{CH}_3\text{COOH}/\text{H}_2\text{O}_2$ -soluble oxidation products is nearly entirely GC-amenable. The ether-insoluble fractions are not. The fraction insoluble in both ether and methylene chloride has the highest proportion of nitrogen and sulfur, but only accounts for 10% of the oxidation product. A structure accounting for the nitrogen and sulfur in this insoluble fraction is proposed. The aromatic fraction of coal extracts contains GC-amenable nitrogen compounds in greater variety and abundance than do the aromatic fraction of the mild pyrolysis products of the same samples. This is in contrast to the polyaromatic hydrocarbons, which are equally complex in the aromatic fractions of both extract and pyrolyzate. Quinoline has been identified and tetraaromatic azaarenes have tentatively been identified as major nitrogen compounds in the aromatic fractions. The ether-soluble peroxyacetic acid oxidation products of the floated coals and their constituent macerals share most of the same principal nitrogen compounds, however the proportions of these compounds vary widely. The greatest variations are seen among the sporinites from the four coals and among the floated coal and macerals of IBCSP #2. Exact mass determinations indicate that the oxidation products of IBCSP #1 floated coal contain compounds with a single nitrogen atom, with two nitrogens and with both a nitrogen and a sulfur atom. The majority of single nitrogen compounds are interpreted to be oxidized derivatives of pyrrole and its benzologues. A variety of derivatives of pyridine and its benzologues are also present in lesser abundance. The majority of N_2 compounds are either derivatives of pyrazole and its benzologues or dinitro compounds. Sulfur and nitrogen coexist principally in heterogeneous polyaromatic systems (e.g., pyridine/thiophene) or, to a lesser extent, as polyaromatic thiazoles. The ether-soluble fraction contained the largest population of nitrogen compounds. In future work, structure determinations would be greatly simplified if the oxidation products were stripped of their incidental oxygen functional groups by reduction procedures prior to analysis by HRMS. Using standard GCMS data collected on oxidation products, it is possible to estimate the retention times where the compounds detected by probe-HRMS may elute and cross-check those results with elution patterns observed on the nitrogen-selective GC-TSD. Major peaks on GC-TSD chromatograms of IBCSP #1 floated coal are tentatively identified as oxidized pyrrolidines. In future work, since the organic nitrogen (and sulfur) compounds are in such low abundance, we recommend the use of GC-HRMS, for greater precision in the detection and identification of heterocompounds.

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

(This project is funded by the U. S. Department of Energy as part of its cost-shared program with Illinois Department of Energy and Natural Resources.)

EXECUTIVE SUMMARY

The combustion of coals containing sulfur and nitrogen leads to the environmentally unacceptable problems associated with acid rain. Ongoing coal characterization research on the nature of organic sulfur in coal is leading to increased understanding of coal chemistry, necessary in the development of clean coal technology. The nature of organic nitrogen in coal is a problem of similar complexity and needs to be addressed. The overall objective of this study was to determine the content and distribution of organic nitrogen in a series of IBCSP coals and their isolated macerals.

Elemental analysis of IBCSP coal samples indicates that, unlike sulfur, there does not appear to be a relationship between maceral type and bulk nitrogen content. In general, oxidation residues, the material insoluble after oxidation with peroxyacetic acid, are much leaner in nitrogen than their unoxidized parent samples and hence they represent part of the coal that has been partially denitrified. The soluble oxidation products show opposite behavior with the large majority of samples having nitrogen contents that are much higher than those of their precursor materials. When nitrogen and sulfur contents are expressed as N/C and S/C atomic ratios, it is apparent that, on a molar basis, nitrogen contents are just as high as sulfur in IBCSP coals. Removal of nitrogen from power plant effluents must be given serious consideration in any attempt to solve the acid rain problem.

The nitrogen-selective thermionic specific detector (TSD) purchased for this project was operating at or above the specifications set by the manufacturer, with a N/C specificity of approximately 80000:1. The mixture of fifteen nitrogen containing standards was used to calibrate the GC-TSD and GCMS system. By comparing the GC-TSD data with that from the GCMS it is clear that there is good agreement.

The methylated $\text{CH}_3\text{COOH}/\text{H}_2\text{O}_2$ -soluble oxidation products may contain high molecular weight organonitrogen compounds that are not volatile in our gas chromatographic systems. To investigate this possibility the methylated oxidation products of the IBCSP #1 floated coal were fractionated based on their solubility in various organic solvents. The ether-soluble fraction is nearly entirely GC-amenable. The ether-insoluble fractions are not. The fraction insoluble in both ether and methylene chloride has the highest proportion of nitrogen and sulfur, but only accounts for 10% of the oxidation product. A structure accounting for the nitrogen and sulfur in this insoluble fraction is proposed.

Solvent extraction and mild pyrolysis have been employed to render coal components GC-amenable. The aromatic liquid chromatographic fractions of the extracts and pyrolyzates should contain the most accessible nitrogen compounds. The GC-TSD chromatograms of the aromatic fractions were compared with GC-TSD data from a standard mixture of azaarene compounds, however there were few matches. This is in part due to the severely limited number of commercially available azaarene compounds. Nevertheless, inferences can be made about the nature of the observed peaks. Quinoline has been identified and tetraaromatic azaarenes have tentatively been identified as major nitrogen compounds in the aromatic fractions. The pyrolyzate aromatics are much simpler in their distributions of nitrogen compounds than are the extract aromatics.

The methylated, ether-soluble products of the peroxyacetic acid oxidation of the floated coals and the isolated macerals (vitrinite, sporinite and inertinite) of four Illinois basin coals (IBCSP #1, #2, #4 and #6) were subjected to GC-TSD analysis. The oxidation procedure has an advantage over the solvent extract and pyrolysis methods in that more of the coal is rendered GC-amenable. The greatest variations are seen among the sporinites from the four coals and among the floated coal

and macerals of IBCSP #2. Whether the oxidation products are examined for nitrogen, sulfur or for oxidized hydrocarbons, a common theme is present, namely that many of the same compounds are seen in all samples, although in widely varying proportions. The variations are a function of maceral type, depositional conditions of the parent peat, and post-depositional alteration history, including rank level. The GC-TSD system successfully detects a large number of nitrogen compounds in the oxidation products and effectively illustrates differences and similarities among samples.

The oxidation product of IBCSP #1 coal was examined by direct probe, low voltage, high resolution mass spectrometry (HRMS), which can resolve the molecular weight of compounds down to 0.0001 daltons. Because of the mass defects of nitrogen, sulfur and oxygen, molecular formulae for the resultant accurate masses can be assigned, with the assistance of a computer. Among nitrogen-containing fragment ions, those with one nitrogen atom are most abundant. Fragments with two and three nitrogens, as well as a nitrogen and a sulfur atom are also encountered. Summation of intensities of all postulated nitrogen-containing fragment ions gives a percentage of nitrogen in the sample in agreement with bulk elemental analysis results. Molecular structures were assigned to the principal nitrogen-containing compounds, which include those compounds with one and two nitrogen atoms and with both a nitrogen and a sulfur atom. The nitrogen compounds in the oxidation product may be grouped into structural classes, which are subdivided into families, whose members differ by the type and amount of oxygen-containing functional groups and hydrocarbon substituents. The pyrrolic class is represented by three structural families. The first has a pyrrolidine core, the second, a tetrahydroindole or indole nucleus, and the third, a benzocarbazole core. The next class of compounds also contains one nitrogen atom, however, in pyridinic form. These are overall less abundant than the pyrrolic types. The pyridinic class is divided into five families according to the number of additional rings. The first is based on simple pyridine, the second has a quinoline nucleus, the third is based on a benzoquinoline, the fourth is derived from naphthoquinoline and is the most abundant of the class, and the fifth is a phenanthroquinoline derivative. In a non-aromatic, non-cyclic class by itself is an oxidized amine, present in very low concentration. The majority of N₂ compounds are either derivatives of pyrazole and its benzologues or dinitro compounds. Sulfur and nitrogen coexist principally in heterogeneous polyaromatic systems (e.g., pyridine/thiophene) or, to a lesser extent, as polyaromatic thiazoles.

The extensive oxidation of the compounds is necessary to liberate them from the macromolecular maceral structure. However, the identification of individual compounds is greatly complicated by the presence of multiple -OH, =O and -COOCH₃ groups. In addition, these oxygen functions decrease the volatility of the compounds, making even those of moderate molecular weight unsuitable for GC analysis, in spite of our derivatization procedure. It is recommended in future work that the oxidation products be stripped of their oxygens by reduction procedures prior to chromatography.

The GC-TSD allows the separation and recognition of individual nitrogen compounds in the peroxyacetic acid oxidation products, but alone it cannot identify them. Probe-HRMS permits identification, although it cannot discriminate between isomers. To bridge this data gap, GCMS is needed. Standard GCMS was run on the peroxyacetic acid oxidation products of IBCSP #1 floated coal, the same sample used for HRMS. Using these data, it is possible to estimate the retention times where the compounds detected by probe-HRMS may elute and cross-check those results with elution patterns observed on the nitrogen-selective GC-TSD. Since the organic nitrogen (and sulfur) compounds are in such low abundance, we recommend the use of GC-HRMS in future work, for greater precision in the detection and identification of heterocompounds.

OBJECTIVES

The overall objective of this study was to determine the content and distribution of organic nitrogen in a series of IBCSP coals and their isolated macerals. The specific objectives were:

- To determine the bulk nitrogen contents for coals, isolated macerals, oxidation products and residues, solvent extracts and their liquid chromatographic fractions, and pyrolyzates.
- To determine the distribution of organic nitrogen in all coal derivatives enumerated in Objective 1 which are GC-amenable. This will be accomplished by GC-TSD.
- To determine the molecular structure of the major nitrogen compounds detected in Objective 2, using mass spectrometry.

INTRODUCTION AND BACKGROUND

The combustion of coals containing sulfur and nitrogen leads to the environmentally unacceptable problems associated with acid rain. Ongoing coal characterization research on the nature of organic sulfur in coal is leading to increased understanding of coal chemistry, necessary in the development of clean coal technology. The nature of organic nitrogen in coal is a problem of similar complexity and needs to be addressed.

The inability to remove organic nitrogen and sulfur from coal, coupled with new state and federal acid rain legislation, prevents the direct and efficient utilization of many important Illinois Basin coal deposits. The intention of this project is to make a significant preliminary step in the development of coal cleaning processes applicable to organic nitrogen.

Nitrogen is contained in the heteroatomic moieties of coal and other fossil fuels. Many of these structures are complex, heterogeneous and of medium to high molecular weight. They are thus difficult to analyze directly by, for example, the gas chromatographic methods which are so effective on hydrocarbon structures in fossil fuels. There has not been a great deal of previous work on the problem. Pyrolysis of nitrogen-rich kerogens indicates that at lower temperatures, carbazole and its homologues (*i.e.*, pyrrole derivatives) are the dominant nitrogen compounds, whereas at higher temperatures, the pyridine derivatives quinoline, benzoquinolines and their homologues predominate, in agreement with X-ray photoelectron spectroscopic data from coal of different ranks. The distribution of azaarenes in fossil fuels is apparently more dependent upon rank than upon organic matter type.

This project is an extension of our previous work on the characterization of organic sulfur in the Illinois Basin Coal Sample Project (IBCSP) coals to incorporate the analysis of organic nitrogen forms. We are documenting the differences in organic nitrogen content and distribution among the various Illinois Basin coals and among the constituent macerals of each single coal.

EXPERIMENTAL PROCEDURES

Elemental Analysis

The elemental analyses were performed on C-H-N and S analyzers, using whole coal, demineralized coal (demineralized by HF/HCl digestion, then floatation in CsCl_2), isolated macerals (separated by density gradient centrifugation), solvent extracts (THF, refluxing 24 hr in a soxhlet apparatus), and the soluble and residual materials from peroxyacetic acid oxidation of coals and macerals. Results are reported on a dry, ash-free basis.

GC Analysis of Azaarene Standards

The azaarene standard compounds were purchased from the Aldrich Co., diluted to concentrations of *ca.* 100 ng/ μL in hexane and injected into a Hewlett Packard 5890A gas chromatograph, coupled with an HP 5970B Mass Selective Detector. The standards were run in full scan mode, using both of the following GC conditions: 1) 25 m HP-1 column (OV-1 equivalent) with a 0.2 mm i.d. and 0.33 μm film thickness, held at 100° C for 10 min., then the temperature was increased to 300° at a rate of 3°/min and 2) 30 m Supelcowax column with a 0.2 mm i.d. and 0.2 μm film thickness, held at 100° C for 10 min., then the temperature was increased to 270° (near the column limit) at a rate of 3°/min. and held for 5 min. The standard compounds were also combined at a concentration of ≈ 7 ng/ μL each and run collectively using the two GCMS conditions outlined above.

GC-TSD runs employed a Varian 3400 gas chromatograph equipped with a Varian thermionic specific detector (TSD) and an HP-1 (OV-1 equivalent) column, 25 m x 0.2 mm i.d. x 0.33 μm film thickness. The column temp was held at 100° C for 10 min., then ramped to 300° at 3° C/min. The injector temperature was held at 270° C, in splitless mode and the detector temperature was 300° C, with attenuation at 128 and range at 12.

Solvent Fractionation of Oxidation Products

The amount of GC amenable nitrogen in the methylated oxidation products was a concern. In an attempt to gain an understanding of the amount being observed, the methylated product from the IBCSP #1 coal was fractionated. About 5 grams of methylated product was dissolved in 100 ml of methylene chloride. The soluble portion was then isolated in a separatory funnel and extracted with 100 ml of diethyl ether. This procedure yielded three fractions: methylene chloride insolubles, MCI, ether insoluble/methylene chloride solubles, MCS, and ether solubles, ES. Each fraction were weighed, analyzed for C, H, N, and S contents. They were also injected into the GC.

Gas Chromatography/Mass Spectrometry

For the analysis of mild oxidation products a Hewlett Packard 5890A gas chromatograph was employed, coupled with an HP 5970B Mass Selective Detector. It were run in full scan mode, using the following GC conditions: 25 m HP-1 column (OV-1 equivalent) with a 0.2 mm i.d. and 0.33 μm film thickness, held at 50° C for 10 min., then the temperature was increased to 300° at a rate of 8°/min. For the analysis of aromatic fractions of extracts and pyrolyzates, the initial temperature was 100°, the ramp was 3°/min., and the final temperature was 300°.

Gas Chromatography with Thermionic Specific Detector

To detect nitrogen compounds, we employed a Varian 3400 gas chromatograph equipped with a Varian thermionic specific detector (TSD) and an HP-1 (OV-1 equivalent) column, 25 m x 0.2 mm i.d. x 0.33 μ m film thickness. For the analysis of extracts and pyrolyzates, the column temperature was programmed from 100 to 300° at 6° C/min. For oxidation products, the initial temperature was 40° and the ramp was 3°/min. The injector temperature was held at 270° C, in splitless mode and the detector temperature was 300° C, with attenuation at 128 and range at 12. Using an analog-to-digital converter, the data were logged to computer disc for later processing. The mixture of azaarene standards was rerun using both temperature programs.

Low-Voltage High-Resolution Mass Spectrometry.

To characterize the peroxyacetic acid oxidation products of IBCSP #1 coal in more detail, methylated oxidation products were examined using low-voltage, high-resolution mass spectrometry (LV-HRMS). Samples of each fraction were introduced into the mass spectrometer using the solid-probe technique. An ionization voltage of 12 eV was used to minimize fragmentation of parent ions. The spectrometer vacuum was better than 10^{-7} Torr and probe temperatures were initially 200° C but were ramped to 350° C in 50° C intervals. Ion masses were recorded in the high resolution mode at each temperature interval. Data were processed to provide empirical formulae for each of the measured masses and a report prepared of the best mass correlations. Dr. Curt White performed the HRMS work at the Pittsburgh Energy Technology Center.

RESULTS AND DISCUSSION

Elemental Analysis

The data obtained from elemental analysis of the samples under investigation in this study are shown in Table 1. In a few instances insufficient sample was available to complete the analyses. Most of the the coals and macerals have nitrogen contents in the range 1.2 to 1.6%. In the case of the IBCSP #1 and #6 coals there is a significant drop in the nitrogen content after the coals have been floated to remove mineral matter. This would indicate that nitrogen containing minerals were removed during the flotation or demineralization processes.

Unlike sulfur, there does not appear to be a relationship between maceral type and nitrogen content. Indeed the two Illinois No.6 coals (IBCSP #1 and #4) have nitrogen content/maceral type relationships that are opposite. In the case of the #1 coal nitrogen contents increase with increasing maceral density while for the #4 coal they decrease with increasing maceral density. Nitrogen contents for THF extracted coals and macerals occupy a range similar to that occupied by the unextracted coals and macerals. In some samples the process of extraction appears to reduce the nitrogen content but for others the opposite is true. No clear pattern is obvious at this time. The exception to this is the nitrogen contents of the THF extracts which are all consistently below those of their parent coal.

When nitrogen and sulfur contents are expressed as N/C and S/C atomic ratios, it is apparent that, on a molar basis, nitrogen contents are just as high as sulfur in IBCSP coals (Figure 1). Removal of nitrogen from power plant effluents must be given serious consideration in any attempt to solve the acid rain problem.

In general, oxidation residues, the material insoluble after oxidation with peroxyacetic acid, are much leaner in nitrogen than their unoxidized parent samples (Figure 1). This is not the result of dilution from added oxygen since the carbon contents of the oxidation residues are comparable to those of the unoxidized samples. Thus these oxidation residue represent part of the coal that has been partially denitrified. However, the soluble oxidation products show opposite behavior. In the large majority of cases the nitrogen contents of the soluble oxidation products are much higher than those of their precursor materials. This is despite the addition of considerable amounts of oxygen during oxidation. Indeed the carbon contents of the oxidation solubles range from around fifty to forty percent, indicating that the N/C ratios are considerably enhanced upon oxidation. This increase in the N/C ratios is probably the result of the oxidation of carbon in the samples to carbon dioxide which escapes detection. The nitrogen however does not form any volatile products and remains in the oxidation products. Since the soluble oxidation products contain the majority of the organic nitrogen in the oxidation products, the characterization study has concentrated on these materials.

Figure 1 — Averaged atomic ratios of the IBCSP samples.

GC-TSD Calibration

After the installation and initial conditioning of the thermionic specific detector (TSD), it was necessary to check that the TSD was operating at the desired levels of sensitivity and specificity. To do this the TSD calibration mixture (provided by Varian) was used. This mixture contains 2.00 ng/ μ L azobenzene, 2.00 ng/ μ L methyl parathion, 4.00 ng/ μ L malathion and 4.00 ng/ μ L of heptadecane dissolved in iso-octane. Upon comparison of the TSD chromatogram of this mixture obtained on our instrument with that shown in the Varian TSD manual as the ideal TSD response, it was quiet clear that the TSD is operating at or above the specifications set by Varian. Not only is the chromatographic resolution better, but the sensitivity and the nitrogen to carbon selectivity is better than specified. According to the Varian TSD manual one can expect a N/C specificity in the order of 50000:1. However from the integration of the azobenzene and the heptadecane peaks, the TSD is operating at a N/C specificity of approximately 80000:1.

Table 2. Nitrogen Containing Standards - GC-TSD Calibration.

Peak	Compound	Ret. Time (min.)	Absolute conc.(ng/ μ l)	Rel. Response Factor *
A	Quinoline	13.47	7.07	1.00
B	Isoquinoline	14.55	7.60	0.59
C	3-Methylisoquinoline	18.26	7.27	0.71
D	6-Methylquinoline	19.41	6.93	0.78
E	2,8-Dimethylquinoline	21.00	6.40	0.90
F	2,6-Dimethylquinoline	22.89	8.93	0.74
G	2,4-Dimethylquinoline	23.83	6.80	0.62
H	2-propylquinoline	25.99	6.73	0.56
I	Benzo[<i>h</i>]quinoline	37.97	8.27	0.44
J	Acridine	38.37	7.67	0.21
K	Phenanthridine	39.20	7.27	-
L	3-Methylcarbazole	39.20	7.80	-
M	Carbazole	39.41	8.47	0.20
N	9-Ethylcarbazole	40.47	8.53	0.47
O	6-Methylphenanthridine	42.29	6.60	0.32

*Relative response factors were calculated based upon the concentration of each compound, not the concentration of nitrogen in each compound.

In addition the mixture of fifteen nitrogen containing standards was used to calibrate the GC-TSD and the GCMS system. Table 2 contains the peak identification, their absolute concentration, their retention times and their relative response factors (based upon the response of quinoline). On both the GCMS and the GC-TSD, complete resolution of phenanthridine and 3-methylcarbazole could not be achieved. Consequently relative response factors for these compounds could not be calculated.

GCMS — Nitrogen Compound Standards

The azaarene standard compounds were analyzed on the GCMS using both a non-polar (OV-1) and a polar (Supelcowax) column. Phenanthridine, methylcarbazole and carbazole were found to be particularly difficult to resolve on the non-polar column. The polar column was used in an attempt to achieve better GC resolution of the azaarenes. The polar column had several distinct disadvantages. The quinoline group compounds elute over a much narrower time range. A natural mixture, such as we expect to find in coal extracts and pyrolyzates, will be much more complicated and the narrower retention time range would likely result in poorer GC resolution. The carbazole group shows improved resolution on the GC column, but the occurrence of carbazole so late in the GC program and the elution of lower molecular weight compounds after higher might confound interpretation unnecessarily when dealing with unknown compounds in coal derivatives (Table 3). The behavior of the available benzoquinoline group compounds shows no improvement on the polar column. In general, all the azaarenes elute much later on the polar column, with the latest-eluting compounds, carbazole in particular, causing the polar column to be pushed to its maximum

temperature. From these experiments, it is concluded that the non-polar column is overall more effective for both the standard compounds and the expected mixtures of azaarenes in coal derivatives. The good separation of major peaks seen in the GC-TSD analysis of oxidation products on OV-1 supports this judgement.

Table 3 — GCMS retention times of azaarene compounds on non-polar and polar columns.

Peak	Compound	HP-1 Ret. Time	Supelcowax Ret. Time	Ret. Time Difference
A	Quinoline	12.85	32.80	19.96
B	Isoquinoline	13.92	34.31	20.39
C	3-Methylisoquinoline	17.72	35.06	17.34
D	6-Methylquinoline	18.86	37.06	18.21
E	2,8-Dimethylquinoline	20.51	33.82	13.31
F	2,6-Dimethylquinoline	22.42	38.01	15.59
G	2,4-Dimethylquinoline	23.34	39.88	16.54
H	2-Propylquinoline	25.52	38.69	13.17
I	Benzo[<i>h</i>]quinoline	37.48	59.07	21.59
J	Acridine	37.85	59.07	21.22
K	Phenanthridine	38.64	59.56	20.92
O	6-Methylphenanthridine	41.79	63.12	21.33
M	Carbazole	38.92	72.21	33.29
L	3-Methylcarbazole	38.74	61.93	23.19
N	9-Ethylcarbazole	40.01	60.28	20.27

Solvent Fractionation of Oxidation Products

The methylated soluble oxidation products may contain high molecular weight organonitrogen compounds that are not volatile in our gas chromatographic systems. To investigate this possibility the methylated oxidation products of the IBCSP #1 floated coal were fractionated based on their solubility in various solvents. The total methylated product was extracted with methylene chloride and soluble and insoluble fractions obtained. The soluble fraction was then isolated and extracted with diethyl ether. This procedure gave three fractions, a methylene chloride insoluble fraction (MCI), a methylene chloride soluble, but ether insoluble fraction (MCS) and an ether soluble fraction (ES). All fractions were weighed, analyzed for elemental content and the soluble fractions injected into the gas chromatograph to see which contained the volatile components. Only the ES fraction gave FID responses. These responses were identical to the whole product response and demonstrated that the only the ES fraction contributes to the GC traces. GC-TSD analyses were not attempted, but it is expected that the results would be comparable. The question remains as to the fraction of the ES being observed by GC analysis. To estimate this a TGA pyrolysis run was conducted. The product was vaporized completely by 250° C. This does not guarantee that all the sample passes through the GC column but it does indicate that it has a high volatility. The MCS and MCI fractions were reoxidized and methylated. The traces of the products from the reoxidation of the MCS was similar to those of the original product. In other words the MCS may just be a less oxidized version of the ES fraction and upon further oxidation degrades to the structures observed

in the ES fraction. The MCI fraction did not provide any additional volatile product upon further oxidation.

The nitrogen and sulfur contents of the MCI fraction are very high and the amount of carbon is small (Table 4). FTIR indicates the presence of NO groups, as well as SO₃. Some CH₂ groups are also present. The hydrogen content is high and may be in error due to the difficulties in measuring moisture content of this sample and the pickup of water during the elemental analysis of this fraction. The fraction is extremely hygroscopic. Thus, the ratio of N/C and S/C may be a more appropriate values to examine. The N/C of the raw coal is 0.010, it decreases to 0.007 in the ether soluble fraction but concentrates to 0.88 in the MCI fraction. The S/C of the starting coal is 0.015, that for the ES fraction is similar, 0.019 but the MCI fraction has a 0.64 S/C. One possible structure which could explain all the data observation so far, including the selective sulfur removal and non selective nitrogen removal, as well as the observed ES oxidation products, is a derivative of cysteine formed during coalification, with a structure like the one shown below:

Methyl sulfonic acid would be the primary product of oxidation (as observed). Some methyl groups might be replaced in the structure by ethyl, phenyl, or other groups (as observed in the ES product). The remaining oxidation product would have an elemental composition very close to the MCI composition. The NOH group observed by FTIR and the SO₃ group observed by XANES and FTIR would be formed.

Table 4 — Ultimate analysis and yields of fractionated IBCSP #1 oxidized and methylated coal products.

Fraction	%C	%H	%N	%S	%Ash	Weight % of Product
ES	48.31	4.90	0.55	2.10	0.0	57
MCS	70.99	6.43	0.82	1.90	0.0	33
MCI	10.61	4.62	10.97	18.22	18.0	10

GC-TSD of Solvent Extracts and Pyrolyzates

Through the use of a gas chromatograph equipped with a nitrogen-selective thermionic specific detector (GC-TSD), nitrogen compounds of sufficient volatility are directly detected. Solvent extraction and mild pyrolysis have been employed to render coal components GC-amenable. The aromatic liquid chromatographic fractions of the extracts and pyrolyzates should contain the most accessible nitrogen compounds. Figure 2 presents GC-TSD chromatograms of the nitrogen compounds in the aromatic fractions thus derived from IBCSP #1 and #2 coals. The extract aromatics have a peak at ≈8 min. matching in retention time peak A of the azarene standards, which is quinoline (Table 3). The chromatogram of the standard compounds in Figure 2 was run under the same GC conditions as the aromatic samples shown in the figure. The moderately tall peak at 19 min. in the IBCSP #2 aromatic fraction matches the K, L, M standard peak, which is a mixture of phenanthridine, carbazole and 3-methylcarbazole. Aside from these two cases, the standard mixture has no other matches with the main peaks in the aromatic fractions. This is in part due to the limited number of commercially available azarenes. Nevertheless, inferences can be made about the nature of the observed peaks. There is a series of tall peaks in sample IBCSP #2, whose regular spacing suggests that they may comprise one or more homologous series. The cluster around 30 min. suggests a group of polyaromatic isomers, perhaps tetraaromatics such as

benzophenanthridine and/or benzocarbazole. This is logical as the diaromatics (quinolines) cluster around a retention time of 10 min. and the triaromatics (e.g., phenanthridine and carbazole) around 20 min.

The pyrolyzate aromatics are much simpler in distribution, as seen in GC-TSD chromatograms of the representative sample IBCSP #2 (Figure 3). The few main peaks present are matched in the aromatic fraction of the extract — quinoline, a moderately strong peak at 16 min. and a doublet in the "tetraaromatic" region at 29 min. This simple distribution is virtually the same in the floated coal and in the maceral separates. At this point we can only speculate as to why the nitrogen compounds are fewer in the pyrolyzate than in the extract, especially since the distributions of polyaromatic hydrocarbons and sulfur compounds are of equivalent complexity in extracts and pyrolyzates.

GC-TSD of Mild Oxidation Products

The methylated, ether-soluble products of the peroxyacetic acid oxidation of floated coals and isolated vitrinites, sporinites and inertinites of four Illinois basin coals (IBCSP #1, #2, #4 and #6) were subjected to GC-TSD analysis. The oxidation procedure has an advantage over the solvent extract and pyrolysis methods in that more of the coal is rendered GC-amenable. However, multiple oxygen-containing functional groups are typically added, increasing the structural complexity of the individual compounds. Figure 4 displays the TSD chromatograms of the oxidation products of the floated coals, Figure 5, the vitrinites, Figure 6, the sporinites, and Figure 7, the inertinites. In general, the traces all show complex distributions of nitrogen compounds, with most of the strongest peaks in the 40 to 70 min. range present in varying proportions in all samples. It should be noted that the retention times in Figures 4-7 cannot be compared with those in Figure 3, as different temperature programs were employed (faster, with a higher initial temperature for Figure 3 traces). More variation is noted in the lower retention time range (<40 min.), for instance, sample #2 among the vitrinites has very strong peaks around 27 min., whereas other vitrinites have no corresponding peaks in that range. The very large peak in the 1 to 5 min. range is coincident with the elution of the solvent. The complex TSD traces of the oxidation products stand in contrast to the sulfur-selective flame photometric detector data collected on the same samples, which show a much simpler distribution of sulfur compounds. The TSD complexity more closely resembles that seen in the results from the flame ionization detector, which is sensitive to all organic compounds. For all three detection systems — nitrogen-selective, sulfur-selective, and general purpose — the data sets have a common theme: many of the same compounds are seen in all samples, but their proportions vary widely. The variations are a function of maceral type, depositional conditions of the parent peat, and post-depositional alteration history, including rank level. The GC-TSD system successfully detects a large number of nitrogen compounds in the oxidation products and dramatically illustrates differences and similarities among samples. It does not, however, by itself, provide identifications for the peaks.

High Resolution Mass Spectrometry — Oxidation Product of Coal.

Curt White at PETC has used direct injection high resolution MS to look for nitrogen and sulfur compounds in the solvent fractionated oxidation product of the floated IBCSP #1 extract residue. The mass spectrometer determines the molecular weight of the MS fragment ions to 0.0001 daltons. The raw data is then fitted to various empirical formulae and the measured mass is compared to the theoretical calculated mass. The original fit routine did not allow for as much oxygen as the oxidation products contain. So, the data was placed into a Lotus-123 spread sheet. Background peaks of the instrument and standards added by PETC were deleted from the set. The best empirical fit was placed in the worksheet. Next peaks were examined for the possibility of ^{13}C presence in the fragment. This was done by looking for peaks exactly 1.0008 daltons lower to see

if a similar formula without the ^{13}C could be found. Approximately 10 out of 12020 peaks were eliminated. At this point many compounds with empirical formulae containing multiple nitrogen and oxygen atoms were found. Attempts were made to substitute other formulae for these. For example S_2CH_4 has the same mass as O_5 with a difference of 0.0007 daltons. By making this and similar substitutions the empirical formulae were adjusted. A CHO formula was accepted if its mass difference between measured and calculated fell within ± 0.002 daltons. Formulae containing N or S atoms were only accepted if the difference in calculated and actual mass was less than 0.0005 daltons or if no CHO formulae could be found within 0.003 daltons. By using these criteria a conservative identification of nitrogen and sulfur compounds were made. The formulae were then sorted by the number of sulfur and nitrogen atoms they contained. The compounds were grouped by N and S content as well as the solvent fraction they were obtained from and the probe temperature under which they volatilized.

There is a large amount of information in this data and space does not permit the full listing of every ion fragment identified. Standard deviations between measured and theoretical mass were calculated for each group. Because of the selection criteria, the standard deviations of the sulfur and nitrogen containing fragments were smaller than fragments assigned to CHO fragments. The range in standard deviations for the various groups ranged from 0.0004 to 0.0014 daltons. All the mean difference were less than 0.00005 daltons from zero and the ranges in mass difference fell within 2 standard deviations of the mean for all but a few (10) ions.

Data are summarized in Table 5, in which fragments are grouped into structures containing 1, 2 and 3 nitrogen atoms ("N1", "N2" and "N3"); 1 nitrogen plus 1 sulfur atom ("N+S"); and 1 sulfur atom ("S"). These structure contain variable C, H and O atoms. A single class was set aside for "CHO" compounds containing no sulfur or nitrogen atoms. Table 5a lists percent intensities of each group for each fraction and probe temperature. As predicted from the GC work the ether soluble fraction yielded the largest intensity totals for all groups with the lower probe temperature yielding the highest intensity for all fractions. This is true for the individual group intensity as well as for the totals. N1 fragments gave higher intensity than N2 fragments which was higher than the S fragments. Intensity of the N+S fragments were higher than the intensity of the N3 fragments but lower than the intensity of S fragments. The CHO intensity was always bigger than any other group. Note that 4% of the fragments contain 2 nitrogens and 9% of the fragments contain 1 nitrogen.

Table 5a — Percent intensities of ion fragments in HRMS of IBCSP #1 coal oxidation products

Fraction	Probe Temp (°C)	N3	N2	N1	N+S	S	CHO	Total
Ether sol.	200	0.04	2.48	4.80	0.69	1.21	40.80	50.02
	300		0.69	3.74	0.32	0.52	38.68	43.95
	350		0.04	0.04	0.00	0.00	0.40	0.48
CH_2Cl_2 sol.	200		0.30	0.00	0.00	0.00	0.41	0.71
	300		0.46	0.052	0.28	0.34	1.45	3.05
	350		0.09	0.30	0.06	0.00	0.10	0.57
CH_2Cl_2 insol.	200		0.00	0.03	0.00	0.00	1.14	1.17
	300		0.00	0.00	0.02	0.00	0.04	0.06
	350		0.00	0.00	0.00	0.00	0.00	0.00
Total Sum		0.04	4.06	9.43	1.37	2.07	83.03	100.00

Table 5b — Number of ions detected by HRMS of IBCSP #1 coal oxidation products

Fraction	Probe Temp (°C)	N 3	N 2	N 1	N+S	S	CHO	Total
Ether sol.	200	1	24	62	10	19	247	363
	300		13	72	9	14	588	696
	350		1	2			14	17
CH ₂ Cl ₂ sol.	200		4	1			11	16
	300		16	17	9	8	47	97
	350		2	7	1		3	13
CH ₂ Cl ₂ insol.	200		1				2	3
	300				1		2	3
<i>Total Sum</i>		<i>1</i>	<i>60</i>	<i>162</i>	<i>30</i>	<i>41</i>	<i>914</i>	<i>1208</i>

Table 5c — Percent N distributed in ions detected by HRMS of IBCSP #1 coal oxidation products.

Fraction	Probe Temp (°C)	N 3	N 2	N 1	N+S	S	CHO	Total
Ether sol.	200	0.67	27.71	26.63	3.84	0.00	0.00	58.86
	300		7.67	20.75	1.78	0.00	0.00	30.20
	350		0.09	0.06	0.00	0.00	0.00	0.15
CH ₂ Cl ₂ sol.	200		0.31	0.04	0.00	0.00	0.00	0.35
	300		2.70	2.86	1.53	0.00	0.00	7.09
	350		1.04	1.68	0.36	0.00	0.00	3.07
CH ₂ Cl ₂ insol.	200		0.00	0.16	0.00	0.00	0.00	0.16
	300		0.00	0.00	0.12	0.00	0.00	0.12
<i>Total sum</i>		<i>0.67</i>	<i>39.52</i>	<i>52.19</i>	<i>7.62</i>	<i>0.00</i>	<i>0.00</i>	<i>100.00</i>

The distribution by number (Table 5b) shows similar trends but the differences in total distribution is not as great. N1, N2, N+S, and S are all larger percentages of the whole on a number basis than on an intensity basis. This indicates that the individual fragments are of lower intensity than the CHO fragments. The data indicate that the N2 compounds tend to be lower masses than the N1 group. The same can be said of S and N+S groups in relation to the N1 group. The CHO group appears to trend to the heaviest masses. Table 5c shows the nitrogen distribution. Nitrogen is nearly evenly split between N2 and N1 structures in the ether soluble fraction, and this fraction contains nearly all of the nitrogen with slightly more in the more volatile (200°) subfraction. Most of the sulfur is in the lightest ether soluble subfraction and appears in single S compounds in greater amounts than in the N+S compounds.

Molecular Structure Assignments — Oxidation Products

As discussed above, low voltage, high resolution mass spectral (LV-HRMS) data have been employed to assign molecular formulae to compounds present in the peroxyacetic acid oxidation product of IBCSP #1 coal. Subsequently, molecular structures were assigned to the principal nitrogen-containing compounds, which include those compounds with one and two nitrogen atoms and with both a nitrogen and a sulfur atom. The high resolution technique is particularly useful in dealing with non-hydrocarbons. For example, because of the mass defects of sulfur and hydrogen, the sulfur compound dibenzothiophene (184.0346 daltons) can be distinguished from the hydrocarbon tetramethylnaphthalene (184.1251 daltons) by reference to exact masses, although

they both have the same nominal mass of 184. Standard GCMS, which only uses nominal masses, may have difficulty in distinguishing such compounds, especially if one is in low concentrations in a complex mixture. HRMS can provide the necessary discrimination, even without chromatographic separation. In the present case, the low voltage mode has been employed, since the soft ionization suppresses fragmentation, yielding more molecular ions, from which formulae for whole molecules can be derived. Finally, in these experiments, the direct probe inlet was employed at three alternate temperatures — 200, 300 and 350 °C. It was decided that the 200 ° run was the most likely to contain those compounds which are volatile enough to be GC-amenable and thus detectable by our other techniques — GC-TSD and standard GCMS. Our molecular structure determination efforts therefore focused on the 200 ° probe data.

Each structure shown is one of many possible isomers. The probe-HRMS technique is blind to isomerism, thus none of the structures presented in this report should be taken to be isomer-specific, but instead taken to represent all possible isomers of the structure, one or more of which may actually be present. Therefore, positions of groups such as -CH₃, -OH, and -COOCH₃ may vary from those illustrated. Also, the arrangement of aromatic rings may differ from that shown, for example, the rings may be in linear, anthracene-like or angular, phenanthrene-like configurations. This however should not detract from the valuable information provided by the technique.

Figure 8 presents alternative structures for the same molecular formulae. In Figure 8a, four possibilities for C₁₂H₁₁N₂O₅ are given, two of which are three-ring pyrrolic structures, another a diaromatic diamine, and another a dinitro tetralin. All structures in Figure 8a are MS fragments, although nearly intact. The pyrrolic types suffer a loss of a hydrogen, whereas the others are interpreted to have the readily-occurring loss of a methoxy group from the esterified carboxyl function. The structures drawn for C₈H₁₁NO₉ are inferred to be molecular ions (Figure 8b). Of the five possibilities presented, one is an *n*-octane derivative with an -NO₂ function, two are cyclohexane derivatives with -NO₂ or -NH₂, and two are cyclic structures with nitrogen incorporated into the rings (oxidized piperidine and pyrrolidine). Figures 8c and 8d show polyaromatic alternatives to two formulae, presenting heterocyclic and exocyclic alternatives for the single nitrogen atom. C₁₂H₁₀N₂O₅ poses a half dozen alternate diaromatic structures (Figure 8e). The "R" factor given for each formula represents the relative intensity of the mass peak, as a percentage of the strongest peak in the data set (a methylated benzenepentacarboxylic acid). At 4.32% relative intensity, the mass of 265.0436 represented in Figure 8b is the most abundant nitrogen-containing peak. The majority of nitrogen peaks in the data set have a relative intensity of <1%. The low abundance of the nitrogen compounds hinders their identification by conventional techniques.

The nitrogen compounds in the oxidation product may be grouped into structural families, whose members differ by the type and amount of oxygen-containing functional groups and hydrocarbon substituents. The oxygens have of course been added during the peroxyacetic acid processing. Oxygen-free moieties must be visualized to arrive at an image of the components as they originally existed within the macromolecular coal structure. The pyrrolic class is represented by Structures I, II and III in Figure 9. Structure I has a pyrrolidine core and is relatively abundant (R=2.68%). Structure II is nearly as abundant and has a tetrahydroindole nucleus. Indoles have also been identified in the data set. Structure III is larger, less abundant and has a benzocarbazole core. Each of these three structures is the most abundant of several similar compounds, thus are chosen to represent their respective structural families.

The next class of compounds also contains one nitrogen atom, however, in pyridinic form. These are overall less abundant than the pyrrolic types represented by Structures I, II and III. The pyridinic class is divided into families according to the number of additional rings (Structures IV through VIII in Figure 9). Again, each structure shown is the most abundant compound in its respective family. Structure IV is derived from simple pyridine and is not very abundant

($R=0.73\%$). Structure V has a quinoline nucleus and is even less abundant than Structure IV. Structure VI is based on a benzoquinoline and the more abundant Structure VII has a partially hydrogenated naphthoquinoline core. The polyaromatic Structure VIII, with a dihydrophenanthroquinoline nucleus, is present as a weak component. In a non-aromatic, non-cyclic class by itself is Structure IX, an oxidized amine, with a very low concentration of 0.14%.

Compounds with two nitrogen atoms are also detected by HRMS. A class of polyaromatic compounds with each nitrogen incorporated into a separate ring is illustrated by Structures X and XI in Figure 9. Structure X represents a heterogeneous arrangement, as it is both pyridinic and pyrrolic. Structure XI is homogeneous, with two fused pyridines. Neither are particularly abundant compounds. Pyrazoles and their benzologues comprise a second class of N_2 compounds, represented by Structures XII, XIII and XIV in Figure 9. Structure XII, a derivative of simple pyrazole, is present in very low abundance. In contrast, Structure XIII represents one of the strongest nitrogen peaks in the data set ($R=3.99\%$) and is characterized by an indazole nucleus. The weak Structure XIV is essentially a naphthoindazole. The last N_2 class of compounds is that in which the nitrogens are not incorporated into rings. One example is shown, Structure XV in Figure 9, an oxidized tetralin with two nitro groups, with a strong relative intensity of 1.85%. This structure is very similar to one in Figure 8a and they both coincidentally have the same nominal mass of 263 daltons. However, the exact mass determinations indicate that their molecular formulae are significantly different. Members of the final class of compounds contain both a nitrogen and a sulfur atom. In Structure XVI these atoms are lodged in different rings, as a pyridine and a dihydrothiophene. The compound is relatively abundant, with a relative intensity of 2.87%. A less abundant family of thiazole derivatives is also present, represented by Structure XVII, which is essentially a dihydronaphthothiazole.

The extensive oxidation of the compounds is necessary to liberate them from the macromolecular maceral structure. However, the identification of individual compounds is greatly complicated by the presence of multiple -OH, =O and -COOCH₃ groups. In addition, these oxygen functions decrease the volatility of the compounds, making even those of moderate molecular weight unsuitable for GC analysis, in spite of our derivatization procedure. It is recommended in future work that the oxidation products be stripped of their oxygens by reduction procedures prior to chromatography.

GCMS of Oxidation Products

The GC-TSD allows the separation and recognition of individual nitrogen compounds in the peroxyacetic acid oxidation products, but alone it cannot identify them. Probe-HRMS permits identification, although it cannot discriminate between isomers. To bridge this data gap, GCMS is needed. Standard GCMS was run on the peroxyacetic acid oxidation products of IBCSP #1 floated coal, the same sample used for probe-HRMS. Standard GCMS chromatograms of the most abundant nitrogen-containing ions in the HRMS data set were compared with the GC-TSD chromatogram. Since these are standard GCMS data, only nominal masses are resolved. Thus, in each chromatogram, it is likely that non-nitrogen peaks are present, possibly as the most abundant compounds. Since the data also was collected using standard high voltage electron impact ionization, there are many fragments ions present, making the picture complex. It is recommended that in future projects of this type, experiments are run using GC-HRMS in soft ionization mode. Nevertheless, with the present conventional GCMS data, the retention time ranges can be determined for each of the ions in question. Partial mass chromatograms m/z 261 and 265, two of the strongest ions in the HRMS data, are shown in figure 10. According to the HRMS results, the abundant pyrrolidine (Structure I, Figure 9) and/or its isomers should be among the peaks on the m/z 261 trace in Figure 10. It is not possible to say which peak in the trace using the standard GCMS data, however the retention time window where it is expected is recognized. Similarly, the m/z 265 chromatogram should display compounds with structures like those in Figure 8b.

Table 6. Nitrogen Containing Standards. Retention Times (Min.)

Peak	Compound	GC-TSD	GCMS
A	Quinoline	25.583	21.117
B	Isoquinoline	26.617	21.577
C	3-Methylisoquinoline	30.033	23.045
D	6-Methylquinoline	31.083	23.496
E	2,8-Dimethylquinoline	32.533	24.083
F	2,6-Dimethylquinoline	34.283	24.803
G	2,4-Dimethylquinoline	35.150	25.184
H	2-propylquinoline	37.150	25.940
I	Benzo[h]quinoline	48.600	30.613
J	Acridine	48.600	30.613
K	Phenanthridine	49.783	31.068
L	3-Methylcarbazole	49.783	31.068
M	Carbazole	49.783	31.068
N	9-Ethylcarbazole	51.033	31.523
O	6-Methylphenanthridine	52.817	32.243

The standard GCMS and probe-HRMS data must be compared to the GC-TSD results. The largest HRMS peaks determined to contain nitrogen should correspond to the tallest peaks on the GC-TSD chromatogram of the same sample, unless the HRMS peak represents the sum of several isomers. Although both chromatographs were equipped with the same column type, the retention times from the GCMS and GC-TSD are not directly comparable, owing to the faster temperature program on the GCMS and to inherent differences in the systems. The azaarene standards used in the calibration of the GC-TSD were run again on both instruments, using the temperature programs employed for the analysis of the oxidation products. The resulting retention times are presented in Table 6 and show excellent correlation: a linear regression gives an r^2 of 1.000, with the following equation:

$$RT_{GCMS} = 0.40673 RT_{GCTSD} + 10.814$$

where RT is retention time, subscripted to indicate the instrument. Thus, it is possible to predict the GCMS retention times of the main nitrogen peaks seen on GC-TSD traces and then check if standard GCMS mass chromatograms of the strongest nitrogen-containing HRMS ions show peaks at or close to any of the predicted times. It is also possible to compute equivalent retention time ranges, useful in denoting the clusters seen in the standard GCMS ion chromatograms. For example, m/z 261 and m/z 265 mass chromatograms in Figure 10 are drawn between 30 and 43 minutes. Using the above regression equation, the equivalent GC-TSD retention time range is 47 to 79 minutes, which is the portion of the GC-TSD chromatogram shown in the center of Figure 10 and includes several of the strongest peaks in that chromatogram (compare the full IBCSP #1 trace in Figure 4). Juxtaposition of the partial chromatograms in Figure 10 shows good alignment of the GC-TSD peak doublet at 60.3 min. with the M/z 261 doublet at 35.3 min., marked with the short double arrow. This peak can be tentatively identified as an oxidized pyrrolidine (tetrahydropyrrole) isomeric to Structure I. The GC-TSD peaks at 66.5 and 68.4 mins. match the m/z 265 peaks at 37.9 and 38.5 mins, marked with long arrows. These are likely to be one of the structures shown in Figure 8b, one of which is also an oxidized pyrrolidine.

CONCLUSIONS AND RECOMMENDATIONS

- When nitrogen and sulfur contents are expressed as N/C and S/C atomic ratios, it is apparent that, on a molar basis, nitrogen contents are just as high as sulfur in IBCSP coals. Removal of nitrogen from power plant effluents must be given serious consideration in any attempt to solve the acid rain problem.
- Unlike sulfur, there does not appear to be a relationship between maceral type and bulk nitrogen content. Nitrogen is concentrated in $\text{CH}_3\text{COOH}/\text{H}_2\text{O}_2$ -soluble oxidation products of the large majority of samples.
- The ether-soluble fraction of the methylated $\text{CH}_3\text{COOH}/\text{H}_2\text{O}_2$ -soluble oxidation products is nearly entirely GC-amenable. The ether-insoluble fractions are not. The fraction insoluble in both ether and methylene chloride has the highest proportion of nitrogen and sulfur, but only accounts for 10% of the oxidation product. A structure accounting for the nitrogen and sulfur in this insoluble fraction is proposed.
- The aromatic fraction of coal extracts contains GC-amenable nitrogen compounds in greater variety and abundance than do the aromatic fraction of the mild pyrolysis products of the same samples. This is in contrast to the polyaromatic hydrocarbons, which are equally complex in the aromatic fractions of both extract and pyrolyzate.
- Quinoline has been identified and tetraaromatic azaarenes have tentatively been identified as major nitrogen compounds in the aromatic fractions.
- The ether-soluble peroxyacetic acid oxidation products of the floated coals and their constituent macerals share most of the same principal nitrogen compounds, however the proportions of these compounds vary widely. The greatest variations are seen among the sporinites from the four coals and among the floated coal and macerals of IBCSP #2.
- Exact mass determinations indicate that the oxidation products of IBCSP #1 floated coal contain compounds with a single nitrogen atom, with two nitrogens and with both a nitrogen and a sulfur atom. The majority of single nitrogen compounds are interpreted to be oxidized derivatives of pyrrole and its benzologues. A variety of derivatives of pyridine and its benzologues are also present in lesser abundance. The majority of N_2 compounds are either derivatives of pyrazole and its benzologues or dinitro compounds. Sulfur and nitrogen coexist principally in heterogeneous polyaromatic systems (e.g., pyridine/thiophene) or, to a lesser extent, as polyaromatic thiazoles. The ether-soluble fraction contained the largest population of nitrogen compounds.
- In future work, structure determinations would be greatly simplified if the oxidation products were stripped of their incidental oxygen functional groups by reduction procedures prior to analysis by HRMS.
- Using standard GCMS data collected on oxidation products, it is possible to estimate the retention times where the compounds detected by probe-HRMS may elute and cross-check those results with elution patterns observed on the nitrogen-selective GC-TSD. Major peaks on GC-TSD chromatograms of IBCSP #1 floated coal are tentatively identified as oxidized pyrrolidines.
- In future work, since the organic nitrogen (and sulfur) compounds are in such low abundance, we recommend the use of GC-HRMS, for greater precision in the detection and identification of heterocompounds.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

TABLE 1. ELEMENTAL COMPOSITION (dry, ash-free basis)

	%C	%H	%N	%S		%C	%H	%N	%S
a) Micronized coals									
IBC 101	82.10	4.89	1.62	4.79	e) Semifusinites	IBC 101	79.61	1.35	2.03
IBC 102	78.87	4.64	1.53	3.52		IBC 102	79.63	1.41	3.75
IBC 104	75.70	4.06	1.39	3.56		IBC 104	83.87	1.24	3.79
IBC 106	87.36	5.45	1.96	4.29		IBC 106	80.96	1.64	3.21
b) Floated coals									
f) Extracted floated coals									
IBC 101	76.86	4.85	1.29	3.27	g) Extracted sporinite	IBC 101	76.61	4.67	1.52
IBC 102	80.64	5.32	1.50	1.57		IBC 102	80.03	5.03	1.40
IBC 104	79.85	4.96	1.31	3.42		IBC 104	77.36	5.08	1.22
IBC 106	80.11	4.98	1.65	2.41		IBC 106	78.70	5.05	1.68
c) Sporinites									
g) Extracted sporinite									
IBC 101	76.18	5.90	1.11	3.87	h) Extracted vitrinite	IBC 101	74.52	-	1.35
IBC 102	79.50	7.17	1.63	1.50		IBC 102	81.42	6.83	1.30
IBC 104	79.62	6.49	1.50	3.51		IBC 104	77.57	6.30	1.53
IBC 106	80.75	6.28	1.29	2.33		IBC 106	76.39	6.76	-
d) Vitrinites									
h) Extracted vitrinite									
IBC 101	77.29	4.82	1.26	2.83		IBC 101	77.41	5.21	1.58
IBC 102	81.15	5.47	1.61	2.03		IBC 102	80.50	5.04	1.63
IBC 104	79.55	5.05	1.42	2.70		IBC 104	75.88	4.90	1.40
IBC 106	77.29	4.82	1.26	2.83		IBC 106	77.19	4.46	1.74

	%C	%H	%N	%S		%C	%H	%N	%S
i) Extracted semifusinites					n) Oxidation Residue - semifusinite				
IBC 101	83.50	3.70	1.14	2.22	IBC 101	83.51	2.81	0.69	0.80
IBC 102	-	-	-	-	IBC 102	73.53	3.66	1.03	1.76
IBC 104	77.93	3.35	1.02	4.60	IBC 104	75.88	3.06	0.83	2.05
IBC 106	80.65	3.72	1.70	3.27	IBC 106	86.41	2.66	0.97	0.90
j) THF extracts					o) Oxidation Soluble Materials - float				
IBC 101	77.89	6.26	1.08	2.15	IBC 101	51.86	3.75	1.16	3.34
IBC 102	79.96	6.74	0.92	1.95	IBC 102	51.61	4.39	1.69	2.20
IBC 104	77.60	6.22	1.03	2.38	IBC 104	53.86	4.17	1.61	4.14
IBC 106	79.50	6.78	1.05	1.40	IBC 106	50.34	4.01	2.09	2.76
k) Oxidation Residue - float					p) Oxidation Soluble Materials - sporinite				
IBC 101	75.30	4.01	0.90	1.25	IBC 101	54.16	5.82	2.68	-
IBC 102	73.18	4.80	1.35	0.80	IBC 102	-	-	-	-
IBC 104	77.54	3.85	0.89	1.33	IBC 104	51.36	5.14	1.44	-
IBC 106	76.80	4.79	0.84	1.14	IBC 106	51.75	4.66	1.65	2.95
l) Oxidation Residue - sporinite					q) Oxidation Soluble Materials - vitrinite				
IBC 101	70.52	6.61	0.60	1.50	IBC 101	37.54	3.40	1.20	6.22
IBC 102	71.02	6.26	0.74	0.98	IBC 102	58.16	4.62	2.28	1.03
IBC 104	70.50	6.66	0.73	1.02	IBC 104	53.81	4.61	1.50	-
IBC 106	69.41	6.27	0.80	2.14	IBC 106	51.34	3.87	1.92	3.09
m) Oxidation Residue - vitrinite					r) Oxidation Soluble Materials - semifusinite				
IBC 101	65.32	4.19	0.96	0.60	IBC 101	47.26	3.34	2.25	3.57
IBC 102	-	-	-	0.45	IBC 102	-	-	-	-
IBC 104	-	-	-	1.34	IBC 104	39.02	4.29	1.38	-
IBC 106	65.79	3.64	1.09	-	IBC 106	49.15	4.08	2.08	4.74

Figure 2 — GC-TSD chromatograms of aromatic fractions and of standard compounds.

Figure 3 — GC-TSD chromatograms of aromatic fractions of mild pyrolysis products of IBCSP #6 floated coal and macerals.

Figure 4 — GC-TSD chromatograms of floated coal oxidation products.

Figure 5 — GC-TSD chromatograms of vitrinite oxidation products.

Figure 6 — GC-TSD chromatograms of sporinite oxidation products.

Figure 7 — GC-TSD chromatograms of inertinite oxidation products.

Figure 8. Possible molecular structures from five of the formulae derived from HRMS data collected on IBCSP #1 floated coal (probe temp. = 200 °C). Exact masses and relative response (R) as a percentage of the strongest peak in the data set are given. See text.

Figure 9. Possible molecular structures from the formulae derived from HRMS data collected on IBCSP #1 floated coal (probe temp. = 200 °C). Exact masses and relative response (R) as a percentage of the strongest peak in the data set are given. See text.

X.
 $C_{18}H_9N_2O$
 269.07148
 $R = 0.63$

XI.
 $C_{14}H_5N_2O_5$
 281.01982
 $R = 0.35$

XII.
 $C_{11}H_{11}N_2O_4$
 315.01641
 $R = 0.25$

XIII.
 $C_{12}H_{10}N_2O_5$
 262.0589
 $R = 3.99$

XIV.
 $C_{22}H_{21}N_2O_2$
 345.16029
 $R = 0.43$

XV.

$C_{11}H_7N_2O_6$
 263.0303
 $R = 1.85$

XVI.
 $C_{14}H_9NSO_5$
 303.02012
 $R = 2.87$

XVII.
 $C_{15}H_{13}NSO_4$
 303.05651
 $R = 0.73$

Figure 9. (Continued)

Figure 10 — Comparison of GCMS and GC-TSD analyses of oxidized IBCSP #1 floated coal. Retention times: GCMS, 30-43 min.; GC-TSD, 47-79 min. See text for explanation of retention time correlation.

PROJECT MANAGEMENT REPORT

1 June through 31 August 1991

Project Title: CHARACTERIZATION OF ORGANIC
NITROGEN IN IBCSP COALS

Principal Investigator: Michael A. Kruege, Assistant Professor
Dept. of Geology
Southern Illinois Univ.
Carbondale, IL 62901 (618) 453-3351

Co-Investigator: Stephen R. Palmer, Adjunct Assistant Professor
Dept. of Geology
Southern Illinois Univ.
Carbondale, IL 62901 (618) 453-7325

Co-Investigator: Edwin J. Hippo, Associate Professor
Dept. of Mechanical Engineering
and Energy Processes
Southern Illinois Univ.
Carbondale, IL 62901 (618) 453-7009

Co-Investigator: John C. Crelling, Professor
Dept. of Geology
Southern Illinois Univ.
Carbondale, IL 62901 (618) 453-3351

Project Monitor: Ken Ho, CRSC

COMMENTS

The project was completed on schedule and on budget.

(This project is funded by the U. S. Department of Energy as part of its
cost-shared program with Illinois Department of Energy and Natural Resources.)

EXPENDITURES - EXHIBIT B

Projected and Estimated Actual Expenditures by Quarter

Quarter*	Types of Cost	Direct Labor	Materials and Supplies	Travel	Major Equipment	Other Direct Costs	Indirect Costs	Total
Sept. 1, 1990	Projected	5,494	950	0	4,300	2,056	2,560	15,360
to Nov. 30, 1990	Estimated Actual	2,175	2,040	0	1,290	1,149	1,331	7,985
Sept. 1, 1990	Projected	10,988	1,900	0	4,300	4,113	4,260	25,561
to Feb. 28, 1991	Estimated Actual	5,891	2,713	0	2,028	2,663	2,659	15,954
Sept. 1, 1990	Projected	16,481	2,850	0	4,300	6,169	5,960	35,760
to May 31, 1991	Estimated Actual	12,392	3,301	0	2,187	2,388	4,054	24,322
Sept. 1, 1990	Projected	42,202	3,800	800	4,300	8,225	11,865	71,192
to Aug. 31, 1991	Estimated Actual	37,523	3,800	800	3,975	8,099	10,679	64,076

*Cumulative by quarter.

COSTS BY QUARTER - EXHIBIT C

Characterization of Organic Nitrogen in IBCSP Coals

Cumulative \$
(thousands)

Months and Quarters

O = Projected Expenditures -----

Δ = Actual Expenditures _____

Total CRSC Award \$ 71,192

SCHEDULE OF PROJECT MILESTONES

COMMENTS:

The project is complete. All milestones have been reached.

END

DATE
FILMED
01/30/92

I