Progress Report Spectroscopic Studies of Hydrogen Collisions

John Kielkopf
Physics Department
University of Louisville, Louisville KY 40292
December 10, 1991

DOE/ER/13332--T4

DE92 008752

Summary

Low energy collisions involving neutral excited states of hydrogen are being studied with vacuum ultraviolet spectroscopy. Atomic hydrogen is generated by focusing an energetic pulse of ArF, KrF, or YAG laser light into a cell of molecular hydrogen, where a plasma is created near the focal point. The H_2 molecules in and near this region are dissociated, and the cooling atomic hydrogen gas is examined with laser and dispersive optical spectroscopy. In related experiments, we are also investigating neutral $H + O$ and $H + metal - atom$ collisions in these laser-generated plasmas.

Diagnostics

Over the past few months we have concentrated on characterizing the plasma conditions in order to select appropriate delay times and densities for the studying the atomic processes we are interested in.

We have produced these plasmas with an excimer laser (193 nm ArF at 400 mJ, or 248 nm KrF at 800 mJ), and with a YAG laser (120mJ or more in the 1064 nm fundamental or 320 mJ in the 532 nm second harmonic). However we found that the difference in beam quality and pulse duration of the lasers results in substantially different threshold conditions, and that without modification the excimer laser does not consistently produce focused light of sufficient field strength to generate a plasma in the target gas. Our goal is to use the excimer laser for this purpose because of the versatility it provides in relative timing of the excitation and the dye laser pulses, but for observations with a delay of about 20 nanoseconds, the same YAG laser can create the plasma and the subsequent tunable four-wave mixed vacuum ultraviolet probe pulse. Emission spectroscopy of this plasma was used by Carlos Aguilar in his thesis research on diagnostics of the laser-generated plasmas. He found that the hydrogen is completely dissociated and highly ionized in the focal spot. The electron density and temperature in the brightest regions exceed 10^{18} cm^{-3} and 10^5 K within nanoseconds of excitation. CCD images of the focal region show an inhomogeneous breakdown, dark at the center where the ionization is total, and bright with intense atomic emission on the incident side. Initially, the plasma radiates recombination radiation and the spectrum of atomic hydrogen modified by collisions with electrons and ions. Microseconds later, neutral collisions dominate.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

Excimer laser upgrade

The excimer laser is being modified to change its cavity to an unstable resonator. The manufacturer claims that this relatively inexpensive improvement will reduce beam divergence by an order of magnitude, and thereby increase by about two orders of magnitude the electric field strength at the laser focus. This will keep the output of the excimer well above threshold for breakdown, even when the energy falls off after several hours of operation.

To provide precise delay timing, a pulse generator with calibrated delays from nanoseconds to milliseconds is being installed on the trigger circuit of the YAG laser. The pulse generator then initiates the probing dye laser pulses after the dissociated hydrogen gas has cooled for a controlled time.

Dissociation of OH

For observations of $H + O$ half-collisions, we worked on ways to produce OH in a quantity sufficient for vacuum ultraviolet laser fluorescence and absorption spectroscopy. Impurities that would quench excited states or contaminate the high vacuum system must be absent. In a study of the possibility of using laser-generated plasmas for this purpose, we searched for emission and fluorescence from OH in the plasma made by the YAG fundamental focussed on H_2 with H_2O added as an impurity. Such plasmas were found to totally dissociate OH. By contrast, a very low density H_2O vapor in the unfocussed beam of a 193 nm excimer laser produces OH and H through multiphoton processes. A measurement of the OH density and the emission from its radiatively dissociating states in collaboration with Frank Tomkins (Argonne National Lab) awaits the modification of the excimer laser.

Dye laser developments

Work has continued on improving the operation of the dual dye laser system. We finished an analysis of dye cell design, and optimized dyes in spectral regions required for four-wave mixing in Hg vapor. The operation of the YAG with simultaneous doubled and tripled outputs was achieved, and preliminary wavelength calibration of the dye lasers allowed us to produce tunable light at 1220 Å in the Lyman- α and OH spectral regions. We implemented an improved I_2 fluorescence system for precise calibration and a CID camera to monitor fluorescence from the two-photon resonance in Hg that is the first stage of vacuum ultraviolet generation. The laser linewidth was further reduced to less than 0.04 cm^{-1} .

Wavelength calibration of the lasers has been a persistent problem, because the monochromators that would be useful for this task are committed to detection in the experiment. There are as yet no commercial wavemeters that permit measurements of ultraviolet wavelengths. To solve this problem we experimented with a vacuum Lloyd's mirror interferometer combined with an array detector and fast Fourier transform analyses. In his thesis research, Lawrence Portaro demonstrated that the prototype device has an accuracy of better than 10 cm^{-1} . We anticipate results of about 2 cm^{-1} in the final version we are working on

now (Kielkopf and Portaro, 1991). That would be sufficient to tune our lasers to resonant frequencies in the experiment, from which they would be self-calibrating.

H_2^+ satellites

Predictions were made several years ago that the continuum emitted by excited $n=2$ states of hydrogen atoms colliding with protons should exhibit greatly enhanced strength in satellite regions near 1240 and 1400 Å. Spectra of the laser plasmas analyzed in August show these effects, and that the satellites are correlated with the ion density (Kielkopf and Aguilar, 1991). Since the H_2^+ potentials are well known, these satellites are excellent test cases for theoretical analyses of far wing line broadening. The laboratory observations also confirm an identification we made of these features in the spectra of hydrogen-rich white dwarf stars (Allard and Kielkopf, 1991).

Experiments continue now on tunable vacuum ultraviolet laser spectroscopic measurements of the regions too close to the line to resolve in the emission spectra. The changing relative effects of H^+ and H perturbers as the discharge evolves is being used to identify radiative continua associated with the H_2^+ and H_2 potentials.

Calculations

On the theoretical side, calculations of the line shape now utilize the exact numerical potential curves, rather than the idealized analytical models we were restricted to previously. This is primarily because we dedicate a workstation to computing the correlation functions from accurate theoretical potentials. These line shapes are being compared to the laser spectroscopic profiles of Lyman- α broadened by H^+ . Work on modeling the laser-generated plasmas, particularly on understanding the time and spatial dependence of the H and H^+ concentrations, is also underway as part of Lisa Xiang's thesis research.

Papers published and in preparation

J. Kielkopf, K. Myneni, and F. Tomkins, *Unusual fluorescence of H_2 excited by multiphoton processes* J. Phys. B **23**, 251-261 (1990).

J. Kielkopf, *Lasing in the aluminum and indium resonance lines following photoionization and recombination in the presence of H_2* J. Opt. Soc. Am. B **8**, 212-219 (1990).

N. Allard and J. Kielkopf, *Temperature and density dependence of the Lyman- α line wing ...* Astron. Astrophys. **242**, 133-141 (1991).

J. Kielkopf and C. Aguilar, *Observations of satellites on Lyman-alpha due to collisions with H^+* in preparation (1991).

J. Kielkopf and L. Portaro, *A Lloyd's mirror wavemeter* in preparation (1991).

END

**DATE
FILMED**

4/10/92