**PORTIONS
OF THIS
DOCUMENT
ARE
ILLEGIBLE**

LA-UR-81-3324

LA-UR--81-3324

DE82 004336

TITLE: ON DUAL SPACES OF DIFFERENTIAL LIE ALGEBRAS

MASTER

AUTHOR(S): B. A. Kupershmidt

NOTICE
PORTIONS OF THIS REPORT ARE REPRODUCED
It has been reproduced from the 1981
Available copy to permit the broadest
possible availability.

SUBMITTED TO: Proceedings of Lie Algebras Conference
University of Rutgers, New Jersey
May 29-31, 1981

University of California

By acceptance of this article, the publisher recognizes that the
U.S. Government retains a nonexclusive, royalty free license
to publish or reproduce the published form of this contribu-
tion, or to allow others to do so, for U.S. Government pur-
poses.

The Los Alamos Scientific Laboratory requests that the pub-
lisher identify this article as work performed under the aus-
pices of the U.S. Department of Energy.

LOS ALAMOS SCIENTIFIC LABORATORY

Post Office Box 1663 Los Alamos, New Mexico 87545

An Affirmative Action/Equal Opportunity Employer

UNITED STATES
DEPARTMENT OF ENERGY
CONTRACT W-7400-ENG-30

100% REPRODUCTION OF THIS DOCUMENT IS UNLIMITED

ON DUAL SPACES OF DIFFERENTIAL LIE ALGEBRAS

B. A. Kupersmidt*

0. Let A be a commutative algebra over a field $\bar{k}$ of characteristic zero, and let $\partial_i: A \rightarrow A$, $i = 1, \dots, m$, be m mutually commuting derivations of A over $\bar{k}$.

Suppose $L = A^N$, $N \leq \infty$, is a differential algebra over $\bar{k}$. That means that if $\bar{x}_i = (0, \dots, x_i, \dots, 0)$ (nonzero element on i^{th} -place), then a multiplication $\cdot$ in L is given by

$$(\bar{x}_i \cdot \bar{y}_j)_k = \sum_{\sigma, \nu \in \mathbb{Z}_+^m} c_{ij, \sigma, \nu}^k \partial^\sigma x_i \cdot \partial^\nu y_j, \quad c_{ij, \sigma, \nu}^k \in A, \quad (0.1)$$

where $\mathbb{Z}_+^m = \{0, 1, 2, \dots\}^m$, $\partial^\sigma = \partial_1^{\sigma_1} \dots \partial_m^{\sigma_m}$ for $\sigma = (\sigma_1, \dots, \sigma_m)$.

If $\mathfrak{g}$ is a finite-dimensional Lie algebra over $\bar{k}$, then σ (understood as a function on $\mathfrak{g}$) one assigns a derivation x_n of the ring $\bar{k}[\text{Sym}(\text{Der}(\mathfrak{g}))]$ is map $\bar{k}[\text{Sym}(\text{Der}(\mathfrak{g}))]$ is Hamiltonian, meaning (see [1]) that

$$x_{(n, f)} = [x_n, x_f] \cdot \nabla n, f \in \text{Sym}(\mathfrak{g}), \quad (0.2)$$

where the Poisson bracket is defined by

$$(n, f) = x_n(f) \quad (0.3)$$

Now I describe what is an analog of this well-known map in the differential case.

1. First we need an analog of functions on L . Let $u_1, \dots, u_m$ be differentially independent variables. Consider a differential ring $A_m = A[u_1, \dots, u_m]$.

*Supported in part by NSF.

with derivations ∂ extended on A_U to act as $\partial_i(u_i^{(\sigma)}) = u_i^{(\sigma+1)}$. If $X \in L$,

$X = (X_1, \dots)$, we denote

$$\langle u, X \rangle = \sum_i u_i X_i \quad (1.1)$$

Denote $\text{Im } \partial = \sum_i \text{Im } \partial_i$ in A_U ; we write $a \equiv b$ if $(a - b) \in \text{Im } \partial$

Definition 1.2. A derivation $V: A_U \rightarrow A_U$ is called evolutionary if $[V, \partial_i] = 0$, $i = 1, \dots, m$.

Note that $V(\text{Im } \partial) \subset \text{Im } \partial$ and V is defined by its action on u_i 's only: $V(u_i^{(\sigma)}) = \partial^\sigma V(u_i)$.

Define the matrix $B \in A_U[[\partial_1, \dots, \partial_m]]$ by

$$\langle u, X \cdot Y \rangle = X^t B Y, \quad \forall X, Y \in L, \quad (1.3)$$

where "t" means "transpose" and elements of L are understood as vectors. From (1.3), (0.1) we get

$$B_{ij} = \sum_{k, c, v} (-1)^{|c|} \partial^c u_k c_{kij, c, v}^v \partial^v \quad (1.4)$$

where $|c| = c_1 + \dots + c_m$.

Finally, we need "functional derivatives." For $P \in A_U$, the vector $\frac{\delta P}{\delta u}$ is defined by $(\frac{\delta P}{\delta u})_i = \frac{\delta P}{\delta u_i}$, where

$$\frac{\delta P}{\delta u_i} = \sum_c (-\partial)^c \left(\frac{\partial P}{\partial u_i^{(c)}} \right) \quad (1.5)$$

We set $\frac{\delta P}{\delta u^t} = \left(\frac{\delta P}{\delta u} \right)^t$.

Proposition 1.6. The matrix B is skew-adjoint iff multiplication in L is skew-commutative.

2. Thereafter we assume B (and L) to be skew. The matrix B generates a map $\Gamma = B\delta$ from A_U into $D^{\text{ev}}(A_U)$ (= evolution derivations of A_U) by

$$x_H(u_i) = \sum_j B_{ij} \left(\frac{\delta H}{\delta u_j} \right), \quad \forall u \in A_U. \quad (2.1)$$

We will write $x_H = B \frac{\delta H}{\delta U}$.

Lemma 2.2. For any $P, Q \in A_U$,

$$\frac{\delta}{\delta U} \left(\frac{\delta P}{\delta U} B \frac{\delta Q}{\delta U} \right) = D \left(\frac{\delta P}{\delta U} \right) B \frac{\delta Q}{\delta U} - D \left(\frac{\delta Q}{\delta U} \right) B \frac{\delta P}{\delta U} + \frac{\delta P}{\delta U} * \frac{\delta Q}{\delta U},$$

where the Frechet derivative $D(\bar{R}), \forall \bar{R} \in (A_U)^N$, is a matrix (differential operator) defined by

$$D(\bar{R})V\bar{U} = V(\bar{R}), \quad \forall V \in D^{\text{ev}}(A_U). \quad (2.3)$$

see e.g., [2], Ch. 1.

The proof is a straightforward analog of the computations in the lemma 7.14, Ch. 1 of [2].

Remark 2.4. One can make A_U into an algebra using the Poisson bracket:

$\{P, Q\} = x_P(Q)$. It is easy to see that, $\forall P, Q, R \in A_U$, $\{P, \{Q, R\}\} + \text{c.p.} \equiv 0$ iff $\cdot$ is Hamiltonian, that is, $x_{\{P, Q\}} = [x_P, x_Q], \forall P, Q \in A_U$.

Now I can formulate the main result.

Theorem 2.5. Γ is Hamiltonian iff L is a Lie algebra.

The proof follows from lemma 2.2.

Remark 2.6. In the case L is an ordinary Lie algebra, $B_{ij} = \sum_k c_{ij}^k u_k$ and the equations of trajectories of the field x_H , $\text{HLS}(L) \in C^1(L^*)$, are familiar

$$\dot{u}_i = \sum_k c_{ij}^k \frac{\partial H}{\partial u_j} \quad (2.7)$$

Thus X_H is tangent to the orbits of the coadjoint representation of a "group G " (whose Lie algebra is L). In the general differential situation, the notion of "orbits" is meaningless. That is why one needs to work algebraically. (The alternative would be either tight restrictions of the "compactness" type or pyramids of functional analysis.)

As a corollary, from the theorem 2.5 one gets

Proposition 2.8. The map $\theta: L \rightarrow D^{ev}(A_U), L \ni X \mapsto X_{-<u, \cdot>}$, is a Lie algebra homomorphism (assuming L is a Lie algebra).

3. Denote by $L_{(s)} = \{X \in L \mid X_i = 0, 1 \leq i \leq s\}$. Consider the matrix B^s defined by

$$B_{ij}^s = \sum_{k > s; \sigma, \nu} (-1)^{|\sigma|} \delta_{ij}^{\sigma, \nu} u_k c_{ij, \sigma, \nu}^k \cdot \partial^\nu, \quad 1, j > s. \quad (3.1)$$

Proposition 3.2. The matrix B^s defines a Hamiltonian structure (i.e., if H, F depend upon u_i with $i > s$ then the same is true for $\{H, F\}$) iff $L_{(s)}$ is a Lie subalgebra in L .

4. Suppose $L_1 = A^M$, $M \leq \infty$, is another algebra over $\bar{k}$, and let $\phi: L \rightarrow L_1$ is a map given by a linear differential operator. Denote by $A_\nu = A[v_j^{(\nu)}]$, $j \leq M$, the differential algebra which plays for L_1 the same role as A_U plays for L . We can define the map $\phi^*: A_\nu \rightarrow A_U$ by

$$\begin{cases} 1) \quad \phi^* \partial_i = \partial_i \phi^*, \quad 2 \leq i \leq m; \\ 2) \quad \langle \phi^* u, X \rangle \equiv \langle u, \phi(X) \rangle, \quad \forall X \in L. \end{cases} \quad (4.1)$$

Suppose L and L_1 are Lie algebras, thus corresponding matrices B_U and B_ν define Hamiltonian structures. We say that ϕ^* is canonical if $\Gamma_U(\phi^* F)$ is ϕ^* -compatible with $\Gamma_\nu(F)$, $\forall F \in A_\nu$ (see motivations in [1], Ch. IV).

Proposition 4.2. ϕ^* is canonical iff ϕ is a Lie algebra homomorphism.

5. Consider some examples.

5.1. Let $A[\xi] = \left\{ \sum_{|\sigma| \in \mathbb{N}^m} a_\sigma \xi^\sigma \mid a_\sigma \in A, \xi^\sigma = \xi_1^{\sigma_1} \dots \xi_m^{\sigma_m} \right\}$ be a ring of differential

operators and $A[\xi, \xi^{-1}]$ the corresponding ring of pseudo-differential operators.

For the Lie algebra generated by elements of $A[\xi, \xi^{-1}]$ of negative order, corresponding matrix B (for $m = 1$) was introduced by Manin ([2], Ch. I) as a formal limit of Gel'fand-Dikii operators. The Lie-algebraic interpretation of that matrix (for $m = 1$) was given in [3].

5.2. Consider $C^\infty(\mathbb{R}^{2m}) = C^\infty(x, \xi)$ and its localization around subspace $\xi_1 = \dots = \xi_m = 0$. The usual Poisson bracket on $\mathbb{R}^{2m} = T^*(\mathbb{R}^m)$ leads to the following algebraic set up: $L = \left\{ \sum_{\mu \in \mathbb{Z}_+^m} x_\mu \xi^\mu \mid x_\mu \in A \right\} = ((x_\mu) \mid x_\mu \in A)$.

The bracket on L is given by

$$(x^*y)_\mu = \sum_{\sigma+\tau=\mu+1} [c_{\sigma, \tau} x_\sigma \partial_\tau (y_\tau) - \tau c_{\tau, \sigma} y_\tau \partial_\sigma (x_\sigma)] \quad (5.3)$$

thus the corresponding Hamiltonian matrix B is

$$B_{\sigma, \tau} = \sum_{\ell=1}^m [c_{\ell, \sigma+\tau-1} \partial_\ell + c_{\ell, \tau} \partial_\ell] \quad (5.4)$$

This matrix appears in m -D hydrodynamics. For $m = 1$ it was introduced in [4]; its Lie-algebraic interpretation is given in [5] (for $m = 1$).

5.5. In $\mathbb{R}^{r+1} = ((x, \xi_1, \dots, \xi_r))$ consider the Lie algebra of functions on $\mathbb{R}^{r+1}$, with the bracket

$$[f, g] = \sum_{Y=1}^r \left(\partial_{\xi_Y} \frac{\partial}{\partial x} f \cdot \partial_Y g - \partial f \cdot \partial_Y \frac{\partial}{\partial \xi_Y} g \right), \quad \partial = \partial_1 = \partial / \partial x \quad (5.6)$$

Localization to the line $\xi = 0$ leads to the algebra $L = \left\{ \sum_{\mu \in \mathbb{Z}_+^r} x_\mu \xi^\mu \mid x_\mu \in A, \partial_1 = \partial / \partial x \right\}$

with Lie bracket

$$(X*Y)_\psi = \sum_{a+b=\psi+1} (a_Y d_Y X_a \partial Y_b - b_Y d_Y Y_a \partial X_b), \quad a, b \in \mathbb{Z}_+^r \quad (5.7)$$

The corresponding matrix B is given by

$$B_{\phi\psi} = \sum_{\gamma=1}^r (\phi_Y d_Y u_{\phi+\psi-1} \partial_\gamma + \psi_Y d_Y u_{\phi+\psi-1} \partial_\gamma) \quad (5.8)$$

which is the multicomponent (for $r > 1$) generalization of the $m=1$ -case (5.4).

5.9. In $\mathbb{R}^{n+p}$ with coordinates $(y_1, \dots, y_p; x_1, \dots, x_n)$ consider Lie algebra $\bar{L}$ consisting of those vector fields $X \in \mathfrak{X}(\mathbb{R}^{n+p})$ which commute with $\frac{\partial}{\partial y_q}$, $q = 1, \dots, p$.

The algebraic version of $\bar{L}$ would be the Lie algebra $\mathfrak{A}(A) \bigoplus_{q=1}^p A \frac{\partial}{\partial y_q}$ acting on $A[y_1, \dots, y_p]$ as derivations. The corresponding matrix B now yields Poisson bracket

$$\begin{aligned} (F, G) &= \sum_{q=1}^p \sum_{\ell=1}^n \left(\frac{\delta F}{\delta u_\ell} \partial_\ell \frac{\delta G}{\delta v_q} - \frac{\delta G}{\delta u_\ell} \partial_\ell \frac{\delta F}{\delta v_q} \right) + \\ &+ \sum_{\ell=1}^n u_\ell \sum_{i=1}^n \left(\frac{\delta F}{\delta u_i} \partial_i \frac{\delta G}{\delta u_\ell} - \frac{\delta G}{\delta u_i} \partial_i \frac{\delta F}{\delta u_\ell} \right), \end{aligned} \quad (5.10)$$

which appeared in [6] (for $n=1, p=2$) in the context of hydrodynamics of compressible fluid.

5.11. On $C^\infty(\mathbb{R}^2)$ consider Lie algebra structure $[f, g] = (\alpha \partial_x f + \beta \partial_y f + \gamma \partial_x g + \delta \partial_y g) \cdot \partial$, $\partial = \partial/\partial x$, $\alpha, \beta \in \mathbb{R}$. Localization on the line $x=0$ leads to the following algebraic set up: $L = A^\infty = \{(X_i)\}_{i=1}^{\infty}$, $m=1$, with multiplication

$$(X*Y)_k = d_k \sum_{n+m=k} (X_n \partial Y_m - Y_m \partial X_n), \quad \partial = \partial/\partial x = \partial_1, \quad (5.12)$$

where $d_k = \alpha k + \beta$, and $\alpha, \beta \in \mathbb{C}$ are arbitrary. The corresponding matrix $B[7]$

$$B_{ij} = d_i u_{i+j} \partial + \partial d_j u_{i+j} \quad (5.13)$$

is the main source of integrable systems in two-space dimensions. Classification table of such systems (see, e.g. [7]) smoothly upon $(\alpha:\beta) \in \mathbb{CP}^1$. Despite of this, one can prove the following

Theorem 5.14. The Lie algebra L with multiplication (5.12) admits a nonzero ad-invariant (modulo $\text{Im } \partial$), bilinear (differential) form (not necessarily symmetric) iff $\frac{3\beta}{\alpha} \in \mathbb{Z}$:

$$(X, Y) = \sum_{i+j+3\beta\alpha-1=0} X_i Y_j \quad (5.15)$$

6. Remark. The existence of Hamiltonian structures on dual spaces of Lie algebras is not the privilege of the differential case only. Analogous results can be found: a) when one substitutes commuting automorphisms $\Delta_1, \dots, \Delta_m: A \rightarrow A$ instead of derivations, as in [8]; b) and moreover, when both derivations and automorphisms are present. The corresponding matrices B behave naturally with respect to "continuum limits" $\Delta_\lambda \rightarrow \exp(\lambda \partial_\lambda)$ [8]. Details will be given elsewhere.

REFERENCES

- [1] B. A. Kupershmidt, "Geometry of Jet Bundles and the Structure of Lagrangian and Hamiltonian Formalisms," in Lect. Notes Math. #775 (1980), 162-218, Springer-Verlag.
- [2] Yu. I. Manin, "Algebraic Aspects of Nonlinear Differential Equations," J. Sov. Math. 11, (1979), 1-122.
- [3] D. R. Lebedev and Yu. I. Manin, "Hamiltonian Operator of Gel'fand-Dikii and Coadjoint Representation of Volterra Group," Funk. Anal. Appl. 13:4 (1979), 40-46.
- [4] B. A. Kupershmidt and Yu. I. Manin, "Equations of Long Waves with a Free Surface II. Hamiltonian Structure and Higher Equations," Funk. Anal. Appl. 12:1 (1978), 25-32.
- [5] D. R. Lebedev, "Benney's Long Waves Equations: Hamiltonian Formalism," Lett. Math. Phys. 3 (1979), 481-488.

-
- [6] P. J. Morrison and J. M. Greene, "Noncanonical Hamiltonian Density Formulation of Hydrodynamics and Ideal Magnetohydrodynamics," *Phys. Rev. Lett.* **45**:10 (1980), 790-794.
- [7] B. A. Kupershmidt, "Deformations of Integrable Systems," *Proc. Roy. Irish Acad.*
- [8] B. A. Kupershmidt, "On Algebraic Models of Dynamical Systems," *Lett. Math. Phys.*
-

Department of Mathematics
The University of Michigan
Ann Arbor, MI 48109