2 ...
DOE/PC/90302--T3

16 1991

DOE/PC/90302--T3

DE92 004860

PRINCIPAL INVESTIGATORS: Robert L. Robinson, Jr.
K. A. M. Gasem

AFFILIATION: School of Chemical Engineering
Engineering North 423
Oklahoma State University
Stillwater, OK 74078

Received by OSTI

DEC 30 1991

TELEPHONE: (405) 744-5280

PROJECT TITLE: "Equilibrium and Volumetric Data and Model Development for
Coal Fluids"

PROJECT PERIOD: April 15, 1991 to July 15, 1991

SUMMARY

During the present reporting period, experimental measurements were obtained on the solubility of carbon monoxide (CO) in selected hydrocarbons using our static type equilibrium cell. Binary mixtures involving CO + n-eicosane (n-C₂₀), n-octacosane (n-C₂₈), and n-hexatricosane (n-C₃₆) were studied at temperatures from 323.2 to 423.2 K and pressures to 104 bar. These data were analyzed using the Scave-Redlich-Kwong and Peng-Robinson equations of state. While a single interaction parameter, C_{ij} , describes the n-C₂₀ and n-C₂₈ measurements within 0.004 in mole fraction over the full temperature range, temperature dependent parameters are required to adequately fit the n-C₃₆ data. The present data are in reasonable agreement with the earlier measurements of Huang and coworkers: in general, deviations within 0.004 in mole fraction are observed.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

DE-FG22-90PC90302

EQUILIBRIUM AND VOLUMETRIC DATA AND MODEL
DEVELOPMENT FOR COAL FLUIDS

Reviewed by OSTI

DEC 30 1991

Report for the Period
April 15, 1991 to July 15, 1991

R. L. Robinson, Jr.
(Principal Investigator)

K. A. M. Gasem
(Principal Investigator)

S. Srivatsan

X. Yi

Oklahoma State University
School of Chemical Engineering
Stillwater, Oklahoma 74078-0537

PREPARED FOR THE UNITED STATES
DEPARTMENT OF ENERGY

EQUILIBRIUM AND VOLUMETRIC DATA AND MODEL

DEVELOPMENT FOR COAL FLUIDS

ABSTRACT

The proper design, operation, and optimization of coal conversion and utilization processes depend heavily on knowledge of phase behavior of the mixtures encountered in these processes. Frequently quoted statistics suggest that in conventional chemical plants 70% of the capital and 90% of the operating expenses are associated with phase separations. Since multiple stages occur in almost all stages of coal conversion process - from feed preparation through conversion reactions to product separation - the proper description of phase behavior is important in each step of these processes.

The long term goal of our efforts is to develop accurate predictive methods for description of equilibrium phase properties for a variety of types of mixtures and operating conditions. The specific objectives of the work specified herein include:

- (1) development of an experimental facility having the capability to provide data on equilibrium phase compositions (solubilities) and liquid densities, and doing so with greater accuracy and speed than our previous facility.
- (2) measurement of equilibrium phase properties for systematically selected mixtures - specifically those containing important solute gases (such as hydrogen, carbon monoxide, carbon dioxide, methane, ethane, carbonyl sulfide, ammonia) in a series of heavy paraffinic, naphthenic and aromatic solvents (e.g., n-decane, n-eicosane, n-triacontane, n-tetracontane, cyclohexane, decalin,

perhydrophenanthrene, perhydropyrene, benzene, naphthalene, phenanthrene, pyrene),

- (3) testing/development of correlation frameworks for representing the phase behavior of fluids of the type encountered in coal conversion processes, and
- (4) generalization of parameters in the correlation frameworks to enable accurate predictions for systems of the type studied, permitting predictions to be made for systems and conditions other than those for which experimental data are available.

During the present reporting period, experimental measurements were obtained for the solubility of carbon monoxide (CO) in selected hydrocarbons using our static type equilibrium cell. Binary mixtures involving CO + n-eicosane (n-C₂₀), n-octacosane (n-C₂₈), and n-hexatricosane (n-C₃₆) were studied at temperatures from 323.2 to 423.2 K and pressures to 104 bar. These data were analyzed using the Soave-Redlich-Kwong (SRK) and Peng-Robinson (PR) equations of state. While a single interaction parameter, C_{ij} , describes the n-C₂₀ and n-C₂₈ measurements within 0.004 in mole fraction over the full temperature range, temperature dependent parameters are required to adequately fit the n-C₃₆ data. The present data are in reasonable agreement with the earlier measurements of Huang and coworkers; in general, deviations within 0.004 in mole fraction are observed.

INTRODUCTION

Essentially all state-of-the-art models for phase behavior contain one [1, 2], two [3] or three [4] interaction parameters to account for unlike molecular pair interactions. These parameters, in general, cannot be predicted a priori from existing theory, but can be established from

experimental measurements on the binary pairs which form the system of interest. These "empirical" interaction parameters have a dramatic effect on the predicted properties of mixtures and are thus required for accurate predictions. In most instances, successful modeling of the behavior of complex multicomponent mixtures requires accurate information on the pure compounds and on the binary interactions that exist between the different molecular species.

Experimental measurements made on systematically chosen mixtures can be used to evaluate interaction parameters and, more importantly, furnish the basis for generalization of the parameters to allow interpolation (and perhaps extrapolation) to other solvents in the same homologous series. Towards this end we have previously reported and analyzed data on the solubility of carbon dioxide, ethane and methane in a series of hydrocarbons [5-10]. In the present report, we complement our previous studies with new solubility, x , (or equivalently, bubble point pressure, p) measurements involving carbon monoxide and a number of heavy hydrocarbons. Specifically, *n*-eicosane, *n*-octacosane, *n*-hexatricosane - all solids at room temperature - were selected from the normal paraffin homologous series for this purpose.

MATERIALS

The carbon monoxide used in this study had a stated purity of 99.99+ mol% and was purchased from Matheson Gas Products. The *n*-eicosane and *n*-hexatricosane were supplied by Aldrich Chemical Company and had a stated purity of 99+ and 98 mol%, respectively. The *n*-octacosane was obtained from Alfa Products with a stated purity of 99 mol%. No further purification of the chemicals was attempted.

DISCUSSION OF TECHNICAL RESULTS

Carbon monoxide solubility in n-eicosane, n-octacosane, and n-hexatriacontane were measured at temperatures from 323.2 to 423.2 K and pressures to 104 bar. The experimental data are presented in Tables I-III. The newly-acquired data were correlated using SRK [1] and PR [2] equations of state (EOS). The uncertainty in measured bubble point pressure, which depends on the p-x profile for each isotherm, is of the order of 0.5 bar. Optimum EOS binary interaction parameters were obtained by minimizing the sum of squares of pressure deviations from the experimental values. Detailed procedures for data reduction are given elsewhere [9]. The input parameters of the pure components (acentric factors, critical temperatures and critical pressures) required by the SRK and PR equations of state, together with the literature sources, are presented in Table IV.

The equation-of-state representation of the carbon monoxide solubilities for the systems studied are documented in Tables V-VII. The SRK equation of state is capable of describing n-C₂₀ and n-C₂₈ data with RMS errors of 0.004 in mole fraction when a single interaction parameter, C_{ij} , is used for each system over the complete temperature range. For these two binaries, only marginal improvements in the equation of state predictions are realized when a second interaction parameter, D_{ij} , is also employed. For the n-C₃₆ system, however, the use of either one or two interaction parameters, over the complete temperature range does not fit the data as well (RMS errors of 0.007 in mole fraction using two interaction parameters C_{ij} and D_{ij}).

From Tables V-VII, it is apparent that the values obtained for the interaction parameters, C_{ij} and D_{ij} , when lumped over the complete temperature range lie outside the range of values obtained for the parameters at each

isotherm. This is possible since two parameters are used while regressing the data set.

When one interaction parameter, C_{ij} , is fitted to each isotherm, for the two systems n-C₂₀ and n-C₂₈, the RMS errors are less than 0.0015 in mole fraction (except for the CO + n-C₂₈ system at 348.15 K) and the two equations of state give comparable representation of data. No significant improvements are obtained for either CO + n-C₂₀ or CO + n-C₂₈ by using two parameters for each isotherm (except for the CO + n-C₂₈ system at 348.15 K) as revealed by the results in Tables V and VI. For the n-C₃₆ system, however, the use of a second interaction parameter, D_{ij} , for each isotherm improves the predictions significantly as shown by the results in Table VII. The need for a second interaction parameter may reflect an inadequacy in the original EOS mixing rules for the highly asymmetric mixtures produced by the larger n-C₃₆ molecules.

The RMS errors in mole fraction are, in general, within the estimated uncertainty of 0.001 when the full potential of EOS model is realized using two interaction parameters (C_{ij} and D_{ij} , per isotherm. The results obtained illustrate both the ability of the equation of state and the precision of our reported data.

Comparisons of our results with those reported by Huang, et al. [12] appear in Figures 1-3. The comparisons are shown in terms of deviations of the solubilities from values predicted using the SRK equation of state. Interaction parameters, C_{ij} and D_{ij} , employed in equation of state predictions were obtained by fitting our data for the isotherm under study. Reasonable agreement (solubility deviations < 0.004) are obtained for the 212°F isotherm

between this work and that of Huang and coworkers for all three systems. As indicated by Figures 1 and 3, higher bubble point pressures are obtained by Huang for the n-C₂₈ and n-C₃₆ systems at 212°F.

CONCLUSIONS

Data have been obtained for the solubility of carbon monoxide in n-eicosane, n-octacosane, and n-hexatriacontane at temperatures from 323.2 to 423.2 K and pressures up to 104 bar. These data are well described by the SRK and PR equations of state. These results will be of value in establishing equation of state interaction parameters for light gases in normal paraffins.

REFERENCES

1. Soave, G., Chem. Eng. Sci., 27, 1197-1203 (1972).
2. Peng, Y. D.; Robinson, D. B., Ind. Eng. Chem. Fund., 15, 59-64 (1976).
3. Turek, E. A.; Metcalfe, R. S.; Yarborough, L.; Robinson, R. L., Jr., J. Soc. Petrol. Eng., 24, 308-324 (1984).
4. Assadipour, H.; Hajela, V., "New Mixing Rules for Parameters of a Three Parameter Equation of State", Paper presented at AIChE meeting, New Orleans, April 6-10 (1986).
5. Robinson, R. L., Jr.; Anderson, J. M.; Barrick, M. W.; Bufkin, B. A.; Ross, C. H., "Phase Behavior of Coal Fluids: Data for Correlation Development," DE-FG22-86PC90523, Final Report, Department of Energy, January (1987).
6. Gasem, K. A. M.; Robinson, R. L., Jr., J. Chem. Eng. Data, 30, 53-56 (1985).
7. Anderson, J. M.; Barrick, M. W.; Robinson, R. L., Jr., J. Chem. Eng. Data, 31, 172-175 (1986).
8. Raff, A. M., M.S. Thesis, Oklahoma State University, Stillwater, Oklahoma (1989).
9. Gasem, K. A. M., Ph.D. Dissertation, Oklahoma State University, Stillwater, Oklahoma (1986).
10. Darwish, N. A., Ph.D. Dissertation, Oklahoma State University, Stillwater, Oklahoma (1990).
11. Reid, R. C.; Prausnitz, J. M.; Sherwood, T. K., "The Properties of Gases and Liquids," 2nd Edition, McGraw Hill, New York (1977).
12. Huang, S. H.; Lin, H. M.; Chao, K. C., Fluid Phase Equilibria, 36, 141-148 (1987).

TABLE I
SOLUBILITY DATA FOR CARBON MONOXIDE IN N-EICOSANE

Mole fraction Carbon Monoxide	Bubble Point Pressure	
	bar	(psia)
- - - - - 323.2 K (50.0°C, 122.0°F) - - - - -		
0.0516	28.0	(407)
0.0645	35.6	(516)
0.0797	45.1	(654)
0.0902	51.6	(749)
0.1024	60.5	(878)
0.1136	67.8	(983)
- - - - - 373.2 K (100.0°C, 212.0°F) - - - - -		
0.0403	19.9	(289)
0.0516	25.8	(373)
0.0715	36.5	(529)
0.0899	47.2	(684)
0.1107	59.3	(860)
0.1308	72.4	(1049)
- - - - - 423.2 K (150.0°C, 302.0°F) - - - - -		
0.0616	28.3	(411)
0.0761	35.7	(518)
0.1004	48.3	(700)
0.1187	58.6	(850)
0.1205	59.4	(861)
0.1390	82.4	(1196)
0.1393	82.7	(1200)
0.1614	83.8	(1216)

TABLE II
SOLUBILITY DATA FOR CARBON MONOXIDE IN N-OCTACOSANE

Mole fraction Carbon Monoxide	Bubble Point Pressure	
	bar	(psia)
- - - - - 348.2 K (75.0°C, 167.0°F) - - - - -		
0.0553	23.3	(338)
0.0604	25.5	(370)
0.0854	38.2	(554)
0.1014	46.2	(671)
0.1320	63.7	(923)
0.1493	75.7	(1097)
- - - - - 373.2 K (100.0°C, 212.0°F) - - - - -		
0.0519	22.5	(326)
0.0718	31.6	(458)
0.0881	39.7	(576)
0.1042	47.6	(690)
0.1157	53.9	(782)
0.1290	60.2	(873)
- - - - - 423.2 K (150.0°C, 302.0°F) - - - - -		
0.0522	20.8	(301)
0.0733	31.3	(455)
0.1038	43.3	(629)
0.1257	53.7	(779)
0.1502	66.3	(961)

TABLE III

SOLUBILITY DATA FOR CARBON MONOXIDE IN N-HEXATRICONTANE

Mole fraction Carbon Monoxide	Bubble Point Pressure	
	bar	(psia)
- - - - - 373.2 K (100.0°C, 212.0°F) - - - - -		
0.0494	18.0	(261)
0.0634	23.4	(340)
0.0899	34.4	(499)
0.1192	47.5	(689)
0.1603	67.6	(980)
0.2002	89.5	(1299)
- - - - - 423.2 K (150.0°C, 302.0°F) - - - - -		
0.0568	18.5	(269)
0.0705	23.7	(344)
0.1040	36.0	(522)
0.1289	46.6	(676)
0.1742	66.1	(958)
0.2099	84.0	(1218)

TABLE IV
CRITICAL PROPERTIES AND ACENTRIC FACTORS USED
IN THE SRK AND PR EQUATIONS OF STATE

Component	Pressure bar	Temperature K	Acentric Factor	Reference
Carbon Monoxide	34.94	132.9	0.0491	11
n-Eicosane	11.17	770.5	0.8738	9
n-Octacosane	8.26	845.4	1.1073	9
n-Hexatria- contane	6.91	901.1	1.2847	9

TABLE V

SRK AND PR EQUATION OF STATE REPRESENTATIONS
OF CARBON MONOXIDE SOLUBILITY IN N-EICOSANE

Temperature K (°F)	Soave Parameters (P-R Parameters)		Error in Mole Fraction (Using P-R Equation)*	
	C_{ij}	D_{ij}	RMS	MAX
323.2 (122.0)	0.2772	-0.0113	0.0003	0.0006
	(0.2598)	(-0.0112)		
	0.1502		0.0006	0.0009
	(0.1386)			
373.2 (212.0)	0.2257	-0.0105	0.0002	0.0004
	(0.2205)	(-0.0123)		
	0.1166		0.0006	0.0009
	(0.1214)			
423.2 (302.0)	0.2292	-0.0140	0.0002	0.0002
	(0.2219)	(-0.0185)		
	0.2373		0.0008	0.0012
	(0.2362)			
323.2, 373.2 423.2	0.1439	-0.0018	0.0040	0.0072
	(-0.1357)	(-0.0323)	(0.0053)	(0.0113)
	0.1253		0.0041	0.0073
	(0.1271)		(0.0052)	(0.0096)

* The RMS and maximum errors in CO mole fraction are essentially the same for both the SRK and PR equations of state whenever not mentioned.

TABLE VI
SRK AND PR EQUATION OF STATE REPRESENTATIONS
OF CARBON MONOXIDE SOLUBILITY
IN N-OCTACOSANE

Temperature K (°F)	Soave Parameters (P-R Parameters)		Error in Mole Fraction (Using P-R Equation)*	
	C_{ij}	D_{ij}	RMS	MAX
348.2 (167.0)	0.4522 (0.4341)	-0.0258 (-0.0270)	0.0005	0.0009
	0.0935 (0.1128)		0.0027	0.0038
373.2 (212.0)	0.0861 (0.1145)	-0.0008 (-0.0019)	0.0004	0.0008
	0.0738 (0.0918)		0.0004	0.0008
423.2 (302.0)	-0.0522 (0.0072)	0.0048 (0.0023)	0.0014	0.0025
	0.0131 (0.0349)		0.0014	0.0029
348.2, 373.2 423.2	0.4495 (0.3541)	-0.0264 (-0.0218)	0.0027 (0.0036)	0.0070 (0.0090)
	0.0743 (0.0916)		0.0036 (0.0042)	0.0079 (0.0089)

* The RMS and maximum errors in CO mole fraction are essentially the same for both the SRK and PR equations of state whenever not mentioned.

TABLE VII

SRK AND PR EQUATION OF STATE REPRESENTATIONS
OF CARBON MONOXIDE SOLUBILITY
IN N-HEXATRIACONTANE

Temperature K (°F)	Soave Parameters (P-R Parameters)		Error in Mole Fraction (Using P-R Equation)*	
	C_{ij}	D_{ij}	RMS	MAX
373.2 (212.0)	0.1966	-0.0115	0.0001	0.0002
	(0.2185)	(-0.0127)		
	0.0073		0.0024	0.0028
	(0.0431)			
423.2 (302.0)	0.1328	-0.0148	0.0005	0.0008
	(0.1688)	(-0.0172)		
	-0.1154		0.0026	0.0034
	(-0.0672)			
373.2, 423.2	0.3102	-0.0212	0.0068	0.0117
	(0.2199)	(-0.0162)	(0.0075)	(0.0116)
	-0.0409		0.0080	0.0144
	(-0.0021)		(0.0085)	(0.0152)

* The RMS and maximum errors in CO mole fraction are essentially the same for both the SRK and PR equations of state whenever not mentioned.

Figure 1. Comparison of Carbon Monoxide Solubilities in n-Eicosane

LEGEND $\square$ - $\square$ - $\square$ This Work: 167 F Δ - Δ - Δ This Work: 212 F $\circ$ - $\circ$ - $\circ$ This Work: 302 F
 $\ast$ - $\ast$ - $\ast$ Huang: 212 F

Figure 2. Comparison of Carbon Monoxide Solubilities in n-Octacosane

LEGEND $\square$ $\square$ $\square$ This Work: 212 F Δ Δ Δ This Work: 302 F
 $\circ$ $\circ$ $\circ$ Huang: 212 F

Figure 3. Comparison of Carbon Monoxide Solubilities in n-Hexatriacontane

END

**DATE
FILMED**

2 / 18 / 92