Models of Fragmentation Phenomena Based on the Symmetric Group S_n and Combinational Analysis

DOE/ER/40561--15
DE92 004368

A.Z. Mekjian

Institute for Nuclear Theory
University of Washington, HN-12, Seattle, Washington 98195
USA

and

S.J. Lee

Department of Physics and Astronomy
Rutgers University, Piscataway, N.J. 08854
USA

PREPARED FOR THE U.S. DEPARTMENT OF ENERGY

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Department of Energy, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the product or process disclosed, or represents that its use would not infringe privately-owned rights.

By acceptance of this article, the publisher and/or recipient acknowledges the U.S. Government's right to retain a nonexclusive, royalty-free license in and to any copyright covering this paper.

MASTER

Models of Fragmentation Phenomena based on the Symmetric Group S_n and Combinational Analysis

A.Z. Mekjian*

*Institute of Nuclear Theory, HN-12, University of Washington, Seattle, WA 98195,
U.S.A.*

and

S.J. Lee

*Dept. of Physics and Astronomy, Rutgers University, Piscataway, New Jersey
08855, U.S.A.*

Abstract

Various models for fragmentation phenomena are developed using methods from permutation groups and combinational analysis. The appearance and properties of power laws in these models are discussed. Various exactly soluble cases are studied.

* Permanent address: Dept. of Physics and Astronomy, Rutgers University, Piscataway, New Jersey 08855, U.S.A.

A.Z.M. : Phone; (206) 685-3633, E-mail; mekjian%bethe@phast.phys.washington.edu

S.J.L. : Phone; (908) 932-4742, E-mail; sjlee@ruthep.rutgers.edu

1 Introduction

Fragmentation phenomena are observed in many different areas of physics. A fragmentation process produces a cluster distribution of objects following a collision or some other method of excitation or deexcitation. The mass or size distribution which results from some type of process is of interest and this distribution gives the number of clusters of a given size as a function of their size. Examples are nuclear fragment distributions following a collision of two heavy ions [1], meteoritic distributions reaching the earth [2], earthquake sizes [3], avalanche [4] and sandpile [5] slides, lunar crater sizes [6], percolation cluster distributions [7], droplet sizes [8] in aerosols and in condensation phenomena. One remarkable feature in these examples is the appearance of a power law in the size distribution. That is, the distribution of sizes falls as a power of the size of the object. Power laws appear in many other areas. Examples are Zipf's law [9] in linguistics for word frequency use, Pareto's law [10] in economics for income distributions, and $1/f$ noise [11]. Moreover, $1/f$ noise appears in many phenomena [5] from starlight flicker to sand flow and traffic flow. A more detailed summary of power laws can be found in Mandelbrot [12] and in Montroll and West [13].

In a previous set of papers [14, 15], a model for a fragmentation process was introduced and developed. A connection of this model with a thermodynamical description of a nuclear fragmentation process is given in Ref.[16]. This paper considers further developments, extensions, and applications of this model. The various cases considered all have hyperbolic power law behavior at some particular point. A link between these models and the symmetric group S_n is developed, and combinational methods are also employed in our studies.

An outline of this paper is as follows. In Section 2, a relationship between frag-

mentation partitions and permutation cycles of the symmetric group S_n is developed. Canonical ensemble partition functions are derived in Section 3 and used to obtain general expressions for the distribution of cluster sizes in fragmentation phenomena. Section 4 considers various exactly soluble cases. This model is also related to probability theory. Section 5 is for the concluding remarks. Some generating functions are given in Appendix.

2 Partitions, Decompositions, Fragmentations, and Permutation Cycles

In a fragmentation process, an object made of A elements is divided into pieces of smaller size. Specifically, the initial A elements end up in clusters of varying sizes characterized by the number of elements j in the cluster. The number of objects of size j is here called n_j and a sum rule

$$A = \sum_{j=1}^A j n_j \quad (1)$$

exists as a constraint. The listing of the partitions of A objects into such groups also appears in number theory as the decomposition of an integer A into integer summands without regard to order. The integer j will be identified with the cluster of size j . The notation

$$\Pi_A = \Pi_A(\{n_j\}) = (1^{n_1}, 2^{n_2}, \dots, A^{n_A}) = (n_1, n_2, \dots, n_A) \quad (2)$$

is used to specify a particular decomposition of an integer A , a particular fragmentation of an object made of A elements, or division of A objects into some groups. The multiplicity of a given partition is

$$m = n_1 + n_2 + \dots + n_A = \sum_{j=1}^A n_j. \quad (3)$$

The possible values of m are $m = 1, 2, \dots, A$. The number of possible partitions of A without regard to order is called $p(A)$ and the number of possible partitions of A into m parts is called $p(A, m)$. Here $p(A) = \sum_{m=1}^A p(A, m)$. Table 1 lists all the partitions for $A = 5$ and 6.

For large A , the total number of partitions of A is the Hardy-Ramanujan asymptotic result [17, 18]

$$p(A) \approx \frac{1}{4\sqrt{3}A} e^{\pi\sqrt{2A/3}}. \quad (4)$$

Also for large A and $m \ll A$ ($m \sim \mathcal{O}(A^{1/3})$),

$$p(A, m) \approx \frac{A^{m-1}}{m!(m-1)!}. \quad (5)$$

Generating functions for $p(A)$ and $p(A, m)$ are given in the appendix.

The $p(A)$ and $p(A, m)$ also appear in level densities. Substitution of $A = g_o E$ turns Eq.(4) into the level density of a Fermi gas [19] and Eq.(5) into the level density for states made of m particles plus holes [20]. Specifically,

$$\begin{aligned} \rho(E) &= g_o p(A = g_o E) = \frac{1}{4\sqrt{3}E} e^{2(\pi^2 g_o E/6)^{1/2}}, \\ \rho_m(E) &= g_o p(A = g_o E, m) = g_o \frac{(g_o E)^{m-1}}{m!(m-1)!}. \end{aligned} \quad (6)$$

Here the g_o is the level density parameter. Moreover a generalization of Eq.(4) can be used to study statistical mechanics of cosmic bosonic strings [21] and the Hagedorn level density of strong interactions [22]. The importance of $p(A)$ here is in counting the number of possible fragmentations. Each fragmentation is specified by $\Pi_A(\{n_i\})$ of Eq.(2) so that $\sum_{\Pi_A} 1 = p(A)$ where the sum is over all partition of A , i.e., all sets of $\{n_i\}$ which satisfy the constraint Eq.(1). For a collision involving $^{235}\text{U}r + ^{235}\text{U}r$, the number of possible fragmentations $p(A)$ is of order 10^{20} .

The partitioning of A also appears in permutation groups. In Ref.[14] a correspondence between clusters in a fragmentation process and cycles of the permutation

group was established. Specifically, a cluster of j objects corresponds to a cycle of length j , and n_j is either the number of clusters or cycles of that size. The Π_A of Eq.(2) is then a classification of a permutation of A objects by its cycle class structure. The group is called symmetric group S_n , with $n = A$ here, and S_n has $A!$ elements. The $p(A)$ counts the total number of cycle classes. The number of permutations of A objects in a specific cycle class with n_1 unit cycles, n_2 cycles of length 2, etc, is given by Cauchy's number:

$$M_2(A, \{n_j\}) = \frac{A!}{\prod_{j=1}^A j^{n_j} n_j!} \quad (7)$$

The number of cycle classes $p(A)$ is also the number of irreducible representations of S_A [23]. The correspondence between the partitioning of an integer, cycles of a permutation, and fragmentation possibilities of an object is shown in Fig.1.

A partition is sometimes specified by a decreasing set of numbers λ_i defined by the transformation

$$\begin{aligned} \lambda_1 &= n_1 + n_2 + \dots + n_A \\ \lambda_2 &= n_2 + \dots + n_A \\ &\vdots \\ \lambda_A &= n_A \end{aligned} \quad (8)$$

Here $A = \sum_i \lambda_i$ and $m = \lambda_1$. Note also $\lambda_1 \geq \lambda_2 \geq \lambda_3 \geq \dots$ and $n_1 = \lambda_1 - \lambda_2$, $n_2 = \lambda_2 - \lambda_3$, etc. For example, the division of 5 into $3 + 2$ which is $\vec{n} = (2^1, 3^1) = (0, 1, 1, 0, 0)_n$ in the n_i representation becomes $(2, 2, 1, 0, 0)_\lambda$ in the λ_i representation or $\vec{\lambda} = (2, 2, 1)_\lambda = (2^2, 1)_\lambda$ where zeros are dropped. The partition $2+2+1$ is represented by $\vec{n} = (2^2, 1)$ in the $\vec{n}$ representation and by $\vec{\lambda} = (3, 2)$ in the representation $\vec{\lambda}$. For each partition in the $\vec{n}$ representation, there is a partition with the same structure in the $\vec{\lambda}$ representation. For a given structure $(2^2, 1)$ as an example, $\vec{\lambda} = (2^2, 1)$ represents the partition of $3 + 2$, and $\vec{n} = (2^2, 1)$ represents the partition of $2 + 2 + 1$. The $A = \sum_i \lambda_i$ can also be considered as the partition of an integer A into summands of smaller integers λ_i . For example, $\vec{\lambda} = (2^2, 1)$ can be considered as a partition of 5

into $\sum_i \lambda_i = 2 + 2 + 1$ and $\vec{\lambda} = (3, 2)$ as $3 + 2$. Thus, as we can see from Table 1, the representation $\vec{\lambda}$ is conjugate to the representation $\vec{n}$.

The conjugate property between two representations $\vec{n}$ and $\vec{\lambda}$ can easily be visualized in terms of block diagram similar to a Ferrer diagram [17] or a Young tableau [23]. Here we can consider four types of diagrams. In Ferrer's block diagram, illustrated in Fig.2a, the partition $(n_1, n_2, \dots, n_j, \dots, n_A) = \vec{n}$ is represented by n_j rows of j horizontal blocks. They are placed in a diagram in descending order with the longest or largest j at the top. The Ferrer's diagram can also be viewed as a block diagram in $\vec{\lambda} = (\lambda_1, \lambda_2, \dots, \lambda_A)$ space. Here λ_i blocks are put vertically in the i -th column. The total number of rows (λ_1) is the multiplicity m and the area of this diagram (the total number of blocks) is A . A Young tableau in permutation groups is a similar type of diagram but, here, λ_i blocks are put horizontally in the i -th row as shown in Fig.2b. We can obtain a Young tableau from Ferrer's block diagram by first turning the block diagram upside down and then by rotating it through 90° clockwise. This transformation between Ferrer's diagram and Young tableau exhibits the conjugate relation between two representations of $\vec{n}$ and $\vec{\lambda}$. Table 1 also shows this conjugate relationship. Again the total number of blocks is A and the number of columns (λ_1) is the multiplicity. A diagram constructed through the upside down version of a Ferrer's block diagram, Fig.2c, is useful in connecting a partition to the one dimensional avalanche and sandpile slide models [4, 5]. Avalanche models are generated by imposing a condition on block diagrams such as a constraint $n_k \leq \sigma$, where σ is a fixed number. Each slide is represented by a roll of a block to a lower column. With the constraint of $n_k \leq 1$, the number of partitions $q(A)$ of an integer A into integer summands without regards to order can be obtained through the generating function of Eq.(91) in the appendix. We hope to develop for this case in future. It is also useful

to picture a nuclear fragmentation distribution in terms of a block representation with the cluster size j the vertical column and with the number of clusters of that size, the $\tilde{n}_j$, given horizontally. The columns are arranged in descending order from left to right as shown in Fig.2d. This fragmentation diagram is the horizontally reflected version of a Young tableau or the 90° counter clockwise rotation of a Ferrer's diagram. The multiplicity is just the number of columns. With this fragmentation diagram, a fragmentation process can be represented as the stepwise descent of blocks from a higher column to a lower one [14]. In these block diagrams, the $p(A)$ simply counts all possible arrangements of A blocks. The $p(A, m)$ counts all possible arrangements with m fixed rows for a Ferrer's diagram or an avalanche diagram and with m fixed columns for a Young tableau or a fragmentation diagram.

3 Canonical Partition Functions for Fragmentation Phenomena and Cycle Indicators of S_n

Consider an ensemble of partitions or cycle classes of A objects into smaller clusters or cycles which are weighted by x_j according to the size j . The weighted sum of all the partition Π_A of A objects becomes the corresponding partition function. Let $W_A(\vec{n}, \vec{x})$ be a weight function for a particular decomposition or partition $\vec{n}$. The $\vec{x} = \{x_j\} = (x_1, x_2, \dots, x_A)$ and $\vec{n} = \{n_j\} = (n_1, n_2, \dots, n_A)$. The canonical partition function $Q_A(\vec{x})$ is given by

$$Q_A(\vec{x}) = \sum_{\Pi_A} W_A(\vec{n}, \vec{x}). \quad (9)$$

When $W_A(\vec{n}, \vec{x})$ is taken as

$$W_A(\vec{n}, \vec{x}) = M_2(A, \vec{n}) \prod_{j=1}^A x_j^{n_j}, \quad (10)$$

with $M_2(A, \vec{n})$ is given by Eq.(7), then $Q_A(\vec{x})$ is also the cycle indicator of the permutation of symmetric group S_A . In Refs.[14, 15], a generating function for the canonical

partition function were developed for fragmentation phenomena. Specifically, a generating function

$$Q(u, \vec{x}) = \exp\left(\sum_{j=1}^{\infty} x_j \frac{u^j}{j}\right) = \sum_{A=0}^{\infty} Q_A(\vec{x}) \frac{u^A}{A!} \quad (11)$$

was used to find the canonical partition function $Q_A(\vec{x})$ and the mean number of clusters of size k . A more general form for the grand canonical partition function $Q(u, \vec{x})$ is given and discussed in Ref.[16].

When all the x_j in $\vec{x}$ are equal, $Q(u, x) = Q(u, \{x_j = x\}) = (1 - u)^{-x}$ and

$$Q_A(x) = Q_A(\{x_j = x\}) = x(x+1) \cdots (x+A-1) = \frac{\Gamma(x+A)}{\Gamma(x)}. \quad (12)$$

The $\Gamma(x)$ is the gamma function. This canonical partition function $Q_A(x)$ has zeros at $x = 0, -1, -2, \dots, -(A-1)$ and the number increases with increasing A . When x is an integer, $x = n$, the $Q_A(n) = n(n+1) \cdots (n+A-1)$ can be written as a binomial or a negative binomial coefficient. In particular,

$$Q_A(n) = A! \binom{n+A-1}{A} = (-1)^A A! \binom{-n}{A}, \quad (13)$$

where the binomial coefficient $\binom{n}{A}$ and the negative binomial coefficient $\binom{-n}{A}$ are

$$\begin{aligned} \binom{n}{A} &= \frac{n(n-1) \cdots (n-A+1)}{A!} = \frac{\Gamma(n+1)}{\Gamma(n-A+1)\Gamma(A+1)}, \\ \binom{-n}{A} &= \frac{(-n)(-n-1) \cdots (-n-A+1)}{A!} = \frac{\Gamma(-n+1)}{\Gamma(-n-A+1)\Gamma(A+1)} \\ &= (-1)^A \binom{n+A-1}{A} = (-1)^A \frac{\Gamma(n+A)}{\Gamma(n)\Gamma(A+1)}. \end{aligned} \quad (14)$$

Here the binomial and the negative binomial are

$$\begin{aligned} (x_1 + x_2)^n &= \sum_{A=0}^n \binom{n}{A} x_1^A x_2^{n-A}, \\ (x_1 + x_2)^{-n} &= \sum_{A=0}^{\infty} \binom{-n}{A} x_1^A x_2^{-(n+A)}, \end{aligned} \quad (15)$$

respectively. At $x = 1$, $Q_A(1) = A!$.

The $Q_A(x)$ is useful in studying permutations and fragmentations by total number of cycles or multiplicity m with $m = \sum_j n_j = \lambda_1$. For example,

$$Q_5(x) = x(x+1)(x+2)(x+3)(x+4) = 24x^1 + 50x^2 + 35x^3 + 10x^4 + x^5$$

and the coefficient in front of x^m , $m = 1$ to 5 , are signless Stirling numbers $(-1)^{A-m} S(A, m)$ of the first kind [18]. In general,

$$Q_A(x) = \frac{\Gamma(x+A)}{\Gamma(x)} = \sum_{m=1}^A (-1)^{A-m} S(A, m) x^m. \quad (16)$$

The sum of the coefficients (signless Stirling numbers) in $Q_A(x)$ is $Q_A(1) = A!$ which is $5! = 120$ for $A = 5$. The signless Stirling numbers appear in the theory of the multiplicity distribution developed in Ref.[14]. When each cycle is tagged with a different x_j , as in $Q_A(\vec{x})$, then

$$Q_5(\vec{x}) = 24x_5 + 30x_4x_1 + 20x_3x_2 + 20x_3x_1^2 + 15x_2^2x_1 + 10x_2x_1^3 + x_1^5,$$

where the subscript i in x_i corresponds to the length of the cycle or the size of the fragment and the power of x_i is n_i . The coefficients of each term are the Cauchy's number M_2 given by Eq.(7) (see Table 1).

We next define a quantity $Q_A^m(\vec{x})$;

$$Q_A^m(\vec{x}) = \sum_{\Pi_{A,m}} W_A(\vec{n}, \vec{x}), \quad (17)$$

where the sum is over all partitions Π_A of A with m fixed. Also, a quantity $Q_A^{n_h}(\vec{x})$ is defined by

$$Q_A^{n_h}(\vec{x}) = \sum_{\Pi_{A,n_h}} W_A(\vec{n}, \vec{x}), \quad (18)$$

with the sum over all partitions Π_A of A with n_h specified. The $Q_A^m(\vec{x})$ and $Q_A^{n_h}(\vec{x})$ are more constrained than $Q_A(\vec{x})$ and

$$Q_A(\vec{x}) = \sum_{m=1}^A Q_A^m(\vec{x}) = \sum_{n_h=0}^{A/h} Q_A^{n_h}(\vec{x}). \quad (19)$$

For a system that has a total energy proportional to m , $Q_A^m(\vec{x})$ becomes a microcanonical partition function. Finally, define p -th order factorial [16] as

$$F_{A,p}^{n_k}(\vec{x}) = n_k(n_k - 1)(n_k - 2) \cdots (n_k - p + 1) Q_A^{n_k}(\vec{x}) = \left[\frac{x_k}{k} \right]^p \frac{A!}{(A - pk)!} Q_{A-pk}^{n_k-p}(\vec{x}). \quad (20)$$

These quantities will be used later.

Once the canonical partition function $Q_A(\vec{x})$ is determined, the distribution of fragments in the canonical ensemble follows using general results obtained in Ref.[15, 16]. Letting $\langle n_k \rangle$ be the mean number of clusters of size k ,

$$\langle n_k \rangle = Y_A(k, \vec{x}) = \frac{\sum_{n_k} F_{A,p=1}^{n_k}(\vec{x})}{Q_A(\vec{x})} = \left[\frac{x_k}{k} \right] \frac{A!}{(A - k)!} \frac{Q_{A-k}(\vec{x})}{Q_A(\vec{x})}. \quad (21)$$

The contribution of partitions with fixed multiplicity m to the mean cluster distribution $\langle n_k \rangle$ of size k is simply

$$\langle n_k \rangle_m = \left[\frac{x_k}{k} \right] \frac{A!}{(A - k)!} \frac{Q_{A-k}^{m-1}(\vec{x})}{Q_A^m(\vec{x})}. \quad (22)$$

The total yield of clusters of size k is then made from a sum of contributions from each multiplicity m

$$\langle n_k \rangle = \sum_{m=1}^A \langle n_k \rangle_m. \quad (23)$$

When the denominator $Q_A(\vec{x})$ in Eq.(22) is replaced by the microcanonical partition function $Q_A^m(\vec{x})$, the resulting mean cluster size distribution of Eq.(22) is that arising from a microcanonical ensemble with fixed multiplicity m and A [16].

Correlations and moments are also easily obtained from $Q_A(\vec{x})$. The correlation $\langle n_k n_j \rangle$ for $k \neq j$ is given by

$$\langle n_k n_j \rangle = \left[\frac{x_k}{k} \frac{x_j}{j} \right] \frac{A!}{(A - k - j)!} \frac{Q_{A-k-j}(\vec{x})}{Q_A(\vec{x})}, \quad (24)$$

while the second factorial moment is

$$\langle n_k(n_k - 1) \rangle = \frac{\sum_{n_k} F_{A,p=2}^{n_k}(\vec{x})}{Q_A(\vec{x})} = \left[\frac{x_k}{k} \right]^2 \frac{A!}{(A - 2k)!} \frac{Q_{A-2k}(\vec{x})}{Q_A(\vec{x})}. \quad (25)$$

In fact, defining $\bar{n}_k^p = n_k(n_k - 1) \cdots (n_k - p + 1)$, the factorial moment of order p is given by

$$\langle \bar{n}_k^p \rangle = \frac{\sum n_k F_{A,p}^{n_k}(\vec{x})}{Q_A(\vec{x})} = \left[\frac{x_k}{k} \right]^p \frac{A!}{(A - pk)!} \frac{Q_{A-pk}(\vec{x})}{Q_A(\vec{x})}. \quad (26)$$

Also, for $k \neq j$,

$$\langle \bar{n}_k^p \bar{n}_j^l \rangle = \left[\frac{x_k}{k} \right]^p \left[\frac{x_j}{j} \right]^l \frac{A!}{(A - pk - lj)!} \frac{Q_{A-pk-lj}(\vec{x})}{Q_A(\vec{x})}. \quad (27)$$

The above results are easily generalized to 3 and more sizes, i.e.,

$$\langle n_k n_j n_l \cdots \rangle = \left[\frac{x_k}{k} \frac{x_j}{j} \frac{x_l}{l} \cdots \right] \frac{A!}{(A - [k + j + l + \cdots])!} \frac{Q_{A-[k+j+l+\cdots]}(\vec{x})}{Q_A(\vec{x})}, \quad (28)$$

where $k, j, l, \dots$, are all different each other. From the factorial moment of Eq.(26), we can find the usual moment $\langle n_k^p \rangle$.

4 Applications to Various Models

The previous section summarizes general expressions for properties associated with cluster sizes arising from fragmentation phenomena. Once the partition function $Q_A(\vec{x})$ is evaluated, the expressions given in Section 3 can be used to obtain the desired quantities. This section gives some applications. Various models for the canonical ensemble partition function $Q_A(\vec{x})$ are discussed. The various models show a rich variety of different possibilities and behaviors.

4.1 Model Based on a Single Tuning Parameter x ; Length Scales and Scale Invariance

When all the x_j 's are equal to x , then the canonical partition function becomes $Q_A(x) = x(x+1) \cdots (x+A-1)$ as given by Eq.(12). Thus, using Eqs.(21) - (25),

$$\langle n_k \rangle = \left[\frac{x}{k} \right] \frac{A!}{(A-k)!} \frac{\Gamma(x+A-k)}{\Gamma(x+A)},$$

$$\begin{aligned}
\langle n_k(n_k - 1) \rangle &= \left[\frac{x}{k} \right]^2 \frac{A!}{(A - 2k)!} \frac{\Gamma(x + A - 2k)}{\Gamma(x + A)}, \\
\langle n_k n_j \rangle &= \left[\frac{x}{k} \right] \left[\frac{x}{j} \right] \frac{A!}{(A - k - j)!} \frac{\Gamma(x + A - k - j)}{\Gamma(x + A)}.
\end{aligned} \tag{29}$$

Properties of the distribution of cluster sizes were discussed in detail in Refs.[14, 15]. Moreover a thermodynamic model for x was developed [14, 16]. It was also recently discovered by one of us (A.Z.M.) that the x model also appears in population genetics. The population genetics model was developed by Ewens [24, 25]. A comparison between the Ewens model and the fragmentation model is developed in Ref.[26]. In population genetics, the cluster size k becomes the number of copies of a given gene, while the number of clusters of size k , the n_k , becomes the number of different types of genes each having k copies as discussed in Ref.[25]. The x is related to the mutation rate [24].

We note the following features of the x model. At $x = 0$, only one cluster of size A exists; which we called the fused mode. At $x = \infty$, A of monomers exist without any other size, i.e., total fragmentation (vaporization) mode. These two extreme cases have zero fluctuations. At $x = 1$,

$$\begin{aligned}
\langle n_k \rangle &= \frac{1}{k}, \\
\langle n_k n_j \rangle &= \left[\frac{1}{k} \right] \left[\frac{1}{j} \right] = \langle n_k \rangle \langle n_j \rangle, \\
\langle n_k(n_k - 1) \rangle &= \frac{1}{k^2} = \langle n_k \rangle^2,
\end{aligned} \tag{30}$$

for $k \leq A$, $k + j \leq A$, and $2k \leq A$ respectively. Otherwise, the quantities in Eq.(30) are zero. The last result of Eq.(30) shows that the fluctuation is the same as the mean number of clusters for $2k \leq A$, i.e.,

$$\langle n_k^2 \rangle - \langle n_k \rangle^2 = \langle n_k \rangle = \frac{1}{k}. \tag{31}$$

The results of Eqs.(30) and (31) are independent of the size A so long as the con-

straints are satisfied. When x is an integer $x = n$, we have

$$\langle n_k \rangle = \left[\frac{n}{k} \right] \frac{(A - k + 1)(A - k + 2) \cdots (A - k + n - 1)}{(A + 1)(A + 2) \cdots (A + n - 1)}. \quad (32)$$

This result shows that $\langle n_k \rangle$ depends on the size A except for $x = n = 1$ case.

Eq.(32) can also be rewritten as

$$\langle n_k \rangle = \left[\frac{n}{k} \right] \frac{\binom{n + A - k - 1}{A - k}}{\binom{n + A - 1}{A}} = (-1)^k \left[\frac{n}{k} \right] \frac{\binom{-n}{A - k}}{\binom{-n}{A}}, \quad (33)$$

and, thus, in terms of binomial or negative binomial coefficient given by Eq.(14). Here x ranges from zero to infinity. A simple transformation to a new variable $t = 1/(1+x)$ changes the domain of x to the interval from 0 to 1: $x = 0$ to $t = 1$, $x = \infty$ to $t = 0$, and $x = 1$ to $t = 1/2$.

As already noted, the cluster size distribution at $x = 1$ is $\langle n_k \rangle = 1/k$ and is thus scale invariant (i.e., independent of A) except for the trivial constraint $k < A$. A distribution having a length scale of length K is characterized by

$$\langle n_k \rangle \propto e^{-k/K}. \quad (34)$$

To obtain a scale invariant behavior such as $1/k$ from Eq.(34), a distribution of length scales must be presents as can be seen from

$$\frac{\Gamma(n+1)}{k^{n+1}} = \int_0^\infty \frac{dK}{K^{n+2}} e^{-k/K}. \quad (35)$$

Thus the hyperbolic behavior $1/k$ can be thought of as arising in a system in which all length scales K are present with a density $1/K^2$. This result is well known in $1/f$ noise where a similar discussion is given for the origin of this noise [11]. Moreover, since $\langle n_k^2 \rangle - \langle n_k \rangle^2 = \langle n_k \rangle = 1/k$, the fluctuation has all length scales present in $\langle n_k \rangle$ at $x = 1$. When $x \gg A$, the behavior of $\langle n_k \rangle$ for large k [14] is

$$\langle n_k \rangle \approx \frac{A}{k} e^{-(k-1)\ln(x/A)} \approx e^{-k\ln(x/A)}, \quad (36)$$

so that a length scale appears to be $K = 1/\ln(x/A)$. Scale invariance also appears in sandpile and avalanche theories [4, 5] and in other power law behaviors.

Before ending, we note that the scale invariant distribution $\langle n_k \rangle = 1/k$ can be rewritten as a $1/f$ spectrum. A given cycle of length k is cyclic with period k . The length 3 permutation $a = \begin{pmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \end{pmatrix}$ is such that $a^3 = e$, the identity element. Note a frequency can be associated with a period k such that $k f_k = 1$, and thus $f_k = 1/k$. In nuclear physics, the harmonic oscillator frequency f_k associated with a nucleus of size k (having k nucleons) is given by $\hbar f_k = \hbar \omega_k = a/k^{1/3}$ where $a \approx 40 \text{ MeV}$ for k large. Using the transformation of $n(f) = \langle n_k \rangle dk/df$, for both of the permutation cycle and nuclear harmonic oscillator, we obtain $n(f) \sim 1/f$ for the case of $\langle n_k \rangle = 1/k$.

4.2 Sum Rules, Cumulative Mass, and Moments and Correlations in Size

This section considers the gross features of the distribution $\langle n_k \rangle$ of cluster sizes by considering sum rules and moments of this distribution. The moments are obtained by multiplying each $\langle n_k \rangle$ by some power of k and summing over k . The zeroth and the first moments are the mean multiplicity and the total number of elements A respectively;

$$\begin{aligned} \sum_{k=1}^A \langle n_k \rangle &= \langle m \rangle, \\ \sum_{k=1}^A k \langle n_k \rangle &= A. \end{aligned} \tag{37}$$

The first moment result follows from the conservation constraint $\sum_k k n_k = A$. The results of Eq.(37) are true for arbitrary x_j 's and not just for the $x_j = x$ case.

Another quantity of interest is the cumulative mass $M(s)$,

$$M(s) = \sum_{k=1}^{[s]+1} k \langle n_k \rangle, \tag{38}$$

where the $[s]$ is the greatest integer in the continuous real value s , i.e., $[s] \leq s < [s] + 1$. Due to the sum rule Eq.(37), the cumulative mass becomes A for $s \geq A - 1$, i.e., $M(s \geq A - 1) = A$. For the $x_j = x = 1$ case, this cumulative mass $M(s)$ becomes a uniform staircase in which the step size is independent of the position k and the size A of the staircase. This result is shown in Fig.3a. For $x = n > 1$, the cumulative mass $M(s)$ is no longer an uniform staircase as can be seen from Eq.(32), i.e., the step size depends on k and A (see Fig.3a for $x = 3$). For x_j chosen such that $x_j = 0$ for some specific j 's and x for others, intermissions or breaks appear in $M(s)$ of Eq.(38). An extreme example in a continuum model is a Devil's staircase obtained by Cantor's rule [14].

Using Eq.(48) in the next subsection, we can obtain all higher order moments for the $x_j = x$ case. Specifically, the second, the third, and the fourth moments are

$$\begin{aligned} \sum_{k=1}^A k^2 < n_k > &= \frac{A(A+x)}{(x+1)}, \\ \sum_{k=1}^A k^3 < n_k > &= \frac{A(A+x)(2A+x)}{(x+1)(x+2)}, \\ \sum_{k=1}^A k^4 < n_k > &= \frac{A(A+x)[6A(A+x) + x(x-1)]}{(x+1)(x+2)(x+3)}. \end{aligned} \quad (39)$$

At $x = 1$ [14], the second moment is a statement that the sum of first A integers is $A(A+1)/2$, while the third moment is the result that the sum of the squares of the first A integers is $A(A+1)(2A+1)/6$ and the fourth is the sum of cubic integers, $A^2(A+1)^2/4$. Moments of a size distribution have been used in Ref.[27] to study the energy dependence of critical exponent of mass distributions in nuclear multifragmentation.

Using the notation $Y_A(k, \vec{x})$ of Eq.(21), Eq.(24) can be rewritten as

$$< n_k n_j > = Y_A(k, \vec{x}) Y_{A-k}(j, \vec{x}) = Y_{A-j}(k, \vec{x}) Y_A(j, \vec{x}).$$

The $\langle n_k n_j \rangle = 0$ for $k + j > A$. Then, we obtain the following results for correlations in cluster sizes:

$$\sum_{k=1}^A \sum_{j=1}^A k^m j^l \langle n_k n_j \rangle = \sum_{k=1}^A k^m Y_A(k, \vec{x}) \sum_{j=1}^{A-k} j^l Y_{A-k}(j, \vec{x}). \quad (40)$$

Specifically, using Eq.(37) for arbitrary x_j 's, we have

$$\begin{aligned} \sum_{k=1}^A \sum_{j=1}^A k j \langle n_k n_j \rangle &= \sum_{k=1}^A k Y_A(k, \vec{x}) \sum_{j=1}^{A-k} j Y_{A-k}(j, \vec{x}) \\ &= \sum_{k=1}^A \langle n_k \rangle k(A-k) = A^2 - \sum_{k=1}^A k^2 \langle n_k \rangle. \end{aligned} \quad (41)$$

When $x_i = x$ and using Eq.(39), the following results are obtained;

$$\begin{aligned} \sum_{k=1}^A \sum_{j=1}^A k j \langle n_k n_j \rangle &= \frac{x A(A-1)}{(x+1)}, \\ \sum_{k=1}^A \sum_{j=1}^A k^2 j^2 \langle n_k n_j \rangle &= \frac{x A(A+x) [A(A+x) - (x+1)]}{(x+1)(x+2)(x+3)}. \end{aligned} \quad (42)$$

Similar types of correlations have been used in Ref.[28] to study critical behavior in nuclear multifragmentation. For the case of $x = 1$, the results of Eq.(42) reduce to $A(A-1)/2$ and $(A+2)(A+1)A(A-1)/24$.

4.3 Fragmentation and Probability Theory

Since the sum rule for the first moment Eq.(37), which is the total number conservation Eq.(1), is independent of the parameter x_j 's, we can define a new probability distribution $P_A(k, \vec{x})$ as

$$P_A(k, \vec{x}) = \frac{k \langle n_k \rangle}{A}. \quad (43)$$

The $P_A(k, \vec{x})$ is the fraction of the total mass in clusters of size k and is thus the probability of a particle belonging to clusters of size k . The $P_A(k, \vec{x})$ also represents the ratio of the step height of a staircase Eq.(38) to the total height A (see Fig.3) at the k -th step. The $A P_A(k, \vec{x})$, in a continuum limit for k , corresponds to the slope

of the cumulative mass $M(s)$. The probability of Eq.(43) is developed and related with various probability theories in Ref.[29] for the $x_j = x$ case. With the probability distribution of Eq.(43), $\sum_k P_A(k, \vec{x}) = 1$ and the $(l+1)$ -th order moment of $\langle n_k \rangle$ becomes the average of l -th moment $\langle k^l \rangle$ of $P_A(k, \vec{x})$ times A :

$$\langle k^l \rangle = \sum_{k=1}^A k^l P_A(k, \vec{x}) = \frac{1}{A} \sum_{k=1}^A k^{l+1} \langle n_k \rangle. \quad (44)$$

Specifically, the second, third, and fourth moments of Eq.(39) are $A \langle k \rangle$, $A \langle k^2 \rangle$, and $A \langle k^3 \rangle$ respectively when $x_i = x$.

When all the x_j 's are x , the $P_A(k, \vec{x})$ is

$$P_A(k, x) = x \binom{A-1}{k-1} B(x+A-k, k). \quad (45)$$

Here, the $\binom{A-1}{k-1}$ is the binomial coefficient given by Eq.(14) and $B(x+A-k, k)$ is a beta function given by

$$B(a, b) = \frac{\Gamma(a)\Gamma(b)}{\Gamma(a+b)} = \int_0^1 dp p^{a-1} (1-p)^{b-1}. \quad (46)$$

Since

$$\begin{aligned} & k(k+1) \cdots (k+n-1) B(x+A-k, k) \\ &= (x+A)(x+A+1) \cdots (x+A+n-1) B(x+A-k, k+n), \end{aligned} \quad (47)$$

the factorial moments in $P_A(k, x)$ are given by

$$\begin{aligned} \langle k(k+1)(k+2) \cdots (k+n-1) \rangle &= \sum_{k=1}^A k(k+1)(k+2) \cdots (k+n-1) P_A(k, x) \\ &= n! \frac{\Gamma(x+1)}{\Gamma(x+1+n)} \frac{\Gamma(x+A+n)}{\Gamma(x+A)}. \end{aligned} \quad (48)$$

Here we have used the integral representation of the beta function. From Eq.(48) or (39), the average of $\langle k \rangle$ and its fluctuation Δk become

$$\langle k \rangle = \sum_{k=1}^A k P_A(k, x) = \frac{(A+x)}{(x+1)},$$

$$\begin{aligned} \langle k^2 \rangle &= \sum_{k=1}^A k^2 P_A(k, x) = \frac{(A+x)(2A+x)}{(x+1)(x+2)}, \\ \Delta k &= \langle k^2 \rangle - \langle k \rangle^2 = \frac{(A+x)(A-1)x}{(x+1)^2(x+2)}. \end{aligned} \quad (49)$$

The average of $\langle k \rangle$ is A for $x = 0$ and 1 for $x = \infty$ with zero fluctuation ($\Delta k = 0$) for both of these cases. When $x = 1$, the average of $\langle k \rangle$ becomes $(A+1)/2$ with its fluctuation $\Delta k = (A^2 - 1)/12$.

Using the integral representation of Eq.(46) for the beta function, the $P_A(k, x)$ can be rewritten as

$$P_A(k, x) = \int_0^1 dp \frac{(A-1)!}{(A-k)!(k-1)!} p^{k-1} (1-p)^{A-k} u(x, p), \quad (50)$$

where

$$u(x, p) = x(1-p)^{x-1}. \quad (51)$$

The $\int_0^1 u(x, p) dp = 1$. Thus this $u(x, p)$ can be considered as a density function weighting each point in p -space, that is, the probability p of a binomial distribution. This expression for $P_A(k, x)$ is a special case of a randomized Bernoulli distribution [30] where the probability of success p for a trial is treated as a random variable with a density $u(x, p)$. The number of successes out of $A-1$ trials becomes $s = k-1$. The density function $u(x, p)$ has the following simple properties: $u(x=1, p) = 1$ which is a uniform distribution giving $P_A(k, x=1) = 1/A$ from Eqs.(46) and (50); $\lim_{x \rightarrow 0} u(x, p) = \delta(1-p)$ which has probability of success equal to unity and $P_A(k, x=0) = \delta_{k,A}$; $\lim_{x \rightarrow \infty} u(x, p) = \delta(p-0)$ so that the probability of success is zero giving $P_A(k, x=\infty) = \delta_{k,1}$.

Using Eq.(43), we can also relate the fragment distribution $\langle n_k \rangle$ to the probability $P_A(k, \vec{x})$ of $k-1$ successes (heads) out of $A-1$ trials (throws) of a skewed coin ($u(\vec{x}, p)$). The $\langle n_k \rangle$ can also be related to the probability that a face with

k dots of a lopsided die with A faces touches the table. Here the size k of a cluster corresponds to the number of dots on a face. The total number of elements A is the number of faces of a die. The probability $P_A(k, \bar{x})$ that the i -th nucleon belongs to one of clusters of size k corresponds to the probability that the face with k dots touches the table at the i -th throw of the total A throws of a die with A faces. The extra factor $u(x, p)$ of Eq.(51) determines the skewness of a coin or the lopsidedness of a die. Inversely, if we know a probability $P_A(k, \bar{x})$, then the distribution of clusters is $\langle n_k \rangle = (A/k)P_A(k, \bar{x})$ through Eq.(43).

If $u(x, p)$ in Eq.(50) had been a delta function at p_o such as $u(x, p) = \delta(p - p_o(x))$, then

$$P_A(k, p_o) = \binom{A-1}{k-1} p_o^{k-1} (1 - p_o)^{A-k} \quad (52)$$

which is a binomial distribution with probability $p_o(x)$. This is the probability having $k-1$ heads in the $A-1$ tosses of a two sided coin which is skewed with probability p_o of head. With this probability, the mean cluster size $\langle k \rangle$ and its fluctuation are

$$\begin{aligned} \langle k \rangle &= \sum_{k=1}^A k P_A(k, p_o) = (A-1)p_o + 1, \\ \Delta k &= (A-1)p_o(1 - p_o). \end{aligned} \quad (53)$$

For this case, we have

$$\langle n_k \rangle = \frac{A!}{(A-k)!k!} [p_o(x)]^{k-1} [1 - p_o(x)]^{A-k}, \quad (54)$$

and thus a scale invariant distribution does not exist. The probability of Eq.(52) is also the same as the Rowlinson urn [31, 32] containing two colored balls, with a fraction p_o of one color and a fraction $1 - p_o$ of another color.

4.4 Models with Two Tuning Parameters; Situations without Monomers

In our next level of complexity, we take $x_k = xy^{\delta_{k,1}}$ that is $\vec{x} = (xy, x, x, \dots, x)$. Even though only a single term in the set of $\{x_i\}$ is treated differently, the expressions are much more complex than the previous case with one parameter. Here the monomer is put in a different footing because it may behave quite differently. For example, quarks and antiquarks are always bound into clusters of size two or more to form color singlets (hadrons). This situation excludes the existence of monomers and corresponds to setting $y = 0$. Here the objects are taken to be of one type and we hope to generalize the model to more complex situations in the future. A monomer is also quite different because of its lack of binding and internal excitations. For the symmetric group S_n , the case $y = 0$ corresponds to the subclass of permutations with no unit cycles. The exponential generating function of Eq.(11) for $Q_A(\vec{x})$ is now

$$Q(u, \vec{x}) = Q(u, x, y) = \exp \left(xyu + \frac{xu^2}{2} + \frac{xu^3}{3} + \dots \right) = \frac{1}{(1-u)^x} e^{x(y-1)u}, \quad (55)$$

where $\vec{x} = (xy, x, x, \dots)$. From $Q(u, \vec{x})$, one can obtain the canonical partition function $Q_A(x, y)$ for this model as

$$Q_A(x, y) = \sum_{r=0}^A \frac{A!}{r!(A-r)!} \frac{\Gamma(x+r)}{\Gamma(x)} [x(y-1)]^{A-r}. \quad (56)$$

When $y = 1$, $Q_A(x, y = 1) = Q_A(x) = x(x-1) \cdots (x+A-1)$. Having obtained $Q_A(x, y)$, $\langle n_k \rangle$ follows Eq.(21):

$$\langle n_k \rangle = \frac{xy^{\delta_{k,1}}}{k} \frac{A!}{(A-k)!} \frac{Q_{A-k}(x, y)}{Q_A(x, y)}.$$

Also the correlations and fluctuations follow from Eqs.(24) - (28). At $x = 1$ and $y = 1$, $\langle n_k \rangle = 1/k$, which is a power law behavior. Further details are studied in Ref.[15] with general y values; here we will study the $y = 0$ case in more detail.

As mentioned above, when $y = 0$, no monomer exists. We call $Q_A(x, y = 0) = D_A(x)$ which can be rewritten as

$$D_A(x) = \sum_{j=0}^A \binom{A}{j} (-x)^j Q_{A-j}(x) \quad (57)$$

from Eq.(56) and is generated by

$$Q(u, x, y = 0) = \frac{1}{(1-u)^x} e^{-xu} = \sum_{A=0}^{\infty} D_A(x) \frac{u^A}{A!}. \quad (58)$$

$Q_{A-j}(x)$ is given by Eq.(12) or (16) and $Q_0(x) = 1$. Expanding $D_A(x)$ in powers of x , we have

$$D_A(x) = \sum_{m=1}^A D(A, m) x^m, \quad (59)$$

$$D(A, m) = \sum_{j=0}^A \binom{A}{j} (-1)^{A-m+j} S(A-j, m-j),$$

where $S(A-j, m-j)$ is a Stirling number of the first kind and $D(A, m)$ is called associated Stirling number of the first kind. This associated number is obtained by summing the Cauchy number $M_2(A, \vec{n})$ of Eq.(7) over all partitions of A with fixed multiplicity m and with $n_1 = 0$. The $D_A(x)$ satisfy a recurrence relationship [17]

$$D_{A+1}(x) = AD_A(x) + Ax D_{A-1}(x), \quad (60)$$

with $D_0(x) = 1$ and $D_1(x) = 0$. For example, $D_2(x) = x$, $D_3(x) = 2x$, $D_4(x) = 3x^2 + 6x$, and $D_5(x) = 20x^2 + 24x$.

For $y = 0$, $\langle n_k \rangle = 0$ for $k = 1$ and $k = A-1$. For all other k ,

$$\langle n_k \rangle \approx \frac{x A}{k(A-k)}, \quad (61)$$

for $x \ll 1$ and $y = 0$. The mean number of clusters at $x = 1$ and $y = 0$ is

$$\langle n_k \rangle = \frac{1}{k} \frac{A!}{(A-k)!} \frac{D_{A-k}}{D_A} = \frac{1}{k} \frac{\sum_{j=0}^{A-k} (-1)^j / j!}{\sum_{j=0}^A (-1)^j / j!}, \quad (62)$$

except for $k = 1$ and $A - 1$ which do not exist at $y = 0$. From Eq.(56) or (57), the D_n is given by

$$D_n = D_n(x = 1) = n! \sum_{j=0}^n \frac{(-1)^j}{j!}, \quad (63)$$

and is called the number of derangements. From the result of Eq.(63), it is easy to show that D_n satisfy a recurrence relationship

$$D_n = nD_{n-1} + (-1)^n.$$

This relation is very similar to the recurrence relationship for factorials; the result differs from it because of the second term $(-1)^n$. The D_n appears in card shuffling and matching question [17] and counts the number of permutations in which no card is in its original position once the deck is shuffled. This problem is known as "*le problème de rencontres*". The $\mathcal{P}_A = D_A/A!$ is then the probability that no card will be in its original position after shuffling. Thus, the mean total mass in clusters of size k , $\langle n_k \rangle$, is the ratio of two such probabilities, consisting of a deck of cards of size $A - k$ and a deck of cards of size A (see Eq.(62)). When n in D_n is large, then

$$D_n \approx n! \frac{1}{e}. \quad (64)$$

Thus for small k and large A , $\mathcal{P}_A \approx 1/e$ and $\langle n_k \rangle \approx 1/k$ for $x = 1$ and $y = 0$. However, Eq.(62) is not a perfect power law in k for all k . Fig.3b shows the cumulative mass $M(s)$ for this case.

Through the relation of Eq.(43), we can relate a fragmentation process without monomers to a dice problem. At $y = 0$ and $x = 1$, the probability $P_A(k, \vec{x}) = \langle n_k \rangle / A$ is

$$P_A(k, x = 1, y = 0) = \frac{1}{A} \frac{\sum_{j=1}^{A-k} (-1)^j / j!}{\sum_{j=1}^A (-1)^j / j!} \quad (65)$$

except for $k = 1$ and $A - 1$ which do not exist, i.e., $P_A(k = 1, x = 1, y = 0) = 0$ and $P_A(k = A - 1, x = 1, y = 0) = 0$. Thus this special case of fragmentation of A objects without monomers is the same as rolling a die with $A - 2$ faces. On the die, there is no face with either $k = 1$ or $A - 1$ dots. Eq.(65) fixes the probability of each face of a lopsided die.

Before ending, we note that the generating function for cases when a certain fragment size (say l) is treated differently as $x_l = xy$ is

$$Q(u, \vec{x}) = \frac{1}{(1-u)^2} e^{(y-1)xy/l}. \quad (66)$$

Then the corresponding canonical partition function $Q_A(\vec{x})$ follows through Eq.(11). The case without monomers discussed above has $l = 1$ and $y = 0$.

4.5 Separation by Even and Odd Permutations; the Appearance of a Glitch at the End of a Uniform Staircase

This section also looks into the question of what happens to the distribution of clusters if we are limited to a subset of partitions which is a fraction of the complete set. The $y = 0$ or no monomer situation discussed in the previous section also considered a subset of all partitions. The subset that will be taken here are either the even or odd permutations. A permutation is even if it contains an even number of transpositions, and odd if it contains an odd number of transpositions. A cycle of even length is odd and one of odd length is even. The cycle indicators of even or odd permutations, Q_A^+ or Q_A^- respectively, are

$$Q_A^\pm(\vec{x}) = \frac{1}{2} \left[Q_A(x_1, x_2, x_3, x_4, \dots, x_A) \pm Q_A(x_1, -x_2, x_3, -x_4, \dots, (-1)^A x_A) \right]. \quad (67)$$

The parity, which is plus for even and minus for odd permutations, is determined by the number of even cycles; specifically, the parity is $(-1)^{n_2+n_4+n_6+\dots}$. Due to Eqs.(1) and (3), the parity of a given multiplicity m for a given A is $(-1)^{A+m}$. Thus, states

of the same multiplicity, even though they contain different classes, have the same parity (see Table 1).

The even permutations form a group [23], the alternating group A_n containing $n!/2$ elements ($n > 1$). In fact, the alternating group A_n is the only invariant subgroup of the symmetric group S_n for $n > 4$. The quotient group or factor group, S_n/A_n , has two one-dimensional representations, one being the identity or symmetric representation and the other the antisymmetric representation with characters $+1$ for even and -1 for odd permutations.

Models based on this distinction between even and odd cycles are easily solved. We consider the case $x_j = x$ for all j . Since $Q(u, \{x_j = (-1)^{j+1}x\}) = (1+u)^n$, we obtain

$$Q_A^\pm(x) = \frac{1}{2} [Q_A(x^+) \pm Q_A(x^-)]. \quad (68)$$

The $Q_A(x^+) = Q_A(x)$ of Eq.(12) which is the partition for $\vec{x} = \{x_j = x\}$ and

$$Q_A(x^-) = x(x-1)\cdots(x-A+1) = \sum_{m=0}^A S(A, m)x^m, \quad (69)$$

which is for $\vec{x} = \{x_j = (-1)^{j+1}x\}$. All properties of the cluster sizes are determined by the results of Section 3. However, due to different sets of x_j 's in $Q_A(x^+)$ and in $Q_A(x^-)$, we should use the proper x_k 's in the corresponding $Q_A(x^\pm)$ in Eqs.(20) - (28). Notice here that $Q_A(x^-)$ has zeros at $x = 0, 1, 2, \dots, A-1$ which are positive integers in contrast to $Q_A(x^+)$ of Eq.(12) which has zeros at negative integers. If we consider the $Q_A(x^-)$ piece only, then the mean cluster distribution $\langle n_k \rangle$ in the ensemble of $Q_A(x^-)$ becomes negative for some k depending on the value of x which is non-physical.

Here we look at the mean number $\langle n_k \rangle^\pm$ of clusters of size k in the ensemble

of even or odd permutations $Q_A^\pm(x)$.

$$\langle n_k \rangle^\pm = \frac{A!}{(A-k)!} \frac{\frac{1}{2} \left[(x/k) Q_{A-k}(x^+) \pm ((-1)^{k+1} x/k) Q_{A-k}(x^-) \right]}{Q_A^\pm(x)}. \quad (70)$$

For integer $x = n < A$, $Q_A(x^-) = 0$ and thus $Q_A^\pm(x) = Q_A(x)$ for $A \geq (n+1)$. Thus, when $x = n$, all properties for clusters of size $k < (A-n)$ are the same as the $Q_A(x)$ case considered in Section 4.1. For example, at $x = 1$, Eq.(70) becomes $\langle n_k \rangle^\pm = 1/k$ for $k = 1$ to $A-2$, while, for $k = A-1$ and $k = A$,

$$\begin{aligned} \langle n_{A-1} \rangle^\pm &= \frac{1 \pm (-1)^A}{A-1}, \\ \langle n_A \rangle^\pm &= \frac{1 \pm (-1)^{A+1}}{A}. \end{aligned} \quad (71)$$

Thus the distributions $\langle n_k \rangle^\pm$ fall exactly as a hyperbolic power law from $k = 1$ to $A-2$ and a glitch appears at either $k = A-1$ or $k = A$ (see Fig.3c). For even A , $\langle n_A \rangle^+ = 0$ and $\langle n_A \rangle^- = 2/A$, and for odd A , $\langle n_A \rangle^+ = 2/A$ and $\langle n_A \rangle^- = 0$. Also, the $k = A-1$ cluster is not present in even permutations ($\langle n_{A-1} \rangle^+ = 0$) for A odd and is not present in odd permutations ($\langle n_{A-1} \rangle^- = 0$) for even A . The results of Eq.(71) can be understood as follows. Only one cluster of size A can exist alone ($A = A$) and only one cluster of size $A-1$ can exist together with only one monomer ($A = (A-1) + 1$). For A even, a cycle of length A is odd; thus the cluster of size A exists only in odd permutation for even A . The cycle of length $A-1$ is even for A even and is odd for odd A . However the unit cycle is always even. Thus a partition into one cluster of size $A-1$ together with one monomer becomes even for even A and odd for odd A . The more remarkable feature is that a hyperbolic power law is maintained until the last two k 's, $k = A-1$ and $k = A$ (see Fig.3c). These results show that the power law is exactly maintained until the last two clusters even when only half of the partitions are used. The previous subsection also used a subset of partitions when $y = 0$ (all those with $n_1 = 0$) and the distribution is approximately

a power law only for small k values (see Fig.3b).

For this model, the probability defined by Eq.(43) for even and odd permutations $Q_A^\pm(x=1)$ becomes $P_A^\pm(k, x=1) = 1/A$ for $k = 1$ to $A-2$. For $k = A-1$ and A , either $P_A^\pm(k = A-1, x=1)$ or $P_A^\pm(k = A, x=1)$ is zero and the other one is $2/A$. In other word,

$$P_A^\pm(k = A-1, x=1) + P_A^\pm(k = A, x=1) = 2/A.$$

Here the probability for the size of $k = A-1$ or A whichever is even (odd) is zero for even (odd) permutations and the other one is nonzero. This situation is the same as a die with $A-1$ faces. The number of dots on the $A-2$ faces are 1 to $A-2$ which are equally weighted with $1/A$. The last face has $A-1$ or A dots whichever is odd (even) for even (odd) permutations. This face is weighted by $2/A$. We also can relate this to a die with A faces, with all faces equally weighted by $1/A$. In this case, the number of dots on the first $A-2$ faces are $k = 1$ to $A-2$. The last two faces have the same number of dots either $k = A-1$ or A which depends on whether A is even or odd and whether even or odd permutations are being considered. This is not a lopsided die, i.e., all the faces are weighted equally. But the numbers of dots on the faces are distributed differently from the usual die.

The cumulative mass (c.f. Eq.(38)), defined by

$$M^\pm(s) = \sum_{j=1}^{[s]+1} j < n_j >^\pm, \quad (72)$$

is shown for $A = 10$ at $x = 1$ in Fig.3c. We note that $M^\pm(s)$ is a uniform staircase function until the last two steps are reached. Then, for A even (Fig.3c), an intermission appears as a double step at $s = A-2$ with a missing rise at $s = A-1$ (i.e., missing A -th step) in $M^+(s)$ and, in $M^-(s)$, as a missing rise at $s = A-2$ with a double step at $s = A-1$. The same situation happens for the A odd case but with $M^+(s)$ and $M^-(s)$ interchanged.

4.6 Cayley's Theorem and Gauss Polynomials; the transformation to a state of dimers only

We next consider a very particular model in which all the x_k 's are different, but the choice for each x_k is generated in a particular way. Namely, we take

$$x_k = \frac{1}{1 - q^k}, \quad (73)$$

where q is a parameter whose range is $-1 < q < 1$ to keep x_k and n_k positive. For this particular choice, the partition function $Q_A(\vec{x})$ is again simple and is obtained by using a theorem due to Cayley [33]. Cayley's theorem for a decomposition is the following:

$$\frac{1}{(1 - q)(1 - q^2) \cdots (1 - q^A)} = \sum_{\vec{n}_A} \frac{M_2(A, \vec{n})/A!}{(1 - q)^{n_1}(1 - q^2)^{n_2} \cdots (1 - q^A)^{n_A}}, \quad (74)$$

with $M_2(A, \vec{n})$ the Cauchy's number of Eq.(7). The sum in Eq.(74) is over all partitions of A . Using this result, the partition function for the model is

$$Q_A(\vec{x}) = A! \prod_{k=1}^A x_k = \frac{A!}{(1 - q)(1 - q^2) \cdots (1 - q^A)}. \quad (75)$$

The ensemble averaged n_k 's can easily be obtained using the results of Section 3 and they are

$$\langle n_k \rangle = \left[\frac{x_k}{k} \right] \frac{x_1 x_2 \cdots x_{A-k}}{x_1 x_2 \cdots x_A} = \frac{x_k}{k \prod_{j=1}^k x_{A-j+1}}. \quad (76)$$

This result can be rewritten in terms of Gauss polynomials [33]. Defining

$$[q]_k = (1 - q)(1 - q^2) \cdots (1 - q^k), \quad (77)$$

then

$$\langle n_k \rangle = \frac{[q]_{k-1}}{k} \left[\frac{A}{k} \right], \quad (78)$$

where $\begin{bmatrix} A \\ k \end{bmatrix}$ is a Gauss polynomial defined as $[q]_A / ([q]_k [q]_{A-k})$.

Properties of the distribution of $\langle n_k \rangle$ are as follows. At $q = 0$, which gives $x_k = 1$ for all k as in Subsection 4.1, $\langle n_k \rangle = 1/k$ and a scale invariant power law emerges. As $q \rightarrow 1$, $x_k = k^{-1} \lim_{q \rightarrow 1} (1 - q)^{-1}$ by Eq.(73) and

$$\langle n_1 \rangle = A. \quad (79)$$

All other n_k 's are zero, so that only monomers are present. The same results can be obtained at $x \rightarrow \infty$ from Subsection 4.1 also. The model has an interesting solution at $q \rightarrow -1$. Eq.(73) gives $x_k = 1/2$ for odd k and $x_k = k^{-1} \lim_{q \rightarrow -1} (1 + q)^{-1}$ for even k . For A even,

$$\langle n_2 \rangle = \frac{A}{2} \quad (80)$$

and all others are zero. For A odd,

$$\begin{aligned} \langle n_1 \rangle &= 1, \\ \langle n_2 \rangle &= \frac{A-1}{2}, \end{aligned} \quad (81)$$

and all others are zero. Thus only clusters of size 2 (dimers) exist for even A . For odd A , only one monomer is present and all other clusters are dimers. The result of Eqs.(80) and (81) is an example of pairing phenomena and is an interesting solution for studying social behavior! A peak appears at the group size of two in a closed Vervet Monkey Troop [34]. Fig.4 illustrates the social behavior of such a troop and some attempts to fit the distribution of group sizes. A negative binomial [34], the x - y model of Section 4.4, and the pairing model of this section are compared with the data. Pairing phenomena appear in various areas of physics.

4.7 Transitions to Modes Made of Clusters with Even Number of Elements only

In the last section, x_k was taken as $1/(1 - q^k)$. Here, we take $x_k = 1 + q^k$ with $-1 \leq q < \infty$ and generate another solution to illustrate the many possibilities which exist. For this choice of x_k , the generating function of Eq.(11) is $Q(u, \vec{x}) = [1/(1-u) - q/(1-qu)]/(1-q)$. The partition function $Q_A(\vec{x})$, with $\vec{x} = (1+q, 1+q^2, \dots)$, can then be shown to be

$$Q_A(\vec{x}) = A! \frac{(1 - q^{A+1})}{(1 - q)}, \quad (82)$$

from a result in Ref.[17]. This gives

$$\langle n_k \rangle = \left[\frac{1 + q^k}{k} \right] \left[\frac{1 - q^{A-k+1}}{1 - q^{A+1}} \right]. \quad (83)$$

At $q = 0$, $x_k = 1$ for all k and $\langle n_k \rangle = 1/k$ as in Subsection 4.1. As $q \rightarrow \infty$, $x_k = \lim_{q \rightarrow \infty} q^k$ and thus $\langle n_k \rangle = 1/k$. This large q limit corresponds to the case in which each constituent is weighted the same, independently of the cluster to which it belongs. Thus the weight $x_k = q^k$ for the cluster of size k has no effect on the cluster distribution and becomes effectively the same as the $x = 1$ case. This behavior has also been shown in Ref.[16].

At $q = -1$, $\vec{x} = (0, 2, 0, 2, 0, \dots)$, so that the x_k for odd k 's are zero and for even k 's are 2. For even A ,

$$\langle n_k \rangle = \frac{2}{k}, \quad (84)$$

for even k and $\langle n_k \rangle = 0$ for odd k . This case shows a scale invariance but with missing clusters of odd sizes. The cumulative mass $M(s)$ shows this situation well with missing rises at each position of even integer s , i.e., missing steps at each odd number of step position (see Fig.3d). For this case, the probability of Eq.(43) becomes

$P_A(k, q = -1) = 2/A$ for even k and zero for odd k . Thus this specific model can describe a die with $A/2$ faces with even number of dots on each faces of $k = 2$ to A . All the $A/2$ faces are weighted equally by $2/A$.

For odd A , as $q \rightarrow -1$,

$$\langle n_k \rangle = \left[\frac{2}{k} \right] \left[\frac{A - k + 1}{A + 1} \right], \quad (85)$$

for even k , and

$$\langle n_k \rangle = \frac{2}{A + 1} \quad (86)$$

for k odd. This case corresponds to a lopsided die and we lose the scale invariance. The cumulative mass $M(s)$ shows the position s dependences (see Fig.3d).

5 Conclusion

We have shown that simple models of fragmentation can be developed which show a rich spectrum of different types of behavior. The models considered here are based on a correspondence between clusters and cycles of the permutation group. Methods from combinational analysis are used to obtain analytic expressions for canonical ensembles. These canonical ensembles are then used to obtain exactly soluble models of fragmentation phenomena.

A wide range of fragmentation behavior can be obtained from these models depending on the choice of tuning parameters. In some cases, exact scale invariant hyperbolic power laws emerge in the distribution of cluster sizes. Situations in which no monomers exist and in which only pairs exist are also studied. One model developed, which is based on a theorem due to Cayley, has the interesting property of having only dimers or pairs present at a particular point. Other situations are

discussed. Various properties of the distribution of cluster sizes are given. Simple expressions are found for moments, correlations, and fluctuations of the distribution of cluster sizes. Fragmentation phenomena are also related to simple probability pictures based on coin and dice models.

The thermodynamical behavior of a nuclear system has been studied using our simple model in Ref.[16]. This model will be compared to the cluster distribution of nuclear fragmentation phenomena in the near future. The model we have considered is based on one type of element only. Extending our model to the more than one type of element, such as neutrons and protons in a nuclear system (rather than just nucleons) or quarks and antiquarks with colors and flavors, also are interesting problems which we hope to develop in the future.

This work was supported in part by National Science Foundation grant number 89-03457 and by a Rutgers University Supercomputer Fellowship and by the U.S. Department of Energy.

Appendix; Generating Functions for Partitions

The generating function for $p(A)$ [18] is

$$P(u) = \frac{1}{(1-u)(1-u^2)(1-u^3)\dots} = \prod_{i=1}^{\infty} \left[\frac{1}{1-u^i} \right] = \sum_{A=0}^{\infty} p(A)u^A. \quad (87)$$

Since $-\ln(1-x) = \sum_{n=1}^{\infty} x^n/n$ and $\sum_{n=1}^{\infty} x^n = x/(1-x)$, the $P(u)$ can be written as $e^{f(u)}$ where

$$f(u) = \sum_{n=1}^{\infty} \frac{u^n}{n(1-u^n)}, \quad (88)$$

for $u^2 < 1$. Using $\sum_{n=1}^{\infty} 1/n = 0.57721 + \lim_{n \rightarrow \infty} \ln n$ and $\sum_{n=1}^{\infty} 1/n^2 = \pi^2/6$, it is easy to show that, near $u = 1$, $f(u \approx 1) \approx (\pi^2/6)(1-u)^{-1} - (1/2)\lim_{n \rightarrow \infty} \ln n + \beta$,

with β a constant of the order of unity. Steepest descent contour integration, i.e., $P(u_0) = e^{f(u_0)} \approx p(A)u_0^A$ where u_0 is such that $d[f(u) - A \ln u]/du = 0$ at $u = u_0$, leads to Eq.(4) up to an unimportant numerical factor.

The enumerating function for $p(A, m)$ is a two variable function [35]

$$P(u, t) = \frac{1}{(1-tu)(1-tu^2)(1-tu^3)\dots} = \prod_{i=1}^{\infty} \left[\frac{1}{1-tu^i} \right] = \sum_{m=0}^{\infty} P(u, m)t^m, \quad (89)$$

with $P(u, m)$ a generating function for partitions with m -parts which is

$$P(u, m) = \sum_{A=0}^{\infty} p(A, m)u^A = \frac{u^m}{(1-u)(1-u^2)\dots(1-u^m)}. \quad (90)$$

Again the steepest descent method leads to Eq.(5).

Other quantities of interest in partition enumeration are the following. The division of A into factors such that no n_i is greater than 1 is defined by a symbol $q(A)$. The $q(A)$ is generated by [18]

$$Q(u) = (1+u)(1+u^2)(1+u^3)\dots = \prod_{i=1}^{\infty} (1+u^i) = \sum_{A=0}^{\infty} q(A)u^A. \quad (91)$$

If different orderings are allowed, as in $5 = 4+1 = 1+4 = \dots$, the number of partitions $r(A)$ is obtained from a generating function $R(u)$ given by [16]

$$R(u) = \frac{1-u}{1-2u} = \sum_{A=0}^{\infty} r(A)u^A. \quad (92)$$

The $p(A)$, $q(A)$, and $r(A)$ all have polynomial generating functions. By contrast, the cycle indicators have exponential generating function as shown by Eq.(11) in Section 3. A more general generating function than the one considered in Section 3 is given in Ref.[16].

References

- [1] J.E. Finn *et al.*, Phys. Rev. Lett. **49**, 1321 (1982).
- [2] G.S. Hawkins, Annu. Rev. Astron. Astrophys. **2**, 149 (1964).
- [3] B. Gutenberg and C.F. Richter, Ann. Geophys. **9**, 1 (1956).
- [4] L. Kadanoff, S. Nagel, L. Wu, and S. Zhou, Phys. Rev. **A39**, 6524 (1989).
- [5] P. Bak, C. Tang, and K. Wiesenfeld, Phys. Rev. Lett. **59**, 381 (1987);
Phys. Rev. **A38**, 364 (1988).
- [6] W.K. Hartmann, Scientific America, January, 84 (1977).
- [7] D. Stauffer, **Introduction to Percolation Theory**, (Tayler & Francis, Philadelphia, 1985).
- [8] M.E. Fisher, Physics (N.Y.) **3**, 255 (1967).
- [9] G. Zipf, **Human Behavior and the Principle of Least Effort**, (Hafner, New York, 1972).
- [10] V. Pareto, **Cours d'Economie Politique**, (Lausanne, 1897).
- [11] W.H. Press, Comm. Mod. Phys. **C7**, 103 (1978);
P. Dutta and P.M. Horn, Rev. Mod. Phys. **53**, 497 (1981).
- [12] B.B. Mandelbrot, **The Fractal Geometry of Nature**, (Freeman, San Francisco, 1982).
- [13] E.W. Montroll and B.J. West, in **Fluctuation Phenomena**, edited by E.W. Montroll and J.L. Lebowitz (North-Holland, Amsterdam, 1976), 61.

- [14] A.Z. Mekjian, *Phys. Rev. C* **41**, 2103 (1990);
Phys. Rev. Letts. **64**, 2125 (1990).
- [15] S.J. Lee and A.Z. Mekjian, *Phys. Lett. A* **149**, 7 (1990).
- [16] S.J. Lee and A.Z. Mekjian, to be published.
- [17] J. Riordan, *An Introduction to Combinational Analysis*, (Wiley, New York, 1958).
- [18] M. Abramowitz and I. Stegun, *Handbook of Mathematical Functions*, Natl. Bur. Stand., Applied Math Series - 55 (U.S. GPO, Washington, DC, 1965).
- [19] A. Bohr and B.R. Mottelson, *Nuclear Structure, Vol. I*, (Benjamin, New York, 1969).
- [20] J.J. Griffin, *Phys. Rev. Lett.* **17**, 478 (1966).
- [21] D. Mitchell and N. Turok, *Phys. Rev. Lett.* **58**, 1577 (1987).
- [22] R. Hagedorn, in *Cargese Lectures in Physics, Vol.6*, ed. E. Schatzman, (Gordon and Breach, New York, 1973), 643.
- [23] M. Hamermesh, *Group Theory and its Application to Physical Problems*, (Addison-Weseley Pub., Reading, 1962).
- [24] W.J. Ewens, *Theoretical Population Biol.*, **3**, 87 (1972).
- [25] W.J. Ewens, *Mathematical Population Genetics*, Springer-Verlag, Berlin (1979).
- [26] A.Z. Mekjian, University of Washington, Institute of Nuclear Theory, preprint.
- [27] K.K. Gudima and Y.A. Murin, *Phys. Lett.* **234B**, 1 (1990).

- [28] H.R. Jaqaman, G. Papp, and D.H.E. Gross, Hahn-Meitner-Institut preprint (1990).
- [29] A.Z. Mekjian, Rutgers University preprint RU-90-07, submitted for publication (1990).
- [30] W. Feller, **An Introduction to Probability Theory and Its Applications**, Volumes 1 and 2, (John Wiley and Sons, Inc., New York, 1957).
- [31] J.S. Rowlinson, *Nature*, **225**, 1196 (1970).
- [32] E.T. Jaynes, in **The Maximum Entropy Formalism**, D. Levine and M. Tribus, editors (MIT Press, Cambridge, MA, 1979), 15.
- [33] G.E. Andrews, **Theory of Partitions**, (Addison-Wesley Publishing Com., Reading, Mass., 1976).
- [34] J.E. Cohen, *Theor. Population Biology* **3**, 119 (1972).
- [35] H. Gupta, **Tables of Partitions**, Royal Society Mathematical Tables, Vol. 4, (Cambridge University press, Cambridge, 1958).

Table 1: Partitions (Π_A) of $A = 5$ and 6. The parity and the Cauchy numbers (M_2) are also given.

A	$p(A)$	m	$p(A, m)$	$\Pi_A(\vec{n})$	$\Pi_A(\vec{\lambda})$	Parity	M_2
5	7	1	1	5	1^5	+	24
			2	1,4	$2,1^3$	-	30
				2,3	$2^2,1$	-	20
		3	2	$1^2,3$	$3,1^2$	+	20
				$1,2^2$	3,2	+	15
			4	$1^3,2$	4,1	-	10
		5	1	1^5	5	+	1
6	11	1	1	6	1^6	-	120
			3	1,5	$2,1^4$	+	144
				2,4	$2^2,1^2$	+	90
				3^2	2^3	+	40
		3	3	$1^2,4$	$3,1^3$	-	90
				$1,2,3$	$3,2,1$	-	120
				2^3	3^2	-	15
		4	2	$1^3,3$	$4,1^2$	+	40
				$1^2,2^2$	4,2	+	45
		5	1	$1^4,2$	5,1	-	15
		6	1	1^6	6	+	1

Figure Captions

Fig.1: Partitions of the integer 5, cycles of permutation group S_5 , and the fragmentations of an object with 5 constituents. The filled circles represent the constituent elements. The large circle indicates the cyclic permutation of elements in the cycle, and the constituents of the cluster are dots inside the circle in the fragmentation scheme.

Fig.2: Block diagrams for a partition Π_A of $A = 50$ into $\vec{n} = (1^4, 4^3, 6^1, 7^2, 9^2)$ which is $\vec{\lambda} = (11, 7^3, 5^2, 4, 2^2)$.

Fig.3: Staircases $M(s)$ (solid and broken lines), cluster distributions $A < n_k >$ (circles and squares), and probabilities $P_A(k, \vec{x})$ (cross and plus) for various models with $A = 10$: (a) One variable x model of Section 4.1 with $x = 1$ (solid line, circle, and cross) and with $x = 3$ (broken line, square, plus); (b) two variable x - y model of Section 4.4 with $x = 1$ and $y = 0$; (c) even (solid line, circle, cross) and odd (broken line, square, plus) permutations of Section 4.5 with $x = 1$; (d) the case of $x_k = 1/(1 - q^k)$ of Section 4.7 with $q = -1$ for $A = 10$ (solid line, circle, cross) and for $A = 9$ (broken line, square, plus).

Fig.4: Group size distribution of a Closed Vervet Monkey Troop. Filled circles are for the observed values [34] and dotted curves are from Ref.[34]. The solid line is the model of Section 4.6 with $q = 0.85$ and the dash and dash-dotted lines are the x - y model of Section 4.4 with $x = 2.0$ and $y = 0.4$ or with $x = 2.5$ and $y = 0.3$ respectively. Here $A = 17$ and the frequencies $\mathcal{N}n_k$ are normalized to $\mathcal{N} \sum_k k n_k = 599$ which is the same as the data.

Fig.1

Partitions of Integer	Cycles of Permutation	Fragmentations of Object
5
4+1
3+2
3+1+1
2+2+1
2+1+1+1
1+1+1+1+1

Fig.2

(a). Ferrer's Diagram :

$$\vec{n} = (1^4, 4^2, 6^1, 7^2, 9^2)$$

(b). Young's Tableau :

$$\vec{\lambda} = (11, 7^3, 5^2, 4, 2^2)$$

(c). Avalanche :

(d). Fragmentation :

Fig.3

Fig.4

END

**DATE
FILMED**
2 / 03 / 92