MASTER

**Pyrochemical and Dry
Processing Methods Program**

A Selected Bibliography

H. F. McDuffie
Duane H. Smith
Park T. Owen

OAK RIDGE NATIONAL LABORATORY
OPERATED BY UNION CARBIDE CORPORATION • FOR THE DEPARTMENT OF ENERGY

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency Thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

Printed in the United States of America. Available from
National Technical Information Service
U.S. Department of Commerce
5285 Port Royal Road, Springfield, Virginia 22161
Price: Printed Copy \$19.00; Microfiche \$3.00

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor any of its employees, nor any of its contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.

Contract No. W-7405-eng-26

Chemistry Division

PYROCHEMICAL AND DRY PROCESSING METHODS PROGRAM
A SELECTED BIBLIOGRAPHY

H. F. McDuffie
Duane H. Smith
Chemistry Division

Park T. Owen
Information Division

March 1979

NOTICE
This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Department of Energy, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.

Prepared by the
OAK RIDGE NATIONAL LABORATORY
Oak Ridge, Tennessee 37830
operated by
UNION CARBIDE CORPORATION
for the
DEPARTMENT OF ENERGY

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

THIS PAGE
WAS INTENTIONALLY
LEFT BLANK

CONTENTS

	<u>Page</u>
ABSTRACT	v
INTRODUCTION	v
SAMPLE REFERENCE	vi
ACKNOWLEDGMENTS	xi
BIBLIOGRAPHIC REFERENCES	1
INDEXES	
Author	407
Subject Descriptors	441
EDB Subject Category	505
Title	509

THIS PAGE
WAS INTENTIONALLY
LEFT BLANK

ABSTRACT

This selected bibliography was compiled to provide information support to the Pyrochemical and Dry Processing Methods (PDPM) Program sponsored by the Department of Energy and administered by the Argonne National Laboratory through work packages in many locations in the United States. Objectives of the PDPM Program are to evaluate nonaqueous methods of reprocessing spent fuel as a route to the development of proliferation-resistant and diversion-resistant methods for widespread use in the nuclear industry. To limit this bibliography to a manageable size for evaluation by the PDPM Program, emphasis was placed on the literature indexed in the ERDA-DOE Energy Data Base (EDB). If the bibliography is acceptable, similar references from Nuclear Science Abstracts (NSA) and Chemical Abstracts (CA) will be combined to form a PDPM Program computerized data base for further program use. The bibliography includes indexes to authors, subject descriptors, EDB subject categories, and titles.

INTRODUCTION

The ORNL Information Division has been building computerized data bases in the nuclear field for many years, with intensified activity since 1971. The proximity to the DOE Technical Information Center and the availability of strong computer support in the Computer Sciences Division led to the establishment of the RECON national network with its base in Oak Ridge. Many of the DOE's established information analysis centers are located in Oak Ridge where they mutually support one another.

When the ORNL work package of the PDPM Program was activated in February 1978, we discussed whether we might undertake to develop an information support function for the program as a whole, in addition to the literature analysis to which we were committed. This selected bibliography is an example, in hard copy, of what might be assembled and made available to program people, either through computer access on RECON, by a continuing series of hard copy documents, by individualized searches and selective dissemination of information (SDI) profiles, or, in cooperation with the information center, the preparation of critical reviews by experts to supplement specialized bibliographies.

We quickly found that our literature capture net when applied to Nuclear Science Abstracts, Chemical Abstracts, and the Energy Data Base was bringing in many more citations than we could conveniently publish in one hard copy specimen bibliography. We decided to restrict the sample to EDB for two reasons: one, the EDB literature is the most recent (nuclear literature from July 1976 when NSA was discontinued) and hence likely to be less familiar to the experienced program people than is the

older literature; and, two, we would intend to model our PDPM data base on EDB with respect to field identification, subject categories, subject descriptors, etc., and therefore would like our program members to know what type of information might become available in a larger base. (We would expect to convert NSA and CA into the EDB format.) We have already ascertained that the PDPM base can, indeed, be "brought up" on RECON where it will be available for individualized searching and printout in work-package locations that have hard-wired RECON lines or that have telephone dial-up capability for terminal access.

From review of this selected bibliography by PDPM people, we hope to learn whether and how it is useful and how it might be improved with respect to both content and format. The data-base building program we are using is sufficiently flexible to permit incorporation of almost any useful suggestions that arise and to change as the program needs or interests change. Particular deficiencies may be apparent in the materials and fabrication areas, since the programs in those areas are less well known to us at this early stage. If the program response is favorable, we should expect to obtain relevant references that other work-package investigators have already gathered and to add them to our mechanically captured set. Other information support activities would include systematic additions from the important earlier literature referenced via sources such as *Reactor Handbook. Vol. II. Fuel Reprocessing*, 2nd ed., edited by S. M. Stoller and R. B. Richards, 1961, and *Nuclear Metallurgy. Vol. 15. Symposium on Reprocessing of Nuclear Fuels* (CONF-690801), August 1969.

SAMPLE REFERENCE

The following is a list of all the field labels and their three-letter mnemonics being used for the numerical tags in the EDB [ERDA *Energy Information Data Base: Magnetic Tape Description* (TID-4581-R3)].

ABS	950	Abstract (when available)
AFA	170	Affiliation (Author of an Analytic)
AFM	190	Affiliation (Author of a Monograph)
ASI	300	Assignee
AUA	060	Personal Author of an Analytic
AUG	620	Title Augmentation
AUM	070	Personal Author of a Monograph
BIB	550	Source of Bibliographic Information Code
CAC	700	Corporate Code (INIS)
CAN	710	Corporate Author (INIS)
COD	250	CODEN
CFD	470	Conference Date
CFP	460	Conference Place
CFT	450	Conference Title
CNO	240	Contract Number

CPC	570	Country of Publication Code
CSC	500	COSATI ID Code
CTY	310	Place (City) of Publication
DAV	430	Availability, Price
DBM	XXX	Original Data Base Mnemonic
DBN	XXX	Original Data Base Name
DPN	440	Drop Notes
DSN	001	Original Document File Number
GEN	801	Subject Descriptors, General
LIT	040	Literary Indicator
LWC	560	Country of Affiliation Code
NCC	610	INIS Category Number
NSC	590	INIS Temporary Record Number
OTM	200	Original Title for a Monograph
PAD	380	Patent Filed Date
PCL	230	International Classification Code
PDD	370	Date
PDI	350	Journal Issue
PDL	420	Language
PDP	360	Journal Pages
PDT	260	Abbreviated Journal Title
PDV	340	Journal Volume
PGM	390	Pages of Books, Reports, or Analytics
PNO	220	Patent Number
PTC	020	Type of Item
PUB	320	Name of Publisher
RNA	530	File(s) Selected for Publication
RNO	150	Primary Report Number
RPO	520	Report Origin
SCC	540	Subject Category Codes
SJA	750	Abstract Number (Secondary)
SJI	011	Issue Number (Secondary)
SPL	802	Subject Descriptors, Splits
SRN	210	Secondary Report Number
THS	490	Thesis Statement
TIA	090	Primary Title of an Analytic
TLM	110	Primary Title of a Monograph
TLS	130	Primary Title of a Series
TRN	410	Translation Note

A complete journal citation from our base, printed with the three-letter labels appropriate to this type of publication, is as follows:

<DSN >76:036237
 <SEC >
 <SJI >32
 <SJA >NSA-33:30133;ERA-01:013612;EDB-76:036237
 <TTL >
 <TLM >DIFFUSION COEFFICIENTS IN MOLTEN METALS
 <ATI >
 <AUM >KONONENKO, V.I.; YATSENKO, S.P.
 <KEY >
 <GEN >ACTIVATION ENERGY;ANTIMONY;BISMUTH;CALCULATION
 METHODS;GALLIUM;INDIUM: T2;INDIUM ALLOYS: T3;LEAD;LIQUID
 METALS;SELF-DIFFUSION: Q1,Q2,Q3,Q4;SILVER;THALLIUM;TIN: T1;TIN ALLOYS:
 T4;USSR;VISCOSITY
 <SJT >
 <DBN >ERDA ENERGY DATA BASE
 <DBM >EEDB
 <PDD >1971
 <JNL >
 <PDT >TR. INST. KHIM., AKAD. NAUK SSSR, URAL. FIL.
 <CDN >TKAUA
 <PDV >23
 <PDP >62-67
 <PTC >TJ
 <ABS >THE SELF-DIFFUSION COEFFICIENTS OF INDIUM AND TIN WERE
 DETERMINED EXPERIMENTALLY IN NOT ONLY THE PURE METALS BUT A WIDE RANGE
 OF BINARY LIQUID INDIUM-TIN ALLOYS OVER THE TEMPERATURE RANGE 800 TO
 1100\$UP 0\$K. ACTIVATION ENERGIES AND PRE-EXPONENTIAL FACTORS WERE
 CALCULATED FOR EACH METAL. A BREAK IN THE ARRHENIUS PLOT WAS FOUND.
 COEFFICIENTS OF DIFFUSION FOR AG, IN, TH, SN, PB, SB, AND BI IN LIQUID
 GALLIUM WERE CALCULATED FROM VISCOSITY DATA. (GRA)
 <SCC >EDB-060100
 <DAV >NTIS \$3.25.
 <RNU >
 <RNO >TT--74-58011
 <SRN >DMDC--14094
 <PGM >10
 <BIB >GRA
 <RNA >EDB;ERA;NSA
 <TRN >TRANSLATED FROM
 <RPO >NP
 <LWC >SU
 <CPC >US

For the selected bibliography we chose not to print the entire
 listing but to show authors, corporate authors, titles, publication
 description, subject descriptors, abstract, subject categories [*ERDA
 Energy Information Data Base: Subject Categories* (TID-4584-R2)], and
 INIS subject categories. The printed citation then looks like the
 following, with some of the fields concatenated for convenience:

<54>

Besmann, T.M.; Lindemer, T.B. (Oak Ridge National Lab., Tenn. (USA))

Thermodynamic assessment of the conversion of plutonium dioxide to plutonium monocarbide AIX-08:334259;EDB-77:137277; J. Nucl. Mater. (JNUMA) 67:(1):. 77-87 (1977)

SUBJECT DESCRIPTORS: CHEMICAL REACTION YIELD:
Q2;EQUILIBRIUM;EVAPORATION;FREE ENERGY;OXYGEN;PHASE
DIAGRAMS;PLUTONIUM;PLUTONIUM CARBIDES: T2;PLUTONIUM DIOXIDE:
T1;REDUCTION: Q1;S CODES;SOLID SOLUTIONS;VERY HIGH TEMPERATURE

ABSTRACT: The equilibrium thermodynamics of the carbothermic reduction of PuO_2 to $\text{PuC}_{0.88}$ are reassessed in light of new thermodynamic values for the Pu-C-O system. Analysis of the thermodynamic data indicates that the PuC_xO_y phase can be successfully modeled by an ideal solid solution of $\text{PuC}_{0.88}$ and PuO . Equilibrium calculations show that oxygen-free monocarbide cannot be practically prepared by the direct carbothermic reduction of PuO_2 because of plutonium vaporization. These results imply that significant plutonium vaporization will occur in similarly reduced UO_2 - PuO_2 mixtures unless the actinides are initially present as a solid solution. Equilibrium calculations also suggest that $\text{PuC}_{0.88}$ can be successfully prepared by a two-step method: the carbothermic reduction of PuO_2 to $\text{PuC}_{1.5}$ followed by the hydrogen reduction of $\text{PuC}_{1.5}$ to $\text{PuC}_{0.88}$.

EEF SUBJECT CATEGORY: EDB-050700;360201;400702
INIS CATEGORY: E12

Naturally, we can select to print, sort, or index on any field of interest to the program people.

THIS PAGE
WAS INTENTIONALLY
LEFT BLANK

ACKNOWLEDGMENTS

Charles H. Bean, PDPM Program Manager, and Kevin M. Myles of the PDPM Program have supported and encouraged this effort so that the program people can evaluate a sample. L. M. Ferris, head of the Inorganic and Organic Section of the ORNL Chemistry Division, assumed a leading role in advocating the information support program and in discussing it with contacts throughout the program.

Many members of the ORNL Information Division and the Union Carbide Corporation-Nuclear Division Computer Sciences Division have been helpful with guidance and advice, as we undertook a very large effort on a very short time scale. In particular, we would like to express our appreciation to Herbert Pomerance, Cathy Fore, Helen Pfuderer, Tim Ensminger, Richard Green, Jackie Gillen, and Allen Singletary for their continuing and patient help. Special thanks must go to Faye Fletcher of the Information Division and Ruth Slusher of the Computer Sciences Division for their outstanding assistance with the management of our data sets and the computer production of this document. The Information Division units concerned with composition, makeup, printing, and distribution, as always, were necessary, efficient, and most cooperative.

<1>

Abdel-Aziz, A.H.K.; Kirshah, M.B. (Alexandria Univ. (Egypt).
Faculty of Engineering)

Density and temperature dependence of the surface tension of
molten bismuth, lead, and bismuth-lead alloys
ERA-03:012843; EDB-78:021748; Z. Metallkd. (ZEMTA)
68: (6):437-439 (Jun 1977)

SUBJECT DESCRIPTORS: BINARY ALLOY SYSTEMS; BISMUTH: T1; BISMUTH
ALLOYS: T2; DENSITY: Q1, Q2, Q3, Q4; EQUILIBRIUM; HIGH
TEMPERATURE; LEAD: T3; LEAD ALLOYS: T4; LIQUID METALS; PHASE
DIAGRAMS; SURFACE TENSION: Q1, Q2, Q3, Q4; TEMPERATURE
DEPENDENCE; VERY HIGH TEMPERATURE; VOLUME

ABSTRACT: The density and the temperature dependence of the
surface tension of molten bismuth, lead, and bismuth-lead
alloys were studied in the temperature range 623 to 953K,
using a maximum bubble pressure apparatus. In the temperature
range covered, the mean density of bismuth is 10,031 kg/m³
and that of lead is 10,530 kg/m³. The surface
tension of bismuth decreases from 0.379 N/m at 653K to 0.359
N/m at 953K, while that of lead decreases from 0.451 N/m at
633K to 0.409 N/m at 923K. The densities of the alloys and
the surface tension isotherms are all below the corresponding
additive straight lines.

EDB SUBJECT CATEGORY: EDB-360104; 360102

<2>

Ackermann, R.J.; Rauh, E.G., Argonne National Lab., Ill. (USA)

In Comparison of the thermodynamic properties and high
temperature chemical behavior of lanthanide and actinide
oxides ** ERA-03:007179; INS-78:000328; EDB-78:010143; (1977)
CONF-770662--3

SUBJECT DESCRIPTORS: ACTINIDE COMPOUNDS: T2; CHEMICAL PROPERTIES:
Q1, Q2, Q3; GASES; MOLECULAR IONS; OXIDES: T3; RARE EARTH
COMPOUNDS:
T1; SOLIDS; SPECTROSCOPY; STOICHIOMETRY; TABLES; THERMODYNAMIC
PROPERTIES: Q1, Q2, Q3; VAPORS

ABSTRACT: The thermodynamic properties of the lanthanide and
actinide oxides are examined, compared, and associated with a
variety of high temperature chemical behavior. Trends are
cited resulting from a number of thermodynamic and
spectroscopic correlations involving solid phases, species in
aqueous solution, and molecules and ions in the vapor phase.
Inadequacies in the data and alternative approaches are
discussed. The characterization of nonstoichiometric phases
stable only at high temperatures is related to a network of

<2>

<2> CONT.

heterogeneous and homogeneous equilibria. A broad perspective of similarity and dissimilarity between the lanthanides and actinides emerges and forms the basis of the projected needs for further study.

EDB SUBJECT CATEGORY: EDB-400702;400201;360204
INIS CATEGORY: B13;B12;B23

<3>

Afanas'ev, Yu.A.; Bayanov, A.P.; Frolov, Yu.A.

Thermodynamic properties of the liquid La--Mg alloys

EDE-76:048717; Izv. Akad. Nauk SSSR, Met. (IZNMA) (1):186-190
(Jan 1975)

SUBJECT DESCRIPTORS: CHEMICAL

COMPOSITION;DIAGRAMS;ENTROPY;EQUATIONS;FREE

ENTHALPY;LANTHANUM ALLOYS: T1;LIQUID METALS;MAGNESIUM ALLOYS:

T2;PRESSURE MEASUREMENT;REACTION KINETICS;TEMPERATURE

DEPENDENCE;THERMODYNAMIC PROPERTIES: Q1,Q2;VAPOR PRESSURE

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-360104
INIS CATEGORY: B22

<4>

Afinogenov, Yu.P.; Dantseva, V.A. (Voronezhskij Gosudarstvennyj Univ. (USSR))

Thermal analysis of $\text{GaCl}_3 + 3\text{In} = \text{InCl} + \text{Ga}$ system

EDE-76:036674; Zh. Neorg. Khim. (ZNOKA) 20:(2):516-518 (Feb 1975)

SUBJECT DESCRIPTORS: CHEMICAL COMPOSITION;CROSS

SECTIONS;DIFFERENTIAL THERMAL ANALYSIS:

Q1,Q2;EQUILIBRIUM;GALLIUM;GALLIUM CHLORIDES: T1;INDIUM;INDIUM

CHLORIDES: T2;PHASE DIAGRAMS

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-360602
INIS CATEGORY: B12

<5>

Ahmad, U.M.; Murr, L.E.

Variation of the solid--solid--nickel--thoria interfacial free energy with temperature
INS-76:014669; ERA-01:020089; EDB-76:054031; Metallography (MEIJA) 9: (1):33-41 (Feb 1976)

SUBJECT DESCRIPTORS: DISPERSION HARDENING; FREE ENERGY; GRAIN BOUNDARIES; INTERFACES; MATRIX MATERIALS; NICKEL ALLOYS: M1; SURFACE PROPERTIES: Q1, Q2; TEMPERATURE DEPENDENCE; THORIUM OXIDES: M2; VERY HIGH TEMPERATURE

ABSTRACT: The interfacial free energy in the nickel--thoria system was measured by comparing grain-boundary groove profiles at the surface and the interface of nickel-thoria bonded couples equilibrated at temperatures of 1300, 1320, 1337, 1373, 1400, and 1420⁰C. The temperature coefficient of interfacial free energy is positive in the range 1300-1400⁰C and negative above 1400⁰C. Results are indicative of a direct influence of changes of particle-matrix interfacial free energy on the coarsening of thoria particles in thoria-dispersed nickel.

EDB SUBJECT CATEGORY: EDB-360204; 360104
INIS CATEGORY: B23; B22

<6>

Alabyshev, A.F.; Morachevskii, A.G.; Kamenetskii, M.V.; Petrov, V.A.; Lisinskii, V.E. (Leningradskii Politekhnikheskij Inst. (USSR))

Thermodynamic properties of diluted solutions of $MnCl_2$ in alkali metal chlorides AIX-08:317661; EDB-77:133372; Zh. Prikl. Khim. (ZPKHA) 49: (1):215-217 (Jan 1976)

SUBJECT DESCRIPTORS: CESIUM CHLORIDES: T1; DILUTION; ELECTROMOTIVE FORCE; EQUATIONS; HIGH TEMPERATURE; MANGANESE CHLORIDES; POTASSIUM CHLORIDES; QUANTITY RATIO; REACTION KINETICS; SOLUTIONS; THERMODYNAMIC PROPERTIES: Q1

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-400201
INIS CATEGORY: B12

<7>

<7>

Alcock, C.B.; Jacob, K.T.; Zador, S. (Toronto Univ., Ontario
(Canada). Dept. of Metallurgy and Materials Science)

1. Thermochemical properties In Zirconium: Physico-chemical properties of its compounds and alloys ** At. Energy Rev., AIX-07:277692;EDB-77:031459; (1976) IAEA

SUBJECT DESCRIPTORS: THERMODYNAMIC PROPERTIES: Q1,Q2;ZIRCONIUM
ALLOYS: T2;ZIRCONIUM COMPOUNDS: T1;ZIRCONIUM COMPOUNDS: T1

ABSTRACT: A critical review of thermodynamic properties of Zr-compounds is presented. Binary and ternary Zr-compounds, pure Zr, and binary Zr-alloys are considered. The thermodynamic properties include heat capacity, enthalpy, gibbs free energy, entropy, vapor pressure, transition heat and heat of solution. Most of the values are given one by one in the text with some comments about its reliability and accuracy.

EDB SUBJECT CATEGORY: EDB-400201;360104
INIS CATEGORY: B12

<8>

Aleksandrovskii, S.V.; Sandler, R.A.; Naumchik, A.N.; Ivanov, I.I. (Leningradskij Gornyj Inst. (USSR))

Standard formation enthalpy of chloride alloys of $ZrCl_3$ sub
4\$--KCl(NaCl) systems AIX-08:298214;EDB-77:062544; Zh. Fiz.
Khim. (ZFKHA) 50:(9):2437 (Sep 1976)

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-400201
INIS CATEGORY: B12

<9>

Alexander, C.A.; Clark, R.B.; Kruger, O.L. (Battelle Columbus
Labs., Ohio (USA)); Robbins, J.L. (California Univ.,
Livermore (USA). Lawrence Livermore Lab.) ** Blank, H.;
Lindner, R. (eds.)

Fabrication and high temperature thermodynamic and transport
properties of Pu in Plutonium 1975 and other actinides **
AIX-08:309967;ERA-02:041838;EDB-77:094416; (1976)
North-Holland

SUBJECT DESCRIPTORS: CHEMICAL
COMPOSITION;DENSITY;ENTROPY;FORMATION HEAT;HOT
PRESSING;MACHINING;PLUTONIUM NITRIDES: T1;POWDERS;PRESSURE
DEPENDENCE;SPECIFIC HEAT;TEMPERATURE DEPENDENCE;THERMAL

<9> CONT.

CONDUCTIVITY; THERMAL DIFFUSIVITY; THERMAL
EXPANSION; THERMODYNAMIC PROPERTIES: Q1; VAPORIZATION HEAT; VERY
HIGH TEMPERATURE

ABSTRACT: Plutonium mononitride continues to be a potentially promising constituent nuclear fuel for the liquid metal fast-breeder reactor due to its high thermal conductivity and high melting point relative to other candidate fuels. A program was undertaken to measure thermodynamic properties of fuel at elevated temperatures: specifically, a pressure-composition diagram was generated for the plutonium-nitrogen system. Transport property measurements were also conducted as a function of sample density and temperature; thermal expansion measurements were made as were thermal diffusivity measurements. In the early stages of the program, it was necessary to spend significant effort developing sample fabrication techniques; this effort consisted of powder synthesis, hot pressing, and machining.

EDB SUBJECT CATEGORY: EDB-360204; C50700
INIS CATEGORY: B24

<10>

Alire, R.M.

Transport of hydrogen in liquid lithium

INS-76:019918; ERA-02:001035; EDB-76:081575; J. Chem. Phys.
(JCPSA) 65:(3):1134-1137 (1 Aug 1976)

SUBJECT DESCRIPTORS: DIFFUSION: Q1; HIGH TEMPERATURE; HYDROGEN:
T1; LIQUID METALS; LITHIUM: T2; LOW PRESSURE; MEDIUM
VACUUM; PERMEABILITY: Q2; SOLUBILITY: Q1; THERMONUCLEAR REACTOR
COOLING SYSTEMS; THERMONUCLEAR REACTORS; VERY HIGH TEMPERATURE

ABSTRACT: The solubilities and diffusion coefficients of $H_{2\%}$ in liquid Li contained in cylinders of Nb and Nb/1%Zr were determined in the temperature range 625--905°C and pressure range 13--1333 Pa. Simultaneous radial and axial diffusion into a finite solid cylinder adequately described the transport of $H_{2\%}$ into liquid Li. The diffusion coefficient was expressed by $D \text{ (cm}^2\text{/sec)} = 13 \pm 1 \exp(-2.5 \pm 0.2 \times 10^4 \text{ K/RT})$. (AIP)

EDB SUBJECT CATEGORY: EDB-360104; 360603; 700201
INIS CATEGORY: B22; B24; A14

<11>

<11>

Allen, C.W.; Liao, K.C.; Miller, A.E. ** Lundin, C.E. (ed.)

Fault structures in rare earth--cobalt intermetallics In
 Proceedings of the 12th rare earth research conference.
 Vol. I ** ERA-03:004343;INS-77:019109;EDB-78:004413; (1976)
 Univ. of Denver

SUBJECT DESCRIPTORS: COBALT ALLOYS: T2;CRYSTAL
 DEFECTS;MATHEMATICAL MODELS;PHASE TRANSFORMATIONS: Q1,Q2;RARE
 EARTH ALLOYS: T1;SHEAR PROPERTIES;TRANSITION ELEMENT ALLOYS

ABSTRACT: In dealing with materials which may undergo a shear transformation or which may contain planar compositional faulting, it is valuable to employ a stacking sequence notation, a structural symbolism, which may be developed for any layer-like structure. Such a conventional notation is presented, as far as possible, consistent for all possible phases in rare earth-transition element systems ranging from $RCo_{2.17}$ through $R_{2.17}Co_{17}$, and application of this notation to shear transformations, shear structure development and the nature of homologous series of phases within such systems is briefly illustrated. (GHT)

EIE SUBJECT CATEGORY: EDB-360102
 INIS CATEGORY: B22

<12>

Allen, G.C.; Crofts, J.A.; Griffiths, A.J. (Central Electricity
 Generating Board, Berkeley (UK). Berkeley Nuclear Labs.)

Infrared spectroscopy of the uranium/oxygen system
 AIX-08:298648;EDB-77:065887; J. Nucl. Mater. (JNUMA)
 62:(2-3):273-281 (Nov 1976)

SUBJECT DESCRIPTORS: ABSORPTION SPECTRA: Q1,Q2,Q3;ABSORPTION
 SPECTROSCOPY;BOND LENGTHS;DATA;INFRARED SPECTRA;SOLID
 SOLUTIONS;URANIUM DIOXIDE: T1;URANIUM OXIDES: T3;URANIUM
 OXIDES U3O8: T2

ABSTRACT: The infrared absorption spectra of six oxides of uranium, including several interstitial compounds, are presented and discussed. These oxides are: $UO_{2.00}$, $UO_{2.05}$, $UO_{2.08}$, $UO_{2.18}$, $U_{4.9}O_{9.9}$ and $U_{3.8}O_{8.8}$. Attention is drawn to certain features and trends of the spectra which lead to their use in the detection of solid solutions of oxygen in uranium dioxide, and in the analysis of uranium oxides of unknown composition.

<12> CONT.

Some inconsistencies are also discussed, which exist in previously published spectra of uranium oxides.

EDE SUBJECT CATEGORY: EDB-050700;400702

INIS CATEGORY: B24

<13>

Amberger, H.D.; Fischer, R.D.; Rosenbauer, G.G.; Spiegl, A.W.
(Erlangen-Nuernberg Univ., Erlangen (F.R. Germany). Inst.
fuer Anorganische Chemie)

Electronic structure of highly symmetric complexes of
lanthanoids and actinoids. Pt. 3. 'Electronic' Raman
spectrum of $\text{Cs}_{2\text{U(IV)Cl}_6}$
AIX-07:277954;EDB-77:031790; Ber. Bunsenges. Phys. Chem.
(BEPKA) 80:(6):495-499 (Jun 1976)

SUBJECT DESCRIPTORS: ABSORPTION SPECTRA;ACTINIDE
COMPLEXES;CESIUM CHLORIDES: T1;CRYSTAL
FIELD;CRYSTALS;ELECTRONIC STRUCTURE: Q1,Q2;ENERGY
LEVELS;MOLECULAR STRUCTURE;RAMAN SPECTRA: Q1,Q2;RARE EARTH
COMPLEXES;SYMMETRY;URANIUM CHLORIDES: T2

ABSTRACT: Besides the pure vibrational Raman bands, the
low-temperature Raman-spectrum of $\text{Cs}_{2\text{U(IV)Cl}_6}$
contains several additional signals. Since their appearance
is widely independent of the choice of the stimulating laser
line, we ascribe them to 'electronic' Raman effects. This
view is supported by the fact that these signals are either
close to bands which have been observed directly by
absorption spectroscopy or lie within narrow energy ranges of
crystal field states, which have been predicted by crystal
field theory. The number as well as the relative grouping of
the electronic Raman-bands show that the U^{4+} -central
ion is exposed to an effective crystal field of low symmetry.

EDE SUBJECT CATEGORY: EDB-400702;400201

INIS CATEGORY: E13

<14>

Amberger, H.D.; Fischer, R.D.; Rosenbauer, G.G.
(Erlangen-Nuernberg Univ., Erlangen (Germany, F.R.). Inst.
fuer Anorganische Chemie)

Electronic structure of highly symmetric complexes of
lanthanoids and actinoids. Pt. 4. The electronic Raman
spectrum of $\text{Cs}_{2\text{NaEu(III)Cl}_6}$
AIX-08:324055;EDB-77:146823; Z. Phys. Chem. (Frankfurt)
(ZPCFA) 102:(56):279-282 (Oct 1976)

<14>

<14> CONT.

SUBJECT DESCRIPTORS: CESIUM CHLORIDES: T1; CRYSTAL
 FIELD; CRYSTALS; ELECTRONIC STRUCTURE: Q1, Q2, Q3; ENERGY
 LEVELS; EUROPIUM CHLORIDES: T3; LOW TEMPERATURE; RAMAN
 EFFECT; RARE EARTH COMPLEXES; SODIUM CHLORIDES:
 T2; SPECTRA; SYMMETRY

ABSTRACT: The low-temperature Raman spectrum of the
 elpasclite-system $\text{Cs}_2\text{NaEu(III)Cl}_6$ exhibits five
 new sharp lines (compared with the room temperature
 spectrum), which are assigned to crystal field (=CF) Raman
 transitions between the CF-ground state GAMMA_1^0 and the octahedral CF-states of the free ion
 terms $7F_0^1$, $7F_2^1$, $7F_3^1$. A first-order
 CF-calculation yields: $E_4^0 = 170 \text{ cm}^{-1}$ and
 $E_6^0 = 3.2 \text{ cm}^{-1}$.

EDE SUBJECT CATEGORY: EDB-400201
 INIS CATEGORY: B13

<15>

Andrew, J.F.; Zocher, R.W.; Kent, R.A., Los Alamos Scientific
 Lab., N.Mex. (USA)

In Thermal expansion of $^{238}\text{PuO}_2$ **
 ERA-01:015780; INS-76:014275; EDB-76:045846; (Feb 1976)
 LA--6232-MS

SUBJECT DESCRIPTORS: PLUTONIUM 238: T2; PLUTONIUM DIOXIDE:
 T1; RADIOISOTOPE HEAT SOURCES: T, Q2; THERMAL EXPANSION:
 Q1; THERMOELECTRIC GENERATORS

ABSTRACT: The linear thermal expansion of stoichiometric $^{238}\text{PuO}_2$ was measured over the range 25 to 1600 K.
 The value of $\Delta L/L_0$ per degree C, calculated from
 the data, is $\Delta L/L_0 = -2.249 \times 10^{-4} + 9.020 \times 10^{-6}T - 1.130 \times 10^{-9}T^2 + 1.525 \times 10^{-12}T^3$.

EDE SUBJECT CATEGORY: EDB-070300
 INIS CATEGORY: D21

<16>

Andrews, L.

Optical spectra of the difluoride, dichloride, and trichloride
 ions in the matrix-isolated M^{2+}F_2 , M^{2+}Cl_2 , and M^{2+}Cl_3 species
 ERA-01:020227; EDB-76:054367; J. Am. Chem. Soc. (JACSA)
 98:(8):2147-2152 (14 Apr 1976)

<16> CONT.

SUBJECT DESCRIPTORS: ARGON;CESIUM CHLORIDES: M3;CHEMICAL BONDS;CHEMICAL REACTION KINETICS;CHLORIDES: M1;ENERGY-LEVEL TRANSITIONS;FLUORIDES: M2;INTERMEDIATE STRUCTURE;LITHIUM;MATRIX ELEMENTS;PHOTOLYSIS;POTASSIUM;SAPPHIRE;SODIUM;STABILITY;ULTRAVIOLET SPECTRA: Q1,Q2,Q3

ABSTRACT: Alkali metal atoms were codeposited with fluorine and chlorine at high dilution in argon on a sapphire plate cooled to 17⁰K. Ultraviolet spectra contained strong absorptions near 300 nm for $M^{+}F_2^{-}$ and 340 nm for $M^{+}Cl_2^{-}$ which are assigned to the $\sigma \rightarrow \sigma^*$ transition of the dihalide anion. This transition for Cl_2^{-} occurs at half of the energy known for Cl_2 . The alkali metal effect on the absorption spectra is consistent with an ionic model of polarized ion pairs. In addition, the strong ultraviolet band of the trichloride anion was observed at 250 nm with the metal-chlorine reactions and with the matrix reactions of $CsCl$ and Cl_2 . (auth)

EDB SUBJECT CATEGORY: EDB-400201

<17>

Andrews, L.

Optical spectra of the dibromide and diiodide ions in the matrix-isolated $M^{+}Br_2^{-}$ and $M^{+}I_2^{-}$ species ERA-01:020228;EDB-76:054368; J. Am. Chem. Soc. (JACSA) 98:(8):2152-2156 (14 Apr 1976)

SUBJECT DESCRIPTORS: ARGON;BROMIDES: M1;CESIUM BROMIDES: M3;CHEMICAL BONDS;CHEMICAL REACTION KINETICS;DIMERS;ENERGY-LEVEL TRANSITIONS;INTERMEDIATE STRUCTURE;IODIDES: M2;LITHIUM;MATRIX ELEMENTS;PHOTOLYSIS;POTASSIUM;SAPPHIRE;SODIUM;ULTRAVIOLET SPECTRA: Q1,Q2,Q3

ABSTRACT: The products of argon matrix reactions of alkali metal atoms and bromine and iodine have been studied optically in thin films on a sapphire window at 17⁰K. The bromine reaction produced a strong band near 360 and a weak band near 640 nm, depending upon the alkali reagent, and the reactions with iodine gave a strong absorption near 380 and a medium intensity absorption near 680 nm. These bands are due to $\sigma \rightarrow \sigma^*$ and $\pi \rightarrow \sigma^*$ transitions, respectively, of the dihalide anions in the $M^{+}X_2^{-}$ species. The small alkali metal effect on the transition energies is in accord with an ionic model of polarizable ion pairs, with the exception of cesium, which is capable of inner shell bonding and red-shifting the

<17>

<17> CONT.

transitions from their expected positions. In addition, a very strong 227 nm absorption in bromine and a strong 254, 286 nm doublet in iodine experiments are attributed to molecular halogen aggregates. (auth)

EDE SUBJECT CATEGORY: EDB-400201

<18>

Angelini, P.; Adair, H.L., Oak Ridge National Lab., Tenn. (USA)

In Preparation of ^{147}Pm metal and the determination of the melting point and phase transformation temperatures **

ERA-01:026076; INS-76:018743; EDB-76:074961; (Jul 1976)

ORNL/TM--4975

SUBJECT DESCRIPTORS: CHEMICAL PREPARATION:

Q1; DISTILLATION; MELTING POINTS: Q1; PHASE TRANSFORMATIONS:

Q1; PROMETHIUM: T1; PROMETHIUM 147; PROMETHIUM

OXIDES; QUARTZ; REDUCTION; THORIUM; VERY HIGH TEMPERATURE

ABSTRACT: The promethium metal used in the determination of the melting point and phase transformation temperatures was prepared by reduction of promethium oxide with thorium metal at $1600 \pm 0^\circ\text{C}$ and distilling the promethium metal into a quartz dome. The melting point and phase transformation temperatures of promethium metal were found to be $1042 \pm 5 \pm 0^\circ\text{C}$ and $890 \pm 5 \pm 0^\circ\text{C}$, respectively. The ratio for the heat of the high-temperature transformation to the heat of fusion was determined to be 0.415.

EDE SUBJECT CATEGORY: EDB-360104; 400703

INIS CATEGORY: B22; D13

<19>

Angerer, G. (Kernforschungszentrum Karlsruhe (Germany, F.R.).
Inst. fuer Neutronenphysik und Reaktortechnik) ** Jaeger,
T.A. (comp.)

Computational simulation of cladding motion after melting
including resolidification processes in Structural mechanics
in reactor technology. Vol. 1 **

AIX-08:287004; ERA-02:024766; EDB-77:049700; (1975)

North-Holland

SUBJECT DESCRIPTORS: COMPUTER CALCULATIONS; DRAG; FLOW BLOCKAGE:

Q2; FUEL CANS; FUEL-CLADDING INTERACTIONS; INTERNAL

FRICTION; LMFBR TYPE REACTORS: T1; LOSS OF FLOW:

M2, Q1; MELTING; MOTION; REACTOR ACCIDENTS; S

CODES; SIMULATION; SOLIDIFICATION; VISCOSITY

<19> CONT.

ABSTRACT: During flow coastdown or reactivity accidents in LWRs the reactor core may reach a condition where the cladding of the fuel pin starts to melt. The following cladding relocation influences reactor reactivity and core thermohydraulics. Plugging of coolant channels, e.g., may reduce dramatically the possibility of post accident heat removal in these channels. For this reason it is highly desirable to simulate and analyze cladding relocation. Due to the complexity of the problem a computer program is required. The code CMOT is an attempt to serve this purpose. Results are presented of code applications to an unprotected flow coastdown accident in a LMFBR. They show that due to the sodium vapor distribution in the voided core, upward flow of molten cladding may occur. This agrees with the observations made in experiments. But if the cladding plugs, the cross-section sodium vapor flow is inhibited and downward motion of additional molten cladding takes place.

EDB SUBJECT CATEGORY: EDB-220900;210500
INIS CATEGORY: C52

<20>

Antonov, B.D. (AN SSSR, Sverdlovsk. Inst. Elektrokhimii)

X-ray diffraction study into molten bromides and iodides of alkali metals AIX-07:270138;EDB-77:019325; Zh. Strukt. Khim. (ZSTKA) 16:(3):501-503 (May 1975)

SUBJECT DESCRIPTORS: ALKALI METAL COMPOUNDS: T1;BROMIDES: T3;CRYSTALLOGRAPHY;IODIDES: T2;ION DENSITY;MOLTEN SALTS;STRUCTURAL CHEMICAL ANALYSIS;X-RAY DIFFRACTION: Q1,Q2,Q3

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-400201
INIS CATEGORY: B12

<21>

Ardashnikova, E.I.; Borzenkova, M.P.; Novoselova, A.V.;
Batsanova, L.R.; Chuvaev, V.F. (Moskovskij Gosudarstvennyj Univ. (USSR))

Study of $\text{KBe}_{25}\text{DyF}_{33}$ cross-section of $\text{KF-BeF}_{33}\text{-DyF}_{33}$ ternary system and some properties of this cross section glasses AIX-08:320577;EDB-77:113818; Izv. Akad. Nauk SSSR, Neorg. Mater. (IVNMA) 11:(12):2169-2174 (Dec 1975)

SUBJECT DESCRIPTORS: BERYLLIUM FLUORIDES: T1;DENSITY;DIFFERENTIAL THERMAL ANALYSIS;DYSPROSIUM

<21>

<21> CONT.

FLUORIDES: T2; FLUORINE 19; GLASS; HYGROSCOPICITY; INFRARED
SPECTRA; NUCLEAR MAGNETIC RESONANCE; PHASE DIAGRAMS:
Q1, Q2, Q3; POTASSIUM FLUORIDES: T3; SOLUBILITY; X-RAY DIFFRACTION

ABSTRACT: A study has been made of the section $\text{KBeF}_{2.5}\text{-DyF}_{3.5}$ of the ternary system $\text{KF-BeF}_{2.5}\text{-DyF}_{3.5}$ by methods of differential-thermal and X-ray diffraction analyses. Glassy phases were studied by the NMR method, infrared spectroscopy and differential-thermal analysis. The phase diagram of the section was plotted. In the study of the section a wide range of glassy state was found (0-22 mol.% $\text{DyF}_{3.5}$) which corresponds to the lowest temperatures of the liquidus and primary crystallization of phases containing beryllium. Most stable (with respect to crystallization) glasses of the section are in the composition range of 10-20 mol.% $\text{DyF}_{3.5}$. It has been found that the maximum of the dependence of the second moment of the ^{19}F -NMR spectrum on the glass composition coincides with the maximum of a similar dependence of the density of glasses. Both maxima correspond to the point with the minimum temperature of the liquidus in the phase diagram of the section. Glasses containing 10-20 mol.% $\text{DyF}_{3.5}$ are most resistant against moisture. Glasses containing agglomerates of $\text{KBeF}_{2.5}$ are more hygroscopic and soluble.

EDB SUBJECT CATEGORY: EDB-360602
INIS CATEGORY: B12

<22>

Arnol'dov, M.N.; Bogdanov, Yu.V.; Ivanovskii, M.N.; Morozov, V.A.

Hydrogen solubility in liquid cesium

AIX-08:283271; EDB-77:043770; Izv. Akad. Nauk SSSR, Met.
(IZNMA) (4):30-32 (Jul 1976)

SUBJECT DESCRIPTORS: ACCURACY; CESIUM:
T1; COOLING; EQUATIONS; HEATING; HIGH TEMPERATURE; HYDROGEN:
T2; LIQUID METALS; LOW PRESSURE; QUANTITY
RATIO; SATURATION; SOLUBILITY: Q2; SOLVENT PROPERTIES:
Q1; THERMAL ANALYSIS

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-360104; 400201
INIS CATEGORY: B22

<23>

Arrance, F.C.; Plizga, M.J., Douglas Aircraft Co., Inc., Newport Beach, Calif. (USA). Astropower Lab.

In Stable inorganic matrix materials for high temperature batteries. Final report No. 1, 2 June 1967--2 February 1968
 ** EDB-76:059523; (Aug 1968) AD--676106

SUBJECT DESCRIPTORS: ALUMINATES; ALUMINIUM NITRIDES; BATTERY SEPARATORS: T2, Q1; CALCIUM COMPOUNDS; DESIGN; ELECTRIC CONDUCTIVITY; ELECTRIC-POWERED VEHICLES; HIGH TEMPERATURE; INORGANIC COMPOUNDS; IONIC CONDUCTIVITY; LITHIUM; LITHIUM CHLORIDES; LITHIUM-SULFUR BATTERIES: T1; MAGNESIUM COMPOUNDS; MAGNESIUM OXIDES; MATERIALS: Q2; MOLTEN SALTS; PHYSICAL PROPERTIES; POTASSIUM CHLORIDES; THORIUM OXIDES; ZIRCONATES

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-250903

<24>

Arsent'ev, P.P.; Vinogradov, B.G. (Moskovskij Inst. Stali i Splavov (USSR))

Viscosity characteristics of the iron-chromium-nickel melts
 EDB-76:048720; Izv. Vyssh. Uchebn. Zaved., Chern. Metall. (IVUMA) (3):14-17 (Mar 1975)

SUBJECT DESCRIPTORS: ACTIVATION ENERGY; CHEMICAL BONDS; CHEMICAL COMPOSITION; CHROMIUM ALLOYS: T1; DENSITY; ENTROPY; FREE ENERGY; IRON BASE ALLOYS: T2; LIQUID METALS; NICKEL ALLOYS: T3; PHASE DIAGRAMS; PHASE TRANSFORMATIONS; QUANTITY RATIO; VISCOSITY: Q1, Q2, Q3

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-360104

INIS CATEGORY: B22

<25>

Asher, R.C.; Davies, D.; Beetham, S.A. (UKAEA Research Group, Harwell. Atomic Energy Research Establishment)

Some observations on the compatibility of structural materials with molten lead AIX-08:332062; ERA-03:012851; EDB-78:021883; Corros. Sci. (CRRSA) 17:(7):545-557 (1977)

SUBJECT DESCRIPTORS: CAST IRON; CERAMICS: T3; COATINGS; COMPATIBILITY; COOLANTS; CORROSION: Q3; CORROSION INHIBITORS; CORROSION PROTECTION; CORROSIVE EFFECTS:

<25>

<25> CONT.

Q1; EMBRITTLEMENT: Q2; FILMS; LEAD: T1; LIQUID METAL COOLED
 REACTORS; LIQUID METALS; MASS TRANSFER; METALLURGICAL EFFECTS:
 Q1; MOLYBDENUM: T2; OXIDES; REACTOR
 COMPONENTS; SOLUBILITY; TITANIUM

ABSTRACT: Exploratory experiments have been carried out relating to the use of molten lead as a coolant in advanced nuclear reactors. It has been found that the attack of molten lead on structural materials is qualitatively related to the solubility of the solid in the lead. The attack can be greatly reduced by the presence of a protective oxide film on the solid, and the ease of formation of such a film depends on the nature of the solid and the oxidizing potential of the lead. The effectiveness as an 'inhibitor' of titanium dissolved in the lead has been confirmed. Mass transfer has been shown to be very inefficient, at least under the experimental conditions used, and reasons for this are discussed. A number of ceramic materials have been shown to be very resistant. A cast iron exhibited 'growth'. With two exceptions a range of coated materials showed unsatisfactory behaviour. An exploratory experiment on the susceptibility of molybdenum to liquid metal embrittlement did not result in failure of the specimen.

FEB SUBJECT CATEGORY: EDB-360105; 220300
 INIS CATEGORY: B15

<26>

Pabayan, G.G.; Mkrtchyan, R.F.; Ter-Arakelyan, K.A. (Erevanskij Gosudarstvennyj Univ. (USSR))

Study of chemical interaction in $Pb_{3}AlF_{6}-LaF_{6}-Ca_{3}AlF_{6}$ system AIX-08:285971; EDB-77:043570;
 Arm. Khim. Zh. (AYKZA) 10::847-849 (1975)

SUBJECT DESCRIPTORS: ALUMINIUM FLUORIDES: T1; CALCIUM FLUORIDES: T2; CHEMICAL REACTIONS: Q1, Q2, Q3, Q4; CRYSTALS; HIGH TEMPERATURE; LANTHANUM FLUORIDES: T4; LEAD FLUORIDES: T3; PHASE DIAGRAMS; THERMAL ANALYSIS; X-RAY SPECTROSCOPY

ABSTRACT: None

FEB SUBJECT CATEGORY: EDB-360101
 INIS CATEGORY: B12

<27>

Babikova, T.F.; Sokolovskii, Yu.S.; Sroelova, M.M.

Interaction of uranyl chloride with zirconium tetrachloride in
the melt of sodium chloride and potassium chloride
AIX-08:324057;EDB-77:146936; At. Energ. (USSR) (AENGA)
40:(5):412 (May 1976)

SUBJECT DESCRIPTORS: CHEMICAL ANALYSIS;CHEMICAL REACTION
KINETICS: Q1,Q2;COMPLEXES;HIGH TEMPERATURE;MOLTEN
SALTS;POTASSIUM CHLORIDES;QUANTITY RATIO;SODIUM
CHLORIDES;SPECTROPHOTOMETRY;URANYL CHLORIDES: T2;ZIRCONIUM
CHLORIDES: T1

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-400702
INIS CATEGORY: B13

<28>

Baer, Y.; Zuercher, C.

Electronic structure of CeN studied by x-ray-photoemission
spectroscopy INS-78:000983;EDB-78:016935; Phys. Rev. Lett.
(PRLTA) 39:(15):956-959 (10 Oct 1977)

SUBJECT DESCRIPTORS: CERIUM;CERIUM NITRIDES: T1;D
STATES;ELECTRONIC STRUCTURE: Q1;F STATES;PHOTOELECTRON
SPECTROSCOPY;PHOTOEMISSION;VALENCE;X-RAY SPECTRA

ABSTRACT: The core levels and the valence-band region CeN have
been studied by x-ray photoemission. Clear evidence is found
that Ce has a mixed valence between 3 and 4 in this compound.
A comparison of the 4d spectra of the Ce pnictides and of
pure Ce at different temperatures leads to the conclusion
that the α phase of this metal corresponds very likely
to an intermediate valence.

EDB SUBJECT CATEGORY: EDB-360202
INIS CATEGORY: B23

<29>

Baernighausen, H.

In Mixed valence rare-earth halides and their unusual crystal
structures ** AIX-07:257159;EDB-77:013440; (1976)
AEE-Ccnf--76-033-001

SUBJECT DESCRIPTORS: COMPARATIVE EVALUATIONS;COORDINATION
NUMBER;CRYSTAL STRUCTURE: Q1,Q2;DYSPROSIUM CHLORIDES:
T1;HALIDES;MONOCRYSTALS;RARE EARTH

<29>

<29> CONT.

COMPOUNDS; SOLIDS; VALENCE; YTTERBIUM CHLORIDES: T2

ABSTRACT: None

EEL SUBJECT CATEGORY: EDB-400201

INIS CATEGORY: B12

<30>

Bagai, R.K.; Warriar, A.V.R. (Solid State Physics Lab., Delhi (India))

Optical absorption spectrum of UO_2^{2+} doped LiF
 AIX-08:338785; EDB-78:022026; J. Phys., C (London) (JPSCA)
 10: (5):L437-L439 (14 Aug 1977)

SUBJECT DESCRIPTORS: ABSORPTION SPECTRA: Q1, Q2; ANIONS; CRYSTAL
 LATTICES; DOPED MATERIALS; ELECTRONIC STRUCTURE; LITHIUM
 FLUORIDES: T1; MOLECULAR ORBITAL METHOD; MOLECULAR
 STRUCTURE; MONOCRYSTALS; URANIUM OXIDES: T2; URANYL COMPOUNDS

ABSTRACT: The absorption spectrum of LiF single crystals doped
 with UO_2^{2+} ion is measured in the region
 extending into the vacuum ultraviolet. It is proposed that
 the UO_2^{2+} ion enters the lattice substitutionally
 at the Li^+ site with the oxygen directed along $\langle 111 \rangle$.
 The observed spectrum is interpreted on the basis of the
 molecular orbital scheme given by McGlynn and Smith (J.
 Molec. Spectrosc.; 6:164 (1961)).

EEL SUBJECT CATEGORY: EDB-360603; 360204

INIS CATEGORY: A13

<31>

Balashov, D.B.; Ikhenov, D.A.

Nuclear quadrupole resonance spectra of HgCl_2 under
 pressure AIX-08:338791; EDB-78:022016; Fiz. Tverd. Tela
 (FTVTA) 17: (9):2693-2696 (Sep 1975)

SUBJECT DESCRIPTORS: ABSORPTION SPECTRA; CHLORINE 35:
 T1; CRYSTAL-PHASE TRANSFORMATIONS: Q2; LOW TEMPERATURE; MEDIUM
 TEMPERATURE; MERCURY CHLORIDES: T2; NUCLEAR QUADRUPOLE
 RESONANCE: Q1; PRESSURE DEPENDENCE; TEMPERATURE DEPENDENCE; VERY
 HIGH PRESSURE

ABSTRACT: The aim of the paper is to study the spectrum of
 nuclear quadrupole resonance (NQR) observed on $\text{Cl}^{35}/\text{H}^5$
 nuclei in a HgCl_2 polycrystalline powder. The spectrum
 has been studied at hydrostatic pressures up to 16.000
 kg/cm² and at temperatures of 256.5, 312.0, and 356.0

<31> CONT.

deg K using a previously developed high-pressure apparatus and a stationary NQR spectrometer. The accuracy of the measurements of the NQR frequency has been about 2KHz and of the pressure, about 50 kg/cm². The temperature stabilized within ± 0.1 deg K. The spectrum consists of two absorption lines due to the crystallographic nonequivalence of the position of chlorine atoms of each molecule in the lattice. Both frequencies decrease with temperature and decrease linearly with pressure. At pressures 6.500 to 6.900 kg/cm² rising with temperature, the frequency isothermals exhibit a sudden change accompanied by a decrease in slope at higher pressures; this fact points to the existence of a phase transition.

EDB SUBJECT CATEGORY: EDB-360602
INIS CATEGORY: A13

<32>

Balej, J. (Ceskoslovenska Akademie Ved, Prague)

Standard potential of sodium amalgam at 25⁰C
AIX-08:286395;EDB-77:068512; Electrochim. Acta (ELCAA)
21:(11):953-956 (Nov 1976)

SUBJECT DESCRIPTORS: ELECTROLYTIC CELLS;ELECTROMOTIVE FORCE:
Q1,Q2;MERCURY ALLOYS: T1;REACTION KINETICS;SODIUM;SODIUM
ALLOYS: T2;THERMODYNAMIC PROPERTIES: Q1,Q2

ABSTRACT: From a set of critically selected literary data on the emf's of concentration cells consisting of sodium metal and sodium amalgam in a sodium salt non-aqueous electrolyte a value of $E^0_{\text{Na(Hg)/Na}^+} = -1.9558 \text{ V}$ was calculated for the standard equilibrium potential of diluted liquid sodium amalgam at 25⁰C. The standard state of the amalgam was derived from an amalgam at infinite dilution where $f_{\text{Na}} \rightarrow 1.0$ for $x_{\text{Na}} \rightarrow 0$. The activity coefficient of sodium in the amalgam at an arbitrary composition, corresponding to this standard state, is obtained from the expression $\lg f_{\text{Na(Hg)}} = 16.393 x_{\text{Na}}$.

EDB SUBJECT CATEGORY: EDB-360104
INIS CATEGORY: B22

<33>

<33>

Balourdet, Marcel; Malmejac, Yves; Desre, Pierre, CEA Centre
d'Etudes Nucleaires de Grenoble, 38 (France). Dept. de
Metallurgie

In Thermodiffusion measurements in liquid alloys and electron
transport heat calculations in the case of almost-free
electrons ** AIX-08:292740;EDB-77:074275; (17 Jun 1976)
CEA-CCNF--3581

ABSTRACT: The thermodiffusion of antimony, indium and silver in
pure liquid tin was studied and the transport heat of these
tracers was determined. The thermodiffusion of silver in
liquid tin presents an inversion at about 600 deg C. This
seems to result from the competition between two phenomena;
the first one is intrinsic and proportional to the absolute
temperature, the other one (electronic) is proportional to
the square of this temperature. The thermodiffusion of Ag in
liquid Ag-Ge of eutectic concentration was also studied.

EDE SUBJECT CATEGORY: EDB-360102

INIS CATEGORY: B22

<34>

Barbanel', Yu.A.; Kotlin, V.P.; Kolin, V.V. (Radievyy Inst.,
Leningrad (USSR)) ** AN SSSR, Moscow; Gosudarstvennyj Komitet
po Ispol'zovaniyu Atomnoj Ehnergii SSSR, Moscow

Absorption spectra of curium (3) chloride complexes In All-union
conference on chemistry of transplutonium elements
(americium, curium, berkelium, californium) **
AIX-08:295437;EDB-77:075240; (1976) INIS-mf--3283

SUBJECT DESCRIPTORS: ABSORPTION SPECTRA: Q1;AMERICIUM
CHLORIDES;CESIUM COMPOUNDS;CHLORIDES;CURIUM COMPLEXES:
T1;HIGH TEMPERATURE;LANTHANUM CHLORIDES;LOW
TEMPERATURE;SODIUM COMPOUNDS

ABSTRACT: None

EDE SUBJECT CATEGORY: EDB-400702

INIS CATEGORY: B13

<35>

Barker, M.G.; Hooper, A.J. (Nottingham Univ. (UK))

Reactions of liquid sodium with transition-metal oxides. VIII.
 The oxides of chromium-(III), -(IV), and -(VI) and disodium
 chromium(VI) tetraoxide AIX-07:272368;EDB-77:026444; J. Chem.
 Soc. (London), Dalton Trans. (JCOTB) (12):1093-1095 (1976)

SUBJECT DESCRIPTORS: CHEMICAL REACTIONS: Q1,Q2,Q3;CHROMATES:
 T2;CHROMIUM OXIDES: T1;CORROSION;HIGH TEMPERATURE;LIQUID
 METALS;MEDIUM TEMPERATURE;OXYGEN;SODIUM: T3;SODIUM
 COMPOUNDS;SODIUM OXIDES;STRUCTURAL CHEMICAL ANALYSIS;X-RAY
 DIFFRACTION

ABSTRACT: The reactions of the chromium oxides CrO_3 , CrO_2 , Cr_2O_3 , and Na_2CrO_4 with liquid sodium have been studied at up to 600°C , in some cases using differential thermal analysis. The reaction products have been examined by powder X-ray crystallography. The ternary oxide NaCrO_2 is formed in each case in which reaction took place, and this compound does not react further with dissolved oxygen in sodium up to the solubility limit. It is shown that the reaction of CrO_3 with liquid sodium does not yield Na_2CrO_3 as previously thought; this error is rationalised in terms of the experimental procedure, and improved techniques are identified.

EDB SUBJECT CATEGORY: EDB-400201

INIS CATEGORY: B12

<36>

Bas'kova, N.A.; Majer, A.A.; Katapet'yants, M.Kh.

Thermodynamic properties of some rare earth carbonates and
 yttrium carbonates EDB-76:036711; Tr. Mosk. Khim.-Tekhnol.
 Inst. (TMKIA) (81):8-10 (1974)

SUBJECT DESCRIPTORS: ENTROPY;EQUATIONS;FORMATION FREE
 ENERGY;FORMATION HEAT;SOLUBILITY;THERMODYNAMIC PROPERTIES:
 Q1,Q2,Q3;THERMODYNAMICS

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-360603

INIS CATEGORY: B12

<37>

<37>

Bauman, H.F., Lockheed Missiles and Space Co., Palo Alto, Calif.
(USA). Materials Sciences Lab.

In Lithium anode limited-cycle battery investigation.
Quarterly technical progress report No. 5 ** EDB-76:047722;
(May 1966) AD--483237

SUBJECT DESCRIPTORS: DESIGN;LITHIUM;METAL-NONMETAL BATTERIES: M1

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-250901;400201

<38>

Bauman, H.F.; Chilton, J.E.; Mauri, R.E., Lockheed Missiles and
Space Co., Palo Alto, Calif. (USA). Materials Sciences Lab.

In Lithium anode limited-cycle battery investigation.
Quarterly technical progress report No. 3 ** EDB-76:047821;
(Nov 1965) AD--474219

SUBJECT DESCRIPTORS: ANODES: Q1;CARBONIC ACID ESTERS;CATHODES:
Q1;COBALT FLUORIDES;COPPER FLUORIDES;ELECTRIC
CURRENTS;ELECTRIC
POTENTIAL;ELECTROLYTES;FABRICATION;LITHIUM;MATERIALS;METAL-NO
METAL BATTERIES: M1;PERFORMANCE;POLARIZATION;REACTION
HEAT;TEMPERATURE DEPENDENCE

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-250903

<39>

Fayonov, A.P.; Frolov, Yu.A.; Afanas'ev, Yu.A.

Investigation into thermodynamic properties of Ce-Mg melts
EDB-76:042263: Izv. Akad. Nauk SSSR, Met. (IZNMA) (3):91-95
(May 1975)

SUBJECT DESCRIPTORS: CERIUM ALLOYS: T1;LIQUID METALS;MAGNESIUM
ALLOYS: T2;PHASE DIAGRAMS;THERMODYNAMIC PROPERTIES:
Q1,Q2;VAPOR PRESSURE;VERY HIGH TEMPERATURE

ABSTRACT: The method of transfer in an argon current was used
for studying the pressure of saturated steam over liquid
alloys of Ce-Mg of nine concentrations in a temperature range
of 1083 to 1133^{sup} 0\$K, for a magnesium content of 0.134 to
0.926 at.%. Calculations are presented of thermodynamic
activities of the cerium and magnesium, their partial-molar
and integral Gibbs energies, and the enthalpy and entropy of

<39> CONT.

the mixture. The values found show negative deviations from the laws of the behavior of ideal solutions. The minimum of the integral thermodynamic values shows a shift towards magnesium.

EDB SUBJECT CATEGORY: EDB-360104
INIS CATEGORY: B22

<40>

Bayanov, A.P.; Ganchenko, E.N.; Afanas'ev, Y.A. (Kubanskiy Gosudarstvennyj Univ., Krasnodar (USSR))

Investigation into neodymium chloride state in equilibrium with metal in eutectic melt of potassium- and lithium chlorides
AIX-08:317665; EDB-77:133380; Zh. Fiz. Khim. (ZFKHA)
49:(9):2452 (Sep 1975)

SUBJECT DESCRIPTORS: CHEMICAL STATE: Q1,Q2,Q3;ELECTRIC BATTERIES;ELECTROMOTIVE FORCE;EQUILIBRIUM;EUTECTICS;HIGH TEMPERATURE;LITHIUM CHLORIDES: T1;MOLTEN SALTS;NEODYMIUM CHLORIDES: T2;POTASSIUM CHLORIDES: T3

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-400201
INIS CATEGORY: B12

<41>

Becker, R.; Lentz, A.; Sawodny, W. (Ulm Univ. (F.R. Germany). Abt. Anorganische Chemie)

Vibrational spectra and force constants of the elpasolites Cs_2KMfF_6 (M=Sc, Y, La, Gd, Yb)
AIX-07:262488; EDB-77:031452; Z. Anorg. Allg. Chem. (ZAACA)
420:(3):210-218 (1976)

SUBJECT DESCRIPTORS: CESIUM FLUORIDES: T1;CHEMICAL PREPARATION;FLUORIDES;GADOLINIUM FLUORIDES: T2;INFRARED SPECTRA: Q1,Q2,Q3,Q4,Q5,Q6,Q7;LANTHANUM FLUORIDES: T3;LATTICE VIBRATIONS;POTASSIUM FLUORIDES: T6;RAMAN EFFECT;SCANDIUM FLUORIDES: T5;VIBRATIONAL STATES;YTTERBIUM FLUORIDES: T4;YTTRIUM FLUORIDES: T7

ABSTRACT: The vibrational spectra of the elpasolites Cs_2KMfF_6 (M=Sc, Y, La, Gd, Yb) have been recorded and assigned including the lattice vibrations. The vibrational frequencies thus obtained were used for the calculation of

<41>

<41> CONT.

MVFF force constants. The values of the stretching force constants are discussed.

EDB SUBJECT CATEGORY: EDB-400201
INIS CATEGORY: B12

<42>

Behl, W.K., Army Electronics Command, Fort Monmouth, N.J. (USA)

In Electrode reactions in molten sodium nitrate--barium nitrate eutectic at 350 degrees C ** EDB-77:073970; (Feb 1970)
AD--703842

SUBJECT DESCRIPTORS: BARIUM NITRATES;ELECTRIC BATTERIES:
T1;ELECTROLYTES: Q1;LEAD;MOLTEN SALTS;SILVER;SODIUM
NITRATES;VOLTAMETRY

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-250903

<43>

Belov, S.F.; Gladneva, A.F. (Moskovskij Inst. Tonkoj
Khimicheskij Tekhnologii (USSR))

Density, viscosity, electric conductivity and meltability of
Na₃AlF₆-La₂O₃ and Na₃AlF₆-
6f-Sm₂O₃ systems AIX-08:295259;EDB-77:056159;
Izv. Vyssh. Uchebn. Zaved., Tsvet. Metall. (IVUTA)
(2):110-113 (1976)

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-400201
INIS CATEGORY: B12

<44>

Belyi, M.U.; Kushnirenko, I.Ya. ** AN Ehstonskoj SSR, Tallin

Spectral properties of impurity molecules of the Csub(3v)
symmetry groups in solid solutions of halide salts In 11.
european congress on molecular spectroscopy. Abstracts **
EDE-76:060598; (1973) INIS-mf--2069

SUBJECT DESCRIPTORS: ABSORPTION SPECTRA;ANIONS;BORATES;CESIUM
CHLORIDES: T1;EMISSION SPECTRA;IMPURITIES;LUMINESCENCE;SOLID
SOLUTIONS;SPECTROSCOPY: Q1;TELLURATES

<44> CONT.

ABSTRACT: None

EDE SUBJECT CATEGORY: EDB-400201

INIS CATEGORY: B12

<45>

Beresneva, T.I.; Denisova, L.A.; Fedorova, N.V.; Ketov, A.N.
(Perm'skij Farmatsevticheskiy Inst. (USSR))

Me₂ZrCl₆--H₂(Me-Na,K,Rb,Cs) system
EDE-76:036676; Zh. Neorg. Khim. (ZNOKA) 20:(2):377-381 (Feb 1975)

SUBJECT DESCRIPTORS: CHEMICAL ANALYSIS;CHEMICAL REACTION
KINETICS;HYDROGEN;PHASE DIAGRAMS:
Q1,Q2,Q3,Q4,Q5;REDUCTION;TEMPERATURE DEPENDENCE;THERMAL
ANALYSIS;X-RAY DIFFRACTION;ZIRCONIUM CHLORIDES: T1

ABSTRACT: None

EDE SUBJECT CATEGORY: EDB-360602

INIS CATEGORY: B12

<46>

Berezhinsky, I.I.; Gnatovskaya, V.N.; Lisitsa, M.P.; Tarasov,
G.G.; Khalimonova, I.N. ** AN Ehstonsoj SSR, Tallin

Temperature effect on vibrational spectra of the impurity
molecular ions in alkali halides In 11. european congress on
molecular spectroscopy. Abstracts ** EDB-76:060599; (1973)
INIS-mf--2069

SUBJECT DESCRIPTORS: ABSORPTION SPECTRA;CESIUM CHLORIDES:
T1;CRYSTAL LATTICES;IMPURITIES;IONS;TEMPERATURE
DEPENDENCE;VIBRATIONAL STATES: Q1

ABSTRACT: None

EDE SUBJECT CATEGORY: EDB-400201

INIS CATEGORY: B12

<47>

<47>

Berezin, B.Y.; Kats, S.A.; Chekhovskoi, V.Y.

Spectral emissivities of molten refractory metals

ERA-02:022873;EDB-77:043793; High Temp. (USSR) (Engl. Transl.) (HTEA) 14:(3):448-452 (Nov 1976)

SUBJECT DESCRIPTORS: EMISSIVITY: Q1,Q2,Q3,Q4,Q5;LIQUID
 METALS;MOLYBDENUM: M1;NIOBIUM: M2;RUTHENIUM: M5;TITANIUM:
 M4;VANADIUM: M3;VISIBLE RADIATION

ABSTRACT: Emissivity measurements at 0.65 μm have been made for liquid molybdenum, niobium, vanadium, titanium, and ruthenium at their melting points; the specimens were heated by levitation, with the brightness-temperature measurement at the melting point and the calculation of ϵ/λ , $1/\lambda$ from Wien's formula. (AIP)

EDB SUBJECT CATEGORY: EDB-360104

<48>

Berndt, K.

In Studies of the electronic structure of simple metals and their alloys with the aid of Compton scattering and positron annihilation using lithium, magnesium, lithium-magnesium and aluminium-zinc as examples ** AIX-07:243024;EDB-76:071212; (1973) Martin-Luther-Universitaet

SUBJECT DESCRIPTORS: ALUMINIUM BASE ALLOYS:
 T1;ANNIHILATION;BINARY ALLOY SYSTEMS;COMPTON EFFECT;COMPTON
 SPECTROMETERS;CROSS SECTIONS;ELECTRONIC STRUCTURE:
 Q1,Q2,Q3;ELECTRONS;FERMI LEVEL;LITHIUM:
 T2;MAGNESIUM;MAGNESIUM BASE ALLOYS: T3;VALENCE

ABSTRACT: The results show that the positron annihilation is an appropriate method for receiving information on the Fermi levels of simple metals. Concerning the global electronic structure of alloys, however, its applicability is limited, particularly in the case of Al-Zn, whereas the local electronic structure can be determined. The Compton profiles were calculated for the metals and compared with experimental results. The influence of the potential Fourier components, of the orthogonalization properties and of interactions on the electronic structure was studied. A first insight into the electronic structure of Zn-Al was gained.

EDB SUBJECT CATEGORY: EDB-360102
 INIS CATEGORY: A13

<49>

Berndt, U.; Maier, D.; Keller, C.

Phase equilibrium in interlanthanide oxides systems

EDE-76:042336; J. Solid State Chem. (JSSCB) 16:(1-2):189-195
(1 Jan 1976)

SUBJECT DESCRIPTORS: LANTHANUM OXIDES: M1;PHASE STUDIES:

Q1,Q2,Q3;RARE EARTH OXIDES: M2;TEMPERATURE DEPENDENCE;VERY
HIGH TEMPERATURE;YTTRIUM OXIDES: M3

ABSTRACT: Investigations in the quasibinary systems $\text{LaO}_{1/2}\text{--REO}_{1/2}$ (Ce--Lu , Y) and $\text{YO}_{1/2}\text{--REO}_{1/2}$ (La--Lu) at 1250 to 1550°C have shown that besides a perovskite compound ABO_3 (for $A = \text{La}$ yields $B = \text{Y}$, Ho--Lu) mixed oxide phases with hexagonal (A-type), monoclinic (B-type) and cubic (C-type) structures of varying phase widths exist. All four phases are only existent in the $\text{LaO}_{1/2}\text{--HoO}_{1/2}$ and $\text{PrO}_{1/2}\text{--LuO}_{1/2}$ systems. The mutual solubility of the individual lanthanide oxides strongly depends on the difference of the ionic radii of the elements, in general, increasing with decreasing difference. Maximum phase widths of B-type mixed oxides at 1550°C occur at $\text{RE} = \text{Tb}$ for the $\text{LaO}_{1/2}\text{--REO}_{1/2}$ systems and at $\text{RE} = \text{Nd}$ for the $\text{YO}_{1/2}\text{--REO}_{1/2}$ systems. At 1400°C up to 9 mole percent $\text{EuO}_{1/2}$ may be incorporated in the perovskite compound PrLuO_3 . For the $\text{LaYO}_3\text{--LaErO}_3$ series a maximum phase width with regard to the $\text{La}:(\text{Y} + \text{Er})$ ratio is to be observed at the 1:1 composition.

EDE SUBJECT CATEGORY: EDE-360202

<50>

Berndt, U.; Tanamas, R.; Keller, C.

Ternary $\text{UO}_2\text{--UO}_3\text{--EuO}_{1/2}$ system and investigation of $\text{Eu(II)--actinide(IV)--perovskites}$ INS-76:013206;EDB-76:042342; J. Solid State Chem. (JSSCB)
17:(1-2):113-120 (Feb 1976)

SUBJECT DESCRIPTORS: BURNABLE POISONS;ELECTROMOTIVE

FORCE;EUROPIUM OXIDES: T3;MIXTURES;PHASE DIAGRAMS:

Q1,Q2,Q3;THERMODYNAMIC PROPERTIES;URANIUM DIOXIDE: T1;URANIUM
TRIOXIDE: T2;VALENCE;VERY HIGH TEMPERATURE

ABSTRACT: At 1250°C, the following single-phase regions have been detected in the $\text{UO}_2\text{--UO}_3\text{--EuO}_{1/2}$ region of the ternary uranium--europium--oxygen system: (a) $\beta\text{-U}_3\text{O}_8$: no solubility for

<50>

<50> CONT.

$\text{EuO}_{1.5}$ could be observed. (b) Fluorite phase: the oxygen limiting compositions are $(\text{U}, \text{Eu})\text{O}_{2.25}$ and $(\text{U}, \text{Eu})\text{O}_{1.75}$, respectively; the range of the stoichiometric composition MO_2 is between UO_2 and $(\text{U}_{0.36}\text{Eu}_{0.64})\text{O}_2$, a fluorite phase with U(VI) is only obtainable for the substoichiometric region from $(\text{U}_{0.32}\text{Eu}_{0.68})$ to $(\text{U}_{0.24}\text{Eu}_{0.76})\text{O}_2$. (c) Rhombohedral phase: it extends from $\text{UO}_{3.6}\text{EuO}_{1.5}$ on the UO_3 -- $\text{EuO}_{1.5}$ side of the system to the 0.7UO_2 · $0.3\text{EuO}_{1.5}$ -- 0.6UO_2 · $0.4\text{EuO}_{1.5}$ line on the UO_2 -- $\text{EuO}_{1.5}$ side, showing an increased phase width with lowering of the uranium valency. (d) Monoclinic B-- $\text{EuO}_{1.5}$: no solubility for UO_2 could be observed. The first lanthanide--actinide perovskite compounds, orthorhombic EuUO_3 , and EuNpO_3 could be prepared by different methods; attempts to prepare similar ternary oxides containing Th(IV), Pu(IV), and Am(IV), however, failed. EMF measurements have proven that cubic C-- $\text{EuO}_{1.5}$ is the thermodynamically stable. $\text{EuO}_{1.5}$ -modification below $1050 \pm 20^\circ\text{C}$.

EDB SUBJECT CATEGORY: EDR-360202

INIS CATEGORY: E23

<51>

Bernstein, E.R.; Meredith, G.R., Colorado State Univ., Fort Collins (USA). Dept. of Chemistry

In Vibrational spectra of transition metal hexafluoride crystals. I. Orthorhombic MoF_6 , WF_6 , UF_6 neat crystals. Technical report ** ERA-03:001204; EDB-77:146753; (Apr 1977) AD-A--039765

SUBJECT DESCRIPTORS: CRYSTALS; MOLECULAR STRUCTURE:

Q1, Q2, Q3; MOLYBDENUM FLUORIDES: T1; RAMAN SPECTRA; TUNGSTEN FLUORIDES: T2; URANIUM HEXAFLUORIDE: T3

ABSTRACT: Neat crystal Raman spectra of orthorhombic MoF_6 , WF_6 , and UF_6 are presented and discussed. The neat crystal spectra are interpreted in terms of an internal-external mode separation and a conventional factor group analysis. In general, these data lead to four essential conclusions concerning hexafluoride crystals: (a) Fermi resonance plays an important role in producing the observed energy and intensity differences in this series; (b) $k = 0$ exciton structures are remarkably similar in the three crystals; (c) similarities and differences between these solids can be directly correlated with molecular parameters such as dipole

<51> CONT.

derivatives, polarizabilities, etc.; and (d) phonons (external modes) can be neatly separated into rotational and translational normal modes.

EDB SUBJECT CATEGORY: EDB-400201;400702

<52>

Bernstein, E.R.; Meredith, G.R., Colorado State Univ., Fort Collins (USA). Dept. of Chemistry

In Vibrational spectra of transition metal hexafluoride crystals. III. Exciton band structures of MoF_6 , WF_6 , and UF_6 . Technical report **
INS-77:017680;ERA-03:001606;EDB-77:148576; (Apr 1977)
AD-A--039793

SUBJECT DESCRIPTORS: BAND THEORY;CRYSTALS;FERMI LEVEL;MOLYBDENUM FLUORIDES: T1;TUNGSTEN FLUORIDES: T2;URANIUM HEXAFLUORIDE: T3;VIBRATIONAL STATES: Q1,Q2,Q3

ABSTRACT: This paper presents the overall vibrational band structure of MF_6 crystals based on the previously discussed two-particle, heavily doped mixed crystal, and neat crystal data in addition to dilute mixed crystal studies. Calculations of $\nu(2)$ and $\nu(5)$ bands based on a quadrupole-quadrupole intermolecular interaction model are determined to be in substantial agreement with the observations. The $\nu(1)$ bands are discussed in detail for a series of concentration studies, and it is possible to demonstrate that MoF_6 and WF_6 conform to the ideal mixed crystal model whereas crystals containing UF_6 do not. A consistent picture of gas-to-crystal shifts, resonance interactions, band shapes and structures, densities of exciton states, Fermi resonance, and simple multipolar interaction models emerges from these studies.

EDB SUBJECT CATEGORY: EDB-640302;400201;400702;360603
INIS CATEGORY: A12;B12;B13;B24

<53>

Bernstein, E.R.; Meredith, G.R., Princeton Univ., N.J. (USA). Dept. of Chemistry

In Interactions in inorganic molecular crystals - electronic spectra of ReF_6 pure and mixed crystals. Technical report, 1 Jul--31 Dec 1975 ** ERA-01:016800;EDB-76:042551; (25 Aug 1975) AD-A--014464

SUBJECT DESCRIPTORS: ABSORPTION SPECTRA: Q1,Q2,Q3,Q4;ELECTRONIC STRUCTURE;JAHN-TELLER EFFECT;MAGNETIC PROPERTIES;RHENIUM

<53>

<53> CONT.

FLUCRIDES: T1

ABSTRACT: Electronic absorption spectra of ReF_6 Quintet $d1$ are obtained for the pure crystal and UF_6 , MoF_6 , and WF_6 mixed crystals. The observed transition arises from an intra-configurational t_{2g} to t_{2g} promotion. The pure crystal is predicted to undergo a magnetic phase transition below ca. 1.5K. Pair spectra, two-molecule transitions, the ground state Jahn-Teller effect, linewidths, and vibrational assignments are discussed. The major mechanism for pair coupling and magnetic ordering is identified as superexchange through low lying delocalized charge transfer bands. These interactions are qualitatively discussed and energy localizing and delocalizing interactions are separated. (GRA)

EIP SUBJECT CATEGORY: EDB-400201;400702

<54>

Besmann, T.M.; Lindemer, T.B. (Oak Ridge National Lab., Tenn. (USA))

Thermodynamic assessment of the conversion of plutonium dioxide to plutonium monocarbide AIX-08:334259;EDB-77:137277; J. Nucl. Mater. (JNUMA) 67:(1):. 77-87 (1977)

SUBJECT DESCRIPTORS: CHEMICAL REACTION YIELD:

Q2;EQUILIBRIUM;EVAPORATION;FREE ENERGY;OXYGEN;PHASE
 DIAGRAMS;PLUTONIUM;PLUTONIUM CARBIDES: T2;PLUTONIUM DIOXIDE:
 T1;REDUCTION: Q1;S CODES;SOLID SOLUTIONS;VERY HIGH TEMPERATURE

ABSTRACT: The equilibrium thermodynamics of the carbothermic reduction of PuO_2 to $\text{PuC}_{0.88}$ are reassessed in light of new thermodynamic values for the Pu-C-O system. Analysis of the thermodynamic data indicates that the $\text{PuC}_{(x)}\text{O}_{(y)}$ phase can be successfully modeled by an ideal solid solution of $\text{PuC}_{0.88}$ and PuO . Equilibrium calculations show that oxygen-free monocarbide cannot be practically prepared by the direct carbothermic reduction of PuO_2 because of plutonium vaporization. These results imply that significant plutonium vaporization will occur in similarly reduced UO_2 - PuO_2 mixtures unless the actinides are initially present as a solid solution. Equilibrium calculations also suggest that $\text{PuC}_{0.88}$ can be successfully prepared by a two-step method: the carbothermic reduction of PuO_2 to $\text{PuC}_{1.5}$ followed by the hydrogen reduction of $\text{PuC}_{1.5}$ to $\text{PuC}_{0.88}$.

EIP SUBJECT CATEGORY: EDB-050700;360201;400702

INIS CATEGORY: B12

<55>

Besmann, T.M.; Lindemer, T.B., Oak Ridge National Lab., Tenn.
(USA)

In Thermodynamics of the conversion of plutonium dioxide to
plutonium monocarbide **

INS-76:019934; ERA-02:001052; EDB-76:087272; (Jul 1976)
ORNL/TM--5510

SUBJECT DESCRIPTORS: CHEMICAL PREPARATION:

Q2; EVAPORATION; HYDROGEN; PHASE DIAGRAMS: Q2, Q3; PLUTONIUM
CARBIDES: T2; PLUTONIUM DIOXIDE: T1; PLUTONIUM OXIDES:
T3; REDUCTION: Q1; THERMODYNAMICS

ABSTRACT: The present study contains an equilibrium
thermodynamic analysis of the Pu--C--O system and a
discussion from an equilibrium thermodynamic point of view of
the direct carbothermic reduction and two-step
carbothermic-hydrogen reduction of PuO_2 to PuC_{1-x} . Included are considerations of the partial pressures of
the various species in the Pu--C--O and Pu--C--H systems, the
process parameters required for conversion of the oxide to
the carbide, and the loss of plutonium due to vapor species.

EDB SUBJECT CATEGORY: EDB-360201; 050700
INIS CATEGORY: B23

<56>

Bhise, V.S.; Bonilla, C.F. ** Cooper, M.H. (ed.), American
Nuclear Society, Hinsdale, Ill.; American Inst. of Mining,
Metallurgical, and Petroleum Engineers, Inc., New York;
Energy Research and Development Administration, Washington,
D.C. (USA); American Nuclear Society, Pittsburgh, Pa.
Pittsburgh Section

Experimental vapor pressure and critical point of sodium In
Proceedings of the international conference on liquid
technology in energy production **

INS-77:015773; ERA-02:059296; EDB-77:138122; (Nov 1976)
CONF-760503--P2

SUBJECT DESCRIPTORS: CRITICAL PRESSURE: Q2; CRITICAL TEMPERATURE:
Q2; LMFER TYPE REACTORS: T1; REACTOR COOLING SYSTEMS:
Q1; SODIUM: T2; THERMODYNAMIC PROPERTIES; VAPOR PRESSURE: Q2

ABSTRACT: The pressure tube method employed previously to
determine the high temperature vapor pressure and the
critical point of cesium, rubidium, and potassium, was
improved and used on sodium. The critical pressure was
determined directly as 256.4 bars (3719 psia) with a standard
deviation of 0.09 percent. Vapor pressure was determined
experimentally from 1255 K (1800 F) up to the critical point.

<56>

<56> CONT.

The value selected for the critical temperature was 2503.3 K (4506 R), which is the lowest temperature at which the critical pressure was observed. Regression analysis showed that the data over the full range were slightly better fitted by $\ln P = 10.627840 - 11936.171/T + 0.03997216 \ln T$ (for P in Bars and T in K) than without the $\ln T$ term. This equation has a standard deviation of 1.1 percent from the data points.

EDB SUBJECT CATEGORY: EDB-210500;360104
INIS CATEGORY: E35;B22

<57>

Bishop, E.C.; Ray, W.E. ** Hill, M.R. (comp.), Oak Ridge National Lab., Tenn. (USA)

Westinghouse Advanced Reactors Division In Mechanical properties test data for structural materials. Quarterly progress report for period ending April 30, 1977 **
ERA-03:004356;EDB-78:004438; (Jun 1977) ORNL--5287

SUBJECT DESCRIPTORS: LMFBR TYPE REACTORS;MECHANICAL PROPERTIES:
Q1,Q2,Q3;REACTOR MATERIALS

ABSTRACT: Test programs are described involving basic specimen testing and tubular specimen testing of LMFBR type reactor materials. Sodium effects on mechanical properties of structural materials are also reported. (GHT)

EDB SUBJECT CATEGORY: EDB-360103;210500

<58>

Blander, M.; Curtiss, L.A.; Hsu, C.C.; Kucera, G.H.; Marr, J.J.; Papatheodorou, G.N.; Renner, T.A.; Saboungi, M.I.; Cafasso, F.A.; Eurriss, L., Argonne National Lab., Ill. (USA)

In Chemical Engineering Division physical inorganic chemistry. Annual report, July 1975--June 1976 **
ERA-03:007120;INS-78:000287;EDB-78:009786; (1976) ANL--76-101

SUBJECT DESCRIPTORS: ANL;RESEARCH PROGRAMS

ABSTRACT: The thermal conductivity of acetonitrile vapor has been measured at temperatures between 338 and 387 K and pressures between 100 and 1050 torr. The pressure dependence of the thermal conductivities indicates the presence of dimeric vapor species with an association constant given by $RT \ln K_{\text{sub}} (\text{atm}^{-1}) = 5002 - 21.15 T$. Molecular orbital studies of hydrogen-bonded methyl alcohol vapor complexes indicate that, the cyclic tetramer is especially stable. Preliminary calculations on trifluoroethanol indicate

<58> CCNT.

that its dimer is more stable than that of methanol. Molecular dynamics calculations have been utilized to calculate the second-order coefficient of conformational ionic solution (CIS) theory. The results indicate that the repulsive part of the pair potential most commonly used for calculations on molten salts is inaccurate and that calculation of the CIS coefficient provides a sensitive test of pair potentials. Laser-Raman spectra of molten solutions of YCl_3 in CsCl , KCl and LiCl were obtained and compared with spectra of solids of known structure. The spectra at low concentrations of YCl_3 (X/YCl_3 is less than 0.25) indicate an octahedrally coordinated YCl_6^{3-} species. Raman and infrared spectra of claudetite and arsenolite were measured and compared with the spectra of vitreous and liquid As_2O_3 , as well as with the As_4O_6 vapor molecules. The spectrum of claudetite has some resemblance to that of vitreous and liquid As_2O_3 . Raman spectra of arsenous oxide indicate that the dimeric As_2O_3 species is present. The Raman spectra of aluminum chloride vapor (AlCl_3 and Al_2Cl_6 vapor molecules), provided nine new bands not observed before and permitted a more complete assignment and analysis of the structure of the species to be made. The Raman spectra of In(I) , In(II) and In(III) chlorides indicate the presence of InCl , In(I)In(III)Cl_4 (In_2Cl_4), InCl_3 , and In_2Cl_6 species in the vapors. (JRD)

EDB SUBJECT CATEGORY: EDB-400100;360100

INIS CATEGORY: B11;B21

<59>

Blasse, G. (Rijksuniversiteit Utrecht (Netherlands). Fysisch Lab.)

Vibrational fine structure in the uranium emission of Mg_3TeO_6 and Y_6WO_{12}

AIX-07:269565;EDB-77:013708; Chem. Phys. Lett. (CHPLB) 41:(2):210-212 (15 Jul 1976)

ABSTRACT: The vibrational fine structure in the emission spectra of the uranium luminescence of Mg_3TeO_6 -U and Y_6WO_{12} -U at 4 K has been analyzed in terms of internal vibrational modes of the UO_6^{3+} octahedron and an external mode. It is concluded that uranium in these compounds is in six-coordination.

EDB SUBJECT CATEGORY: EDB-400702

INIS CATEGORY: A13

<60>

<60>

Bober, M.; Breitung, W.; Karow, H.U.; Schretzmann, K.
 {Kernforschungszentrum Karlsruhe (F.R. Germany). Inst. fuer
 Neutronenphysik und Reaktortechnik}

Investigation of fuel vapor pressure for fast breeder safety
 analysis AIX-07:270780; ERA-02:020150; EDB-77:035931;
 Atomwirtsch., Atomtech. (AWAKA) 21:(7):365-367 (Jul 1976)

SUBJECT DESCRIPTORS: EQUATIONS OF STATE; EXCURSIONS: Q1; FBR TYPE
 REACTORS: T1; KERNFORSCHUNGSZENTRUM KARLSRUHE; LASERS; MIXING
 RATIO; MIXTURES; PLUTONIUM DIOXIDE: T2; REACTOR SAFETY; SOLID
 FUELS; TRANSIENTS; URANIUM DIOXIDE: T3; VAPOR PRESSURE:
 Q2, Q3; VERY HIGH TEMPERATURE

ABSTRACT: The investigation of hypothetical over-prompt critical
 excursions for the safety analysis of fast reactors requires
 the knowledge of the equation of state of the oxidic fuel in
 the temperature region up to about 5,000 K. Conventional
 methods for measuring the vapour pressure, which were applied
 to oxide fuels up to 3,000 K, fail at higher temperatures.
 For some time now, experiments have been carried out in the
 Nuclear Research Centre, Karlsruhe, to measure fuel vapour
 pressures with open surface evaporation in the temperature
 region above 3,000 K, where the samples are locally heated
 without crucible by means of special laser techniques. The
 evaporation behaviour of
 ($U_{sub}(.80)Pu_{sub}(.20)O_{sub}(1.95...2.00)$) and of $UO_{sub}2$ and
 the connected vapour pressure build-up and its deviation from
 the sought equation of state were quantitatively investigated.

EDE SUBJECT CATEGORY: EDB-220900; 210500
 INIS CATEGORY: C52

<61>

Bode, D.D.; Wild, J.F.; Hulet, E.K. (California Univ., Livermore
 (USA). Lawrence Livermore Lab.)

Moessbauer studies of americium compounds
 AIX-07:270229; EDB-77:031457; J. Inorg. Nucl. Chem. (JINCA)
 38:(7):1291-1297 (1976)

SUBJECT DESCRIPTORS: AMERICIUM 243: T1; AMERICIUM
 CARBONATES; AMERICIUM COMPOUNDS: M2; AMERICIUM
 FLUORIDES; AMERICIUM OXIDES; ELECTRONIC STRUCTURE; GAMMA
 RADIATION; HYPERFINE STRUCTURE; ISOMER SHIFT; LINE
 WIDTHS; MOESSBAUER EFFECT: Q1; PLUTONIUM 243; STRUCTURAL
 CHEMICAL ANALYSIS: Q2; VALENCE

ABSTRACT: Moessbauer studies of several compounds of Am were
 made with ^{243}Pu as a source and ^{243}Am as an
 absorber. Isomer shifts, line widths, and resonance

<61> CONT.

intensities were measured with various sources and absorbers for recoilless emission of the 84-keV γ -ray of ^{243}Am . The pentavalent Am compounds KAmO_2CO_3 and KAmO_2F_2 gave five-peak absorption spectra attributable to electric hyperfine interaction. A complex spectrum, interpreted as a result of magnetic hyperfine interaction, as also observed for the system $\text{PuFe}_2/\text{AmO}_2$. The changes in isomer shifts with the Am oxidation state are attributed to increases in the electron density in the region of the nucleus caused by removal of 5f electrons, and consequently, an increase in the effective Coulombic nuclear attraction for the outer s and p electrons.

EDB SUBJECT CATEGORY: EDB-400201
INIS CATEGORY: B13

<62>

Boivineau, J.C.

In X-ray study of the plutonium--oxygen phase diagram at ambient temperatures up to 1100°C **
ERA-02:017802;EDB-77:026211; J. Nucl. Mater. (JNUMA) 60::31-38 (1976) LA-tr--76-30

SUBJECT DESCRIPTORS: PHASE DIAGRAMS: Q1; PLUTONIUM OXIDES: T1; VERY HIGH TEMPERATURE; X-RAY DIFFRACTION

ABSTRACT: A monochromatic x-ray diffraction technique was used to establish several new features in the Pu--O phase diagram, such as a wide stability region for the α phase, a narrow ($\alpha + \gamma$) phase region, a eutectoidal point at 670°C for an O/Pu ratio of 1.78 and several γ -type phases. Unfortunately, because of the complexity of the phases and the existence of metastable equilibria, details of the diagram could not be established for low oxygen contents. 4 tables, 5 figs.

EDB SUBJECT CATEGORY: EDB-360202

<63>

Borgstedt, H.U.; Frees, G.; Schneider, H. (Kernforschungszentrum Karlsruhe (Germany, F.R.). Inst. fuer Material- und Festkoerperforschung)

Characterisation of the chemical activities of non-metals dissolved in liquid alkali metals AIX-08:334265;EDB-77:141001; Ber. Bunsenges. Phys. Chem. (BBPCA) 80:(8):740-745 (Aug 1976)

SUBJECT DESCRIPTORS: ALKALI METALS; CARBON; CHEMICAL ACTIVATION: Q1; DIAGRAMS; ELECTROCHEMISTRY: T; ELECTROLYTIC

<63>

<63> CONT.

CELLS;ELECTROMOTIVE FORCE;EQUILIBRIUM;LIQUID METALS;OXYGEN:
T1;SOLUTIONS;TEMPERATURE DEPENDENCE;THERMODYNAMICS;THORIUM
OXIDES;TIME DEPENDENCE;VANADIUM;VANADIUM OXIDES;YTTRIUM OXIDES

ABSTRACT: The oxygen activity in molten alkali metals can be measured by the use of solid electrolyte emf cells. The emf cells use yttria doped thorium as the electrolyte and a gaseous oxygen atmosphere or a solid oxide as the reference system. Unfortunately the emf output of the cells differs from values to be expected from theory. To receive the emf of a well defined oxygen potential in the liquid alkali metals we have used exchange reactions like $\text{Na}_2\text{O} + \text{V} = \text{VO} + 2\text{Na}$, $\text{K}_2\text{O} + \text{V} = \text{VO} + 2\text{K}$. In both cases we have got the same final values of the emf corresponding to the free energy of formation of VO. The analytical determination of the oxygen fixed by the vanadium allows to relate the initial emf values to a known oxygen concentration. The oxygen exchange calibration method enables the emf cells to measure oxygen activities resp. concentrations in liquid sodium or potassium. The exchange of carbon between solid alloys and liquid sodium can be applied to measure carbon activities. At temperatures of 700 to 750°C the exchange of carbon between thin foils of 18/8 CrNi steel and sodium reaches equilibrium after 120 hours. By the application of this reaction and analyses of the final carbon concentrations in the steel specimens it could be demonstrated that in sodium circuits constructed of austenitic steels the carbon contents are on a level of lower than one w-ppm. The aim of the work is to enable the analytical techniques to measure these rather small carbon contents in alkali metal circuits.

EDB SUBJECT CATEGORY: EDB-400201

INIS CATEGORY: F12

<64>

Borgstedt, H.U.; Frees, G.; Schneider, H., Kernforschungszentrum
Karlsruhe (Germany, F.R.). Inst. fuer Material- und
Festkoerperforschung

In Characterisation of the chemical activities of non-metals
dissolved in liquid alkali metals **

AIX-07:272374;EDB-77:056343; (1976) AED-Conf--76-087-006

SUBJECT DESCRIPTORS: ALKALI METALS: T3;CALIBRATION:

T1;CARBON;CERAMICS;CHEMICAL COMPOSITION;CHEMICAL REACTION
KINETICS;CORROSION;DIAGRAMS;DISSOLUTION;ELECTROLYTES;ELECTRO-
MOTIVE FORCE;EQUILIBRIUM;GETTERING;LIQUID METALS;MONITORING:
Q1,Q2,Q3;OXIDES;OXYGEN: T2;THORIUM OXIDES;TIME
DEPENDENCE;YTTRIUM OXIDES

ABSTRACT: The oxygen activity in molten alkali metals can be

<64> CONT.

measured by the use of solid electrolyte emf cells. The emf cells use yttria doped thoria as the electrolyte and a gaseous oxygen atmosphere or a solid oxide as the reference system. Unfortunately the emf output of the cells differs from values to be expected from theory. To receive the emf of a well defined oxygen potential in the liquid alkali metals we have used exchange reactions like $\text{Na}_2\text{O} + \text{V} = \text{VO} + 2 \text{Na}$ and $\text{K}_2\text{O} + \text{V} = \text{VO} + 2 \text{K}$. In both cases we have got the same final values of the emf corresponding to the free energy of formation of VO. The analytical determination of the oxygen fixed by the vanadium allows to relate the initial emf values to a known oxygen concentration. The oxygen exchange calibration method enables the emf cells to measure oxygen activities or concentrations in liquid sodium or potassium. The exchange of carbon between solid alloys and liquid sodium can be applied to measure carbon activities. At temperatures of 700 to 750°C the exchange of carbon between thin foils of 18/8 CrNi steel and sodium reaches equilibrium after 120 hours. By the application of this reaction and analyses of the final carbon concentrations in the steel specimens it could be demonstrated that in sodium circuits constructed of austenitic steels the carbon contents are on a level of lower than one w-ppm. The aim of the work is to enable the analytical techniques to measure these rather small carbon contents in alkali metal circuits.

EDB SUBJECT CATEGORY: EDB-400400

INIS CATEGORY: B12

<65>

Borsese, A.; Borzone, G.; Calabretta, A.; Capelli, R.; Ferro, R.
 ** Lundin, C.E. (ed.)

Advances on thermochemistry of rare earths alloys In Proceedings of the 12th rare earth research conference. Vol. II **
 ERA-02:060383; INS-77:017515; EDB-77:140547; (1976) Univ. of Denver

SUBJECT DESCRIPTORS: FORMATION HEAT: Q1; RARE EARTH ALLOYS:
 T1; REVIEWS

ABSTRACT: The values of the heats of formation so far obtained on alloys of rare earths with V group elements are summarized. 5 figures, 17 references. (GHT)

EDB SUBJECT CATEGORY: EDB-360104

INIS CATEGORY: B22

<66>

<66>

Bougon, R.; Charpin, P.; Desmoulin, J.P.; Malm, J.G.

Structural study and properties of the alkali metal, nitrosyl,
and ammonium hepta- and octafluorouranates(VI)
INS-77:005831; EDB-77:044532; Inorg. Chem. (INOCA)
15:(10):2532-2540 (Oct 1976)

SUBJECT DESCRIPTORS: AMMONIUM COMPOUNDS: T8; BOND LENGTHS; CESIUM
COMPOUNDS: T4; CHEMICAL PROPERTIES:
Q1, Q2, Q3, Q4, Q5, Q6, Q7, Q8, Q9; CHEMICAL REACTION
KINETICS; FLUORINE COMPOUNDS: T6; HIGH TEMPERATURE; INFRARED
SPECTRA; MEDIUM TEMPERATURE; NITROSO COMPOUNDS: T7; NUCLEAR
MAGNETIC RESONANCE; POTASSIUM COMPOUNDS: T2; PYROLYSIS; RAMAN
SPECTRA; RUBIDIUM COMPOUNDS: T3; SODIUM COMPOUNDS:
T1; STRUCTURAL CHEMICAL ANALYSIS:
Q1, Q2, Q3, Q4, Q5, Q6, Q7, Q8, Q9; SYNTHESIS:
Q1, Q2, Q3, Q4, Q5, Q6, Q7, Q8, Q9; TEMPERATURE EFFECTS; URANIUM
COMPOUNDS: T5; URANIUM FLUORIDES: T9; X-RAY DIFFRACTION

ABSTRACT: The thermal decomposition of the
heptafluorouranates(VI) of the alkali metals is shown to take
place in two steps. The first step gives the
octafluorouranates(VI) and UF_6 , and the decomposition
rate is noticeable at temperatures above 100, 130, 150, and
210°C for the Na, K, Rb, and Cs salts, respectively.
The second step for Na_2UF_8 yields pure NaF and
 UF_6 above 300°C, whereas the decomposition
temperatures for the K, Rb, and Cs salts are above 300, 350,
and 400°C, respectively. Depending on the decomposition
conditions, F_2 and M_2UF_7 ($M = K, Rb, Cs$) or
 F_2 , UF_6 , and M_3UF_8 are formed. The
heptafluorouranates(VI) of all the cations studied, except
for ammonium, were shown to exhibit dimorphism. The
parameters of their cubic form were obtained and are as
follows: KUF_7 ; $a = 5.22$ Å; $RbUF_7$; $a = 5.385$ Å;
 $CsUF_7$; $a = 5.517$ Å; $NOUF_7$; $a = 5.334$ Å; NH_4UF_7 ; $a = 5.393$ Å; $NaUF_7$ (fccub), $a = 8.511$ Å, $Z = 4$. The x-ray pattern of the low-symmetry form of $CsUF_7$
just below the solid transition temperature (15 ± 1°C) was indexed with a tetragonal cell where $a = 5.50$ Å and
 $c = 5.37$ Å. The x-ray diagrams of the low symmetry form of
the other MUF_7 salts were not indexed, whereas those of
the octafluorouranates were indexed with orthorhombic cells.
The vibrational spectra of the hepta- and octafluorouranates
were found to be very dependent on the temperature, and for
the same temperature on the cation size. Some trends observed
in this study, like the thermal decomposition temperatures or
the relative symmetries, are thought to arise from the
differences in the cation--anion interaction. This
interaction is stronger with smaller cations, which results

<66> CONT.

in more distorted anions, less ionic U--F bonds, and paradoxically less stable complexes.

EEB SUBJECT CATEGORY: EDB-400702;400104

INIS CATEGORY: B13;B11

<67>

Bourgeois, M.; Le Bouhellec, J.; Miquel, P.; Sauteron, J.; Boundry, J.C., Kernforschungszentrum Karlsruhe (Germany, F.R.). Internationales Buero/Uebersetzergruppe

In Reprocessing of fast breeder fuels in France **

AIX-07:272604;ERA-02:026001;EDB-77:053617; (Jun 1976)

KFK-tr--471

ABSTRACT: The CEA has for a long time been making considerable efforts in the field of the construction of sodium-cooled fast breeders, in particular regarding the reprocessing of spent fuels from this construction: operation of the plant AT\$sub 1\$ from the Hague, in which the fuel irradiated in Rapsodie is reprocessed, laboratory investigations and chemical engineering tests in Fontenay-aux-Roses etc. Numerous problems are yet to be solved for the reprocessing to be sufficiently matured for industry: high specific activity, large release of heat, insoluble residues, handling of large quantities of plutonium, waste, treatment etc. An extensive research and development programme is thus being carried out so that these problems can be solved in time (laboratory, chemical engineering, plant AT\$sub 1\$ pilot plant Marcoule, testing of prototypes true-to-scale, layout studies). The experience gained and the continuous investigations in the following fields are described: the work on the fuel prior to reprocessing (deactivation, handling, sodium extraction, transport), the wet-chemical reprocessing (initial treatment, solvation and preparation of the solution, solvent extractions), the dry reprocessing (vaporization of the hexafluorides), the particular problems of wastes (solid, liquid, gaseous). To conclude, an explanation is given as to why it is still too early to work out a strategy for the reprocessing on industrial scale for this reactor construction.

EEB SUBJECT CATEGORY: EDB-050800

INIS CATEGORY: B16

<68>

<68>

Bouteillan, J.; Barbier, M.J. (Ecole Nationale Supérieure
d'Electrochimie et d'Electrometallurgie, 38 - Grenoble
(France))

Study of the Ni-Ni (II) equilibrium exchange, by radiotracer
techniques in fused LiCl-KCl eutectic
AIX-08:298231; EDB-77:062577; Rev. Int. Hautes Temp. Refract.
(RIHTA) 13: (1):43-47 (Jul 1976)

SUBJECT DESCRIPTORS: CATIONS; CHEMICAL REACTION KINETICS:
Q1; ELECTRODES; ELECTROLYTIC CELLS; EUTECTICS; HIGH
TEMPERATURE; ISOTOPIC EXCHANGE: T1; LITHIUM CHLORIDES; NICKEL
65; NICKEL COMPOUNDS; POTASSIUM CHLORIDES; TRACER TECHNIQUES

ABSTRACT: Radiotracer techniques were applied to fused LiCl-KCl
eutectic to study the equilibrium kinetics of the Ni-Ni (II)
exchange. A relation was established which gives the time
dependent theoretical variation of the radioactivity of the
solution, containing labelled Ni(II) species, in contact
with an inactive nickel metallic electrode; theoretical
curves were compared with the experimental one. This
comparison corroborates the velocity of the Ni-Ni (II)
electrochemical system previously observed by means of
electrical methods; it induces to assume a number of nickel
atoms involved in the isotopic interfacial exchange greater
than the quantity calculated on the basis of a surface
monolayer.

EDB SUBJECT CATEGORY: EDB-400203
INIS CATEGORY: B12

<69>

Howersox, D.F., Los Alamos Scientific Lab., N.Mex. (USA)

In Review of partitioning proposals for spent nuclear fuels **
INS-77:001898; ERA-02:015896; EDB-77:029158; (Jul 1976)
LA--6442

SUBJECT DESCRIPTORS: COMPARATIVE EVALUATIONS; FISSION
PRODUCTS; FLUORIDE VOLATILITY PROCESS; PUREX
PROCESS; PYROMETALLURGY; RECOVERY; REPROCESSING: Q1; SPENT FUELS:
T1

ABSTRACT: The initial phase of a study about recovery of
valuable fission products from spent nuclear fuels has been
to review various partitioning proposals. This report briefly
describes the aqueous Purex process, the salt transport
process, melt refining, fluoride volatility process, and
gravimetric separations. All these processes appear to be
possible technically, but further research will be necessary
to determine which are most feasible. This review includes

<69> CONT.

general recommendations for experimental research and development of several partitioning options.

EDB SUBJECT CATEGORY: EDB-050800
INIS CATEGORY: B16

<70>

Brandtberg, S.

In Method for recovering uranium from residual solution **
AIX-08:298504; ERA-02:032139; EDB-77:072669; (21 Jun 1976)

SUBJECT DESCRIPTORS: AMMONIA; AQUEOUS SOLUTIONS; CALCIUM CARBONATES; CARBON DIOXIDE; FLUORIDE VOLATILITY PROCESS; FUEL FABRICATION PLANTS: M2; RECOVERY: Q1, Q2; REPROCESSING; URANIUM: M1; URANIUM HEXAFLUORIDE

ABSTRACT: A method is given for recovering uranium from a residual solution provided by an installation for the manufacture of a nuclear fuel from uranium hexafluoride in the presence of ammonia by eliminating the fluorides of the residual solution. The method is characterized by the steps of precipitating the fluorides of the residual solution through adding a powdered alkaline earth carbonate, and of recovering uranium from the residual solution thus free from fluorides.

EDB SUBJECT CATEGORY: EDB-052001; 050700
INIS CATEGORY: B16

<71>

Braunstein, J.; Bronstein, H.R.; Cantor, S.; Heatherly, D.; Vallet, C.E., Oak Ridge National Lab., Tenn. (USA)

In Molten carbonate fuel cell research at ORNL **
ERA-02:049055; EDB-77:106734; (May 1977) ORNL/TM--5886

SUBJECT DESCRIPTORS: ALUMINIUM OXIDES; BORON NITRIDES; CARBONATES; COMPATIBILITY; HIGH-TEMPERATURE FUEL CELLS: T1; MASS TRANSFER; MATERIALS; MOLTEN SALTS: T2; RESEARCH PROGRAMS: Q1; THERMODYNAMIC PROPERTIES: Q2

ABSTRACT: The activities at ORNL during the period July 1976 to February 1977 on the molten carbonate fuel cell program, funded by the ERDA Division of Conservation Research and Technology, are summarized. This period marks the initiation of molten carbonate fuel cell research at ORNL, making use of the extensive background of expertise and facilities in molten salt research. The activities described include a literature survey on molten carbonates, design, acquisition

<71>

<71> CONT.

and installation of apparatus for experimental studies of molten carbonates, initial experiments on materials compatibility with molten carbonates, electrolysis experiments for the determination of transference numbers, and theoretical studies of transport behavior and the coupling of mass flows in molten carbonate mixtures. Significant accomplishments were the theoretical prediction of a possibly appreciable change in the alkali ion ratio at molten carbonate fuel cell electrodes, operated at high current densities, as a result of mobility differences of the alkali ions; design, construction and assembly of an electrolysis cell, and initiation of measurements of composition profiles in mixed alkali carbonate electrolytes; initiation of differential scanning calorimetry of pure alkali carbonates for quantitative measurement of transition enthalpies, eventually leading to new, more reliable values of the enthalpies and free energies of formation of the pure and mixed carbonates.

EDB SUBJECT CATEGORY: EDB-300501;300503;300505

<72>

Breitung, W., Kernforschungszentrum Karlsruhe (Germany, F.R.).
Inst. fuer Neutronenphysik und Reaktortechnik

In Calculation of vapor pressures of oxide fuels up to 5000 K in equilibrium and non-equilibrium evaporation **
ERA-02:000789;EDB-76:086430; (Jun 1975) EURPNE--1283

SUBJECT DESCRIPTORS: EXCURSIONS: Q1;FUEL PINS;LMPBR TYPE
REACTORS: T1;MELTDOWN;PLUTONIUM OXIDES: T3;ULTRAHIGH
TEMPERATURE;URANIUM DIOXIDE: T2;URANIUM OXIDES: T4;VAPOR
PRESSURE: Q2,Q3,Q4;VERY HIGH TEMPERATURE

ABSTRACT: The evaporation kinetics of multi-component systems is studied with UO_2 as the example. The evaporation, which is generally incongruent, implies that two opposing types of steady-state evaporation must be distinguished: equilibrium evaporation and "forced congruent" evaporation. In the first case, a homogeneous composition of the material is maintained despite incongruent evaporation from the surface. This is ensured by sufficiently large diffusion flows towards the evaporating surface within the substance. By contrast, forced congruent evaporation occurring in the case of intensive evaporation gives rise to concentration gradients in the material, which are due to the high evaporation rates. The evaporating surface changes its composition such that congruent evaporation of the material takes place. The two types of evaporation indicated entail different vapor pressures. In some prompt critical reactor incidents forced congruent evaporation must be anticipated. The calculation of

<72> CONT.

the vapor pressures of UO_2 and $(\text{U}, \text{Pu})\text{O}_2$ for both types of evaporation up to temperature of 5000 K is presented. The calculating procedures are based on the method of Rand and Markin (1967) incorporating the recent thermodynamic data. The agreement between the measured and calculated total pressures is good for the ranges of temperature and stoichiometry for which experimental results are available. This supports the results calculated for higher temperature ranges.

EDE SUBJECT CATEGORY: EDB-220900;210500;360204;400702

<73>

Breitung, W., Kernforschungszentrum Karlsruhe (Germany, F.R.).
Inst. fuer Neutronenphysik und Reaktortechnik;
Kernforschungszentrum Karlsruhe (Germany, F.R.). Projekt
Schneller Brueter

In Calculation of vapor pressures of oxide fuels up to 5,000 K
for equilibrium and nonequilibrium evaporation **
AIX-07:231065;ERA-02:003044;EDB-76:091352; (Jun 1975)
RFR--2091

SUBJECT DESCRIPTORS: BINARY MIXTURES;CHEMICAL
COMPOSITION;COMPARATIVE EVALUATIONS;COMPUTER
CALCULATIONS;DATA;DIAGRAMS;ENTHALPY;EQUILIBRIUM;EVAPORATION:
Q1;NUCLEAR FUELS: T1;PLUTONIUM DIOXIDE: T2;ULTRAHIGH
TEMPERATURE;URANIUM DIOXIDE: T3;VAPOR PRESSURE: Q2,Q3

ABSTRACT: The evaporation kinetics of multicomponent systems is studied with UO_2 as the example. The evaporation, which is generally incongruent, implies that two opposing types of steady-state evaporation must be distinguished: equilibrium evaporation and forced congruent evaporation. The two types of evaporation indicated entail different vapor pressures. In some prompt critical reactor incidents forced congruent evaporation must be anticipated. Vapor pressures of UO_2 and $(\text{U}, \text{Pu})\text{O}_2$ are calculated for both types of evaporation up to a temperature of 5,000 K. The calculating procedures are based on the method of Rand and Markin (1967) incorporating recent thermodynamic data. The agreement between the measured and calculated total pressures is good for the ranges of temperature and stoichiometry for which experimental results are available. This supports the results calculated for higher temperature ranges.

EDE SUBJECT CATEGORY: EDB-220300;360202
INIS CATEGORY: E23

<74>

<74>

Breitung, W., Kernforschungszentrum Karlsruhe (Germany, F.R.).
 Inst. fuer Neutronenphysik und Reaktortechnik;
 Kernforschungszentrum Karlsruhe (Germany, F.R.). Projekt
 Schneller Brueter

In Calculation of the oxygen potential of $(U,Pu)O_{2-x}$ with
 defect models and measured oxygen potential data in the
 temperature range from 1,000 to 1,700 K **
 AIX-08:297306;EDB-77:072618; (Oct 1976) KFK--2363

SUBJECT DESCRIPTORS: BINARY MIXTURES;CHEMICAL
 COMPOSITION;ELECTROCHEMISTRY;FORMATION FREE
 ENTHALPY;INTERSTITIALS: Q1,Q2;OXYGEN;PLUTONIUM DIOXIDE:
 T1;URANIUM DIOXIDE: T2;VACANCIES: Q1,Q2;VERY HIGH TEMPERATURE

ABSTRACT: The dominating defects in $(U,Pu)O_{2-x}$ are oxygen vacancies and
 oxygen interstitials. It is assumed that an oxygen vacancy
 and an oxygen interstitial in the (U,Pu) -mixed oxide, is bound
 to a Pu_{2O_3} - and U_{2O_5} -complex respectively. The model used to describe the oxygen potential
 $\Delta G(C_{O_2})$ is based on the equilibrium between these
 defects and gaseous oxygen. The equilibrium constants needed,
 were evaluated from measured $\Delta G(O_{2O})$ -data. The
 calculated oxygen potentials are given as function of x,y and
 T . Especially the model provides $\Delta G(O_{2O})$ -data of
 nearly stoichiometric (U,Pu) mixed oxides, for which no
 measured data are available. For the free energy of formation
 of oxygen Frenkel defects $\Delta G(FO)$ in (U,Pu) mixed
 oxides, values between 2.0 and 2.3 eV were determined for $y =$
 0.3 ... 0.1.

EDB SUBJECT CATEGORY: EDB-050700;360202
 INIS CATEGORY: A13

<75>

Presesti, A.M.; Bresesti, M.; Cuypers, M.; D'Adamo, D.; Lezzoli,
 L.; Pannella, R.; Giuriato, A.; Paoletti Gualandi, M.;
 Peroni, P., Commission of the European Communities, Ispra
 (Italy). Joint Research Centre

In Investigations on radioactive fission product correlations:
 gamma spectrometry measurements on spent fuel assemblies
 discharged from the Trino Vercellese reactor at the end of
 the 2nd irradiation cycle ** ERA-01:019092;EDB-76:058135;
 (Jun 1975) EUR--5334

SUBJECT DESCRIPTORS: B CODES;BURNUP;CESIUM 134;CESIUM
 137;CORRELATIONS;EUROPIUM 154;FISSION PRODUCTS;FUEL
 ASSEMBLIES;FUEL ELEMENTS: Q1;GAMMA FUEL
 SCANNING;PLUTONIUM;RADIOMETRIC ANALYSIS;SELNI REACTOR:

<75> CONT.

T1; SPENT FUEL ELEMENTS

ABSTRACT: In the framework of a cooperation program between ENEL and EURATOM, gamma spectrometry measurements were performed on 12 fuel assemblies of the Trino Vercellese PWR to study the correlations existing between radioactive fission products and the burnup and plutonium content. A mechanical and collimation system permitted measurements to be performed at different levels on the four corners of the fuel assemblies in the spent fuel pool of the Trino Vercellese plant. Gamma spectrometry measurements were also carried out on solutions of the Trino Vercellese fuel, taken at the input of the Eurcochemic reprocessing plant. The gamma ray activities of the fission products were correlated with values of burnup and plutonium content generated by BURSQUID, a two-dimensional diffusion burnup code, and with values of plutonium content determined by means of chemical analyses at the input of the reprocessing plant. A proportionality has been observed between Cs 137 activity and burnup and between Cs 134/Cs 137 activity ratio and Pu/U mass ratio. This result indicates the possibility of non-destructive measurements of burnup and plutonium content distributions in PWR fuel assemblies. The experimental results are supported by theoretical predictions carried out by means of computer code calculations.

EDB SUBJECT CATEGORY: EDB-210200

INIS CATEGORY: E32

<76>

Broadhead, K.G.; Heady, H.H., Bureau of Mines, Reno, Nev. (USA).
Reno Metallurgy Research Center

In Radiotracer applications in electrometallurgical processing

** EDB-77:120034; (Sep 1962) BM-RI--6195

SUBJECT DESCRIPTORS: ELECTROMETALLURGY:

T1,Q2;RADIOISOTOPES;TRACER TECHNIQUES: Q1

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-360101

<77>

<77>

Broers, G.H.J.; Van Ballegoy, H.J.J.

Phase equilibria in Li--Na--K carbonate/aluminate systems In
 Third international symposium on fuel cells **
 ERA-01:019721;EDB-76:053277; (1969) Presses Academiques
 Europeennes

SUBJECT DESCRIPTORS: ALUMINATES: T3; CARBONATES: T2; ELECTROLYTES:
 Q1; HIGH-TEMPERATURE FUEL CELLS: T1; LITHIUM COMPOUNDS:
 T4; MOLYBDENUM SALTS; PHASE DIAGRAMS: Q2, Q3, Q4, Q5, Q6; POTASSIUM
 COMPOUNDS: T6; SODIUM COMPOUNDS: T5

ABSTRACT: Molten carbonate fuel cells on basis of liquid
 carbonate/solid aluminate paste electrolytes are far more
 stable in long term operation than similar cells on basis of
 carbonate/magnesium oxide pastes. Phase equilibria between
 the carbonate melt and the (insoluble) solid aluminates of
 Li, Na and K have been studied both theoretically and
 experimentally, by means of thermal and electrical
 conductance analyses.

EDB SUBJECT CATEGORY: EDB-300503;360602

<78>

Bron, J. (Natal Univ., Durban (South Africa). Dept. of Chemistry)

Isotope effect on the thermodynamic quantities of gaseous
 uranium hexafluoride AIX-07:257165;EDB-77:007647; Can. J.
 Chem. (CJCHA) 54: (1):160-165 (1 Jan 1976)

SUBJECT DESCRIPTORS: ENTHALPY: Q2; ENTROPY: Q2; FREE ENTHALPY:
 Q2; GASES; ISOTOPE EFFECTS: Q1; MOLECULAR MODELS; PARTITION
 FUNCTIONS; QUANTUM MECHANICS; SPECIFIC HEAT: Q2; URANIUM 235:
 T1; URANIUM 238; URANIUM HEXAFLUORIDE: T2

ABSTRACT: Isotopic effects on thermodynamic properties of
 gaseous uranium hexafluoride are calculated from the
 logarithm of the ratio of the partition function. Changes in
 the molar quantities Gibbs' free energy, enthalpy, specific
 heat and entropy on substituting ^{235}U for ^{238}U in
 UF_6 are tabulated as functions of temperature. It is
 concluded that the approximation known as the first quantum
 correction has little quantitative value.

EDB SUBJECT CATEGORY: EDB-400702;400202
 INIS CATEGORY: B12

<79>

Brown, C.H. Jr.; Hightower, J.R. Jr.; Klein, J.A., Oak Ridge National Lab., Tenn. (USA)

In Measurement of mass transfer coefficients in a mechanically agitated, nondispersing contactor operating with a molten mixture of $\text{LiF--BeF}_2\text{--ThF}_4$ and molten bismuth **
INS-77:003220; ERA-02:016585; EDB-77:023552; (Nov 1976)
ORNL--5143

SUBJECT DESCRIPTORS: BERYLLIUM FLUORIDES; BISMUTH; BREEDER REACTORS; DATA; EXTRACTION APPARATUSES: T2; HIGH TEMPERATURE; LIQUID METALS; LITHIUM FLUORIDES; MASS TRANSFER: Q2, Q3; MOLTEN SALT REACTORS: T1; MOLTEN SALTS; REPROCESSING: T3, Q1; SPENT FUELS; THORIUM FLUORIDES

ABSTRACT: A mechanically agitated, nondispersing contactor in which molten fluoride salt and molten bismuth phases were contacted has been built and operated. The mass transfer performance of the contactor was evaluated over a range of agitator speeds under conditions in which the major resistance to mass transfer was in the salt phase. The measured mass transfer rates were compared with rates predicted by literature correlations. The equipment necessary to contain the salt and bismuth at approximately 600°C is described along with the complete set of experimental data obtained during operation. 11 figures, 12 tables.

EDB SUBJECT CATEGORY: EDB-210500; 050800
INIS CATEGORY: E35; B16

<80>

Brown, D.; De Paoli, G.; Whittaker, B. (UKAEA Research Group, Harwell. Atomic Energy Research Establishment)

Conversion of protactinium monocarbide to the penta- and tetra-halides AIX-07:274902; EDB-77:019525; J. Chem. Soc. (London), Dalton Trans. (JCOTB) (14):1336-1338 (1976)

SUBJECT DESCRIPTORS: CHEMICAL PREPARATION: Q1, Q2, Q3; CHEMICAL REACTIONS: Q4; INFRARED SPECTRA; MOLECULAR STRUCTURE; PROTACTINIUM BROMIDES: T1; PROTACTINIUM CARBIDES: T4; PROTACTINIUM CHLORIDES: T2; PROTACTINIUM IODIDES: T3; RAMAN SPECTRA; STRUCTURAL CHEMICAL ANALYSIS; X-RAY DIFFRACTION

ABSTRACT: Protactinium monocarbide has been prepared by carbothermic reduction of the pentaoxide. It reacts with iodine (400°C), bromine (350°C), and thionyl chloride (200°C) in evacuated sealed vessels to yield the appropriate pentahalide. The tetraiodide, PaI_4 , is obtained by interaction of the monocarbide with either protactinium pentaiodide (600°C) or mercury(II) iodide

<80>

<80> CONT.

(500⁰C) in vacuo. X-ray powder-diffraction results and vibrational spectral data are reported for the halides.

EDB SUBJECT CATEGORY: EDB-400702

INIS CATEGORY: B13

<81>

Brown, D.; De Paoli, G.; Whittaker, B., UKAEA Research Group, Harwell. Atomic Energy Research Establishment

In Some chemical reactions of uranium hexachloride and the chlorination of uranium tetrachloride **

AIX-07:248008;EDB-77:019508; (Jan 1976) AERE-R--8260

SUBJECT DESCRIPTORS: ABSORPTION SPECTRA;CHEMICAL PREPARATION;CHEMICAL REACTIONS: Q1;CHLORINATION;ELECTRONIC STRUCTURE;INFRARED SPECTRA;LATTICE PARAMETERS: Q1;MOLECULAR STRUCTURE;STRUCTURAL CHEMICAL ANALYSIS;TRICLINIC LATTICES;URANIUM CHLORIDES: T1;URANIUM COMPLEXES

ABSTRACT: Uranium hexachloride has been prepared by chlorination of the tetrachloride and its chemical properties investigated. It decomposes slowly in anhydrous, oxygen-free carbon tetrachloride to yield $\beta\text{-UCl}_5$ which exhibits triclinic symmetry with $a = 7.09$, $b = 9.66$, $c = 6.36$ angstroms $\alpha = 88.5^\circ$, $\beta = 117.6^\circ$, $\gamma = 108.5^\circ$ in space group $P1$. Infrared spectra and solid state optical transmission spectra are reported for $\beta\text{-UCl}_5$, and electronic transitions are assigned from the γ_7 ground state. Attempts to prepare uranium hexachloride complexes with donor ligands have yielded pentavalent complexes of the types $\text{UCl}_5 \cdot 2\text{L}$ (L = α , α -dipyridyl, benzanthrone and dimethylformamide) and $\text{UCl}_5 \cdot \text{L}$ (L = triphenylphosphine oxide), and the tetravalent complex $\text{UCl}_4 \cdot 2\text{L}$ (L = anthrone). The hexachloride is immediately reduced to soluble pentavalent species on the addition of anhydrous methyl cyanide; evaporation of the resulting solution yields $\text{UCl}_5 \cdot 2\text{CH}_3\text{CN}$. Electronic spectra are reported for the solid and for its solutions in CH_3CN .

EDB SUBJECT CATEGORY: EDB-400702

INIS CATEGORY: B13

<82>

Brown, D.; Hurtgen, C. (UKAEA Research Group, Harwell. Atomic Energy Research Establishment); Fuger, J. (Liege Univ. (Belgium))

Preparation and some properties of actinide(III) and lanthanide(III) oxide halides, and the enthalpies of formation of $\text{PuBr}_3(\text{s})$ and $\text{PuI}_3(\text{s})$
 AIX-08:346226;EDB-78:028028; Rev. Chim. Miner. (RVCMA) 14:(2):189-198 (1977)

SUBJECT DESCRIPTORS: CHEMICAL PREPARATION:

Q1,Q2,Q3,Q4,Q5,Q6,Q7,Q8;DYSPROSIUM COMPOUNDS:
 T6;ENTHALPY;ERBIUM COMPOUNDS: T8;FORMATION HEAT:
 Q9,Q10;GADOLINIUM COMPOUNDS: T5;LANTHANUM COMPOUNDS:
 T3;NEODYMIUM COMPOUNDS: T4;NEPTUNIUM COMPOUNDS:
 T1;OXYBROMIDES;OXYCHLORIDES;OXYIODIDES;PLUTONIUM BROMIDES:
 T9;PLUTONIUM COMPOUNDS: T2;PLUTONIUM IODIDES: T10;TERBIUM
 COMPOUNDS: T7

ABSTRACT: Reactions between actinide (Np, Pu) and lanthanide (La, Nd, Gd, Dy, Tb, Er) trihalides and antimony sesquioxide at 400-700 deg C in vacuo yield the corresponding oxide halides, MOX (X=variously Cl, Br, I). Attempts to prepare UOCl and UOI by similar reactions have been unsuccessful, the identified products being uranium dioxide and unchanged trihalide. Infrared spectral data are reported for the oxide halides together with unit cell dimensions for ErOI . Enthalpies of formation determined for $\text{PuBr}_3(\text{s})$ and $\text{PuI}_3(\text{s})$ from solution calorimetry are -793.3 ± 1.9 kJ mole⁻¹ and -579.8 ± 2.7 kJ mole⁻¹, respectively.

EDB SUBJECT CATEGORY: EDB-400702;400201
 INIS CATEGORY: B13

<83>

Browning, P.; Phillips, B.A.; Potter, P.E.; Rand, M.H. (UKAEA Research Group, Harwell. Atomic Energy Research Establishment) ** Blank, H.; Lindner, R. (eds.)

Segregation and vapour pressure studies on the uranium-plutonium-carbon system In Plutonium 1975 and other actinides ** AIX-08:309982;ERA-02:041839;EDB-77:094417; (1976) North-Holland

SUBJECT DESCRIPTORS: CARBON;FORMATION FREE ENTHALPY;LATTICE PARAMETERS;PHASE DIAGRAMS;PLUTONIUM: T3;PLUTONIUM CARBIDES: T2;SEGREGATION: Q1,Q2,Q3,Q4;SOLID SOLUTIONS;URANIUM: T4;URANIUM CARBIDES: T1;VAPOR PRESSURE: Q1,Q2,Q3,Q4;VERY HIGH TEMPERATURE

ABSTRACT: Some experimental studies are reported on the

<83>

<83> CONT.

segregation of U and Pu, and the plutonium vapour pressures, at 1400-1600°C, in the MC + M₂C₃ (M = (U+Pu)) phase field in which mixed uranium-plutonium 'monocarbide' fuels normally lie. Comparison of the segregation results with calculations based on ideal solution behaviour in the MC and M₂C₃ phases suggests the deviations from ideality are quite small. Vapour pressure measurements confirm this, indicating that any deviations are likely to be positive. Regular solution calculations and vapour pressure data are also reported for the phase fields containing graphite and sesquicarbide or dicarbide.

EDB SUBJECT CATEGORY: EDB-360204;360202;050700
 INIS CATEGORY: E24

<84>

Browning, P.; Potter, P.E.; Phillips, B.A.; Rand, M.H. **
 Commission of the European Communities, Karlsruhe (Germany, F.R.). European Inst. for Transuranium Elements

Segregation and vapour pressure studies in the uranium-plutonium-carbon system In 5th international conference on plutonium and other actinides **
 AIX-07:233104;EDB-76:087288; (1975) AED-Conf--75-417-019

SUBJECT DESCRIPTORS: PLUTONIUM CARBIDES: T1; SEGREGATION: Q1,Q2; URANIUM CARBIDES: T2; VAPOUR PRESSURE: Q1,Q2; VERY HIGH TEMPERATURE

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-360202
 INIS CATEGORY: B23

<85>

Bryan, G.H.; Thompson, J.K.; Van Tuyl, H.H.; Brown, C.L.; Ryan, J.L., Battelle Pacific Northwest Labs., Richland, Wash. (USA)

In Results of research to evaluate solid plutonium nitrate as a safe shipping form **
 ERA-02:021786; INS-77:005524; EDB-77:041187; (Nov 1976)
 BNWL--1941

SUBJECT DESCRIPTORS: FEASIBILITY STUDIES; PLUTONIUM NITRATES: T1; PYROLYSIS; RADIOLYSIS; SOLUBILITY; TRANSPORT: Q1

ABSTRACT: The results of the research described indicate that the compound Pu(NO₃)₃·4xH₂O is a viable material for consideration as a solid shipping form. It meets the criteria set forth to define a suitable solid form of

<85> CONT.

plutonium. Several plutonium compounds were considered for study as a shipping form before selecting $\text{Pu}(\text{NO}_3)_4 \cdot x\text{H}_2\text{O}$ as having the most desirable characteristics. It is reasonably easy to prepare on a kilogram scale with relatively simple equipment using vacuum evaporation techniques. Plutonium nitrate does begin to thermally decompose at about 100°C ; however, proper equipment design and instrumentation can overcome this problem. Although $\text{Pu}(\text{NO}_3)_4 \cdot x\text{H}_2\text{O}$ does produce some radiolytic gases, such as oxygen and nitrogen, an explosive mixture is not produced as only minor amounts of hydrogen are formed except during extended storage. Excessive pressurization in the shipping mode should not be a problem if the transport container is not closed for protracted periods of time. However, some form of thermal insulation would be required to prevent decomposition of the compound and to preclude over-pressurization during a fire. Plutonium nitrate has been found to be completely soluble in nitric acid even after storage for extended periods of time. After the equivalent of 25 years of storage time of first-cycle reactor-grade plutonium from a radiation damage standpoint, $\text{Pu}(\text{NO}_3)_4 \cdot x\text{H}_2\text{O}$ prepared from 85 percent ^{238}Pu was found to be completely soluble in 4M HNO_3 . Plutonium nitrate was also found to be soluble in water, but it rapidly converted to plutonium polymer.

EDE SUBJECT CATEGORY: EDB-050900
INIS CATEGORY: E15

<86>

Bukhalova, G.A.; Burlakova, V.M.; Sementsova, D.V.
(Rostovskij-na-Donu Inzhenerno-Stroitel'nyj Inst. (USSR))

Ternary systems from lithium, sodium, strontium and barium chlorides EDB-76:042590; Zh. Neorg. Khim. (ZNOKA) 20:(6):1657-1659 (Jun 1975)

SUBJECT DESCRIPTORS: BARIUM CHLORIDES:
T4;CRYSTALLIZATION;DIFFERENTIAL THERMAL
ANALYSIS;EUTECTICS;LITHIUM CHLORIDES: T1;MELTING;PHASE
DIAGRAMS: Q1,Q2,Q3,Q4;SODIUM CHLORIDES: T2;SOLID
SOLUTIONS;STRONTIUM CHLORIDES: T3

ABSTRACT: None

EDE SUBJECT CATEGORY: EDB-400201
INIS CATEGORY: B12

<87>

<87>

Bukhalova, G.A.; Litvinova, G.N.; Berezhnaya, V.T.; Burlakova, V.M. (Rostovskij-na-Donu Inzhenerno-Stroitel'nyj Inst. (USSR))

Phase equilibria in the system of sodium chlorides, cesium chlorides and strontium chlorides EDB-77:003056; Zh. Neorg. Khim. (ZNOKA) 20:(3):837-839 (Mar 1974)

SUBJECT DESCRIPTORS: CESIUM CHLORIDES: T2; DIFFERENTIAL THERMAL ANALYSIS; EQUILIBRIUM; PHASE STUDIES: Q1, Q2, Q3; SODIUM CHLORIDES: T1; STRONTIUM CHLORIDES: T3

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-360602
INIS CATEGORY: B12

<88>

Bukhalova, G.A.; Litvinova, G.N.; Berezhnaya, V.T.; Burlakova, V.M.

Phase equilibria in the system of sodium chlorides, cesium chlorides and strontium chlorides EDB-77:062251; Zh. Neorg. Khim. (ZNOKA) 20:(3):837-839 (Mar 1974)

SUBJECT DESCRIPTORS: CESIUM CHLORIDES: T2; DIFFERENTIAL THERMAL ANALYSIS; EQUILIBRIUM; PHASE STUDIES: Q1, Q2, Q3; SODIUM CHLORIDES: T1; STRONTIUM CHLORIDES: T3

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-360602

<89>

Bukhalova, G.A.; Mirsoyanova, N.N.; Yagub'yan, E.S. (Rostovskij-na-Donu Inzhenerno-Stroitel'nyj Inst. (USSR))

Quaternary mutual system from fluorides and chlorides of lithium, sodium and strontium AIX-08:298234; EDB-77:062511; Zh. Neorg. Khim. (ZNOKA) 20:(8):2214-2218 (Aug 1975)

SUBJECT DESCRIPTORS: EUTECTICS; LITHIUM CHLORIDES: T6; LITHIUM FLUORIDES: T1; MIXTURES; MOLTEN SALTS; PHASE DIAGRAMS: Q1, Q2, Q3, Q4, Q5, Q6; SODIUM CHLORIDES: T5; SODIUM FLUORIDES: T3; STRONTIUM CHLORIDES: T4; STRONTIUM FLUORIDES: T2; THERMAL ANALYSIS; THERMODYNAMICS

ABSTRACT: Simplified partition of a prism constituted by a quaternary Li, Na, Sr//F, Cl mutual system has been made. Visual-polythermal and thermographic methods were used to study the liquidus surface of three tetrahedron-forming

<89> CONT.

sections. Four points of non-variant equilibria have been found, one of these points being a transition point and the other three are eutectic ones (the lowest melting point being 434⁰C). A phase diagram of the Li, Na, Sr//F, Cl system has been constructed. Thermodynamic calculations have been made for reactions occurring at the center points of the conversion lines.

EDB SUBJECT CATEGORY: EDB-400201

INIS CATEGORY: B12

<90>

Bukhalova, G.A.; Yagub'yan, E.S.; Zaporozhets, E.G.; Buryakov, A.V. (Rostovskij-na-Donu Inzhenerno-Stroitel'nyj Inst. (USSR))

Quaternary system of potassium halogenides EDB-76:036689; Zh. Neorg. Khim. (ZNOKA) 20:(4):1099-1102 (Apr 1975)

SUBJECT DESCRIPTORS: PHASE DIAGRAMS: Q1,Q2,Q3,Q4;POTASSIUM BROMIDES: T1;POTASSIUM CHLORIDES: T2;POTASSIUM FLUORIDES: T3;POTASSIUM IODIDES: T4;THERMAL ANALYSIS

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-360602;400201

INIS CATEGORY: B12

<91>

Bulatova, V.F.; Morachevskii, A.G.; Demidov, A.I. (Leningradskij Politekhnikeskij Inst. (USSR))

Interaction of the lithium-tin alloys with the LiCl-KCl-CsCl melted eutectic mixture AIX-08:280594;EDB-77:037402; Elektrokhimiya (ELKKA) 12:(8):1339 (Aug 1976)

SUBJECT DESCRIPTORS: BINARY ALLOY SYSTEMS;CESIUM CHLORIDES: T4;CHEMICAL REACTION KINETICS;CHEMICAL REACTIONS: Q1,Q2,Q3,Q4,Q5;ELECTROLYSIS;ELECTROLYTES;EUTECTICS;LITHIUM ALLOYS: T1;LITHIUM CHLORIDES: T2;MOLTEN SALTS;POTASSIUM CHLORIDES: T3;QUANTITY RATIO;TIN ALLOYS: T5

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-400201

INIS CATEGORY: B12

<92>

<92>

Bulatova, V.F.; Morachevskii, A.G. (Leningradskij
Politekhnicheskij Inst. (USSR))

Equilibrium between Li-K-Te alloys and LiCl-KCl molten eutectic
mixture AIX-08:336897;EDB-78:017302; Zh. Prikl. Khim. (ZPKHA)
50:(1):183-184 (Jan 1977)

SUBJECT DESCRIPTORS: CATHODES;EQUILIBRIUM:
Q1,Q2,Q3,Q4;EUTECTICS;HIGH TEMPERATURE;LIQUID METALS;LITHIUM
CHLORIDES: T3;LITHIUM TELLURIDES: T2;MOLTEN
SALTS;POLARIZATION;POTASSIUM;POTASSIUM CHLORIDES:
T4;TELLURIDES: T1

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-400201
INIS CATEGORY: B12

<93>

Buraya, I.T.; Polyachenok, O.G.; Komshilova, O.M.; Novikov, G.I.
(Belorusskij Tekhnologicheskij Inst., Minsk)

Tensimetric study of pressure and composition of saturated vapor
in TlCl-InCl₃ system AIX-08:298235;EDB-77:062547; Zh.
Fiz. Khim. (ZPKHA) 50:(9):2441 (Sep 1976)

SUBJECT DESCRIPTORS: CHEMICAL COMPOSITION:
Q1,Q2;EQUATIONS;INDIUM CHLORIDES: T1;MOLECULES;MOLTEN
SALTS;SUBLIMATION;THALLIUM CHLORIDES: T2;VAPOR PRESSURE: Q1,Q2

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-400201
INIS CATEGORY: B12

<94>

Burch, W.D.; Feldman, M.J.; Groenier, W.S.; Vondra, B.L., Oak
Ridge National Lab., Tenn. (USA)

In LMFBR Fuel Reprocessing Program progress report for period
April 1 to June 30, 1976 ** ERA-02:004572;EDB-76:095940; (Aug
1976) ORNL/TM--5556

SUBJECT DESCRIPTORS: ADSORPTION;CONTROL
SYSTEMS;DESIGN;DISSOLUTION: Q4;ENGINEERING;FUEL REPROCESSING
PLANTS: T2;IODINE: T3;IODOX PROCESS: T;LMFBR TYPE REACTORS:
T1;MEASURING INSTRUMENTS;NITROGEN OXIDES: T6;OFF-GAS
SYSTEMS;PALLADIUM IODIDES: T5;PILOT PLANTS;PLUTONIUM DIOXIDE:
T4;RELIABILITY;REMOVAL: Q3;REPROCESSING: Q7;RESEARCH
PROGRAMS: Q2;SCRUBBING: Q6;SOLUBILITY: Q5;SPENT FUELS:

<94> CONT.

T7,Q1;VOLOXIDATION PROCESS;ZEOLITES

ABSTRACT: The program is aimed at providing the technology for the design of commercial reprocessing facilities that will be required by the late 1990s. The process and equipment development phase should culminate in a hot pilot-plant process demonstration in 1985-90 for the feed from FFTF and CRER. Criteria were developed for this hot pilot plant. Laboratory and hot-cell studies included the study of base metal zeolites for I^{2+} trapping, dissolution of PuCl_2 in HI, solubilities of PdI^{2+} , and fixation of iodine waste in concrete. Engineering research studies are being conducted on voloxidation, dissolution, feed preparation (nitrogen oxide scrubbing), fluorocarbon absorption, reliability, Iodex process, and instrumentation and control systems. Component testing was also conducted. The status of the Integrated Prototype Equipment Test Facility is reported. (DLC)

EDB SUBJECT CATEGORY: EDB-050800

<95>

Burmeister, R.A.; Hiskes, R.; Felmler, T.L., Hewlett-Packard Co., Palo Alto, Calif. (USA)

In Environmental sensitivity of structural metals: liquid metal embrittlement. Annual technical progress report No. 2, 25 Jun 1970--31 May 1971 ** EDB-77:094367; (Jun 1971) AD--885334

SUBJECT DESCRIPTORS: BARIUM FLUORIDES;BARIUM OXIDES;BORON OXIDES;CRYSTAL GROWTH: Q1,Q2,Q3,Q4,Q5,Q6;EPITAXY;MAGNETIC MATERIALS;SOLVENTS

ABSTRACT: A steady-state crystal growth system suitable for both homoepitaxial and heteroepitaxial growth of uniaxial magnetic rare earth compounds has been constructed and used for the growth of both YFeO_3 and EuErGaO_{12} . A very stable solvent, comprised of $\text{BaO-B}_2\text{O}_3\text{-BaF}_2$ has been developed. This solvent is nonvolatile and nonreactive, and is well suited to the solution growth of magnetic oxides. Arrays of bubbles have been generated in YFeO_3 and the coercivity and mobility of crystals grown from this solvent have been measured and found to be about 0.05 Oe and about 3500 cm/s-Ce, comparable to the values for crystals grown in the PbO -based solvents. Stable arrays of mobile bubbles have also been generated in heteroepitaxially grown EuErGaO_{12} . Emission spectrographic analysis indicates that barium

<95>

<95> CONT.

incorporation in the crystals grown from the BaO-based solvent is less than lead incorporated in those grown from the PbO-based solvent. (auth)

EDB SUBJECT CATEGORY: EDB-360201

<96>

Burylev, B.P.; Gershunina, V.Ya. (Krasnodarskiy Politekhnikheskiy Inst. (USSR))

Investigation into thermodynamic properties of FeCl_2 -CsCl system melts AIX-08:320595; EDB-77:133391; Zh. Fiz. Khim. (ZFKHA) 49:(9):2211-2215 (Sep 1975)

SUBJECT DESCRIPTORS: CESIUM CHLORIDES: T1; ELECTRIC BATTERIES; ELECTROMOTIVE FORCE; EQUATIONS; HIGH TEMPERATURE; IRON CHLORIDES: T2; MOLTEN SALTS; QUANTITY RATIO; REACTION KINETICS; THERMODYNAMIC PROPERTIES: Q1, Q2

ABSTRACT: Activities of iron chloride in the melt of the system FeCl_2 -CsCl have been studied by the emf method. Various expressions describing the dependence of the activity on the concentration have been examined. A most general equation has been derived from which all the known relationships can be deduced.

EDB SUBJECT CATEGORY: EDB-400201

INIS CATEGORY: B12

<97>

Burylev, B.P.; Mironov, V.L. (Krasnodarskiy Politekhnikheskiy Inst. (USSR))

Saturation vapor pressure in iron chloride-alkaline earth metal chlorides systems AIX-07:232716; EDB-76:054341; Izv. Vyssh. Uchebn. Zaved., Chern. Metall. (IVUMA) 12::9-11 (Dec 1974)

SUBJECT DESCRIPTORS: BINARY MIXTURES; EQUATIONS; IRON CHLORIDES: T1; ISOTHERMAL PROCESSES; MOLTEN SALTS; QUANTITY RATIO; SATURATION; STRONTIUM CHLORIDES: T2; TEMPERATURE DEPENDENCE; VAPOR PRESSURE: Q1, Q2; VERY HIGH TEMPERATURE

ABSTRACT: The boiling-point method was used to investigate the temperature and concentration dependence of the saturated vapour pressure in ferric chloride systems with alkali earth metal chlorides at 1000⁰K. Saturated vapour pressure isotherms were constructed for different solution ratios of these systems. In the FeCl_2 - MgCl_2 system slight positive deviations from additivity were observed, while in ferric chloride systems with calcium, strontium and barium

<97> CONT.

chlorides there were negative deviations increasing on transition from calcium to barium, i.e. with increasing cationic radius of the alkali earth metal. The vapour composition of weakly interacting systems is discussed.

EDE SUBJECT CATEGORY: EDB-400201;360603
INIS CATEGORY: B12

<98>

Burylev, B.P.; Mironov, V.L.; Tsemekhman, L.Sh.; Sryvalin, I.T.
(Krasnodarskij Politekhnikheskij Inst. (USSR))

Study of saturated vapor pressure above salt melts in CoCl_2 - MCl systems EDB-76:042455; Izv. Vyssh. Uchebn. Zaved., Khim. Khim. Tekhnol. (IVUKA) 18:(4):663-665 (1975)

SUBJECT DESCRIPTORS: BINARY MIXTURES;CESIUM CHLORIDES: T1;COBALT CHLORIDES: T2;COMPUTER CALCULATIONS;DIAGRAMS;EQUATIONS;LEAST SQUARE FIT;MOLTEN SALTS;QUANTITY RATIO;VAPOR PRESSURE: Q1,Q2

ABSTRACT: None

EDE SUBJECT CATEGORY: EDB-360603
INIS CATEGORY: B12

<99>

Buryleva, E.B.; Mironov, V.L.; Garanina, I.A.; Ignat'eva, L.N.
(Krasnodarskij Politekhnikheskij Inst. (USSR))

Temperature dependence of vapor pressure of rare earth halides and their alloys AIX-08:289193;EDB-77:069112; Zh. Fiz. Khim. (ZFKHA) 50:(8):2174-2175 (Aug 1976)

SUBJECT DESCRIPTORS: ALLOYS;HALIDES;RARE EARTH COMPOUNDS: T1;TEMPERATURE DEPENDENCE;VAPOR PRESSURE: Q1

ABSTRACT: None

EDE SUBJECT CATEGORY: EDB-400201
INIS CATEGORY: B12

<100>

<100>

Putskii, E.V.; Grigoryan, V.A.; Fillipov, A.F.; Topilin, V.V.;
Krasnova, I.A. (Moskovskij Inst. Stali i Splavov (USSR))

Solubility of nitrogen in multicomponent nickel base alloys
EDB-76:036267; Izv. Vyssh. Uchebn. Zaved., Chern. Metall.
(IVUMA) (1):47-51 (Jan 1975)

SUBJECT DESCRIPTORS: LIQUID METALS; MOLYBDENUM ALLOYS: T1; NICKEL
BASE ALLOYS: T2; NIOBIUM ALLOYS: T3; NITROGEN: T5; SOLUBILITY:
Q5; SCIENTIFIC PROPERTIES: Q1, Q2, Q3, Q4; TEMPERATURE
DEPENDENCE; THERMODYNAMIC MODEL; TUNGSTEN ALLOYS: T4; VERY HIGH
TEMPERATURE

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-360102
INIS CATEGORY: E22

<101>

Bykhovskij, A.I.; Pashchenko, A.V. (Ukrainskaya
Sel'skokhozyajstvennaya Akademiya, Kiev)

Nonisothermal flow of liquid metals over the surface of solid
ones EDB-76:036462; Fiz. Met. Metalloved. (FMMEA)
38: (6):1209-1217 (Dec 1974)

SUBJECT DESCRIPTORS: ABSORPTION HEAT; CAPILLARY FLOW:
Q5; DIAGRAMS; DIFFUSION; ENTHALPY; ISOTHERMAL PROCESSES; KINETIC
EQUATIONS; MERCURY: T5; SOLUBILITY; TEMPERATURE
DEPENDENCE; WETTABILITY: Q1, Q2, Q3, Q4

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-360104
INIS CATEGORY: E22

<102>

Cape, T.W.; Maroni, V.A.; Cunningham, P.T. (Argonne National
Lab., Ill. (USA)); Bates, J.B.

Raman and i.r.-emission studies of some tungstate- and
molybdate-containing melts AIX-07:262497; EDB-77:069054;
Spectrochim. Acta, Part A (SAMCA) 32A: (5):1219-1223 (1976)

SUBJECT DESCRIPTORS: INFRARED SPECTRA: Q1, Q2; MOLECULAR
STRUCTURE; MOLYBDENUM SALTS; MOLYBDATES: T1; RAMAN SPECTRA:
Q1, Q2; SALTS; SOLUTIONS; STRUCTURAL CHEMICAL
ANALYSIS; TUNGSTATES: T2; VIBRATIONAL STATES

ABSTRACT: Melts containing tungstate and molybdate have been studied

<102> CONT.

using Raman and i.r.-emission spectroscopy. Vibrational data obtained for MoO_4^{2-} and MoO_4^{3-} containing melts firmly support a tetrahedral structure for MoO_4^{2-} (and by implication for WO_4^{2-} as well) and establish the correct assignment of ν_2 and ν_4 for this species. Acidification of these MoO_4^{2-} and WO_4^{2-} melts by addition of the corresponding metal trioxides produced no significant change in the observed spectra, although significant solubilities of the trioxides in the WO_4^{2-} and MoO_4^{2-} melts were observed.

EDB SUBJECT CATEGORY: EDB-400201

INIS CATEGORY: B12

<103>

Cha, Y.S. (Argonne National Lab., IL); Costello, T.J.; Hopenfeld, J.; Miller, R.L. ** Cooper, M.H. (ed.), American Nuclear Society, Hinsdale, Ill.; American Inst. of Mining, Metallurgical, and Petroleum Engineers, Inc., New York; Energy Research and Development Administration, Washington, D.C. (USA); American Nuclear Society, Pittsburgh, Pa. Pittsburgh Section

Experimental investigation of cavitation inception and damage in a flowing sodium environment In Proceedings of the international conference on liquid technology in energy production ** INS-77:015776; ERA-02:059299; EDB-77:138125; (Nov 1976) CONF-760503--P2

SUBJECT DESCRIPTORS: CAVITATION: Q2; DATA; FLUID FLOW; LMFBR TYPE REACTORS: T1; MASS TRANSFER; MOCKUP; PRESSURE DROP; REACTOR COOLING SYSTEMS: M2, Q1; SIMULATION; SODIUM

ABSTRACT: The paper describes the interim results of cavitation inception and damage tests in a flowing sodium loop typical of that encountered in an LMFBR. Venturis were used as the cavitators in both the hot leg (538°C) and the cold leg (371°C) test sections. Cavitation inception data obtained in the present test showed no significant difference from those in water system reported previously. The venturi throat velocity for the damage test was equal to approximately 1.2 times that at inception for the same downstream pressure. Severe damage was observed in both venturis for a testing duration of 2000 hours with cavitation damage in the cold leg being more severe than that in the hot leg. Weight loss due to mass transfer and other effects was much smaller than that due to cavitation. Pressure drop across the venturis was observed to decrease with time during

<103>

<103> CONT.

the test, presumably due to the increase in venturi throat diameter.

EDB SUBJECT CATEGORY: EDB-210500;420400
INIS CATEGORY: E35;E11

<104>

Chandola, L.C.; Machado, I.J. (Bhabha Atomic Research Centre, Bombay (India). Spectroscopy Div.)

Spectrographic determination of trace impurities in reactor grade aluminium AIX-07:241781;EDB-76:071559; Indian J. Technol. (IJOTA) 13:(10):471-743 (Oct 1975)

SUBJECT DESCRIPTORS: ACCURACY;ALUMINIUM;ALUMINIUM OXIDES:
T1;ANTIMONY: T16;BISMUTH: T4;BORON: T3;CADMIUM: T5;CHEMICAL ANALYSIS: Q1;CHROMIUM: T7;COBALT: T6;COPPER: T8;EMISSION SPECTROSCOPY:
Q2,Q3,Q4,Q5,Q6,Q7,Q8,Q9,Q10,Q11,Q12,Q13,Q14,Q15,Q16,Q17,Q18,Q19,Q20,Q21;GALLIUM: T10;IMPURITIES;INDIUM: T11;IRON: T9;LEAD: T15;MAGNESIUM: T12;MOLYBDENUM: T13;NICKEL: T14;QUANTITATIVE CHEMICAL ANALYSIS;REACTOR MATERIALS;SAMPLE PREPARATION;SENSITIVITY;SILICON: T17;SILVER: T2;TIN: T18;TITANIUM: T19;TRACE AMOUNTS;VANADIUM: T20;ZINC: T21

ABSTRACT: A spectrographic method enabling the determination of 21 trace impurities in aluminium oxide is described. The technique involves mixing the sample with graphite buffer in the ratio 1:1, loading it in a graphite electrode and arcing it for 30 sec. in a dc arc to 10 A current against a pointed graphite cathode. The spectra are photographed on Ilford N.30 emulsion employing a large quartz spectrograph. The aluminium line at 2669.2 Å serves as the internal standard. The impurities determined are Ag, B, Bi, Cd, Co, Cr, Cu, Fe, Ga, In, Mg, Mo, Ni, Pb, Sb, Si, Sn, Ti, V and Zn. The sensitivity varies from 5 to 100 ppm and the precision from ± 5 to $\pm 22\%$ for different elements. A method for converting aluminium metal to aluminium oxide is described. It is found that boron is not lost during this conversion.

EDB SUBJECT CATEGORY: EDB-400104
INIS CATEGORY: B11

<105>

Chaplygin, G.V. (AN SSSR, Moscow. Inst. Kristallografii)

Experimental determination of formation heat of gaseous Ga_2O and In_2O_3 AIX-08:312148; EDB-77:094787; Zh. Fiz. Khim. (ZFKHA) 49: (11):2767-2771 (Nov 1975)

SUBJECT DESCRIPTORS: CHEMICAL PREPARATION; CHEMICAL REACTIONS; FORMATION HEAT: Q1, Q2; GALLIUM OXIDES: T2; INDIUM; INDIUM OXIDES: T1; LIQUID METALS; VERY HIGH TEMPERATURE; WATER VAPOR

ABSTRACT: The flow method was used to study the reaction $2\text{In}(\text{liq.}) + \text{H}_2\text{O}(\text{gas}) \rightleftharpoons \text{In}_2\text{O}_3(\text{gas}) + \text{H}_2(\text{gas})$ at temperatures of 1100-1300 deg K. The vapor pressure of water was 6.1×10^{-3} atm. The experimental value of the enthalpy of the reaction was determined by taking into account possible errors present in the standard energy function. This made it possible to obtain a more precise value for the standard enthalpy for the formation of gaseous In_2O_3 : $\Delta H_f(298 \text{ deg K}) = -15.7 \pm 1.0 \text{ kcal/mole}$ (according to the Third law of thermodynamics) and -14.6 kcal/mole (according to the second law of thermodynamics). These values are close to the corresponding average values reported in literature and calculated according to the Third law of thermodynamics. The use of a simple chemical reaction made it possible to increase the precision of determining the standard enthalpy of $\text{In}_2\text{O}_3(\text{gas})$ formation.

EDB SUBJECT CATEGORY: EDB-400201

INIS CATEGORY: B12

<106>

Cheburkov, V.I.; Rozanov, A.N.

In Kinetics of reaction of liquid lithium with oxygen and nitrogen ** ERA-02:049275; EDB-77:107370; Metall. Metalloved. Chist. Met., Sb. Nauchn. Rabot (MMIFA) (7):168-173 (1968) ENWL-tr--247

SUBJECT DESCRIPTORS: ACTIVATION ENERGY; CHEMICAL REACTION KINETICS: Q2, Q3; CHEMICAL REACTIONS; HIGH TEMPERATURE; LIQUID METALS; LITHIUM: M1; LOW PRESSURE; MEDIUM VACUUM; SOLUBILITY; SOLVENT PROPERTIES

ABSTRACT: The kinetics of the reaction of liquid lithium with oxygen was investigated in the temperature range 210 to 640°C at pressures of 130 to 1 mm Hg. The kinetics of the reaction of liquid lithium with nitrogen was investigated at temperatures ranging from 295 to 409°C at 20 to 0.5 mm Hg. The observed deviations of the kinetic curves of

<106>

<106> CONT.

oxidation and nitriding of liquid lithium from a linear relation are due to the solution of oxygen and nitrogen in the liquid metal. The solubility of oxygen in lithium in the temperature range 210 to 635°C (0.003 to 0.57 at. percent respectively) was determined from the kinetic curves. The activation energies of the processes of oxidation ($Q_{\text{lin}} = 15.6$ kcal/mole and $Q_{\text{log}} = 15.3$ kcal/mole) and nitriding ($Q = 23$ kcal/mole) of lithium and the heat of solution of oxygen in lithium ($\Delta H = 11.7$ kcal/mole) were calculated.

EDB SUBJECT CATEGORY: EDB-400201;360105

<107>

Chereau, P.; Dean, G.; de Franco, M. (CEA Centre d'Etudes Nucleaires de Pontenay-aux-Roses, 92 (France). Dept. d'Etudes des Combustibles a Base de Plutonium); Gerdanian, P.

High-temperature microcalorimetric measurements of partial molar enthalpy of solution, $\Delta H(O_{\text{sub } 2})$, in plutonium + oxygen AIX-08:305647;EDB-77:107493; J. Chem. Thermodyn. (JCTDA) 9: (3):211-219 (Mar 1977)

SUBJECT DESCRIPTORS: CALORIMETRY;CHEMICAL COMPOSITION;OXYGEN;PHASE DIAGRAMS: Q1;PLUTONIUM;PLUTONIUM OXIDES: T1;SOLUTION HEAT: Q1;VARIATIONS;VERY HIGH TEMPERATURE

ABSTRACT: Direct measurements of $\Delta H(O_{\text{sub } 2})$, the partial molar enthalpy of solution of oxygen, have been performed, at 1373 K, with a high-temperature microcalorimeter, in the plutonium + oxygen system. The variation of $\Delta H(O_{\text{sub } 2})$ with composition gives new information, especially in the $\text{PuO}_{\text{sub } (1.5)}$ to $\text{PuO}_{\text{sub } (1.75)}$ region, which are in agreement with the new phase diagram proposed by Boivineau (J. Nucl. Mater.; 31:38 (1976)), from high-temperature X-ray studies. This system is similar to cerium + oxygen.

EDB SUBJECT CATEGORY: EDB-400702
INIS CATEGORY: B13

<108>

Chernov, V.S.; Busol, F.I.

Mechanism of metals modifying AIX-07:236361;EDB-76:053654; Izv. Akad. Nauk SSSR, Met. (IZNMA) (2):71-77 (Mar 1975)

SUBJECT DESCRIPTORS: ADSORPTION;FORON;CRYSTAL GROWTH;CRYSTAL-PHASE TRANSFORMATIONS: Q1;CRYSTALLIZATION;DENDRITES;DIFFUSION;EQUATIONS;EUTECTICS;GERMANIUM;INTERFACES;METALS: T1;PHASE

<108> CONT.

DIAGRAMS;PGTASSIUM;QUANTITY RATIO;SODIUM;SOLID
SOLUTIONS;SOLUBILITY;SURFACE TENSION;TEMPERATURE DEPENDENCE

ABSTRACT: Data from the literature are cited to show that in several cases, the mechanism of modification of metals and alloys by additives soluble in the melt cannot be explained by the adsorption hypothesis based on the surface activity of these additives. It is suggested that the mechanism of modification by soluble additives can be explained by the presence of a layer of liquid enriched (or depleted) in the additives preceding the crystallization front (densification by concentration), formed under actual conditions of crystallization as a result of the different solubilities of the additive in the solid and liquid phases of the base metal, regardless of its surface activity. This densification by concentration leads to the inhibition of crystal growth in the base metal (barrier action) and to concentration overcooling. On the basis of this theory it is suggested that the modifying action of additives can be predicted from some parameters of the phase diagrams.

EDB SUBJECT CATEGORY: EDB-360102
INIS CATEGORY: B22

<109>

Chilton, G.R.; Kirkham, I.A. ** Commission of the European Communities, Karlsruhe (Germany, F.R.). European Inst. for Transuranium Elements

Determination of oxygen potentials of hyperstoichiometric U--Pu dioxides in the temperature range 1,500-1,850 K In 5th international conference on plutonium and other actinides. Book of abstracts ** AIX-07:232722;EDB-76:087400; (1975) AEE-Conf--75417-013

SUBJECT DESCRIPTORS: CHEMICAL COMPOSITION:
Q1,Q2;EQUILIBRIUM;PLUTONIUM DIOXIDE: T1;URANIUM DIOXIDE:
T2;VERY HIGH TEMPERATURE

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-400102;050700
INIS CATEGORY: B12

<110>

<110>

Chilton, G.R.; Kirkham, I.A. (UKAEA Reactor Group, Windscale.
Reactor Development Labs.) ** Plank, H.; Lindner, R. (eds.)

Determination of oxygen potentials of hyperstoichiometric U-Pu
dioxides in the temperature range 1500-1850 K In Plutonium
1975 and other actinides **

AIX-(8:309989; ERA-02:041826; EDB-77:094384; (1976)
North-Holland

SUBJECT DESCRIPTORS: CHEMICAL COMPOSITION: Q1,Q2;FREE
ENTHALPY;OXYGEN;PHASE STUDIES;PLUTONIUM DIOXIDE:
T1;TEMPERATURE DEPENDENCE;THERMODYNAMICS;URANIUM DIOXIDE:
T2;VERY HIGH TEMPERATURE

ABSTRACT: Oxygen potentials for $U_{sub}(0.85)Pu_{sub}(0.15)$ and
 $U_{sub}(0.69)Pu_{sub}(0.31)$ oxides have been measured at 1518, 1623
and 1823K using a transpiration technique. The results show
that the oxygen potentials for the oxide become progressively
less negative with increasing plutonium content over the
temperature range used. Differences exist between
extrapolation of previously published low temperature data
and the results of this work.

EDB SUBJECT CATEGORY: EDB-360202;050700
INIS CATEGORY: B24

<111>

Chilton, J.E. Jr., Lockheed Missiles and Space Co., Palo Alto,
Calif. (USA)

In Lithium--silver chloride secondary battery investigation.
Quarterly technical progress report No. 2 ** EDB-76:047750;
(Jul 1964) AD--450428

SUBJECT DESCRIPTORS: ADDITIVES;ALUMINATES;ALUMINIUM;ALUMINIUM
CHLORIDES;BROMIDES;CARBONIC ACID
ESTERS;CATHODES;CHLORATES;CHLORINE COMPOUNDS;CONTAINERS:
Q1;ELECTROCHEMISTRY:
Q1;ELECTRODEPOSITION;ETHERS;FABRICATION;GRAPHITE;LITHIUM;LITHIUM
CHLORIDES;LITHIUM COMPOUNDS;METAL-NONMETAL BATTERIES:
M1;ORGANIC SOLVENTS;ORGANOMETALLIC
COMPOUNDS;SEALS;SILVER;SILVER CHLORIDES;SILVER
COMPOUNDS;SOLUBILITY;TOLUENE

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-250902;250903

<112>

Cho, K. (Korea Univ., Kodaira, Tokyo (Japan). Faculty of Engineering)

Study on conductivity, density, and viscosity of molten salt systems AIX-08:300825;EDB-77:089225; Denki Kagaku (DNKKA) 44:(5):335-338 (May 1976)

SUBJECT DESCRIPTORS: CORRELATIONS; DENSITY: Q1,Q2,Q3,Q4,Q5; DYSPROSIUM CHLORIDES: T1; ELECTRIC CONDUCTIVITY: Q1,Q2,Q3,Q4,Q5; GADOLINIUM CHLORIDES: T2; LANTHANUM CHLORIDES: T3; MELTING POINTS; MOLTEN SALTS; NEODYMIUM CHLORIDES: T4; PRASEODYMIUM CHLORIDES: T5; VISCOSITY: Q1,Q2,Q3,Q4,Q5

ABSTRACT: A relation between the equivalent conductivity and density for molten salts is deduced with the aid of significant structures theory, and the solid state density at melting point is evaluated approximately for some rare-earth metal chlorides and the other chlorides. Furthermore, the relation among the equivalent conductivity, density, and viscosity for some molten salts is discussed.

EDB SUBJECT CATEGORY: EDB-400201

INIS CATEGORY: B12

<113>

Chudinov, E.G.; Cdinov, B.V.

Investigation into reactions between trioctylammonium nitrates and rare earth nitrates by X-ray electron spectroscopy AIX-08:286002;EDB-77:044226; Koord. Khim. (KOKHD) 2:(6):768-771 (1976)

SUBJECT DESCRIPTORS: BINDING ENERGY; CHEMICAL REACTIONS: Q1,Q2,Q3; COMPLEXES; ELECTRON SPECTROSCOPY; NITRATES: T2; NITRIC ACID; RARE EARTH COMPOUNDS: T3; TOA: T1; X-RAY SPECTROSCOPY

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-400201;400301

INIS CATEGORY: B12

<114>

<114>

Chupin, A.N.; Krylov, E.I. ** AN SSSR, Sverdlovsk. Inst. Khimii;
 Vostochnyj Nauchno-Issledovatel'skij i Proektnyj Inst.
 Ogneupornoj Promyshlennosti, Sverdlovsk (USSR); Ural'skij
 Nauchno-Issledovatel'skij Khimicheskij Inst., Sverdlovsk
 (USSR)

Synthesis and properties of rare earth titanotungstates In
 Summaries of reports of the all-union meeting on solid body
 chemistry. 3--5 June 1975. III. Synthesis, phase
 equilibrium and use of solid oxide compounds **
 AIX-08:289196;EDB-77:094481; (1975) INIS-mf--3246

SUBJECT DESCRIPTORS: CALCINATION;CHEMICAL PREPARATION:
 Q1,Q2,Q3;CRYSTAL STRUCTURE;PHYSICAL PROPERTIES: Q1,Q2,Q3;RARE
 EARTH COMPOUNDS: T1;TEMPERATURE DEPENDENCE;TITANIUM
 COMPOUNDS: T3;TUNGSTATES: T2;VERY HIGH TEMPERATURE

ABSTRACT: None

EDE SUBJECT CATEGORY: EDB-360601
 INIS CATEGORY: B12

<115>

Chuprikov, G.E.; Epifanova, K.I.; Malyutina, T.M.; Samokhin, G.A.

Compatibility of molten rare earth metals with refractory
 materials In Nauchnye trudy GIREDMETA. Vol. 52 **
 AIX-08:317938;EDB-77:120044; (1974) GIREDMET

SUBJECT DESCRIPTORS: CAVITIES;CHEMICAL COMPOSITION;CHROMIUM:
 T7;COMPATIBILITY:
 Q1,Q2,Q3,Q4,Q5,Q6,Q7,Q8,Q9,Q10;CRYSTALLIZATION;MELTING;METALL
 GRAPHY;MICROHARDNESS;MOLYBDENUM: T10;MOLYBDENUM ALLOYS:
 T6;NIOBIUM: T8;OXYGEN;PLASTICITY;POROSITY;RARE EARTHS:
 T1;SOLUBILITY: Q1,Q2,Q3,Q4,Q5,Q6,Q7,Q8,Q9,Q10;TANTALUM
 ALLOYS: T5;TUNGSTEN: T9;TUNGSTEN ALLOYS: T4;VANADIUM:
 T3;WEAR;ZIRCONIUM: T2

ABSTRACT: Interaction of rare-earth metal and yttrium melts with
 Zr, V, Nb, Cr, W, Mo refractory metals as well as alloys of
 W-Mo, W-Ta systems has been studied. W /2-3%/ Mo alloy proved
 to be most chemically resistant with respect to the melts
 under consideration. It is shown that the W /2-3%/ Mo alloy
 should be used as the crucible material for growing the
 specimens from the melt intended for precision physical
 measurements at a temperature of liquid helium since tungsten

<116> CONT.

tantalum which becomes a superconductor at a temperature of 4.5 deg K.

EDB SUBJECT CATEGORY: EDB-360101;360104

INIS CATEGORY: E22

<116>

Clark, R.P.; Elucher, R.L.; Goldsmith, H.J.

Phase diagram of system $\text{LiCl--KCl--CaCrO}_4$ EDB-77:131934;
J. Chem. Eng. Data (JCEAA) 14:(4):465-470 (Oct 1969)

SUBJECT DESCRIPTORS: CALCIUM COMPOUNDS: T4;CHEMICAL
COMPOSITION;CHROMATES: T5;ELECTROLYTES: Q1;EUTECTICS;LITHIUM
CHLORIDES: T2;MELTING POINTS;PHASE DIAGRAMS:
Q2,Q3,Q4,Q5;POTASSIUM CHLORIDES: T3;THERMAL BATTERIES: T1

ABSTRACT: The phase diagram was determined for the system $\text{LiCl--KCl--CaCrO}_4$. Binary eutectics were located at 60 wt % $\text{KCl--40 wt % CaCrO}_4$ (mp 651 C) and 62 wt % $\text{LiCl--38 wt % CaCrO}_4$ (m 638 C); ternary eutectic was found at 41 wt % $\text{LiCl--50 wt % KCl--9 wt % CaCrO}_4$ (mp 342 C). It was verified by x-ray diffraction that $\text{LiCl--KCl--CaCrO}_4$ is stable diagonal section in quaternary reciprocal system K, Li, CaCrO_4 , Cl. The salt mixture $\text{LiCl--KCl--CaCrO}_4$ is electrolyte--depolarizer combination used in thermal cells (voltaic cells employing molten salt electrolyte).

EDB SUBJECT CATEGORY: EDB-250903;360602

<117>

Clark, R.P.; Reinhardt, F.W.

Phase diagram for the ternary system $\text{CaCl}_2\text{--KCl--CaCrO}_4$
4\$ FRA-01:016809;EDB-76:042604; Thermochim. Acta (THACA)
14: (1/2): 113-129 (1976)

SUBJECT DESCRIPTORS: CALCIUM CHLORIDES: T1;CALCIUM COMPOUNDS:
T3;CHROMATES: T4;CRYSTALLIZATION;DIFFERENTIAL THERMAL
ANALYSIS;EUTECTICS;PHASE DIAGRAMS: Q1,Q2,Q3,Q4;POTASSIUM
CHLORIDES: T2;THERMAL BATTERIES;X-RAY DIFFRACTION

ABSTRACT: The phase diagram for the system $\text{CaCl}_2\text{--KCl--CaCrO}_4$ has been determined using differential thermal analysis in conjunction with x-ray diffraction. $\text{CaCl}_2\text{--KCl--CaCrO}_4$ has been shown to be a stable diagonal section of the Ca,K//Cl, CrO_4 reciprocal ternary system. The 3 binary systems are: $\text{CaCl}_2\text{--KCl}$ which exhibits a congruently melting compound (CaKCl_3)

<117>

<117> CONT.

melting at 741°C with eutectics at 24.0 mole percent KCl (m.p. 615°C) and 74.3 mole percent KCl (m.p. 594°C); CaCl_2 -- CaCrO_4 which shows a eutectic at 23.4 mole percent CaCrO_4 (m.p. 660°C); and KCl-- CaCrO_4 with a eutectic at 24.2 mole percent CaCrO_4 (m.p. 651°C). The binary congruently melting compound separates the ternary system into 2 pseudoternary subsystems. A binary eutectic exists in the pseudobinary system CaKCl_3 -- CaCrO_4 at 12.9 mole percent CaCrO_4 (m.p. 672°C). Ternary eutectics exist at 71.8 mole percent CaCl_2 --17.3 percent KCl--10.9 percent CaCrO_4 (m.p. 575°C) and 23.2 mole percent CaCl_2 --71.2 percent KCl--5.6 percent CaCrO_4 (m.p. 573°C). Isotherms are shown for the liquidus surface (primary crystallization) and for the secondary crystallization surface. Isothermal and vertical sections through the phase diagram are also shown. (auth)

EDE SUBJECT CATEGORY: EDB-400201

<118>

Clinton, J.E. Jr.; Conner, W.J.; Cook, G.M.; Holsinger, R.W.,
Lockheed Missiles and Space Co., Sunnyvale, Calif. (USA)

In Lithium--silver chloride secondary battery investigation.
Final report, December 1963--November 1964 ** EDB-77:061339;
(Feb 1965) AD--612189

SUBJECT DESCRIPTORS: ALUMINIUM CHLORIDES; CARBONIC ACID
ESTERS; CHEMICAL
REACTIONS; CONTAINERS; DESIGN; ENTROPY; ETHERS; FABRICATION; FREE
ENERGY; LITHIUM; LITHIUM CHLORIDES; METAL-NONMETAL BATTERIES;
M1; PERFORMANCE; Q1; POLYETHYLENES; REACTION HEAT; SILVER
CHLORIDES; THERMODYNAMIC PROPERTIES; TOLUENE

ABSTRACT: None

EDE SUBJECT CATEGORY: EDB-250901

<119>

Cogley, D.R.; Butler, J.N.; Synnott, J.C., Tyco Labs., Inc.,
Waltham, Mass. (USA)

In Composition of nonaqueous solutions of potential use in high
energy density batteries. Final report, 15 Nov 1969--14 Nov
1972 ** EDE-76:052748; (1972) AD--763140

SUBJECT DESCRIPTORS: DESIGN; DMSO; METAL-NONMETAL BATTERIES:

<119> CONT.

M1;SOLUBILITY: Q2,Q3,Q4,Q5,Q6;STORAGE LIFE

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-250901;400201

<120>

Collins, J.L.; Osborne, M.F.; Malinauskas, A.P.; Lorenz, R.A.;
Manning, S.R., Oak Ridge National Lab., Tenn. (USA)

In Knudsen cell--mass spectrometer studies of cesium--urania
interactions ** INS-77:000334;ERA-02:010656;EDB-77:007630;
(Jun 1976) ORNL/NUREG/TM--24

SUBJECT DESCRIPTORS: CESIUM;CESIUM CARBONATES: T2;CESIUM
HYDROXIDES: T3;CESIUM URANATES: M4;CHEMICAL REACTION YIELD:
Q4;CHEMICAL REACTIONS: Q1,Q2,Q3;FISSION PRODUCTS;KNUDSEN
FLOW;MASS SPECTROSCOPY;PHASE STUDIES;URANIUM DIOXIDE:
M1;VAPOR PRESSURE;VERY HIGH TEMPERATURE

ABSTRACT: Limited Knudsen cell--mass spectrometer studies were
made of the partial pressures of cesium-containing species
[assumed to be primarily Cs(g)] over Cs_2CO_3 and
over phase equilibria involving UO_2 and probable Cs-U-O
compounds formed from mixtures that initially contained
either $\text{Cs}_2\text{CO}_3\text{-UO}_2$ or CsOH-UO_2 .
Although additional work is required to further define the
equilibria involved, the data demonstrate unambiguously a
significant reduction in cesium partial pressures due to
probable Cs-U-O compound formation and indicate essentially
identical behavior with either CsOH or Cs_2CO_3 as
the starting material with UO_2 .

EDB SUBJECT CATEGORY: EDB-400702;400201
INIS CATEGORY: B13;B12

<121>

Coninck, R. de; Lierde, W. van; Gijs, A. (Centre d'Etude de
l'Energie Nucleaire, Mol (Belgium))

Uranium carbide: thermal diffusivity, thermal conductivity and
spectral emissivity at high temperatures EDB-76:048840; J.
Nucl. Mater. (JNUMA) 57:(1):69-76 (Jul 1975)

SUBJECT DESCRIPTORS: EMISSIVITY;THERMAL CONDUCTIVITY;THERMAL
DIFFUSIVITY;THERMODYNAMIC PROPERTIES: Q1;URANIUM CARBIDES:
T1;VERY HIGH TEMPERATURE

ABSTRACT: The thermal diffusivity, thermal conductivity, and
spectral emissivity at $0.65 \mu\text{m}$ of uranium carbide have

<121>

<121> CONT.

been measured with the aid of a modulated electron-beam apparatus. For hypostoichiometric samples, a reversible, time-dependent oxidation-reduction of the surface interfered with the measurements between 1100 and 1400\$sup 0\$C, without affecting the results. Expressions are derived representing the thermal diffusivity, thermal conductivity, and spectral emissivity for stoichiometric and for slightly hypo- and hyperstoichiometric uranium carbide.

EDB SUBJECT CATEGORY: EDB-360202

INIS CATEGORY: B24

<122>

Courths, R. (Freie Univ. Berlin (F.R. Germany). Inst. fuer Atom- und Festkoerperphysik)

Optical absorption Zeeman spectroscopy of rare earth orthochromites. II. Theory AIX-07:261722;EDB-77:019349; Z. Phys., B (ZPBBD) 22:(3):253-259 (Nov 1975)

SUBJECT DESCRIPTORS: ABSORPTION SPECTROSCOPY;ANISOTROPY;CHROMIUM COMPOUNDS: M1;ELECTRONIC STRUCTURE: Q1,Q2;MAGNETIC FIELDS;MAGNETO-OPTICAL EFFECTS;RARE EARTH COMPOUNDS: M2;RARE EARTHS;TEMPERATURE DEPENDENCE;ZEEMAN EFFECT

ABSTRACT: Using a single ion molecular field model, expressions for the energy of Kramers ion states of rare earths in rare earth orthochromites (RECrO_3) have been derived as a function of temperature and applied magnetic fields. These microscopic expressions reflect the various RE-RE and RE-Cr coupling constants of the isotropic-symmetric and antisymmetric interactions of an RE ion in the various magnetic states of RECrO_3 .

EDB SUBJECT CATEGORY: EDB-400201

INIS CATEGORY: A13

<123>

Courths, R.; Huefner, S. (Freie Univ. Berlin (F.R. Germany). Inst. fuer Atom- und Festkoerperphysik)

Optical absorption Zeeman spectroscopy of rare earth orthochromites. I. ErCrO_3 AIX-07:261721;EDB-77:019348; Z. Phys., B (ZPBBD) 22:(3):245-251 (Nov 1975)

SUBJECT DESCRIPTORS: ABSORPTION SPECTROSCOPY: Q1,Q2;CATIONS;CHROMIUM COMPOUNDS: T1;ERBIUM COMPOUNDS: T2;MAGNETIC DIPOLES;MAGNETO-OPTICAL EFFECTS;RARE EARTH COMPOUNDS;SPECIFIC HEAT;TEMPERATURE DEPENDENCE;ZEEMAN EFFECT

<123> CONT.

ABSTRACT: The optical absorption spectrum of Er^{3+} in ErCrO_3 in the phases γ_1 , γ_2 and γ_4 has been studied as a function of temperature and applied magnetic field. The experimental data for the Er^{3+} ground doublet splitting are analyzed using a single ion molecular field approach. The analysis shows that the Cr-Er interaction is predominantly antisymmetric exchange, the magnetic dipole-dipole interaction being an order of magnitude smaller. The Er-Er interaction is of exchange as well as of dipolar origin.

EDE SUBJECT CATEGORY: EDB-400201
INIS CATEGORY: A13

<124>

Cox, D.L.; Cronenberg, A.W. (New Mexico Univ., Albuquerque (USA))

Theoretical prediction of the thermal conductivity of uranium carbide vapor AIX-08:343057; EDB-78:021943; J. Nucl. Mater. (JNUMA) 67:(3):326-331 (Aug 1977)

SUBJECT DESCRIPTORS: CHAPMAN-ENSKOG THEORY; CHEMICAL REACTIONS; DISSOCIATION; ENTHALPY; LMFBR TYPE REACTORS; MOLECULE-MOLECULE COLLISIONS; REACTOR CORE DISRUPTION; THERMAL CONDUCTIVITY: Q1, Q2; URANIUM CARBIDES: T1; VAPORS: T2

ABSTRACT: Theoretical predictions are made for the thermal conductivity of uranium carbide vapor. The calculations take into account collisional, dissociation, and chemical reaction effects.

EDE SUBJECT CATEGORY: EDB-360204
INIS CATEGORY: B24

<125>

Dalvi, A.G.I.; Deodhar, C.S.; Joshi, B.D. (Bhabha Atomic Research Centre, Bombay (India). Radiochemistry Div.)

Chemical separation and spectrographic estimation of rare earths and yttrium in PuO_2 and $(\text{U}, \text{Pu})\text{O}_2$ AIX-08:312068; EDB-77:094725; Talanta (TLNTA) 24:(2):143-145 (Feb 1977)

SUBJECT DESCRIPTORS: CERIUM: T1; CHEMICAL ANALYSIS: Q15, Q16; DYSPROSIUM: T2; EMISSION SPECTROSCOPY: Q1, Q2, Q3, Q4, Q5, Q6, Q7, Q8, Q9, Q10, Q11, Q12, Q13, Q14; ERBIUM: T3; EUROPIUM: T4; GADOLINIUM: T5; HOLMIUM: T6; IMPURITIES; LANTHANUM: T7; LUTETIUM: T8; MIXTURES; NEODYMIUM: T9; PLUTONIUM DIOXIDE: T15; QUANTITATIVE CHEMICAL

<125>

<125> CONT.

ANALYSIS;SAMARIUM: T10;SOLVENT EXTRACTION;TERBIUM:
 T11;THULIUM: T12;TOA;TRACE AMOUNTS;URANIUM DIOXIDE:
 T16;XYLENES;YTTERBIUM: T13;YTTRIUM: T14

ABSTRACT: An emission spectrographic method for the estimation of rare-earth impurities in plutonium oxide and the mixed oxide (U,Pu) $O_{2\frac{1}{2}}$ (30% $PuO_{2\frac{1}{2}}$ - 70% $UO_{2\frac{1}{2}}$) has been developed. Rare-earth impurities are separated from the matrix by solvent extraction with tri-n-octylamine in xylene and are estimated after concentration. Thulium is used as internal standard and LiF-AgCl as carrier. The spectra are excited in a d.c. arc. The detection limits lie in the range 0.004 to 0.1 μg and the coefficient of variation ranges from 5 to 23%.

EDE SUBJECT CATEGORY: EDB-400104
 INIS CATEGORY: B11

<126>

Danan, J. (CEA Centre d'Etudes Nucleaires de Fontenay-aux-Roses, 92 (France))

Specific heat of thorium monocarbide from 2 to 300K
 EDB-76:054020; J. Nucl. Mater. (JNUMA) 57: (3):280-282 (Sep 1975)

SUBJECT DESCRIPTORS: DEBYE TEMPERATURE;ELECTRONIC SPECIFIC HEAT;LATTICE VIBRATIONS;LOW TEMPERATURE;SPECIFIC HEAT: Q1;TEMPERATURE DEPENDENCE;THORIUM CARBIDES: T1;ULTRALOW TEMPERATURE;VERY LOW TEMPERATURE

ABSTRACT: The specific heat of ThC has been measured in the range 1.5 to 300 $^{\circ}K$. From the results, the thermodynamic functions were calculated at 298.15 $^{\circ}K$; the Debye and Einstein temperatures are linked with the acoustic and optical branches, respectively, of the lattice vibrations, and the coefficient ($\gamma=2.12mJ/mol.K^2$) of electronic specific heat.

EDE SUBJECT CATEGORY: EDB-360204
 INIS CATEGORY: B24

<127>

Dao, N.Q.; Knidiri, M. (Ecole Centrale des Arts et Manufactures, 92 - Chatenay-Malabry (France). Inst. de Chimie)

Study of the compounds $M\text{UO}_2\text{F}_5$ ($M = \text{K}, \text{Rb}, \text{Cs}, \text{NH}_4$) by i.r. absorption and Raman diffusion spectrophotometry. III. Effect of cation on vibration frequency of the ion $\text{UO}_2\text{F}_5^{3-}$
 AIX-07:262500;EDB-77:019519; Spectrochim. Acta, Part A (SAMCA) 32A:(5):1113-1118 (1976)

SUBJECT DESCRIPTORS: AMMONIUM COMPOUNDS: T1;CATIONS;CESIUM FLUORIDES: T2;CHEMICAL BONDS;INFRARED SPECTRA;MOLECULAR STRUCTURE;POTASSIUM FLUORIDES: T3;RAMAN SPECTRA;RUBIDIUM COMPOUNDS: T4;STRUCTURAL CHEMICAL ANALYSIS;URANYL FLUORIDES: T5;VIBRATIONAL STATES: Q1,Q2,Q3,Q4,Q5

ABSTRACT: Normal coordinate analysis of the $\text{UO}_2\text{F}_5^{3-}$ ion using the Wilson-Decius GF matrix method was carried out. Good agreement with the observed spectra was obtained. Discussions are given on the force constants obtained for U-O and U-F bonds and on the outer-sphere interaction in the series $M\text{UO}_2\text{F}_5$.

EEB SUBJECT CATEGORY: EDB-400702
 INIS CATEGORY: B12

<128>

Davies, B.S.J.; Polley, M.V.; Skyrme, G., Central Electricity Generating Board, Berkeley (UK). Berkeley Nuclear Labs.

In Physical models of mass transport of iron and nickel in liquid sodium systems ** AIX-07:248041;EDB-77:018848; (Dec 1975) CEGE-RD/B/M--3548

SUBJECT DESCRIPTORS: CORROSION: Q1,Q2;CORROSIVE EFFECTS: Q3;DIFFUSION;IRON: T1;LIQUID FLOW;LIQUID METALS;MASS TRANSFER;NICKEL: T2;SODIUM: T3;SOLUBILITY

ABSTRACT: Experimental observations on corrosion of pure iron and nickel specimens in non-isothermal loops containing flowing sodium have been used to derive values of the concentration of dissolved material at the entrance to the test section and diffusion coefficients of the test material in sodium. The former values differ from the saturation value by only 10^{-3} ppm, which is small compared to currently recommended solubility values. The phenomenon cannot be explained in terms of circulating particles. Two other possible explanations are also dismissed. The diffusion coefficient values are consistent with the corroding species being atoms, or molecules containing a few atoms. It is also shown that the observations are better explained in terms of

<128>

<128> CONT.

boundary layer controlled mass transfer, rather than a surface controlled process. A computer model based on an alternative solubility relationship is shown to produce results which describe well the observed variation of corrosion rate with oxygen concentration, sodium velocity and downstream position.

EDB SUBJECT CATEGORY: EDB-360105
INIS CATEGORY: E15

<129>

Davydov, A.D.; Danil'tseva, N.A.; Kamkin, A.N.

Electrochemical machining of the molybdenum VM-1 alloy
AIX-08:295619;EDB-77:055195; Elektron. Obrab. Mater. (ECBMA)
(4):8-9 (Jul 1976)

SUBJECT DESCRIPTORS: ALLOY-VM-1; T1; ELECTROCHEMICAL MACHINING;
Q1; ELECTROLYTES; METALLOGRAPHY; POTASSIUM
BROMIDES; PRODUCTIVITY; QUANTITY RATIO; SODIUM CARBONATES; SODIUM
CHLORIDES; SODIUM HYDROXIDES; SODIUM
NITRATES; SOLUBILITY; SURFACES

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-360101
INIS CATEGORY: E21

<130>

de Beer, R.; van Ormondt, D.; Homs, M.H. (Technische Hogeschool Delft (Netherlands)); den Hartog, H.W. (Rijksuniversiteit Groningen (Netherlands))

^{139}La ENDOR of $\text{LaCl}_3 \cdot \text{Gd}^{3+}$
AIX-08:308023;EDB-77:082742; Physica, B, C (PHBCD)
84:(3):367-370 (Oct 1976)

SUBJECT DESCRIPTORS: CATIONS; DOPED MATERIALS; ENDOR;
Q2; GADOLINIUM IONS; HAMILTONIANS; HYPERFINE STRUCTURE;
Q1; LANTHANUM: T1; LANTHANUM 139: T2; LANTHANUM
CHLORIDES; LIGANDS; MAGNETIC FIELDS; MICROWAVE
RADIATION; MULTICHARGED IONS; QUADRUPOLES; SPECTRA; SPIN; ULTRALOW
TEMPERATURE

ABSTRACT: The magnetic transferred h.f.i. and the electric quadrupole interaction of the two ^{139}La nuclei nearest to Gd^{3+} in LaCl_3 have been determined with ENDOR, using Ka-band (34.5 GHz) microwaves. For both lanthanum positions it was found that $A_{\text{sub}}(s)/A_{\text{sub}}(p)$ approximately 1 and $A_{\text{sub}}(p)-A_{\text{sub}}(d)$ approximately 0. The ENDOR data give no

<130> CONT.

evidence for significant local distortion around Gd^{3+} . This implies that ligand displacements due to the difference in size between Gd^{3+} and La^{3+} cannot account for the disagreement between the observed and calculated B^{2+} of Gd^{3+} .

EDB SUBJECT CATEGORY: EDB-360603
INIS CATEGORY: A13

<131>

de Bievre, P.; van Audenhove, J. (Commission of the European Communities, Geel (Belgium). Central Bureau for Nuclear Measurements)

Accurate procedure to safeguard the fissile material content of input and output solutions of reprocessing plants in Safeguarding nuclear materials. Vol. 2 **
AIX-08:284444; ERA-02:019449; EDB-77:035054; (1976) IAEA

SUBJECT DESCRIPTORS: ACCOUNTING: Q2; FISSILE MATERIALS: T2; FUEL REPROCESSING PLANTS: T1; FUEL SCANNING; ISOTOPE DILUTION; MASS SPECTROSCOPY; MATERIAL BALANCE; PLUTONIUM; QUANTITY RATIO; SAFEGUARDS: Q1; SPENT FUELS; URANIUM

ABSTRACT: A procedure is described by which to assay accurately fissile material contents of input and output solutions of reprocessing plants, using well-defined solid spikes and isotope dilution mass spectrometry. The procedure has been used in measurements performed at the request of the Directorate of Euratom Safeguards in the 1972-1975 period and some results are given. The procedure allows the fissile element contents to be guaranteed at the time and place of sample-taking (i.e. of safeguards inspection), which is an improvement over guaranteeing such contents at the time and place of analysis.

EDB SUBJECT CATEGORY: EDB-055001; 050800
INIS CATEGORY: F41

<132>

De Coninck, R.; De Batist, R.; Gijss, A. (Centre d'Etude de l'Energie Nucleaire, Mol (Belgium))

Thermal diffusivity, thermal conductivity and spectral emissivity of uranium dicarbide at high temperatures
AIX-08:295891; EDB-77:068695; High Temp. - High Pressures (HTHPA) 8: (2): 167-176 (1976)

SUBJECT DESCRIPTORS: ELECTRON BEAMS; EMISSION SPECTRA; EMISSIVITY: Q1; HIGH TEMPERATURE; TEMPERATURE DEPENDENCE; TEMPERATURE

<132>

<132> CONT.

MEASUREMENT; THERMAL CONDUCTIVITY: Q1; THERMAL DIFFUSIVITY:
Q1; URANIUM CARBIDES: M1; VERY HIGH TEMPERATURE; X-RAY
DIFFRACTION

ABSTRACT: Thermal diffusivity and thermal conductivity between 600 and 2070^{°C}, and spectral emissivity at 0.65 and at 2.3 micron between 1000 and 2070^{°C} have been determined for UC₂ samples with C/U approximately equal to 1.8 with a modulated-electron-beam thermal-diffusivity apparatus. Some associated thermophysical properties are discussed in accordance with metallographic microscopic examinations and x-ray diffraction analysis. The transformation of the samples into the (UC + UC₂) cubic solid-solution phase is clearly observable through the four measured quantities. Formation of U₂C₃, between 1100 and 1600^{°C}, was accompanied by an increase of the diffusivity and, indirectly, through cracks and holes, of the emissivities, but the conductivity remained unchanged. A simple but reliable procedure for correcting temperature measurements with a two-color pyrometer when the sample cannot be regarded as a gray body, as is the case for UC₂, is described.

EDF SUBJECT CATEGORY: EDF-360204; 050700
INIS CATEGORY: B23

<133>

de Franco, M.; Gatesoupe, J.P. (CEA Centre d'Etudes Nucleaires de Fontenay-aux-Roses, 92 (France)) ** Blank, H.; Lindner, R. (eds.)

Thermodynamic properties of (U,Pu)O_{2-x} described by a cluster model In Plutonium 1975 and other actinides **
AIX-(8:309996; ERA-02:041840; EDF-77:094418; (1976)
North-Holland

SUBJECT DESCRIPTORS: CHEMICAL COMPOSITION; CRYSTAL
MODELS; ENTHALPY; ENTROPY; FORMATION FREE
ENTHALPY; INTERSTITIALS; MIXTURES; PLUTONIUM OXIDES:
T1; TEMPERATURE DEPENDENCE; THERMODYNAMIC PROPERTIES:
Q1, Q2; URANIUM OXIDES: T2; VACANCIES

ABSTRACT: A simple model is presented which allows the easy computation of ΔG_{O2} and ΔH_{O2} , at any temperature and oxygen and plutonium content for U₂PuO_{2-x}. Some implications on the nature of the defects

<134> CONT.

are drawn, values for their energy of formation are given and their concentrations may be computed.

EDB SUBJECT CATEGORY: EDB-360204;050700

INIS CATEGORY: B24

<134>

De Keroulas, P.; Calais, D.; Marcon, J.P.

In Distribution coefficient for U and Pu in the system (UPu)C + (UPu)₂C₃ ** ERA-01:016692;EDB-76:042331; J. Nucl. Mater. (JNUMA) 44::64-70 (1972) ORNL-tr--4133

SUBJECT DESCRIPTORS: FREE ENERGY;PHASE STUDIES;PLUTONIUM CARBIDES: T2;THERMODYNAMICS: Q1,Q2;URANIUM CARBIDES: T1

ABSTRACT: As part of a research program on mixed uranium/plutonium carbide, which is a possible fuel for fast reactors, a microprobe analyzer was used to examine the distribution of uranium and plutonium between the phases (U, Pu)C and (U, Pu)₂C₃. Two types of samples were available, with Pu/(U + Pu) ratios of 0.20 and 0.50. The proportion of (U, Pu)₂C₃ varied from 10 to 70 percent. Two equilibria temperatures, 1500 and 1700°C, were used. Measurements of the distribution coefficients $k_{U/} = \text{amount of uranium in (U, Pu)C} / \text{amount of uranium in (U, Pu)}_{2}C_{3}$, $k_{Pu/} = \text{amount of plutonium in (U, Pu)C} / \text{amount of plutonium in (U, Pu)}_{2}C_{3}$ for each of the 14 samples led to values of the coefficients independent of temperature and proportions of the two phases. It is shown that with the help of two relationships (one defined by the proportion of phases (z and (1 - z)) and by the total ratio $\text{GAMMA} = U/(U + Pu)$, $\text{GAMMA} = xz + y(1 - z)$, where $x/y = k_{U/}$ and $(1 - x)/(1 - y) = k_{Pu/}$; the other defined by the free energies ΔG of the two phases $\Sigma \Delta G = [\Delta G(UC) - \Delta G(PuC)] - [\Delta G(UC/\text{sub } 1.5/) - \Delta G(PuC/\text{sub } 1.5/)] = RT \log [y/(1 - y) \times (1 - x)/x]$, one can determine a priori values of x and y from a knowledge of GAMMA and z. By using the experimentally determined distribution coefficients and values of x and y = f(x) deduced from the second equation for arbitrary values of the various free energies, one can determine summation ΔG experimentally. The result is consistent with recent measurement of ΔG for stoichiometric PuG. (auth) one can determine $\Sigma \Delta G$ experimentally. The result is consistent with recent measurement of ΔG for stoichiometric PuC. (auth)

EDB SUBJECT CATEGORY: EDB-360202

<135>

<135>

Dem'yanovich, M.A.; Ivanov, P.M.; Novoselov, G.P.; Prusakov, V.N.; Renard, Eh.V.; Skiba, O.V.; Serov, A.V.; Tsarenko, A.F.; Shevchenko, V.B.; Shmidt, V.S., International Atomic Energy Agency, Vienna (Austria)

In Basic problems of fast reactor fuel reprocessing **
 AIX-C8:302956; ERA-02:037117; EDB-77:079691; (1977)
 IAEA-CN--36/318

SUBJECT DESCRIPTORS: BOR-60 REACTOR:

T1;CHEMISORPTION;DECLADDING;ENTRAINMENT;FBR TYPE REACTORS:
 T3;FLUORIDE VOLATILITY PROCESS;PURIFICATION;RADIATION
 PROTECTION;RADIOACTIVITY;REPROCESSING: Q4;SBR-5 REACTOR:
 T2;SOLVENT EXTRACTION;SPENT FUEL ELEMENTS: T4;SPENT FUELS:
 Q1,Q2,Q3;URANIUM DIOXIDE

ABSTRACT: The paper considers two major directions in the recovery of fuel elements of nuclear power plants operated by fast reactors that are being developed in the USSR: a) aqueous procedures based on dissolution of fast reactor fuel with the following isolation and complete purification of valuable fuel components by a solvent extraction method; b) non-aqueous procedures based on fluorination of valuable fuel components and their purification from fission products by sublimation of higher fluorides. The section of the report dealing with aqueous procedures considers the problems of dissolution, radiation chemistry of aqueous and organic solutions, extraction and separation of uranium and plutonium macroamounts (equilibrium and kinetics), mathematical modelling of extraction process, use of water soluble neutron poisons, valent state of plutonium macroconcentrations, etc. The section of the report dealing with fluorination procedures considers the results of the experimental reprocessing of reactor BOR-60 irradiated uranium fuel with the burn-up above 10% cooled for 3-6 months. The paper gives the results of decladding experiments carried out on BOR-60 and BR-5 reactor fuels by the method of melting, as well as data on the distribution of fission-product elements in the apparatuses of experimental flowsheet and in the products of fuel recovery. The paper also gives data on amounts, corrosion properties, gas evolution, and activity of wastes and discusses the conditions of their long-term safe storage.

EDB SUBJECT CATEGORY: EDB-C50800
 INIS CATEGORY: B16

<136>

Denisova, N.D.; Bystrova, C.N.; Safronov, E.K.

Properties of zirconium and hafnium tetrachlorides and their saturated solutions In Proceedings of GIREDMET **
AIX-08:320603;EDB-77:113887; (1975) GIREDMET

SUBJECT DESCRIPTORS: EQUATIONS OF STATE; HAFNIUM CHLORIDES: T1; PHASE STUDIES: Q1, Q2; TEMPERATURE DEPENDENCE; VAPOR PRESSURE; ZIRCONIUM CHLORIDES: T2

ABSTRACT: The properties of saturated vapor of zirconium tetrachloride and hafnium tetrachloride were determined by investigating the P-V-T-N dependence of these compounds in gas phase. For each compound measurements were made of the molar volume and pressure of saturated vapor and the dependence of these parameters on the temperature. The saturated vapor of zirconium tetrachloride and hafnium tetrachloride obeys the ideal gas law starting with the following values of pressure, molar volume and temperature respectively: 2.5 kg/cm², 20600 cm³/mole, 360 deg C and 2.6 kg/cm², 19600 cm³/mole, 340 deg C. Phase diagram within P-N coordinates of the ZrCl₄-HfCl₄ system, calculated by using the state equation and Eicon equation, was approximately the same as reported in previous papers.

EDB SUBJECT CATEGORY: EDB-360603;400201
INIS CATEGORY: B12

<137>

Deodhar, S.S., Syracuse Univ., N.Y. (USA)

In Thermodynamics and kinetics of the formation of rare earth intermetallics ** INS-76:018149;EDB-76:071210; (1975) Syracuse Univ.

SUBJECT DESCRIPTORS: ACTIVATION ENERGY; BINARY ALLOY SYSTEMS; DIFFUSION; FORMATION HEAT; FORMATION HEAT: Q1, Q2, Q3, Q4; INTERMETALLIC COMPOUNDS; LAVES PHASES; MAGNETIC PROPERTIES; RARE EARTH ALLOYS: T1; THERMODYNAMIC PROPERTIES

ABSTRACT: Heats of reaction of rare earth intermetallics with iron, cobalt and nickel were determined using Differential Thermal Analysis technique. The intermetallic compounds studied were of MgCu₂ type Laves phases and the rare earth elements studied were praseodymium, gadolinium, dysprosium and erbium. The reactions were exothermic and the heats of reaction were generally high. They varied from the low of -2.5 kcal/g mole for Fe₂Gd to the high of -35.3 kcal/g mole for Ni₂Er. The magnitudes of heats of reaction were always greater for the intermetallics of heavy

<137>

<137> CONT.

rare earth elements. The rare earth intermetallics studied were either ferromagnetic or antiferromagnetic. The variations in the magnetic moments and the heats of reaction with respect to the atomic number of the rare earth elements followed certain trends. The similarities were observed in the trends of two properties. Electronic configuration for the $\text{MgCu}_{2\text{ type}}$ rare earth intermetallics is proposed using Engel--Brewer correlation for metallic structures and the structural features of the Laves phase compounds. Kinetics of the reactions between the rare earth elements and iron, cobalt, and nickel was studied. The rate of reaction was diffusion controlled in each case. The Valensi--Carter equation for the diffusion mechanism satisfactorily described the kinetic behavior. The magnitudes of activation energies and frequency factors were determined. The reactions can be characterized by their reaction temperatures since they always begin at definite temperatures. It was observed that the reaction began at a higher temperature if the activation energy for the reaction was high.

EDB SUBJECT CATEGORY: EDB-360102
INIS CATEGORY: B22

<138>

Dergacheva, M.B.; Kozin, L.F.; Imangaziev, E.K. (AN Kazakhskoj SSR, Alma-Ata. Inst. Organicheskogo Kataliza i Ehlektrokhimii)

Determination of vibrational contribution into excessive mixing entropy of liquid amalgams. Alkali metal amalgams
AIX-08:337211; EDB-77:140518; Zh. Fiz. Khim. (ZFKHA)
51:(2):500-503 (Feb 1977)

SUBJECT DESCRIPTORS: CESIUM ALLOYS: T3; COMPUTER CALCULATIONS; COORDINATION NUMBER; ENTROPY: Q1, Q2, Q3; LIQUID METALS: T1; MERCURY ALLOYS: T2; MIXING HEAT: Q1, Q2, Q3; QUANTITY RATIO

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-360104
INIS CATEGORY: B22

<139>

Desyatnik, V.N.; Dubinin, B.V.; Mel'nikov, Yu.T.; Raspopin, S.P.

Interaction of uranium trichloride with alkaline earth metal
chlorides EDB-76:037147; Zh. Neorg. Khim. (ZNOKA)
20:(4):1085-1087 (Apr 1975)

SUBJECT DESCRIPTORS: BARIUM CHLORIDES: T5; BINARY
MIXTURES; CALCIUM CHLORIDES: T3; CHEMICAL REACTIONS:
Q1, Q2, Q3, Q4, Q5; DIFFERENTIAL THERMAL
ANALYSIS; EUTECTICS; MAGNESIUM CHLORIDES: T2; PHASE DIAGRAMS:
Q1, Q2, Q3, Q4, Q5; STRONTIUM CHLORIDES: T4; URANIUM CHLORIDES: T1

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-400702
INIS CATEGORY: B13

<140>

Desyatnik, V.N.; Dubinin, B.V.; Kurbatov, N.N.

Thermographic investigation into ternary systems containing
sodium and potassium chlorides, uranium trichloride and
thorium tetrafluoride AIX-07:277958; EDB-77:031784; Zh. Neorg.
Khim. (ZNOKA) 21:(2):575-578 (Feb 1976)

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-400703; 400201
INIS CATEGORY: B13

<141>

Desyatnik, V.N.; Katyshev, S.F.; Raspopin, S.P.; Chervinskii, Yu.F.

Density, surface tension, and viscosity of uranium
trichloride-sodium chloride melting
AIX-07:262686; EDB-77:031122; At. Energ. (USSR) (AENGA)
39:(1):70-72 (Jul 1975)

SUBJECT DESCRIPTORS: ACTIVATION
ENERGY; DATA; DENSITY; MELTING; PHYSICAL PROPERTIES:
Q1, Q2; QUANTITY RATIO; SODIUM CHLORIDES: T1; SURFACE
TENSION; TEMPERATURE DEPENDENCE; URANIUM CHLORIDES: T2; VERY
HIGH TEMPERATURE; VISCOSITY

<141>

<142> CONT.

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-360603

INIS CATEGORY: P22

<142>

Desyatnik, V.N.; Katyshev, S.F.; Raspopin, S.P.; Chervinskii, Y.F. (Ural'skij Politekhnikheskij Inst., Sverdlovsk (USSR))

Density, surface tension and viscosity of melts of uranium trichloride-lithium chloride AIX-09:349320; EDB-78:028043; Zh. Prikl. Khim. (ZPKHA) 50:(4):765-769 (Apr 1976)

SUBJECT DESCRIPTORS: COMPLEXES; DENSITY: Q1, Q2; LITHIUM CHLORIDES: T1; MOLTEN SALTS; QUANTITY RATIO; SURFACE TENSION: Q1, Q2; TEMPERATURE DEPENDENCE; URANIUM CHLORIDES: T2; VISCOSITY: Q1, Q2

ABSTRACT: The results are presented of studying the density, viscosity, and surface tension of the melts of the system $\text{UCl}_3\text{-LiCl}$ in a wide temperature (888-1282°K) and concentration ranges. For all the melts of the system in the studied temperature range, linear dependences of density and surface tension and exponential dependence of viscosity on temperature have been obtained. It has been established that at 1123°K the density of the melts increases monotonously when passing from LiCl to UCl_3 . In the whole concentration range the density isotherm has a positive deviation from linearity. The surface tension isotherm deviates from linearity towards the negative side in the whole concentration range. In the region 20-40 mol% of UCl_3 , a minimum is observed on the isotherm due to the formation of complexes surface-active with respect to both components. The isotherms of molar viscosity and a change in the activation energy of viscous flow depending on the composition of melted mixtures also point to the complex formation in the system.

EDB SUBJECT CATEGORY: EDB-400702; 400201

INIS CATEGORY: B13

<143>

Desyatnik, V.N.; Kurbatov, N.N.; Raspopin, S.P. (Ural'skij Politekhnikheskij Inst., Sverdlovsk (USSR). Kafedra Metallurgii Redkikh Metallov)

Phase diagram of the $\text{ThCl}_4\text{-LiCl-PbCl}_2$ system AIX-07:274976; EDB-77:031759; Izv. Vyssh. Uchebn. Zaved., Tsvet. Metall. (IVUTA) (3):137-138 (1975)

<143> CONT.

SUBJECT DESCRIPTORS: DIFFERENTIAL THERMAL ANALYSIS; LEAD
CHLORIDES: T1; LITHIUM CHLORIDES: T2; MOLTEN SALTS; PHASE
DIAGRAMS: Q1, Q2, Q3; THORIUM CHLORIDES: T3; X-RAY DIFFRACTION

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-400702
INIS CATEGORY: B16

<144>

Desyatnik, V.N.; Raspopin, S.P.

Interaction of uranium- and thorium tetrachlorides with sodium-
and potassium chlorides EDB-76:037150; Zh. Neorg. Khim.
(ZNOKA) 20: (5):1385-1388 (May 1975)

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-400702
INIS CATEGORY: B16

<145>

Desyatnik, V.N.; Raspopin, S.P.; Trifonov, K.I. (Ural'skij
Politekhnikeskij Inst., Sverdlovsk (USSR). Kafedra
Metallurgii Redkikh Metallov)

Viscosity of UCl_3 - MgCl_2 binary system melts
AIX-07:272483; EDB-77:031758; Izv. Vyssh. Uchebn. Zaved.,
Tsvet. Metall. (IVUTA) (4):168-170 (1975)

SUBJECT DESCRIPTORS: BINARY MIXTURES; MAGNESIUM CHLORIDES:
T1; MOLTEN SALTS; QUANTITY RATIO; TEMPERATURE DEPENDENCE; URANIUM
CHLORIDES: T2; VISCOSITY: Q1, Q2

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-400702
INIS CATEGORY: B13

<146>

Desyatnik, V.N.; Raspopin, S.P.; Trifonov, K.I.

Viscosity of molten CaCl_2 - UCl_3 mixtures
AIX-07:270244; EDB-77:031760; Zh. Prikl. Khim. (ZPKHA)
48: (5):1157-1158 (May 1975)

SUBJECT DESCRIPTORS: ACTIVATION ENERGY; BINARY MIXTURES; CALCIUM
CHLORIDES: T1; CHEMICAL COMPOSITION; MOLTEN SALTS; QUANTITY
RATIO; TEMPERATURE DEPENDENCE; URANIUM CHLORIDES: T2; VISCOSITY:

<146>

<146> CONT.

Q1,Q2

ABSTRACT: None

EED SUBJECT CATEGORY: EDB-400702

INIS CATEGORY: B13

<147>

Desyatnik, V.N.; Trifonov, K.I.; Raspopin, S.P.

Calculation of thermodynamic activity and liquidus lines of components in binary systems of plutonium trichloride with alkali and alkaline earth metal chlorides

AIX-08:334409;EDB-77:133658; Radiokhimiya (RADKA)

18:(1):105-108 (Jan 1976)

SUBJECT DESCRIPTORS: CHEMICAL COMPOSITION;COMPARATIVE EVALUATIONS;CRYSTALLIZATION;MOLTEN SALTS;PHASE DIAGRAMS: Q1,Q2,Q3,Q4,Q5;PLUTONIUM CHLORIDES: T1;THERMODYNAMICS

ABSTRACT: Comparison of experimental crystallization lines for plutonium trichloride and chlorides of alkaline and alkaline-earth metals (Na, K, Cs, and Sr) with theoretical ones has shown a possibility of forming complex ions of different composition in melts. As the ionic radii of alkaline and alkaline-earth metals increase so does the deviation of activity of components from ideal.

EED SUBJECT CATEGORY: EDB-400702;400201

INIS CATEGORY: B13

<148>

Desyatnik, V.N.; Vorobei, M.P.; Kurbatov, N.N.

Phase diagrams of ternary systems containing sodium and potassium chlorides, thorium tetrachloride and plutonium trichloride AIX-07:272482;EDB-77:031408; At. Energ. (USSR) (AENGA) 38:(3):173-174 (Mar 1975)

ABSTRACT: None

EED SUBJECT CATEGORY: EDB-400201;400702

INIS CATEGORY: B13

<149>

Dewally, D.; Perrot, P. (Lille-1 Univ., 59 - Villeneuve-d'Ascq (France))

Contribution to the thermodynamic study of uranium and nickel mixed oxides AIX-08:317692; EDB-77:132977; Rev. Chim. Miner. (RVCMA) 13:(6):605-613 (1976)

SUBJECT DESCRIPTORS: DIFFERENTIAL THERMAL ANALYSIS: Q1,Q2; ENTROPY; FREE ENTHALPY; NICKEL OXIDES: T1; PHASE DIAGRAMS; THERMAL GRAVIMETRIC ANALYSIS: Q1,Q2; URANIUM OXIDES: T2

ABSTRACT: The study by differential thermal analysis and by thermogravimetric analysis under controlled atmospheres of UO_2 , U_3O_8 and the mixed oxides $\text{NiU}_3\text{O}_{10}$ and NiU_2O_6 have led to the enthalpy of formation and to the standard entropy. $\Delta H_f^\circ = -942150$ et -610820 cal. $S^\circ = 85,6$ and $48,5$ cal/mole. deg. respectively for $\text{NiU}_3\text{O}_{10}$ and NiU_2O_6 . The free enthalpy of formation of $\text{NiU}_3\text{O}_{10}$ from its oxides ($\text{NiO} + 3 \text{UO}_2$) is given by the expression $\Delta G^\circ = -7350 - 2.3T$. The oxygen potential-temperature diagrams relative to the system U-Ni-O has been established and also the nature of phases in equilibrium. From the eleven monovariant equilibria, four have been determined by experience, four extracted from the literature and three calculated. Domains of stability of oxides have been precised.

EDB SUBJECT CATEGORY: EDB-360204
INIS CATEGORY: B12

<150>

dos Santos, A.M., Kernforschungsanlage Juelich G.m.b.H. (Germany, F.R.). Inst. fuer Chemische Technologie; Technische Hochschule Aachen (Germany, F.R.)

In Formation of actinides in irradiated HTGR fuel elements **
AIX-07:270367; ERA-02:021774; EDB-77:041176; (Mar 1976)
Juel--1284

ABSTRACT: Actinide nuclide concentrations of 11 spent AVR fuel elements were determined experimentally. The burnup of the spheres varied in the range between 10% and 100% fima, the Th:U ratio was 5:1. The separation procedures for actinide isolation were tested with highly irradiated ThO_2 . Separation and decontamination factors are presented. Actinide nuclide formation can be described by exponential functions of the type $\ln m_{\text{nuclide}} = A + B \times \text{fima}$. The empirical factors A and B were calculated performing a least squares analysis. Build-up of ^{232}U was discussed.

<150>

<150> CONT.

According to the experimental results, ^{232}U is mainly produced from ^{230}Th , a certain amount (e.g. about 20% at a 10^{55} MWd/t burnup) originated from a $(n,2n)$ reaction of ^{233}U ; a formation from ^{233}Th by a $(n,2n)$ followed by a (n,γ) reaction was not observed. The AVR breeding rate was ascertained to be 0.5. The hazard potential of high activity waste was calculated. After a 1,000 years' storage time, the elements Pa, Am and Cm will no longer influence the total hazard index. Actinide recovery factors were proposed in order to reduce the hazard potential of the waste by an actinide removal in consideration of the reprocessing technology which is available presently.

EDB SUBJECT CATEGORY: EDB-C50800;400103
INIS CATEGORY: B16

<151>

Drobacheva, T.I.; Skoropad, T.S. (Rostovskij-na-Donu
Inzhenerno-Stroitel'nyj Inst. (USSR))

Binary system of sodium, rubidium and cesium tungstates
AIX-C8:292292;EDB-77:069063; Zh. Fiz. Khim. (ZPKHA)
50:(4):1044 (Apr 1976)

SUBJECT DESCRIPTORS: BINARY MIXTURES;CESIUM TUNGSTATES:
T1;EUTECTICS;PHASE DIAGRAMS: Q1,Q2,Q3;RUBIDIUM TUNGSTATES:
T3;SODIUM;TUNGSTATES: T2

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-400201
INIS CATEGORY: B12

<152>

Drotning, W.D., Sandia Labs., Albuquerque, N.Mex. (USA)

In Thermal expansion of molten materials by the gamma
attenuation technique **
INS-78:000262;ERA-03:008785;EDB-78:009351; (1977)
SAND--77-0626C

SUBJECT DESCRIPTORS: ABSORPTION;ACCURACY;LIQUID METALS:
T1;LIQUIDS: T2;MEASURING METHODS: Q3;RESOLUTION;THERMAL
EXPANSION: T3,Q1,Q2

ABSTRACT: An apparatus was constructed for measurement of the density and volumetric thermal expansion of high temperature molten materials using the gamma attenuation technique. It was found that significant improvements in overall experimental precision could be realized by employing an

<152> CONT.

automatic gain control in the linear section of the gamma counting system. This device reduced errors due to system gain shifts and thereby increased the precision of the gamma counting. Corrections were also made for the dead time of the electronics system. A preliminary experimental investigation of the precision of the method was made by measuring small dimensional steps of a tool steel block at a single temperature. A fractional length increment $\Delta l/l$ of 1% was determined to 1%, equivalent to a length change precision of 10^{-4} cm/cm. In addition, the precision of the experimental technique was analytically examined for both absolute and relative density determinations. Three analytical expressions used to reduce data for liquid density determinations were evaluated for their precision. Each allows use of a different set of input data parameters, which can be chosen based on experimental considerations. Using experimentally reasonable values for the precision of the parameters yields a similar resultant density precision from the three methods, on the order of 0.2%.

EEB SUBJECT CATEGORY: EDB-360104;360204

INIS CATEGORY: B22;B23

<153>

Dudareva, A.G.; Bogatov, Yu.Eh.; Lityagov, V.Ya.; Molodkin, A.K.
(Universitet Druzhby Narodov, Moscow (USSR))

Investigation into interaction of indium tribromide with dibromides of manganese, cobalt and nickel in melt
AIX-08:292295;EDB-77:069045; Zh. Neorg. Khim. (ZNOKA)
20: (9):2566-2567 (Sep 1975)

ABSTRACT: State diagrams of $\text{InBr}_3\text{-MnBr}_2$, $\text{InBr}_3\text{-CoBr}_2$, $\text{InBr}_3\text{-NiBr}_2$ were studied with differential thermal analysis, roentgenophase and crystalloptical methods. Indium tribromide and manganese dibromide produce limited solid solutions. $\text{InBr}_3\text{-CoBr}_2$ and $\text{InBr}_3\text{-NiBr}_2$ are the systems with simple eutectics. The eutectic mixture containing 14 molar % of CoBr_2 and 86 molar % of InBr_3 melts at 408 deg C. In the system $\text{InBr}_3\text{-NiBr}_2$ the eutectic point is sharply shifted towards InBr_3 and corresponds to 98 molar % of InBr_3 and 2 molar % of NiBr_2 . The data of thermal analysis were confirmed by crystalloptical and roentgenophase analysis.

EEB SUBJECT CATEGORY: EDB-400201

INIS CATEGORY: B12

<154>

<154>

Dumoulin, E.; Tinelli, G.; Hatterer, A. (Ecole Supérieure de Chimie, 68 - Mulhouse (France). Lab. de Chimie Minérale Appliquée)

Solubility of sodium iodide in liquid sodium between 140 and 500^{°C} AIX-08:280610; EDB-77:030802; J. Inorg. Nucl. Chem. (JINCA) 38:(10):1876-1877 (1976)

SUBJECT DESCRIPTORS: HIGH TEMPERATURE; LIQUID METALS; SODIUM: T2; SODIUM IODIDES: T1; SOLUBILITY: Q1; SOLUTIONS; SOLVENT PROPERTIES: Q2

ABSTRACT: Results are presented for the solubility of sodium iodide in liquid sodium between 140 and 500^{°C}. The results are discussed.

EEB SUBJECT CATEGORY: EDB-360104; 400201
INIS CATEGORY: B12

<155>

Dutchak, Ya.I.; Klym, N.M.; Frenchko, V.S. (L'vivskij Gosudarstvennyj Univ. (Ukrainian SSR))

Atomic structure of melts of complex system with eutectic transformations AIX-08:335890; EDB-78:021663; Izv. Vyssh. Uchebn. Zaved., Fiz. (IVUFA) (7):88-92 (1976)

ABSTRACT: Short-range order structure of compound eutectic melts In-Sb-Sn, In-Sb and Al-Sb-Bi is considered. The investigations are carried out with a high-temperature X-ray diffractometer. The intensity curves of X-ray coherent scattering are used for obtaining functions of radial distribution of atoms (FRDA). The analysis of FRDA is performed for different structural models. It has been established that a most appropriate structural model of the In-Sb-Sn melt is a complex "quasieutectic" distribution corresponding to atom coordination in melted InSb and in liquid In and Sn. In a eutectic alloy of the In-Sb system in the range of complete melting there are microinhomogeneous regions enriched with InSb atom complexes and pure Sb. The structure of the Al-Sb-Bi melt is that typical for doped structures. The melts studied with temperature increasing tend to formation of a homogeneous solution.

EEB SUBJECT CATEGORY: EDB-360102
INIS CATEGORY: A13

<156>

Dybkov, V.I.; Eremenko, V.H.; Natanzon, Ya.V.

Molybdenum and tungsten interaction with liquid aluminium
 AIX-08:292770; EDB-77:061999; Izv. Akad. Nauk SSSR, Met.
 (IZNMA) (6):124-128 (Nov 1975)

SUBJECT DESCRIPTORS: ALUMINIUM: T1; BIMETALS; CHEMICAL REACTION
 KINETICS: Q1, Q2, Q3; DIFFUSION; DISSOLUTION: Q2, Q3; LIQUID
 METALS; MOLYBDENUM: T2; QUANTITY RATIO; SOLUBILITY; SOLVENT
 PROPERTIES: Q1; TUNGSTEN: T3

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-360104; 400201
 INIS CATEGORY: B22

<157>

Eagles, D.M.

Revised model for 5d-electrons in the heavy rare-earth metals
 EDE-76:065323; Z. Phys., B (ZPBBD) 21B:(1):69-78 (1975)

SUBJECT DESCRIPTORS: BAND THEORY; ELECTRONIC STRUCTURE:
 Q1; ELECTRONS; FERROMAGNETISM; RARE EARTHS: T1; STRUCTURAL
 MODELS; WAVE FUNCTIONS

ABSTRACT: A revised version of a recently published model for 5d-electrons in the ferromagnetic state of the heavy rare-earth metals is described. The model involves the broadening of local 5d-states into overlapping bands with individual widths, W . In the new approach it is assumed that the local 5d-wave functions lie at some point between those for atomic $4f$ and $5d$ configurations and those calculated for such configurations subject to the restriction that the 4f-shell is kept with its moment rigidly fixed in some given direction. The admixture of nonaligned 4f-states as in the atom lowers the local energy, but it also lowers the 5d-bandwidth due to misfit of the 4f-states which occur with and without the presence of a 5d-electron. This second effect raises the energy of the low-lying states in the band. The best local states are determined by minimizing the total electronic energy of the system, using approximations which are most suitable for 4f-shells with large excitation energies. Bandwidths are found by fitting the observed saturation magnetic moments in Gd and Tm, and satisfy W is approximately equal to 1 eV.

EDB SUBJECT CATEGORY: EDB-360102

<158>

<158>

Eckert, E.R.G.

Heat transfer bibliography EDB-76:049391; Int. J. Heat Mass
Transfer (IJHMA) 18::319-327 (Feb 1975)

SUBJECT DESCRIPTORS: BIBLIOGRAPHIES: Q1; FLUID FLOW; FLUIDIZED
FED; HEAT TRANSFER: T1; LIQUID METALS; MAGNETOHYDRODYNAMICS; MASS
TRANSFER; THERMODYNAMIC PROPERTIES

ABSTRACT: None

EDE SUBJECT CATEGORY: EDB-420400

<159>

Eder, O.J.; Kunsch, B.; Pressl, G.

Structure factor of liquid aluminum and calculation of transport
coefficients EDB-77:043630; Aluminium (Duesseldorf) (ALUMA)
52:(5):2-6 (1976)

SUBJECT DESCRIPTORS: ALUMINIUM: T1; ATOM
TRANSPORT; DIFFUSION; FOURIER TRANSFORMATION; LIQUID
METALS; MOLECULAR STRUCTURE: Q1; PAIRING ENERGY; SCATTERING; SLOW
NEUTRONS; VERY HIGH TEMPERATURE; VISCOSITY

ABSTRACT: The structure factors measured by slow neutron scatter
in liquid aluminum in the temperature range around 750^oC
are compared with the hard core structure factor. The
statistical pair distribution function for liquid aluminum
are determined by Fourier transformation, and the nearest
neighbor coordination numbers are calculated. The hard core
model is then used for approximating transport coefficients
such as the diffusion and viscosity coefficients for several
metals and metalloids. The results are compared with
experimental data. A demonstration is given how the
macroscopic transport coefficients are related to microscopic
variables.

EDE SUBJECT CATEGORY: EDB-360102

<160>

Efremov, V.A.; Trunov, V.K. (Moskovskij Gosudarstvennyj Univ.
(USSR))

Phase diagrams of $\text{Li}_2\text{MoO}_4\text{--ZnMoO}_4$, $\text{K}_2\text{MoO}_4\text{--ZnMoO}_4$ and $\text{K}_2\text{WO}_4\text{--ZnWO}_4$
systems AIX-08:298250; EDB-77:062513; Zh. Neorg. Khim. (ZNOKA)
20:(8):2200-2203 (Aug 1975)

SUBJECT DESCRIPTORS: BINARY MIXTURES; DIFFERENTIAL THERMAL

<160> CONT.

ANALYSIS;MOLTEN SALTS;PHASE DIAGRAMS:
Q1,Q2,Q3,Q4,Q5,Q6,Q7;X-RAY DIFFRACTION

ABSTRACT: Phase diagrams of $\text{Li}_2\text{MoO}_4\text{-ZnMoO}_4$, $\text{K}_2\text{MoO}_4\text{-ZnMoO}_4$, and $\text{K}_2\text{WO}_4\text{-ZnWO}_4$ systems have been plotted from the results of X-ray phase and differential thermal analyses. All the double molybdates and tungstates formed in these systems melt incongruently. General regularities as to the type of the phase diagrams of Me_2EO_4 (Me-Li, Na, K; E-Mo, W)- ZnEO_4 system are considered, as well as some features of the structure and properties of the intermediate compounds formed in these systems.

EDS SUBJECT CATEGORY: EDB-400201
INIS CATEGORY: E12

<161>

Egorov, R.F.; Kuznetsov, E.V.; Shirokovskij, V.P.; Ganin, G.V.
(AN SSSR, Sverdlovsk. Inst. Fiziki Metallov)

Electron spectrum of light alkali metals EDB-76:042257; Fiz.
Met. Metalloved. (FMMEA) 39:(2):241-249 (Feb 1975)

SUBJECT DESCRIPTORS: ALKALI METALS; T2; BRILLOUIN ZONES; CRYSTAL
FIELD; D STATES; ELECTRON DENSITY; ELECTRONIC STRUCTURE;
Q1,Q2; ENERGY SPECTRA; FERMI LEVEL; GREEN FUNCTION; LITHIUM;
T1; MUFFIN-TIN POTENTIAL; POISSON EQUATION; POTASSIUM; S
STATES; SLATER METHOD; SODIUM; TABLES; VALENCE; WAVE
FUNCTIONS; WIGNER-SEITZ METHOD

ABSTRACT: None

EDS SUBJECT CATEGORY: EDB-360104
INIS CATEGORY: B22

<162>

Ehlert, G.V.; Slovyanskikh, V.K. (AN SSSR, Moscow. Inst.
Obshchej i Neorganicheskoi Khimii)

On interaction in U--S--Te system EDB-76:048904; Zh. Neorg.
Khim. (ZNOKA) 20:(2):504-508 (Feb 1975)

SUBJECT DESCRIPTORS: CRYSTALLIZATION; DIFFERENTIAL THERMAL
ANALYSIS; PHASE DIAGRAMS: Q1,Q2; SULFUR; TELLURIUM; TELLURIUM
SULFIDES: T1; TERNARY ALLOY SYSTEMS; URANIUM; URANIUM SULFIDES;
T2; VERY HIGH TEMPERATURE; X-RAY DIFFRACTION

<162>

<162> CONT.

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-360602

INIS CATEGORY: B13

<163>

Elliott, G.R.B., Los Alamos Scientific Lab., N.Mex. (USA)

In Some considerations in establishing thermodynamic activities
from high temperature EMF measurements **

ERA-01:020203;EDB-76:054333; (1976) LA-UR--76-1070

SUBJECT DESCRIPTORS: ELECTROMOTIVE FORCE; ERRORS; MEASURING
METHODS; MOLTEN SALTS: M1; THERMODYNAMIC PROPERTIES: Q1

ABSTRACT: Various sources of uncertainty and error in
establishing thermodynamic properties from experimental data
are discussed and evaluated. These problems reflect
experimental difficulties, the frequent absence of necessary
data, some dubious assumptions in the absence of data,
certain fallacies when simple models are combined with
thermodynamic manipulations to correlate limited or imprecise
data, and some ill-chosen methods of describing data. Effects
of metal solubility in molten salts, of reactions between
electrolyte and metal, of experimental errors, and of poor
models are considered.

EDB SUBJECT CATEGORY: EDB-400201

<164>

Elovskikh, E.M.; Krylov, E.I.; Kasimov, G.G. (Ural'skiy
Politekhnicheskij Inst., Sverdlovsk (USSR))

Stability of rare earth metaniobates in hydrogen atmosphere
AIX-08:282855; EDB-77:037336; Izv. Akad. Nauk SSSR, Neorg.
Mater. (IVNMA) 11:(8):1524 (Aug 1975)

SUBJECT DESCRIPTORS: CALCINATION; CERIUM NITRATES: T2; CONTROLLED
ATMOSPHERES; ELECTRIC CONDUCTIVITY; ELECTRONIC
STRUCTURE; HYDROGEN; LANTHANUM NITRATES: T1; NEODYMIUM NITRATES:
T4; NIOBATES: T; PHASE DIAGRAMS; PRASEODYMIUM NITRATES:
T3; STABILITY: Q1, Q2, Q3, Q4; VERY HIGH TEMPERATURE; X-RAY
DIFFRACTION

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-400201

INIS CATEGORY: B12

<165>

Eren, A., Technische Univ. Darmstadt (Germany, F.R.)

In Zeeman effect and satellite carrier structure in absorption spectra of pure and doped praseodymium and europium double nitrates ** EDB-77:007299; (27 Jan 1975) N--75-27933

SUBJECT DESCRIPTORS: ABSORPTION SPECTRA; CHARGE CARRIERS; COBALT CARBIDES: T3; ELECTRONIC STRUCTURE; EUROPIUM NITRATES: T1; MIXTURES; MONOCRYSTALS; PRASEODYMIUM NITRATES: T2; ZEEMAN EFFECT: Q1, Q2, Q3

ABSTRACT: Absorption spectra of the mixed crystals $\text{Pr}_{2-x}\text{Mg}_x\text{Co}_3(\text{NO}_3)_{12}\text{H}_2\text{O}$ exhibited secondary lines from impurity ion spins. $\text{PrZn}(\text{Co})\text{-DN-}$, $\text{Pr}(\text{Eu})\text{Mg-Dn}$, and $\text{Pr}(\text{Bi})\text{Mg-DN}$ -crystals showed only minor crystal field defects. Zeeman spectra for the transformations of 7F_0 yields a, b, 5D_1 ; a, b, a 5D_2 in single EuMg-DN crystals were obtained; magnetic dipole radiation was found in the 7F_0 yields 5D_0 transfer, and temperature- and additive-dependent oscillator strength changes were observed in the transfers 7F_0 yields 5D_0 ; a, b 5D_1 ; a, b, c 5D_2 .

EDB SUBJECT CATEGORY: EDB-360602

<166>

Esin, Yu.O.; Valishev, M.G.; Gel'd, P.V. (Ural'skiy Politekhnikheskiy Inst., Sverdlovsk (USSR))

Formation enthalpy of liquid tin-vanadium alloys
AIX-08:337219; EDB-77:140520; Zh. Fiz. Khim. (ZPKHA)
51: (2):471-472 (Feb 1977)

SUBJECT DESCRIPTORS: ENTHALPY: Q1, Q2, Q3; FORMATION HEAT: Q1, Q2, Q3; LIQUID METALS: T3; QUANTITY RATIO; TIN ALLOYS: T2; VANADIUM ALLOYS: T1; VERY HIGH TEMPERATURE

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-360104
INIS CATEGORY: B22

<167>

Evans, R.; Kumaravadivel, R. (Bristol Univ. (UK). H.H. Wills Physics Lab.)

Thermodynamic perturbation theory for the surface tension and ion density profile of a liquid metal
AIX-07:262688; EDB-77:030819; J. Phys., C (London) (JPCFA)
9: (10):1891-1906 (28 May 1976)

<167>

<167> CONT.

SUBJECT DESCRIPTORS: ALUMINIUM: T1; FREE ENERGY; INTERATOMIC FORCES; ION DENSITY; IONS; LIQUID METALS; MATHEMATICAL MODELS; PERTURBATION THEORY; POTASSIUM: T2; SODIUM: T3; SURFACE TENSION: Q1, Q2, Q3

ABSTRACT: A simple scheme for determining the ion density profile and the surface tension of a liquid metal is described. Assuming that the interaction between metallic pseudoc-ions is of the form introduced by Evans, an approximate expression for the excess free energy of the system is derived using the thermodynamic perturbation theory of Weeks, Chandler and Anderson. This excess free energy is then minimized with respect to a parameter which specifies the ion density profile, and the surface tension is given directly. From a consideration of the dependence of the interionic forces on the electron density it is predicted that the ions should take up a very steep density profile at the liquid metal surface. This behaviour is contrasted with that to be expected for rare-gas fluids in which the interatomic forces are density-independent. The values of the surface tension calculated for liquid Na, K and Al from a simplified version of the theory are in reasonable agreement with experiment.

EEB SUBJECT CATEGORY: EDB-360104
INIS CATEGORY: B22

<168>

Evdokimov, A.A.; Trunov, V.K. (Moskovskij Gosudarstvennyj Univ. (USSR). Kafedra Neorganicheskoy Khimii)

Phase relations in $\text{PbWO}_4\text{-TR}_{2(\text{WO}_4)_3}$ systems AIX-08:336916; EDB-77:146841; Zh. Neorg. Khim. (ZNCKA) 21: (1):200-203 (Jan 1976)

SUBJECT DESCRIPTORS: DIFFERENTIAL THERMAL ANALYSIS; LATTICE PARAMETERS; LEAD COMPOUNDS: M4; LITHIUM COMPOUNDS; PHASE DIAGRAMS: Q1, Q3, Q4; PRASEODYMIUM COMPOUNDS; PYROLYSIS; RARE EARTH COMPOUNDS: M1; TEMPERATURE DEPENDENCE; TERBIUM COMPOUNDS; TUNGSTATES: M3; X-RAY DIFFRACTION

ABSTRACT: Phase ratios in the systems $\text{PbWO}_4\text{-TR}_{2(\text{WO}_4)_3}$ ($\text{TR}=\text{Pr-Lu}$) have been studied and phase diagrams of the systems $\text{PbWO}_4\text{-Pr}_{2(\text{WO}_4)_3}$ and $\text{PbWO}_4\text{-Tb}_{2(\text{WO}_4)_3}$ have been plotted using the methods of X-ray phase and differential-thermal analysis. The study of the systems $\text{PbWO}_4\text{-TR}_{2(\text{WO}_4)_3}$ has shown that in the series of rare-earth elements the phase ratios are close in the following three groups of elements: $\text{TR}=\text{La}$; $\text{TR}/\text{Pr}=\text{Pr-Eu}$; $\text{TR}=\text{d-Lu}$. Between the components with the structure of

<168> CONT.

scheelite and tungstate of europium the intermediate phases are formed with wide homogeneity regions. The fact that the phase $\text{Pb}_{0.5}\text{La}_3(\text{WO}_4)_5$ is isostructural to the phase $\text{LiLa}_3(\text{WO}_4)_5$ has been proved by the monocrystal study. In the systems $\text{PbWO}_4\text{-TR}$ $\text{Pb}_{0.5}\text{TR}_3(\text{WO}_4)_5$ (TR=Pr-Eu) highly temperature intermediate phases are isostructural to $\text{BaLa}_4(\text{WO}_4)_7$ since the lines on the X-ray patterns of these samples are satisfactorily induced in the assumption of the cell $\text{BaLa}_4(\text{WO}_4)_7$. The compounds of the composition $\text{PbTR}_2(\text{WO}_4)_4$ are formed in the systems $\text{PbWO}_4\text{-TR}$ $\text{Pb}_{0.5}\text{TR}_2(\text{WO}_4)_4$ (TR=Pr-Lu). They all crystallize in one structural type, have no noticeable homogeneity regions, and decompose irreversibly in a solid phase with increasing temperature. The decomposition temperature for the compounds $\text{PbTR}_2(\text{WO}_4)_4$ is given as a function of the order number. Interplanar spacings in the sample of the composition $\text{PbTb}_2(\text{WO}_4)_4$ are presented.

EDB SUBJECT CATEGORY: EDB-400201

INIS CATEGORY: B12

<169>

Ewart, F.T.; Taylor, R.G. (UKAEA Research Group, Harwell. Atomic Energy Research Establishment); Horspool, J.M.; James, G. (UKAEA Reactor Group, Sellafield)

Chemical effects of composition changes in irradiated oxide fuel materials. II. Fission product segregation and chemical equilibria AIX-07:279173; ERA-02:019367; EDB-77:034973; J. Nucl. Mater. (JNUMA) 61:(3):254-270 (Sep 1976)

SUBJECT DESCRIPTORS: CHEMICAL COMPOSITION; CHEMICAL RADIATION EFFECTS: Q1, Q2; CORROSION; ELECTRON MICROPROBE ANALYSIS; FISSION PRODUCTS; FUEL CANS; FUEL PINS; FUEL-CLADDING INTERACTIONS: Q1, Q2; IRRADIATION; NEUTRON FLUX; PLUTONIUM OXIDES: T1; URANIUM OXIDES: T2

ABSTRACT: Four miniature fuel pins containing uranium/plutonium oxide, covering a wide range of oxygen-to-metal ratios, have been irradiated in a thermal reactor. These pins were subjected to a detailed examination, by electron probe microanalysis and ceramography. The results of this analysis have been interpreted in the light of the results of a computer model of the pin behaviour. The widely different fuel chemistry in each of the fuel pins produced different fission-product phases and different clad corrosion phenomena. The phases found are in general agreement with the model which predicts increasing oxygen-to-metal ratio with increasing burn-up and oxygen redistribution in the

<169>

<169> CCNT.

temperature gradient. Clad corrosion is seen to be inhibited at low oxygen potentials but the distinction between intergranular and 'broad front' corrosion is not attributable simply to the local oxygen potential.

EEB SUBJECT CATEGORY: EDB-050700;220300;360206;400702
INIS CATEGORY: E23

<170>

Ezhov, V.K.

Some properties of hydrogen fluoride compounds with group 5 metal pentafluorides AIX-08:282857;EDB-77:037408; Zh. Neorg. Khim. (ZNOKA) 21:(8):2263-2264 (Aug 1976)

SUBJECT DESCRIPTORS: COMPLEXES;HYDRIODIC ACID;NIOBIUM FLUORIDES: T1;TANTALUM FLUORIDES: T2;TEMPERATURE DEPENDENCE;VAPOR PRESSURE: Q1,Q2

ABSTRACT: None

EEB SUBJECT CATEGORY: EDB-400201
INIS CATEGORY: B12

<171>

Fahey, J.A. (Bronx Community Coll., New York (US)); Turcotte, R.P.: Chikalla, T.D.

Decomposition, stoichiometry and structure of neptunium oxides AIX-07:241900;EDB-76:092570; J. Inorg. Nucl. Chem. (JINCA) 38:(3):495-500 (1976)

ABSTRACT: The neptunium-oxygen system for $2.00 < O/Np < 2.7$ was studied between room temperature and 900°C and under controlled oxygen pressure. Thermogravimetry, gas manometry and X-ray diffraction were applied to decomposition of Np(V) and Np(VI) hydroxides and Np(V) nitrate. The oxide region of apparent stoichiometry $2.50 < O/Np < 2.67$ was emphasized, and evidence is presented showing that the only higher oxide in this region is monoclinic Np_2O_5 . Decomposition to the dioxide is achieved by heating to above 700°C , for which the O_2 pressure dependence is given. Oxide having the Np(VI) hydroxide as a precursor is shown to decompose through an additional intermediate near Np_4O_9 .

EEB SUBJECT CATEGORY: EDB-360202
INIS CATEGORY: B13

<172>

Fedorchenko, V.I.; Averin, V.V.

Kinetic peculiarities of nitrogen dissolving in iron-chromium and iron-chromium-nickel melts EDB-76:036457; Izv. Akad. Nauk SSSR, Met. (IZNMA) (3):57-63 (May 1974)

SUBJECT DESCRIPTORS: CHROMIUM ALLOYS: T3; IRON ALLOYS: T2; LIQUID METALS; NICKEL ALLOYS: T4; SOLVENT PROPERTIES: Q1, Q2, Q3, Q4

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-360104

INIS CATEGORY: B21

<173>

Fedorov, N.F.; Mel'nikova, O.V.; Saltykova, V.A. (Leningradskij Tekhnologicheskij Inst. (USSR))

Physicochemical investigation of rare earth titanoniobates, LnTiNbO_6 AIX-08:320614; EDB-77:113937; Izv. Akad. Nauk SSSR, Neorg. Mater. (IVNMA) 12:(4):773-774 (Apr 1976)

SUBJECT DESCRIPTORS: CRYSTAL STRUCTURE: Q1, Q2, Q3; CRYSTALLIZATION; DENSITY; LATTICE PARAMETERS; MELTING POINTS: Q1, Q2, Q3; NIOBATES: T1; RARE EARTH COMPOUNDS: T2; REFRACTIVITY: Q1, Q2, Q3; TITANIUM COMPOUNDS: T3; X-RAY DIFFRACTION

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-360603; 400201

INIS CATEGORY: B12

<174>

Fedorov, P.I.; Malova, N.S.; Petrova, G.K.; Vorob'eva, G.V. (Moskovskij Inst. Tonkoj Khimicheskoy Tekhnologii (USSR))

Interaction of lower indium chlorides with chloride and chloroindates of potassium EDB-76:048906; Zh. Neorg. Khim. (ZNOKA) 20:(3):637-640 (Mar 1975)

SUBJECT DESCRIPTORS: CHEMICAL REACTIONS; INDIUM CHLORIDES: T1; PHASE DIAGRAMS: Q1, Q2; POTASSIUM CHLORIDES: T2; SOLID SOLUTIONS

<174>

<175> CONT.

ABSTRACT: None

EDE SUBJECT CATEGORY: EDB-360602

INIS CATEGORY: B12

<175>

Fee, D.C.; Johnson, I.; Davis, S.A.; Shinn, W.A.; Staahl, G.E.;
Johnson, C.E., Argonne National Lab., Ill. (USA)

In Cs--U--O phase diagram and its application to
uranium--plutonium oxide fast reactor fuel pins **
NTS-78:001713; ERA-03:011718; INS-78:002138; EDB-78:025371; (Aug
1977) ANL--76-126

SUBJECT DESCRIPTORS: BREEDING

BLANKETS; CESIUM; DEFORMATION; DIFFUSION; FAST REACTORS; FISSION
PRODUCTS; FUEL CANS: T1; FUEL PINS; HIGH
TEMPERATURE; OXYGEN; PHASE DIAGRAMS; PLUTONIUM DIOXIDE:
T3; STABILITY; THERMODYNAMIC PROPERTIES:
Q2, Q3, Q1; URANIUM; URANIUM DIOXIDE: T2; VERY HIGH TEMPERATURE

ABSTRACT: Portions of the cesium-uranium-oxygen system have been
investigated between 873 and 1273\$sup 0\$K and a phase diagram
has been constructed using our data and the data of other
workers in the field. Thermodynamic and kinetic data have
been used to examine the reactions that occur in fast-reactor
fuel pins between fission-product cesium and the uranium
oxide blanket. It was concluded that at the low oxygen
potentials existing at the interface between the
uranium-plutonium mixed-oxide and the uranium oxide blanket,
Cs\$sub 2\$UO\$sub 4\$ is the only Cs-U-O compound expected to be
formed in the uranium oxide blanket.

EDE SUBJECT CATEGORY: EDB-050700;360202

INIS CATEGORY: B23

<176>

Ferris, L.M. ** McNeese, L.E., Oak Ridge National Lab., Tenn.
(USA)

Chemistry In Molten-salt reactor program. Semiannual progress
report for period ending February 29, 1976 **
INS-77:003086; ERA-02:015877; EDB-77:029145; (Aug 1976)
CRNL--5132

SUBJECT DESCRIPTORS: BERYLLIUM FLUORIDES; BORON COMPOUNDS:

A4; BORO FLUORIDES: A6; CHEMISTRY: Q1, Q8; CHROMIUM
COMPOUNDS; DECOMPOSITION: Q2; DENSITY:
Q6; ELECTROCHEMISTRY; EQUILIBRIUM; FLUORINE COMPOUNDS:
A5; HYDROGEN; HYDROLYSIS: Q4, Q5; LITHIUM FLUORIDES; LITHIUM

<176> CONT.

TELLURIDES: A2;MOLTEN SALT REACTORS: A1;MOLTEN SALTS:
A8;OXYGEN COMPOUNDS;REDUCTION: Q3;SODIUM
COMPOUNDS;SOLUBILITY: Q2;TELLURIDES;TELLURIUM
COMPOUNDS;THORIUM FLUORIDES;TRITIUM;URANIUM COMPOUNDS:
A7;URANIUM TETRAFLUORIDE: A3;VALENCE: Q7;VAPORS;ZIRCONIUM
FLUORIDES

ABSTRACT: The chemistry of Te in molten salts was studied (solubility of Li_2Te , Te species in LiF--BeF_2 , decomposition pressure of LiTe_3 , $\text{UF}_4 + \text{H}_2$ reversible $\text{UF}_3 + \text{HF}$ equilibrium in molten fluorides). Work on coolant-salt chemistry was continued (hydrolysis of Na_3BF_6 , vapor density in $\text{BF}_3\text{--H}_2$). Monitoring of the $\text{U}^{4+}/\text{U}^{3+}$ ratio in loops and creep-test machines was continued. Data were obtained during the third tritium injection experiment at the Coolant-Salt Technology Facility. Electrochemical studies indicated that LiTe_3 and Cr_3Te_4 are insoluble in MSBR fuel salt. The species $\text{Te}^{n-1}/\text{Te}^n$ and oxide species were studied. (DLC)

EDB SUBJECT CATEGORY: EDB-050700;400702;210500;400201
INIS CATEGORY: B13;E35;B12

<177>

Filippov, E.S. (Moskovskij Inst. Stali i Splavov (USSR))

Discrete variation of the structure and texture of liquid metals formed by components with discrete packing
AIX-08:289572;EDB-77:043637; Izv. Vyssh. Uchebn. Zaved., Chern. Metall. (IVUMA) (3):125-132 (1976)

SUBJECT DESCRIPTORS: BCC LATTICES;CRYSTAL STRUCTURE:
Q1,Q2,Q3,Q4;DENSITY: Q1,Q2,Q3,Q4;EUTECTICS;FCC
LATTICES;INDIUM: T1;INDIUM ALLOYS: T2;LIQUID METALS;MELTING
POINTS;PHASE DIAGRAMS: Q2,Q4;QUANTITY RATIO;TEMPERATURE
DEPENDENCE;TEXTURE;TIN: T3;TIN ALLOYS: T4

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-360102
INIS CATEGORY: B22

<178>

<178>

Finch, A.; Gates, P.N.; Peake, S.J. (Royal Holloway Coll., London (UK))

Thermochemistry of polyhalides. Pt.1. Caesium and rubidium dichloroiodates(I) AIX-08:305594; EDB-77:133353; J. Chem. Soc. (London), Dalton Trans. (JCDTB) (4):397-400 (1977)

SUBJECT DESCRIPTORS: CESIUM CHLORIDES: T1; CESIUM IODIDES: T2; CRYSTALS; FORMATION HEAT: Q1, Q2, Q3, Q4; RUBIDIUM CHLORIDES: T3; RUBIDIUM IODIDES: T4

ABSTRACT: The standard enthalpies of formation of crystalline $\text{Cs}[\text{ICl}_2]$ and $\text{Rb}[\text{ICl}_2]$ have been determined as -511.1 ± 4.1 and -491.4 ± 4.1 kJ mol⁻¹ respectively, based on enthalpies of reaction with aqueous silver(I) nitrate. The results are discussed in relation to various thermodynamic parameters.

EDB SUBJECT CATEGORY: EDB-400201
INIS CATEGORY: B12

<179>

Fischer, D.F.; Leibowitz, L. (Argonne National Lab., Ill. (USA))

Enthalpy of uranium-plutonium carbide from 298K to the melting point AIX-08:340257; EDB-78:016962; J. Nucl. Mater. (JNUMA) 67: (3):244-248 (Aug 1977)

SUBJECT DESCRIPTORS: CALORIMETERS; ENTHALPY: Q1, Q2; HIGH TEMPERATURE; MEDIUM TEMPERATURE; NUCLEAR FUELS; PLUTONIUM CARBIDES: T2; PYROMETERS; SPECIFIC HEAT; URANIUM CARBIDES: T1; VERY HIGH TEMPERATURE

ABSTRACT: An induction-heated drop-calorimetric system was used to measure the enthalpy of $(\text{U}_{0.8}\text{Pu}_{0.2})\text{C}$ from 1039 to 2421 K. By combining the measured enthalpy values with previously estimated low-temperature enthalpy data (900 K and below), an equation was derived that is applicable over the temperature range from 298 to 2548 K. Thermodynamic functions of $(\text{U}_{0.8}\text{Pu}_{0.2})\text{C}$ as a function of temperature are also given.

EDB SUBJECT CATEGORY: EDB-360204
INIS CATEGORY: B23

<180>

Fischer, E.A. (Kernforschungszentrum Karlsruhe (F.R. Germany).
Inst. fuer Angewandte Systemanalyse); Kinsman, P.R.; Ohse,
R.W. (Commission of the European Communities, Karlsruhe (F.R.
Germany). European Inst. for Transuranium Elements)

Critical assessment of equation of state data for UO_2
AIX-07:243897; ERA-02:007618; EDB-77:003016; J. Nucl. Mater.
(JNUMA) 59:(2):125-136 (Feb 1976)

SUBJECT DESCRIPTORS: CRITICAL PRESSURE; CRITICAL
TEMPERATURE; ENTHALPY; EQUATIONS OF STATE: Q1; FREE
ENERGY; GASES; LIQUIDS; SOLIDS; URANIUM DIOXIDE: T1; VAPOR PRESSURE

ABSTRACT: A new equation of state for UO_2 has been derived using the theory of significant liquid structures, which predicts a critical temperature of 7560K, a critical pressure of 1210 atm, and a critical density of 1.66 g/cm³. The resulting critical compressibility of 0.316 is in good agreement with the values predicted by the theory of corresponding states. Consistency between literature data on enthalpy of solid UO_2 , recent vapour pressure measurements over liquid UO_2 , and the pressure data over solid UO_2 could only be obtained by the inclusion of an electronic excitation term in the gaseous partition function. A new set of free energy functions for gaseous and solid UO_2 is presented.

EDE SUBJECT CATEGORY: EDB-360202; 050700
INIS CATEGORY: E12

<181>

Fisher, S.B. (Central Electricity Generating Board, Berkeley
(UK). Berkeley Nuclear Labs.)

Atomic diffusion in alloys exposed to liquid sodium
AIX-08:295720; EDB-77:055608; J. Nucl. Mater. (JNUMA)
62:(2-3):247-256 (Nov 1976)

ABSTRACT: Reasons for the observed enhancement of atomic diffusion coefficient in alloys exposed to sodium are explored. The diffusion coefficient in the presence of a chemical potential gradient is derived from a simple atomistic model. It is concluded that two mechanisms could contribute to the enhancement. (a) A non-zero gradient of partial molar enthalpy. (b) An increase in the atomic mobility due to the presence of high diffusivity paths. It is shown that the effect of (a) is not of sufficient magnitude to account for the increase in coefficient. However the results can be analysed successfully in terms of grain boundary diffusion and the coefficients are determined for iron in Ni/Cr ($1.2 \times 10^{-9} \text{ cm}^2/\text{s}$) at $650 \pm 0^\circ\text{C}$ and

<181>

<181> CONT.

nickel in iron ($5 \times 10^{-8} \text{ cm}^2/\text{s}$) at 730°C . The total absorbed diffusant is also calculated for the case of iron into Ni/Cr, and a general expression for the absorbed diffusant is given.

EDB SUBJECT CATEGORY: EDB-360105;360104
INIS CATEGORY: B22

<182>

Fitzner, K.; Jacob, K.T. (Toronto Univ., Ontario (Canada). Dept. of Metallurgy and Materials Science)

Solubility and activity of oxygen in liquid indium and copper-indium alloys AIX-08:324210;EDB-77:132606; J. Less-Common Met. (JCOMA) 52:(2):279-291 (Apr 1977)

SUBJECT DESCRIPTORS: COPPER ALLOYS: T1;ELECTROMOTIVE FORCE;FORMATION FREE ENERGY;HIGH TEMPERATURE;INDIUM: T2;INDIUM ALLOYS: T3;INDIUM OXIDES;LIQUID METALS;OXYGEN: T4;PHASE STUDIES;SOLUBILITY: Q1,Q2,Q3,Q4;SOLUTION HEAT;TEMPERATURE DEPENDENCE;VERY HIGH TEMPERATURE

ABSTRACT: The solubility of oxygen in liquid indium in the temperature range $650-820^\circ \text{C}$ and in liquid copper-indium alloys at 1100°C in equilibrium with indium sesquioxide has been measured by a phase equilibration technique. The solubility of oxygen in pure indium is given by the relation $\log(\text{at.}\%) = -4726/T + 3.73 (+0.08)$. Using the recently measured values for the standard free energy of formation of In_2O_3 and assuming that oxygen obeys Sievert's law up to saturation, the standard free energy of solution of molecular oxygen in liquid indium is calculated. The effect of indium additions on the activity coefficient of oxygen dissolved in liquid copper was measured by a solid oxide galvanic cell. The experimentally determined variation of the activity coefficient of oxygen in dilute solution in Cu-In alloys is in fair agreement with that predicted by a quasichemical model in which each oxygen atom is assumed to be interstitially coordinated to four metal atoms and the nearest neighbour metal atoms are assumed to lose approximately half their metallic cohesive energies.

EDB SUBJECT CATEGORY: EDB-360104
INIS CATEGORY: B22

<183>

Fleury, P.A.; Lyons, K.B. ** Lucovsky, G. (ed.)

Anomalous quasielastic light scattering from metal oxide glasses
 In Structure and excitations of amorphous solids **
 INS-77:000264; ERA-02:010523; EDB-77:007242; (1976) American
 Inst. of Physics

SUBJECT DESCRIPTORS: ANOMALOUS DIMENSION; BORON OXIDES:
 T3; EXCITATION: Q1, Q2, Q3, Q4, Q5, Q6; GLASS; LANTHANUM OXIDES:
 T6; LIGHT SCATTERING: Q1, Q2, Q3, Q4, Q5, Q6; NIOBIUM OXIDES:
 T4; OXIDES; RARE EARTHS; TANTALUM OXIDES: T2; TERBIUM OXIDES:
 T1; THORIUM OXIDES: T5; TRANSITION ELEMENTS

ABSTRACT: An anomalous quasielastic component has been observed in the light scattering spectra of several heavy metal oxide glasses. This component consists of a depolarized central peak of frequency width less than approximately 0.5 cm^{-1} , depending on temperature and composition. Observations are reported for eight different compositions, all containing either transition metal or rare earth oxides. The integrated intensity for glasses whose major constituent is ThO_2 correlates well with Tb concentration, while the linewidth is insensitive to Tb concentration as well as host composition.

EDS SUBJECT CATEGORY: EDR-360204
 INIS CATEGORY: B23

<184>

Flotow, H.E.; Osborne, D.W.; Fried, S.M.; Malm, J.G.

Heat capacity of PuO_2 from 12 to 350°K and of PuO_2 from 4 to 25°K . Entropy, enthalpy, and Gibbs energy of formation of PuO_2 at 298.15°K INS-76:019939; ERA-02:001078; EDB-76:081660; J. Chem. Phys. (JCPSA) 65: (3):1124-1129 (1 Aug 1976)

SUBJECT DESCRIPTORS: ENTHALPY: Q1; ENTROPY: Q1; FORMATION FREE ENERGY: Q1; LOW TEMPERATURE; MEDIUM TEMPERATURE; PLUTONIUM 242; PLUTONIUM 244; PLUTONIUM DIOXIDE: T1; SPECIFIC HEAT: Q1; VERY LOW TEMPERATURE

ABSTRACT: Heat capacity measurements have been made on a 22.5 g sample of PuO_2 from 12 to 350°K and on a 3.6 g sample of PuO_2 from 4 to 25°K . The heat capacity curve has the normal sigmoid shape. At 298.15°K the heat capacity C_p , entropy S , enthalpy $H - H_0$, and tempered Gibbs energy $(G - H_0)/T$ are, respectively, $(66.25 \pm 0.26) \text{ J } ^\circ\text{K}^{-1} \text{ mole}^{-1}$, $(66.13 \pm 0.26) \text{ J } ^\circ\text{K}^{-1} \text{ mole}^{-1}$, $(10784 \pm 43) \text{ J } ^\circ\text{K}^{-1} \text{ mole}^{-1}$, $(10784 \pm 43) \text{ J } ^\circ\text{K}^{-1} \text{ mole}^{-1}$.

<184>

<184> CONT.

and $-(29.96 \pm 0.12)$ J degreeK⁻¹xmole⁻¹. This value of the entropy is significantly lower than the one previously derived from heat capacity measurements on PuO_2 . At 298.15degreeK the standard entropy of formation and the standard Gibbs energy of formation are $-(195.2 \pm 0.4)$ J degreeK⁻¹xmole⁻¹ and $-(998.0 \pm 0.7)$ kJ mole⁻¹, respectively. (AIP)

EDB SUBJECT CATEGORY: EDB-360204

INIS CATEGORY: E23

<185>

Fomichev, V.V.; Alikeev, B.D.; Reznik, E.M.; Ivanova, M.M.; Petrov, K.I. (Moskovskij Inst. Tonkoj Khimicheskoy Tekhnologii (USSR))

Vibration spectra of rare earth oxytungstates- Ln_2WO_6 composition AIX-08:295289;EDB-77:069089; Zh. Neorg. Khim. (ZNCKA) 21:(7):1733-1737 (Jul 1976)

SUBJECT DESCRIPTORS: CRYSTAL STRUCTURE;FREQUENCY MEASUREMENT;FREQUENCY RANGE;INFRARED SPECTRA;OXYGEN COMPOUNDS;RARE EARTH COMPOUNDS: T1;TABLES;TUNGSTATES: T2;VIBRATIONAL STATES: Q1,Q2

ABSTRACT: None

EIP SUBJECT CATEGORY: EDB-400201

INIS CATEGORY: B12

<186>

Forney, J.J. (Ecole Polytechnique Federale, Lusanne, Switzerland); Quattropani, A.; Bassani, F. ** Pilkuhn, M.H. (ed.)

Structure of the biexciton molecule and optical transition selection rules in zincblende and wurtzite materials In Physics of semiconductors ** ERA-01:016804;EDB-76:042565; (1974) B. G. Teubner

SUBJECT DESCRIPTORS: BINDING ENERGY: Q1,Q2;CADMIUM SULFIDES: T2;COPPER CHLORIDES: T1;ELECTRON EXCHANGE;HOLES;SELECTION RULES

ABSTRACT: The states of the biexciton molecules are related to the band structure of the crystal and effects due to

<186> CONT.

electron-hole exchange are taken into account. For biexcitons in CdS and CuCl optical selection rules are derived and the electron hole exchange corrections to the energy is computed.

EDB SUBJECT CATEGORY: EDB-400201

<187>

Foster, M.S.; McCloud, G.H.; Cairns, E.J.

Electrochemical studies of the sodium--bismuth system

EDE-77:043222; Am. Chem. Soc., Div. Fuel Chem., Prepr. (ACFPA) 11:(1):276-281 (1967)

SUBJECT DESCRIPTORS: BISMUTH BASE ALLOYS; ELECTRIC POTENTIAL: Q1; FREE ENERGY; METAL-METAL BATTERIES: M1; SODIUM; SODIUM ALLOYS; SODIUM CHLORIDES

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-250901

<188>

Fournier, J.M. (CEA Centre d'Etudes Nucleaires de Grenoble, 38 (France). Dept. de Recherche Fondamentale)

Bonding and the electronic structure of the actinide metals

AIX-07:236376; EDB-76:092474; J. Phys. Chem. Solids (JPCSA) 37:(2):235-244 (1976)

SUBJECT DESCRIPTORS: ACTINIDES:

M1; ACTINIUM; AMERICIUM; BERKELIUM; BINDING ENERGY; CALIFORNIUM; CURIUM; EINSTEINIUM; ELECTRONIC STRUCTURE: Q1; FERMIUM; LAWRENCIUM; MELTING POINTS; MENDELEVIUM; NEPTUNIUM; NOBELIUM; PLUTONIUM; PROTACTINIUM; SUBLIMATION HEAT; THORIUM; URANIUM

ABSTRACT: The electronic configurations of the 15 actinide metals from actinium to lawrencium have been determined using a phenomenological model based on the metallic radius, the crystal structure, the melting temperature and the enthalpy of sublimation. The above properties can be related to the electronic configuration which accounts for their anomalies. It can be shown that transberkelium metals are divalent and that from protactinium to plutonium, 5f electrons are band

<189>

<189> CONT.

electrons. The hitherto unknown values of the enthalpies of sublimation and melting temperatures have been calculated.

EDB SUBJECT CATEGORY: EDB-360104

INIS CATEGORY: P22

<189>

Freeman, A.J. (Northwestern Univ., Evanston, Ill. (USA). Dept. of Physics); Koelling, D.D. (Argonne National Lab., Ill. (USA))

Electronic structure and properties of the actinides

AIX-08:332145; ERA-03:012847; EDB-78:021808; Physica, B, C (PHBCD) 86-88::16-25 (1977)

SUBJECT DESCRIPTORS: ACTINIDE ALLOYS: T1; ACTINIDES: T2; BAND THEORY; CORRELATIONS; COULOMB FIELD; DE HAAS-VAN ALPHEN EFFECT; ELECTRONIC STRUCTURE: Q1, Q2; F STATES; FERMI LEVEL; INTERMETALLIC COMPOUNDS; MAGNETIC PROPERTIES: Q1, Q2; REVIEWS; WAVE FUNCTIONS

ABSTRACT: The authors review the present state of understanding of the electronic structure and physical properties of actinide metals and intermetallic compounds as derived from relativistic APW energy band studies of some of the light (Th, U, Np and Pu) and heavy (Am, Bk and Cm) metals, the intermetallics $\text{URh}_{1/3}$ and $\text{UIr}_{1/3}$ and (NaCl structure compounds such as) UC. Emphasis is placed on the importance of Coulomb correlation and the role of actinide-actinide separation in determining the itinerancy (as in the light metals) or localization (as in the heavy metals) of the 5f electrons and in turn their resulting magnetic and other properties. The $\text{UX}_{1/3}$ systems considered are significant because the uranium sites are sufficiently separated to fall in the local moment range of the Hill superconductivity/magnetism plot but show no magnetic ordering. Comparisons to recent optical, de Haas-van Alphen and other data are given when available. The contrasting cases of the transition and rare-earth systems make it clear that the 5f electrons are a unique species which offer exciting challenges to both experimentalists and theorists.

EDB SUBJECT CATEGORY: EDB-360104

INIS CATEGORY: B22

<190>

Freyland, W.; Steinleitner, G., Marburg Univ. (Germany, F.R.).
Fachbereich Physikalische Chemie

In Magnetic susceptibility and electronic structure in metallic and non-metallic liquid cesium-gold and cesium-antimony alloys ** AIX-07:263989;EDB-77:036694; (1976)
AED-Conf--76-087-013

SUBJECT DESCRIPTORS: ANTIMONY ALLOYS: T3;BINARY ALLOY SYSTEMS;CESIUM ALLOYS: T1;CHEMICAL COMPOSITION;COMPARATIVE EVALUATIONS;DIAGRAMS;ELECTRONIC STRUCTURE;EQUATIONS;GOLD ALLOYS: T2;INTERMETALLIC COMPOUNDS;LIQUID METALS;MAGNETIC SUSCEPTIBILITY: Q1,Q2,Q3;METALS;NONMETALS;QUANTITY RATIO;TEMPERATURE DEPENDENCE

ABSTRACT: Measurements of the concentration and temperature dependence of the magnetic susceptibility κ of liquid Cs-Au and Cs-Sb alloys are reported. At the stoichiometric compositions CsAu and Cs₃Sb, respectively, κ exhibits a pronounced diamagnetic minimum of -81×10^{-6} emu x mole⁻¹ for CsAu and -85×10^{-6} emu x mole⁻¹ for Cs₃Sb. It is shown that magnetic susceptibility provides a good means for the study of chemical bonding and electronic structure in these liquids. In the nonmetallic electrical transport regime the data are discussed in terms of various models for the electronic structure of metal atoms dissolved in a more ionic or more semiconducting liquid alloy. In the case of Cs-Au alloys a complete description of the change of the nature of electronic states from nonmetallic to metallic behaviour can be deduced from the measured κ -values. The temperature dependence of κ for Cs-Sb alloys near the composition Cs₃Sb gives indication for a transport mechanism typical of liquid semiconductors.

EDB SUBJECT CATEGORY: EDB-360104
INIS CATEGORY: A13

<191>

Friederich, E.; Sittig, L.

In Production and properties of high-melting lower oxides ** ERA-02:060439;EDB-77:140654; Z. Anorg. Allg. Chem. (ZAACA) 145: (1/2):127-140 (1925) UCRL-Trans--11251

SUBJECT DESCRIPTORS: CERIUM OXIDES: T5;CHEMICAL PREPARATION: Q1,Q2,Q3,Q4,Q5,Q6,Q7,Q8;COLOR: Q1,Q2,Q3,Q4,Q5,Q6,Q7,Q8;ELECTRIC CONDUCTIVITY: Q1,Q3,Q4,Q5,Q6,Q7,Q8;HARDNESS: Q1,Q3,Q4,Q5,Q6;MELTING POINTS: Q1,Q3,Q4,Q5,Q6,Q8;MOLYBDENUM OXIDES: T7;NIOBIUM OXIDES: T4;TITANIUM OXIDES: T1;TUNGSTEN OXIDES: T6;URANIUM OXIDES:

<191>

<191> CONT.

T8;VANADIUM OXIDES: T3;ZIRCONIUM OXIDES: T2

ABSTRACT: A number of lower oxides were produced, and their properties, such as color, hardness, melting point, and electrical conductivity, were investigated. No lower oxides with a specific formula could be obtained from the elements titanium and zirconium by reduction of the saturated oxides TiO_2 and ZrO_2 with hydrogen or carbon. A dark blue, current-conducting product with a maximum weight increase of 7.7% on calcining in air was produced by annealing TiO_2 in hydrogen. ZrO_2 was not reduced by hydrogen, but with coal it could be reduced to a bluish-white product which was far removed from the zirconium content of Zr_2O_3 and formed a carbide when heated further with an excess of coal. It oxidized to white ZrO_2 in the presence of concentrated sulfuric oxide. The lower vanadium oxide V_2O_3 was formed by heating V_2O_5 in hydrogen; V_2O_4 was obtained by calcining V_2O_5 in nitrogen with the amount of coal necessary for reduction. Nb_2O_3 and Ce_2O_3 were obtained by reduction of Nb_2O_5 and CeO_2 with hydrogen. Ta_2O_5 could not be reduced to lower oxides with hydrogen or carbon. It is doubtful whether lower tantalum oxides with the formulas Ta_2O_4 and Ta_2O_3 can be produced at all. The tungsten and molybdenum oxides WO_2 , W_2O_5 , MoO_2 , and Mo_2O_5 were obtained by reduction of the corresponding acids WO_3 and MoO_3 with coal or tungsten and molybdenum metal in a stream of nitrogen. The blue uranium oxide UO_2 was produced by annealing U_3O_8 in a nitrogen stream, and the brown modification of UO_2 was obtained in a hydrogen stream. U_3O_8 itself was obtained by calcining UO_3 in air. The resistance of all these oxides had a negative temperature coefficient. (RWR)

EDB SUBJECT CATEGORY: EDB-360201;360204

<192>

Fromm, E.; Jehn, H.; Hoerz, G. (Max-Planck-Institut fuer Metallforschung, Stuttgart (Germany, F.R.). Inst. fuer Werkstoffwissenschaften)

In Gases and carbon in metals (thermodynamics, kinetics, and properties). I. Alkali metals, alkaline earth metals, light metals (Li, Na, K, Rb, Cs; Ca, Sr, Ba; Be, Mg, Al) ** Physik Daten/Physics Data, AIX-08:298559;EDB-77:068253; (1976) ZAED

SUBJECT DESCRIPTORS: ALKALI METALS: T1;ALKALINE EARTH METALS: T2;DATA: Q1,Q2,Q3;GASES;METALS: T3

<192> CONT.

ABSTRACT: This issue is part of a series of data on 'Gases and Carbon in Metals' which supplements the data compilation in the book 'Gase und Kohlenstoff in Metallen' (Gases and Carbon in Metals), edited by E. Fromm and E. Gebhardt, Springer-Verlag, Berlin 1976. The present survey includes results from papers published after the copy deadline and recommends critically selected data. Furthermore it comprises a bibliography of relevant literature. Each part of this series refers to metals belonging to one or several groups of the periodic table of elements and compiles data and references on the interaction of the metals with hydrogen, carbon, nitrogen, and oxygen.

EDE SUBJECT CATEGORY: EDB-360100

INIS CATEGORY: B22

<193>

Fromm, E.; Kirchheim, R. (Max-Planck-Institut fuer Metallforschung, Stuttgart (F.R. Germany). Inst. fuer Werkstoffwissenschaften)

Electronic conductivity and polarization effects of solid $\text{ThO}_{0.2}\text{-Y}_{0.2}\text{O}_{0.3}$ electrolytes and of oxygen containing Va-metal electrodes AIX-07:262512; EDB-77:019311; Z. Metallkd. (ZEMTA) 66:(4):196-202 (Apr 1975)

SUBJECT DESCRIPTORS: DOPED MATERIALS; ELECTRIC CONDUCTIVITY: Q1, Q2, Q3; ELECTRODES; ELECTROLYTES; ELECTROMOTIVE FORCE; ELECTRON TRANSFER; INTERSTITIALS; MASS TRANSFER; OXYGEN ADDITIONS; POLARIZATION; QUANTITY RATIO; SOLIDS; SURFACE PROPERTIES; TANTALUM; TEMPERATURE DEPENDENCE; THORIUM OXIDES: T1; TIME DEPENDENCE; VANADIUM: T3; YTTRIUM OXIDES: T2

ABSTRACT: An apparatus for e.m.f.-measurements, using a solid state electrolyte, is described and results of galvanostatic measurements are presented. Electrodes of Va-metals with interstitially dissolved oxygen produce overvoltages of some mV when only small current densities of some $\mu\text{A}/\text{cm}^2$ are applied. The dependence of the overvoltage upon time, oxygen concentration, and temperature was determined. It can be concluded that the oxygen transfer through the surface of the electrodes and the roughness of the surface cause the polarization. Diffusion effects can be neglected for short times. A remarkable influence of the electronic conductivity of the $\text{ThO}_{0.2}\text{-Y}_{0.2}\text{O}_{0.3}$ electrolyte on the e.m.f. appeared at oxygen activities of 10^{-28} to 10^{-29} atm. The effect of polarization on the determination of electronic transference numbers is discussed and a mean value

<193>

<193> CONT.

to $t_{\text{sub}}(e) = 0.01$ is estimated for a vanadium electrode
 $(p_{\text{sub } 02} = 10^{\text{sup } 29} \text{ atm.})$

EDF SUBJECT CATEGORY: EDB-400201

INIS CATEGORY: B12

<194>

Franczek, F.R.; Halstead, G.W.; Raymond, K.N. (California Univ.,
 Berkeley (USA). Dept. of Chemistry)

Actinide metallocarborane complex; synthesis and x-ray
 structure determination of the bis[n^{sup}
 5-(3)-1,2-dicarbollyl]dichlorouranium(IV) dianion
 AIX-07:257230; EDB-77:007650; J. Chem. Soc. (London), Chem.
 Commun. (JCCCA) (8):279-280 (21 Apr 1976)

SUBJECT DESCRIPTORS: BORANES;CHEMICAL PREPARATION:

Q1,Q2;CHEMICAL REACTIONS;CRYSTAL STRUCTURE:

Q1,Q2;FURANS;LATTICE PARAMETERS;LIGANDS;LITHIUM

COMPOUNDS;MOLECULAR STRUCTURE;ORGANIC BORON COMPOUNDS:

T2;URANIUM CHLORIDES;URANIUM COMPLEXES: T1;X-RAY DIFFRACTION

ABSTRACT: The first actinide metallocarborane complex,
 $[U(C_{\text{sub } 2}B_{\text{sub } 9}H_{\text{sub } 11})_{\text{sub } 2}Cl_{\text{sub } 2}]^{\text{sup } 2-}$, has
 been obtained by reaction of the 1,2-dicarbollide ion with
 $UCl_{\text{sub } 4}$ in tetrahydrofuran (THF), and a crystal structure
 determination of its lithium salt has shown the complex anion
 to have a distorted tetrahedral geometry with
 pentahapto-bonded dicarbollide ligands.

EDF SUBJECT CATEGORY: EDB-400702

INIS CATEGORY: B13

<195>

Fukushima, Y.; Suzuki, K. (Tohoku Univ., Sendai (Japan).
 Research Inst. for Iron, Steel and Other Metals)

Structure of $NO_{\text{sub } 3}^{\text{sup } -}$ radical in molten monovalent metal
 nitrates AIX-08:294245; EDB-77:068745; Kakuriken Kenkyu Hokoku
 (KKYHB) 8: (2):288-295 (Dec 1975)

SUBJECT DESCRIPTORS: DISTRIBUTION FUNCTIONS;LITHIUM NITRATES:

T1;MOLECULAR STRUCTURE: Q1,Q2,Q3,Q4,Q5,Q6;MOLTEN

SALTS;NEUTRON DIFFRACTION;NITRATES: T6;POTASSIUM NITRATES:

T2;RADICALS;RUBIDIUM NITRATES: T3;SILVER NITRATES: T4;SODIUM

NITRATES: T5;STRUCTURAL CHEMICAL ANALYSIS;TEMPERATURE

DEPENDENCE

ABSTRACT: The T-O-F pulsed neutron diffraction was
 experimentally studied on various kinds of monovalent metal

<195> CONT.

nitrites in the liquid state near the melting point. The T-O-F neutron diffractometer installed on the 300 MeV Tohoku University electron linac was used to measure the total structure factors for molten MNO_3 , where M represents Li, Na, K, Rb or Ag. Samples were sealed in vacua in silica tubes with 0.3 mm thick wall and 10 mm inner diameter. Before sealing, all samples were dried at 120 deg C to 150 deg C in vacua for 24 to 72 hours. During the measurements, the temperature of liquid samples were kept by an electric furnace at 280, 340, 390 and 260 deg C for Li-, Na-, K-, Rb- and $AgNO_3$, respectively. As a result of the experiment, there was big difference in the structure factors in the range of Q 5 Å⁻¹. In this region, a lot of informations must be included about $M^{+}NO_3^{3-}$ correlations. On the other hand, the structure factors in high Q region still showed that oscillatory behavior was very similar with one another. The radial distribution functions for the molten nitrites are illustrated. Least-squares analysis, the effect of intra-radical vibration, and the correction for the superposition of a close next peak for O-O separation are discussed. Discussion are focused on the behavior of the structure factors in high Q region.

EDB SUBJECT CATEGORY: EDB-360602
INIS CATEGORY: A13

<196>

Gamanovich, N.M.; Glybin, V.P.; Dobrotin, R.B. (Belorusskij Tekhnologicheskij Inst., Minsk)

Study of lanthanum, yttrium and erbium oxychlorides in melts of alkali metal chlorides AIX-08:339889; EDB-78:017320; Zh. Neorg. Khim. (ZNOKA) 22:(1):193-197 (Jan 1977)

SUBJECT DESCRIPTORS: CESIUM CHLORIDES: T1; COMPARATIVE EVALUATIONS; ERBIUM COMPOUNDS: T6; EUTECTICS; LANTHANUM COMPOUNDS: T7; LITHIUM CHLORIDES: T2; MOLTEN SALTS; OXYCHLORIDES; PHASE DIAGRAMS: Q1, Q2, Q3, Q4, Q5, Q6, Q7, Q8; POLYMERIZATION; POTASSIUM CHLORIDES: T4; RADIUM CHLORIDES: T3; RUBIDIUM CHLORIDES: T5; YTTRIUM COMPOUNDS: T8

ABSTRACT: The system $MeCl-(La, Y)OCl$ ($Me=Li, Na, K, Rb, Cs$) and $MeCl-ErOCl$ ($Me=Li, Rb, Cs$) have been studied by the methods of differential-thermal and X-ray phase analyses. The system $NaCl-LaOCl$ has been studied by cryoscopic method as well. It has been established that the systems refer to the type of fusibility diagrams with a simple eutectics. A comparison of the fusibility diagrams with calculations for ideal systems, under the assumption that the association degree of ions is different, makes it possible to assume a considerable

<196>

<196> CONT.

polymerizations degree of oxychloride in the melts
 MeCl-(Y,Er)OCl.

EDB SUBJECT CATEGORY: EDB-400201

INIS CATEGORY: B12

<197>

Garashina, L.S.; Vishnyakov, Y.S.

Structural changes in the series LnFeO_3 and LnF_3
 ERA-G3:008820;EDB-78:009638; Sov. Phys. - Crystallogr. (Engl.
 Transl.) (SPHCA) 22:(3):313-315 (May 1977)

SUBJECT DESCRIPTORS: ANISOTROPY;COORDINATION NUMBER;CRYSTAL
 STRUCTURE: Q1;INTERATOMIC DISTANCES;IONIC CRYSTALS;LATTICE
 PARAMETERS;RARE EARTH COMPOUNDS: M1

ABSTRACT: A structural analogy is established between two
 classes of rare earth compounds: fluorides LnF_3
 ($\text{Ln}=\text{Sm}--\text{Lu}$) and oxides LnFeO_3 ($\text{Ln}=\text{Pr}--\text{Lu}$). A reduction
 in the coordination number of the Ln^{3+} cation takes
 place in these compounds as a result of the displacement of
 the O or F atoms, leading to an increase in certain
 cation--anion distances. Anisotropy in the changes of
 individual interatomic distances is responsible for the
 nonmonotonic nature of the curve representing the crystal
 lattice parameters in the LnF_3 and LnFeO_3 series.

EDB SUBJECT CATEGORY: EDB-360602

INIS CATEGORY: B24

<198>

Gavrilin, I.V.; Ershov, G.S.

Self-diffusion in liquid metals AIX-08:286424;EDB-77:043780;
 Izv. Akad. Nauk SSSR, Met. (IZNMA) (5):87-94 (Sep 1976)

SUBJECT DESCRIPTORS: ACTIVATION ENERGY;BCC LATTICES;CADMIUM:
 T1;CLUSTER MODEL;PCC LATTICES;LIQUID METALS: T2;MELTING
 POINTS;SELF-DIFFUSION: Q1,Q2,Q3;TUNGSTEN: T3

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-360104

INIS CATEGORY: B22

<199>

Gawel, W.; Terpilowski, J. (Akademia Medyczna, Wroclaw (Poland).
Instytut Chemii i Technologii Srodkow Leczniczych)

Cryometric studies of the system $\text{CdCl}_2\text{-UCl}_4$
AIX-08:337064; EDB-77:141211; Rocz. Chem. (ROCHA)
48: (10):1683-1686 (1974)

SUBJECT DESCRIPTORS: CADMIUM CHLORIDES:

T1; ENTHALPY; EUTECTICS; FORMATION FREE ENERGY; PHASE STUDIES:
Q1, Q2; PHASE TRANSFORMATIONS; URANIUM CHLORIDES: T2

ABSTRACT: The phase equilibria in the system $\text{CdCl}_2\text{-UCl}_4$ have been examined. On the basis of liquidus curve, a thermodynamic description of the liquid phase in this system has been given.

EDB SUBJECT CATEGORY: EDB-400702
INIS CATEGORY: B13

<200>

Gee, R.; Shelton, R.A.J.

Thermodynamic properties of Cr, Mn, Co and Ni dichlorides from
emf measurements on cells with solid electrolytes
EDB-77:128149; Inst. Min. Metall., Trans., Sect. C (TMECA)
85::C208-C210 (Dec 1976)

SUBJECT DESCRIPTORS: CHROMIUM CHLORIDES: T1; COBALT CHLORIDES:
T2; ELECTROLYTIC CELLS; ELECTROMOTIVE FORCE; MANGANESE
CHLORIDES: T4; NICKEL CHLORIDES: T3; ORE
PROCESSING; THERMODYNAMIC PROPERTIES: Q1, Q2, Q3, Q4

ABSTRACT: Emf measurements on solid electrolyte cells were used
to determine thermodynamic properties of chromium, cobalt,
manganese, and nickel dichlorides.

EDB SUBJECT CATEGORY: EDB-360603; 320303

<201>

Gel'd, P.V.; Esin, Yu.O.; Petrushevskii, M.S.; Ryss, G.M.;
Strigancov, A.I. (Ural'skij Politekhnikheskij Inst.,
Sverdlovsk (USSR); Chelyabinskij Politekhnikheskij Inst.
(USSR))

Enthalpies of formation of melts of yttrium with aluminium
AIX-08:283293; EDB-77:068481; Dokl. Akad. Nauk SSSR (DANKA)
228: (2):386-388 (11 May 1976)

SUBJECT DESCRIPTORS: ALUMINIUM ALLOYS: T1; BINARY ALLOY
SYSTEMS; ENTHALPY: Q1, Q2; LIQUID METALS; QUANTITY RATIO; VERY

<201>

<201> CONT.

HIGH TEMPERATURE; YTTRIUM ALLOYS: T2

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-360104

INIS CATEGORY: B22

<202>

Giacchetti, G.; Sari, C.

Behavior of molybdenum in mixed-oxide fuel

INS-77:000020; ERA-02:009200; EDB-77:005582; Nucl. Technol.
(NUTYPE) 31: (1): 62-69 (Oct 1976)

SUBJECT DESCRIPTORS: BARIUM: T3; BARIUM OXIDES; CHEMICAL
 REACTIONS: Q2, Q3, Q4, Q5, Q6, Q7; FISSION PRODUCTS:
 Q1; MIXTURES; MOLYBDENUM: T2; MOLYBDENUM ALLOYS; MOLYBDENUM
 OXIDES; NUCLEAR FUELS: T1; OXIDATION; PALLADIUM ALLOYS; PLUTONIUM
 OXIDES: T7; RHODIUM ALLOYS; RUTHENIUM
 ALLOYS; SOLUBILITY; TEMPERATURE GRADIENTS; TUNGSTEN: T5; TUNGSTEN
 OXIDES; URANIUM OXIDES: T6; VAPORS; ZIRCONIUM: T4

ABSTRACT: Metallic molybdenum, Mo--Ru--Rh--Pd alloys, barium, zirconium, and tungsten were added to uranium and uranium--plutonium oxides by coprecipitation and mechanical mixture techniques. This material was treated in a thermal gradient similar to that existing in fuel during irradiation to study the behavior of molybdenum in an oxide matrix as a function of the $O/(U + Pu)$ ratio and some added elements. Result of ceramographic and microprobe analysis shows that when the overall $O/(U + Pu)$ ratio is less than 2, molybdenum and Mo--Ru--Rh--Pd alloy inclusions are present in the uranium--plutonium oxide matrix. If the $O/(U + Pu)$ ratio is greater than 2, molybdenum oxidizes to MoO_2 , which is gaseous at a temperature approximately $1000^\circ C$. Molybdenum oxide vapor reacts with barium oxide and forms a compound that exists as a liquid phase in the columnar grain region. Molybdenum oxide also reacts with tungsten oxide (tungsten is often present as an impurity in the fuel) and forms a compound that contains approximately 40 wt percent of actinide metals. The apparent solubility of molybdenum in uranium and uranium--plutonium oxides, determined by electron microprobe, was found to be less than 250 ppm both for hypo- and hyperstoichiometric fuels.

EDB SUBJECT CATEGORY: EDB-050700; 220300

INIS CATEGORY: E23

<203>

Gilbert, B. (Univ. of Tennessee, Knoxville); Mamantov, G.;
 Begun, G.M.

Raman spectra of Al_2O_3 solutions in molten cryolite
 and other aluminum fluoride containing melts
 ERA-03:012892; EDB-78:022305; Inorg. Nucl. Chem. Lett. (INUCA)
 12::415-424 (1976)

SUBJECT DESCRIPTORS: ALUMINIUM FLUORIDES: T1; ALUMINIUM OXIDES:
 T2; CHEMICAL ANALYSIS: Q1; RAMAN SPECTRA: Q2

ABSTRACT: Experimental results are reported that show that oxide
 concentrations as low as 2 mole % can be detected in aluminum
 fluoride melts using Raman spectroscopy. Results of attempts
 to obtain further information on the complex species observed
 on dissolution of Al_2O_3 were not definitive, but
 species involving nonbridging Al--O bonds have been
 eliminated from consideration in concentrated Al_2O_3
 solutions in aluminum fluoride containing melts. All new
 bands observed associated with the oxide were below 600
 cm^{-1} . (ELM)

EDF SUBJECT CATEGORY: EDB-400201; 400104

<204>

Gilbert, B.; Mamantov, G. (Univ. of Tennessee, Knoxville);
 Begun, G.M.

Simple Raman cell and furnace usable at temperatures higher than
 1000°C for corrosive melts ERA-01:013809; EDB-76:036965;
 Appl. Spectrosc. (APSPA) 29:(3):276-278 (1975)

SUBJECT DESCRIPTORS: ALUMINIUM FLUORIDES:
 T1; DESIGN; FURNACES; MOLTEN SALTS: T3; RAMAN SPECTRA:
 Q1, Q2, Q3; SODIUM FLUORIDES: T2; SPECTROSCOPY; VERY HIGH
 TEMPERATURE

ABSTRACT: An extremely simple experimental setup was designed
 which can be built in a few days. A windowless cell was
 chosen; it was comprised of graphite. The furnace and cell
 were built in such a way that no surface capable of emitting
 thermal radiation can be seen by the spectrometer. To
 illustrate the performance possible, Raman spectra are
 presented for AlF_3 --NaF at 740°C and molten
 cryolite at 1015°C. (DLC)

EDF SUBJECT CATEGORY: EDB-400201

<205>

<205>

Girifalcc, L.A.

Criteria for solid solution formation in NFE alloys

ERA-02:022819;INS-77:005719;EDB-77:043652; Acta Metall.

{AMETA} 24:(8):759-791 (Aug 1976)

SUBJECT DESCRIPTORS: BINARY ALLOY SYSTEMS;DENSITY;ELECTRONIC
STRUCTURE: Q1;ELECTRONS;METALS;MIXING HEAT;RARE EARTH ALLCYS:
T1;SOLID SOLUTIONS;VALENCE;VEGARD LAW

ABSTRACT: The first order, nearly free electron (NFE) theory of metals is applied to homovalent, random, binary alloys. It is found that the volume dependent energy of mixing for such alloys can never be negative, though it can be quite small. The fundamental parameter that determines the volume dependent properties of metals and alloys is an ionic core radius (related to the cut-off radius of the empty core potential) which can be computed from the density of the pure metals. It was found that systems for which the theoretical energy of mixing was less than $10\frac{1}{2} - 3$ Rydbergs per valence electron form solid solutions, while others do not. This result is easily expressed as a requirement that the ion core radii differ by less than approximately 13% for divalent alloys and approximately 20% for monovalent alloys. Application of these criteria to a large number of alloys shows that they predict whether or not extensive solid solution will take place with reasonable success. The theory also shows that Vegard's law is very nearly valid in NFE random alloys. This is in general agreement with experiment.

EDB SUBJECT CATEGORY: EDB-360102

INIS CATEGORY: B22

<206>

Glushkova, V.B.; Panova, T.I.; Keler, Eh.K.; Fedorov, N.F. (AN
SSSR, Leningrad. Inst. Khimii Silikatov)

Thermodynamics of reactions of rare earth oxides with niobium
oxide AIX-08:292317;EDB-77:050536; Izv. Akad. Nauk SSSR,
Neorg. Mater. (IVNMA) 12:(9):1596-1599 (Sep 1976)

SUBJECT DESCRIPTORS: ALLOTROPY;CHEMICAL PREPARATION;CHEMICAL
REACTIONS: Q1,Q2;FORMATION HEAT: Q1,Q3;FREE ENERGY;NIOBATES:
T3;NIOBIUM OXIDES: T2;OXIDES;RARE EARTH COMPOUNDS:
T1;SPECIFIC HEAT;TEMPERATURE DEPENDENCE

<207> CONT.

ABSTRACT: None

EDE SUBJECT CATEGORY: EDB-360204;400201

INIS CATEGORY: B12

<207>

Golub, A.M.; Perepelitsa, A.P.; Slobodyanik, N.S.; Popel', P.P.
(Kievskij Gosudarstvennyj Univ. (Ukrainian SSR))

Binary molybdates of rare earths and silver (I)

AIX-08:289228;EDB-77:050844; (ZNOKAZh. Neorg. Khim.)

21: (4):1142-1144 (Apr 1976)

SUBJECT DESCRIPTORS: CHEMICAL PREPARATION: Q1,Q2,Q3;CRYSTAL

STRUCTURE: Q1,Q2,Q3;LATTICE PARAMETERS;MELTING

POINTS;MOLYBDATES: T1;RARE EARTH COMPOUNDS: T2;SILVER

COMPOUNDS: T3;THERMAL GRAVIMETRIC ANALYSIS;X-RAY DIFFRACTION

ABSTRACT: None

EDE SUBJECT CATEGORY: EDB-400201

INIS CATEGORY: B12

<208>

Golub, M.A.; Khajmenov, A.P. (AN SSSR, Sverdlovsk. Inst.
Elektrokhimii)

High-temperature unit to the IKS-21 spectrometer for the
detection of molten salts absorption spectra

AIX-08:322633;EDB-77:121461; Priib. Tekh. Eksp. (PRTEA)

18: (4):231 (Jul 1975)

SUBJECT DESCRIPTORS: ABSORPTION SPECTRA:

Q1;FOCUSING;FURNACES;INFRARED SPECTROMETERS: T;LENSES;MOLTEN

SALTS: T1;SPECTROPHOTOMETERS: T;VERY HIGH TEMPERATURE

ABSTRACT: A high-temperature attachment to the
batch-manufactured IKS-21 infrared spectrometer is described.
The instrument is equipped with a heating oven that is placed
between the illuminator and monochromator in such a way that
the flux of light energy from the global lamp is focused at
the center of the cell. An additional collecting lens is used
to focus the beam onto the monochromator slit.

EDE SUBJECT CATEGORY: EDB-440300

INIS CATEGORY: E42

<209>

<209>

Gcmczov, L.I.; Dedjurin, A.J.; Titov, S.G.; Ivanov, O.S. (AN SSSR, Moscow. Inst. Metallurgii) ** Blank, H.; Lindner, R. (eds.)

Physical and mechanical properties of some uranium-containing carbides In Plutonium 1975 and other actinides **
AIX-08:310001; ERA-02:041837; EDB-77:094406; (1976)
North-Holland

SUBJECT DESCRIPTORS: BINDING ENERGY; CRYSTAL LATTICES; ELECTRONIC SPECIFIC HEAT; INTERSTITIALS; LATTICE PARAMETERS; MECHANICAL PROPERTIES: Q1; MICROHARDNESS; PHASE DIAGRAMS; PHASE STABILITY; PHYSICAL PROPERTIES: Q1; POISSON RATIO; SOLID SOLUTIONS; STRAIN HARDENING; URANIUM CARBIDES: T1; VICKERS HARDNESS; YIELD STRENGTH; YOUNG MODULUS

ABSTRACT: The alloying of uranium carbides considerably deteriorates their nuclear characteristics. The change in physical and mechanical properties on formation of solid solutions of carbides, however, is of essential interest from the view-point of formulating rules for their strengthening. Properties of various uranium containing carbides are discussed.

EDB SUBJECT CATEGORY: EDB-360203; 360204; 050700
INIS CATEGORY: B24

<210>

Gopala Rao, R.V.; Sen, D. (Jadavpur Univ., Calcutta (India))

Perturbation treatment of the Percus-Yevick equation and structures of liquid magnesium and cadmium
AIX-08:300995; EDB-77:088841; Chem. Phys. Lett. (CHFLB) 43: (3):455-456 (1 Nov 1976)

SUBJECT DESCRIPTORS: CADMIUM: T; CORRELATION FUNCTIONS; FOURIER TRANSFORMATION; HIGH TEMPERATURE; LIQUID METALS; MAGNESIUM: T; PERCUS-YEVICK EQUATION; SQUARE-WELL POTENTIAL

ABSTRACT: An analytical expression for the structure factor has been derived by using a square well potential as a perturbation over the hard sphere potential. The calculated values for the structure factors of liquid magnesium and cadmium are in good agreement with experiments.

EDB SUBJECT CATEGORY: EDB-360102
INIS CATEGORY: B22

<211>

Gordon, P.; Breyer, N.N.; Albright, D.L.; Warke, W.R., Illinois
Inst. of Tech., Chicago (USA)

In Environmental sensitivity of structural metals: liquid metal
embrittlement. Annual technical progress report No. 4, 1 Jun
1972--31 May 1973 ** EDB-77:094287; (31 May 1973) AD--912439

SUBJECT DESCRIPTORS: ALUMINIUM;ANTIMONY;ARSENIC;CADMIUM;COPPER
ALLOYS;CRACKS;EMBRITTLEMENT: Q1;FAILURES;FRACTURE
PROPERTIES;FRACTURES;GRAIN
BOUNDARIES;IMPURITIES;INDIUM;LEAD;LIQUID METALS:
T2;MERCURY;METALLURGICAL EFFECTS: Q2;METALS:
T1;PHOSPHORUS;SOLDERING;STEELS;SURFACE PROPERTIES:TIN;ZINC

ABSTRACT: The phenomenon of LME is being investigated on levels
from the atomic through bulk specimen and structural
properties, and is being considered from both experimental
and theoretical viewpoints. The research is aimed at
elucidating the three important aspects of LME, namely, the
mechanism by which embrittlement takes place at a crack, or
potential crack, site; the mechanism by which the embrittling
species is transported to this site; and various
metallurgical, physical, and mechanical factors which have a
significant influence on the severity of the embrittlement.
(GRA)

EDB SUBJECT CATEGORY: EDB-360105

<212>

Gordon, P.; Breyer, N.N.; Broutman, L.J.; Dally, J.W.; Albright,
D.L., Illinois Inst. of Tech., Chicago (USA)

In Environmental sensitivity of structural metals: liquid metal
embrittlement. Annual technical progress report No. 2, 25
Jun 1970--31 May 1971 ** EDB-77:094057; (Jun 1971) AD--886441

SUBJECT DESCRIPTORS: ALUMINIUM ALLOYS;BUILDING MATERIALS:
T1;CADMIUM;EMBRITTLEMENT: Q1;FRACTURES;INDIUM;LEAD;LIQUID
METALS: T2;METALLURGICAL EFFECTS: Q2;STEELS;STRESSES:TIN;ZINC

ABSTRACT: Liquid metal embrittlement (LME) is being investigated
on levels from the atomic through bulk specimen and
structural properties, and is being considered from both
experimental and theoretical viewpoints. The research is
aimed at elucidating the three important aspects of LME,
namely, the mechanism by which the embrittlement takes place
at a crack, or potential crack, site, the mechanism by which
the embrittling species is transported to this site, and

<212>

<212> CONT.

various metallurgical, physical and mechanical factors which have a significant influence on the severity of the embrittlement. (auth)

EDB SUBJECT CATEGORY: EDB-360103

<213>

Gorecki, T. (Wyzsza Szkola Inzynierska, Opole (Poland). Lab. of Physics)

Vacancies und melting curves of metals at high pressure

AIX-08:327774;EDB-77:140506; Z. Metallkd. (ZEMTA)

68: (3):231-236 (Mar 1977)

SUBJECT DESCRIPTORS: BINDING ENERGY;DATA COMPILATION;MELTING: Q1,Q2,Q3,Q4,Q5;MELTING POINTS;METALS: T5;NEPTUNIUM: T1;PLUTONIUM: T2;PRESSURE DEPENDENCE;RARE EARTHS: T3;URANIUM: T4;VACANCIES;VERY HIGH PRESSURE

ABSTRACT: The vacancy mechanism of the melting process is utilized as a starting point in derivation of the pressure dependence of melting temperature for metals. The results obtained for the initial slope of the melting curve are compared with experimental data for 45 metals (including U, Np, Pu, rare earths) and in most cases the agreement is very good. An on-linearity of the fusion curve and appearance of the maximum on the melting curve at a pressure approximately equal to the bulk modulus is also predicted with qualitative agreement with existing experimental data.

EDB SUBJECT CATEGORY: EDB-360104

INIS CATEGORY: B22

<214>

Gorcneck, K.; Falta, G.; Pichlmayer, F., Oesterreichische Studiengesellschaft fuer Atomenergie G.m.b.H., Vienna

In Determination of the burn up and differentiation of the fission yields of uranium-thorium-fuels **

AIX-08:315028;ERA-02:054499;EDB-77:127184; (Feb 1975)

SGAF--2425

SUBJECT DESCRIPTORS: BURNUP: T2,Q1;CESIUM 137;FISSION YIELD;FUEL ELEMENTS: T1;GAMMA SPECTROSCOPY;HIGHLY ENRICHED URANIUM;ISOTOPE RATIO;MASS SPECTROSCOPY;MEASURING METHODS: Q2;MOLYBDENUM;MOLYBDENUM OXIDES;NUCLEAR FUELS;THORIUM OXIDES;URANIUM;URANIUM DIOXIDE

ABSTRACT: The burnup of highly enriched and highly burned up (U, Th) O₂ fuels has been determined by two different and

<214> CONT.

unrelated methods. The results for the fuel burnup calculated from the ratio of the uranium isotopes is compared with the results from the method of the stable molybdenum isotopes. The number of Mo-95 atoms enables the calculation of the total burnup; the respective fission yields of U-233 and U-235 are obtained by the ratio of Mo-95 to Mo-100 isotopes. This method requires fewer assumptions than the former but it is more laborious because one has to determine the content of uranium and molybdenum of the samples. Moreover the burnup data obtained from the gamma-spectroscopic determination of Cs-137 are given.

EDB SUBJECT CATEGORY: EDB-220300
INIS CATEGORY: B16

<215>

Goryachev, Yu.M.; Kovenskaya, B.A.; Dudnik, E.M.; Severyanina, E.N.; Arabei, B.G.

Electron structure of rare-earth metal tetraborides
EDE-77:031006; Zh. Strukt. Khim. (ZSTKA) 16: (6):1036-1040
(1975)

SUBJECT DESCRIPTORS: BORIDES: T2; ELECTRICAL PROPERTIES; ELECTRONIC STRUCTURE: Q1, Q2, Q3, Q4, Q5, Q6; GADOLINIUM BORIDES: T4; HALL EFFECT; HOLMIUM BORIDES: T6; NEODYMIUM BORIDES: T3; PHYSICAL PROPERTIES; RARE EARTH COMPOUNDS: T1; TERBIUM BORIDES: T5

ABSTRACT: The electron structure of the tetraborides of rare-earth metals, in particular NdB_4 , Gd_2B_4 , Td_2B_4 , and HoB_4 , is discussed theoretically and the results are compared with the experimentally-measured electrical and thermophysical properties. On the basis of a linear-combination-of-atomic-orbits approach, the electron energy spectrum of these four tetraborides is calculated, and the principal parameters of the electron energy structures are deduced. A scheme is proposed for the nature of the interatomic bonds in the tetraborides. Experimental Hall effect data, together with the theory, indicate that the conduction band of these compounds is formed by a group of electron states of B and metal atoms, the B electron states playing a minor role.

EDB SUBJECT CATEGORY: EDB-360204

<216>

<216>

Gotkis, I.S.; Gusarov, A.V.; Gorokhov, L.N. (AN SSSR, Moscow.
Inst. Vysokikh Temperatur; Ivanovskij
Khimiko-Tekhnologicheskij Inst. (USSR))

Mass spectra and vapor composition of niobium pentafluoride
EDB-76:049131; Zh. Neorg. Khim. (ZNOKA) 20:(5):1250-1253 (May
1975)

SUBJECT DESCRIPTORS: CHEMICAL COMPOSITION: Q1; MASS SPECTRA:
Q1; NIOBIUM FLUORIDES: T1; SUBLIMATION HEAT; VAPORS

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-400201
INIS CATEGORY: B12

<217>

Goudy, A.J., Pittsburgh Univ., Pa. (USA)

In Hydrogen solubility in rare earth intermetallic compounds **
INS-78:001581; EDB-78:021867; (1976) Univ. of Pittsburgh

SUBJECT DESCRIPTORS: ABSORPTION: Q4; CRYSTAL STRUCTURE; HYDROGEN:
T4; RARE EARTH ALLOYS: T1; SORPTIVE PROPERTIES: Q1, Q2, Q3; VOLUME

ABSTRACT: A study of the hydrogen sorption characteristics of
some $RFe_{3\%}$ and $R_{2\%}Co_{7\%}$ intermetallic compounds
was conducted. These compounds are of interest since new
materials are being sought which absorb and release large
quantities of hydrogen rapidly and reversibly at ambient
temperatures and pressures. These materials were found to be
very receptive to hydrogen. The proton densities in most
cases were as high as that in liquid hydrogen. Accordingly,
measurements were made pertaining to the thermodynamics of
hydride formation, specifically pressure-composition
isotherms from which ΔH , ΔG and ΔS for the
hydrogenations could be determined. The data provide
information as to the hydride stabilities.

EDB SUBJECT CATEGORY: EDB-360104
INIS CATEGORY: B22

<218>

Green, D.W.; Gabelnick, S.D.; Reedy, G.T., Argonne National
Lab., Ill. (USA)

In Fine definition of ir spectra from high temperature
interactions of U + O₂. Part II. Final report, 1
July 1974--30 June 1975 **
INS-77:002399; ERA-02:017902; EDB-77:031754; (Aug 1975)
AD-A--020534

SUBJECT DESCRIPTORS: CHARGE EXCHANGE: Q1,Q2,Q3,Q4; INFRARED
SPECTRA; NITRIC OXIDE: M4; NITROGEN DIOXIDE: M3; URANIUM
DIOXIDE: M1; URANIUM OXIDES: M2; VERY LOW TEMPERATURE

ABSTRACT: The reactions of UO and UO₂ with NO and NO₂
were studied by ir spectroscopy using the matrix-isolation
technique. Codeposition of vaporized UO and UO₂ with
NO₂ and with NO gases in an argon matrix at 14⁰K
resulted in the production of the UO₂⁺ molecular
ion with either a NO₂⁺ or NO⁺ anion. Ir
absorption frequencies were measured and interpreted as the
stretching modes of a linear UO₂⁺ ion and a bent
(bond angle = 109⁰) NO₂⁺ ion. Three
different reactions were observed to yield a UO₂⁺
cation product: (1) UO₂ + NO₂; (2) UO₂ +
NO; and (3) UO + NO₂.

EDB SUBJECT CATEGORY: EDB-400702; 400201; 360204
INIS CATEGORY: E13; B12; B23

<219>

Green, D.W.; Reedy, G.T.

Identification of UN in Ar matrices

INS-76:020261; ERA-02:001820; EDB-76:083240; J. Chem. Phys.
(JCPA) 65:(7):2921-2922 (1 Oct 1976)

SUBJECT DESCRIPTORS: ABSORPTION SPECTRA; INFRARED SPECTRA:
Q1; ISOTOPE EFFECTS: Q3; NITROGEN 14; NITROGEN 15: T3; SPECTRAL
SHIFT; URANIUM NITRIDES: T1; VIBRATIONAL STATES

ABSTRACT: Measurements of vibrational frequencies of the
matrix-isolated UN molecule are reported, leading to the
identification of a linear N-U-N molecule. (AIP)

EDB SUBJECT CATEGORY: EDB-640302
INIS CATEGORY: A12

<220>

<220>

Green, D.W.; Reedy, G.T.; Leibowitz, L., Argonne National Lab.,
Ill. (USA)

In Thermodynamic functions and vapor pressures of uranium and
plutonium oxides at high temperatures **

INS-77:014996; ERA-02:055867; EDB-77:130893; (1977)
CONF-770537--2

SUBJECT DESCRIPTORS: ELECTRONIC STRUCTURE; ENERGY LEVELS; INFRARED
SPECTRA; ISOTOPE EFFECTS; OXYGEN 16; OXYGEN 18; PLUTONIUM
DIOXIDE; PLUTONIUM OXIDES: M1; ROTATIONAL STATES; THERMODYNAMIC
PROPERTIES: Q1, Q2; URANIUM DIOXIDE; URANIUM OXIDES: M2; URANIUM
TRIOXIDE; VIBRATIONAL STATES

ABSTRACT: The total energy release in a hypothetical reactor
accident is sensitive to the total vapor pressure of the
fuel. Thermodynamic functions which are accurate at high
temperature can be calculated with the methods of statistical
mechanics provided that needed spectroscopic data are
available. This method of obtaining high-temperature vapor
pressures should be greatly superior to the extrapolation of
experimental vapor pressure measurements beyond the
temperature range studied. Spectroscopic data needed for
these calculations are obtained from infrared spectroscopy of
matrix-isolated uranium and plutonium oxides. These data
allow the assignments of the observed spectra to specific
molecular species as well as the calculation of
anharmonicities for monoxides, bond angles for dioxides, and
molecular geometries for trioxides. These data are then
employed, in combination with data on rotational and
electronic molecular energy levels, to determine
thermodynamic functions that are suitable for the calculation
of high-temperature vapor pressures.

EDB SUBJECT CATEGORY: EDB-050700; 360204
INIS CATEGORY: R23

<221>

Grishchenko, V.F.; Lutsenko, V.G. (AN Ukrainskoj SSR, Kiev.
Inst. Obshchej i Neorganicheskij Khimii)

Electrochemical behavior of tellurium (4) in oxychloride melts
AIX-08:334295; EDB-77:141085; Elektrokhimiya (ELKKA)
12: (12): 1863-1865 (Dec 1976)

SUBJECT DESCRIPTORS: ELECTROCHEMISTRY:
Q1; ELECTROLYSIS; ELECTROMOTIVE FORCE; MOLTEN
SALTS; OXYCHLORIDES; REACTION KINETICS; TELLURIUM COMPOUNDS: T1

<221> CONT.

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-400400

INIS CATEGORY: B12

<222>

Gruenberg, H., Zentralinstitut fuer Kernforschung, Rossendorf
bei Dresden (German Democratic Republic)

In Calculation and discussion of state diagrams in the system
Pu-F-O-H **

AIX-08:329845; ERA-03:010750; NTS-78:000938; EDB-78:017483; (Oct
1976) ZfK--316

SUBJECT DESCRIPTORS: ENTHALPY; FLUORIDE VOLATILITY
PROCESS; FLUORINATION; PHASE DIAGRAMS: Q1, Q2, Q3; PLUTONIUM
COMPOUNDS; PLUTONIUM FLUORIDES: T1; PLUTONIUM HYDRIDES:
T2; PLUTONIUM OXIDES: T3; THERMODYNAMICS

ABSTRACT: In the system Pu-F-O-H state diagrams at a pressure of
reacting gases of 1 atm have been calculated and discussed
together with state diagrams of the system U-F-O-H.
Conclusions are drawn for the separation of uranium and
plutonium and about the conversion of fluorides of uranium
and plutonium to the oxides.

EDB SUBJECT CATEGORY: EDB-400702; 050800

INIS CATEGORY: D12

<223>

Gubanov, V.A. (AN SSSR, Sverdlovsk. Inst. Khimii); Rosen, A.;
Ellis, D.E.

Electron structure and bonding in ThO_2 and UO_2
AIX-08:321137; EDB-77:113737; Solid State Commun. (SSCOA)
22: (4): 219-223 (Apr 1977)

SUBJECT DESCRIPTORS: CHARGE DENSITY; CHEMICAL BONDS; DIRAC
APPROXIMATION; ELECTRONIC STRUCTURE: Q1, Q2; ENERGY
LEVELS; HARTREE-FOCK METHOD; PHOTOELECTRON SPECTROSCOPY; SLATER
METHOD; THORIUM OXIDES: T1; URANIUM DIOXIDE: T2; X RADIATION

ABSTRACT: Electron energy levels and charge densities for
 ThO_2 and UO_2 are calculated in a molecular
cluster approximation, using spin unrestricted
Hartree-Fock-Slater and relativistic Dirac-Slater models.
Results compare favourably with X-ray photoelectron spectra

<223>

<223> CONT.

and reveal similarities in chemical bonding with rare earth oxides.

EDB SUBJECT CATEGORY: EDB-360202

INIS CATEGORY: B23

<224>

Guentherodt, H.-J.; Hauser, E.; Kuenzi, H.W. (Basel Univ. (Switzerland)); Evans, R.; Evers, J.; Kaldis, E.

Negative temperature coefficients of electrical resistivity: the divalent liquid metals Eu, Yb and Ba

AIX-07:275060;EDB-77:018827; J. Phys., F (London) (JPFMA) 6:(8):1513-1522 (Aug 1976)

SUBJECT DESCRIPTORS: BARIUM: T1;ELECTRIC CONDUCTIVITY: Q1,Q2,Q3;EUROPIUM: T2;HIGH TEMPERATURE;INTERATOMIC FORCES;LIQUID METALS;MELTING;MELTING POINTS;ORDER-DISORDER TRANSFORMATIONS;TEMPERATURE DEPENDENCE;VALENCE;VERY HIGH TEMPERATURE;YTTERBIUM: T3

ABSTRACT: The electrical resistivities ρ , of Eu, Yb and Ba have been measured at high temperatures. In solid Yb and Ba ρ varies linearly with temperature. This is contrary to some previous results. Both metals show a large percentage increase of ρ on melting. The temperature dependence of ρ in Eu is nonlinear and there is a smaller percentage change on melting. The resistivities of liquid Eu and Ba are larger than those of any other liquid metals. Negative temperature coefficients of resistivity are observed for all three metals in the liquid state. These results and those obtained in earlier measurements on alloys of rare earth metals are discussed in the context of existing theories. It is suggested that even in the liquid rare earth metals and their alloys an interpretation involving the liquid structure factors and the effective valences of the components may still provide a reasonable qualitative explanation of the behavior of the resistivity. Comments are also made on the occurrence of negative temperature coefficients in other disordered metallic systems.

EDB SUBJECT CATEGORY: EDB-360104

INIS CATEGORY: B22

<225>

Gunnerson, F.S.; Cronenberg, A.W. (New Mexico Univ., Albuquerque (USA). Dept. of Nuclear Engineering)

Prediction of the inert gas solubilities in stoichiometric molten UO_2 AIX-07:239562; ERA-02:012470; EDB-77:016973; J. Nucl. Mater. (JNUMA) 58:(3):311-320 (Dec 1975)

SUBJECT DESCRIPTORS: FAST NEUTRONS;FISSION PRODUCTS;LIQUID FUELS;LIQUIDS;RARE GASES: T1;SOLUBILITY: Q1;SOLVENT PROPERTIES: Q2;URANIUM DIOXIDE: T2

ABSTRACT: To analyze the effect of fission gas behaviour on fast reactor fuels during a hypothetical overpower transient, the solubility characteristics of the noble gases in molten UO_2 have been assessed. To accomplish this, a theoretical estimation of such solubilities is made by determining the reversible work required to introduce a hard sphere, the size of the gas atom, into the liquid solvent. Results indicate that the solubility of the noble gases in molten UO_2 is quite low, the molar fraction of gas-to-liquid being approximately 10^{-6} . Such a low solubility of fission gases suggests that for preirradiated fuels, added swelling or formation may occur upon melting. In addition, such low solubility potential indicates that the fission gases do not play an appreciable role in the fragmentation of molten UO_2 upon quenching in sodium coolant.

EDB SUBJECT CATEGORY: EDB-050700;360104;360603
INIS CATEGORY: B12

<226>

Hadari, Z.; Mintz, M.H., Israel Nuclear Society, Yavne

In Uranium dioxide as a ceramic nuclear fuel ** Israel Nuclear Society symposium on 'Nuclear fuel in Israel', Weizmann Institute of Science, Rehovot, 1.7.75, EDB-76:033599; (1975)
INIS-mf--1966/5

SUBJECT DESCRIPTORS: CERAMICS;DENSITY;ENRICHED URANIUM;FABRICATION: Q1;FISSION PRODUCTS;FUEL PELLETS;NATURAL URANIUM;NUCLEAR FUELS;PHASE DIAGRAMS;PHYSICAL PROPERTIES: Q1;POWDER METALLURGY;SINTERING;THERMAL CONDUCTIVITY;URANIUM DIOXIDE: T1

ABSTRACT: The advantages and disadvantages of the use of UO_2 as a nuclear fuel are shown. The two stages of the process of manufacture of UO_2 nuclear fuel, UO_2 powder and UC_2 pelletizing, are described as well as the factors influencing the sintering process. A phase diagram for the uranium-oxygen system below 1500°C is presented

<226>

<226> CCNT.

and the influence of O/U on thermal conductivity is exemplified. It is concluded that in the manufacture of pellets a density should be attained that would allow both a good thermal conductivity and the emission of the fission products.

EDB SUBJECT CATEGORY: EDB-050700;360201
INIS CATEGORY: B23

<227>

Haines, H.R.; Mardon, P.G.; Potter, P.E. (UKAEA Research Group, Harwell. Atomic Energy Research Establishment) ** Blank, H.; Lindner, R. (eds.)

Some constitutional studies on uranium-carbon- and plutonium-carbon-rhenium and technetium systems In Plutonium 1975 and other actinides **
AIX-08:310003;ERA-02:041827;EDB-77:094385; (1976)
North-Holland

ABSTRACT: Some new experimental data on the phase diagrams U-Re-C, U-Te-C, Pu-Re-C, and Pu-Tc-C are presented. The data obtained for U-Re-C are in agreement with previous studies. The observations on the Tc systems confirm the formation of a Tc carbide with the upper limit of carbon content corresponding to the formula $Tc_{0.3}C$. In the U-Tc-C system, the $UTcC_{0.2}$ is present having an orthorhombic crystal structure, and is isomorphous with $URcC_{0.2}$. In the Pu-Re-C system there is no compound analogous to $URcC_{0.2}$, however, in this system and in the Pu-Tc-C system there is a ternary phase with a composition C ca. 30 at.%, and the Pu:Re or Tc ratio ca. 1.25. The Pu-Tc-C system is also characterised by a ternary compound $PuTc_{0.2}(2-y)$ which has a monoclinic crystal structure. Tentative phase diagrams are presented for U-Re-C, and the Tc systems.

EDB SUBJECT CATEGORY: EDB-360202
INIS CATEGORY: B24

<228>

Haines, H.R.; Mardon, P.G.; Potter, P.E. ** Commission of the European Communities, Karlsruhe (Germany, F.R.). European Inst. for Transuranium Elements

Some constitutional studies on uranium--carbon and plutonium--carbon--rhenium and technetium systems In 5th international conference on plutonium and other actinides **
AIX-07:233120;EDB-76:087287; (1975) AED-Conf--75-417-017

SUBJECT DESCRIPTORS: CRYSTAL LATTICES;PHASE DIAGRAMS: Q1,Q2,Q3,Q4

<228> CONT.

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-360202

INIS CATEGORY: B23

<229>

Hall, D.G., Oak Ridge National Lab., Tenn. (USA)

In Spectral density functions for disordered systems **

INS-77:002176; ERA-02:017710; EDB-77:030648; (Nov 1976)

CRNL--5184

SUBJECT DESCRIPTORS: AMORPHOUS STATE: T3,Q1; HAMILTONIANS; LIQUID METALS: T2; MONTE CARLO METHOD; ONE-DIMENSIONAL CALCULATIONS; ORDER PARAMETERS; SOLIDS: T1; SPECTRAL DENSITY: Q2,Q3

ABSTRACT: This work describes a Monte Carlo calculation of the spectral density function $A(k,E)$ for a one-dimensional model of an amorphous solid. The model contains an adjustable short-range order parameter and $A(k,E)$ is obtained for several values of that parameter. The Hamiltonians considered are: (1) the Hamiltonian describing an electron moving in a potential consisting of randomly placed delta functions, (2) the Hamiltonian describing a system of coupled harmonic oscillators, and (3) a tight-binding Hamiltonian describing in a simplified way either electrons or spin-waves. This numerical calculation was used to interpret recent neutron scattering data on amorphous systems. The spectral density $A(k,E)$ gives the single-particle contribution to the dynamic form factor of neutron scattering theory. In another application of the model calculation, the spectral density is obtained in the quasicrystalline approximation (QCA) for the problem of an electron in a one-dimensional liquid metal. The QCA and exact spectral densities are compared and it is shown that the QCA fails in the strong scattering regime. 14 figures. (DIC)

EDB SUBJECT CATEGORY: EDB-360102

INIS CATEGORY: B22

<230>

Harder, J.M.; Young, W.H. (East Anglia Univ. (UK). School of Mathematics and Physics)

Entropies of liquid rare earth metals

AIX-08:342975; EDB-78:021744; Phys. Lett., A (PYLAA)

61:(7):468-470 (27 Jun 1977)

SUBJECT DESCRIPTORS: ANGULAR MOMENTUM; CERIUM:

<230>

<230> CONT.

T1;COUPLING;ELECTRONIC STRUCTURE:

Q1,Q2,Q3,Q4,Q5,Q6,Q7,Q8,Q9;ENTROPY:

Q1,Q2,Q3,Q4,Q5,Q6,Q7,Q8,Q9;EUROPIUM: T2;GADOLINIUM: T3;GROUND STATES;LANTHANUM: T4;LIQUID METALS: T9;NEODYMIUM:

T5;PRASEODYMIUM: T6;TERBIUM: T7;YTTERBIUM: T8

ABSTRACT: The entropies of the liquid rare earths are explained using hypotheses about the electronic structure which closely parallel some believed to be correct for the corresponding solids.

EDB SUBJECT CATEGORY: EDB-360104

INIS CATEGORY: B22

<231>

Hardy, R.J. (Univ. of Nebraska, Lincoln); Karo, A.M.

Lattice-dynamical calculation of the thermodynamic properties of NaCl, KCl and RbCl ERA-02:043811;EDB-77:101169; J. Phys. Chem. Solids (JPCSA) 38:(4):335-344 (1977)

SUBJECT DESCRIPTORS: CALCULATION METHODS;CRYSTAL

LATTICES;GRUENEISEN CONSTANT;POTASSIUM CHLORIDES:

T2;PREDICTION EQUATIONS;RUBIDIUM CHLORIDES: T3;SODIUM

CHLORIDES: T1;SPECIFIC HEAT;THERMAL EXPANSION;THERMODYNAMIC PROPERTIES: Q1,Q2,Q3

ABSTRACT: Results of a lattice-dynamical calculation of the coefficient of thermal expansion, the heat capacity, and the macroscopic Grueneisen ratio for NaCl, KCl and RbCl are presented. Good agreement with the experimental results is obtained without adjustable parameters. The needed normal mode frequencies and mode-Grueneisen parameters were calculated with a deformation dipole model with nearest- and next-nearest-neighbor central-force interactions. Predicted values for the static elastic constants, the strain derivatives of the static potential energy, the mechanical equilibrium value of the nearest-neighbor distance, and the mode-Grueneisen parameters are given.

FILE SUBJECT CATEGORY: EDB-400201

<232>

Hashimoto, T.; Yamamoto, M.; Kobayashi, K.

Preparation of uranium-233 with low uranium-232 content
 EDB-77:098238; J. Nucl. Sci. Technol. (Tokyo) (JNSTA)
 13: (3):119-124 (Mar 1976)

SUBJECT DESCRIPTORS: FUEL CYCLE; IRRADIATION: Q2; ISOTOPE
 RATIO; NEUTRON FLUENCE; PRODUCTION: Q1; PURIFICATION:
 Q1; SPECTROSCOPY; THORIUM OXIDES: T2; URANIUM 233: T1

ABSTRACT: Thorium dioxide, previously freed of uranium and protactinium contamination by means of ion-exchange, was irradiated for 70.5 and 140 hr respectively in the core and graphite thermal column positions of the KUR (Kyoto University Reactor). After being cooled for more than 1 1/2 yr, the ^{233}U produced was chemically purified by ion-exchange, and the resulting ^{233}U specimens were electro-deposited on a stainless steel plate for submission to α -spectrometric measurement with a silicon surface barrier detector. From the α spectra thus obtained, the ^{232}U to ^{233}U atomic ratios of the specimens were evaluated and compared with the calculated values. This revealed that the rate of formation of ^{232}U depended largely on the $^{232}\text{Th}(n, 2n)^{231}\text{Th}$ reaction by fast neutrons, and consequently, on the neutron energy distribution prevailing at the position of irradiation. The purest ^{233}U sample obtained from the thorium target, irradiated with pure thermal neutrons in the graphite thermal column, yielded a ^{232}U to ^{233}U atomic ratio of 3.02×10^{-9} , which corresponded to 6.62×10^{-6} in α activity ratio. 11 refs.

EDS SUBJECT CATEGORY: EDB-050700;360206

<233>

Hausman, J., Israel Atomic Energy Commission, Yavne. Soreq
 Nuclear Research Center; Tel Aviv Univ. (Israel). Dept. of
 Engineering Sciences

In Measurement of fluorescence of rare earths in molten salts
 (Nd^{3+} in $\text{NaNO}_3\text{-KNO}_3$ eutectic) **
 EDB-76:054300; (Oct 1975) IA--1318

SUBJECT DESCRIPTORS: COMPARATIVE
 EVALUATIONS; CONVERSION; EUTECTICS; FLUORESCENCE:
 Q1, Q2; FLUORESCENCE SPECTROSCOPY; GLASS; LIFETIME; MOLTEN
 SALTS; NEODYMIUM: T2; NEODYMIUM NITRATES: M1; POTASSIUM
 NITRATES; SODIUM NITRATES; TEMPERATURE DEPENDENCE

ABSTRACT: The existence of fluorescence of $\text{Nd}(\text{NO}_3)_3$ in $\text{NaNO}_3\text{-KNO}_3$ eutectic was proved and measured

<233>

<233> CONT.

with a type S-1 photomultiplier tube connected to an oscilloscope. The fluorescence of a 5% Nd: glass rod was also measured and its lifetime of $303 \pm 30 \mu\text{s}$ agrees with the data given in the literature. Fluorescence lifetimes of $\text{Nd}(\text{NO}_3)_3$ in NaNO_3 - KNO_3 were measured for some concentrations of $\text{Nd}(\text{NO}_3)_3$. The measured lifetime was nearly $10 \mu\text{s}$ for all the concentrations, 30 times lower than the value for glass. This low value indicates the existence of some quenchers in the sample, probably water that had remained after drying. An experimental system for measuring fluorescence spectra of the system was set up. The transitions $^4\text{F}_{3/2} \rightarrow ^4\text{I}_{9/2}$ and $^4\text{F}_{3/2} \rightarrow ^4\text{I}_{11/2}$ were seen and recorded. Measurements of lifetime in a sample containing 2% $\text{Nd}(\text{NO}_3)_3$ in NaNO_3 - KNO_3 were performed once at close to 250°C and twice one hour later at nearly room temperature. After cooling the fluorescence lifetime was close to $6 \mu\text{s}$ instead of $10 \mu\text{s}$ for the hot sample. The fluorescence conversion efficiency (η) of Nd: glass and of neodymium in NaNO_3 - KNO_3 eutectic was evaluated and for 5% Nd : glass a value of 0.25 ± 0.04 was obtained.

FILE SUBJECT CATEGORY: EDB-400201

INIS CATEGORY: B11

<234>

Heatherly, D.E.; Vallet, C.E., Oak Ridge National Lab., Tenn.
(USA)

In Composition gradient during electrolysis in mixtures
analogous to molten salt battery electrolytes **
ERA-03:004593; NTS-77:001090; EDB-77:146902; (1977)
CONF-770814--5

SUBJECT DESCRIPTORS: CHEMICAL

COMPOSITION; ELECTRODES; ELECTROLYSIS: Q1, Q2; ELECTROLYTES:
Q3; INTERFACES; LITHIUM CHLORIDES; LITHIUM-SULFUR BATTERIES:
T3; MASS TRANSFER; POTASSIUM CHLORIDES; POTASSIUM NITRATES:
T1; SILVER NITRATES: T2

ABSTRACT: Molten salt mixtures are attractive electrolytes for more efficient batteries. Li/S batteries with LiCl - KCl mixture as electrolyte have recently been developed. Calculations based on mass transport at electrode--electrolyte interface and in the electrolyte lead to a prediction of a concentration gradient between the electrodes. Experiments in the analogous potassium nitrate--silver nitrate system that confirm the results of

<234> CONT.

the calculations are reported. The material consists of the descriptions of slides given at a slide presentation. The slides, however, are not included. (RWR)

EDE SUBJECT CATEGORY: EDE-400400;250902

<235>

Heiss, A.

In Thermodynamic study of the U-O-C system by measurement of the carbon monoxide pressures at equilibrium **

ERA-02:049153;EDB-77:107005; J. Nucl. Mater. (JNUMA) :207-233 (1975) ORNL-tr--4326

SUBJECT DESCRIPTORS: CARBON: T3;CHEMICAL REACTIONS;GRAPHITE;OXYGEN: T2;PHASE STUDIES: Q1,Q2,Q3;URANIUM: T1;URANIUM DIOXIDE

ABSTRACT: Reaction in $\text{UC}_{0.2}$ /graphite mixtures, leading to carbon monoxide evolution, was investigated. Isotherms were established in the range 1720--1870°C, representing CO pressure as a function of O/U and C/U ratios. Mono- and bivariant equilibrium pressures and phase diagrams are established. This has made it possible to deduce the composition of the solid phases at high temperatures. The $\text{UC}_{0.2}$ phase contains a few vacancies while $\text{UC}_{0.2}$ (in equilibrium with $\text{UC}_{0.2}$ and C) does not, but contains 0.06 at percent of oxygen substituted for carbon $\text{UC}_{0.2}$ in equilibrium with $\text{UC}_{0.97}$ and U(C,O) is strongly hypostoichiometric and contains little oxygen ($\text{UC}_{1.03}$); the monoxycarbide, hyperstoichiometric, has the formula $\text{UC}_{0.95}\text{O}_{0.10}$. A maximum oxygen solubility of 25; in UC is observed. From these results, the free energies of formation, ΔG° , were calculated for $\text{UC}_{0.97}$, $\text{UC}_{0.94}\text{O}_{0.06}$, $\text{UC}_{0.91}$, $\text{UC}_{0.92}\text{O}_{0.06}$, $\text{UC}_{0.91}$, $\text{UC}_{0.63}\text{O}_{0.03}$, $\text{UC}_{0.66}$, $\text{UC}_{0.95}\text{O}_{0.1}$, UC and $\text{UC}_{0.75}\text{O}_{0.25}$.

EDE SUBJECT CATEGORY: EDE-360102

<236>

<236>

Herranz, J.; Bloxom, S.R.; Keeler, J.B.; Roth, S.R.,
Massachusetts Inst. of Tech., Oak Ridge, Tenn. (USA). School
of Chemical Engineering Practice

In Determination and correlation of mass transfer coefficients
in a stirred cell **

ERA-01:018709;INS-76:015991;EDB-76:057823; (17 Dec 1975)

ORNL/MIT--221

SUBJECT DESCRIPTORS: BREEDER REACTORS;EXTRACTION APPARATUSES:
T1;MASS TRANSFER: Q1;MERCURY;MOLTEN SALT
REACTORS;POLAROGRAPHY;REPROCESSING: Q2;SPENT FUELS: M2;WATER

ABSTRACT: In the proposed Molten Salt Breeder Reactor flowsheet,
a fraction of the rare earth fission products is removed from
the fuel salt in mass transfer cells. To obtain design
parameters for this extraction, the effect of cell size,
blade diameter, phase volume, and agitation rate on the mass
transfer for a high density ratio system (mercury/water) in
nondispersing square cross section contactors was determined.
Aqueous side mass transfer coefficients were measured by
polarography over a wide range of operating conditions.
Correlations for the experimental mass transfer coefficients
as functions of the operating parameters are presented.
Several techniques for measuring mercury-side mass transfer
coefficients were evaluated and a new one is recommended.
(auth)

EDB SUBJECT CATEGORY: EDB-050800;210500

INIS CATEGORY: E16;E35

<237>

Hightower, J.R. Jr. ** McNeese, L.E., Oak Ridge National Lab.,
Tenn. (USA)

Fuel processing for molten-salt reactors In Molten-salt reactor
program. Semiannual progress report for period ending
February 29, 1976 **

INS-77:003090;ERA-02:015898;EDB-77:029160; (Aug 1976)

ORNL--5132

SUBJECT DESCRIPTORS: BISMUTH;EXTRACTION
APPARATUSES;FLUORINATION;FUEL CYCLE;MASS TRANSFER;MOLTEN SALT
FUELS: T1;MOLTEN SALT REACTORS;NEODYMIUM
COMPOUNDS;PROTACTINIUM;RARE EARTHS;REPROCESSING: Q1;RESEARCH
PROGRAMS;SEPARATION PROCESSES;URANIUM

ABSTRACT: During this report period, work on the chemistry of
fuel reconstitution was resumed, and engineering development
progressed on the metal transfer process for rare-earth
element removal, on continuous fluorinators for uranium

<237> CONT.

removal, on fuel reconstitution, and on salt-bismuth contactors to be used in reductive extraction. (DLC)

EDB SUBJECT CATEGORY: EDB-050800;210500

INIS CATEGORY: B16;E35

<238>

Hildenbrand, D.L.

Thermochemistry of gaseous UF_5 and UF_4

ERA-02:039725;INS-77:010735;EDB-77:095045; J. Chem. Phys.

(JCPSCA) 66:(11):4788-4794 (1 Jun 1977)

SUBJECT DESCRIPTORS: ENTROPY;EQUILIBRIUM;FORMATION HEAT:

Q1,Q2;THERMOCHEMICAL PROCESSES;THERMODYNAMIC

PROPERTIES;URANIUM FLUORIDES: T2;URANIUM TETRAFLUORIDE: T1

ABSTRACT: Thermochemical data for gaseous UF_5 and UF_4 were obtained from equilibrium and electron impact measurements made by high temperature mass spectrometry. Equilibrium measurements yielded for the gaseous reaction $\text{Ag} + \text{UF}_5 = \text{AgF} + \text{UF}_4$ the enthalpy change $\Delta H_{298}^\circ = 16.9 \pm 1.5$ kcal by second law analysis. Additionally, the sublimation of $\text{UF}_4(\text{s})$ was studied by the torsion-effusion method, leading to $\Delta H_{298}^\circ = 76.2 \pm 0.5$ kcal/mol. These results were combined with auxiliary thermochemical data to derive the standard heats of formation ΔH_f° for $\text{UF}_5(\text{g}) = -460.6 \pm 3.4$ kcal/mol and ΔH_f° for $\text{UF}_4(\text{g}) = -377.8 \pm 0.6$ kcal/mol, and the bond dissociation energies $D_0(\text{F}-\text{U}-\text{F}) = 68.3 \pm 3.5$ kcal and $D_0(\text{F}-\text{U}-\text{F}) = 101.0 \pm 3.4$ kcal. Threshold energies for formation of UF_5^+ and UF_4^+ by dissociative ionization of UF_6 were determined along with the ionization potentials of UF_6 (14.00 eV), UF_5 (11.29 eV), and UF_4 (9.96 eV), leading to the BDE values $D_0(\text{F}-\text{U}-\text{F}) = 68.0 \pm 3.2$ kcal and $D_0(\text{F}-\text{U}-\text{F}) = 102.4 \pm 3.2$ kcal, in agreement with the equilibrium data. Uncertainties associated with the measurements and the interpretation of the data are discussed. The entropy of $\text{UF}_5(\text{g})$ inferred from the equilibrium measurements was used in a third law treatment of published UF_5 sublimation data, giving thermochemical results in good agreement with the above. The UF_4 sublimation measurements lead to a vapor entropy that seems more

<238>

<239> CONT.

compatible with a molecular configuration of lower symmetry than the obvious tetrahedral structure.

EDB SUBJECT CATEGORY: EDB-400702
INIS CATEGORY: B13

<239>

Hildenbrand, D.L., Stanford Research Inst., Menlo Park, Calif.
(USA)

In Thermochemistry of gaseous metal oxides. Final report, 1
Apr 1973--31 Dec 1975 ** ERA-02:014307;EDB-77:019279; (Mar
1976) AD-A--025661

SUBJECT DESCRIPTORS: DISSOCIATION ENERGY:

Q1,Q2,Q3,Q4,Q5;EUROPIUM OXIDES: T4;IONIZATION POTENTIAL:

Q1,Q2,Q3,Q4,Q5;THORIUM OXIDES: T1;TITANIUM OXIDES: T3;URANIUM
OXIDES: T5;ZIRCONIUM OXIDES: T2

ABSTRACT: The dissociation energies and ionization potentials of gaseous oxides in the Th-O, Zr-O, Ti-O, Eu-O, and U-O systems have been determined by high temperature mass spectrometry. Dissociation energies were derived from gaseous equilibrium measurements, while the ionization potentials were obtained from electron impact threshold measurements. Specifically, data were obtained for gaseous ThO, ThO₂, ZrO, ZrO₂, TiO, TiO₂, EuO, UO₂, and NO₃. Attempts to characterize the higher oxide of aluminum, AlO₂, were unsuccessful. New experimental information on the ionization potentials of Th, Zr, and Eu was also obtained. The results make it possible to accurately define the energetics of a number of ion-neutral and neutral-neutral reactions of the metals which are of importance in the analysis of nuclear burst effects. (GRA)

EDB SUBJECT CATEGORY: EDB-400201;360204;400702

<240>

Hildenbrand, D.L., Stanford Research Inst., Menlo Park, Calif.
(USA)

In Thermochemistry of gaseous compounds of metals. Annual
summary report 5 Dec 74--4 Dec 75 **
ERA-03:004547;EDB-78:009952; (Feb 1976) AD-A--040222

SUBJECT DESCRIPTORS: BARIUM BROMIDES: T1;BARIUM CHLORIDES:
T2;CALCIUM BROMIDES: T3;DISSOCIATION ENERGY;FORMATION
HEAT;STRONTIUM BROMIDES: T4;THERMODYNAMIC PROPERTIES:
Q1,Q2,Q3,Q4

ABSTRACT: Thermochemical studies of gaseous IIA monohalides

<240> CONT.

CaBr, SrBr, BaBr, and BaCl were carried out by high temperature mass spectrometry. The monohalides were generated by the reaction of HBr or HCl with appropriate metal oxides in molybdenum effusion cells. Dissociation energies and standard heats of formation were evaluated from various equilibrium measurements at temperatures of 1800 to 2000 K. Preliminary thermochemical data were obtained for gaseous BaI. In the course of the work, the gaseous oxybromide OBr was also observed, and thermochemical data were derived. From studies of the vaporization of CaBr₂, SrBr₂, and BaBr₂ by the torsion-effusion methods, values were derived for the heats of sublimation/vaporization, heats of formation, and dissociation energies of the gaseous dibromides. The derived dissociation energies of the monohalides are compared with values obtained from flame equilibria, chemiluminescence, ionic model calculations, and Birge-Sponer extrapolations; the results provide an interesting comparison of the experimental methods, and give further information about chemical bonding in the monohalides. (Author)

EDB SUBJECT CATEGORY: EDB-400201

<241>

Hildenbrand, D.L., Stanford Research Inst., Menlo Park, Calif. (USA). Physical Sciences Div.

In Thermochemistry of gaseous compounds of metals. Annual summary report, 5 Dec 1975--4 Dec 1976 **
ERA-02:045983;EDB-77:107363; (Jan 1977) AD-A--C35704

SUBJECT DESCRIPTORS: BARIUM IODIDES: T3; CALCIUM IODIDES: T1; CHEMILUMINESCENCE; DISSOCIATION ENERGY; EUROPIUM FLUORIDES: T4; FORMATION HEAT; SAMARIUM OXIDES: T5; STRONTIUM IODIDES: T2; THERMODYNAMIC PROPERTIES: Q1, Q2, Q3, Q4, Q5

ABSTRACT: Thermochemical studies of a number of gaseous metal halides and oxides have been carried out by high temperature mass spectrometry, including CaI, SrI, BaI, EuF, EuF₂, and SmO. Dissociation energies and heats of formation are derived from the data, and the results are compared with the predictions of several theoretical models where possible. For the IIA metal iodides, the experimental dissociation energies follow trends established by the fluorides, chlorides and bromides; the results are compatible with corrected Birge-Sponer extrapolations and with electrostatic model calculations. The data for BaI suggest that chemiluminescent spectra resulting from the Ba + I₂ reaction may be due in part to the reaction of electronically-excited Ba atoms. Sublimation data for CaI₂(s), and vaporization data for SrI₂(l) and BaI₂(l) obtained by the torsion-effusion method are consistent with estimated molecular parameters of the

<241>

<241> CONT.

gaseous diiodide molecules, leading to accurate thermochemical data for these molecules. The results for SmO resolve a long-standing discrepancy between the dissociation energies of SmO and EuO . And finally, torsion-effusion studies of the dissociative vaporization of $\text{PbF}_2(\text{s})$ yielded new thermochemical data for $\text{PbF}_4(\text{g})$.

EDE SUBJECT CATEGORY: EDB-400201

<242>

Hirata, K.; Moriya, K.; Waseda, Y. (Tohoku Univ., Sendai (Japan). Research Inst. of Mineral Dressing and Metallurgy)

High temperature thermal expansion of ThO_2 , MgO and Y_2O_3 by X-ray diffraction AIX-08:312372; EDB-77:107199; J. Mater. Sci. (JMTSA) 12:(4):838-839 (Apr 1977)

SUBJECT DESCRIPTORS: EQUATIONS; HIGH TEMPERATURE; LATTICE PARAMETERS; MAGNESIUM OXIDES: T1; TEMPERATURE DEPENDENCE; THERMAL EXPANSION: Q1, Q2, Q3; THORIUM OXIDES: T2; VERY HIGH TEMPERATURE; X-RAY DIFFRACTION; YTTRIUM OXIDES: T3

ABSTRACT: Using apparatus described previously (Waseda et al., High Temp. High Press.; 7:221 (1975)) X-ray diffraction measurements were made on ThO_2 , MgO and Y_2O_3 samples over the temperature range 500 to 1800°C. Lattice parameters over the range of temperature are shown. The data were converted into % linear expansion and plotted. Equations are presented.

EDE SUBJECT CATEGORY: EDB-360204
INIS CATEGORY: B23

<243>

Hoch, M.; Vernardakis, T. (Cincinnati Univ., Ohio (USA). Dept. of Materials Science and Metallurgical Engineering)

Thermodynamic properties of solid and liquid metals and ceramics AIX-08:298632; EDB-77:068455; High Temp. - High Pressures (HTHPA) 8:(3):241-246 (1976)

SUBJECT DESCRIPTORS: ALUMINIUM OXIDES: T3; DEBYE TEMPERATURE; ELECTRONIC SPECIFIC HEAT; ENTHALPY; FUSION HEAT; HIGH TEMPERATURE; LEAD: T2; LIQUID METALS; SODIUM: T1; SOLIDS; SPECIFIC HEAT; TEMPERATURE DEPENDENCE; THERMODYNAMIC PROPERTIES: Q1, Q2, Q3, Q4, Q5; URANIUM DIOXIDE: T4; VANADIUM: T5; VERY HIGH TEMPERATURE

ABSTRACT: It has been shown earlier that the high-temperature specific heat of solids can be expressed as $C_{\text{sub}}(p) =$

<243> CONT.

$3nRF(\theta/D)/T + bT + dT^3$ where $F(\theta/D)/T$ is the Debye function, b is equivalent to the electronic specific heat, and anharmonic vibrations within the lattice contribute only to d ; n is the number of atoms per molecule. The specific heat of liquid metals at high temperature can be expressed as $C_{sub}(p) = 3RF(\theta/D)/T + gT + hT^2$ where g is the electronic term, and h the anharmonic term. These two equations are used after integration to express the enthalpy of Na, Pb, V, $Al_{sub} 2O_{sub} 3$, and $UO_{sub} 2$, over a wide temperature range, in both solid and liquid phases, where heat content data are available; they are found to represent the data very well. In addition, heats of fusion calculated with the use of the equations are in very good agreement with experimental values. The second of the two equations can also be used to estimate the high-temperature specific heat for other liquid metals and ceramics where available data are scarce.

EDB SUBJECT CATEGORY: EDB-360104;360204
INIS CATEGORY: B23

<244>

Hoch, M.; Vernardakis, T. (Cincinnati Univ., Ohio (USA). Dept. of Materials Science and Metallurgical Engineering)

Specific heat of liquid metals and ceramics

AIX-08:323990;EDB-78:009985; Rev. Int. Hautes Temp. Refract. (RIHTA) 3:(2):75-82 (Nov 1976)

SUBJECT DESCRIPTORS: ALUMINIUM OXIDES:

T1; CERAMICS; ENTHALPY; IRON: T2; LEAD: T3; LIQUID METALS; LITHIUM: T4; MERCURY: T5; MOLTEN SALTS; POTASSIUM: T6; SODIUM: T7; SPECIFIC HEAT: Q1, Q2, Q3, Q4, Q5, Q6, Q7, Q8, Q9, Q10; TEMPERATURE DEPENDENCE; TITANIUM: T8; URANIUM DIOXIDE: T9; VANADIUM: T10; VERY HIGH TEMPERATURE

ABSTRACT: An analysis of the specific heat of liquid metals, where accurate heat content data are available over a large temperature range (lithium, potassium, sodium, lead, mercury), show that the specific heat of a liquid at high temperature can be expressed as $C_{sub}(l) = 3RF(\theta/D)/T + gT + hT^2$ where g is the electronic specific heat and h the anharmonic term. This equation was applied to Fe, V, Ti, $Al_{sub} 2O_{sub} 3$, and $UO_{sub} 2$ where heat content data are available over a large temperature range; the fit is very good. This equation can thus be used to estimate the high temperature specific heat

<244>

<244> CONT.

for other liquid metals and ceramics where the available data are scarce.

EDB SUBJECT CATEGORY: EDB-400201;360104;360204;400702
INIS CATEGORY: B12

<245>

Holleck, H. (Kernforschungszentrum Karlsruhe (Germany, F.R.).
Inst. fuer Material- und Festkoerperforschung)

Structure of the thorium-(zirconium, niobium, ruthenium,
rhodium)-carbon systems AIX-08:324226;EDB-77:132591; J. Nucl.
Mater. (JNDMA) 66:(3):273-282 (May 1977)

SUBJECT DESCRIPTORS: EQUILIBRIUM;FORMATION HEAT:
Q1,Q2,Q3,Q4,Q5;PHASE DIAGRAMS:
Q1,Q2,Q3,Q4,Q5;SOLUBILITY;TERNARY ALLOY SYSTEMS;THORIUM
CARBIDES: T1

ABSTRACT: With the aid of arc-melted and subsequently homogenised samples which were examined by X-rays, micrography and (sometimes) microanalysis, it was possible to establish isothermal sections in the ternary systems Th-Zr-C, Th-Nb-C, Th-Ru-C and Th-Rh-C. In the Th-Zr-C and Th-Nb-C systems, the stable carbides of zirconium and niobium are in equilibrium with thorium and thorium carbides. There are no ternary carbides. Three complex ternary carbides determine the phase-fields in the Th-Ru-C system. The reaction of thorium carbides with rhodium generates stable intermetallic phases, plus free carbon. $\text{ThRh}_{3\%}$ dissolves about 5-7 at.% carbon. This stabilizes $\text{ThRh}_{3\%}$ relative to neighbouring Th/Rh phases, and this suppresses $\text{ThRh}_{2\%}$ and $\text{ThRh}_{5\%}$ in the ternary region. The free enthalpies of formation of the ternary compounds were estimated from binary data, taking into account observed phase equilibria, and the estimates are discussed.

EDB SUBJECT CATEGORY: EDB-360104;360102
INIS CATEGORY: B22

<246>

Hseuh, L.; Nelson, A.G.; Mirza, Z.I.; Bennion, D.N., California Univ., Los Angeles (USA). School of Engineering and Applied Science

In Nonaqueous electrolyte systems: ionic transport in nonaqueous media. Interim technical report No. 7 **
EDE-77:049808; (Jul 1970) AD--714680

SUBJECT DESCRIPTORS: ALUMINIUM OXIDES;BORATES;ELECTRIC

<246> CCNT.

BATTERIES: T2;ELECTROLYTES: Q1,Q2;FEASIBILITY STUDIES;FUEL
CELLS: T1;ION MOBILITY;IONIC CONDUCTIVITY;LITHIUM
COMPOUNDS;LITHIUM FLUORIDES;ORGANIC SULFUR
COMPOUNDS;PERCHLORATES;SODIUM OXIDES

ABSTRACT: The purpose of this work was to determine the general suitability of beta-alumina for use in electrochemical systems and the suitability of lithium perchlorate--dimethyl sulfite and lithium tetrafluoroborate--dimethyl sulfite electrolyte solutions in electrochemical fuel cells or batteries. The emphasis is on the nature of ionic conduction in beta-alumina and the nature of ionic transport in the dimethyl sulfite electrolytes.

EDB SUBJECT CATEGORY: EDB-250903;300503

<247>

Hubberstey, P. (Nottingham Univ. (UK)) ** Evans, R.; Greenwood, D.A. (eds.)

Aspects of the solution chemistry of liquid alkali metals as elucidated from electrical resistivity studies In Liquid metals, 1976 ** AIX-08:320631;EDB-77:114220; (1977) Institute of Physics

SUBJECT DESCRIPTORS: CHEMICAL COMPOSITION;CHEMICAL REACTION KINETICS;CHEMICAL REACTIONS;ELECTRIC CONDUCTIVITY;LIQUID METALS;LITHIUM: T1;PHASE STUDIES;SODIUM: T2;SOLUBILITY;SOLUTIONS;SOLVATION;SOLVENT PROPERTIES: Q1,Q2;SOLVENTS;TEMPERATURE DEPENDENCE

ABSTRACT: Resistivity techniques have been applied to a study of the behavior of liquid alkali metals as solvents and reaction media; phase equilibria, solute reaction chemistry and the nature of the solute species have been investigated. Lithium and sodium are generally chosen as solvents and both non-metals and metals are studied as solutes. Phase relationships in alkali metal systems have been investigated using both resistivity-concentration and resistivity-temperature methods; these techniques are particularly effective for the determination of solubility data (i.e. the hypereutectic liquidus) and of liquid-liquid miscibility gaps, phase boundaries which are difficult to examine using conventional thermal methods. The stoichiometry and kinetics of chemical reactions in alkali metal solutions have been studied using resistivity techniques to monitor solute concentration. Thus, resistivity changes have shown the role of the solvent to be important when considering solute interaction in liquid metals; whereas association between $H^{\text{sup}} - \text{and } O^{\text{sup}} 2 -$ occurs in sodium at 723 K according to the equilibrium H^{sup}

<247>

<247> CONT.

$-e + 0e^{sup 2} = 0H^{sup -} + 2e$, $H^{sup -}$ and $N^{sup 3-}$ act independently in lithium at 693 K. Furthermore, resistivity changes have been used to identify solvated species; dissolution of nitrogen in solutions of barium in liquid sodium at 573 K gives rise to strongly bonded solvated unit of stoichiometry close to $Ba^{sub 4}N$.

EDB SUBJECT CATEGORY: EDB-400201
INIS CATEGORY: B12

<248>

Huijben, M.J.; van der Lugt, W. (Rijksinstituut voor Zuivering van Afvalwater, Voorburg (Netherlands))

Static structure factor of liquid caesium

AIX-07:274108; EDB-77:018822; J. Phys., F (London) (JPFMA)
6: (8):L225-L229 (Aug 1976)

SUBJECT DESCRIPTORS: CESIUM: T1; DISTRIBUTION FUNCTIONS; ELECTRIC CONDUCTIVITY; INTERATOMIC FORCES: Q1; LIQUID METALS; MEDIUM TEMPERATURE; X-RAY DIFFRACTION

ABSTRACT: X-ray transmission measurements of the structure factor of liquid caesium have been carried out for temperatures between $30^{sup 0}C$ and $150^{sup 0}C$. From these measurements, the radial distribution function, the pair potential and the electrical resistivity are derived.

EDB SUBJECT CATEGORY: EDB-360104
INIS CATEGORY: A13

<249>

Huijben, M.J.; van der Lugt, W. (Rijksuniversiteit Groningen (Netherlands)) ** Evans, R.; Greenwood, D.A. (eds.)

Structure factor of liquid caesium and sodium-caesium alloys In Liquid metals, 1976 ** AIX-08:319480; EDB-77:120333; (1977) Institute of Physics.

SUBJECT DESCRIPTORS: CESIUM: T1; CESIUM ALLOYS: T2; ELECTRONIC STRUCTURE: Q1, Q2, Q3, Q4, Q5; INTERATOMIC DISTANCES; INTERATOMIC FORCES; LIQUID METALS; POTASSIUM: T5; SCALING LAWS; SODIUM: T4; SODIUM ALLOYS: T3; TRANSMISSION; X-RAY DIFFRACTION

ABSTRACT: X-ray transmission diffraction experiments have been carried out for liquid sodium, potassium, caesium and some sodium-caesium alloys. The structure factor of liquid pure

<249> CONT.

caesium at the melting point can be scaled down to those of sodium and potassium with reasonable accuracy.

EDE SUBJECT CATEGORY: EDE-360104
INIS CATEGORY: A13

<250>

Huijben, M.J.; van der Lugt, W. (Rijksuniversiteit Groningen (Netherlands)) ** Nederlandse Natuurkundige Vereniging, Amsterdam

Structure factors of liquid alkali metals In Spring meeting, 26 and 27 April 1976, Amsterdam ** AIX-08:283295;EDB-77:068427; (1976) INIS-mf--3227

SUBJECT DESCRIPTORS: CESIUM: T1;GONIOMETERS;LIQUID METALS;POTASSIUM: T2;SMALL ANGLE SCATTERING;SODIUM;X-RAY DIFFRACTION: Q1,Q2

ABSTRACT: None

EDE SUBJECT CATEGORY: EDE-360104
INIS CATEGORY: B22

<251>

Ikeda, Y.; Tamaki, M.; Matsumoto, G. (Nagoya Univ. (Japan). Faculty of Engineering)

Vaporization of uranium-carbon-nitrogen system
AIX-07:244137;EDB-77:003017; J. Nucl. Mater. (JNUMA) 59:(2):103-111 (Feb 1976)

SUBJECT DESCRIPTORS: ENTHALPY;EVAPORATION: Q1,Q2;SOLID SOLUTIONS;TEMPERATURE DEPENDENCE;URANIUM CARBIDES: T1;URANIUM NITRIDES: T2;VAPOR PRESSURE;VERY HIGH TEMPERATURE

ABSTRACT: The vaporization of species from the uranium-carbon-nitrogen system has been investigated by mass spectrometry with the Knudsen cell technique. After calibration with gold, the vapor pressures of U and $N_{2(g)}$ have been obtained over $UC_{(1-x)}N_{(x)}$ in the temperature range of 1900-2300K. For uranium mono-nitride, values of enthalpies ΔH^0_{298} for overall reaction (1) $UN(s) = U(g) + 0.5N_{2(g)}$ and the partial reactions (2) $UN(s) = U(L) + 0.5N_{2(g)}$ and $U(L) = U(g)$ have been obtained with both second- and third-law treatments. For $UC_{(1-x)}N_{(x)}$ the experimental vapor pressures were used to calculate activity of UN in the solutions. For three

<251>

<251> CONT.

compositions ($x=0.69, 0.48, 0.30$) measured, activity coefficients were found to be slightly smaller than unity.

EDB SUBJECT CATEGORY: EDB-360202

INIS CATEGORY: B23

<252>

Iksanov, E.A.; Martynov, S.Z.; Minaev, Y.A.; Pavlov, Y.A.
(Moskovskij Inst. Stali i Splavov (USSR))

Spreading of iron-nickel alloys on ZrC and nickel-cobalt alloys on ZrN AIX-09:349475;EDB-78:027704; Izv. Vyssh. Uchebn. Zaved., Chern. Metall. (IVUMA) (1):12-13 (1976)

ABSTRACT: Spreading of iron-nickel melts on ZrC has been studied, and the diffusion film thickness is estimated alongside with the mobility of components in nickel alloys. A fairly linear character of the squared drop radius dependence on the time of wetting indicates that the process is of diffusion nature, namely, the wetting film is spreading according to the diffusion growth mechanism.

EDB SUBJECT CATEGORY: EDB-360204;360104

INIS CATEGORY: B22

<253>

Il'in, V.K.; Chervonnyi, A.D.; Krenev, V.A. (AN SSSR, Moscow.
Inst. Obshchej i Neorganicheskoy Khimii)

Kinetics and mechanism into preparation of samarium-, europium- and ytterbium dichlorides and their some properties
AIX-08:329852;EDB-78:010005; Zh. Neorg. Khim. (ZNOKA)
21:(3):614-618 (Mar 1976)

ABSTRACT: A kinetic study has been made of the reaction of hydrogen with anhydrous trichlorides of samarium, europium and ytterbium in the temperature ranges 20-710, 20-480 and 20-660 deg and hydrogen pressures 0.02-1.0 atm, as well as of the process of oxidation of dichlorides of these elements by hydrogen chloride in the temperature ranges 560-660 deg, 280-360 and 450-585 deg. Activation energies of the process of reduction of trichlorides of samarium, europium and ytterbium were found to be 18.9; 28.9; 25.7 kcal/mole, respectively. The results of kinetic studies allowed conditions of reduction of trichlorides to be proposed to obtain anhydrous dichlorides of high purity. The pressure of saturated vapours of dichlorides obtained was determined in the temperature range 620-1005 deg. Data obtained on the rate of evaporation of dichlorides allowed calculation of the

<253> CONT.

dependence of the vapour pressure of these compounds on temperature in the form of a two-member equation.

EDB SUBJECT CATEGORY: EDB-400201

INIS CATEGORY: B12

<254>

Il'yasov, I.I.; Dunaeva, T.I. (Rostovskij-na-Donu Filial Zaochnogo Inst. Sovetskoy Torgovli (USSR))

Ternary system of potassium, cesium and calcium nitrates
AIX-08:339901; EDB-78:017324; Zh. Neorg. Khim. (ZNOKA)
22: (1):265-267 (Jan 1977)

SUBJECT DESCRIPTORS: CALCIUM NITRATES: T1; CESIUM NITRATES:
T2; CRYSTALLIZATION; EUTECTICS; PHASE DIAGRAMS:
Q1, Q2, Q3; POTASSIUM NITRATES: T3

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-400201

INIS CATEGORY: B12

<255>

Il'yasov, I.I.; Dunaeva, T.I.; Litvinov, Yu.G.; Volchanskaya, V.V. (Zaochnyj Inst. Sovetskoy Torgovli, Rostov-na-Donu (USSR))

$\text{LiNO}_3\text{-CsNO}_3\text{-Ca(NO}_3)_2$ and $\text{NaNO}_3\text{-CsNO}_3\text{-Ca(NO}_3)_2$ system
AIX-08:282887; EDB-77:037410; Zh. Neorg. Khim. (ZNOKA)
21: (8):2298-2301 (Aug 1976)

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-400201

INIS CATEGORY: B12

<256>

Il'yasov, I.I.; Iskandarov, K.I.; Davranov, M.; Berdieva, R.N.

Phase diagrams of LiCl , CsCl , CaCl_2 and LiBr , CsBr , CaBr_2 ternary systems EDB-76:036670; Zh. Neorg. Khim. (ZNOKA)
20: (1):250-253 (Jan 1975)

SUBJECT DESCRIPTORS: EUTECTICS; HIGH TEMPERATURE; MELTING
POINTS; MOLTEN SALTS; PHASE DIAGRAMS: Q1, Q2, Q3, Q4, Q5, Q6

<256>

<256> CONT.

ABSTRACT: None

EIE SUBJECT CATEGORY: EDB-360602

INIS CATEGORY: B12

<257>

Il'yasov, I.I.; Iskandarov, K.I. (Samarkandskiy Meditsinskiy
Inst. (USSR))Ternary mutual system of chlorides and iodides of rubidium and
cadmium AIX-08:329853; EDB-78:010006; Zh. Neorg. Khim. (ZNCKA)
21: (3):792-794 (Mar 1976)SUBJECT DESCRIPTORS: CADMIUM CHLORIDES: T1; CADMIUM IODIDES:
T2; CHEMICAL REACTIONS; CRYSTALLIZATION; PHASE DIAGRAMS:
Q1, Q2, Q3, Q4; RUBIDIUM CHLORIDES: T3; RUBIDIUM IODIDES: T4

ABSTRACT: A study has been made of the ternary systems of rubidium and cadmium chlorides and iodides by visual polythermal method. To avoid decomposition of iodides a stream of dry carbon dioxide has been permanently passed through the melt. It has been found that the crystallization surface of the system studied is determined by existence of three compounds $\text{CdJ}_{\text{sub } 2} \times 2\text{RbJ}$, $\text{CdCl}_{\text{sub } 2} \times \text{RbCl}$ and $\text{CdCl}_{\text{sub } 2} \times 4\text{RbCl}$ on sides $\text{CdJ}_{\text{sub } 2} - \text{Rb}_{\text{sub } 2}\text{J}_{\text{sub } 2}$ and $\text{CdCl}_{\text{sub } 2} - \text{Rb}_{\text{sub } 2}\text{Cl}_{\text{sub } 2}$, as well as eutectics on two other sides. The system is a belt-like adiaxonal mutual system with clear-cut formation of complexes between components of binary systems. The crystallization surface is complex and consists of seven fields: four fields of components and three fields of compounds. In the field of $\text{CdCl}_{\text{sub } 2}$ the homomorphous transformation at 460 deg obtained earlier has been found. The above-mentioned fields coincide in five non-variant points.

EDE SUBJECT CATEGORY: EDB-400201

INIS CATEGORY: B12

<258>

Il'yasov, I.I.; Litvinov, Yu.G. (Rostovskiy-na-Donu Filial
Zachnogo Inst. Sovetskoy Torgovli (USSR))TlCl-RbCl-CsCl and NaI-TlI-CsI systems EDB-76:048905; Ukr. Khim.
Zh. (UKZHA) 41: (2):210-211 (Feb 1975)

SUBJECT DESCRIPTORS: PHASE DIAGRAMS: Q1, Q2, Q3, Q4, Q5, Q6

<258> CONT.
ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-360602;400201
INIS CATEGORY: B12

<259>
Imoto, S.; Adachi, H.; Hori, T. (Osaka Univ., Suita (Japan).
Faculty of Engineering)

Band structure and chemical bond in ThX compounds
AIX-08:280630;EDB-77:031011; J. Nucl. Sci. Technol. (Tokyo)
(JNSTA) 12:(11):711-716 (Nov 1975)

SUBJECT DESCRIPTORS: BAND THEORY;CHEMICAL BONDS;COMPUTER
CALCULATIONS;ELECTRONIC STRUCTURE;MOLECULAR STRUCTURE:
Q1,Q2,Q3,Q4;THORIUM;THORIUM CARBIDES: T1;THORIUM NITRIDES:
T2;THORIUM PHOSPHIDES: T3;THORIUM SULFIDES: T4

ABSTRACT: The band structure of metallic Th and NaCl-type ThX
(X=C, N, P, S) compounds were calculated by tight-binding
method, with the parametrization improvement. Values of
two-center parameters chosen so as to fit the result for Th
given by Gupta and Loucks were found to be very close to
two-center integrals with a suitable integration range.
Applying this method for evaluating parameters, the band
structures for ThX compounds were also computed. Population
analysis to determine the electron populations in each
orbital based on the results obtained by the method indicated
that for all compounds calculated the occupied states consist
mainly of sigma type bonding between metalloid p and Th d
sub(\$gamma\$) as well as \$pi\$ bonding between p and Th
dsut(\$epsilon\$).

EDB SUBJECT CATEGORY: EDB-360204;400702
INIS CATEGORY: B12

<260>
Inaba, H.; Matsui, T.; Naito, K. (Nagoya Univ. (Japan). Faculty
of Engineering)

Entropy change for the lambda-type transition in
nonstoichiometric U₄O_(9-y) EDB-76:053987; J. Nucl.
Mater. (JNUMA) 58:(1):115-116 (Oct 1975)

SUBJECT DESCRIPTORS: ELECTRIC CONDUCTIVITY;ENTROPY:
Q1;ORDER-DISORDER TRANSFORMATIONS;URANIUM OXIDES: T1

ABSTRACT: The Z-type jump of the electrical conductivity for
UO_{2.250}, UO_{2.240} and UO_{2.228} has been measured

<260>

<260> CONT.

by the conventional four probe method. The relative jump widths and entropy changes are presented.

EDE SUBJECT CATEGORY: EDB-360202

INIS CATEGORY: B24

<261>

Inaba, H.; Ono, S.; Naito, K. (Nagoya Univ. (Japan). Faculty of Engineering)

Phase transition in doped U_{4-y} . I. Heat capacity measurement AIX-08:310007; EDB-77:094397; J. Nucl. Mater. (JNUMA) 64:(1):189-194 (Jan 1977)

SUBJECT DESCRIPTORS: CATIONS; DOPED

MATERIALS; ENTHALPY; ENTROPY; HIGH TEMPERATURE; LOW

TEMPERATURE; MEDIUM TEMPERATURE; MULTICHARGED IONS; NIOBIUM

IONS; PHASE TRANSFORMATIONS; SPECIFIC HEAT: Q1; THORIUM

IONS; TRANSITION TEMPERATURE; URANIUM OXIDES: T1

ABSTRACT: Heat capacities of U_{4-y} doped with Th^{4+} and Nb^{5+} ions as impurities in the temperature range from 200 to 505 K by means of adiabatic calorimetry and the effect of doping on the phase transition of U_{4-y} was investigated. The lambda-type peak in the heat capacity curve becomes broadened on doping, resulting from U^{4+} and U^{5+} sites being disordered by the presence of impurity ions. The enthalpy change during the transition is proportional to the transition temperature for both doped and undoped samples, but at the same enthalpy change the transition temperature of doped samples is higher than that of undoped samples. The entropy changes during the transition are interpreted in terms of the average valence of the uranium.

EDE SUBJECT CATEGORY: EDB-360202

INIS CATEGORY: B24

<262>

Inaba, H.; Shimizu, H.; Naito, K. (Nagoya Univ. (Japan). Faculty of Engineering)

Lambda-type heat capacity anomalies in $U_{3-x}O_8$

AIX-08:310006; ERA-02:041852; EDB-77:094438; J. Nucl. Mater.

(JNUMA) 64:(1):66-70 (Jan 1977)

SUBJECT DESCRIPTORS: ENTHALPY; ENTROPY; HIGH TEMPERATURE; MEDIUM

TEMPERATURE; PHASE TRANSFORMATIONS; SPECIFIC HEAT:

Q1; TEMPERATURE DEPENDENCE; TRANSITION TEMPERATURE; URANIUM

OXIDES UO_2 : T1; X-RAY DIFFRACTION

<262> CONT.

ABSTRACT: The heat capacity of U_{3O_8} was measured in the temperature range from 310 to 970 K using an adiabatic scanning calorimeter. Three lambda-type heat capacity anomalies were observed at 483, 568 K and 850 K, indicating second order phase transitions. The enthalpy and entropy changes for these anomalies were 135, 148 and 314 J mol⁻¹, and 0.28, 0.26 and 0.37 J mol⁻¹ K⁻¹, respectively. A high temperature X-ray diffraction study has also been undertaken to confirm these phase transitions. The first anomaly at 483 K involves the structural change from orthorhombic to pseudohexagonal (strictly speaking orthorhombic). The origin of the other two phase transitions is not yet clear, but some order-disorder process would be involved in these phase transitions.

EDB SUBJECT CATEGORY: EDB-360204;050700

INIS CATEGORY: B24

<263>

Inman, D.; Graves, A.D.; Nobile, A.A.

Electrochemistry of molten salts EDB-77:044399; Electrochemistry (ECHME) (2):61-116 (1972)

SUBJECT DESCRIPTORS: DIFFUSION;ELECTRIC

CONDUCTIVITY;ELECTROCHEMISTRY:

Q1;ELECTRODES;ELECTROLYSIS;ELECTROLYTES:

Q2,Q3;HIGH-TEMPERATURE FUEL CELLS: M2;MASS

TRANSFER;METAL-NONMETAL BATTERIES: M3;MOLTEN SALTS: M1;REVIEWS

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-400400;300505;250903;300503

<264>

Inyushkina, T.L.; Marenkova, I.N.; Dzyubo, L.N.

Titanium dioxide solubility in melts of lithium- and alkaline earth metal chlorides AIX-08:317710;EDB-77:133385; Zh. Fiz. Khim. (ZFKHA) 49:(9):2458-2459 (Sep 1975)

SUBJECT DESCRIPTORS: ENTHALPY;MOLTEN SALTS: T2;SOLUBILITY:

Q1;SOLVENT PROPERTIES: Q2;STRONTIUM CHLORIDES;TEMPERATURE DEPENDENCE;TITANIUM OXIDES: T1

<264>

<265> CONT.

ABSTRACT: None

EED SUBJECT CATEGORY: EDB-400201

INIS CATEGORY: B12

<265>

Inyushkina, T.I.; Petukhova, L.P.; Kornilova, V.T. (Ural'skij
Politekhnikheskij Inst., Sverdlovsk (USSR))

Solubility of magnesium oxide in alkali metal chloride melt
EDE-76:049116; Zh. Neorg. Khim. (ZNOKA) 20:(4):1058-1060 (Apr
1975)

SUBJECT DESCRIPTORS: CESIUM CHLORIDES: T2;CHEMICAL REACTION
KINETICS;MAGNESIUM OXIDES: T1;MOLTEN SALTS;SOLUBILITY:
Q1;SOLVENT PROPERTIES: Q2;TEMPERATURE DEPENDENCE

ABSTRACT: None

EED SUBJECT CATEGORY: EDB-400201

INIS CATEGORY: B12

<266>

Inzenhofer, A. (Kraftwerk Union A.G., Erlangen (Germany, F.R.).
Abt. Brennelement-Technologie)

Influence of small additions to the preparation and the
properties of UO_2 ceramics. Part A. $\text{UO}_2/\text{Gd}_2\text{O}_3$
AIX-08:337296;ERA-03:000270;EDB-78:007611;
Staatsanz. Baden-Wuerttemb. (SAZRA) 109:(9):510-513 (1976)

SUBJECT DESCRIPTORS: BINARY MIXTURES;CHEMICAL PREPARATION:
Q1,Q2;GADOLINIUM OXIDES: T2;LATTICE PARAMETERS;MELTING
POINTS: Q1;PHASE DIAGRAMS: Q1,Q2;REVIEWS;SINTERING;SOLID
SOLUTIONS;THERMAL EXPANSION;URANIUM DIOXIDE: T1

ABSTRACT: The effects of oxide additions on the preparation and
burn-up behaviour of UO_2 fuels is presented within the
framework of a literature survey. In the present part A, the
addition of Gd_2O_3 (max. 5 wt.%) is dealt with.
Firstly a few crystallographic data are given: phase
diagrams, lattice parameters, coefficient of expansion. The
preparation of UO_2 - Gd_2O_3 sinter bodies is
then treated. The melting point of UO_2 is somewhat

<267> CONT.

lowered by the addition of Gd_{2O_3} , the lattice constants become smaller.

EEB SUBJECT CATEGORY: EDB-050700;360201;360202

INIS CATEGORY: E23

<267>

Inzenhofer, A. (Kraftwerk Union A.G., Erlangen (Germany, F.R.).
Abt. Brennelement-Technologie)

Influence of small additions on the preparation and the properties of UO_2 ceramics. Part B. The UO_2 - TiO_2 system

AIX-08:337297;ERA-03:000267;EDB-78:013800; Staatsanz.
Baden-Wuerttemb. (SAZRA) 109:(12):728-731 (Dec 1976)

SUBJECT DESCRIPTORS: BINARY MIXTURES;CHEMICAL
PREPARATION;CRYSTAL
DEFECTS;DENSITY;DIFFUSION;EUTECTICS;FLEXURAL STRENGTH;GRAIN
GROWTH;MELTING POINTS: Q1;PHASE DIAGRAMS:
Q1,Q2;SINTERING;SOLID SOLUTIONS;TITANIUM OXIDES: T2;URANIUM
DIOXIDE: T1

ABSTRACT: In the present part B of the literature survey on the effect of oxide additions on the behaviour of UO_2 , the influence of titanium oxide is dealt with. Firstly, a few important basic data of the two-phase system UO_2 - TiO_2 are given: phase diagram, diffusion constants, flexural strength. The sinter behaviour and the grain growth are then treated. In particular, the question of at which temperature a molten phase forms and this then contributes to an increase of the sintering rate is answered.

EEB SUBJECT CATEGORY: EDB-050700;360204;360202

INIS CATEGORY: B23

<268>

Iskandarov, K.I.; Litvinov, Yu.G.; Il'yasov, I.I. (Zaochnyj
Inst. Sovetskoy Torgovli, Rostov-na-Donu (USSR))

LiBr, RbBr, CsBr ternary system AIX-08:295305;EDB-77:056216; Zh.
Necrg. Khim. (ZNOKA) 21:(7):1990-1992 (Jul 1976)

SUBJECT DESCRIPTORS: CESIUM BROMIDES: T1;COMPLEXES;HIGH
TEMPERATURE;LITHIUM BROMIDES: T2;MELTING POINTS;PHASE
DIAGRAMS: Q1,Q2,Q3;QUANTITY RATIO;RUBIDIUM BROMIDES: T3;SOLID
SOLUTIONS

<268>

<268> CONT.

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-400201

INIS CATEGORY: B12

<269>

Ito, Y.; Matsunaga, M.; Yoshizawa, S. ** Selman, J.R. (ed.)

Studies on the negative alloy electrode In Proceedings of the symposium and workshop on advanced battery research and design, March 22--24, 1976 ** ERA-02:003289; EDB-76:091634; (1976) ANL--76-8

SUBJECT DESCRIPTORS: ALKALI METALS; ALUMINIUM ALLOYS; ANODES: Q1; EFFICIENCY; ELECTRIC POTENTIAL; HIGH TEMPERATURE; LITHIUM ALLOYS; LITHIUM-CHLORINE BATTERIES: M1, Q2; MOLTEN SALTS; OFF-PEAK ENERGY STORAGE: T2; PHASE DIAGRAMS; TIN ALLOYS; ZINC ALLOYS

ABSTRACT: Alkali metal alloy is a promising negative electrode for a high-performance battery. Energy density and power density of such alloy electrodes can be discussed in relation to the phase diagram. Liquid alloy is preferable when cycle life is exaggerated, and solid alloy is preferable from the cell design aspect. Experimental data to support such conclusions are shown in this paper. 15 figures, 3 tables.

EDB SUBJECT CATEGORY: EDB-250903

<270>

Ivanova, I.A.; Petrova, M.A.; Podkolzina, I.G.

LiF-ErF₃ system AIX-08:286055; EDB-77:044213; Zh. Neorg. Khim. (ZNOKA) 20:(8):2292-2293 (Aug 1975)

SUBJECT DESCRIPTORS: BINARY MIXTURES; CRYSTALLIZATION; DIFFERENTIAL THERMAL ANALYSIS; ERBIUM FLUORIDES: T1; EUTECTICS; LITHIUM FLUORIDES: T2; MICROSCOPY; PHASE DIAGRAMS: Q1, Q2; REFRACTION; X-RAY DIFFRACTION

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-400201

INIS CATEGORY: B12

<271>

Iyer, R.H.; Natarajan, P.R. (eds.), Bhabha Atomic Research Centre, Bombay (India)

In Radiochemistry Division annual progress report for 1973 **
AIX-07:270256;EDB-77:044468; (1975) BARC--821

ABSTRACT: The R and D work carried out in the Radiochemistry Division of the Bhabha Atomic Research Centre, Bombay, during 1973 is reported under the following topical headings: process chemistry (of transuranic elements), radioanalytical chemistry and services, chemical quality control of Pu fuels, heavy element chemistry, nuclear chemistry and instrumentation. The major highlights are: preparation of ^{238}Pu , non-destructive estimation of Pu by X-ray fluorescence and gamma counting, determining impurities in trace amounts in uranium and plutonium fuels, determination of solubility of PuF_3 in molten fluoride mixtures as a part of the chemical development program for the molten salt reactor concept, studies on correlation between average total kinetic energy, fission asymmetry and shell structure.

EDB SUBJECT CATEGORY: EDB-400700;400103
INIS CATEGORY: B13

<272>

Jacobs, H. ** Commission of the European Communities, Brussels (Belgium)

The calculation of the consequences of sodium/fuel interactions with allowance for temperature gradients in the sodium in Second specialist meeting on sodium/fuel interaction in fast reactors, Ispra, 21-23 November 1973. Sponsored by the Committee on the Safety of Nuclear Installations, OECD Nuclear Energy Agency, Paris, France ** EDB-76:052697; ([nd]) EUR--5309

SUBJECT DESCRIPTORS: B CODES: Q3;COMPUTER CODES: T3;FUEL-COOLANT INTERACTIONS: T,Q1;FUSION HEAT;HEAT TRANSFER;LMFBR TYPE REACTORS: T1;NUCLEAR FUELS;SODIUM;TEMPERATURE DEPENDENCE;THERMAL CONDUCTIVITY

ABSTRACT: The state of the code BRENDA is described together with an outlook on its further development. The code is applied to study the effect of various assumptions used in sodium fuel interaction codes in connection with sodium thermal conductivity, heat transfer during phase B, fuel

<272>

<273> CONT.

latent heat of fusion, decoupled heat transfer calculations,
and incompressible sodium slugs.

EDB SUBJECT CATEGORY: EDB-220900;210500
INIS CATEGORY: F51

<273>

Jacucci, G.; Taylor, R. (Paris-11 Univ., 91 - Orsay (France).
Centre de Calcul); Klein, M.L.

Static structure factor of liquid Li

AIX-07:269600;EDB-77:013024; Solid State Commun. (SSCOA)
19: (7):657-659 (Jul 1976)

SUBJECT DESCRIPTORS: BOUNDARY CONDITIONS;INTERATOMIC FORCES:
Q1;INTERMOLECULAR FORCES;IONS;LIQUID METALS;LITHIUM:
T1;MATHEMATICAL MODELS;NEUTRON DIFFRACTION;POTENTIAL ENERGY

ABSTRACT: The classical Molecular Dynamics method has been used to calculate the structure factor $S(Q)$ of liquid Li and this quantity is compared with that measured by neutron scattering. A system of 1458 Li atoms has been employed interacting, via periodic boundary conditions, with a recently derived interionic pair potential. The good agreement between the calculation and experiment is taken to indicate that the explicit many body interionic forces are not important in liquid Li.

EDB SUBJECT CATEGORY: EDB-360102
INIS CATEGORY: A13

<274>

James, S.D.; DeVries, L.E.

Structure and anodic discharge behavior of lithium-boron alloys in the liCl-KCl eutectic melt. EDB-76:080765; J. Electrochem. Soc. (JESOA) 123: (3):321-327 (Mar 1976)

SUBJECT DESCRIPTORS: ANODES: Q1;FORON ALLOYS;ELECTRIC BATTERIES:
T1;ELECTRODES;EUTECTICS;HIGH TEMPERATURE;LITHIUM BASE
ALLCYS;LITHIUM CHLORIDES: T2;PHASE DIAGRAMS;POTASSIUM
CHLORIDES: T3;THERMODYNAMIC PROPERTIES: Q2,Q3

ABSTRACT: Anodic discharge curves in the LiCl--KCl eutectic melt have been obtained from two lithium--boron alloys (70 and 80 weight per cent Li) between 673°K and 873°K (400°K and 600°K and from 2 to 8 A/cm². The behavior of Li contained in wicks of porous Ni was also studied for comparison. Calculated Li contents for Li--B alloy anodes, at the main break in curvature of discharge

<274> CONT.

curves, clustered closely around 0.67 mole fraction Li. The authors consider this strong evidence for the existence of a compound Li_2B in this alloy system. Between 748⁰ and 873⁰ (475⁰ and 600⁰) and up to at least 8 A/cm², there is no diffusional limitation to the anodic extraction of Li from the Li--B alloys.

EDB SUBJECT CATEGORY: EDB-250903;360102

<275>

James, S.D.; DeVries, L.E., Naval Surface Weapons Center, Silver Spring, Md. (USA)

In Structure of lithium--boron alloys and their performance as potential anodes for high power molten salt batteries.

Progress report, 1 Aug 1974--30 Jun 1975 **

ERA-02:013693;EDB-77:018209; (1 Mar 1976) AD-A--025202

SUBJECT DESCRIPTORS: ANODES: Q1,Q2; BORON ALLOYS; ELECTRIC BATTERIES: T1; ELECTROCHEMISTRY; EUTECTICS; HIGH TEMPERATURE; INTERMETALLIC COMPOUNDS; LITHIUM BASE ALLOYS; LITHIUM CHLORIDES; LITHIUM-SULFUR BATTERIES: T2; MOLTEN SALTS; NICKEL; PERFORMANCE; POTASSIUM CHLORIDES

ABSTRACT: Anodic discharge curves in the LiCl--KCl eutectic melt have been obtained for two lithium--boron alloys (70 and 80 wt % Li) between 673 and 873 K (400 and 600 C) and from 2 to 8 A/sq cm. The behavior of Li contained in wicks of porous Ni was also studied for comparison. Calculated Li contents for Li--B alloy anodes, at the main break in curvature of discharge curves, clustered closely around 0.67 mole fraction Li. This is considered strong evidence for the existence of a compound Li_2B in this alloy system. Between 748 and 873 K (475 and 600 C) and up to at least 8 A/sq cm, there are no diffusional limitations to the anodic extraction of Li from the Li--B alloys. The shape of discharge curves, the lack of anodic diffusional limitation, and the longer low-V plateaus of the alloy richer in Li are consistent with the following picture of the discharge process. The discharging material behaves like a sponge of solid Li_2B compound containing a Li-rich, Li--B liquid alloy, probably a conjugate phase to the Li_2B compound. Lithium anodizes into the melt at constant V out of this liquid alloy until it is wholly transformed to solid Li_2B . This process constitutes the main plateau of alloy discharge curves. At this point (the main break in discharge curves) voltage rises to values where tighter-bound Li can be anodized out of the decomposing compound. The Li--B alloy electrodes are markedly superior to the Li--Ni wicks, both in gravimetric coulombic capacity and in the ability to retain Li at high temperatures. Useful

<275>

<275> CONT.

coulombic capacities of the Li--B alloys ranged up to 7700 ccoulombs per g of (Li^{sup} + \$B) at about 3 V versus a chlorine reference electrode. (GRA)

EDB SUBJECT CATEGORY: EDB-250903

<276>

Janz, G.J.; Allen, C.B.; Downey, J.R. Jr.; Tomkins, R.P.T.,
Rensselaer Polytechnic Inst., Troy, N.Y. (USA). Cogswell Lab.

In EUTECTIC DATA: safety, hazards, corrosion, melting points,
compositions, and bibliography **

ERA-U2:034550;EDB-77:073948; (Jul 1976) TID--27163-P1

SUBJECT DESCRIPTORS: BIBLIOGRAPHIES;CHEMICAL
COMPOSITION;CORROSIVE EFFECTS;DATA COMPILATION:
Q1,Q2;EUTECTICS: T1;HAZARDS;HEAT STORAGE: T;MELTING
POINTS;MOLTEN SALTS: T2;SAFETY

ABSTRACT: The present compilation provides an authoritative compendium of properties relative to safety, hazards, corrosion, melting points, and compositions of molten salt eutectic mixtures. The scope of the single salts data in this compilation encompasses the end-components of the eutectic mixtures reported. Each section contains a systems index and bibliography. The bibliographies in Pt. 1 and Pt. 2 total approximately 4000 entries. More than 6000 eutectic entries are included in the two parts.

EDB SUBJECT CATEGORY: EDB-250600

<277>

Janz, G.J.; Reeves, R.D.

Molten-salt electrolytes--transport properties EDB-76:052816;
Adv. Electrochem. Electrochem. Eng. (AEEEE) 5::137-204 (1967)

SUBJECT DESCRIPTORS: DENSITY;DIFFUSION;ELECTRIC BATTERIES:
T1;ELECTRIC CONDUCTIVITY;ELECTROCHEMISTRY: Q2;ELECTROLYTES:
Q1;MASS TRANSFER;MOLTEN SALTS: T2;SOLVENTS;VISCOSITY

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-250902;400400

<278>

Jayadevan, N.C.; Chadha, A.; Chackraburttty, D.M., Bhabha Atomic Research Centre, Bombay (India)

In Preparation and characterization of higher uranates(VI) of thallium(I) ** AIX-08:292339;EDB-77:075233; (1976) BARC--866

SUBJECT DESCRIPTORS: CHEMICAL COMPOSITION;CHEMICAL PREPARATION: Q1,Q2;CRYSTAL STRUCTURE: Q1,Q2;INFRARED SPECTRA;STRUCTURAL CHEMICAL ANALYSIS;THALLIUM COMPOUNDS: T1;THERMAL ANALYSIS;URANATES: T2;URANIUM OXIDES U3O8;URANIUM TRIOXIDE;X-RAY DIFFRACTION

ABSTRACT: Two new thallium(I) uranates, Tl_2UO_4 and $\text{Tl}_2\text{U}_7\text{O}_{22}$ have been prepared by heating Tl_2CO_3 with UO_3 or U_3O_8 at 600°C in air. The chemical compositions were confirmed from the analysis of uranium and thallium on solutions of the pure compounds in very dilute nitric acid. The tetrauranate is hexagonal with cell constants of $a = 14.315$ (5) Å and $c = 14.280$ (5) Å while the heptauranate is indexed on an orthorhombic cell of dimension $a = 6.950$ (4) Å, $b = 19.609$ (7) Å. Both are isostructural with the corresponding potassium and rubidium compounds. The compounds were further characterized by infrared and thermal methods. $\text{Tl}_2\text{U}_7\text{O}_{22}$ decomposes in air above 850°C to form Tl_2UO_4 and U_3O_8 which decomposes into the components Tl_2O and U_3O_8 only above 1150°C. Reaction of $\text{Tl}_2\text{U}_4\text{O}_{13}$ with excess UO_3 gives $\text{Tl}_2\text{U}_7\text{O}_{22}$ at 600°C.

EDB SUBJECT CATEGORY: EDB-400702

INIS CATEGORY: B12

<279>

Jefferson, J.H.; Stevens, K.W.H. (Nottingham Univ. (UK))

Statistical mechanics of intermediate-valence rare earth compounds AIX-07:266496;EDB-77:007340; J. Phys., C (London) (JFSOA) 9: (11):2151-2165 (14 Jun 1976)

SUBJECT DESCRIPTORS: CURIE-WEISS LAW;FREE ENTHALPY;MAGNETIC SUSCEPTIBILITY;PERTURBATION THEORY;RARE EARTH COMPOUNDS: M1;STATISTICAL MECHANICS: Q1;VALENCE

ABSTRACT: Using thermodynamic perturbation theory the statistical mechanics of intermediate-valence rare earth compounds is investigated. The Gibbs' free energy is calculated to second order in the Anderson mixing interaction and is shown to diverge logarithmically as T tends to 0. At high temperatures the susceptibility is approximately of the

<279>

<279> CONT.

Curie-Weiss form $\chi = C/(T + T_{\text{sub}}(F))$, where $kT_{\text{sub}}(F)$ approximately δ , the 'width parameter' associated with interconfigurational fluctuations.

EDB SUBJECT CATEGORY: EDB-360603

INIS CATEGORY: A13

<280>

Joergensen, C.K. (Geneva Univ. (Switzerland). Dept. de Chimie Minerale et Analytique) ** Dunitz, J.D. (ed.) (Eidgenossische Technische Hochschule, Zurich (Switzerland). Lab. fuer Organische Chemie); Hemmerich, P. (ed.) (Konstanz Univ. (F.R. Germany). Fachbereich Biologie); Holm, R.H. (ed.) (Massachusetts Inst. of Tech., Cambridge (USA). Dept. of Chemistry); Ibers, J.A. (ed.) (Northwestern Univ., Evanston, Ill. (USA). Dept. of Chemistry); Joergensen, C.K. (ed.) (Geneva Univ. (Switzerland). Dept. de Chimie Minerale et Analytique); Neillands, J.B. (ed.) (California Univ., Berkeley (USA). Dept. of Biochemistry); Reinen, D. (ed.) (Marburg Univ. (F.R. Germany). Fachbereich Chemie); Williams, R.J.F. (ed.) (Oxford Univ. (UK). Inorganic Chemistry Lab.)

Partly filled shells constituting anti-bonding orbitals with higher ionization energy than their bonding counterparts in Rare earths ** Structure and Bonding,
AIX-07:257186;EDB-77:007477; (1975) Springer

SUBJECT DESCRIPTORS: CHEMICAL BONDS;COMPARATIVE EVALUATIONS;COVALENCE;DATA;ELECTRONIC STRUCTURE: Q1;EXCITED STATES;F STATES;FERMI LEVEL;IONIZATION;LCAO METHOD;PHOTOELECTRON SPECTROSCOPY;RARE EARTH COMPOUNDS;RARE EARTHS: T1;REVIEWS;SPECTRA;TABLES

ABSTRACT: In a first part, one-electron substitutions in the lowest configuration are treated by monatomic entities, 4f-group compounds, and comparison with iodide, xenon and cesium, in a second part, the Copenhagen principle of final states are described by photo-electron spectra of metallic lanthanides and of lanthanide compounds, and by the question of the number of 4f electrons as an integer, and in a third part, the partly covalent bonding is treated by means of the variation principle and the LCAO model, the importance of kinetic energy, and the comparison with copper(II) complexes.

EDB SUBJECT CATEGORY: EDB-400201

INIS CATEGORY: B12

<281>

Johnson, C.E.; Blackburn, P.E.; Carroll, K., Argonne National
Lab., Ill. (USA)

In Chemical Engineering Division fuels and materials chemistry.
Annual report, July 1975--June 1976 **
ERA-02:042665;EDB-77:098236; (1976) ANL--76-103

SUBJECT DESCRIPTORS: BURNUP;CERIUM OXIDES;CESIUM OXIDES;CHEMICAL
REACTIONS: Q1,Q2,Q3;DATA COMPILATION:
Q5;ENTHALPY;ENTROPY;EVAPORATION;FISSION PRODUCTS:
T3;GASES;LITHIUM;LMFBR TYPE REACTORS: T4;MOLYBDENUM
OXIDES;NUCLEAR FUELS: T5,Q4;PLUTONIUM CARBIDES: T7;PLUTONIUM
DIOXIDE: T1;STOICHIOMETRY;THERMODYNAMIC PROPERTIES:
Q6,Q7,Q1,Q2,Q3;TRAPPING: Q8;TRITIUM IONS: T8;URANIUM
CARBIDES: T6;URANIUM DIOXIDE: T2;URANIUM OXIDES;VERY HIGH
TEMPERATURE

ABSTRACT: Measurements of the oxygen potential for
urania-plutonia mixtures show an increase with increasing
plutonium content. The oxygen potential model for
uranium-plutonium oxide has been extended to include the
effects of burnup on oxygen potential. The model predicts an
increase in oxygen potential with burnup for both enriched
and nonenriched hypostoichiometric fuel, with the nonenriched
fuel exhibiting a higher rate of increase in oxygen potential
with burnup than does the enriched fuel. Knudsen
effusion-mass spectrometric studies of the urania-molybdena
system have shown that molybdena has limited solubility in
urania at high temperatures and that the solid solutions
exhibit positive deviations from ideal behavior. The kinetics
of the reaction of gaseous cesium with hyperstoichiometric
urania is being studied. The reaction rate has been found to
depend linearly on the initial U/O ratio of the urania.
Measurements of the carbon activity above a U--Pu--C system
have shown segregation to monocarbide and sesquicarbide
phases, with the sesquicarbide phase exhibiting a higher
plutonium content. A handbook of thermophysical and
mechanical properties of carbide and nitride fuels has been
compiled to serve as an authoritative source of properties
data for persons working on advanced fuels. A limited number
of measurements were made to determine the trapping
efficiency of lithium for tritium ions. A trapping efficiency
of lithium for tritium was determined to be about 90 percent.

EDB SUBJECT CATEGORY: EDB-050700;360200

<282>

<282>

Johnsen, C.E.; Foster, M.S.

Phase equilibrium studies of lithium halide-containing electrolytes EDB-77:131874; J. Electrochem. Soc. (JESOA) 116:(11):1612-1613 (Nov 1969)

SUBJECT DESCRIPTORS: CHEMICAL PROPERTIES; ELECTRIC BATTERIES: T1; ELECTROCHEMICAL CELLS; ELECTROLYTES: Q1; LITHIUM; LITHIUM CHLORIDES; LITHIUM HALIDES: T2; LITHIUM IODIDES; MELTING POINTS; PHASE STUDIES: Q2; PHYSICAL PROPERTIES; POTASSIUM IODIDES

ABSTRACT: The application of fused-salt mixtures as electrolytes in electrochemical cells has resulted in increased interest in their physical and chemical properties. A knowledge of these properties is important in selecting appropriate electrolytes, since the choice depends on a variety of factors. One of the salt systems being considered for use as an electrolyte in lithium-anode galvanic cells is LiCl--LiI--KI. This system was chosen for investigation after a preliminary examination had indicated that it had a minimum melting point of 264 C, one of the lowest observed for ternary alkali halide mixtures. The good agreement between the calculated and experimental values and the few experiments required to obtain the necessary data makes it clear that the conformal ionic solution theory offers significant saving of experimental effort in the selection of low-melting-point electrolyte mixtures.

EDB SUBJECT CATEGORY: EDB-250901;360602

<283>

Kabankova, N.N.; Moskvitina, E.N.; Kuzyakov, Yu.Ya. (Moskovskii Gosudarstvennyi Univ. (USSR))

Absorption spectrum of HfCl molecule

AIX-07:265060; EDB-77:013460; Vestn. Mosk. Univ., Ser. II. Khim. (VBUKA) 16:(2):232-233 (Mar 1975)

SUBJECT DESCRIPTORS: ABSORPTION SPECTRA: Q1; DATA; HAFNIUM CHLORIDES: T1; VIBRATIONAL STATES

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-400201
INIS CATEGORY: B12

<284>

Kadcchnikov, V.A.; Lebedev, V.A.; Nichkov, I.F.; Raspopin, S.P.

Thermodynamic properties of the U-Sn-alloys riched with tin
 AIX-08:283300;EDB-77:043771; Izv. Akad. Nauk SSSR, Met.
 (IZNMA) (4):67-69 (Jul 1976)

SUBJECT DESCRIPTORS: BINARY ALLOY SYSTEMS;EQUATIONS;HIGH
 TEMPERATURE;LIQUID METALS;MEDIUM TEMPERATURE;REACTION
 KINETICS;SOLUBILITY;TABLES;TEMPERATURE
 DEPENDENCE;THERMODYNAMIC PROPERTIES: Q1,Q2;THERMODYNAMICS;TIN
 ALLCOYS: T1;URANIUM ALLOYS: T2;VERY HIGH TEMPERATURE

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-360104

INIS CATEGORY: B22

<285>

Kakihana, H.; Yato, Y. (Tokyo Inst. of Tech. (Japan). Research
 Lab. of Nuclear Reactor)

Calculation of the reduced partition function ratio for uranium
 hexafluoride AIX-07:259569;EDB-77:013689; Bull. Tokyo Inst.
 Technol. (Engl. Ed.) (BTITA) (127):71-81 (Mar 1975)

SUBJECT DESCRIPTORS: CORIOLIS FORCE;ISOTOPE EFFECTS;ISOTOPE
 SEPARATION;MOLECULAR STRUCTURE;PARTITION FUNCTIONS:
 Q1;SPECTRAL SHIFT;SYMMETRY;URANIUM 235;URANIUM 238;URANIUM
 HEXAFLUORIDE: T1;VIBRATIONAL STATES

ABSTRACT: The reduced partition function ratio of UF_6 is
 calculated by determining the complete general valence force
 field from the Coriolis zeta constant data together with the
 observed vibrational frequencies. The disagreement in the
 previous calculations is due to the ambiguity that arises in
 the force field determination for the octahedral molecules.
 The calculation leads to the $\ln(s/s') f(^{238}UF_6 / ^{235}UF_6)$
 value of $(2.346 \pm 0.134) \times 10^{-3}$ at $300^\circ K$ in reasonable agreement with a previous
 estimate by Kakihana who used the Badger's relation to
 determine the force field.

EDB SUBJECT CATEGORY: EDB-400702

INIS CATEGORY: B12

<286>

<286>

Kalina-Khorokhorova, L.A.; Simonova, M.V.; Kalinin, S.N.
(Rostovskij-na-Donu Inzhenernc-Stroitel'nyj Inst. (USSR))

Reaction between sodium, rubidium and strontium chlorides
AIX-08:286060;EDB-77:044249; Khim. Khim. Tekhnol. (Alma-Ata)
(KKTKA) 19:(4):664 (Apr 1976)

SUBJECT DESCRIPTORS: CHEMICAL COMPOSITION;CHEMICAL REACTIONS:
Q1,Q2,Q3;ENTROPY;EUTECTICS;FUSION HEAT;MELTING
POINTS;RUBIDIUM CHLORIDES: T1;SODIUM CHLORIDES: T2;STRONTIUM
CHLORIDES: T3

ABSTRACT: None

ELP SUBJECT CATEGORY: EDB-400201
INIS CATEGORY: B12

<287>

Kalinyuk, N.N. (AN Ukrainskoj SSR, Kiev. Inst. Ehlektrosvarki)

Hydrogen solubility in liquid zirconium
AIX-08:340196;EDB-78:016857; Izv. Vyssh. Uchebn. Zaved.,
Tsvet. Metall. (IVUTA) (1):134-138 (Jan 1976)

SUBJECT DESCRIPTORS: HYDROGEN: T1;LIQUID METALS;MAGNETIC
FIELDS;SOLUBILITY: Q1,Q2;TEMPERATURE DEPENDENCE;THERMODYNAMIC
PROPERTIES;VERY HIGH TEMPERATURE;ZIRCONIUM: T2

ABSTRACT: The hydrogen solubility in liquid zirconium was studied in the temperature range of 1965 to 2485 deg C with the hydrogen partial pressure of 7.9 to 43.8 mm Hg. The 'liquid zirconium - hydrogen' system has been found to obey the Sieverts law within the temperature range studied. A comparison of the free energies of hydrogen dissolving in liquid and solid zirconium at the melting point shows that the free energy value, and the hydrogen solubility as well, in the solid-to-liquid zirconium transition increases due to an abrupt increase of the system entropy. This process is accompanied by some decrease in the heat of hydrogen dissolving in zirconium. The data obtained indicate that in liquid zirconium, as well as in molten titanium and niobium, the hydrogen solubility decreases with the temperature growth and the boiling point it will be equal to zero.; in the process of zirconium melting the hydrogen solubility in it is growing in steps. At the melting point the hydrogen

<288> CONT.

solubility in liquid zirconium is 1.7 to 2.5 times as high as in solid zirconium.

EDB SUBJECT CATEGORY: EDB-360104;360105

INIS CATEGORY: B22

<288>

Kaloev, N.I.; Egerev, O.I.; Dzeranova, K.B.; Kulova, I.K.
(Gruzinskij Politekhnikheskij Inst., Tbilisi (USSR))

Phase diagrams of $\text{BiBr}_{\text{sub } 3}\text{-BiI}_{\text{sub } 3}$ system

AIX-08:336946;EDB-77:146483; Zh. Neorg. Khim. (ZNOKA)
21:(1):290-292 (Jan 1976)

SUBJECT DESCRIPTORS: BINARY MIXTURES;BISMUTH BROMIDES:

T1;BISMUTH IODIDES: T2;DIFFERENTIAL THERMAL

ANALYSIS;ENTHALPY;MELTING POINTS;MOLTEN SALTS;PHASE DIAGRAMS:
Q1,Q2;SOLID SOLUTIONS

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-360602

INIS CATEGORY: B12

<289>

Kanashvili, N.V.; Morozov, A.I. (AN SSSR, Moscow. Inst. Obshchej
i Neorganicheskoy Khimii)

Thermal investigation of $\text{NdCl}_{\text{sub } 3}\text{-AlCl}_{\text{sub } 2}\text{-KCl}$ and
 $\text{NdCl}_{\text{sub } 3}\text{-AlCl}_{\text{sub } 3}\text{-CsCl}$ system

AIX-08:336948;EDB-77:146787; Zh. Neorg. Khim. (ZNOKA)
22:(2):464-470 (Feb 1977)

ABSTRACT: The systems $\text{NdCl}_{\text{sub } 3}\text{-AlCl}_{\text{sub } 3}\text{-MCl}$ (M-K, Cs) have been studied by the DTA method with the use of the visual-polythermal technique. On the liquidus surface of the system $\text{NdCl}_{\text{sub } 3}\text{-AlCl}_{\text{sub } 3}\text{-KCl}$ the authors found seven fields of primary crystallization corresponding to primary isolation of KCl, $\text{KAlCl}_{\text{sub } 4}$, $\text{AlCl}_{\text{sub } 3}$, $\text{NdCl}_{\text{sub } 3}$, $\text{K}_{\text{sub } 2}\text{NdCl}_{\text{sub } 5}$ and $\text{K}_{\text{sub } 3}\text{NdCl}_{\text{sub } 6}$. The system has five four-phase equilibria occurring at three points of triple eutectics and at two peritectic points. In the system $\text{NdCl}_{\text{sub } 3}\text{-AlCl}_{\text{sub } 3}\text{-CsCl}$ there are six fields of initial crystallization, corresponding to the isolation of $\text{NdCl}_{\text{sub } 3}$, $\text{NdCl}_{\text{sub } 3}\text{x AlCl}_{\text{sub } 3}$, $\text{AlCl}_{\text{sub } 3}$, $\text{CsAlCl}_{\text{sub } 4}$,

<289>

<290> CONT.

CsCl , $\text{Cs}_{3\text{NdCl}_6}$ from the melt and four four-phase equilibria at three eutectic and one peritectic points.

EDB SUBJECT CATEGORY: EDB-400201

INIS CATEGORY: B12

<290>

Kang, T.; Castanet, R. (Centre National de la Recherche Scientifique, 13 - Marseille (France). Centre de Recherches de Microcalorimetrie et de Thermochimie)

Thermodynamic study of the binary system copper-indium. I.
Calorimetric study AIX-08:309827;EDB-77:093958; J.
Less-Common Met. (JCOMA) 51:(1):125-135 (Jan 1977)

SUBJECT DESCRIPTORS: BINARY ALLOY SYSTEMS; CALORIMETRY; CHEMICAL COMPOSITION; COPPER ALLOYS: T1; FORMATION HEAT: Q1, Q2; HIGH TEMPERATURE; INDIUM ALLOYS: T2; LIQUID METALS; PHASE DIAGRAMS; VERY HIGH TEMPERATURE

ABSTRACT: The integral enthalpy of formation $h_{\text{sup}}(f)$ of Cu-In liquid alloys has been measured by direct reaction calorimetry for $0 < x_{\text{sub}}(\text{In}) < 1$ and $903 < T(\text{K}) < 1348$. The excess heat capacity $C_{\text{sub}}(p)_{\text{sup}}(E)$ which has been deduced is temperature independent. The $h_{\text{sup}}(f)$ minimum and the $C_{\text{sub}}(p)_{\text{sup}}(E)$ maximum, at $x_{\text{sub}}(\text{In})$ approximately 0.75, are interpreted in terms of a local order in the melts. The integral enthalpy of formation of the β , γ and ϵ solid phases has also been measured by dissolution calorimetry at 904 and 941 K. The entropy of melting of the γ phase has been determined. It is pointed out that many systems do not have zero excess heat capacity; thus the mean values of $h_{\text{sup}}(f)$ as calculated from measurements at different temperatures are incorrect.

EDB SUBJECT CATEGORY: EDB-360101;360104

INIS CATEGORY: B22

<291>

Kang, T.; Kehiaian, H.V.; Castanet, R. (Centre National de la Recherche Scientifique, 13 - Marseille (France). Centre de Recherches de Microcalorimetrie et de Thermochimie)

Thermodynamic study of the binary system copper--indium. II.
Potentiometric study AIX-08:340197;EDB-78:016828; J.
Less-Common Met. (JCOMA) 53:(2):153-166 (Jun 1977)

SUBJECT DESCRIPTORS: BINARY ALLOY SYSTEMS; COPPER ALLOYS: T1; ELECTROLYTIC CELLS; ELECTROMOTIVE FORCE; ENTROPY: Q1, Q2; FORMATION FREE ENERGY; FORMATION FREE ENTHALPY:

<291> CONT.

Q1,Q2;INDIUM ALLOYS: T2;LIQUID METALS;POTENTIOMETRY;SPECIFIC
HEAT: Q1,Q2;THERMODYNAMIC ACTIVITY;VERY HIGH TEMPERATURE

ABSTRACT: The electrochemical cell with solid electrolytes
-Mo/In(l), In₂O₃(s)/ZrO₂, CaO/air, Pt/Pt,
air/CaO, ZrO₂/In-Cu(l), In₂O₃(s)/Mo+ was
used to measure the activity of indium in liquid Cu-In alloys
at nine concentrations in the temperature range from 1020 to
1389 K. The procedure was checked by determining the enthalpy
of formation of indium oxide In₂O₃. Combining the
present results with those obtained by direct calorimetry,
thermodynamically consistent values were proposed for the
free energy, entropy, enthalpy and heat capacity of liquid
Cu-In alloys.

EDE SUBJECT CATEGORY: EDB-360104
INIS CATEGORY: B22

<292>

Kanno, M. (Tokyo Univ. (Japan). Faculty of Engineering);
Yamagami, S.

Chronopotentiometry of U(III) and U(IV) in molten LiCl-KCl
eutectic AIX-07:243901;EDB-76:075888; Denki Kagaku (DNKKA)
43:(3):131-137 (Mar 1975)

SUBJECT DESCRIPTORS: DIFFUSION;EUTECTICS;HIGH
TEMPERATURE;LITHIUM CHLORIDES;MIXTURES;MOLTEN SALTS;POTASSIUM
CHLORIDES: T1;POTENTIOMETRY: Q1;QUANTITATIVE CHEMICAL
ANALYSIS;TIME DEPENDENCE;URANIUM CHLORIDES

ABSTRACT: For the needs to kinetic studies of the transfer
processes in liquid metal-molten salt system, a
chronopotentiometric investigation for the quantitative
analysis of U(III) and U(IV) in LiCl-KCl eutectic salt has
been investigated. It was found that chronopotentiometry of
U(III) and U(IV) obeyed Sand's equation, and U(III) and U(IV)
have been determined with an accuracy of $\pm 3\%$ when there is
no oxygen and water contamination at 450°C both in
their individual solutions and mixed solutions. According to
Sand's equation, the product of current density i and $1/2$
power of transition time τ should be constant with
different i 's. However, it was observed to hold within a
limited range of current densities. Positive and negative
deviations from the constancy were explained. The diffusion
coefficients of U(III) and U(IV) in LiCl-KCl eutectic salt at
 400°C have been determined by chronopotentiometry as
follows, $D_{\text{sub}}(u(i i i)) = 4.38 \cdot 10^{\text{sup } -4} \exp(-5,940/RT)$

<292>

<292> CONT.

$\text{cm}^2/\text{sec} \text{ D}_{\text{sub}}(u(i, v)) = 1.73 \cdot 10^{-3} \exp(-7,430/RT)$
 $\text{cm}^2/\text{sec}.$

EDE SUBJECT CATEGORY: EDB-400702
 INIS CATEGORY: B12

<293>

Kanno, M. (Tokyo Univ. (Japan). Faculty of Engineering);
 Yamagami, S.

Electrochemical behavior of U(III) and U(IV) in molten LiCl-KCl
 eutectic AIX-07:262534; EDB-77:019438; Denki Kagaku (DNKKA)
 43: (5):253-257 (May 1975)

SUBJECT DESCRIPTORS: ELECTROCHEMISTRY: Q1; ELECTROMOTIVE
 FORCE; EUTECTICS; HIGH TEMPERATURE; LITHIUM
 CHLORIDES; MIXTURES; MOLTEN SALTS; POTASSIUM
 CHLORIDES; POTENTIOMETRY; REDOX POTENTIAL; URANIUM: T1; URANIUM
 CHLORIDES

ABSTRACT: The investigation of the reversibilities of the
 electrode reactions for the U(III)-U(IV) system in LiCl-KCl
 eutectic melt mainly based on the potential-time
 relationships by chronopotentiometry has been presented. A
 theoretical chronopotentiogram for a reversible redox system
 has been derived and U(III) and U(IV) have been confirmed to
 form a reversible redox system. Standard potentials for the
 couple of U(IV)/U(III) in LiCl-KCl eutectic melt have been
 determined against the Ag/AgCl reference electrode over the
 temperature range from 390 to 530°C, and to be 0.595 V
 at 450°C.

EDE SUBJECT CATEGORY: EDB-400400
 INIS CATEGORY: B12

<294>

Karasev, N.M.; Sidorov, L.N.; Korenev, Yu.M. (Moskovskij
 Gosudarstvennyj Univ. (USSR))

Molecular vapor composition and mass spectra in KF-ZrF₄,
 CsF-ZrF₄ systems AIX-08:298279; EDB-77:069077; Zh. Fiz.
 Khim. (ZFKHA) 50: (6):1630 (Jun 1976)

ABSTRACT: None

EDE SUBJECT CATEGORY: EDB-400201
 INIS CATEGORY: B12

<295>

Karasev, N.M.; Sidorov, L.N.; Korenev, Yu.M. (Moskovskij Gosudarstvennyj Univ. (USSR))

Partial pressures, activities and evaporation heat in the
 KF-ZrF₄ system AIX-08:298280; EDB-77:069078; Zh. Fiz.
 Khim. (ZFKHA) 50:(6):1629 (Jun 1976)

SUBJECT DESCRIPTORS: CHEMICAL REACTION KINETICS:
 Q1,Q2; DISSOCIATION; MASS SPECTRA; POTASSIUM FLUORIDES: T1; VAPOR
 PRESSURE: Q1,Q2; VAPORIZATION HEAT: Q1,Q2; ZIRCONIUM FLUORIDES:
 T2

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-400201
 INIS CATEGORY: B12

<296>

Karasev, N.M.; Sidorov, L.N.; Korenev, Y.M. (Moskovskij Gosudarstvennyj Univ. (USSR))

Interval of congruent and incongruent sublimation of 3KFxZrF₄
 compound in the KF-ZrF₄ system
 AIX-08:339906; EDB-78:017326; Zh. Fiz. Khim. (ZFKHA)
 51:(1):274 (Jan 1977)

SUBJECT DESCRIPTORS: CHEMICAL COMPOSITION; CRYSTALS; PHASE
 DIAGRAMS; POTASSIUM FLUORIDES: T2; SUBLIMATION:
 Q1,Q2; TEMPERATURE DEPENDENCE; ZIRCONIUM FLUORIDES: T1

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-400201
 INIS CATEGORY: B12

<297>

Karow, H.U., Kernforschungszentrum Karlsruhe (Germany, F.R.).
 Inst. fuer Neutronenphysik und Reaktortechnik

In Thermodynamic state, specific heat, and enthalpy function of
 saturated UO₂ vapor between 3000 K and 5000 K **
 ERA-03:012442; EDB-78:021168; (Feb 1977) EURFNR--1460

SUBJECT DESCRIPTORS: ENTHALPY; EVAPORATION; LMFBR TYPE REACTORS:
 T1; MELTDOWN; REACTOR ACCIDENTS: T,Q1; SPECIFIC
 HEAT; THERMODYNAMIC PROPERTIES: Q2; ULTRAHIGH
 TEMPERATURE; URANIUM DIOXIDE: T2; VAPORS; VERY HIGH TEMPERATURE

ABSTRACT: Reactor safety analysis requires knowledge of the
 thermophysical properties of molten oxide fuel and of the

<297>

<297> CONT.

thermal equation-of-state of oxide fuel in thermodynamic liquid-vapor equilibrium far above 3000\$sup 0\$K. The thermodynamic state of saturated UO\$sub 2\$ fuel vapor, its internal energy U(T), specific heats C/sub v/(T) and C/sub p/(T), and enthalpy functions H\$sup 0\$ (H) and H\$sup 0\$ (T)-H\$sub 0\$\$sup 0\$ have been determined by means of statistical mechanics in the temperature range 3000 to 5000\$sup 0\$K. The discussion of the thermodynamic state includes the evaluation of the plasma state and its contribution to the caloric variables-of-state of saturated oxide fuel vapor. The resulting enthalpy functions and specific heats for saturated UO\$sub 2\$ vapor of equilibrium composition and that for pure UO\$sub 2\$ gas are compared with the enthalpy and specific heat data of gaseous UO\$sub 2\$ at lower temperatures known from literature. Tentatively also experimental enthalpy data of the condensed phases of UO\$sub 2\$ have been converted into the vapor phase using the Clausius-Clapeyron equation. However, the correlation based on the Clausius-Clapeyron equation turns out to fail because this equation is not applicable to the chemically reactive oxide fuel system.

EDE SUBJECT CATEGORY: EDB-220900;210500

<298>

Kartsev, V.E.; Kovalev, P.V.; Korshunov, B.G. (Gosudarstvennyj Inst. Redkikh Metallov, Moscow (USSR); Moskovskij Inst. Stali i Splavov (USSR))

Fusibility in K\$sub 2\$Nb(Ta)F\$sub 7\$-LiF-NaF systems
AIX-07:277791;EDB-77:031398; Izv. Vyssh. Uchebn. Zaved.,
Tsvet. Metall. (IVUTA) (3):150-152 (1975)

ABSTRACT: None

EDE SUBJECT CATEGORY: EDB-400201

INIS CATEGORY: D12

<299>

Kartsev, V.E.; Kovalev, P.V.; Korshunov, B.G.

Meltability in system of K\$sub 2\$TaF\$sub 7\$--NaF--NaCl--KCl
AIX-08:298281;EDB-77:069023; Zh. Neorg. Khim. (ZNOKA)
20: (8):2204-2208 (Aug 1975)

SUBJECT DESCRIPTORS: MIXTURES;MOLTEN SALTS;PHASE DIAGRAMS:
Q1,Q2,Q3,Q4,Q5;POTASSIUM CHLORIDES: T1;POTASSIUM FLUORIDES:
T2;QUANTITY RATIO;SODIUM FLUORIDES: T3;SODIUM FLUORIDES:
T5;TANTALUM FLUORIDES: T4;THERMAL ANALYSIS

<299> CONT.

ABSTRACT: Thermographic and visual-polythermal techniques were used to study the meltability in $K_{2}TaF_{7}$ -NaF-NaCl-KCl system. The tetrahedron-forming sections NaF-NaCl- $K_{2}TaF_{7}$ -KCl and NaF- $K_{2}TaF_{7}$ - $K_{2}TaF_{7}$ -KCl divide the concentration tetrahedron into three particular tetrahedra: NaF- $K_{2}TaF_{7}$ - $K_{2}TaF_{7}$ -KCl, NaF-NaCl- $K_{2}TaF_{7}$ -KCl, and NaF-NaCl-KCl- $K_{2}TaF_{7}$ -KCl. Non-variant equilibrium points in all of the particular four-component systems have been determined.

EDB SUBJECT CATEGORY: EDB-400201
INIS CATEGORY: B12

<300>

Kartsev, V.E.; Kovalev, F.V.; Korshunov, B.G.

Meltability in $K_{2}NbF_{7}$ -NaF-NaCl-KCl system
AIX-07:277792; EDB-77:031466; Zh. Neorg. Khim. (ZNOKA)
21: (2):536-540 (Feb 1976)

SUBJECT DESCRIPTORS: EUTECTICS; MELTING; NIOBIUM FLUORIDES:
T1; PHASE DIAGRAMS: Q1, Q2, Q3, Q4, Q5; POTASSIUM CHLORIDES:
T2; POTASSIUM FLUORIDES: T3; SODIUM CHLORIDES: T4; SODIUM
FLUORIDES: T5; THERMAL ANALYSIS

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-400201
INIS CATEGORY: B12

<301>

Kazanjan, A.R., Atomics International Div., Golden, Colo. (USA)

In Radiolytic gas generation in plutonium contaminated waste materials ** INS-77:004217; ERA-02:019404; EDB-77:035009; (29 Oct 1976) RFP-2469

SUBJECT DESCRIPTORS: ALPHA SOURCES; AUTORADIOLYSIS: Q1; CARBON
TETRACHLORIDE: T6; CHEMICAL REACTION YIELD: Q2; GLOVES:
T10; HYDROGEN: T2; MASS SPECTROSCOPY; OILS: T5; ORGANIC ION
EXCHANGERS: T7; ORGANIC SOLVENTS: T9; PLEXIGLAS: T3; PLUTONIUM
238; PLUTONIUM 239; PLUTONIUM OXIDES; POLYETHYLENES: T8; PVC:
T4; RADIOACTIVE WASTES: T1; RADIOLYSIS:
Q3, Q4, Q5, Q6, Q7, Q8, Q9, Q10; SOLID WASTES

ABSTRACT: Many plutonium contaminated waste materials decompose into gaseous products because of exposure to alpha radiation. The gases generated (usually hydrogen) over long-storage

<301>

<301> CONT.

periods may create hazardous conditions. To determine the extent of such hazards, knowing the gas generation yields is necessary. These yields were measured by contacting some common Rocky Flats Plant waste materials with plutonium and monitoring the enclosed atmospheres for extensive periods of time. The materials were Plexiglas, polyvinyl chloride, glove-box gloves, machining oil, carbon tetrachloride, chloroethene VG solvent, Kimwipes (dry and wet), polyethylene, Dowex-1 resin, and surgeon's gloves. Both ^{239}Pu oxide and ^{238}Pu oxide were used as radiation sources. The gas analyses were made by mass spectrometry and the results obtained were the total gas generation, the hydrogen generation, the oxygen consumption rate, and the gas composition over the entire storage period. Hydrogen was the major gas produced in most of the materials. The total gas yields varied from 0.71 to 16 cm^3 (standard temperature pressure) per day per curie of plutonium. The oxygen consumption rates varied from 0.0088 to 0.070 millimoles per day per gram of plutonium oxide-239 and from 0.0014 to 0.0051 millimoles per day per milligram ^{238}Pu .

EDB SUBJECT CATEGORY: EDB-052001;400600
 INIS CATEGORY: E51;B14

<302>

Kazanskii, K.A.; Novikov, V.M.

Thermophysical and electrophysical properties of uranium hexafluoride at temperatures of $(1-11) \times 10^3$ $^\circ\text{K}$ and pressures of 0.1-100 atm ERA-02:023066;EDB-77:044535; High Temp. (USSR) (Engl. Transl.) (HITEA) 14:(3):410-415 (Nov 1976)

SUBJECT DESCRIPTORS: ELECTRIC CONDUCTIVITY: Q1;MEDIUM
 PRESSURE;PARTIAL MOLAL VOLUME;THERMODYNAMIC PROPERTIES:
 Q1;URANIUM HEXAFLUORIDE: T1;VERY HIGH TEMPERATURE

ABSTRACT: The partial composition of uranium hexafluoride at temperatures of $(1-11) \times 10^3$ $^\circ\text{K}$ and pressures of 0.1-100 atm is calculated with due regard to all the uranium fluoride dissociation processes and single ionization of uranium atoms. These data are used to calculate the thermodynamic functions and electrical conductivity of UF_6 . The region in which the electronegativity of fluorine has little effect on the conductivity is determined. Qualitative explanations of the variation of the thermodynamic quantities with temperature are given. (AIP)

EDB SUBJECT CATEGORY: EDB-400702

<303>

Keller, C.; Berndt, U. (Kernforschungszentrum Karlsruhe (F.R. Germany). Inst. fuer Radiochemie) ** Mueller, W.; Lindner, R. (eds.)

Reaction of americium oxides with threevalent metal oxides In Transplutonium 1975 **

AIX-07:270260; ERA-02:020582; EDB-77:036885; (1976)
North-Holland

SUBJECT DESCRIPTORS: ALUMINIUM OXIDES: T4; AMERICIUM OXIDES; BINARY MIXTURES; BCRON OXIDES: T7; LATTICE PARAMETERS; LUTETIUM OXIDES: T2; NEPTUNIUM OXIDES: T9; PHASE STUDIES: Q1, Q2, Q3, Q4, Q5, Q6, Q7, Q8, Q9, Q10; PLUTONIUM DIOXIDE: T10; SAMARIUM OXIDES: T1; SCANDIUM OXIDES: T6; SOLUBILITY; THERMODYNAMIC PROPERTIES; THORIUM OXIDES: T8; VANADIUM OXIDES: T5; YTTRIUM OXIDES: T3

ABSTRACT: In accordance with its electronic configuration and the resulting position in the periodic table, the 5f element americium is the heavy homologue of the 4f element europium. But according to the ionic radii it is anticipated that Am^{III} ($r = 0.99\text{\AA}$) shows solid state reactions more similar to Nd^{III} ($r = 0.995\text{\AA}$) than to Eu^{III} ($r = 0.95\text{\AA}$). Differences in the solid state chemical behaviour of Am^{III} and Nd^{III} (or Eu^{III}) are observed because of the higher oxidation states $\text{Am}(\text{IV})$, $\text{Am}(\text{V})$ and $\text{Am}(\text{VI})$, especially being stabilized in ternary and polynary oxides. Only a few investigations have been published concerning Am^{III} in ternary oxides. This report gives a comprehensive review on the solid state reaction of $\text{AmO}_{1.5}$ ($\text{Am}^{\text{II}}\text{O}$ $\text{Am}^{\text{III}}\text{O}$ O_2).

EDB SUBJECT CATEGORY: EDB-360202
INIS CATEGORY: B13

<304>

Kent, R.A.; Zocher, R.W., Los Alamos Scientific Lab., N.Mex. (USA)

In Reduction of PuO_2 by CO and equilibrium partial pressures above plutonia **

INS-77:010092; ERA-02:039272; EDB-77:087060; (Nov 1976)
LA--6534

SUBJECT DESCRIPTORS: CARBON MONOXIDE: T2; CHEMICAL PREPARATION; CHEMICAL REACTIONS: Q2; PLUTONIUM DIOXIDE: T1; RADIOISOTOPE HEAT SOURCES; REDUCING AGENTS; REDUCTION: Q1; THERMODYNAMICS; VERY HIGH TEMPERATURE

ABSTRACT: The reduction of plutonia in flowing carbon monoxide has been determined over the range of 1200 to 1800°K.

<304>

<304> CONT.

The experimental data confirm recently published thermodynamic functions for the plutonium--oxygen system. Equilibrium partial pressures are listed for vapor species above stoichiometric and reduced plutonia in the range 1200 to 1800\$sup 0\$K.

EDB SUBJECT CATEGORY: EDB-070300;360201
INIS CATEGORY: D21;B23

<305>

Khaldcyanidi, K.A.; Grankina, Z.A. (AN SSSR, Novosibirsk. Inst. Neorganicheskij Khimii)

Some properties of molybdenum pentoxide

AIX-08:320642;EDB-77:114198; Izv. Sib. Otd. Akad. Nauk SSSR, Ser. Khim. Nauk (IZSKA) 2:(1):47-50 (Jan 1976)

SUBJECT DESCRIPTORS: CHEMICAL PREPARATION: Q1;CHEMICAL PROPERTIES: Q1;DIFFERENTIAL THERMAL ANALYSIS: Q1,Q2;INFRARED SPECTRA: Q1,Q2;MAGNETIC SUSCEPTIBILITY;MOLTEN SALTS;MOLYBDENUM COMPLEXES;MOLYBDENUM OXIDES: T1;POTASSIUM FLUORIDES: T2;SOLUBILITY;X-RAY DIFFRACTION: Q1,Q2

ABSTRACT: The molybdenum (V) pentoxide has been synthesized by the dehydration of $\text{MoO}(\text{OH})_3$, some of its physico-chemical properties have been investigated. The system $\text{KF}-\text{Mo}_2\text{O}_5$ containing from 0 to 60 mol.% KF was studied by thermography, X-ray and IR-spectroscopy methods.

EDB SUBJECT CATEGORY: EDB-400201;360204
INIS CATEGORY: B12

<306>

Khanaev, E.I.; Afanas'ev, Y.A.; Storozhenko, T.P. (Kubanskiy Gosudarstvennyy Univ., Krasnodar (USSR))

Formation enthalpy of di- and trifluorides of samarium

AIX-08:317721;EDB-77:133387; Zh. Fiz. Khim. (ZFKHA) 49:(9):2454 (Sep 1975)

SUBJECT DESCRIPTORS: CALORIMETRY;DISSOLUTION;FORMATION HEAT: Q1;HYDROFLUORIC ACID;MEDIUM TEMPERATURE;SAMARIUM;SAMARIUM FLUORIDES: T1

<307> CONT.

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-400201

INIS CATEGORY: B12

<307>

Khomyakov, V.I.; Shishkev, Yu.D. ** AN SSSR, Moscow;
Gosudarstvennyj Komitet po Ispol'zovaniyu Atomnoj Ehnergii
SSSR, Moscow

Use of fluoride melts for separation of some fission products and americium from mixtures of different fluorides being the simulators of residues from fast reactor fuel fluorination In All-union conference on chemistry of transplutonium elements (americium, curium, berkelium, californium) **
AIX-08:295460;EDB-77:072630; (1976) INIS-mf--3283

SUBJECT DESCRIPTORS: AMERICIUM: T2;AMERICIUM FLUORIDES;CALCIUM FLUORIDES;CESIUM FLUORIDES;DISTRIBUTION FUNCTIONS;EUROPIUM FLUORIDES;EUTECTICS;FISSION PRODUCTS: T1;LANTHANUM FLUORIDES;MOLTEN SALTS;OXIDES;SEPARATION PROCESSES: Q1,Q2;SILICON FLUORIDES;SODIUM FLUORIDES;STRONTIUM FLUORIDES;URANIUM FLUORIDES

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-050800

INIS CATEGORY: B13

<308>

Khomyakov, V.I.; Shishkev, Yu.D.; Galkin, N.P. ** AN SSSR, Moscow; Gosudarstvennyj Komitet po Ispol'zovaniyu Atomnoj Ehnergii SSSR, Moscow

Investigation into americium extraction process from residues of irradiated nuclear fuel after its fluorination by SiF-NaF and SiF-CaF melts of eutectic composition In All-union conference on chemistry of transplutonium elements (americium, curium, berkelium, californium) **
AIX-08:295583;ERA-02:032905;EDB-77:089179; (1976)
INIS-mf--3283

SUBJECT DESCRIPTORS: AMERICIUM: T1;AMERICIUM FLUORIDES;CALCIUM FLUORIDES;DISTRIBUTION FUNCTIONS;EUTECTICS;MOLTEN SALTS;NUCLEAR FUELS;PLUTONIUM FLUORIDES;RARE EARTH COMPOUNDS;SEPARATION PROCESSES: Q1;SILICON FLUORIDES;SODIUM FLUORIDES

<308>

<308> CONT.

ABSTRACT: None

EDE SUBJECT CATEGORY: EDB-400105

INIS CATEGORY: B16

<309>

King, R.T.; DeVan, J.H.; Goodwin, G.M.; Isaacs, H.; Heestand,
R.L.; McCurdy, H.C., Oak Ridge National Lab., Tenn. (USA)

In Report of task force on alternate structural materials for
Liquid Metal Fast Breeder Reactors **

ERA-01:019155;EDB-76:052411; (May 1976) ORNL--5076

SUBJECT DESCRIPTORS: COMPARATIVE EVALUATIONS: Q2;CORROSION;LMFBR
TYPE REACTORS: T1;MECHANICAL PROPERTIES: Q3,Q4;REACTOR
MATERIALS: T2,Q1;STAINLESS STEEL-304: T3;STAINLESS STEEL-316:
T4

ABSTRACT: The findings and recommendations of the Task Force on Alternate Structural Materials for Liquid Metal Fast Breeder Reactors are summarized. The objectives and results of this activity were: (1) Identify existing and potential problems with types 304 and 316 stainless steel as LMFBR structural alloys for vessels, piping, and components. The advantages and disadvantages of types 304 and 316 stainless steels for these applications are summarized. Sufficient disadvantages were found to warrant exploring alternate materials. (2) Assess commercial alloys and their properties that may solve problems identified in the Phase I section. For the expected designs and operating temperatures of future LMFBRs, no commercial alloys were found that solve all problems without introducing others. However, the advantages of certain stabilized ferritic alloys containing about 9--14 percent Cr and 1 percent Mo, austenitic 15 Cr-15 Ni alloys, subclasses of types 304 and 316 stainless steels that offer possible improved stress corrosion resistance, and nickel-bearing alloys warrant further evaluation and development for their service as LMFBR structural alloys. (3) Identify specific objectives of a program plan to develop alternate structural alloy(s) to the point where an LMFBR system based thereon may be designed by 1985. Objectives of development programs involving the testing and evaluation of progressively larger heats of material are identified. Fabrication and testing of trial components in sodium test loops are included. Relevant industrial experience and literature information that form the basis of the report are identified.

EDE SUBJECT CATEGORY: EDB-210500;360103

<310>

Kirbiyik, M.; Garner, P.L.; Refling, J.G.; Reynolds, A.B.
(Virginia Univ., Charlottesville (USA). Dept. of Nuclear
Engineering)

Hydrodynamics of post-disassembly fuel expansions

AIX-07:245060; ERA-02:007339; EDB-77:002390; Nucl. Eng. Des.
(NEDEA) 35:(3):441-460 (Dec 1975)

SUBJECT DESCRIPTORS: BOUNDARY

CONDITIONS; ENTHALPY; ENTROPY; EQUATIONS OF STATE; F CODES; FINITE
DIFFERENCE METHOD; HYDRODYNAMICS: Q2; LAGRANGIAN FUNCTION; LMFBR
TYPE REACTORS: M1; MATHEMATICAL MODELS; NUCLEAR FUELS; REACTOR
CORE DISRUPTION: M2, Q1; REACTOR CORES; SODIUM; THERMAL EXPANSION

ABSTRACT: The early expansion of the fuel following disassembly
in an LMFBR core disruptive accident is modeled. Spherical
expansion in the sodium is assumed. A Lagrangian,
finite-difference hydrodynamic code (FEXPAN) describes the
motion. Disassembly employs VENUS-II, and a consistent
equation of state for fuel was used throughout disassembly
and FEXPAN. Time-dependent mechanical work and fuel vaporized
without fuel mixing are obtained. FEXPAN is compared with
time-independent expansion for the effect of fuel mixing. For
example, for a particular accident analysed 75 MW sec
mechanical work was calculated for expansion with no mixing
versus 10 MW sec with complete core mixing.

EDB SUBJECT CATEGORY: EDB-220900; 210500

INIS CATEGORY: E35

<311>

Kiriolovich, A.P.; Dem'yanovich, M.A.; Lavrinovich, Yu.G.;
Skiba, O.V.; Gazizov, R.K.; Vorobey, M.P.,
Nauchno-Issledovatel'skij Inst. Atomnykh Reaktorov, Melekes
(USSR)

In Study of the behavior of high activity waste produced during
the regeneration of fast reactor fuel elements by the gaseous
fluoride method ** INS-77:000068; ERA-02:009302; EDB-77:005682;
(1975) ERDA-tr--204

SUBJECT DESCRIPTORS: BOR-60 REACTOR; CARBON STEELS: M2; CHEMICAL
PROPERTIES: Q5; COMPATIBILITY; COMPATIBILITY: Q5; CORROSION
RESISTANCE: Q2, Q3, Q4; DEGASSING; FLUORIDE VOLATILITY
PROCESS; NICKEL: M4; RADIOACTIVE WASTE DISPOSAL:
M, Q1; RADIOACTIVE WASTES: M5; REPROCESSING: M1; SPENT FUEL
ELEMENTS; STAINLESS STEELS: M3

ABSTRACT: The composition and certain physicochemical and
radiation properties of waste obtained from reprocessing
BOR-60 fuel by the gaseous fluoride method were studied on an

<311>

<311> CONT.

experimental stand. It was found that from 80 to 85 percent of the total activity is concentrated in the fluorination cakes. From thermographic analysis, measurement of radiation-induced gas evolution and composition of the gaseous phase, it is concluded that the waste must be canned. Compatibility of carbon steel, austenitic steel, and nickel with low-activity fluoride cakes was studied. It is shown that the corrosion of these materials increases with temperature and moisture content of the medium. In the first stage of waste disposal, hermetic storage in special containers of stainless steel or nickel is recommended. From their corrosion resistance in low-activity fluoride cakes, the wall temperatures of containers made of St. 3 should not exceed 100°C , Kh18N10T 230°C , and nickel 300°C . Later, after cooling and a decrease in the γ activity of the waste (10 to 15 years), they can be reprocessed to recover valuable components (Am, Cm, etc.) and then buried. 5 tables. (DLC)

EDB SUBJECT CATEGORY: EDB-052002;050800

INIS CATEGORY: E52;B16

<312>

Kiselev, Y.M.; Sevast'yanov, V.G.; Spitsyn, V.I. (AN SSSR, Moscow. Inst. Obshchej i Neorganicheskoy Khimii; Moskovskij Gosudarstvennyj Univ. (USSR))

Stability of rare earth tetrafluorides

AIX-08:320647;EDB-77:114207; Izv. Akad. Nauk SSSR, Ser. Khim. (IASKA) (5):959-962 (May 1976)

SUBJECT DESCRIPTORS: ACTIVATION ENERGY;CERIUM FLUORIDES:

T1;PRASEODYMIUM FLUORIDES: T2;PYROLYSIS:

Q1,Q2,Q3;STABILITY;TERBIUM FLUORIDES: T3;THERMAL GRAVIMETRIC ANALYSIS;THERMODYNAMICS;VAPOR PRESSURE;X-RAY DIFFRACTION

ABSTRACT: A study was made of thermal stability of tetrafluorides of praseodymium, cerium and terbium. The study was carried out thermogravimetrically in the nitrogen atmosphere with the heating rate of 3-6 grad./min, and manometrically. The products of thermolysis were investigated by the X-ray diffraction method.

EDB SUBJECT CATEGORY: EDB-400201

INIS CATEGORY: B12

<313>

Kisun'ko, V.Z.; Novokhatskij, I.A.; Arkharov, V.I.; Belov, B.P.

Effect of various additions on the temperature of structural transformation in liquid iron AIX-07:232998; EDB-76:059951; Izv. Akad. Nauk SSSR, Met. (IZNMA) (2):176-179 (Mar 1975)

SUBJECT DESCRIPTORS: ACTIVATION ENERGY; CARBON ADDITIONS; CHROMIUM ADDITIONS; IRON: T1; LIQUID METALS; MANGANESE ADDITIONS; NICKEL ADDITIONS; PHASE TRANSFORMATIONS: Q1; SILICON ADDITIONS; STABILITY; TEMPERATURE DEPENDENCE; TRANSITION TEMPERATURE; VISCOSITY

ABSTRACT: The crucible torsional vibration method was used for demonstrating that both the absolute values of the kinematic viscosity (η) and the activation energy of viscous flow (E_{η}) of molten iron (melting temp. 1750°C) change abruptly at 1635 $\pm$ 10°C. This phenomenon is explained in terms of a polymorphic transformation of ordered microregions (clusters) of the melt. The characteristic features of cluster polymorphism in liquid cobalt and nickel are not observed in the melting temperature range up to 1700°C. A study is made of the effect of C, Si, Mn, Cr, Ni, and W, in a concentration of 1.0 at.%, on the polymorphic transformation temperature (t_{pol}) of clusters of liquid iron at which the abrupt change of η and E_{η} was recorded.

EED SUBJECT CATEGORY: EDB-360102

INIS CATEGORY: B22

<314>

Kitamura, T.; Kojima, T. (Government Industrial Research Inst., Takamatsu (Japan)); Ichise, Mitsunojo

Comparison of the tendencies of complex formation of various metal ions in fused LiF-NaF-KF (eut.) and in fused LiCl-KCl (eut.) AIX-07:243905; EDB-76:075656; Denki Kagaku (DNKKA) 43: (3):119-126 (Mar 1975)

SUBJECT DESCRIPTORS: ANIONS; CADMIUM COMPLEXES:

T1; CATIONS; CHEMICAL PREPARATION:

Q1, Q2, Q3, Q4, Q5, Q6, Q7, Q8, Q9; CHEMICAL REACTIONS; COBALT

COMPLEXES: T7; COPPER COMPOUNDS: T2; CYANIDES; ELECTROMOTIVE

FORCE; EQUILIBRIUM; EUTECTICS; FREE ENERGY; IODIDES; IRON

COMPLEXES: T8; LEAD COMPLEXES: T4; LITHIUM CHLORIDES; LITHIUM

FLUORIDES; MANGANESE COMPLEXES: T9; MOLTEN SALTS; NICKEL

COMPLEXES: T6; POTASSIUM CHLORIDES; POTASSIUM FLUORIDES; SODIUM

FLUORIDES; STABILITY; THALLIUM COMPLEXES: T3; VOLTAMETRY; ZINC

COMPLEXES: T5

ABSTRACT: Complex formation of Cu^{+} , Tl^{+} , Cd^{++} ,

<314>

<314> CONT.

Pb⁺⁺, Zn⁺⁺, Ni⁺⁺, Co⁺⁺, Fe⁺⁺ and Mn⁺⁺ with I⁻ and CN⁻ in LiF-NaF-KF (eut.) was studied by emf method and potential sweep voltammetry. Stability constants and free energies of complex formation were determined for Cu⁺-I⁻, Cu⁺-CN⁻, Tl⁺-I⁻, Cd⁺⁺-CN⁻, Ni⁺⁺-CN⁻, Co⁺⁺-CN⁻, and Fe⁺⁺-CN⁻ systems. The tendencies of complex formation in LiF-NaF-KF were compared with those in LiCl-KCl. The free energies of complex formation of Cu⁺ and Tl⁺ were greater in LiF-NaF-KF than in LiCl-KCl, but those for divalent metal ions were greater in LiCl-KCl than in LiF-NaF-KF. Cd⁺⁺ and Pb⁺⁺ formed iodide complexes in LiCl-KCl, but did not in LiF-NaF-KF. Zn⁺⁺, Ni⁺⁺, Co⁺⁺, Fe⁺⁺, and Mn⁺⁺ did not form any iodide complex in both solvents. Prediction of these tendencies was tried by calculating free energy changes of complex formation from the standard electromotive forces for single metal halides given by Hamer. Though the predicted values were substantially smaller than the experimental values, they showed qualitatively the tendencies of the complex formation.

EDP SUBJECT CATEGORY: EDB-400201

INIS CATEGORY: B12

<315>

Kleykamp, H., Kernforschungszentrum Karlsruhe (F.R. Germany)

In Microprobe study of fission product behavior in high-burnup HTR fuels ** ERA-01:019121; INS-76:016066; EDB-76:058152; (1 Sep 1975) ORNL-tr--4108

SUBJECT DESCRIPTORS: BURNUP; COATED FUEL PARTICLES:

Q1, T2; COATINGS; CORROSION; DIFFUSION; FISSION PRODUCT RELEASE:

Q2; FISSION PRODUCTS; HTGR TYPE REACTORS:

T1; PERFORMANCE; PYROLYTIC CARBON; SILICON

CARBIDES; SOLUBILITY; URANIUM CARBIDES; URANIUM DIOXIDE

ABSTRACT: Electron microprobe analysis of irradiated coated particles with high burnup (greater than 50 percent fima) gives detailed information on the chemical state and the transport behavior of the fission products in UO₂ and UC₂ kernels and in the coatings. In oxide fuel kernels, metallic inclusions and ceramic precipitations are observed. The solubility behavior of the fission products in the fuel matrix has been investigated. Fission product inclusions could not be detected in carbide fuel kernels; post irradiation annealed UC₂ kernels, however, give information on the element combinations of some fission product phases. Corresponding to the chemical state in the

<315> CONT.

kernel, Cs, Sr, Ba, Pd, Te and the rare earths are released easily and diffuse through the entire pyrocarbon coating. These fission products can be retained by a silicon carbide layer. The initial stage of a corrosive attack of the SiC coating by the fission products is evidenced.

EDB SUBJECT CATEGORY: EDB-210300
INIS CATEGORY: E33

<316>

Kleykamp, H. (Kernforschungszentrum Karlsruhe (Germany, F.R.).
Inst. fuer Material- und Festkuerperforschung)

Radial oxygen distribution in the fuel of irradiated mixed oxide fuel elements with different final stoichiometries
AIX-08:325220; ERA-03:003611; EDB-77:145592; J. Nucl. Mater. (JNUMA) 66:(3):292-300 (May 1977)

SUBJECT DESCRIPTORS: BURNUP; CHEMICAL COMPOSITION; ENTHALPY; FAST NEUTRONS; FUEL PINS; IRRADIATION; MOLYBDENUM; MOLYBDENUM OXIDES: T1; PLUTONIUM OXIDES; QUANTITY RATIO; SPATIAL DISTRIBUTION; STOICHIOMETRY: Q1, Q2; URANIUM OXIDES: T2

ABSTRACT: The radial gradients of the oxygen potential and the oxygen/metal ratio O/M in the fuel of irradiated mixed oxide pins at the end of life have been determined from the Mo and MoO₂ concentrations in the metallic and ceramic inclusions. The measurements were carried out on uranium-plutonium oxides with different initial stoichiometries (O/M) of 1.96, 1.98 and 1.99, resp., after fast neutron irradiation under identical conditions. Using the measured overall final stoichiometries and the stoichiometry shifts during burnup, the initial stoichiometries of the three investigated oxides have been recalculated, giving (O-M) values of 1.961, 1.978, and > 1.985, resp.

EDB SUBJECT CATEGORY: EDB-220300
INIS CATEGORY: E23

<317>

Klinkova, L.A.; Ukshe, E.A.

Dissolution of corundum in molten vanadates EDB-76:036560; Zh. Neorg. Khim. (ZNOKA) 20:(2):481-484 (Feb 1975)

SUBJECT DESCRIPTORS: ALKALI METAL COMPOUNDS: T1; CORUNDUM: T2; CRYSTALS; DIFFERENTIAL THERMAL ANALYSIS; DISSOLUTION; EUTECTICS; MOLTEN SALTS; PHASE DIAGRAMS: Q1, Q2, Q3; VANADATES: T3; X-RAY DIFFRACTION

<317>

<317> CONT.

ABSTRACT: None

EIE SUBJECT CATEGORY: EDB-360202

INIS CATEGORY: B12

<318>

Kmetko, E.A.; Hill, H.H. (Los Alamos Scientific Lab., N.Mex. (USA))

Anomalous melting of f electron metals (with attention to Pu)
 AIX-07:270409;EDB-77:030681; J. Phys., F (London) (JPPMA)
 6: (6):1025-1037 (Jun 1976)

SUBJECT DESCRIPTORS: ACTINIDES: T1;BCC LATTICES;ELECTRONS:F
 STATES;LIQUID METALS;MELTING POINTS: Q1,Q2;PLUTONIUM:
 T2;PLUTONIUM-EPSILON;SELF-DIFFUSION

ABSTRACT: An explanation is offered for the anomalously low melting points of f electron metals, especially the early actinides. It is known that f electron participation in bonding is responsible for the crystal structures of unusually low symmetry found in the early actinides at room temperature. It is suggested that f electrons, because of the angular dependence of their wavefunctions, are able to bond only to a minimal extent in the high temperature BCC phase out of which most of the metals of concern melt. Indeed the f electrons bond better in the liquid than in the BCC phase. Attention is directed to Pu, and several of the anomalous properties of its BCC epsilon phase are discussed in detail. A role for f electron participation in self-diffusion is proposed.

EDB SUBJECT CATEGORY: EDB-360102

INIS CATEGORY: B22

<319>

Kniseley, R.N.; Butler, C.C.; Fassel, V.A. ** Dean, J.A. (ed.)

Rare earth elements In Flame emission and atomic absorption spectrometry. III. Elements and matrices **
 ERA-01:016787;EDB-76:042531; (1975) Marcel Dekker, Inc.

SUBJECT DESCRIPTORS: ABSORPTION SPECTROSCOPY: Q1;CHEMICAL
 ANALYSIS;EMISSION SPECTROSCOPY: Q1;RARE EARTHS: T1;SENSITIVITY

ABSTRACT: Methods are surveyed for the determination of the rare earth elements by flame and atomic absorption spectroscopy. Flame conditions necessary for the flame excitation of the line spectra of the rare earths are specified. Data for the

<319> CONT.

emission and atomic absorption detection limits for the rare earths are tabulated. Analytical applications are discussed for both methods. 82 references. (BLM)

EDB SUBJECT CATEGORY: EDB-400104

<320>

Kober, V.I.; Nichkov, I.E. (Ural'skiy Politekhnikheskiy Inst., Sverdlovsk (USSR))

Interaction of atoms of rare earths-metal alloys

AIX-08:298575;EDB-77:068255; Zh. Fiz. Khim. (ZFKHA)
50:(1):281 (Jan 1976)

SUBJECT DESCRIPTORS: BINARY ALLOY SYSTEMS;CHEMICAL REACTION KINETICS;ELECTRONIC STRUCTURE;MELTING POINTS;OREIT STABILITY;PHASE DIAGRAMS;RARE EARTH ALLOYS: T

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-360100

INIS CATEGORY: B22

<321>

Kobzenko, G.F.; Chernogorenko, V.B.; L'vov, S.N.; Lesnaya, M.I.; Lynchak, K.A.

Phase diagram and properties of Nd--As alloys

AIX-08:298286;EDB-77:068799; Zh. Neorg. Khim. (ZNOKA)
20:(8):2169-2173 (Aug 1975)

SUBJECT DESCRIPTORS: ARSENIC: T2;ARSENIC COMPOUNDS;ARSENIDES: T1;BINARY ALLOY SYSTEMS;CHEMICAL COMPOSITION;ELECTRIC CONDUCTIVITY;MELTING POINTS: Q1;N-TYPE CONDUCTORS;NEODYMIUM;NEODYMIUM ALLOYS;PHASE DIAGRAMS: Q1;QUANTITY RATIO;SOLUBILITY: Q2;SOLVENT PROPERTIES

ABSTRACT: The phase diagram of Nd--As system has been constructed, and the electrophysical properties of the alloys belonging to this system have been studied. It has been established that the system includes three arsenides: NdAs_3 , NdAs and NdAs_2 . When heated, the latter dissociates into NdAs and gaseous arsenic. The melting point of NdAs is 2.250 ± 15 deg C. The solubility of arsenic in

<321>

<322> CONT.

neodymium is less than 1%. In all alloys containing up to 50% of arsenic, n-conduction is prevalent.

EDB SUBJECT CATEGORY: EDB-360603;360104

INIS CATEGORY: B12

<322>

Koch, G. (Kernforschungszentrum Karlsruhe (Germany, F.R.). Inst. fuer Heisse Chemie)

Chemical reprocessing of spent nuclear fuels

AIX-08:334490;ERA-02:058490;EDB-77:137286; Chem. Ztg. (CMKZA) 101: (2):64-81 (Feb 1977)

ABSTRACT: The reprocessing of nuclear fuels from atomic power stations has a twofold goal. On the one hand it is serving for fuel supply by recovering the fissile materials which have not been consumed or which have been freshly generated in the reactor. On the other hand the radioactive waste products from nuclear power generation are pretreated for long-term safe disposal. The core element of the chemical processing is the PUREX Process, a counter-current solvent extraction procedure using tributyl phosphate (TRP) as the solvent for uranium and plutonium. The chemical basis and the technological performance of the process are discussed.

EDB SUBJECT CATEGORY: EDB-050800

INIS CATEGORY: B16

<323>

Koelling, D.D., Argonne National Lab., Ill. (USA)

In Band structure studies of actinide systems **

INS-77:004829;ERA-02:022817;EDB-77:043619; (1976)
CONF-760956--2

SUBJECT DESCRIPTORS: ACTINIDE ALLOYS: T3;ACTINIDE COMPOUNDS:

T2;ACTINIDES: T1;BAND THEORY;ELECTRONIC STRUCTURE:

Q1,Q2,Q3,Q4,Q5,Q6;F STATES;URANIUM ALLOYS: T5;URANIUM

CARBIDES: T4;URANIUM HEXAFLUORIDE: T6

ABSTRACT: The nature of the f-orbitals in an actinide system plays a crucial role in determining the electronic properties. It has long been realized that when the actinide separation is small enough for the f-orbitals to interact directly, the system will exhibit itinerant electron properties: an absence of local moment due to the f-orbitals and sometimes even superconductivity. However, a number of systems with the larger actinide separation that should imply local moment behavior also exhibit itinerant properties.

<323> CONT.

Such systems (URh_{3f} , UIr_{3f} , UGe_{3f} , UC) were examined to learn something about the other f-interactions. A preliminary observation made is that there is apparently a very large and anisotropic mass enhancement in these systems. There is very good reason to believe that this is not solely due to large electron--electron correlations but to a large electron--phonon interaction as well. These features of the 'non-magnetic', large actinide separation systems are discussed in light of our results to date. Finally, the results of some recent molecular calculations on actinide hexafluorides are used to illustrate the shielding effects on the intra-atomic Coulomb term U_{f-f} which would appear in any attempt to study the formation of local moments. As one becomes interested in materials for which a band structure is no longer an adequate model, this screened U_{f-f} is the significant parameter and efforts must be made to evaluate it in solid state systems.

EDB SUBJECT CATEGORY: EDB-360102;400702;360202
INIS CATEGORY: E22;B13;B23

<324>

Kolster, B.H., Nijverheidsorganisatie TNO, The Hague (Netherlands). Projectgroep Kernenergie TNO

In Study of mass transfer by liquid sodium in closed loop systems. Sodium technology development programme **
EDE-76:036501; (Dec 1974) INIS-mf--2001

SUBJECT DESCRIPTORS: CARBON;CHROMIUM ADDITIONS;CORROSION: Q1;CORROSION PRODUCTS;CORROSIVE EFFECTS: Q2;DEPOSITION;HIGH TEMPERATURE;LIQUID METALS;MASS TRANSFER: Q2;NIOBIUM ADDITIONS;PRIMARY COOLANT CIRCUITS;SECONDARY COOLANT CIRCUITS;SODIUM: T2;SODIUM COOLED REACTORS;STAINLESS STEEL-316: T1

ABSTRACT: Results of a research program on mass transfer phenomena by liquid sodium are reported. Concerning corrosion, results are presented of investigations of the influence of temperature, oxygen level, and downstream position on the corrosion behavior of stainless steel 316 specimens. A brief summary of a model and a derivation of the corrosion rate expression is given. For deposition the same parameters as for corrosion are studied. Specimens were extensively investigated by means of a combination of metallographic techniques. In order to eliminate the influence of the loop material on the mass transport, a loop will be built using molybdenum for construction material. Decarburization studies are carried out and it is concluded that heat treatment and an addition of certain amounts of

<324>

<324> CONT.

niobium and chromium have a stabilizing effect on the decarburization of specimen materials.

EDF SUBJECT CATEGORY: EDB-360105

INIS CATEGORY: B15

<325>

Komarevtseva, N.I.; Kuz'micheva, E.U.; Kovba, L.M.

New oxide of uranium EDB-76:066381; Sov. Radiochem. (Engl. Transl.) (SVRDA) 17:(3):432-433 (1975)

SUBJECT DESCRIPTORS: BOND ANGLE; BOND LENGTHS; CHEMICAL PREPARATION; PHASE STUDIES: Q1; STRUCTURAL CHEMICAL ANALYSIS: Q1; URANIUM OXIDES: M1; VERY HIGH PRESSURE; VERY HIGH TEMPERATURE; X-RAY DIFFRACTION

ABSTRACT: The formation of the phase $U_{16}O_{39}$ in the interaction of the uranium oxides UO_2 and U_3O_8 by heating for 5 min at $1000^\circ C$ and a pressure of 87 kbar was detected. The parameters of the unit cell were determined by an x-ray diffraction method.

EDF SUBJECT CATEGORY: EDB-400702; 400104; 360202

<326>

Komm, T.Z.; Kaul, A.R.; Tret'yakov, Yu.D.

Investigation of formation thermodynamic of rare earth aluminates with perovskite structure
AIX-08:334306; EDB-77:140993; Vestn. Mosk. Univ., Ser. II. Khim. (VMUKA) 17:(5):636 (Sep 1976)

SUBJECT DESCRIPTORS: ALUMINATES: T1; CRYSTAL STRUCTURE; FORMATION FREE ENTHALPY: Q1, Q2; LANTHANUM COMPOUNDS; NEODYMIUM COMPOUNDS; PRASEODYMIUM COMPOUNDS; RARE EARTH COMPOUNDS: T2; SAMARIUM COMPOUNDS

ABSTRACT: None

EDF SUBJECT CATEGORY: EDB-400201

INIS CATEGORY: B12

<327>

Kononenko, V.I.; Lomovtsev, V.I.; Sukhman, A.L.

Adhesion properties of melted Al, Ga, In on solid titanium
 AIX-08:286443;EDB-77:043782; Izv. Akad. Nauk SSSR, Met.
 (IZNMA) (5):104-107 (Sep 1976)

SUBJECT DESCRIPTORS: ADHESION: Q1,Q2,Q3;ALUMINIUM: T1;GALLIUM:
 T2;GRAPHITE;HIGH TEMPERATURE;INDIUM: T3;LIQUID
 METALS;SOLUBILITY;SUBSTRATES;TEMPERATURE DEPENDENCE;TITANIUM:
 T4;VERY HIGH TEMPERATURE;VISCOSITY;WETTABILITY: Q4

ABSTRACT: None

EDE SUBJECT CATEGORY: EDB-360104

INIS CATEGORY: P22

<328>

Kononenko, V.I.; Yatsenko, S.P.

In Diffusion coefficients in molten metals **

NSA-33:30133;ERA-01:013612;EDB-76:036237; Tr. Inst. Khim.,
 Akad. Nauk SSSR, Ural. Fil. (TKAUA) 23::62-67 (1971)
 TT--74-58011

SUBJECT DESCRIPTORS: ACTIVATION

ENERGY;ANTIMONY;BISMUTH;CALCULATION METHODS;GALLIUM;INDIUM:
 T2;INDIUM ALLOYS: T3;LEAD;LIQUID METALS;SELF-DIFFUSION:
 Q1,Q2,Q3,Q4;SILVER;THALLIUM;TIN: T1;TIN ALLOYS:
 T4;USSR;VISCOSITY

ABSTRACT: The self-diffusion coefficients of indium and tin were
 determined experimentally in not only the pure metals but a
 wide range of binary liquid indium-tin alloys over the
 temperature range 800 to 1100⁰K. Activation energies
 and pre-exponential factors were calculated for each metal. A
 break in the Arrhenius plot was found. Coefficients of
 diffusion for Ag, In, Th, Sn, Pb, Sb, and Bi in liquid
 gallium were calculated from viscosity data. (GRA)

EDE SUBJECT CATEGORY: EDB-360102

<329>

Koroleva, G.N.; Gava, S.A.; Poluehktov, N.S.; Kirillov, A.I.;
 Kornelli, M.E.. (AN Ukrainskoj SSR, Odessa. Inst. Obshchej i
 Neorganicheskoj Khimii)

Interaction of lanthanide ions with fluorine-ions

AIX-08:295321;EDB-77:056246; Dokl. Akad. Nauk SSSR (DANKA)
 228:(6):1384-1386 (21 Jul 1976)

<329>

<329> CONT.

SUBJECT DESCRIPTORS: FLUORIDES; MEDIUM TEMPERATURE; PH
 VALUE; POTENTIOMETRY; QUANTITY RATIO; RARE EARTH COMPLEXES:
 T1; SOLUBILITY: Q1; STABILITY

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-400201
 INIS CATEGORY: B12

<330>

Košćurkov, A.A.; Korenev, Yu.M.; Novoselova, A.V.

Preparation of lithium octafluorohafnates EDB-76:036678; Zh.
 Neorg. Khim. (ZNOKA) 20:(2):543-544 (Feb 1975)

SUBJECT DESCRIPTORS: CHEMICAL COMPOSITION; CHEMICAL PREPARATION:
 Q1, Q2; CRYSTALS; HAFNIUM FLUORIDES: T1; LATTICE
 PARAMETERS; LITHIUM FLUORIDES: T2; STRUCTURAL CHEMICAL
 ANALYSIS; TETRAGONAL LATTICES

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-360602
 INIS CATEGORY: B12

<331>

Kotlin, V.P.; Barbanel, Y.A.

Use of the SF-8 detection spectrophotometer for measuring
 absorption spectra of molten salts
 AIX-08:322641; EDB-77:121463; Prih. Tekh. Eksp. (PRTEA)
 18:(4):232-233 (Jul 1975)

SUBJECT DESCRIPTORS: ABSORPTION SPECTRA:
 Q1; FOCUSING; FURNACES; LENSES; MOLTEN SALTS:
 T1; SPECTROPHOTOMETERS: T; VERY HIGH TEMPERATURE

ABSTRACT: Three versions of the use of the SF-8
 spectrophotometer for high-temperature measurements are
 described: 1) based on the standard optical scheme at
 temperatures of up to 400°C; 2) based on a modified
 one-wave scheme; 3) based on a modified twin-wave scheme. The
 external heater used in the schemes 2 and 3 ensures that
 measurements can be performed at temperatures of up to
 1000°C. In order to obtain the spectra of thin-layer
 samples vessels with an insert are used. For operation of the
 spectrophotometer according to the schemes 2 and 3 the heater

<331> CONT.

may in principle be replaced by a cryostat or some other external device.

EDB SUBJECT CATEGORY: EDB-440300

INIS CATEGORY: E42

<332>

Kouhsen, C.

In Preparation and thermochemical stability of uranium-zirconium-carbonitrides. **

AIX-07:270464; ERA-02:027039; EDB-77:055710; (1975) Technische Hochschule

SUBJECT DESCRIPTORS: CARBON NITRIDES; CHEMICAL PREPARATION; CHEMICAL REACTION KINETICS: Q2, Q3, Q4, Q5; ENTROPY; FORMATION HEAT; LATTICE PARAMETERS; METALLOGRAPHY: Q2, Q3, Q4, Q5; METALLURGY; PHASE DIAGRAMS; QUANTITATIVE CHEMICAL ANALYSIS; QUATERNARY ALLOY SYSTEMS; REDUCTION; SEGREGATION; SOLID FUELS; SOLID SOLUTIONS; STABILITY; URANIUM CARBIDES: T2; URANIUM NITRIDES: T3; ZIRCONIUM CARBIDES: T4; ZIRCONIUM NITRIDES: T5

ABSTRACT: This investigation deals with the preparation and the thermochemical stability of uranium-zirconium-carbonitrides as well as with the mechanism of (U,Zr) (C,N)-preparation by carbothermic reduction of uranium-zirconium-oxide. Single-phase (U,Zr) (C,N)-solid solutions with U:Zr-proportions of 3:1, 1:1, and 1:3 were prepared from oxide powder, obtained using the KFA-H-Process. The thermochemical stability of the (U,Zr) (C,N)-solid solutions against carbon was measured for varying Zr- and N-contents and for several temperatures; the results indicate an increase of the uranium carbide stability potential by the formation of (U,Zr) (C,N)-solid solutions. The thermodynamic properties ΔG^0 , ΔH^0 , and ΔS^0 were calculated and the correlation between the M(C,N)-lattice constant and the N-content was evaluated. Through an intensive investigation of the reaction mechanism, several different reaction paths were found; for each of them the characteristic diffusion of matter was explained by means of the microsections. It was shown that the Zr-concentration of the oxide reactant and the heating rate during the carbothermic reduction influence the species of the reaction product, especially the homogeneity of the (U,Zr) (C,N)-solid solution.

EDB SUBJECT CATEGORY: EDB-360204

INIS CATEGORY: B23

<333>

<333>

Kouhsen, C., Kernforschungsanlage Juelich G.m.b.H. (Germany, F.R.). Inst. fuer Reaktorwerkstoffe und Heisse Zellen; Technische Hochschule Aachen (Germany, F.R.)

In Preparation and thermochemical stability of uranium-zirconium-carbonitrides. A contribution to nuclear fuels for advanced high-temperature reactors **
AIX-07:265247; ERA-02:020575; EDB-77:036863; (Aug 1975)
Juel--1228

SUBJECT DESCRIPTORS: CARBON NITRIDES: T3; CHEMICAL PREPARATION: Q1, Q2, Q3, Q4, Q5; CHEMICAL REACTION KINETICS; ENTROPY; FORMATION HEAT; LATTICE PARAMETERS; METALLOGRAPHY; PHASE DIAGRAMS; QUANTITATIVE CHEMICAL ANALYSIS; QUATERNARY ALLCY SYSTEMS; REDUCTION; SEGREGATION; SOLID FUELS; SOLID SOLUTIONS; STABILITY: Q1, Q2, Q3, Q4, Q5; URANIUM CARBIDES: T1; URANIUM NITRIDES: T2; ZIRCONIUM CARBIDES: T4; ZIRCONIUM NITRIDES: T5

ABSTRACT: This investigation deals with the preparation and the thermochemical stability of uranium-zirconium-carbonitrides as well as with the mechanism of (U,Zr) (C,N)-preparation by carbothermic reduction of uranium-zirconium-oxide. Single-phase (U,Zr) (C,N)-solid solutions with U:Zr- proportions of 3:1, 1:1, and 1:3 were prepared from oxide powder. The thermochemical stability of the (U,Zr) (C,N)-solid solutions against carbon was measured for varying Zr- and N-contents and for several temperatures; the results indicate an increase of the uranium carbide stability potential by the formation of (U,Zr) (C,N)-solid solutions. The thermodynamic properties ΔG^0 , ΔH^0 , and ΔS^0 were calculated and the correlation between the M(C,N)-lattice constant and the N-content was evaluated. Through an intensive investigation of the reaction mechanism, several different reaction paths were found; for each of them the characteristic diffusion of matter was explained by means of the microsections. It was shown that the Zr-concentration of the oxide reactant and the heating rate during the carbothermic reduction influence the species of the reaction product, especially the homogeneity of the (U,Zr) (C,N)-solid solution.

EDB SUBJECT CATEGORY: EDB-360201; 050700
INTS CATEGORY: E23

<334>

Kouhsen, C.; Naoumidis, A.; Nickel, H. (Kernforschungsanlage Juelich G.m.b.H. (F.R. Germany). Inst. fuer Reaktorwerkstoffe)

Preparation and thermochemical stability of uranium-zirconium-carbonitrides
AIX-07:270478; ERA-02:015888; EDB-77:029153; J. Nucl. Mater. (JNUMA) 61:(1):88-98 (Jul 1976)

SUBJECT DESCRIPTORS: COATED FUEL PARTICLES: T1; ENTHALPY; HTGR TYPE REACTORS; LATTICE PARAMETERS; PRODUCTION: Q1, Q2, Q3, Q4, Q5; SOLID SOLUTIONS; STABILITY; TEMPERATURE DEPENDENCE; URANIUM CARBIDES: T2; URANIUM NITRIDES: T3; VEGARD LAW; VERY HIGH TEMPERATURE; X-RAY DIFFRACTION; ZIRCONIUM CARBIDES: T4; ZIRCONIUM NITRIDES: T5

ABSTRACT: Although the currently available coated particle fuel developed for use in gas-cooled HTRs can nominally be used up to at least 1250°C, in recent years for safety reasons the need to develop a kernel having greater compatibility with the PyC coating has increased. Uranium carbide is a potential high temperature fuel if it can be stabilized against reaction with PyC of the coating. One method is to combine the UC with nitrogen and zirconium. In this paper possible ways of preparing (U,Zr)(C,N)-solid solutions are described. The range of stability of (U,Zr)(C,N) against carbon and the thermodynamic properties (ΔG° , ΔH° , and ΔS°) were determined by heating the solid solution-carbon mixture to equilibrium. X-ray analysis was used to determine the (U,Zr)(C,N) lattice parameters as a function of nitrogen content, and it was found that there was a positive deviation from Vegard's law.

EDB SUBJECT CATEGORY: EDB-050700;360201
INIS CATEGORY: B24

<335>

Kovalevskii, A.V.; Lebedev, V.A.; Nichkov, I.F.; Raspopin, S.P.

Corrosion of liquid alloys of zinc with yttrium and with rare earth elements in LiCl--KCl melts EDB-77:132851; J. Appl. Chem. USSR (Engl. Transl.) (JAPUA) 46:(11):2743-2745 (Nov 1973)

SUBJECT DESCRIPTORS: BINARY ALLOY SYSTEMS; CORROSIVE EFFECTS: Q10, Q11; ELECTRIC POTENTIAL; ELECTROCHEMICAL CORROSION: Q1, Q2, Q3, Q4, Q5, Q6, Q7, Q8, Q9; LIQUIDS; LITHIUM CHLORIDES: M10; PHASE STUDIES; POTASSIUM CHLORIDES: M11; TEMPERATURE DEPENDENCE; ZINC BASE ALLOYS: M9

ABSTRACT: The corrosion of liquid alloys of zinc with yttrium and with rare earth elements (RE) in an LiCl--KCl melt was

<335>

<335> CONT.

investigated, and their stationary potentials were measured. The corrosion rates were calculated from these data, and the calculated values were compared with corrosion rates found experimentally. The experiments were conducted in the electrolytic cell in an atmosphere of purified argon. The alloys were prepared directly in the cell by fusion of Zn with freshly prepared shavings of the following metals: Ce, Nd, La, Dy, Er, Y, Sm, and Yb. The metal contents (8 to 10 wt.%) in the zinc corresponded to the two-phase state (saturated solution-compound) of the alloys. It is shown from experimental data that under equal conditions the corrosion of most of the alloys is approximately equal, and increases regularly with rise of temperature. The corrosion of alloys of zinc with samarium and especially with ytterbium is greater. Experimental rates of lanthanum and cerium corrosion in zinc alloys are compared with values calculated from the stationary potentials of the alloys. The good agreement between the values confirms the electrochemical mechanism of corrosion of Ce--Zn and La--Zn alloys.

EDB SUBJECT CATEGORY: EDB-360105

<336>

Kozachenko, E.G.; Safonov, V.V.; Ksenzenko, V.I. (Moskovskij Inst. Tonkoj Khimicheskoy Tekhnologii (USSR))

On interaction of tellurium tetrachloride with copper (1), mercury (2), bismuth and iron (3) chlorides EDB-76:036679; Zh. Neorg. Khim. (ZNOKA) 20:(2):562-563 (Feb 1975)

SUBJECT DESCRIPTORS: DIFFERENTIAL THERMAL ANALYSIS; MELTING; PHASE DIAGRAMS: Q1, Q2, Q3, Q4; QUANTITY RATIO; SOLID SOLUTIONS; TELLURIUM CHLORIDES: T1

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-360602

INIS CATEGORY: B12

<337>

Kozachenko, E.G.; Safonov, V.V.; Levina, N.A.; Ksenzenko, V.I. (Moskovskij Inst. Tonkoj Khimicheskoy Tekhnologii (USSR))

Phase diagrams of $\text{BiCl}_3\text{-CuCl-TeCl}_4$ and $\text{CuCl-FeCl}_3\text{-TeCl}_4$ systems AIX-08:336958; EDB-77:146484; Zh. Neorg. Khim. (ZNOKA) 21:(1):219-221 (Jan 1976)

ABSTRACT: Interaction in the systems $\text{TeCl}_4\text{-CuCl-BiCl}_3$ and $\text{FeCl}_3\text{-CuCl-TeCl}_4$ has been studied by differential-thermal analysis. In the system $\text{BiCl}_3\text{-CuCl-TeCl}_4$ the

<337> CONT.

nonvariant point E has a melting point 150 deg C; in the system $\text{CuCl}-\text{FeCl}_3-\text{TeCl}_4$ eutectic mixtures (points E_1 , E_2 and E_3) melt at 113 deg, 115 deg and 140 deg, respectively. Experimental and literature data on interaction of TeCl_4 with chlorides of impurities (BiCl_3 , CuCl , and FeCl_3) and with NaCl allow a conclusion to be made about possible separation of the main amounts of impurities from TeCl_4 with the aid of salt chromatography.

EDB SUBJECT CATEGORY: EDB-360602

INIS CATEGORY: B12

<338>

Kozeeva, L.P.; Konstanchuk, I.G.; Klevtsov, P.V.

Double tungstate $\text{K}_5\text{La}(\text{WO}_4)_4$ EDB-76:066305;
Inorg. Mater. (USSR) (Engl. Transl.) (INOMA) 11:(11):1800
(Nov 1975)

SUBJECT DESCRIPTORS: CHEMICAL PREPARATION: Q1,Q2,Q3,Q4;CRYSTAL
STRUCTURE; LANTHANUM TUNGSTATES: T2; LATTICE PARAMETERS; MELTING
POINTS; POTASSIUM TUNGSTATES: M3; RARE EARTH COMPOUNDS:
A4; TUNGSTATES: M1

ABSTRACT: The possibility of formation of $\text{K}_5\text{Ln}(\text{WO}_4)_4$ was investigated for $\text{Ln} = \text{La}, \text{Ce}, \text{Pr}, \text{Nd}, \text{Sm}, \text{Eu}, \text{Gd},$ and Y ; solid-phase reaction and crystallization from $\text{K}_2\text{W}_2\text{O}_7$ melt were used for the synthesis. X-ray phase analysis showed that only the compound $\text{K}_5\text{La}(\text{WO}_4)_4$ was formed; the others formed a mixture of two phases, $\text{KLn}(\text{WO}_4)_2$ and K_2WO_4 . $\text{K}_5\text{La}(\text{WO}_4)_4$ melts at $820 \pm 20^\circ\text{C}$; it is isostructural with palmierite, and its hexagonal lattice constants are $a = 6.030 \pm 0.008 \text{ \AA}$ and $c = 20.89 \pm 0.04 \text{ \AA}$. (DLC)

EDB SUBJECT CATEGORY: EDB-400201

<339>

Kozin, L.P.; Dergacheva, M.B.; Almazova, N.G.

Cs-Hg amalgam thermodynamic properties
AIX-08:283309; EDB-77:043773; Izv. Akad. Nauk SSSR, Met.
(IZNMA) (4):178-184 (Jul 1976)

SUBJECT DESCRIPTORS: BINARY ALLOY SYSTEMS; CESIUM ALLOYS:
T2; ELECTROMOTIVE FORCE; HIGH TEMPERATURE; LIQUID METALS; MERCURY
ALLOYS: T1; QUANTITY RATIO; REACTION
KINETICS; TABLES; THERMODYNAMIC PROPERTIES: Q1,Q2

<339>

<339> CONT.

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-360104

INIS CATEGORY: B22

<340>

Kozin, L.F.; Nigmatova, R.Sh.; Dairova, A.M.; Amangel'diev, B.R.

Thermodynamic properties of Cd-Zn-Hg liquid alloys

AIX-08:286444;EDB-77:043783; Izv. Akad. Nauk SSSR, Met.

(IZNMA) (5):224-228 (Sep 1976)

SUBJECT DESCRIPTORS: CADMIUM ALLOYS: T1; ENTHALPY; HIGH
TEMPERATURE; ISOTHERMAL PROCESSES; LIQUID METALS; MERCURY
ALLOYS: T2; PHASE STUDIES; QUANTITY RATIO; TEMPERATURE
DEPENDENCE; TERNARY ALLOY SYSTEMS; THERMODYNAMIC PROPERTIES:
Q1, Q2, Q3; VAPOR PRESSURE; ZINC ALLOYS: T3

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-360104

INIS CATEGORY: B22

<341>

Kozin, V.F.; Sheka, I.A. (AN Ukrainskoj SSR, Kiev. Inst.
Obshchej i Neorganicheskoj Khimii)

Study of kinetics of monovalent indium ions formation in the
In⁰-InCl₃-LiCl-H₂O system EDB-76:037013;
Ukr. Khim. Zh. (UKZHA) 41:(5):486-490 (May 1975)

SUBJECT DESCRIPTORS: ACTIVATION ENERGY; CATIONS; CHEMICAL REACTION
KINETICS: Q1, Q2, Q3, Q4; FORMATION HEAT; INDIUM: T1; INDIUM
CHLORIDES: T2; LITHIUM CHLORIDES: T3; QUANTITY
RATIO; TEMPERATURE DEPENDENCE; WATER: T4

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-400201

INIS CATEGORY: B12

<342>

Kozina, L.N.; Revyakin, A.V.; Romanova, V.P.

Solubility of oxygen in liquid niobium EDB-76:042266; Izv. Akad. Nauk SSSR, Met. (IZNMA) (3):28-33 (May 1975)

SUBJECT DESCRIPTORS: AFFINITY;EQUILIBRIUM;FREE ENERGY;LIQUID METALS;NIOBIUM: T1;NIOBIUM OXIDES;OXIDATION;OXYGEN: T2;SOLUBILITY: Q2;SOLVENT PROPERTIES: Q1;VERY HIGH TEMPERATURE

ABSTRACT: A method has been developed for determining the solubility of oxygen in liquid refractory metals possessing a strong affinity for oxygen and having volatile oxides. The equilibrium constant $K_{\text{sub}}(p)$ and the change of standard free energy ΔF^0 have been determined for the reaction which occurs when oxygen is dissolved in liquid niobium at 2800°K : $K_{\text{sub}}(p)=1.08 \times 10^4$ wt.%/at.; $\Delta F^0 = -52\ 150$ cal/mole.

EDB SUBJECT CATEGORY: EDB-360104

INIS CATEGORY: B22

<343>

Kozlov, V.B.; Karpenko, I.P.

Approximation of tabulated thermodynamic data for purposes of optimization of two-phase liquid-metal power cycles
ERA-02:034000;EDB-77:073116; Heat Transfer - Sov. Res.
(HTSRA) 8: (2):49-54 (1976)

SUBJECT DESCRIPTORS: ALKALI METALS: T3;CESIUM;DESIGN;ENTHALPY;ENTROPY;LIQUID METALS;LITHIUM;OPTIMIZATION;POTASSIUM;RUBIDIUM;SODIUM;THERMAL POWER PLANTS: T1;THERMODYNAMIC CYCLES: Q1;THERMODYNAMIC PROPERTIES: Q3;TWO-PHASE FLOW

ABSTRACT: Methods of utilization of tabulated thermodynamic properties of liquid-metal working fluids in computer calculations for purposes of optimization of power cycles are presented. Over a given temperature range of the two-phase flow the properties are represented as linear functions of the saturation temperature. Errors in determining the properties over a given temperature range are evaluated. Given the allowable error in determination of the properties, the maximum temperature range over which the method is applicable is determined.

EDB SUBJECT CATEGORY: EDB-200102

<344>

<344>

Kozlova, O.B. (Vsesoyuznyj Inst. Ogneuporov, Leningrad (USSR));
 Suvorov, S.A. (Leningradskij Tekhnologicheskij Inst. (USSR))

Wetting of certain refractories of the system $\text{MgO-Al}_2\text{O}_3\text{-ZrO}_2$ with metallic melts
 AIX-08:337300;EDB-77:140700; Ogneupory (OGNPA) (12):41-45
 (1976)

ABSTRACT: The wettability of refractory bricks ($\text{Al}_2\text{O}_3\text{-ZrO}_2$ and $\text{MgAl}_2\text{O}_4\text{-ZrO}_2$) by iron and aluminium has been investigated at 1550 and 1100 deg C, respectively. A study of the dependencies of the wetting angle THETA upon the retention time at the test temperature and the analysis of the brick microstructure subsequent to wetting by iron and aluminium show the materials examined to be wetted by iron from the moment of melting, the THETA value in corundum refractory bricks being two times less than that in spinel-based bricks. Least wettability by iron is characteristic of compositions comprising 85% $\text{MgAl}_2\text{O}_4 + 15\% \text{ZrO}_2$. Refractory bricks comprising 90% $\text{Al}_2\text{O}_3 + 10\% \text{ZrO}_2$ are wetted less by aluminium than bricks of different composition. The application of bricks comprising MgAl_2O_4 with a ZrO_2 admixture is recommended in the manufacture of insertfeeders for continuous steel teeming.

EDF SUBJECT CATEGORY: EDB-360204
 INIS CATEGORY: B23

<345>

Krascella, N.L., United Technologies Research Center, East
 Hartford, Conn. (USA)

In The spectral properties of uranium hexafluoride and its
 thermal decomposition products **
 ERA-02:037512;INS-77:010118;EDB-77:088303; (Sep 1976)
 N--76-33030

SUBJECT DESCRIPTORS: ABSORPTION;DECOMPOSITION: Q2;GAS FUELED
 REACTORS: M1;HELIUM;NUCLEAR FUELS: Q1;TEMPERATURE
 EFFECTS;ULTRAVIOLET RADIATION;ULTRAVIOLET SPECTRA: Q2;URANIUM
 HEXAFLUORIDE: M2;VERY HIGH TEMPERATURE

ABSTRACT: This investigation was initiated to provide basic spectral data for gases of interest to the plasma core reactor concept. The attenuation of vacuum ultraviolet (VUV) radiation by helium at pressures up to 20 atm over path lengths of about 61 cm and in the approximate wavelength range between 80 and 300 nm was studied. Measurements were also conducted to provide basic VUV data with respect to UF_6 and UF_6 /argon mixtures in the wavelength range between 80 and

<345> CONT.

120 nm. Finally, an investigation was initiated to provide basic spectral emission and absorption data for UF₆ and possible thermal decomposition products of UF₆ at elevated temperatures. (Author) (GRA)

EDB SUBJECT CATEGORY: EDB-220100;050700
INIS CATEGORY: E21

<346>

Krasnii, Yu.P.; Shvets, V.T. (Odesskij Gosudarstvennyj Univ. (Ukrainian SSR))

Electron contribution to the liquid metal viscosity
AIX-08:334577;EDB-77:132689; Ukr. Fiz. Zh. (Russ. Ed.)
(UFIZA) 21:(4):679-680 (Apr 1976)

SUBJECT DESCRIPTORS: CESIUM: T2; DIFFRACTION MODELS; ELECTRIC CONDUCTIVITY; ELECTRON DENSITY; FERMI LEVEL; LEAD; LIQUID METALS: T1; LITHIUM: T3; RELAXATION; SODIUM: T4; THERMAL CONDUCTIVITY; THERMOELECTRIC PROPERTIES; VISCOSITY: Q1,Q2,Q3,Q4

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-360104
INIS CATEGORY: B22

<347>

Krasnoperova, A.P. (Khar'kovskij Gosudarstvennyj Univ. (Ukrainian SSR)) ** Khar'kovskij Gosudarstvennyj Univ. (Ukrainian SSR)

Investigation of solubility and thermodynamics of dissolution of salts in mixed solvents In 8th radiochemical conference **
AIX-08:286087;EDB-77:069006; (1975) INIS-mf--3228

SUBJECT DESCRIPTORS:
DISSOLUTION; ENTHALPY; ENTROPY; ETHANOL; LABELLED COMPOUNDS; MEDIUM TEMPERATURE; METHANOL; MIXED SOLVENTS; SOLUBILITY: Q1; STRONTIUM 89; STRONTIUM NITRATES: T1; TEMPERATURE DEPENDENCE; TRACER TECHNIQUES

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-400201
INIS CATEGORY: B12

<348>

<348>

Krasulin, Yu.L.

Ceramic-base heat resisting constructional material

EDB-76:036582; Fiz. Khim. Obrab. Mater. (FKOMA) (5):62-69
(1974)

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-360203;360202

INIS CATEGORY: E23

<349>

Krivanek, M.; Krtil, J.; Sus, F.; Moravec, J. (Ustav Jaderneho
Vyzkumu, Rez (Czechoslovakia))Experience of the Central Control Laboratory (CCL) in accounting
for and controlling nuclear material in Czechoslovakia In
Safeguarding nuclear materials. Vol. 2 **

AIX-08:284456; ERA-02:019456; EDB-77:035061; (1976) IAEA

SUBJECT DESCRIPTORS: ACCOUNTING:

Q2; ACCURACY; BURNUP; CZECHOSLOVAKIA: T1; DISSOLUTION; FUEL
SCANNING; INVENTORIES; ISOTOPE RATIO; MASS SPECTROSCOPY; NUCLEAR
FUELS; NUCLEAR MATERIALS MANAGEMENT:
T2, Q1; PLUTONIUM; QUANTITATIVE CHEMICAL ANALYSIS; QUANTITY
RATIO; SAFEGUARDS; THORIUM; TITRATION; URANIUM; URANIUM DIOXIDEABSTRACT: The four years of experience obtained in accounting
for and controlling nuclear materials within the framework of
the Czechoslovak safeguards system and with the co-operation
of the IAEA is discussed. The methods for determining
uranium, thorium, and plutonium in various nuclear materials,
from the aspect of both chemical and isotopic composition,
have been developed and introduced into routine use. The
transport, storing, and preparation of samples for analysis,
as well as experimental devices, calibration, and comparative
tests, are involved. The errors evaluated are given. Other
methods have also been established at the CCL for solving
some problems in nuclear fuel control such, for example, as
the determination of burnup and nuclear spectrometry.
Participation in the PAFEX-I and PAFEX-II experiments is
noted.

EDB SUBJECT CATEGORY: EDB-055001

INIS CATEGORY: F41

<350>

Krohn, B.J.; Person, W.B.; Overend, J.

Predicted infrared spectrum and the structure of the isolated
 UF_5 molecule INS-76:020004; ERA-02:001214; EDB-76:081892;
 J. Chem. Phys. (JCPA) 65:(3):969-976 (1 Aug 1976)

SUBJECT DESCRIPTORS: ABSORPTION SPECTRA; INFRARED SPECTRA:
 Q1; MOLECULAR STRUCTURE: Q1; URANIUM FLUORIDES: T1; VIBRATIONAL
 STATES

ABSTRACT: The infrared absorption spectrum by the fundamental
 modes of UF_5 is calculated using force constants
 transferred from UF_6 to predict the wave numbers and
 using the F atom polar tensor from CH_3F to predict the
 intensities. Calculations are made for several assumed
 geometrical configurations of UF_5 . Comparison of the
 predicted spectra with the recently observed spectrum, in the
 UF-stretching region, of a photolysis product of UF_6
 isolated in an Ar matrix suggests strongly that the structure
 is square pyramidal (C_{4v}) with the U atom above
 the F atom equatorial plane. Although the model for the
 intensities is simple, the calculated spectrum is expected to
 predict the correct order of magnitude of the intensities of
 the fundamentals, first overtones, and binary combination
 bands of UF_5 , and the approximate wavenumbers for the
 expected absorption. Some comparisons are given between the
 calculations for the several structures of UF_5 and
 observed spectra of other XF_5 molecules, including
 PF_5 , BrF_5 , and IF_5 . (AIP)

EDB SUBJECT CATEGORY: EDB-400702
 INIS CATEGORY: B13

<351>

Kryukova, A.I.; Korshunov, I.A. (Gor'kovskij Gosudarstvennyj
 Univ. (USSR))

Solubility of $\text{Sr}(\text{NO}_3)_2$ in the LiNO_3 - KNO_3
 melt of variable composition EDB-76:037021; Zh. Neorg. Khim.
 (ZNOKA) 20:(5):1438-1440 (May 1975)

SUBJECT DESCRIPTORS: CRYSTALLIZATION; EUTECTICS; ISOTHERMAL
 PROCESSES; LITHIUM NITRATES: T2; MIXTURES; MOLTEN SALTS; PHASE
 DIAGRAMS: Q1, Q2, Q3; POTASSIUM NITRATES:
 T3; SOLUBILITY; STRONTIUM NITRATES: T1

<351>

<352> CONT.

AESTRACT: None

EDB SUBJECT CATEGORY: EDB-400201

INIS CATEGORY: B12

<352>

Kucera, G.H.; Sabounji, M.L.

Solubility of magnesium oxide in calcium oxide--calcium chloride mixtures ERA-02:005456:EDB-76:097750; Metall. Trans., B (MTTEC) 156:(7B):213-215 (Jun 1976)

SUBJECT DESCRIPTORS: CALCIUM CHLORIDES: T2; CALCIUM OXIDES: T3; MAGNESIUM OXIDES: T1; SOLUBILITY: Q1; SOLVENT PROPERTIES: Q2, Q3; VERY HIGH TEMPERATURE

ABSTRACT: Measurements and theoretical calculations of the solubility product of magnesium oxide in calcium chloride-calcium oxide mixtures at 1133⁰K are reported. A filtration technique in an inert atmosphere was used to obtain samples for analysis by atomic absorption for magnesium content. At low concentrations of 0²%, the data are in consonance with the very low solubility product of 10⁻⁶% predicted by a simple thermodynamic cycle first proposed by Flood, Foerland and Grjotheim.

EDB SUBJECT CATEGORY: EDB-360603

<353>

Kugel, R.; Williams, C.; Fred, M.; Malm, J.G.; Carnall, W.T.; Hindman, J.C.; Childs, W.J.; Goodman, L.S.

Isotope effects in the molecular spectrum of plutonium hexafluoride ERA-02:008299; INS-77:000577; EDB-77:009372; J. Chem. Phys. (JCPA) 65:(9):3486-3492 (1 Nov 1976)

SUBJECT DESCRIPTORS: ABSORPTION SPECTRA: Q1; EXCITED STATES; INFRARED SPECTRA; ISOTOPE EFFECTS: Q2, Q3; PLUTONIUM 239: T2; PLUTONIUM 242: T3; PLUTONIUM FLUORIDES: T1; VIBRATIONAL STATES

ABSTRACT: The first evidence for an isotope shift in the near infrared region of the spectrum of an actinide molecule, PuF₆, is reported. By use of samples of ²³⁹PuF₆ and ²⁴²PuF₆, isotope shifts were found in the absorption region near 8000 Å. Shifts of 0.3 to 1.2 cm⁻¹ were observed. The data are interpreted as indicating that some of the absorption bands in this region are vibronically induced electronic transitions which result from coupling of one or more quantum changes in the

<353> CONT.

asymmetric $\nu_{3}(F_{1u})$ and $\nu_{4}(F_{1u})$ vibrations to electronic states of the nonbonding 5f electrons in the molecule. (AIP)

EDB SUBJECT CATEGORY: EDB-640302;400202

INIS CATEGORY: A12;B12

<354>

Kuliasha, M.A.; Cronenberg, A.W. (New Mexico Univ., Albuquerque (USA))

Theoretical prediction of the thermal conductivity of uranium nitride vapor AIX-08:343064;EDB-78:021944; J. Nucl. Mater. (JNOMA) 67:(3):320-325 (Aug 1977)

SUBJECT DESCRIPTORS: CHEMICAL

REACTIONS;DISSOCIATION;ENTHALPY;LMFBR TYPE

REACTORS;MOLECULE-MOLECULE COLLISIONS;REACTOR CORE

DISRUPTION;THERMAL CONDUCTIVITY: Q1;URANIUM NITRIDES:

T1;VAPORS

ABSTRACT: A theoretical calculation of the thermal conductivity of uranium nitride vapor has been carried out. The calculational procedure is based on consideration of collisional, polyatomic, dissociation and reaction effects and only effects which do not perturb values by more than a factor of two are considered. Parameters such as the thermal conductivity of vapors are necessary for safety analyses of potential nuclear fuels for LMFBR reactors, such as uranium nitride.

EDB SUBJECT CATEGORY: EDB-360204

INIS CATEGORY: B24

<355>

Kuntz, L.K. (Ames Lab., IA); Bautista, R.G.

Heat capacities and heat content of molten cerium by levitation calorimetry INS-76:022924;EDB-76:097491; Metall. Trans., B (MTTBC) 7B:(1):107-113 (Mar 1976)

SUBJECT DESCRIPTORS: CALORIMETRY;CERIUM:

T1;CORRELATIONS;ENTHALPY: Q1;LEVITATION;LIQUID

METALS;SPECIFIC HEAT: Q1;VERY HIGH TEMPERATURE

ABSTRACT: A 15 kVa, 450 kHz radio frequency generator was used to levitate and melt cerium samples ranging in size from 0.4 to 1.7 g. The molten cerium samples were then dropped into a copper block calorimeter contained in isothermal surroundings. The cerium data between 1400 and 1700°K

<355>

<355> CONT.

indicated a nonlinear heat content function as described by the equation: $H/\text{sub } T/ \text{ -- } H_{\text{sub } 2}^{\text{sub } 9} \text{sub } 8^{\text{sub } 15} = \text{--}0.01255 T^{\text{sup } 2} + 75.305 T \text{ -- } 32,995.401 \text{ J/mol}$. The correlation coefficient for the curve fitting process was 0.81. The data between 1700 and 2500⁰K indicated a linear heat content function as described by the equation: $H/\text{sub } T/ - H_{\text{sub } 2}^{\text{sub } 9} \text{sub } 8^{\text{sub } 15} = [31.616 \text{ -- } 0.596](T - 1077) + [38,917.267 \text{ -- } 575.049] \text{ J/mol}$. The average percent deviation between this and the experimental values was 0.67. Experimental heat contents were corrected for convection and radiation heat losses during the fall of the sample from the levitation chamber into the calorimeter. The maximum estimated error for the levitation calorimetry work was ± 2.5 percent.

EDB SUBJECT CATEGORY: EDB-360104
INIS CATEGORY: B22

<356>

Kutsenko, S.A.; Amirova, S.A.; Chizhov, N.N. (Permskiy Politekhnikheskiy Inst. (USSR))

Influence of intermediate products on chlorination kinetics of vanadium oxides in salt melt AIX-07:274857;EDB-77:019304;
Izv. Vyssh. Uchebn. Zaved., Tsvet. Metall. (IVUTA) (3):75-78 (1975)

SUBJECT DESCRIPTORS: CHEMICAL REACTION KINETICS;CHEMICAL REACTION YIELD;CHLORINATION: Q1;MASS TRANSFER;MOLTEN SALTS;OXYCHLORIDES;VANADIUM OXIDES: T1

ABSTRACT: This study deals with the kinetics of chlorination of vanadium oxides in a melt of salts KCl, NaCl, KCl-NaCl. The rate-determining step of the process is the mass transfer of the intermediate product formed as a result of interaction of chlorine with the oxide ($K_{\text{sub } 2}^{\text{sub } 9} \text{sub } 8^{\text{sub } 15}$). Adding a carbonic reducer to the melt is conducive to a sharp increase in the rate of chlorination of the oxide stock. In this case, the rate-determining step of the process is the mass transfer of oxygen from the oxide to the carbon, which is aided by oxochlorides. The diffusion nature of the process is particularly manifested in the case of chlorination in an equimolar mixture of KCl and NaCl. In this case, the yield of $\text{VOCl}_{\text{sub } 3}$ is much higher than that of pure salts, which can be accounted for by the improved hydrodynamic characteristics

<357> CONT.

of the melt, owing to the fact that the KCl-NaCl mixture is less viscous as compared to pure salts.

EDB SUBJECT CATEGORY: EDB-400201

INIS CATEGORY: B12

<357>

Kuz'min, N.M.; Pelipas, V.P.; Tikhonova, A.I.; Khromov, A.D.

Investigation of some rare earth aluminates by the method of infrared spectroscopy AIX-08:283314;EDB-77:044046; Izv. Akad. Nauk SSSR, Neorg. Mater. (IVNMA) 12:(8):1424-1426 (Aug 1976)

SUBJECT DESCRIPTORS: ALUMINATES;COMPARATIVE EVALUATIONS;CRYSTAL STRUCTURE;FILMS;INFRARED SPECTRA: Q1,Q2;NEODYMIUM COMPOUNDS: T1;RARE EARTH COMPOUNDS: T2

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-360603;400201

INIS CATEGORY: P22

<358>

Kyrs, M.; Malek, Z.; Vesely, V.

In Twenty years of research in the processing of radioactive fuel and neutralization of radioactive waste at the Nuclear Research Institute (NRI) ** ERA-02:024089;EDB-77:047078; Jad. Energ. (JADEA) 21:(5):181-183 (1975) RFP-Trans--217

SUBJECT DESCRIPTORS: CZECHOSLOVAKIA;FLUORIDE VOLATILITY PROCESS;ION EXCHANGE;RADIOACTIVE WASTE PROCESSING: T;REPROCESSING: Q1;SPENT FUELS: T1

ABSTRACT: A complete review of the activities of workers in the field of radicchemistry over the past twenty years is given. It is subdivided into problems of the aqueous methods, fluoride method, and problems encountered in processing radioactive waste.

EDB SUBJECT CATEGORY: EDB-052001;050800

<359>

<359>

Lamparter, P.; Knoll, W.; Steeb, S., Max-Planck-Institut fuer
Metallforschung, Stuttgart (Germany, F.R.). Inst. fuer
Werkstoffwissenschaften

In Structure of molten Bi--Sb alloys by means of neutron
diffraction ** AIX-07:276420;EDB-77:061862; (1976)
AED-Conf--76-087-009

SUBJECT DESCRIPTORS: ANTIMONY ALLOYS: M1;BISMUTH ALLOYS:
M2;CHEMICAL COMPOSITION;COMPARATIVE EVALUATIONS;CRYSTAL
STRUCTURE;DIAGRAMS;INTERMETALLIC COMPOUNDS;LIQUID
METALS;MICROSTRUCTURE: Q1,Q2;NEUTRON DIFFRACTION;PHASE
STUDIES;QUANTITY RATIO;SPECTRA;STRUCTURAL CHEMICAL ANALYSIS

ABSTRACT: The structural investigations with melts can be
subdivided into two groups: The first group contains molten
metals and molten alloys, and one can state that the
structure of molten metals and of molten alloys nowadays is
rather well understood. Interference functions of molten
metals may be described by a hard sphere model. This is valid
also for molten alloys with statistical distribution. For the
second group, namely molten non-metals and molecular melts,
the interference functions as well as the pair correlation
functions are very often rather complicated and not well
understood. The present study is concerned with the
transition region between these two groups. It is shown that
the melts of the Bi-Sb system exhibit a change from metallic
to non-metallic structure. Regarding the experimental
details: the experiments were done with the two-axes
spectrometer D 4 at the high-flux reactor at Grenoble. The
containers consisted of cylindrical quartz tubes with a wall
thickness of 0.1 cm. The furnace consisted of a direct-heated
vanadium tube. The wavelength of the neutrons was 0.695 Å.
The final result is that the structure in molten Bi-Sb-alloys
consists of primitive tetrahedra with coordination number 3.
There are less tetrahedra in molten Bi than in molten Sb.
Also with rising temperature the number of tetrahedra
decreases. It is shown how to compose the coordination
numbers of two structures to get the observed coordination
number. The observed values are always the mean values of the
two structures.

EDB SUBJECT CATEGORY: EDB-360102
INIS CATEGORY: A13

<360>

Landresse, G.; Martinot, L.; Lysy, R.; Duyckaerts, G. (Liege Univ. (Belgium)) ** Mueller, W.; Lindner, R. (eds.)

Redox and chemical properties of U, Np and Pu in molten alkali chlorides In Transplutonium 1975 **

AIX-07:270268; ERA-02:020660; EDB-77:037517; (1976)
North-Holland

SUBJECT DESCRIPTORS: CHEMICAL PROPERTIES: Q1,Q2,Q3; CHEMICAL REACTION KINETICS; HIGH TEMPERATURE; MOLTEN SALTS; NEPTUNIUM: T1; NEPTUNIUM CHLORIDES: T5; OXIDATION; PLUTONIUM: T2; PLUTONIUM CHLORIDES: T4; REDOX REACTIONS: Q4,Q5,Q6; REDUCTION; SPECTROPHOTOMETRY; URANIUM: T3; URANIUM CHLORIDES: T6

ABSTRACT: Besides their fundamental interest, the chemical properties of uranium and the transuranium elements in molten alkali chlorides are of importance for the development of nuclear reactor technology. Their behaviour in redox reactions are best predicted by the potential versus pO_{H_2} ($-\log /C_{\text{H}_2}^{--}$) diagrams. In order to work out these diagrams, the standard potential of the various half-cell reactions corresponding to the different oxidation stages, were measured. Two methods have been used: (a) e.m.f. measurements were carried out in various alkali halides melts and at different temperatures and (b) spectrophotometric methods.

EEB SUBJECT CATEGORY: EDB-400702
INIS CATEGORY: B13

<361>

Laptev, D.M.; Kulagin, N.M.; Astakhova, I.S. (Sibirskii Metallurgicheskii Inst., Novokuznetsk (USSR))

Melting point of europium dichloride

AIX-07:265077; EDB-77:013485; Zh. Neorg. Khim. (ZNOKA)
20:(7):1987-1989 (Jul 1975)

SUBJECT DESCRIPTORS: ALLOTROPY; DIFFERENTIAL THERMAL ANALYSIS; EUROPIUM CHLORIDES: M1; MELTING POINTS: Q1; PHASE TRANSFORMATIONS; TRANSITION TEMPERATURE

ABSTRACT: Europium dichloride produced by hydrogen reduction of waterless EuCl_3 when exposed to $580-590^\circ\text{C}$ during 7-8h was studied by DTA method. Polymorphous transformation

<361>

<362> CONT.

was discovered with temperature equal to $747 \pm 5^\circ \text{C}$.
 Melting temperature of EuCl_2 is $254 \pm 2^\circ \text{C}$.

EDE SUBJECT CATEGORY: EDB-400201
 INIS CATEGORY: B12

<362>

Larin, A.A.; Vlasov, V.G.

Effect of oxygen content on the structural characteristics of
 uranium oxycarbides EDB-77:018938; Sov. Powder Metall. Met.
 Ceram. (Engl. Transl.) (SPMCA) 13:(9):761-763 (Sep 1974)

SUBJECT DESCRIPTORS: CRYSTAL STRUCTURE: Q1,Q2;PHASE
 STUDIES;URANIUM CARBIDES: T2;URANIUM OXIDES: T1

ABSTRACT: A study was made of the effect of oxygen content upon
 the structural characteristics of uranium oxycarbides. It is
 shown that uranium oxycarbides are deficient phases with an
 ordered disposition of vacancies. The fact that, with rise in
 the oxygen content of the oxycarbide, the number of vacant
 sites in both the sublattices increases and the crystal
 lattice parameter decreases is explained in terms of the
 occurrence of Me-Me and Me-X reactions. 14 refs.

EDE SUBJECT CATEGORY: EDB-360202;400702

<363>

Lau, K.F., California Inst. of Tech., Pasadena (USA)

In Pulse NMR in solids: chemical shift, lead fluoride, and
 thorium hydride ** INS-77:008225;EDB-77:069327; (1976)
 California Inst. of Tech.

SUBJECT DESCRIPTORS: ANISOTROPY;CALCIUM FLUORIDES: T1;CHEMICAL
 SHIFT: Q1;CRYSTAL STRUCTURE: Q4;DOPED MATERIALS;FLUORINE 19:
 T2;LEAD FLUORIDES: T3;MONOCRYSTALS;NUCLEAR MAGNETIC
 RESONANCE: Q2;PHASE TRANSFORMATIONS;PULSE TECHNIQUES;SODIUM
 FLUORIDES;SPECTRAL SHIFT;SPIN-LATTICE RELAXATION:
 Q3;TEMPERATURE DEPENDENCE;THORIUM HYDRIDES: T4;VERY HIGH
 PRESSURE

ABSTRACT: The fluorine chemical shift of a single crystal
 CaF_2 was measured up to 4 kilobar at room temperature
 using multiple pulse NMR. The pressure dependence of the
 shift is found to be -1.7 ± 1 ppm/kbar, while an overlap
 model predicts a shift of -0.46 ppm/kbar. The chemical shift
 tensor is separated into 'geometrical' and 'chemical'
 contributions, and comparison of the proposed model
 calculations with recent data on hydroxyl proton chemical

<363> CONT.

shift tensors shows that the geometrical portion accounts for the qualitative features of the measured tensors. A study of fluoride ion motion in β -PbF₂ doped with NaF was conducted by measurement of the ¹⁹F transverse relaxation time (T_2), spin lattice relaxation time (T_1) and the spin lattice relaxation time in the rotating frame ($T_{1\rho}$). Two samples of Th₄H₁₅, prepared under different conditions but both having the proper ratio of H/Th (to within 1 percent), were studied. The structure of the Th₄H₁₅ suggested by x-ray measurements is confirmed through a moment analysis of the rigid lattice line shape.

EDB SUBJECT CATEGORY: EDB-400702;400201

INIS CATEGORY: B13;B12

<364>

Lazarev, V.M.; Proshina, O.P.; Suponitskii, Yu.L.; Mayer, A.I.; Karapet'yants, M.Kh. (Moskovskij Khimiko-Tekhnologicheskij Inst. (USSR))

High-temperature calorimeter for determination of reaction enthalpy AIX-08:289600;EDB-77:044970; Zh. Fiz. Khim. (ZFKHA) 50:(8):2160-2161 (Aug 1976)

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-440300;400201

INIS CATEGORY: B22

<365>

Le Roy, J.; Moreau, J.M.; Paccard, D.; Parthe, E. (Geneva Univ. (Switzerland). Lab. Crystallographie aux Rayons X)

R₃T₂ compounds (R = rare earth or Y; T = Rh, Pd, Pt) with the rhombohedral Er₃Ni₂ structure type AIX-08:342994;EDB-78:021626; Acta Crystallogr., Sect. B (ACECA) 33:(8):2414-2417 (15 Aug 1977)

ABSTRACT: Sm₃Rh₂ crystallizes with the Er₃Ni₂ structure type. Space group R $\bar{3}$, a=8.701 (2), c=16.526 (7) Å, Z=9, D_{calc}(x)=9.06 g cm⁻³; μ (Mo K α)=430 cm⁻¹. Direct methods, absorption correction, least-squares refinement. R=0.11 for 451 independent reflexions. Ce₃Rh₂, Pr₃Rh₂, Nd₃Rh₂, La₃Pt₂, Ce₃Pt₂, Pr₃Pt₂, Nd₃Pt₂ and Y₃Pd₂ are isostructural with Sm₃Rh₂. The Er₃Ni₂ structure type is compared with the U₃Si₂ structure type which has been found with

<365>

<365> CONT.

rare-earth-palladium compounds. Both structures are characterized by transition-metal-centered trigonal double prisms and rare-earth-centred cubes.

EDS SUBJECT CATEGORY: EDB-360102

INIS CATEGORY: P22

<366>

Lee, H.J.; Kim, C.W.

Thermophysical properties of UO_2 fuel materials

ERA-02:028909;EDB-77:060199; J. Korean Nucl. Soc. (KNSJA)
8: (2):81-88 (Jun 1976)

SUBJECT DESCRIPTORS: DOPED MATERIALS; FLASH HEATING; GADOLINIUM
OXIDES; SPECIFIC HEAT; TEMPERATURE DEPENDENCE; THERMAL
CONDUCTIVITY; THERMAL DIFFUSIVITY; THERMODYNAMIC PROPERTIES:
Q1; URANIUM DIOXIDE: M1

ABSTRACT: A flash method for measuring the unknown thermal property (density, specific heat, or thermal diffusivity could be chosen as unknown) is described. The thermal diffusivity of UO_2 fuel samples is obtained from room temperature (300 K) to high temperature (1400 K). The specific heat is measured using a commercially available differential scanning calorimeter from room temperature to 500 K. The thermal conductivity of UO_2 fuel samples is calculated from the density, thermal diffusivity, and specific heat at constant pressure. Results are in complete agreement with the usual trends for the thermal conductivity of dielectric materials, in which impurity levels are very important at low temperatures but become relatively unimportant at high temperatures. In addition, the thermal diffusivity values at room temperature are reexamined by measuring the thermal diffusivity of several UO_2 fuel samples with the same level of dopant Gd_2O_3 .

EDS SUBJECT CATEGORY: EDB-050700;360204

<367>

Leibowitz, L.; Blomquist, R.; Fink, J.K., Argonne National Lab.,
Ill. (USA)

In Chemical Engineering Division, reactor safety and physical
property studies annual report, July 1974--June 1975 **
ERA-01:019331;EDB-76:059345; (1975) ANL--75-45

SUBJECT DESCRIPTORS: AFTER-HEAT REMOVAL: Q2; CHEMICAL REACTION
KINETICS: Q9; CORE CATCHERS; ENTHALPY; ENTROPY; LMFBR TYPE
REACTORS: T1; MATHEMATICAL MODELS; PHYSICAL PROPERTIES:

<367> CONT.

Q3,Q4,Q5,Q6,Q7,Q8;PLUTONIUM CARBIDES: T4;PLUTONIUM DIOXIDE:
T6;REACTOR CORE DISRUPTION: M2,Q1;REACTOR MATERIALS:
M10,Q1;REACTOR SAFETY;RESEARCH PROGRAMS: Q10;SODIUM:
T9;SPECIFIC HEAT;STAINLESS STEEL-304: T7;STAINLESS STEEL-316:
T8;THERMAL CONDUCTIVITY;URANIUM CARBIDES: T3;URANIUM DIOXIDE:
T5;VAPOR PRESSURE;VISCOSITY

ABSTRACT: Included are the following: a progress report on preparations to measure the enthalpies of (U, Pu)C and of sodium by drop calorimetry; progress on measurements of the speed of sound in sodium at high temperature; study of the vapor pressures of uranium and plutonium oxides by matrix isolation spectroscopy; evaluations and extrapolations of thermophysical property data for uranium carbide and steel; a discussion of studies aimed at measuring the thermal diffusivity of molten UO₂; an investigation of the interactions with sodium of various candidate sacrificial barrier materials; and an analysis of some phases of post-accident heat removal following a hypothetical core-disruptive accident.

EDB SUBJECT CATEGORY: EDB-220900;210500;360104;360204

<368>

Lemmer, R.H. (Rand Afrikaans Univ., Johannesburg (South Africa)); Viljoen, J.duP; (Atomic Energy Board, Pelindaba, Pretoria (South Africa))

Spin structure and elastic properties of uraniummononitride In Eleventh annual seminar on theoretical physics, Johannesburg, 5-9 July 1976 ** AIX-08:329695;NTS-78:000928;EDB-78:016954; (1976) INIS-mf--3670

SUBJECT DESCRIPTORS: ANTIFERROMAGNETISM;ELASTICITY:
Q1;HEISENBERG MODEL;MAGNETIC PROPERTIES: Q1;RÜDERMAN-KITTEL COUPLING;SPIN;SPIN WAVES;URANIUM NITRIDES: T1

ABSTRACT: The magnetic properties of uraniummononitride (UN) and the effect the magnetic properties have on the elasticity of the crystals are discussed in this lecture. The theoretical calculations of the longitudinal velocity of sound in the (001)-direction are compared with the experimental results.

EDB SUBJECT CATEGORY: EDB-360204
INIS CATEGORY: A34

<369>

<369>

Lengyel, T. (Magyar Tudományos Akadémia Izotop Intézete,
Budapest)

Investigations on the solvent extraction from molten salts by
tracer technique AIX-07:277828; EDB-77:026390; J. Radioanal.
Chem. (JRACE) 30:(2):433-440 (1976)

SUBJECT DESCRIPTORS: DISTRIBUTION

FUNCTIONS; DMSO; EUTECTICS; GALLIUM; MOLTEN SALTS; QUANTITY
RATIO; SOLVENT EXTRACTION: Q1; TEP; TERBIUM: T1; TERBIUM
160; TRACER TECHNIQUES; TTA

ABSTRACT: The distribution of terbium in a three-phase system of
an organic extractant, a fused salt and metallic gallium, as
well as in the corresponding two-phase subsystems has been
studied. Changing the concentrations in the organic and ionic
phases the distribution ratios can favorably be varied and so
the selectivity of extraction improved. The lithium
nitrate-ammonium nitrate eutectic (melting at 366[°]C) was
used as the ionic phase and gallium, the only multivalent
metal besides mercury forming a liquid at moderate
temperature represented the metallic one. Tri-n-butyl
phosphate, thenoyl-trifluoroacetone and dimethyl sulfoxide at
various concentrations diluted with decalin were used as
solvent extractants while terbium was chosen as the
representative of the lanthanide group, in the form of ¹⁶⁰Tb
tracer of high specific activity, enabling to keep
solute concentrations as low as 10⁻⁶ M. The equilibrium
measurements were performed at 383 K by shaking sealed glass
ampoules containing the three immiscible phases and the
solute in a thermostat for one hour. The distribution ratios
were calculated by determining the concentration of the
solute in aliquots of the samples radiometrically.

EDB SUBJECT CATEGORY: EDB-400105

INIS CATEGORY: F12

<370>

Lengyel, T. (Magyar Tudományos Akadémia Izotop Intézete,
Budapest)

Studies on some binary solvent systems for the extraction of
gadolinium AIX-08:346248; EDB-78:027829; Radiochem. Radioanal.
Lett. (RRAIA) 28:(5):453-458 (1977)

SUBJECT DESCRIPTORS: BINARY MIXTURES; DISTRIBUTION

FUNCTIONS; EUTECTICS; GADOLINIUM: T1; MOLTEN SALTS; SOLVENT
EXTRACTION: Q1; SYNERGISM

ABSTRACT: In addition to previous work regarding the extraction
of terbium from a molten eutectic several binary solvent

<370> CONT.

systems have been studied with respect to the efficiency of extraction of gadolinium. Though terbium and gadolinium as neighbouring rare earths behave rather similarly, due to synergistic effects some differences can be found allowing a selective separation of the lanthanons.

EDB SUBJECT CATEGORY: EDB-400105

INIS CATEGORY: B13

<371>

Lengyel, T. (Magyar Tudományos Akadémia Izotop Intézete, Budapest) ** Magyar Tudományos Akadémia Izotop Intézete, Budapest

Investigations of the solvent extraction from molten salts with the use of tracer technique In 8th radiochemical conference
** AIX-08:282946;EDB-77:056060; (1975) INIS-mf--3228

SUBJECT DESCRIPTORS: CHEMICAL COMPOSITION;EUTECTICS;MOLTEN SALTS: T2;ORGANIC SOLVENTS;QUANTITY RATIO;SOLVATION;SOLVENT EXTRACTION: Q1;SOLVENT PROPERTIES: Q2;SYNERGISM;TERBIUM: T1;TERBIUM 160;TERBIUM COMPLEXES;TRACER TECHNIQUES

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-400105

INIS CATEGORY: B12

<372>

Levin, Y.B.; Filippov, S.I. (Moskovskij Inst. Stali i Splavov (USSR))

Effect of basicity on ferrous oxide activity in complex oxide systems AIX-08:320662;EDB-77:113748; Izv. Vyssh. Uchebn. Zaved., Chern. Metall. (IVUMA) (7):5-11 (1975)

SUBJECT DESCRIPTORS: ALUMINIUM OXIDES: T5;CALCIUM OXIDES: M1;ELECTROMOTIVE FORCE;ENTHALPY;ENTROPY;IRON OXIDES: M2;MELTING;MOLTEN SALTS;REACTION KINETICS: Q1,Q2,Q3,Q4,Q5,Q6;SILICON OXIDES: M3;TEMPERATURE DEPENDENCE;THERMODYNAMICS;TITANIUM OXIDES: T6;VANADIUM OXIDES: M4

ABSTRACT: An empirical study was made of the thermodynamic properties of melts in relation to the CaO/SiO_2 ratio in the following systems: FeO-CaO-SiO_2 , $\text{FeO-CaO-SiO}_2\text{-Al}_2\text{O}_3$, $\text{FeO-CaO-SiO}_2\text{-TiO}_2$, $\text{FeO-CaO-SiO}_2\text{-V}_2\text{O}_5$. The activity of FeO in the solution of the oxides was determined by the electrochemical method in concentrated circuits with solid

<372>

<372> CONT.

electrolyte $\text{ZrO}_2(+\text{Y}_2\text{O}_3)$. From the obtained data calculations were made of the following: the relative partial molar and excessive partial molar isobar-isothermal potentials, enthalpy and entropy of mixing of FeO with the melt of the oxides. A comparison of $\alpha\text{-FeO}$ in oxide systems was made at constant CaO/SiO_2 ratio.

EDE SUBJECT CATEGORY: EDB-360204;400201

INIS CATEGORY: E12

<373>

Levinskii, Yu.V.

p--t diagram of the system uranium--nitrogen EDB-77:088914; Sov. At. Energy (Engl. Transl.) (SATEA) 37:(3):929-932 (Sep 1974)

SUBJECT DESCRIPTORS: NITROGEN;PHASE DIAGRAMS: Q1;URANIUM;URANIUM NITRIDES: T1

ABSTRACT: A more complete and detailed version of the p--T phase diagram for the uranium-nitrogen system is proposed.

EDE SUBJECT CATEGORY: EDB-360104;400702

<374>

Li, K.C., Iowa Univ., Iowa City (USA). Dept. of Chemistry

In Lines in the spectrum of ^7LiH (4728--5298 Å) **
INS-77:007315;ERA-02:030368;EDB-77:062485; (1976)
COO--2326-19

SUBJECT DESCRIPTORS: CRYSTALS;EMISSION SPECTRA: Q1;EXCITED STATES;ISOTOPE ENRICHED MATERIALS;LITHIUM;LITHIUM HYDRIDES: T1;WAVELENGTHS

ABSTRACT: The emission spectra of the $A^{1\Sigma^+}$ $+\text{X}^{1\Sigma^+}$ bands of ^7LiH were photographed in the 4728 Å - 5298 Å region with a 3.4 meter Ebert Spectrograph of theoretical resolution of about 0.07 cm^{-1} . High purity ^7LiH crystals were obtained from Oak Ridge National Laboratory. The atomic percent of ^7Li in ^7LiH and ^7LiD was 99.93 percent. The discharge source was a demountable stainless steel hollow cathode lamp. The lithium hydride crystals were packed into the cathode. Pressure in the discharge tube was about 10 to 20 torr of H_2 . The discharge was run at about 600 volts and 1.25 to 1.75 amperes. Acceptable spectra were obtained with exposure time of 6 hours. A Westinghouse iron hollow cathode was used to produce the iron spectrum for calibration. The plates were measured on the Gaertner

<374> CONT.

photoplate comparator with an encoder system and on-line computer service at Argonne National Laboratory. The measured lines in the spectra of ${}^7\text{LiH}$ are given in this report (${}^9\text{COO-2326-19}$). Similar spectra for ${}^6\text{LiH}$ and ${}^6\text{LiD}$ are given in companion reports (${}^9\text{COO-2326-17}$) and (${}^9\text{COO-2326-18}$), respectively. The relative intensities of the lines are applicable only to short regions and do not extend over the whole spectrum.

EDB SUBJECT CATEGORY: EDB-400201;640302;400104
INIS CATEGORY: B12;A12;B11

<375>

Liang, W.W.; Elliott, F.

Standard free energy of formation of liquid Na_2CrO_4 , 1100°K to 1220°K. ERA-01:020230;EDB-76:054371; J. Electrochem. Soc. (JESOA) 123:(5):617-620 (May 1976)

SUBJECT DESCRIPTORS: CHROMATES: T2;FORMATION FREE ENERGY: Q1,Q2;MOLTEN SALTS;SODIUM COMPOUNDS: T1;VERY HIGH TEMPERATURE

ABSTRACT: The standard free energy of sodium chromate was determined employing a reversible cell. Results give $2\text{Na(l)} + 5/4 \text{O}_2\text{(g)} + 1/2 \text{Cr}_2\text{O}_3\text{(s)} = \text{Na}_2\text{CrO}_4\text{(l)}$; $\Delta G^\circ = -648,900 + 106.94T$ (J/mole) and $2\text{Na(l)} + 2 \text{O}_2\text{(g)} + \text{Cr(s)} = \text{Na}_2\text{CrO}_4\text{(l)}$; $\Delta G^\circ = -1,209,900 + 233.1T$ (J/mole). These results are valid in the vicinity of 900°K. (auth)

EDB SUBJECT CATEGORY: EDB-400201;360603

<376>

Lindgard, P.

Theory of rare-earth alloys INS-77:017972;EDB-77:146249; Phys. Rev., B (PLRBA) 16:(5):2168-2176 (1 Sep 1977)

SUBJECT DESCRIPTORS: CRITICAL TEMPERATURE;EXCHANGE INTERACTIONS;FIELD THEORIES;PHASE DIAGRAMS: Q1;RARE EARTH ALLOYS: T1;SCALING LAWS

ABSTRACT: A mean-field random alloy theory combined with a simple calculation of the exchange interaction $J(c,Q)$ is shown to quantitatively account for the phase diagrams for alloys of rare-earth metals with Y, Lu, Sc, and other rare-earth metals. A concentration-dependent $J(c,Q)$ explains the empirical $2/3$ law and reconstitutes the idea of de Gennes scaling. The phase diagrams exhibit examples of both bicritical and tetracritical points. No significant

<376>

<376> CONT.

deviations from the mean-field calculation can be detected with the present experimental accuracy. A linear interpolation of $J(Q)$ for Gd and Er is found to account for all alloys except the Sc based. The exceptional behavior of the Sc alloys is due to a low density of states for Sc. A brief discussion is given of the effect on the mean-field results of changes in volume or c/a ratio and of critical fluctuations. Since the physical mechanisms of these ideal alloys are well documented they may serve as good candidates for studies of statistical effects such as multicritical phenomena or spin-glass phenomena.

EDB SUBJECT CATEGORY: EDB-360102

INIS CATEGORY: B22

<377>

Litvincv, Yu.G.; Il'yasov, I.I.; Savva, V.I. (Zaochnyj Inst. Sovetskoy Torgovli, Rostov-na-Donu (USSR))

Phase diagram of the ternary system of lithium-, rubidium-, and cesium nitrates AIX-08:292394; EDB-77:069048; Zh. Neorg. Khim. (ZNOKA) 20:(9):2560-2562 (Sep 1975)

SUBJECT DESCRIPTORS: CESIUM NITRATES: T1; EUTECTICS; LITHIUM NITRATES: T2; PHASE DIAGRAMS: Q1, Q2, Q3; RUBIDIUM NITRATES: T3

ABSTRACT: Two congruently melting compounds: LiNO_3 and RbNO_3 located at the two edges LiNO_3 - RbNO_3 and LiNO_3 - CsNO_3 define the character of the system. The solid solutions $(\text{Rb}, \text{Cs})\text{NO}_3$ with minimum 295 deg C inside the system decompose at near 260 deg C. Two triangular sections separate the system into three phase triangles producing two eutectic and one transition points.

EDB SUBJECT CATEGORY: EDB-400201

INIS CATEGORY: B12

<378>

Loechner, U.; Baernighausen, H.; Corbett, J.D.

Rare earth metal--metal halide systems. 19. Structural characterization of the reduced holmium chloride HoCl_2 ERA-03:010736; EDB-78:017287; Inorg. Chem. (INCCA) 16:(8):2134-2135 (Aug 1977)

SUBJECT DESCRIPTORS: CHEMICAL COMPOSITION; CRYSTAL STRUCTURE: Q1; HOLMIUM CHLORIDES: T1; X-RAY DIFFRACTION

ABSTRACT: The reduced holmium chloride HoCl_2 , with

<378> CONT.

the correct composition $\text{HoCl}_{2.20}$, is isomorphous with the monoclinic $\text{Dy}_{0.5}\text{Cl}_{1.1}$. The space group is $P2_1/m$. The crystal structure can be considered as a one-dimensional superstructure of the fluorite type built from five basic fluorite units with four additional anions per unit cell. The reason for the relatively large difference between the x-ray stoichiometry 2.20 and the analytical stoichiometry 2.14 found previously is tentatively attributed to contamination by monochloride. (JSR)

EDB SUBJECT CATEGORY: EDB-400201

<379>

Lovell, G.H.B.; van Tets, A. (Atomic Energy Board, Pelindaba, Pretoria (South Africa))

Solution studies in the $\text{US-UC}_{2.2}$ system

AIX-08:310011; EDB-77:094376; J. Nucl. Mater. (JNOMA) 64:(1):. 44-48 (Jan 1977)

SUBJECT DESCRIPTORS: CHEMICAL COMPOSITION; CRYSTALLOGRAPHY; CUBIC LATTICES; LATTICE PARAMETERS; NEUTRON DIFFRACTION; SOLID SOLUTIONS; SOLUBILITY: Q1, Q2; TETRAGONAL LATTICES; URANIUM CARBIDES: T1; URANIUM SULFIDES: T2; VERY HIGH TEMPERATURE; X-RAY DIFFRACTION

ABSTRACT: The mutual solubility of US and $\text{UC}_{2.2}$ was studied just below 1500°C to avoid a change of composition by $\text{CS}_{2.2}$ development. After cooling, the products were analysed by X-ray and neutron diffraction. Their essentially unchanged compositions were checked chemically. $\text{UC}_{2.2}$ was found to be soluble to a large extent in US. To a lesser extent, US is also soluble in the tetragonal $\text{UC}_{2.2}$ phase. A miscibility gap occurs which covers at least the 20 to 50 mol% US range.

EDB SUBJECT CATEGORY: EDB-360201

INIS CATEGORY: B24

<380>

Lufkin, D.A.; Paez, O.A.; Remanick, A.H.; Shaw, M., Whittaker Corp., San Diego, Calif. (USA). Narmco Research and Development Div.

In Electrochemical characterization of nonaqueous systems for secondary battery application. Quarterly report, February--April 1968 ** EDB-77:061394; (17 May 1968) N--68-28554

SUBJECT DESCRIPTORS: CALCIUM; CHEMICAL

<380>

<380> CONT.

REACTIONS;CHLORIDES;ELECTRIC BATTERIES:
 T1;ELECTROCHEMISTRY;ELECTRODES: Q1;ELECTROLYTES:
 Q1;FLUORIDES;LITHIUM;MAGNESIUM;PERCHLORATES;SOLUBILITY;TESTIN
 ;WIRES

ABSTRACT: None

EIE SUBJECT CATEGORY: EDB-250903

<381>

Lyall, A.E.; Seiger, H.N.; Crshich, J.; Michaels, R.G., Gulton
 Industries, Inc., Metuchen, N.J. (USA)

In Lithium--nickel fluoride secondary battery investigation.
 Quarterly technical progress report No. 1, 27 Apr--27 Jul
 1966 ** EDB-76:047839; (Aug 1966) AD--487828

SUBJECT DESCRIPTORS: CARBON BLACK;CARBONIC ACID ESTERS;CATHODES:
 Q1;DENSITY;DMSO;ELECTRIC CONDUCTIVITY;ELECTROLYTES:
 Q1;FABRICATION;GRAPHITE;LITHIUM;METAL-NONMETAL BATTERIES:
 T1;NICKEL FLUORIDES;PERFORMANCE;PHOSPHORUS
 FLUORIDES;POTASSIUM FLUORIDES;SODIUM
 PHOSPHATES;STARCH;VISCOSITY

ABSTRACT: None

EIE SUBJECT CATEGORY: EDB-250903

<382>

Lynch, D.W.; Olson, C.G. ** Koch, E.E. (ed.)

Rare earth 4d spectra in rare earth trifluorides In Vacuum
 ultraviolet radiation physics ** ERA-01:024881;EDB-76:067922;
 (1974) Friedr. Vieweg and Sohn Verlagsgesellschaft mbH

SUBJECT DESCRIPTORS: ABSORPTION SPECTRA:
 Q1,Q2,Q3,Q4,Q5,Q6,Q7,Q8,Q9,Q10,Q11,Q12;CERIUM FLUORIDES:
 T2;DYSPROSIUM FLUORIDES: T8;ERBIUM FLUORIDES: T10;FAR
 ULTRAVIOLET RADIATION;GADOLINIUM FLUORIDES: T6;HOLMIUM
 FLUORIDES: T9;LANTHANUM FLUORIDES: T1;NEODYMIUM FLUORIDES:
 T4;PRASEODYMIUM FLUORIDES: T3;SAMARIUM FLUORIDES: T5;TERBIUM
 FLUORIDES: T7;THULIUM FLUORIDES: T11;YTTERBIUM FLUORIDES: T12

ABSTRACT: Absorption spectra were measured for LaF_3 , 11
 rare earth trifluorides, and SmF_2 in the 80 to 260 eV
 range. The data for some of the compounds are shown. A

<382> CONT.

detailed comparison of the spectra with those of the metals will be published elsewhere; the agreement and differences in the data are given in this work. 7 figures. (GHT)

EDB SUBJECT CATEGORY: EDB-640302

<383>

Lyon, W.G.; Osborne, D.W.; Flotow, H.E.; Hoekstra, H.R. (Argonne National Lab., Ill. (USA))

Sodium uranium(V) trioxide, NaUO_3 : heat capacity and thermodynamic properties from 5 to 350 K
AIX-68:305658; EDB-77:107494; J. Chem. Thermodyn. (JCTDA) 9: (3):201-210 (Mar 1977)

SUBJECT DESCRIPTORS: CALORIMETRY; ENTHALPY: Q1, Q2; ENTROPY: Q1, Q2; FREE ENTHALPY: Q1, Q2; LOW TEMPERATURE; MEDIUM TEMPERATURE; SODIUM OXIDES: T1; SPECIFIC HEAT: Q1, Q2; ULTRALOW TEMPERATURE; URANIUM OXIDES: T2; VERY LOW TEMPERATURE

ABSTRACT: The heat capacity of a sample of NaUO_3 was determined in the temperature range 5 to 350 K by aneroid adiabatic calorimetry. The heat capacity at constant pressure $C_{p,m}(298.15 \text{ K})$, the entropy $S_m(298.15 \text{ K})$, the enthalpy $(H_m(298.15 \text{ K}) - H_m(0))$ and the function $-(G_m(298.15 \text{ K}) - H_m(0))/298.15 \text{ K}$ were found to be $(108.87 \pm 0.11) \text{ J K}^{-1} \text{ mol}^{-1}$, $(132.84 \pm 0.13) \text{ J K}^{-1} \text{ mol}^{-1}$, $(20176 \pm 20) \text{ J mol}^{-1}$, and $(65.17 \pm 0.06) \text{ J K}^{-1} \text{ mol}^{-1}$, respectively. A non-isothermal peak in the heat capacity, probably correlated with magnetic ordering, was noted at $(31 \pm 1) \text{ K}$.

EDB SUBJECT CATEGORY: EDB-400702
INIS CATEGORY: B13

<384>

Lyon, W.G.; Osborne, D.W.; Flotow, H.E.; Hoekstra, H.R.

Sodium uranium(V) trioxide, NaUO_3 : heat capacity and thermodynamic properties from 5 to 350 K
ERA-03:004500; EDB-77:146513; J. Chem. Thermodyn. (JCTDA) 9: 201-210 (1977)

SUBJECT DESCRIPTORS: LOW TEMPERATURE; MEDIUM TEMPERATURE; SODIUM OXIDES: T1; SPECIFIC HEAT; THERMODYNAMIC PROPERTIES: Q1, Q2; ULTRALOW TEMPERATURE; URANIUM OXIDES: T2

ABSTRACT: The heat capacity of a sample of NaUO_3 was determined in the temperature range 5 to 350 K by aneroid adiabatic calorimetry. The heat capacity at constant pressure

<384>

<384> CONT.

$C_p^0(298.15 \text{ K})$, the entropy $S^0(298.15 \text{ K})$, the enthalpy $[H^0(298.15 \text{ K}) - H^0(0)]$ and the function $-[G^0(298.15 \text{ K}) - H^0(0)]/298.15 \text{ K}$ were found to be $(108.87 \pm 0.11) \text{ J K}^{-1} \text{ mol}^{-1}$, $(132.84 \pm 0.13) \text{ J K}^{-1} \text{ mol}^{-1}$, $(20176 \pm 20) \text{ J mol}^{-1}$, and $(65.17 \pm 0.06) \text{ J K}^{-1} \text{ mol}^{-1}$, respectively. A non-isothermal peak in the heat capacity, probably correlated with magnetic ordering, was noted at $(31 \pm 1) \text{ K}$.

EDE SUBJECT CATEGORY: EDB-360603

<385>

Lysy, R.; Duyckaerts, G. (Liege Univ. (Belgium))

Quantitative study of the chemical equilibria in solution in molten LiCl-KCl (70-30 mol.%) by absorption spectrophotometry. Application to neptunium
AIX-07:239659; EDB-77:013693; Inorg. Nucl. Chem. Lett. (INUCA) 12: (2):205-215 (1976)

SUBJECT DESCRIPTORS: ABSORPTION SPECTRA; CHEMICAL REACTION KINETICS; Q1; CHEMICAL REACTIONS; EQUILIBRIUM; HIGH TEMPERATURE; LITHIUM CHLORIDES; MOLTEN SALTS; NEPTUNIUM: T1; NEPTUNIUM OXIDES; POTASSIUM CHLORIDES; RADICALS; SOLUTIONS; SPECTROPHOTOMETRY

ABSTRACT: A quantitative study has been made, in a LiCl-KCl (70 to 30 mol. %) melt, of the following reactions: $\text{NpO}_2 + 4\text{HCl}$ reversible $\text{Np}^{4+} + 2\text{H}_2\text{O} + 1/2\text{Cl}_2 + 3\text{Cl}^-$; and $\text{NpO}_2 + \text{Cl}_2$ reversible $\text{NpO}_2^{2+} + 1/2\text{Cl}_2 + 2\text{Cl}^-$, in the temperature range of 450 to 600°C, by sparging through the melt gas mixtures (H_2 , HCl , Cl_2 , N_2) of known composition. The concentrations of the neptunium species were measured by a spectrophotometric method. The equilibrium constants, at 450°C, are respectively: $K_1 = (4.1 \pm 0.8) \times 10^{-6} \text{ atm}^{-3/2}$; and $K_2 = (52 \pm 13) \text{ atm}^{1/2}$. The standard potential $E^\circ(\text{NpO}_2^{2+}(\text{VI})/\text{NpO}_2^{2+}(\text{V}))$ at 450°C equals $(0.25 \pm 0.02) \text{ V}$ against the standard chlorine potential.

EDE SUBJECT CATEGORY: EDB-400702

INIS CATEGORY: B13

<386>

Lysy, R.; Duyckaerts, G. (Liege Univ. (Belgium))

Determination of the Np(IV)--Np(III) normal potential in fused
LiCl-CsCl (55-45%) by absorption spectrophotometry
AIX-08:346251;EDB-78:028033; Rev. Chim. Miner. (RVCMA)
14:(2):119-126 (1977)

SUBJECT DESCRIPTORS: ABSORPTION SPECTRA; CATIONS; CESIUM
CHLORIDES; ENTHALPY; ENTROPY; HIGH TEMPERATURE; LITHIUM
CHLORIDES; MOLTEN SALTS; NEPTUNIUM: T1; REDOX POTENTIAL:
Q1; SPECTROPHOTOMETRY

ABSTRACT: The following reaction $\text{Np}^{4+} + \frac{1}{2} \text{H}_2 + \text{Cl}^- \rightleftharpoons \text{Np}^{3+} + \text{HCl}$ has been studied quantitatively in fused LiCl-CsCl (55-45 mol %) in the temperature range of 450-600°C, by visible and near IR absorption spectrophotometry. At 450 deg C, the equilibrium constant equals $K = 7.4 \text{ atm}^{1/2}$. The mean values of ΔH and ΔS in the temperature range of 450-600 deg C are: $\Delta H = (35.6 \pm 10.3) \text{ kJ mol}^{-1}$, $\Delta S = (65.5 \pm 13.0) \text{ J mol}^{-1} \text{ K}^{-1}$.

EDE SUBJECT CATEGORY: EDE-400702

INIS CATEGORY: B13

<387>

MacDonald, P.E.; Thompson, L.B. (eds.), EG and G Idaho, Inc.,
Idaho Falls (USA)

In MATPRO: Version 09. A handbook of materials properties for
use in the analysis of light water reactor fuel rod behavior
** FRA-02:028904; INS-77:007752; EDB-77:065884; (Dec 1976)
TREE-NUREG--1005

ABSTRACT: This handbook describes the materials properties
correlations and computer subcodes (MATPRO-Version 09)
developed for use with various LWR fuel rod behavior
analytical programs at the Idaho National Engineering
Laboratory. Documentation and formulations that are generally
semiempirical in nature are presented for uranium dioxide and
mixed uranium-plutonium dioxide fuel, zircaloy cladding, gas
mixture, and LWR fuel rod material properties.

EIB SUBJECT CATEGORY: EDB-050700;360204;360203;210100;210200

INIS CATEGORY: E23;E31;E32

<388>

<388>

Maiorcva, E.A.; Morachevskii, A.G. (Leningradskij
Politekhnikheskij Inst. (USSR))

Thermodynamic properties of diluted sodium solutions in liquid
indium AIX-08:334317;EDB-77:140533; Elektrokhiimiya (ELKKA)
12:(12):1836-1838 (Dec 1976)

SUBJECT DESCRIPTORS: BINARY ALLOY SYSTEMS;ELECTROMOTIVE
FORCE;HIGH TEMPERATURE;INDIUM ALLOYS: T1;LIQUID
METALS;REACTION KINETICS;SODIUM ALLOYS: T2;THERMODYNAMIC
PROPERTIES: Q1,Q2

ABSTRACT: None

EDE SUBJECT CATEGORY: EDB-360104
INIS CATEGORY: B12

<389>

Maiti, H.S.; Subbarao, E.C.

Electrical conduction in CaO-doped thoria electrolytes
INS-77:006417;EDB-77:055746; J. Electrochem. Soc. (JESCA)
123:(11):1713-1718 (Nov 1976)

SUBJECT DESCRIPTORS: CALCIUM OXIDES: T2;CHEMICAL
COMPOSITION;CRYSTAL DEFECTS;DOPED MATERIALS;ELECTRIC
CONDUCTIVITY: Q1,Q2;ELECTROLYTES: Q3;HIGH-TEMPERATURE FUEL
CELLS: T3;HOLES;IONIC CONDUCTIVITY;OXYGEN;PRESSURE
DEPENDENCE;QUANTITY RATIO;TEMPERATURE DEPENDENCE;THORIUM
OXIDES: T1;VACANCIES

ABSTRACT: The electrical conductivity of sintered specimens of
nominally pure thoria and thoria doped with 1-15 mole percent
(m/o) CaO was measured between 600⁰ and 1400⁰C in
the P₀₂ range of 1-10⁻²⁵ atm. The measured
conductivity decreases initially with decreasing P₀₂
(mixed conduction region where electron holes and oxygen ions
contribute), until it becomes independent of P₀₂ (ionic
conduction region). The P₀₂ corresponding to the
transition between these regions is a function of temperature
and composition. The variation of conductivity with
temperature gives rise to three regions: the high temperature
region where the conductivity is attributed to dissociated
vacancies, an intermediate temperature range where associated
vacancy-impurity complexes dominate, and a low temperature
region where grain boundary effects appear to become
significant. The activation energy for electrical conduction

<389> CONT.

is a function of composition and P_{02} . (7 figures, 31 references)

EDB SUBJECT CATEGORY: EDB-360204;300503
INIS CATEGORY: B23;A16

<390>

Makarcv, L.L. (Leningradskij Gosudarstvennyj Univ. (USSR))

Origin of excess entropy of alkali metals halides solid solutions AIX-08:320671;EDB-77:113904; Zh. Fiz. Khim. (ZFKHA) 49:(9):2220-2223 (Sep 1975)

ABSTRACT: The formation of solid solutions of alkali metal halides has been considered in the case of KI-RbI and CsCl-CsBr systems. The difference is determined of the sublattice of the common ion and that of miscible ions one where ion vibration frequencies may vary only under the influence of the force field factor. The kinetic factor $S_{sup}(E)$ (f_{mu}) in $S_{sup}(E)$, the latter being presented as a sum of two items corresponding to kinetic and potential effects has been calculated. The calculated data are compared with the known experimental values of $S_{sup}(E)$. An agreement has been observed for all the systems over the whole range of compositions. A change in $S_{sup}(E)$ reflects correctly a transition from system to system when the common ion is changed and from structure to structure, it allows the origin to be understood of the peculiarities in the behaviour of the excessive entropy of solid solutions due to the reduced mass effect.

EDB SUBJECT CATEGORY: EDB-360603;400201
INIS CATEGORY: B12

<391>

Makhmadmurodov, A.; Dudchik, G.P.; Polyachenok, O.G.
(Belcrusskij Tekhnologicheskij Inst., Minsk)

Pressure of saturated vapor of rare earth and yttrium bromides AIX-08:282952;EDB-77:037347; Zh. Fiz. Khim. (ZFKHA) 49:(10):2714-2715 (Oct 1975)

SUBJECT DESCRIPTORS: ENTHALPY;ENTROPY;EQUATIONS;EVAPORATION;HIGH TEMPERATURE;RARE EARTH COMPOUNDS: T1;SUBLIMATION;TEMPERATURE DEPENDENCE;VAPOR PRESSURE: Q1,Q2;YTTRIUM BROMIDES: T2

<391>

<392> CONT.

ABSTRACT: None

EDE SUBJECT CATEGORY: EDB-400201

INIS CATEGORY: B12

<392>

Mal, H.H. van; Buschow, K.H.J.; Miedema, A.R. (Philips Research Labs., Eindhoven (The Netherlands))

Hydrogen absorption of rare-earth (3d) transition intermetallic compounds AIX-07:278235; EDB-77:030837; J. Less-Common Met. (JCOMA) 49: (1-2):473-475 (Sep 1976)

SUBJECT DESCRIPTORS: CRYSTAL STRUCTURE; HYDRIDES; HYDROGEN: T4; INTERMETALLIC COMPOUNDS; RARE EARTH ALLOYS: T1; SOLUBILITY: Q4; SOLVENT PROPERTIES: Q1, Q2, Q3; THORIUM ALLOYS: T2; TRANSITION ELEMENT ALLOYS: T3

ABSTRACT: It is shown that the absorption of large amounts of hydrogen by rare-earth 3d metal compounds and thorium 3d metal compounds is a general phenomenon, not depending much on crystal structure or on the relative concentration of the rare-earth metal.

EDE SUBJECT CATEGORY: EDB-360104; 080201

INIS CATEGORY: F22

<393>

Mal'kevich, N.V.; Grishin, I.A.

In Properties of yttrium and scandium basic salts **

ERA-01:026214; INS-76:018800; EDB-76:075637; Uch. Zap. Gor'k. Gos. Univ. (UZGUA) (105):34-38 (1970) ORNL-tr-4199

SUBJECT DESCRIPTORS: PHASE STUDIES; PYROLYSIS: Q1, Q2, Q3, Q4, Q5, Q6; SCANDIUM HYDROXIDES: M1; SCANDIUM SULFATES: M2; THERMAL GRAVIMETRIC ANALYSIS; X-RAY DIFFRACTION; YTTRIUM CHLORIDES: M5; YTTRIUM HYDROXIDES: M3; YTTRIUM NITRATES: M6; YTTRIUM SULFATES: M4

ABSTRACT: Thermographic and x-ray phase analysis methods were used to study the basic salts $\text{Sc}(\text{OH})_{2.32}(\text{SO}_4)_{0.34}$, $\text{Y}(\text{OH})_{2.5}(\text{SO}_4)_{0.5}$, $\text{Y}(\text{OH})_{2.6}(\text{Cl})_{0.4}$, and $\text{Y}(\text{OH})_{2.6}(\text{NO}_3)_{0.4}$, schemes for their thermal decay

<394> CONT.

are presented. It is noted that the basic salts have a weak crystalline structure.

EDB SUBJECT CATEGORY: EDB-400201

INIS CATEGORY: B12

<394>

Mal'tsev, V.T.; Cherpillo, V.P.; Machavariani, Z.I.
(Rostovskij-na-Donu Inzhenernc-Stroitel'nyj Inst. (USSR))

Phase equilibria in the $\text{PbO-GeO}_{0.5}\text{-V}_{0.5}$ ternary system AIX-07:274861; EDB-77:018943; Zh. Neorg. Khim. (ZNOKA) 20: (7):1961-1964 (Jul 1975)

SUBJECT DESCRIPTORS: EQUILIBRIUM;GERMANIUM OXIDES: T1;LEAD OXIDES: T2;MOLTEN SALTS;PHASE DIAGRAMS: Q1,Q2,Q3;PHASE STUDIES;VANADIUM OXIDES: T3;X-RAY DIFFRACTION

ABSTRACT: Visual-polythermal and X-ray phase analysis techniques have been used to study the ternary system $\text{PbO-GeO}_{0.5}\text{-V}_{0.5}$. There have been defined the fields of primary crystallization of the starting component PbO , binary phases and three ternary congruently melting compounds of the following composition: $15\text{PbO}\cdot 4\text{GeO}_{0.5}\cdot 3\text{V}_{0.5}$; $19\text{PbO}\cdot 2\text{GeO}_{0.5}\cdot 2\text{V}_{0.5}$; $17\text{PbO}\cdot 2\text{GeO}_{0.5}\cdot \text{V}_{0.5}$. It has been established that two compounds with approximately the following compositions: $5\text{PbO}\cdot 6\text{GeO}_{0.5}\cdot 9\text{V}_{0.5}$ and $6\text{PbO}\cdot \text{GeO}_{0.5}\cdot 2\text{V}_{0.5}$ are formed in the subsolidus region. The region of stratification in the liquid phase of the ternary melts, adjacent to the binary system $\text{GeO}_{0.5}\text{-V}_{0.5}$ is limited. The melts adjacent, by composition, to the binary system $\text{PbO-GeO}_{0.5}$ and rich in $\text{GeO}_{0.5}$, as well as those adjacent to the middle portion of the binary system $\text{PbO-V}_{0.5}$, show a tendency to vitrification.

EDB SUBJECT CATEGORY: EDB-360202

INIS CATEGORY: B12

<395>

Malinin, G.V.; Tolmachev, Yu.M. (Radievyyj Inst., Leningrad (USSR))

Thermal decomposition of solid metal oxides EDB-76:037148; Usp. Khim. (USKHA) 44: (5):782-792 (May 1975)

SUBJECT DESCRIPTORS: ACTIVATION ENERGY;CHEMICAL REACTION KINETICS;DISSOCIATION;PRASEODYMIUM OXIDES: M2;PYROLYSIS: Q1,Q2;REACTION HEAT;REVIEWS;SOLIDS;URANIUM OXIDES: M1

<395>

<395> CONT.

ABSTRACT: None

EED SUBJECT CATEGORY: EDB-400702;400201

INIS CATEGORY: B12

<396>

Malinin, S.D.; Dernov-Pegarev, V.F.

Investigation of calcite solubility in K_2CO_3 and Na_2CO_3 solutions at temperatures of 200 to 350°C EDB-76:033940; Geokhimiya (GEOKA) (3):454-462 (Mar 1974)

SUBJECT DESCRIPTORS: CALCITE: T1; HIGH TEMPERATURE; POTASSIUM CARBONATES; POTASSIUM CHLORIDES; SODIUM CARBONATES; SOLUBILITY: Q1; SOLUTIONS

ABSTRACT: As a result of an experimental study of $CaCO_3$ (Iceland spar) in K_2CO_3 , Na_2CO_3 solutions, and mixed solutions- $K_2CO_3 + KCl$ and $K_2CO_3 + Na_2CO_3$ with concentration from 1 to 15 molal, the existence of the $Ca(CO_3)_2$ complex has been established. $PK_c = -\log K_c$ is equal to 4.5 $\pm$ 0.5 at 200°C, 3.8 $\pm$ 0.3 at 300°C and 3.3 $\pm$ 0.2 at 350°C. The high solubility of $CaCO_3$ in solutions of alkali carbonates attaining 24 g/1000 g in 75 % of K_2CO_3 solution at 350°C together with data of mineralogical observations gives ground to consider natural carbonate solutions as probable agents giving rise to carbonatite body formation.

EED SUBJECT CATEGORY: EDB-152001;400201

<397>

Malyshev, V.V. (Gosudarstvennyj Komitet po Ispol'zovaniyu Atomnoj Ehnergii SSSR, Moscow. Inst. Atomnoj Ehnergii)

Thermodynamic similarity and universal equations of state of hexafluorides AIX-08:334318; EDB-77:141015; Teplofiz. Vys. Temp. (TVYTA) 14:(1):47-55 (Jan 1976)

SUBJECT DESCRIPTORS: COMPARATIVE EVALUATIONS; DENSITY; EQUATIONS OF STATE; MOLYBDENUM FLUORIDES: T1; THERMODYNAMIC PROPERTIES: Q1, Q2, Q3; THERMODYNAMICS; TUNGSTEN FLUORIDES: T2; URANIUM HEXAFLUORIDE: T3; VAPOR PRESSURE

ABSTRACT: There have been generalized the experimental data on the thermodynamic properties of hexafluorides of sulfur, molybdenum, tungsten, uranium in Van der Waals region. The similarity criteria have been checked up both graphically by

<397> CONT.

plotting the curves with the respective parameters and analytically by determining the coefficients of the appropriate universal equations of state. Seven criteria of thermodynamic similarity have been shown to be realized. There have been found the universal equations for the state of the hexafluorides in the region of the rarefied and thick gas, as well as for vapour saturation, equilibrium densities of vapour and liquid, heat of evaporation.

EDB SUBJECT CATEGORY: EDB-400201;400702
INIS CATEGORY: B12

<398>

Manes, L.; Naegele, J. (Commission of the European Communities, Karlsruhe (Germany, F.R.). European Inst. for Transuranium Elements) ** Blank, H.; Lindner, R. (eds.)

Electronic properties of actinide oxides In Plutonium 1975 and other actinides ** AIX-08:310014; ERA-02:041829; EDB-77:094387; (1976) North-Holland

SUBJECT DESCRIPTORS: BINDING ENERGY; CHEMICAL BONDS: Q1, Q2, Q3; CHEMICAL COMPOSITION; COVALENCE; CRYSTAL STRUCTURE; ELECTRICAL PROPERTIES; ELECTRONIC STRUCTURE; ENERGY GAP; MAGNETIC PROPERTIES; OPTICAL PROPERTIES; PLUTONIUM OXIDES: T1; SPECTROSCOPY; THERMODYNAMIC PROPERTIES; THORIUM OXIDES: T2; URANIUM OXIDES: T3

ABSTRACT: The available information of a direct nature on the properties of electrons in actinide oxides is critically surveyed. Some critical assessment is made on the new ideas about the bonding of these systems, ideas which have emerged mostly from thermodynamical and structural considerations. The aim in doing this is to a) judge how many of these ideas can be supported by direct experimental evidence on the electronic properties; b) attempt, on the other hand, to imagine which experimental results are needed to confirm or refute them.

EDB SUBJECT CATEGORY: EDB-360202
INIS CATEGORY: B24

<399>

<399>

Mangelmann, M.; Keller, C. (Kernforschungszentrum Karlsruhe
(Germany, F.R.). Inst. fuer Radiochemie)

Thermal decomposition of intermetallic compounds of platinum
with alkaline earth and lanthanide metals at high
temperatures AIX-08:303033;EDF-77:082573; J. Inorg. Nucl.
Chem. (JINCA) 39:(2):305-309 (1977)

SUBJECT DESCRIPTORS: CHEMICAL

PREPARATION;DECOMPOSITION;DYSPROSIUM;DYSPROSIUM ALLOYS:
T2;EUROPIUM;EUROPIUM ALLOYS: T3;FREE
ENTHALPY;GADOLINIUM;GADOLINIUM ALLOYS: T4;HIGH
TEMPERATURE;INTERMETALLIC COMPOUNDS;KNUDSEN FLOW;MELTING
POINTS;OXIDES;PLATINUM ALLOYS: T1;PYROLYSIS;REDUCTION:
Q1,Q2,Q3,Q4,Q5,Q6;SAMARIUM;SAMARIUM ALLOYS: T5;VAPOR
PRESSURE;VERY HIGH TEMPERATURE;YTTERBIUM;YTTERBIUM ALLOYS: T6

ABSTRACT: The thermal decomposition of intermetallic compounds
of platinum with alkaline earth ($A = \text{Mg, Ca, Ba}$) and
lanthanide ($A = \text{Sm, Eu, Gd, Dy, Yb}$) elements obtained by the
coupled reduction method, was investigated by using the
Knudsen effusion technique. It was demonstrated that the
 $\text{APt}_{\text{sub } 2}(\text{APt}_{\text{sub } 3})$ intermetallic compounds are thermally
decomposed to yield $\text{APt}_{\text{sub } 5}(\text{APt}_{\text{sub } 7})$ and $\text{A}_{\text{sub } (g)}$, the
condensation of which may be used to prepare pure metal A.
Furthermore, it was shown that the free molar solution
enthalpy of A in $\text{APt}_{\text{sub } (x)}$: $\Delta H_{\text{sub } 1}^{\text{G}}(\text{A})$ yields
 $\text{APt}_{\text{sub } (x)}$ has smaller absolute values for the alkaline earth
compounds than for the lanthanide compounds, europium,
however, showing a behaviour more characteristic for divalent
metals. Data for the free molar solution enthalpy and the
vapour pressure of A over $\text{APt}_{\text{sub } (x)}$ for temperatures up to
2000°C are given together with the melting points of
the compounds investigated and a critical description for the
procedure to prepare the metal A by hydrogen reduction of the
oxide $\text{AO}_{\text{sub } (y)}$ vs $\text{APt}_{\text{sub } (x)}$ -compounds as intermediates.

EDB SUBJECT CATEGORY: EDB-360104

INIS CATEGORY: E22

<400>

Mar, R.W.; Hastings, J.C., Sandia Labs., Livermore, Calif. (USA)

In Synthesis of UN coatings on uranium **

INS-77:013089;ERA-02:049201;EDB-77:107128; (Apr 1975)
SAND--75-8008

SUBJECT DESCRIPTORS: CHEMICAL REACTIONS: Q1,Q3;CRACKS;FREE
ENERGY;MAGNESIUM;MAGNESIUM NITRIDES: M3;NITRIDATION: Q1;PHASE
STUDIES;PROTECTIVE COATINGS;SYNTHESIS: Q2;TEMPERATURE
DEPENDENCE;THICKNESS;TIME DEPENDENCE;URANIUM: M1;URANIUM

<400> CONT.

NITRIDES: M2;VERY HIGH TEMPERATURE

ABSTRACT: A single-displacement reaction approach was used to form uranium mononitride as a protective coating for uranium; the source of the nitrogen was a solid nitride, and magnesium nitride was chosen as the most attractive candidate for the experiment. While the goal of synthesizing a single-phase uranium mononitride coating at temperatures lower than 1405 K was met, a number of problems inherent in the process were identified. 6 figures, 2 tables.

EDS SUBJECT CATEGORY: EDB-360105;360201;400201

INIS CATEGORY: B15;B23;B12

<401>

Marchant, D.D.; Bowen, H.K. ** Cooper, A.R.; Heuer, A.H. (eds.)

Oxygen redistribution in UO_2 due to a temperature gradient
In Mass transport phenomena in ceramics **

INS-77:009204;ERA-02:034778;EDB-77:074806; ([nd]) Plenum
Publishing Corporation

SUBJECT DESCRIPTORS: ATCM TRANSPORT: Q1; BOUNDARY
CONDITIONS; DIFFUSION: Q2; KINETIC EQUATIONS; MASS
TRANSFER; MIGRATION; NUCLEAR FUELS; OXYGEN:
T2; STOICHIOMETRY; TEMPERATURE EFFECTS; TEMPERATURE
GRADIENTS; THERMAL DIFFUSION; URANIUM DIOXIDE: T1

ABSTRACT: Uranium dioxide exhibits a wide range of stoichiometry, with the deviation from stoichiometry, x , ranging from -0.31 to $+0.24$. The observation of oxygen redistribution in uranium dioxide-based fuels in a nuclear reactor and the consequent effect on operational performance parameters has prompted investigation of the migration of oxygen in a temperature gradient. Laboratory studies consist in annealing the samples in a temperature gradient, quenching to room temperature, and measuring the resulting composition profiles. The results have been interpreted in terms of an effective heat of transport relating concentration changes to the temperature gradient for the particular experimental conditions.

EDS SUBJECT CATEGORY: EDB-360204;050700

INIS CATEGORY: B23

<402>

<402>

Markov, B.F.; Khorlbekk, V.; Lojchenko, V.Ya. (AN Ukrainskoj SSR, Kiev. Inst. Obshchej i Neorganicheskoy Khimii)

Diagram of excess free energy of the $PbCl_2$ - $CsCl$ - $ZnCl_2$ system EDB-76:042587; Ukr. Khim. Zh. (UKZHA) 41:(3):245-248 (Mar 1975)

SUBJECT DESCRIPTORS: CESIUM CHLORIDES: T1; DIAGRAMS; ELECTRIC POTENTIAL; FREE ENERGY: Q1, Q2, Q3; LEAD CHLORIDES: T2; MOLTEN SALTS; ZINC CHLORIDES: T3

ABSTRACT: None

EDE SUBJECT CATEGORY: EDB-400201
INIS CATEGORY: B12

<403>

Markov, E.G.

Speed of ultrasound and the thermophysical properties of the liquid metals Sn, Pb, Cd and of their binary alloys Pb:Sn and Pb:Cd ERA-01:023779; EDB-76:065751; High Temp. (USSR) (Engl. Transl.) (HITEA) 13:(5):1027-1030 (Sep 1975)

SUBJECT DESCRIPTORS: BINARY ALLOY SYSTEMS; CADMIUM: T1; CADMIUM ALLYS: T2; COMPRESSIBILITY: Q1, Q2, Q3, Q4, Q5, Q6; DENSITY; LEAD: T4; LEAD ALLOYS: T6; LIQUID METALS; SPECIFIC HEAT: Q1, Q2, Q3, Q4, Q5, Q6; TEMPERATURE DEPENDENCE; THERMAL EXPANSION: Q1, Q2, Q3, Q4, Q5, Q6; TIN: T3; TIN ALLOYS: T5; ULTRASONIC WAVES: Q1, Q2, Q3, Q4, Q5, Q6

ABSTRACT: The temperature dependence of the speed of sound in the liquid metals Sn, Pb, and Cd, as well as in binary mixtures of them was determined. The compressibility and specific heat of the metals was also studied. (AIP)

EDE SUBJECT CATEGORY: EDB-360104

<404>

Maroni, V.A.; Calaway, W.F.; Veleckis, E.; Yonco, R.M., Argonne National Lab., Ill. (USA)

In Solution behavior of hydrogen isotopes and other non-metallic elements in liquid lithium **
INS-76:018261; ERA-01:026208; EDB-76:075630; (1976)
CCNF-760503--5

SUBJECT DESCRIPTORS: HIGH TEMPERATURE; IMPURITIES: Q4; LITHIUM; LITHIUM: T4; LITHIUM DEUTERIDES: T1; LITHIUM HYDRIDES: T2; LITHIUM NITRIDES: T5; LITHIUM TRITIDES: T3; PHASE

<404> CONT.

DIAGRAMS; REMOVAL; SEPARATION PROCESSES; SOLUBILITY: Q1, Q2, Q3, Q5

ABSTRACT: Results of experimental studies to measure selected thermodynamic properties for systems of lithium with non-metallic elements are reported. Investigations of the Li-H, Li-D, and Li-T systems have led to the elucidation of the dilute solution behavior and the H/D/T isotope effects. In the case of the Li-H and Li-D systems, the principal features of the respective phase diagrams have been delineated. The solubility of Li-D in liquid lithium has been measured down to 200^{sup} 0°C. The solubility of Li₃N in liquid lithium and the thermal decomposition of Li₃N have also been studied. From these data, the free energy of formation of Li₃N and the Sieverts' constant for dissolution of nitrogen in lithium have been determined. Based on studies of the distribution of non-metallic elements between liquid lithium and selected molten salts, it appears that molten salt extraction offers promise as a means of removing these impurity elements (e.g., H, D, T, O, N, C) from liquid lithium.

FDB SUBJECT CATEGORY: EDB-400201; 700209; 060200
INIS CATEGORY: B12; A14; B11

<405>

Maroni, V.A.; Calaway, W.F.; Veleckis, E.; Yonco, R.M.; Cafasso, F.A.; Burris, L., Argonne National Lab., Ill. (USA)

In Physical research on liquid-metal systems. Annual report, July 1975--June 1976 **
ERA-03:00 1146; INS-77:018009; EDB-78:004542; (1976) ANL--76-106

SUBJECT DESCRIPTORS: HIGH TEMPERATURE; ISOTOPE EFFECTS; LIQUID METALS; LITHIUM: T1; LITHIUM DEUTERIDES: T3; LITHIUM HYDRIDES: T2; LITHIUM TRITIDES: T4; PRESSURE DEPENDENCE; SOLUBILITY: Q1, Q2, Q3, Q4; THERMONUCLEAR REACTOR MATERIALS; VERY HIGH TEMPERATURE

ABSTRACT: Physicochemical and thermodynamic studies of liquid-lithium-containing systems have continued. Pressure-composition isotherms for the Li-H system measured by a gravimetric technique are found to be in reasonably good agreement with earlier tensimetric results. Plateau-pressure studies of the Li-H, Li-D, and Li-T systems have been made at temperatures above and below the monotectic. The observed isotope effects were found to be near the predicted values. Measurements of the solubility of LiD in liquid lithium gave results in good agreement with prior data for the solubility of LiH in lithium and confirmed that the limits on cold trapping of lithium hydrides from lithium are too high for projected fusion reactor applications. Determination of the

<405>

<405> CONT.

solubility of Li_2O in liquid lithium is nearing completion. Solubility values measured to date are slightly lower than, but in the range of, prior measurements.

EDB SUBJECT CATEGORY: EDB-360105;700209

INIS CATEGORY: B15;A14

<406>

Martinot, L. (Liege Univ. (Belgium))

Some thermodynamic properties of dilute solutions of actinide chlorides in (Li-K)Cl and in (Na-K)Cl eutectics

AIX-07:236205;EDB-76:092883: J. Inorg. Nucl. Chem. (JINCA) 37: (12):2525-2528 (Dec 1975)

SUBJECT DESCRIPTORS: ELECTROMOTIVE

FORCE;ENTROPY;EUTECTICS;FORMATION HEAT;FREE ENTHALPY;LITHIUM CHLORIDES: M2;NEPTUNIUM CHLORIDES;PLUTONIUM CHLORIDES: M3;POTASSIUM CHLORIDES;REACTION KINETICS;SODIUM CHLORIDES;SOLUTIONS;THERMODYNAMIC PROPERTIES: Q1,Q2,Q3;URANIUM CHLORIDES: M1

ABSTRACT: The activity coefficients at infinite dilution of UCl_3 , UCl_4 , NpCl_3 , NpCl_4 , PuCl_3 in (Li-K)Cl and of UCl_3 and UCl_4 in (Na-K)Cl are reported. The Gibbs free energy and entropy of formation of these compounds in dilute solution are calculated from e.m.f. data. A comparison between the cited thermodynamic quantities and the activation energy for the diffusion of the + 3 and + 4 actinide ions is also presented.

EDB SUBJECT CATEGORY: EDB-400702

INIS CATEGORY: B13

<407>

Martinot, L.; Bohet, J.; Duyckaerts, G. (Liege Univ. (Belgium)); Muller, W.

Thermodynamic properties of dilute solutions of ThCl_4 in (Li-K)Cl and (Na-K) eutectics AIX-08:332016;EDB-78:022423; Inorg. Nucl. Chem. Lett. (INUCA) 13: (7):315-319 (1977)

SUBJECT DESCRIPTORS: DILUTION;ELECTROMOTIVE

FORCE;EUTECTICS;FORMATION FREE ENTHALPY: Q1;LITHIUM CHLORIDES;MOLTEN SALTS;POTASSIUM CHLORIDES;SODIUM CHLORIDES;SOLUTIONS;THORIUM CHLORIDES: T1

ABSTRACT: The Gibbs free energy of formation of ThCl_4 in (Li-K)Cl is reported and a comparison is made with similar data obtained in (Na-K)Cl. The thermodynamic data have been

<407> CONT.

obtained from e.m.f. measurements extrapolated down to infinite dilution. The standard potentials of the Th(IV-O) system at 400, 450 and 500\$sup 0\$C are respectively equal to 2.52, 2.50 and 2.47 volts vs. Cl\$sub 2\$/2Cl\$sup -\$ in the m.f. scale.

EDE SUBJECT CATEGORY: EDB-400702
INIS CATEGORY: B13

<408>

Martinot, L.; Duyckaerts, G. (Liege Univ. (Belgium))

Diffusion of uranium species in several molten chlorides
AIX-08:332017;EDB-78:022424; Inorg. Nucl. Chem. Lett. (INUCA)
13:(7):321-327 (1977)

SUBJECT DESCRIPTORS: CESIUM CHLORIDES: T2;DIFFUSION:
Q1;EUTECTICS;IONS;LITHIUM CHLORIDES: T3;MOLTEN
SALTS;PERMEABILITY: Q2,Q3,Q4;RUBIDIUM CHLORIDES: T4;URANIUM:
T1;URANIUM COMPOUNDS

ABSTRACT: The chronopotentiometric diffusion coefficients of U(III) and U(IV) have been measured in (Li-Cs)Cl and in (Rb-Cs)Cl eutectics. Taking into account previous data it is concluded that the molten salts 'holes' model proposed for the self-diffusion in pure molten salts is also able to interpret qualitatively interdiffusion in eutectic melts.

EDE SUBJECT CATEGORY: EDB-400702
INIS CATEGORY: B13

<409>

Masson, J.P.; Desmoulin, J.P.; Charpin, P.; Bougon, R.

Synthesis and characterization of a new uranium(V) compound:
H\$sub 3\$O\$sup +\$UF\$sub 6\$ \$sup -\$ INS-77:005830;EDB-77:044531;
Inorg. Chem. (INOCA) 15:(10):2529-2531 (Oct 1976)

SUBJECT DESCRIPTORS: CHEMICAL PREPARATION: Q1,Q2;COMPARATIVE
EVALUATIONS;CUBIC LATTICES;ELECTRON SPIN RESONANCE;EMISSION
SPECTRA;FORMATION HEAT;INFRARED SPECTRA;OXONIUM IONS:
T;PHYSICAL PROPERTIES: Q1,Q2;RAMAN SPECTRA;URANIUM COMPOUNDS:
T2;URANIUM FLUORIDES: T1;X-RAY DIFFRACTION

ABSTRACT: The reaction of equimolar amounts of UF\$sub 5\$ and H\$sub 2\$O in hydrogen fluoride results in the partial dissolution of UF\$sub 5\$, yielding a blue-green solution from which the new salt oxonium hexafluorouranate (V) (H\$sub 3\$O\$sup +\$UF\$sub 6\$ \$sup -\$) could be isolated as a green crystalline solid. Calorimetric measurements showed H\$sub 3\$O\$sup

<409>

<409> CONT.

+ UF_6 - F to decompose at about 68°C and its heat of formation to be equal to -628 ± 2 kcal/mol. Its ionic nature in the solid state and in HF solutions was demonstrated from vibrational and electronic spectra. The electronic spectrum is closely similar to those of LiUF_6 , NaUF_6 , and CsUF_6 and differs from those of RbUF_6 and KUF_6 . This adduct shows a strong ESR signal, with $g = -0.78 \pm 0.10$, characteristic of UF_6 salts. Based on its x-ray powder diffraction pattern, $\text{Hf}_3\text{O}_7 \cdot \text{UF}_6$ is cubic with $a = 5.2229 \pm 0.0005$ Å.

EDB SUBJECT CATEGORY: EDB-400702

INIS CATEGORY: B13

<410>

Matthews, G.E. Jr.; Crawford, J.H. Jr.

Ionic-thermoelectric study of the dipole relaxation and equilibrium in Gd-doped SrF_2

INS-77:003555; ERA-02:017832; EDB-77:026274; Phys. Rev., B (PLRBA) 15:(1):55-60 (1 Jan 1977)

SUBJECT DESCRIPTORS: DIPOLES; Q1; ENTHALPY; GADOLINIUM

ICNS; INTERSTITIALS; LCF

TEMPERATURE; POLARIZATION; RELAXATION; STRONTIUM FLUORIDES;

T1; TEMPERATURE DEPENDENCE

ABSTRACT: By means of thermal depolarization (ionic thermoelectric) measurements, the population of nearest-neighbor (type-I) and next-nearest-neighbor (type-II) Gd^{3+} interstitial fluoride ion dipoles have been observed as functions of temperature from 180 to 210 K, the region of the type-II dipole relaxation peak. The ratio of type-II to type-I concentration varies exponentially with reciprocal polarization temperature, indicating that at each temperature the two dipole species reach equilibrium within the polarization period, chosen to be at least six times the type-II dipole relaxation time. The enthalpy difference between the two species was determined to be 0.046 eV. The shift in the populations with temperature suggests that reorientation of the type-II dipole proceeds by means of the fluoride passing through a nearest-neighbor (type-I) site, in accordance with the accepted model. By assuming that the total number of dipoles remains constant, the ratio of the dipole moments of the two species is found to be 2.4. (AIP)

EDB SUBJECT CATEGORY: EDB-360603

INIS CATEGORY: B24

<411>

McClelland, J.F., Ames Lab., Iowa (USA)

In Electromodulation spectroscopy of sc and fcc phase TlCl and TlBr ** ERA-01:026159;EDB-76:075467; (Jun 1976) IS-T--721

SUBJECT DESCRIPTORS: BAND THEORY;CARRIERS;CRYSTAL LATTICES;ELECTRONIC STRUCTURE: Q1,Q2;EXCITED STATES;EXCITONS;FCC LATTICES;MODULATION;THALLIUM BROMIDES: T2;THALLIUM CHLORIDES: T1

ABSTRACT: Electromodulation measurements were made on these compounds and the spectra were reduced to the electric field induced changes in the dielectric function. The results indicate the importance of photocarrier effects in both theory and experiment in the electromodulation of exciton states. In the future, calculations should include the effect of photocarriers on the field seen by the exciton and experimentally samples should be developed with known and reproducible photocarrier properties with temperature control between liquid helium and nitrogen temperatures and bipolar modulation fields. The abnormal (fcc) phase electrobabsorption (EA) measurements have demonstrated the usefulness of the modulation method in resolving exciton states by determining the $n = 2$ energy in TlBr. This has enabled a number of quantities to be calculated from the Wannier exciton model. The resolution of the $n = 2$ energy in TlCl is probably also possible with an EA measurement and patience with the signal to noise problem. The α and β features are still unassigned but the unusual EA lineshape and sample preparation sensitivity found in this investigation may prove useful in making definitive assignments in conjunction with future work.

EDE SUBJECT CATEGORY: EDB-360603

<412>

McClure, D.S. (Princeton Univ., N.J. (USA). Dept. of Chemistry)

Progress in the spectroscopy of rare earth and transition metal ions AIX-07:253199;ERA-02:023428;EDB-77:096450; J. Lumin. (JLUMA) 12-13::67-78 (Mar 1976)

SUBJECT DESCRIPTORS: BIBLIOGRAPHIES;ELECTRON TRANSFER;ELECTRONIC STRUCTURE;ENERGY SPECTRA;ION SPECTROSCOPY;IONS;LIGANDS;LUMINESCENCE: Q1,Q2;MOLECULES;RARE EARTHS: T1;REVIEWS;TRANSITION ELEMENTS: T2

ABSTRACT: Several areas in which there has been progress in spectroscopy of transition metal and rare earth ions of relevance to luminescence are reviewed. These are, structure of impurity centers in crystals, radiationless processes,

<412>

<412> CONT.

molecular spectra, charge transfer spectra and high energy spectra. The aim of the reviews is not to be comprehensive, but is to note unusual things and to suggest directions where further progress would be most valuable.

EDE SUBJECT CATEGORY: EDB-640302

INIS CATEGORY: A12

<413>

McDiarmid, R.

Assignments in the electronic spectrum of $\text{UF}_{6\text{g}}$

INS-76:016568; EDB-76:061836; J. Chem. Phys. (JCP3A)

65: (1):168-173 (1 Jul 1976)

SUBJECT DESCRIPTORS: ABSORPTION SPECTRA; ELECTRONIC
STRUCTURE; ENERGY-LEVEL TRANSITIONS; ULTRAVIOLET SPECTRA:
Q1; URANIUM HEXAFLUORIDE: T1

ABSTRACT: The absorption spectrum of gaseous $\text{UF}_{6\text{g}}$ was recorded from 1200--4200 Å (83 300--23 800 cm^{-1}). Eight electronic transitions were observed with intensities ranging from 0--20 000 $1/\text{mole}\cdot\text{cm}$. Based on correlations between this spectrum and those of $\text{UF}_{6\text{g}}$, $\text{UCl}_{6\text{g}}$, $\text{MoF}_{6\text{g}}$, and $\text{WF}_{6\text{g}}$ and the photoelectron spectrum of $\text{UF}_{6\text{g}}$, the experimental spectrum was assigned. (AIP)

EDE SUBJECT CATEGORY: EDB-640302

INIS CATEGORY: A12

<414>

McElroy, D.L. (Oak Ridge National Lab., Tenn. (USA)); Haines, H.R.; Hall, R.O.A.; Lee, J.A. (UKAEA Research Group, Harwell. Atomic Energy Research Establishment) ** Blank, H.; Lindner, R. (eds.)

Specific heat of $\text{PuC}_{(1-x)}$ and $\text{Pu}_{2\text{C}}\text{Pu}_{3\text{C}}$ from 10 to 300 K in Plutonium 1975 and other actinides **

AIX-08:310020; ERA-02:041841; EDB-77:094419; (1976)

North-Holland

SUBJECT DESCRIPTORS: CHEMICAL COMPOSITION; DEBYE
TEMPERATURE; ENTROPY; LAMBDA POINT; LATTICE PARAMETERS; LOW
TEMPERATURE; MEDIUM TEMPERATURE; ORDER PARAMETERS; PLUTONIUM
CARBIDES: T1; SCHOTTKY EFFECT; SPECIFIC HEAT: Q1; TEMPERATURE
DEPENDENCE; ULTRALOW TEMPERATURE; VERY LOW TEMPERATURE

ABSTRACT: The specific heat, $C_{\text{sub}}(p)$, of $\text{PuC}_{(0.82)}$, $\text{PuC}_{(0.90)}$, and $\text{PuC}_{(1.51)}$ was determined from 10 to 300K using adiabatic technique. Characterization of the arc-melted

<414> CONT.

and heat-treated samples included chemical, X-ray and metallographic analyses. The $C_{sub}(p)$ -T values for $Pu_{sub}(0.82)$ and $Pu_{sub}(0.90)$ were accurate to $\pm 1.3\%$ and conclusively reveal anomalous $C_{sub}(p)$ behaviour below 150K where neutron diffraction has shown magnetic ordering to occur before rising in a normal manner to values of 46.06 J.mol⁻¹K⁻¹ ($Pu_{sub}(0.82)$), which is 0.3% below the one literature value for $Pu_{sub}(0.81)$ and 47.92 ($Pu_{sub}(0.90)$) at 298.15K. The normal behaviour above 150K was fitted to better than 1% for both $Pu_{sub}(0.82)$, and $Pu_{sub}(0.90)$, by a simple Debye model, with a Debye temperature of 400K and electronic coefficients of 0.015 J.mol⁻¹K⁻² and 0.135 J.mol⁻¹K⁻², respectively. The difference between the calculated and measured curves below 150K show a basic Schottky transition for the $Pu_{sub}(0.82)$ with a superimposed lambda-type contribution in the case of $Pu_{sub}(0.90)$: i.e. the transition initiates as a nonco-operative excitation, becoming co-operative in its later stages.

EDE SUBJECT CATEGORY: EDB-360204

INIS CATEGORY: B24

<415>

McElroy, D.L.; Haines, H.R.; Hall, R.O.A.; Lee, J.A. **
Commission of the European Communities, Karlsruhe (Germany,
F.R.). European Inst. for Transuranium Elements

Specific heat of $Pu_{sub}(1-x)$ and $Pu_{sub} 25C_{sub} 3f$ from 10 to
300 K In 5th international conference on plutonium and other
actinides ** AIX-07:233144; EDB-76:087311; (1975)
AEE-Conf--75-417-020

SUBJECT DESCRIPTORS: DEBYE TEMPERATURE; HIGH
TEMPERATURE; MATHEMATICAL MODELS; MEDIUM TEMPERATURE; PLUTONIUM
CARBIDES: T1; SPECIFIC HEAT: Q1; TEMPERATURE DEPENDENCE

ABSTRACT: None

EDE SUBJECT CATEGORY: EDB-360204

INIS CATEGORY: B23

<416>

<416>

McElroy, W.N.; Ulseth, J.A.; Lippincott, E.P.; Matsumoto, W.Y.;
Kellogg, L.S., Hanford Engineering Development Lab.,
Richland, Wash. (USA)

In Fission rates, burnup, and neutron flux-fluence-spectra
characterization for mixed oxide fuel experiments in the
EBR-II ** FRA-01:013010;EDB-76:052407; (Jan 1976)
HEDL-TME--75-74

SUBJECT DESCRIPTORS: BURNUP;EBR-2 REACTOR: T1;FPTF REACTOR;FUEL
PINS;IRRADIATION;LMFBR TYPE REACTORS;NEUTRON FLUENCE;NEUTRON
FLUX: Q1;NEUTRON SPECTRA;PERFORMANCE TESTING;PLUTONIUM
DIOXIDE;URANIUM DIOXIDE

ABSTRACT: Analysis of a mixed oxide fuel experiment (PNL-11) has
been completed and results for fission rates, burnup, neutron
flux, fluence, anti-E and S.S. damage correlation parameters
have been derived from dosimetry measurements. The results
demonstrated successful dosimetry characterization at the +-
5 percent (1sigma) uncertainty level or better for fission
rates, burnup, and fluence, and at the +- 10 percent (1sigma)
level or better for flux-spectra values. Values of burnup
based on activation dosimeters (^{140}Ba) were in good
agreement with ^{148}Nd measurements made on neighboring
fuel pins using recommended ENDF/B-IV fission yield values.
Good agreement, to within 3 percent, between the reactor
midplane multiple foil derived value of total flux and an ANL
estimate was obtained. Based on this study, current
uncertainty estimates are set at +- 5 percent (1sigma) for
fission rate, burnup, and fluence measurements for short term
controlled "test sample" irradiations of mixed oxide fuel
in EBR-II. It is concluded, however, that additional tests
and standardizations are required to reduce current
uncertainties of 5 to 10 percent to the 2 to 5 percent level.

EDB SUBJECT CATEGORY: EDB-210500
INIS CATEGORY: E35

<417>

McLachlan, A.D., La Trobe Univ., Bundoora (Australia)

In Photoelectron spectroscopic studies of some transition metals
and alloys ** AIX-07:256684;EDB-77:013002; (1974) La Trobe
University

SUBJECT DESCRIPTORS: CADMIUM;COPPER: T1;ELECTRONIC
STRUCTURE;GOLD: T3;GOLD ALLOYS: T6;PALLADIUM ALLOYS:
T5;PHOTOELECTRON SPECTROSCOPY: Q1,Q2,Q3,Q4,Q5,Q6;SAMPLE
PREPARATION;SILVER: T2;SILVER BASE ALLOYS: T4;ZINC

ABSTRACT: Photoelectron spectra of polycrystalline samples of

<417> CONT.

Cu, Ag and Au at photon energies of 21.22, 40.81 eV and 1487 eV were measured. The corrected 40.81 eV results were compared to theoretical band structure calculations and monochromatized x-ray photoelectron results. Correlation of hitherto unresolved peaks in the 40.81 eV spectra was observed. Comparison of the relative intensities of the spectral d bands and the theoretical calculations revealed discrepancies which were assigned to matrix element modulation effects in the photoelectron emission process. Experimental measurements and theories of the electronic structure of disordered alloy systems were reviewed. The 21.22 eV and 40.81 eV photoelectron spectra of some AgPd and AgAu alloys were measured. The spectra were compared with previous x-ray photon results, and with theoretical calculations based on the Coherent Potential Approximation (CPA) model of disordered alloy systems. The present results were found to give more clearly defined spectral details, with differences in the comparison reflecting the simplifying assumptions of the CPA calculation.

EDB SUBJECT CATEGORY: EDB-360102;400104
INIS CATEGORY: A13

<418>

McQuillan, A.E. (Birmingham Univ. (UK). Dept. of Physical Metallurgy and Science of Materials)

Comparative study of systems of heavier rare-earth metals with hydrogen [Gd, Tb, Dy, Lu] AIX-07:278251;EDB-77:031167; J. Less-Common Met. (JCOMA) 49:(1-2):431-438 (Sep 1976)

SUBJECT DESCRIPTORS: CHEMICAL COMPOSITION:
Q1,Q2,Q3,Q4;COMPARATIVE EVALUATIONS;DYSPROSIUM;DYSPROSIUM
HYDRIDES: T3;GADOLINIUM HYDRIDES: T1;HYDROGEN
ADDITIONS;LUTETIUM HYDRIDES: T4;SOLID
SOLUTIONS;SOLUBILITY;SOLUTION HEAT;TEMPERATURE
DEPENDENCE;TERBIUM HYDRIDES: T2

ABSTRACT: Hydrogen equilibrium pressures as a function of hydrogen content of the metal and temperature have been measured for gadolinium, terbium, dysprosium and lutetium, in the primary solid-solution range up to and including the two-phase region with the dihydride phase. All systems show strong similarities, and all are marked by a decrease in heat of solution of hydrogen in the metal with decreasing hydrogen concentration.

EDB SUBJECT CATEGORY: EDB-360603;080800
INIS CATEGORY: B22

<419>

<419>

Mehlhorn, B.; Hoppe, R. (Giessen Univ. (F.R. Germany). Inst. fuer Anorganische und Analytische Chemie)

New hexafluorozirconates(IV): BaZrF_6 , PbZrF_6 , EuZrF_6 , SrZrF_6 AIX-08:289289; EDB-77:069132; Z. Anorg. Allg. Chem. (ZAACA) 425:(2):180-188 (Sep 1976)

SUBJECT DESCRIPTORS: BARIUM FLUORIDES: T1; CHEMICAL PREPARATION; COORDINATION NUMBER; EUROPIUM FLUORIDES: T2; INTERATOMIC DISTANCES; LATTICE PARAMETERS; LEAD FLUORIDES: T3; MONOCRYSTALS; ORTHORHOMBIC LATTICES; SPACE GROUPS; STRONTIUM FLUORIDES: T4; STRUCTURAL CHEMICAL ANALYSIS: Q1, Q2, Q3, Q4, Q5; X-RAY DIFFRACTION; ZIRCONIUM FLUORIDES: T5

ABSTRACT: The new compounds BaZrF_6 , PbZrF_6 , EuZrF_6 and SrZrF_6 (all colorless) crystallize according to single crystal investigations orthorhombic. From the determination of crystal structure of BaZrF_6 results, that similar to K_2ZrF_6 , there are $\text{sub}(\infty)\text{sup}(1)[\text{ZrF}_6(4/2)\text{F}(4/1)]$ -chains. Zr^{4+} is therefore surrounded by eight fluorine atoms. The Madelung part of lattice energy as well as the effective coordination numbers are calculated and discussed.

EDB SUBJECT CATEGORY: EDB-400201
INIS CATEGORY: B12

<420>

Mehrotra, G.M.; Froberg, M.G.; Kapoor, M.L. (Technische Univ. Berlin (F.R. Germany). Inst. fuer Metallforschung)

Activities of lead and bismuth in their molten alloys
EDE-76:081857; Z. Metallkd. (ZEMTA) 67:(3):186-188 (Mar 1976)

SUBJECT DESCRIPTORS: BISMUTH: T2; BISMUTH ALLOYS; BISMUTH OXIDES; CHEMICAL COMPOSITION; ELECTROMOTIVE FORCE; EQUATIONS; EQUILIBRIUM; LEAD: T1; LEAD ALLOYS; LEAD OXIDES; LIQUID METALS; METAL-METAL OXIDE BATTERIES; SLAGS; THERMODYNAMIC PROPERTIES: Q1, Q2; VERY HIGH TEMPERATURE

ABSTRACT: From distribution equilibria and emf-measurements of cells of scheme $[\text{Pb-Bi}]/(\text{PbO-Bi}_2\text{O}_3)/\text{ZrO}_2(+\text{CaO})/\text{FeO, Fe}$, the activities of lead and bismuth in their molten alloys at 900°C have been determined. The results of the present investigation agree well with the

<421> CONT.

known data. The method presented here can be used to determine and to simultaneously obtain the activities in the alloy phase in equilibrium with the oxide melt.

EDB SUBJECT CATEGORY: EDB-400400;360104

<421>

Merrill, L.

Behavior of the AB-type compounds at high pressures and high temperatures INS-78:002256;EDB-78:027693; J. Phys. Chem. Ref. Data (JPCRB) 6:(4):1205-1252 (Oct 1977)

SUBJECT DESCRIPTORS: CRYSTAL STRUCTURE;CRYSTAL-PHASE TRANSFORMATIONS: Q1,Q2;MELTING: Q1,Q2;PHASE DIAGRAMS;VERY HIGH PRESSURE;X-RAY DIFFRACTION

ABSTRACT: The results of the published work on the high pressure-high temperature properties of the AB-type compounds have been compiled and evaluated. All pressure studies above the range of 1 kilobar have been included with an emphasis on the accurate characterization of the solid-solid phase boundaries and the experimental melting curves. Whenever x-ray diffraction data are available for the high pressure phases, they have also been reviewed. Phase diagrams are included for all compounds in which measurement of more than one point along the phase boundary was made. This review discusses a total of 87 compounds and 212 distinct high pressure-high temperature phases.

EDB SUBJECT CATEGORY: EDB-360202
INIS CATEGORY: B23

<422>

Meyer, A.; Stott, M.J.; Young, W.H. (Queen's Univ., Kingston, Ontario (Canada). Dept. of Physics)

Densities of electronic states in liquid transition metals
AIX-07:245437;EDB-76:071328; Philos. Mag. (PHMAA)
33:(2):381-385 (Feb 1976)

SUBJECT DESCRIPTORS: CHROMIUM;COBALT;CRYSTAL STRUCTURE;DENSITY;ENERGY-LEVEL DENSITY: Q1;ENTROPY;EQUATIONS OF STATE;FERMI LEVEL;IRON;LIQUID METALS;MANGANESE;NICKEL;PALLADIUM;PLATINUM;TITANIUM;TRANSITION ELEMENTS: T1;VANADIUM;ZIRCONIUM

ABSTRACT: Measured structure factors and entropies are used to obtain, very directly, the electronic contributions to be

<422>

<422> CONT.

entropies and hence the densities of states at the Fermi levels in some liquid transition metals.

EDB SUBJECT CATEGORY: EDB-360104

INIS CATEGORY: A13

<423>

Meyer, A.S. ** Lyon, W.S. (comp.), Oak Ridge National Lab., Tenn. (USA)

Energy programs: molten-salt reactor program In Analytical Chemistry Division annual progress report for period ending November 30, 1975 **

ERA-01:013775; INS-76:014730; EDB-76:049027; (Feb 1976)

CRNL--5100

SUBJECT DESCRIPTORS: CATIONS; CHEMICAL ANALYSIS: Q1; CHEMICAL REACTIONS; MOLTEN SALTS: T1; VALENCE

ABSTRACT: Progress is reported in studies on electrochemical in-line measurements in fuel systems. Results of studies of the electrochemical behavior of Fe, and Te in fuel melts are given. At the Coolant-Salt Technology Facility (CSTF), first measurements were made on the tritium distribution within the system. Spectrophotometric methods were used to identify and characterize various oxidation states and compounds of actinide ions. (JGB)

EDB SUBJECT CATEGORY: EDB-400104; 400400

INIS CATEGORY: B11

<424>

Michaille, P.; Kikindai, T. (Ecole Centrale des Arts et Manufactures, 92 - Chatenay-Malabry (France))

Effect of fluoride ions on the kinetics of the oxidation of uranium(IV) by cerium(IV) AIX-08:312240; EDB-77:095034; J. Inorg. Nucl. Chem. (JINCA) 39:(3):493-497 (1977)

SUBJECT DESCRIPTORS: ANIONS; CERIUM COMPLEXES; CERIUM FLUORIDES; CERIUM HYDROXIDES; CHEMICAL REACTION KINETICS; OXIDATION: Q1; SPECTROPHOTOMETRY; URANIUM: T1; URANIUM COMPOUNDS

ABSTRACT: The equilibrium constants of the two fluoro-complexes of the cerium(IV) ion were calculated from spectrophotometric measurements; at $[HClO_4] = 0.5\text{ M}$, $I = 2.0\text{ M}$ (adjusted by $NaClO_4$) and $t = 20^\circ\text{C}$, the values $K_1 = \frac{[CeOHf^{2+}][H^{+}]}{[CeOHf^{3+}][HF]} = 41000 + 2000$ and $K_2 = \frac{[CeF^{2+}]}{[CeOHf^{2+}][HF]} = 3120$

<424> CONT.

+ 50 M⁻¹ were found. These values were used to interpret the effect of the fluoride ions on the kinetics of the Ce(IV)-U(IV) reaction studied in the same conditions with a stopped-flow spectrophotometer: a maximum of the rate constant is obtained when two fluoride ions are involved, but the reaction cannot proceed through reaction paths involving more fluoride ions; the reaction paths that involve one or two fluoride ions also involve one hydroxyl ion; the rate law that takes into account the predominant reactive species may be written: $-d[\text{Ce}(\text{IV})]/dt = v_{\text{0}}[\text{CeOH}^{3+}][\text{U}^{4+}] + v_{\text{I}}[\text{CeOHF}^{2+}][\text{U}^{4+}] + v_{\text{II}}[\text{CeOHF}^{2+}][\text{UF}^{3+}]$, where $v_{\text{0}} = 14600 \text{ M}^{-1}\text{sec}^{-1}$, $v_{\text{I}} = 38670 \text{ M}^{-1}\text{sec}^{-1}$ and $v_{\text{II}} = 70570 \text{ M}^{-1}\text{sec}^{-1}$. A bridge mechanism is discussed from the comparison of these results with those obtained at low sulfate concentrations.

EDB SUBJECT CATEGORY: EDB-400702

INIS CATEGORY: B13

<425>

Miedema, A.R. (Philips Research Labs., Eindhoven, Netherlands); de Boer, F.R.; Boom, R. ** Lundin, C.E. (ed.)

Heat of formation of solid and liquid alloys of transition metals In Proceedings of the 12th rare earth research conference. Vol. II **
ERA-02:060317; INS-77:017495; EDB-77:140388; (1976) Univ. of Denver

SUBJECT DESCRIPTORS: CRYSTAL STRUCTURE: Q1; FORMATION HEAT: Q1; LIQUID METALS; MATHEMATICAL MODELS; TRANSITION ELEMENT ALLOYS: T1

ABSTRACT: Energy effects in alloys of transition metals are shown to be predominantly nearest-neighbor effects whereas both crystal structure and entropy effects are relatively unimportant. The model description predicts the sign of ΔH (is the interaction repulsive or attractive) for binary systems with 99% accuracy. Progress is made toward the prediction of numerical values for both heats of formation of solid compounds and heats of mixing of liquid alloys. (GHT)

EDB SUBJECT CATEGORY: EDB-360102

INIS CATEGORY: B22

<426>

<426>

Mikhailichenko, A.I.; Abramov, L.A.; Dozhdeva, N.M.

Tetrad effect in the extraction of the lanthanides with neutral oxygen-containing compounds EDB-76:066299; Sov. Radiochem. (Engl. Transl.) (SVRDA) 17:(3):349-351 (1975)

SUBJECT DESCRIPTORS: ATOMIC NUMBER;CHEMICAL PROPERTIES:
Q1;CHLORIDES;DISTRIBUTION FUNCTIONS: Q1;NITRATES;PETROLEUM
PRODUCTS;RARE EARTHS: M1;SOLVENT EXTRACTION:
Q1;SULFOXIDES;TRP;THIOCYANATES

ABSTRACT: The distribution of the lanthanides in the extraction of their nitrates, chlorides, and thiocyanates by neutral oxygen-containing extraction reagents (in 12 systems) was studied. In all the systems on the curves of the distribution coefficient versus atomic number of the element, four subgroups of the elements (four tetrads) appear: La, Ce, Pr, Nd; Pm, Sm, Eu, Gd; Gd, Tb, Dy, Ho; and Er, Tm, Yb, Lu. Between Nd and Pm there is a junction of the first and second tetrads, and between Ho and Er that of the third and fourth. The second and third tetrads are in contact on Gd, and moreover, Gd is the last element of the second tetrad and simultaneously the first element of the third tetrad.

EDE SUBJECT CATEGORY: EDB-400201;400105

<427>

Mikhajlov, M.A.; Ehpov, D.G.; Shchetinina, G.P.; Barinova, M.V.
(AN SSSR, Vladivostok. Inst. Khimii)

Thermogravimetric investigation of interaction of some rare earth oxides with ammonium bifluoride EDB-76:036914; Izv. Sib. Otd. Akad. Nauk SSSR, Ser. Khim. Nauk (IZSKA) (9):114-119 (Jul 1974)

SUBJECT DESCRIPTORS: AMMONIUM COMPOUNDS: T6;CHEMICAL REACTION
KINETICS: Q1,Q2,Q3,Q4,Q5,Q6;CHEMICAL REACTIONS;DYSPROSIUM
OXIDES: T5;FLUORIDES;FORMATION HEAT;GADOLINIUM OXIDES:
T4;INFRARED SPECTRA;NEODYMIUM OXIDES: T2;OXIDES;PRASEODYMIUM
OXIDES: T1;RARE EARTH COMPOUNDS;SAMARIUM OXIDES: T3;THERMAL
GRAVIMETRIC ANALYSIS;X-RAY DIFFRACTION

ABSTRACT: None

EDE SUBJECT CATEGORY: EDB-400201
INIS CATEGORY: B12

<428>

Mintz, M.H.; Hadari, Z. (Israel Atomic Energy Commission, Beersheba. Nuclear Research Center-Negev); Bixon, M. (Tel Aviv Univ. (Israel). Dept. of Chemistry)

Thermodynamic study of the neptunium-hydrogen system
AIX-08:280946;EDB-77:037068; J. Less-Common Met. (JCOMA)
48: (1):183-186 (Jul 1976)

SUBJECT DESCRIPTORS: FORMATION FREE ENERGY;FORMATION
HEAT;HYDROGEN;NEPTUNIUM;NEPTUNIUM HYDRIDES: T1;PHASE STUDIES:
Q1

ABSTRACT: The neptunium-hydrogen system has been investigated previously by Mulford and Wiewandt (1965) who found two hydride phases. One phase, NpH_{2+x} ($0 < x < 0.7$) has a fcc crystal structure and the other phase, NpH_{3-x} , has a hcp structure. An anomalous behavior of the cubic NpH_{2+x} phase was found at the lower composition limit as the hydrogen content increases with increasing temperature. Another anomalous trend may be shown to exist in the thermodynamic values calculated from the pressure-composition isotherms. When partial molal enthalpies and entropies are calculated for the homogeneity range of the cubic, non-stoichiometric NpH_{2+x} (by plotting $\log P$ vs. $1/T$, for a fixed value of x), it is found that ΔH values are independent of hydrogen composition (x), whereas ΔS decreases (becomes more negative) with increasing x . Both results contradict conventional statistical-thermodynamic derivations which are applied to metal-hydrogen systems. The present note reinvestigates the neptunium-hydrogen system and establishes whether such anomalous thermodynamic behavior really exists.

EDE SUBJECT CATEGORY: EDB-360603
INIS CATEGORY: B22

<429>

Montgomery, R.L., Bureau of Mines, Reno, Nev. (USA)

In Electronegativities of the rare-earth elements **
EDB-77:120756; (May 1959) BM-RI--5567

SUBJECT DESCRIPTORS: ELECTRONEGATIVITY: Q1;FORMATION
HEAT;HALIDES;HYDRIDES;NITRIDES;OXIDES;RARE EARTHS:
T1;SULFATES;SULFIDES

ABSTRACT: None

EDE SUBJECT CATEGORY: EDB-400201

<430>

<430>

Morachevskii, A.G.; Demidov, A.I.; Gerasimenko, L.N.; Bikina, G.V.

In Current yield during separation of lithium on a liquid bismuth cathode ** ERA-02:001179; EDB-76:081853; Izv. Vyssh. Uchebn. Zaved., Tsvet. Metall. (IVUTA) 17:(3):94-95 (1974) UCRL-Trans--11105

SUBJECT DESCRIPTORS: CURRENT DENSITY; EFFICIENCY: Q2; ELECTRODEPOSITION: Q1; ELECTROLYTIC CELLS: T2; EUTECTICS; LITHIUM: T1; LITHIUM CHLORIDES; LITHIUM FLUORIDES; MOLTEN SALTS

ABSTRACT: The current efficiency in a fused eutectic LiCl--LiF electrolytic cell was studied as a function of current density, volumetric concentration of Li, and surface concentration of Li. The curve for the current density dependence passes through a maximum and therefore at high current densities the current yield decreases. For both the volumetric and surface Li concentrations there is first a slow decrease in current yield followed by a sharp decrease. (JSR)

EDB SUBJECT CATEGORY: EDB-400400; 400105

<431>

Moravec, J.; Sara, V.; Sourkova, L.

Quantitative determination of UO_2 , and ThO_2 in powdery mixtures by infrared spectrometry EDB-76:036856; J. Radicanal. Chem. (JRACB) 28:(1-2):119-123 (1975)

SUBJECT DESCRIPTORS: ABSORPTION SPECTROSCOPY: Q1, Q2; COMPUTER CODES; INFRARED SPECTRA; NUCLEAR FUELS; PLUTONIUM DIOXIDE; POWDERS; QUANTITATIVE CHEMICAL ANALYSIS; THORIUM OXIDES: T1; URANIUM DIOXIDE: T2

ABSTRACT: Infrared spectra of powdery mixtures of UO_2 and ThO_2 were quantitatively evaluated with a computer. The computer program was tested using the spectra of sulfates. Preliminary results were given and a possible use of this method for analyzing the mixtures of UO_2 with PuO_2 was suggested. (auth)

EDB SUBJECT CATEGORY: EDB-400104

<432>

Moret, J.M.; Lacroix, R. (Geneva Univ. (Switzerland))

Effect of temperature on the fine structure of Eu^{2+} in KCl and KI AIX-07:276246; EDB-77:027679; Helv. Phys. Acta (HFACA) 49: (3):313-323 (18 Jun 1976)

SUBJECT DESCRIPTORS: DOPED MATERIALS; ELECTRON SPIN RESONANCE; ENERGY-LEVEL TRANSITIONS; EUROPIUM IONS: T1; FINE STRUCTURE: Q1, Q2, Q3; LOW TEMPERATURE; MEDIUM TEMPERATURE; ORTHORHOMBIC LATTICES; POTASSIUM CHLORIDES: T2; POTASSIUM IODIDES: T3; TEMPERATURE DEPENDENCE; ULTRALOW TEMPERATURE; VERY LOW TEMPERATURE

ABSTRACT: The effects of temperature variation (between 5 and 296 $^{\circ}$ K) and isotropic pressure variation (between 1 and 75 kgf/mm 2) on the fine structure parameters $b_{\text{sub}}(k)_{\text{sup}}(q)$ of Eu^{2+} in KCl and KI has been studied by electron paramagnetic resonance. The symmetry of the centre is orthorhombic (C_{2v}) and only variations of b_{sub}^2 , b_{sub}^2 and b_{sub}^4 are important. A microscopic description of $b_{\text{sub}}(k)_{\text{sup}}(q)$ is proposed, the temperature dependence is attributed partly to lattice dilatation and partly to two local modes which are revealed by vibronic structure of optical (f-d) transitions of Eu^{2+} .

EDE SUBJECT CATEGORY: EDB-640302

INIS CATEGORY: A12

<433>

Morozov, A.I. (AN SSSR, Moscow. Inst. Obshchej i Neorganicheskoy Khimii)

Pressure and composition of vapors over alkali metal tetrachloroferrates in chlorine atmosphere EDB-76:036733; Zh. Neorg. Khim. (ZNOKA) 20: (4):955-960 (Apr 1975)

SUBJECT DESCRIPTORS: CESIUM CHLORIDES: T1; CHEMICAL COMPOSITION; CHLORINE; IRON CHLORIDES: T2; MOLTEN SALTS; TEMPERATURE DEPENDENCE; THERMODYNAMIC PROPERTIES: Q1, Q2; VAPOR PRESSURE; VAPORS

ABSTRACT: None

EDE SUBJECT CATEGORY: EDB-360603

INIS CATEGORY: B12

<434>

<434>

Mcrozov, A.I.; Kanashvili, N.V.; Mcrozov, I.S.; Gaprindashvili, V.N. (AN SSSR, Moscow. Inst. Obshchej i Neorganicheskoi Khimii)

$\text{NdCl}_3\text{-AlCl}_3\text{-NaCl}$ system AIX-08:295345; EDB-77:056225; Zh. Neorg. Khim. (ZNOKA) 21:(7):1909-1912 (Jul 1976)

SUBJECT DESCRIPTORS: ALUMINIUM CHLORIDES:

T1;CRYSTALLIZATION;HIGH TEMPERATURE;NEODYMIUM CHLORIDES:

T2;PHASE DIAGRAMS: Q1,Q2,Q3;QUANTITY RATIO;SODIUM CHLORIDES:

T3;TEMPERATURE DEPENDENCE

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-400201

INIS CATEGORY: B12

<435>

Mcrozov, A.I.; Maksyukova, E.V. (AN SSSR, Moscow. Inst. Obshchei i Neorganicheskoi Khimii)

$\text{FeCl}_3\text{-MeCl}$ systems AIX-07:265084; EDB-77:013486; Zh. Neorg. Khim. (ZNOKA) 20:(7):2001-2003 (Jul 1975)

SUBJECT DESCRIPTORS: BINARY MIXTURES;CESIUM CHLORIDES: T1;IRON

CHLORIDES: T2;LITHIUM CHLORIDES: T3;MELTING POINTS;MOLTEN

SALTS;PHASE DIAGRAMS: Q1,Q2,Q3,Q4;RUBIDIUM CHLORIDES:

T4;THERMAL ANALYSIS

ABSTRACT: Phase diagrams of $\text{FeCl}_3\text{-MeCl}$ (Me-Li, Rb, Cs) systems was studied by thermal analysis. Investigated systems belong to the congruence melted type. Compound LiFeCl_4 is melted at 163°C, RbFeCl_4 -330°C, CsFeCl_4 -383°C.

EDB SUBJECT CATEGORY: EDB-400201

INIS CATEGORY: B12

<436>

Moser, Z.; Zakulski, W.

Interaction parameters in the Zn--Pb--Sn system at low zinc concentration ERA-01:016577; EDB-76:042162; J. Electrochem. Soc. (JESOA) 122:(5):691-698 (May 1973)

SUBJECT DESCRIPTORS: ACTIVATION ENERGY;ELECTROMOTIVE FORCE;LEAD

ALLOYS: T2;LIQUID METALS;TERNARY ALLOY SYSTEMS;THERMODYNAMIC

PROPERTIES: Q1,Q2,Q3;TIN ALLOYS: T3;ZINC ALLOYS: T1

ABSTRACT: The thermodynamic properties of liquid dilute zinc

<436> CONT.

solutions at a lead concentration of $X_{\text{Pb}} = 0.01$ to 0.7 molar fraction were determined by means of emf measurements of concentration cells. These investigations combined with previous analogical experiments at a tin concentration of $X_{\text{Sn}} = 0.01$ to 0.1 molar fraction enabled the calculation of ternary interaction parameters. The binary end-points were determined from separate measurements carried out on Zn--Sn and Zn--Pb systems at concentrations up to 0.1 molar fraction of zinc. Experimental results were interpreted by means of Krupkowski's formalism and attempts were made to calculate the interaction parameters on the basis of limiting values of activity coefficients of components of binary systems Zn--Sn, Zn--Pb, and Pb--Sn.

EDE SUBJECT CATEGORY: EDB-360102;360104

<437>

Moshnerova, I. (Statni Vyzkumny Ustav pro Stavbu Stroj, Pechovice (Czechoslovakia)) ** AN Belorusskoj SSR, Minsk. Inst. Teplo- i Massoobmena

Periodically variable wall temperature effect on heat transfer in liquid metals In Abstracts of papers and communications submitted to the 5. all-union heat and mass transfer conference ** AIX-08:299402;EDB-77:075738; (1976) INIS-mf--3284

SUBJECT DESCRIPTORS: ANALYTICAL SOLUTION;BOUNDARY CONDITIONS;HEAT TRANSFER: Q1;LIQUID METALS: T1;TEMPERATURE EFFECTS;TIME DEPENDENCE;TUBES;WALLS

ABSTRACT: None

EDE SUBJECT CATEGORY: EDB-420400

INIS CATEGORY: E11

<438>

Mueller-Buschbaum, H., Kiel Univ. (Germany, F.R.). Inst. fuer Anorganische Chemie

In Structural or solid state chemistry of rare earth sesquioxides from Goldschmidt to the present time ** AIX-07:262557;EDB-77:050763; (1976) AED-Conf--76-033-004

SUBJECT DESCRIPTORS: CHEMISTRY;CRYSTAL STRUCTURE: Q1,Q2;LANTHANUM OXIDES: T1;PHASE DIAGRAMS;PHASE TRANSFORMATIONS;RARE EARTH COMPOUNDS: T2;SOLIDS;VALENCE

<438>

<438> CONT.

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-400201

INIS CATEGORY: B12

<439>

Mukhina, I.Yu.

Some regularities of the corrosion behavior of binary magnesium alloys EDB-77:132810; Izv. Akad. Nauk SSSR, Met. (IZNMA) (2):206-213 (1976)

SUBJECT DESCRIPTORS: BINARY ALLOY SYSTEMS;CORROSION:

Q1,Q3,Q4,Q5,Q6,Q7,Q8,Q9,Q10,Q11,Q12,Q13,Q14,Q15,Q16,Q17,Q18,Q9,Q20,Q21,Q22,Q23;CORROSIVE EFFECTS:

Q2;ELECTROCHEMISTRY;MAGNESIUM ALLOYS: T1;SODIUM CHLORIDES:

T2;SOLUBILITY;STABILITY

ABSTRACT: A study is made of the influence of 26 alloying elements Li, Cu, Ag, Be, Ca, Zn, Cd, Al, Ga, In, Tl, Sc, Y, La, Ce, Pr, Nd, Sm, Si, Sn, Pb, Sb, Bi, Ti, Zr and Mn on: (a) the corrosion rate of magnesium in a 3% solution of sodium chloride and in a moist atmosphere; (b) the change in the potentials of polarization characteristics and hydrogen overvoltage in a 3% sodium chloride solution. A regular change in the corrosion properties of Mg-Me alloys was found as a function of atomic number of the alloying metal and its position in the periodic table. The corrosion resistance of the alloys studied improves in each group of the periodic table with increasing atomic number of the alloying element, decreasing electronegativity, increasing solubility in magnesium and decreasing stability of intermetallic compounds formed with magnesium. The electrochemical characteristics also change regularly in all groups and periods according to the change in corrosion resistance. These regularities in the change in corrosion properties of alloys in binary systems are explained by three factors: the electrochemical factor, valence of the dissolved element and the volumetric factor. 5 References.

EDB SUBJECT CATEGORY: EDB-360105

<440>

Munir, Z.A.; Veerkamp, G.R., California Univ., Davis (USA).
Dept. of Engineering

In Investigation of the parameters influencing the
microstructure of hot-pressed boron carbide **
INS-08:019475; ERA-02:001058; EDB-76:087293; (Nov 1975)
UCRL--13673

SUBJECT DESCRIPTORS: ALUMINIUM; ALUMINIUM FLUORIDES; BORON
CARBIDES; T1; COMPACTS; DENSITY; ELECTRON PROBES; FRACTURE
PROPERTIES; GRAIN SIZE; HOT PRESSING; MAGNESIUM; MECHANICAL
PROPERTIES; MICROHARDNESS; MICROSTRUCTURE: Q1; PHASE
DIAGRAMS; SOLUBILITY; VERY HIGH TEMPERATURE

ABSTRACT: An investigation was conducted to determine the
influence of various parameters on the microstructure of
hot-pressed boron carbide compacts. It was determined that a
critical temperature (approximately 2200°C) was
necessary to suppress exaggerated grain growth, and that
exaggerated growth was also enhanced at higher pressures
within the pressure range investigated. Microhardness
examinations and twin density analyses were also performed,
both of which demonstrated a primary dependence on grain
size, with a secondary dependence occurring between
microhardness and pressure and twin density and temperature.
Fracture analysis indicated a mixed
intergranular-transgranular fracture mode, with transgranular
fracture predominating at smaller grain sizes. Also, observed
bond strengths of fracture specimens are in reasonable
agreement with reported values. Electron microprobe scanning
was performed on B_{4C} , $B_{4C} + AlF_3$, and $B_{4C} + MgF_2$. The results indicated a solubility of Al
and Mg in the lattice. X-ray analysis was also conducted on
 $B_{4C} + 10$ percent AlF_3 specimens, and tended to
confirm the formation of a $B_{12}C_2Al$ compound,
which apparently assumes an expanded B_4C unit cell
configuration.

EDB SUBJECT CATEGORY: EDB-360202
INIS CATEGORY: B23

<441>

Murch, G.E. (Flinders Univ. of South Australia, Bedford Park.
School of Physical Sciences)

Monte Carlo analysis of diffusion and thermodynamics in UO_2
2+x/ AIX-07:236459; EDB-76:092581; Philos. Mag. (PHMAA)
32:(6):1129-1140 (Dec 1975)

SUBJECT DESCRIPTORS: CHEMICAL COMPOSITION; CRYSTAL
EFFECTS; DIFFUSION: Q1, Q2; ENTROPY; MATHEMATICAL MODELS; MONTE

<441>

<441> CONT.

CARLO METHOD; OXYGEN: T2; TRACER TECHNIQUES; URANIUM DIOXIDE:
T1; VARIATIONS; VERY HIGH TEMPERATURE

ABSTRACT: The variation of the tracer oxygen diffusion coefficient with composition has been investigated using a modified Monte Carlo method. A model which assumes a distribution of Willis 2:2:2 defect clusters subject to a short-range defect-defect interaction gives reasonable agreement with available tracer diffusion data at 800^oC. Satisfactory agreement with entropy data can also be obtained with the same model provided that a realistic electronic contribution is included.

EDB SUBJECT CATEGORY: EDB-360204

INIS CATEGORY: B23

<442>

Murch, G.E.; Thorn, R.J. (Argonne National Lab., Ill. (USA))

Thermodynamics of non-stoichiometry. A model for UC/sub x/
AIX-07:272434; EDB-77:026229; Philos. Mag. (PHMAA)
34: (2):299-309 (Aug 1976)

SUBJECT DESCRIPTORS: CARBON; CRITICAL TEMPERATURE; CRYSTAL
MODELS; ENTHALPY; ENTROPY; FCC LATTICES; MONTE CARLO
METHOD; SQUARE-WELL POTENTIAL; TEMPERATURE
DEPENDENCE; THERMODYNAMICS; Q1; URANIUM CARBIDES: T1

ABSTRACT: The partial molar properties of carbon in UC_x,
1 greater than or equal to x less than or equal to 2.0
described in terms of an f.c.c. lattice gas model in which
the C₂ groups attract only when they are nearest
neighbours. A modified Monte Carlo method is used to
calculate directly the chemical potential as a function of x
and temperature. Satisfactory agreement is obtained between
the calculated and experimental values of the partial molar
entropy and enthalpy of carbon. The nearest neighbour
interaction energy (attractive) is -1.91 kcal/mole and
was calculated from the critical temperature.

EDB SUBJECT CATEGORY: EDB-360204; 400702

INIS CATEGORY: B12

<443>

Nafikova, S.Kh.; Reshetnikova, L.P.; Novoselova, A.V.
(Moskovskij Gosudarstvennyj Univ. (USSR). Kafedra
Neorganicheskoy Khimii)

RbF-NdF₃ system AIX-08:286120; EDB-77:044316; Zh. Neorg.
Khim. (ZNOKA) 21:(9):2521-2524 (Sep 1976)

SUBJECT DESCRIPTORS: BINARY MIXTURES; CHEMICAL
COMPOSITION; DIFFERENTIAL THERMAL ANALYSIS; LATTICE
PARAMETERS; NEODYMIUM FLUORIDES: T1; PHASE DIAGRAMS:
Q1, Q2; RUBIDIUM FLUORIDES: T2; X-RAY DIFFRACTION

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-400201
INIS CATEGORY: B12

<444>

Nafikova, S.Kh.; Reshetnikova, L.P.; Efremov, V.A.; Novoselova,
A.V.

Interaction of rubidium and cesium fluorides with lutetium
trifluorides AIX-08:292423; EDB-77:050879; (VMUKAVestn. Mosk.
Univ., Ser. II. Khim.) 17:(4):506 (Jul 1976)

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-400201
INIS CATEGORY: B12

<445>

Nagai, S.; Hatakeyama, F.; Furuya, H.; Koizumi, M. (Power
Reactor and Nuclear Fuel Development Corp., Tokai, Ibaraki
(Japan). Tokai Works)

Vaporization behavior of UO₂
AIX-08:315109; ERA-02:045119; EDB-77:105137; Koon Gakkai-Shi
(KGAKI) 2:(3):144-147 (May 1976)

SUBJECT DESCRIPTORS: CATIONS; CHEMICAL REACTION
KINETICS; EQUILIBRIUM; EVAPORATION; IONIZATION; TEMPERATURE
DEPENDENCE; URANIUM DIOXIDE: T1; VAPORIZATION HEAT: Q1; VERY
HIGH TEMPERATURE

ABSTRACT: The vaporization of UO₂ was investigated by mass
spectrometric method over the range of temperature 1836^{sup}
°C to 2036^{sup} °C. From the temperature dependence of the

<445>

<445> CONT.

ion intensities of UO_2^{2+} , heat of vaporization of UO_2 was 136.7 $\pm$ 6 kcal/mole.

EDB SUBJECT CATEGORY: EDB-050700;360204
INIS CATEGORY: B23

<446>

Nagibina, L.E.; Vakhobov, A.V.; Dzhuraev, T.D. (AN Tadzhikskoj SSR, Dushanbe. Inst. Khimii)

Calculation of thermodynamic activity of components in strontium liquid alloys AIX-08:283331;EDB-77:036738; Zh. Fiz. Khim. (ZPKHA) 50:(3):806 (Mar 1976)

SUBJECT DESCRIPTORS: BINARY ALLOY
SYSTEMS;EQUILIBRIUM;INTERMETALLIC COMPOUNDS;LIQUID
METALS;PHASE DIAGRAMS;STRONTIUM ALLOYS: T1;THERMODYNAMIC
PROPERTIES: Q1

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-360104
INIS CATEGORY: B22

<447>

Nair, V.R.; Mishra, S.L.; Murthy, T.K.S., Bhabha Atomic Research Centre, Bombay (India). Chemical Engineering Div.

In Determination of thorium in rare earths chloride:
coprecipitation with cerium (IV) iodate and
spectrophotometric determination with thoron **
ERA-01:020178;EDB-76:054243; (1975) BARC--814

SUBJECT DESCRIPTORS: CARRIERS;CHEMICAL ANALYSIS:
Q2;CHLORIDES;COPRECIPITATION;RARE EARTH COMPOUNDS:
T2;SPECTROPHOTOMETRY: Q1;THORIUM: T1;THORIUM 234;TRACER
TECHNIQUES

ABSTRACT: A method for the determination of small amounts of thorium in rare earth chloride is described. This is based on the coprecipitation of thorium as iodate using cerium(IV) as carrier and final spectrophotometric determination with thoron. The efficiency of the precipitation has been checked by using ^{234}Th as tracer. (auth)

EDB SUBJECT CATEGORY: EDB-400104

<448>

Naito, K.; Kamegashira, N. ** Henley, E.J. (ed.)

High temperature chemistry of ceramic nuclear fuels with emphasis on nonstoichiometry In Advances in nuclear science and technology. Vol. 9 **

INS-77:009011; ERA-02:035846; EDB-77:072620; (1976) Academic Press Inc.

SUBJECT DESCRIPTORS: BIBLIOGRAPHIES; CRYSTAL DEFECTS; CRYSTAL STRUCTURE; DIFFUSION; ELECTRIC CONDUCTIVITY; EVAPORATION; NUCLEAR FUELS: T3; PHASE DIAGRAMS: Q1, Q2; PHASE TRANSFORMATIONS; STOICHIOMETRY; THERMODYNAMIC PROPERTIES: Q3; URANIUM CARBIDES: T2; URANIUM OXIDES: T1

ABSTRACT: Phase equilibria of ceramic nuclear fuels are reviewed. Sections are included on phase equilibria and defect structures, diffusion, and vaporization. (JRD)

EDB SUBJECT CATEGORY: EDB-050700; 360202

INIS CATEGORY: B23

<449>

Nakagawa, I. (Government Industrial Research Inst., Nagoya (Japan))

Electrode potentials of tantalum chlorides in a molten NaCl-KCl eutectic AIX-07:254856; EDB-76:071599; Nippon Kagaku Kaishi (NKAKE) 1975: (6): 958-962 (Jun 1975)

SUBJECT DESCRIPTORS: EQUILIBRIUM; EUTECTICS; MOLTEN SALTS; POTASSIUM CHLORIDES; REDOX POTENTIAL: Q1; SODIUM CHLORIDES: T1; TANTALUM CHLORIDES; TEMPERATURE DEPENDENCE

ABSTRACT: The electrode potentials of tantalum chlorides in an eutectic mixture of molten sodium and potassium chlorides were measured against a silver-silver chloride reference electrode. Potential of the reference electrode was referred to a chlorine electrode at the temperature range between 750 and 850⁰C. In equilibrium state with Ta metal, $\text{Ta(IV)} + \text{Ta} = 2\text{Ta(II)}$ was considered at relatively low concentration of tantalum ions in the melt. The standard potentials, $E^{\circ}_{\text{Ta/Ta(II)}}$ and $E^{\circ}_{\text{Ta/Ta(IV)}}$ and the redox potential, $E^{\circ}_{\text{Ta(II)/Ta(IV)}}$ were determined as follows: $E^{\circ}_{\text{Ta/Ta(II)}}/V_{\text{sub}} 0(+)} = -1.81 + 5.6 \times 10^{-4} T/^\circ\text{K}$ $E^{\circ}_{\text{Ta/Ta(IV)}}/V_{\text{sub}} 0(+)} = -1.80 + 7.7 \times 10^{-4} T/^\circ\text{K}$ $E^{\circ}_{\text{Ta(II)/Ta(IV)}}/V_{\text{sub}} 0(+)} = -1.79 + 9.8 \times 10^{-4} T/^\circ\text{K}$ where T is absolute temperature. The equilibrium constant, K, of the reaction $\text{Ta(IV)} + \text{Ta} = 2\text{Ta(II)}$ in the melt

<449>

<449> CONT.

was expressed in the following equation, $\log K = -(13/T) + 0.54$
(auth.).

EED SUBJECT CATEGORY: EDB-400201

INIS CATEGORY: B12

<450>

Nakagawa, I. (Government Industrial Research Inst., Nagoya
(Japan))

Electrochemistry of niobium and tantalum, (5). Anodic
dissolution of tantalum and the average valence of tantalum
ions in molten KCl AIX-08:300854; EDB-77:089305; Nagoya Kogyo
Gijutsu Shikensho Hokoku (NKGSA) 25:(2-3):64-69 (Feb 1976)

SUBJECT DESCRIPTORS: ANODES; CATIONS; CURRENT
DENSITY; DISSOLUTION; ELECTROCHEMISTRY:
Q1; ELECTROLYSIS; EQUILIBRIUM; MOLTEN SALTS; POTASSIUM
CHLORIDES; QUANTITY RATIO; TANTALUM: T1; TANTALUM
CHLORIDES; VALENCE

ABSTRACT: The average valence of the dissolved Ta ions was
determined by coulometry. It was 3--4 in case of the mole
fraction of the Ta being below 10^{-3} , the anodic current
density less than 80 mA/cm^2 and the temperature 800°C . The average valence increased with increase in the
quantity of electricity, and decreased with increase in the
surface area ratio of the tantalum anode to the tantalum
cathode. At tantalum mole fraction concentrations of above
 10^{-3} and anodic current densities of more than 80 mA/cm^2 , Ta(V) ion was predominantly generated. The
equilibrium between tantalum ions and tantalum metal can be
described as: $\text{Ta(IV)} + \text{Ta(S)} = 2\text{Ta(II)}$ (auth.).

EDB SUBJECT CATEGORY: EDB-400400

INIS CATEGORY: B12

<451>

Nakagawa, I. (Government Industrial Research Inst., Nagoya
(Japan)); Kiriara, T.

Electrochemistry of niobium and tantalum. VI. Electrode
potentials of tantalum chlorides in molten KCl
AIX-08:305613; EDB-77:107434; Nagoya Kogyo Gijutsu Shikensho
Hokoku (NKGSA) 25:(2):70-75 (Feb 1976)

SUBJECT DESCRIPTORS: CURRENT DENSITY; ELECTROCHEMISTRY:
Q2; ELECTROLYSIS; EQUILIBRIUM; MOLTEN SALTS; POTASSIUM
CHLORIDES; REDOX POTENTIAL: Q1; TANTALUM: T2; TANTALUM
CHLORIDES: T1; TEMPERATURE DEPENDENCE; VALENCE

<451> CONT.

ABSTRACT: The electrode potentials of tantalum chlorides in molten potassium chloride were measured by using a silver-silver chloride reference electrode. Potentials of the reference electrode were referred to a chlorine electrode in the temperature range from 775 to 875°C. The standard potentials, $E^{\circ}_{\text{Ta/Ta(II)}}$, and $E^{\circ}_{\text{Ta/Ta(IV)}}$, and the redox potential $E^{\circ}_{\text{Ta(II)/Ta(IV)}}$ were determined as follows: (A) $E^{\circ}_{\text{Ta/Ta(II)}}$ / $V^{\circ}(+) = -1.67 + 6.4 \times 10^{-4} T/K$ (up to 4×10^{-4} mol. fract.), $E^{\circ}_{\text{Ta/Ta(IV)}}$ / $V^{\circ}(+) = -2.01 + 7.7 \times 10^{-4} T/K$, and $E^{\circ}_{\text{Ta(II)/Ta(IV)}}$ / $V^{\circ}(+) = -2.35 + 9.0 \times 10^{-4} T/K$; (B) $E^{\circ}_{\text{Ta/Ta(II)}}$ / $V^{\circ}(+) = -1.74 + 7.2 \times 10^{-4} T/K$ (above 4×10^{-4} mol. fract.), $E^{\circ}_{\text{Ta/Ta(IV)}}$ / $V^{\circ}(+) = -1.65 + 5.0 \times 10^{-4} T/K$, and $E^{\circ}_{\text{Ta(II)/Ta(IV)}}$ / $V^{\circ}(+) = -1.56 + 2.8 \times 10^{-4} T/K$. The equilibrium constants (K) of the reaction $\text{Ta(IV)} + \text{Ta} = 2\text{Ta(II)}$ in the melt were expressed in the following equations, (A) $\log K = -2.6 - 6817/T$, and (b) $\log K = -4.4 + 1804/T$.

EDB SUBJECT CATEGORY: EDB-400400

INIS CATEGORY: B12

<452>

Nakajima, T.; Nakanishi, K.; Watanabe, N. (Kyoto Univ. (Japan). Faculty of Engineering)

Dispersion of metallic lithium in various molten salts

AIX-07:259592; EDB-77:030776; Nippon Kagaku Kaishi (NKAKB) 1975: (4):617-621 (Apr 1975)

SUBJECT DESCRIPTORS: CALCIUM CHLORIDES; DISPERSIONS: Q1; EUTECTICS; GASES; LITHIUM: T1; LITHIUM CHLORIDES; LITHIUM FLUORIDES; LITHIUM NITRIDES; LITHIUM OXIDES; MOLTEN SALTS; NITROGEN; OXYGEN; POTASSIUM CHLORIDES; POTASSIUM FLUORIDES; SODIUM CHLORIDES; SOLUBILITY

ABSTRACT: The dispersion of metallic lithium in various molten salts was studied at constant partial pressures of nitrogen or oxygen gas. The increase of dispersed lithium amounts in the presence of nitrogen gas was more rapid than that in the presence of oxygen gas. A large amounts of lithium dispersed in LiCl and LiCl-KCl eutectic mixture where the solubilities of lithium oxide or nitride were almost equal to those in liquid lithium. On the other hand, the amounts of dispersed lithium were small in LiF-LiCl, LiF-KF and NaCl-CaCl₂.

<452>

<452> CONT.

systems, where the solubilities of oxide or nitride were much different from those in lithium.

EDB SUBJECT CATEGORY: EDB-360104

INIS CATEGORY: E12

<453>

Natsume, H.; Matsuura, S.; Okashita, H.; Umezawa, H.; Ezure, H.,
International Atomic Energy Agency, Vienna (Austria)

In Collection of gamma spectra data of irradiated light water moderated reactor spent fuel and study of the applicability of the methods for fuel identification. Final report for the period 1 February 1972--October 1975 **

AIX-07:242718; ERA-01:025733; EDB-76:074101; (Oct 1975)

IAEA-B--1119-F

SUBJECT DESCRIPTORS: BURNUP; COMPARATIVE

EVALUATIONS; DATA; DESTRUCTIVE TESTING; FISSION PRODUCTS; GAMMA

FUEL SCANNING: T, Q1, Q2; GAMMA SPECTRA; ISOTOPE RATIO; JPDR

REACTOR: T1; MASS SPECTROSCOPY; NONDESTRUCTIVE ANALYSIS; REACTOR

OPERATION; SPENT FUEL ELEMENTS: T2; SPENT FUELS; TRANSURANIUM

ELEMENTS; URANIUM

ABSTRACT: The gamma spectrometric data are obtained non-destructively on 96 points of a spent fuel assembly of the JPDR-1 by using an on-site gamma scanning facility in the fuel storage pond. Destructive gamma spectrometric, alpha spectrometric, chemical and mass spectrometric analyses are also performed for uranium, transuranium elements and some fission products on the selected 24 samples among the 96 non-destructively measured points. The feasibility of the non-destructive measurements is examined by comparing those results with the results of destructive analyses. Isotopic ratios of various combinations of fission products or heavy nuclides are studied in relation to burn-up and Pu/U ratios. The results show that the relationships are affected by reactor characteristics of BWR and the operation history. The applicability of the isotopic correlations technique is discussed from the standpoint of analysing reactor characteristics and applying them to safeguarding of nuclear materials.

EDB SUBJECT CATEGORY: EDB-210100; 400103

INIS CATEGORY: E36

<454>

Nazare, S.; Ondracek, G.; Schulz, B. (Kernforschungszentrum Karlsruhe (Germany, F.R.). Inst. fuer Material- und Festkoerperforschung) ** Kernforschungszentrum Karlsruhe (Germany, F.R.). Projekt Nukleare Sicherheit

Properties of corium melts In Semiannual report 1975/2 **
AIX-08:293632; ERA-02:034485; EDB-77:073861; (Jun 1976)
KFK--2262

SUBJECT DESCRIPTORS: CORIUM: T1; DATA; MELTDOWN: T, Q2; MELTING POINTS; PHYSICAL PROPERTIES: Q1; REACTOR CORES: T2; REACTOR MATERIALS; SPECIFIC HEAT; STEELS; THERMAL CONDUCTIVITY; THERMODYNAMIC PROPERTIES; URANIUM DIOXIDE; VISCOSITY; ZIRCALOY.

ABSTRACT: In order to describe the behaviour of the molten core as a consequence of a LOCA, data on properties of such melts are required. The properties are however dependent on the chemical composition and structural characteristics of the melts which in turn depend on the oxidation potential of the environment. Available experimental evidence was used to define 4 types of melts based on interaction products between varying compositions of UO_2 -Zr-Steel in either inert, partially or totally oxidizing conditions. A critical review of theoretical and experimental results on properties such as viscosity, heat capacity and thermal conductivity (in the liquid state) was used as a basis to estimate these properties for the various types of Corium melts.

EDF SUBJECT CATEGORY: EDB-220900; 360104
INIS CATEGORY: E11

<455>

Nicholson, E.L. Jr., Oak Ridge National Lab., Tenn. (USA)

In PuO_2 dissolution problem for LWR plutonium recycle and LMFBR fuels: fabrication and reprocessing problems and their resolution ** ERA-02:058488; EDB-77:137284; (Jul 1977)
ORNL/TM--5903

SUBJECT DESCRIPTORS: BWR TYPE REACTORS; CRITICALITY; DISSOLUTION: Q1; FUEL CYCLE: Q2, Q3; FUEL FABRICATION PLANTS; FUEL REPROCESSING PLANTS; LMFBR TYPE REACTORS: T3; PLUTONIUM DIOXIDE: T1; PWR TYPE REACTORS; SOLUBILITY; URANIUM DIOXIDE; WATER COOLED REACTORS: T2

ABSTRACT: A survey was made of the information reported to date for laboratory-scale dissolution experiments on PuO_2 - UO_2 fuels, of reprocessing plant problems that might be encountered with these fuels, and of the fabrication methods for producing these fuels. The possibility of

<455>

<455> CONT.

producing fuels that will be highly soluble in pure nitric acid without resorting to the use of corrosive fluorides for complete dissolution is examined. The report concludes that production of highly soluble fuel is possible, that it is probably economically justifiable, and that fluorides are not necessary for dissolution. Highly soluble fuel would likely have minimal impact on reprocessing criticality and waste disposal problems. Reactor specifications for PuO_2 fuels seemingly permit a greater degree of nonhomogeneity than is desirable for achieving high solubility. A fuel solubility criteria is therefore proposed that would limit the amount of insoluble PuO_2 in irradiated fuels to less than or equal to 0.01 percent of the total amount of plutonium present.

EDB SUBJECT CATEGORY: EDB-050800;050700

<456>

Nickerson, G.M.; Kastenber, W.E. (California Univ., Los Angeles (USA). Dept. of Energy and Kinetics)

Preliminary assessments of carbide fuel pins during mild overpower transients in LMFBRs
 AIX-07:257871;ERA-02:010052;EDB-77:006616; Nucl. Eng. Des. (NEDEA) 32:(2):209-233 (Feb 1976)

SUBJECT DESCRIPTORS: CHEMICAL COMPOSITION;FUEL PINS: T2;GRAIN BOUNDARIES;H CODES;LMFBR TYPE REACTORS; T1:PLUTONIUM CARBIDES;SPECIFIC HEAT;THERMAL CONDUCTIVITY;THERMAL EXPANSION;TRANSIENTS: Q1,Q2;URANIUM CARBIDES

ABSTRACT: Mixed carbide fuel has been proposed for advanced liquid metal fast breeder reactors (LMFBRs). The general characteristics of carbide fuel are reviewed and a survey of available data for use in safety studies is presented. A preliminary investigation of the unprotected transient overpower accident sequence postulated for a large carbide fueled reactor is also presented. The HOPE computer code is the computational tool used to analyse a conceptual core design fueled with mixed carbides of uranium and plutonium. The details of this design are presented and compared with those of the Clinch River Breeder Reactor Plant. Results of HOPE calculations for both helium and sodium bonded carbide fueled cores are presented and compared with a similar study performed on an oxide core. Conclusions are drawn based on

<457> CONT.

the findings of the preliminary transient overpower studies conducted.

EDB SUBJECT CATEGORY: EDB-220900;210500
INIS CATEGORY: E35

<457>

Nisel'son, L.A.; Lyzlov, Y.N.; Solov'ev, S.I.

Phase equilibrium in NdCl_3 -KCl system
AIX-08:339952;EDB-78:017337; Zh. Neorg. Khim. (ZNOKA)
22: (1):180-183 (Jan 1977)

SUBJECT DESCRIPTORS: CRYSTALLIZATION;EQUILIBRIUM:
Q1,Q2;LIQUIDS;NEODYMIUM CHLORIDES: T1;PHASE
TRANSFORMATIONS;POTASSIUM CHLORIDES: T2;QUANTITY
RATIO;VAPORS;VOLATILITY

ABSTRACT: Liquid-steam equilibrium in the NdCl_3 -KCl system has been studied by the boiling point method. A considerable negative deviation of the system from ideal conditions has been established. Concentration dependence of the relative volatility coefficients α on liquid phase composition has been found. α_{real} decreases monotonously from 4.2 to 1.5 with increasing NdCl_3 concentration. The equilibrium distribution coefficient has been determined for equilibrium crystals-liquid in the system studied with the aid of directed crystallization. In the region of pure neodymium trichloride the coefficient is equal to 0.24.

EDB SUBJECT CATEGORY: EDB-400201
INIS CATEGORY: B12

<458>

Nisel'son, L.A.; Sokolova, T.D.; Leonov, A.M. (Gosudarstvennyj Nauchno-Issledovatel'skij i Proektnyj Inst. Redkometallicheskoj Promyshlennosti, Moscow (USSR))

Density and viscosity of distillable products from ZrCl_4 and HfCl_4 interaction with POCl_3
AIX-08:295353;EDB-77:056166; Izv. Vyssh. Uchebn. Zaved., Tsvet. Metall. (IVUTA) (2):102-103 (1976)

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-400201
INIS CATEGORY: B12

<459>

<459>

Noda, Y.; Naito, K. (Nagoya Univ. (Japan). Faculty of Engineering)

Thermal conductivity and diffusivity of $U_{4O_{9-y}}$ from 250 to 415 K AIX-08:324324; EDB-77:132966; J. Nucl. Mater. (JNUMA) 66: (1):. 17-22 (1977)

SUBJECT DESCRIPTORS: CRYSTAL LATTICES; INTERSTITIALS; MEAN FREE PATH; MEDIUM TEMPERATURE; PHASE TRANSFORMATIONS; PHONONS; POROSITY; SPECIFIC HEAT; THERMAL CONDUCTIVITY: Q1; THERMAL DIFFUSIVITY: Q1; URANIUM OXIDES: T1

ABSTRACT: The thermal conductivity and diffusivity of $U_{4O_{9-y}}$ with various O/U ratios (2.227, 2.234, 2.254) were measured by means of a scanning temperature method in the temperature range from 250 to 415 K. An anomaly of the second-order phase transition type was revealed by the thermal diffusivity measurements, and the peak temperature for each O/U ratio was in good agreement with that previously obtained from heat capacity measurements. The anomaly was not as clearly observed in the thermal conductivity data as in the thermal diffusivity data. The measured values for thermal conductivity of $U_{4O_{9-y}}$ were very low: from 10^{-2} to 2×10^{-2} W K⁻¹ cm⁻¹ over the temperature range investigated, and varied little with temperature. The phonon mean free path values calculated from the thermal conductivity data were very small, and all fell within the range 2.5 to 5.5 Å, being higher in the specimens with the lower oxygen content. Mean free path values decreased with increase in temperature, and the rate of decrease was greater in the region of the transition.

EDB SUBJECT CATEGORY: EDB-360204
INIS CATEGORY: B24

<460>

Nevikov, I.I.; Proskurin, V.B.; Spiridonova, T.N. (AN SSSR, Moscow. Inst. Metallurgii)

Some structural characteristics of rubidium and cesium in the vicinity of their melting points AIX-07:278273; EDB-77:030829; Zh. Fiz. Khim. (ZPKHA) 50: (7):1657-1660 (Jul 1976)

SUBJECT DESCRIPTORS: CESIUM: T1; COMPARATIVE EVALUATIONS; COMPRESSIBILITY; DENSITY; ELECTRONIC STRUCTURE: Q1, Q2; LIQUID METALS; MELTING POINTS; RUBIDIUM: T2; SOUND WAVES; VELOCITY

<460> CONT.

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-360104

INIS CATEGORY: B22

<461>

Novikov, N.I.; Novichkov, V.Kh.; Rybkina, R.D. ** Poletaev, I.F.; Rubajlova, K.M. (eds.), Gosudarstvennyj Nauchno-Issledovatel'skij i Proektnyj Inst. Redkometallicheskoj Promyshlennosti, Moscow (USSR)

Electric conductivity of melts of the LiF-misch metal fluoride system in rare earth metals ** AIX-08:292433; EDB-77:074898; (1976) INIS-mf--3315

SUBJECT DESCRIPTORS: CERIUM FLUORIDES: T1; ELECTRIC CONDUCTIVITY: Q1, Q2, Q3, Q4, Q5; LANTHANUM FLUORIDES: T2; LITHIUM FLUORIDES: T3; MISCH METAL; MOLTEN SALTS; NEODYMIUM FLUORIDES: T4; PHASE DIAGRAMS; PRASEODYMIUM FLUORIDES: T5; QUANTITY RATIO; VERY HIGH TEMPERATURE

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-360603

INIS CATEGORY: B12

<462>

Novikov, N.I.; Rybkina, R.D.; Novichkov, V.Kh. ** Poletaev, I.F.; Rubajlova, K.M. (eds.), Gosudarstvennyj Nauchno-Issledovatel'skij i Proektnyj Inst. Redkometallicheskoj Promyshlennosti, Moscow (USSR)

Misch metal solubility in melt of LiF-BaF₂ system in rare earth metals ** AIX-08:292434; EDB-77:074899; (1976) INIS-mf--3315

SUBJECT DESCRIPTORS: BARIUM FLUORIDES: T2; LITHIUM FLUORIDES: T3; MISCH METAL: T1; MOLTEN SALTS; NEODYMIUM ALLOYS; PRASEODYMIUM ALLOYS; QUANTITY RATIO; SOLUBILITY: Q1; SOLVENT PROPERTIES: Q2, Q3; VERY HIGH TEMPERATURE

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-360603; 360104

INIS CATEGORY: B12

<463>

<463>

Novokhatskii, I.A.; Arkharov, V.I.; Prokopenko, V.K. (AN
Ukrainskoj SSR, Donetsk. Fiziko-Tekhnicheskij Inst.)

NMR study on structural microinhomogeneity of fluoride-oxide
melts AIX-07:271652;EDB-77:018942; Fiz. Tverd. Tela (PTVTA)
17:(5):1520-1522 (May 1975)

SUBJECT DESCRIPTORS: FLUORIDES: T1;FLUORINE 19;HIGH
TEMPERATURE;LEAD FLUORIDES;MICROSTRUCTURE: Q1,Q2;NUCLEAR
MAGNETIC RESONANCE;ORDER-DISORDER TRANSFORMATIONS;OXIDES:
T2;QUANTITY RATIO;SILICON OXIDES;SOLID
CLUSTERS;SPECTRA;TEMPERATURE DEPENDENCE;VERY HIGH TEMPERATURE

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-360202
INIS CATEGORY: A13

<464>

Nowotny, H.; Seifert, K. (Vienna Univ. (Austria). Inst. fuer
Physikalische Chemie)

3. Zirconium compounds with non-metals In Zirconium:
Physico-chemical properties of its compounds and alloys **
At. Energy Rev., AIX-07:277855;EDB-77:031075; (1976) IAEA

ABSTRACT: The crystal structure and density of various
Zr-compounds with non-metals (H, B, C, Si, Ge, N, P, As, Sb,
O, S, Se, Te, F, Cl, Br, I) are listed in table form. The
values are taken from the literature (Chemical Abstracts up
to 1973, references given). Structural type and atomic
parameters, crystal system, unit cell dimensions and density
of the compounds are given. Lattice parameters are given for
some of the compounds as a function of temperature or
chemical composition (diagrams). The tables are arranged in
the order binary compounds, ternary compounds and complex
compounds.

EDB SUBJECT CATEGORY: EDB-360602
INIS CATEGORY: B12

<465>

O'Donnell, T.A.; Waugh, A.B.; Randall, C.H. (Melbourne Univ.,
Parkville (Australia))

Reactivity of transition metal fluorides. IX. Protactinium
pentafluoride and tetrafluoride AIX-08:346257;EDB-78:028035;
J. Inorg. Nucl. Chem. (JINCA) 39:(9):1597-1600 (1977)

SUBJECT DESCRIPTORS: CHLORIDES;OXIDATION: Q1;PROTACTINIUM

<465> CONT.

FLUORIDES: T1;REDUCTION: Q1;THERMODYNAMIC ACTIVITY

ABSTRACT: In a continuing survey of the relative reactivities of higher fluorides of d- and f-transition elements, redox reactions of protactinium pentafluoride with selected reductants have been studied. Halogen-exchange reactions with covalent chlorides have been investigated for both the pentafluoride and tetrafluoride. The results of these reactions are correlated with those from previous investigations. Special attention was given to preparative and reaction techniques because of the radioactive nature and because of the small quantities of the compounds available for this work compared with the scale of earlier related studies in this series.

EDE SUBJECT CATEGORY: EDB-400702

INIS CATEGORY: B13

<466>

O'Hare, P.A.G.; Boerio, J.

Cesium nitrate. Standard enthalpies of solution and formation and the standard entropy at 298.15 K

ERA-01:011161;EDB-76:036956; J. Chem. Thermodyn. (JCTDA) 7::937-941 (1975)

SUBJECT DESCRIPTORS: CESIUM NITRATES: T1;ENTROPY: Q1;FORMATION HEAT: Q1;MEDIUM TEMPERATURE;MONOCRYSTALS;SOLUTION HEAT: Q1

ABSTRACT: Calorimetric measurements on single-crystal CsNO_3 have yielded the standard enthalpy of solution at 298.15 K, $\Delta H^\circ_{\text{soln}} = (9639 \pm 10) \text{ cal/sub th/mol}^{-1}$. Based on this value, the standard enthalpy of formation, $\Delta H^\circ_f(\text{CsNO}_3, \text{c}, 298.15 \text{ K})$ was calculated to be $-(120.88 \pm 0.12) \text{ kcal/sub th/mol}^{-1}$. Solubility and activity coefficient data for CsNO_3 gave $779 \text{ cal/sub th/mol}^{-1}$ for the standard Gibbs energy of solution, $\Delta G^\circ_{\text{soln}}$. Thence, the standard entropy, $S^\circ(\text{CsNO}_3, \text{c}, 298.15 \text{ K})$, was deduced to be $(37.1 \pm 0.2) \text{ cal/sub th/K}^{-1} \text{ mol}^{-1}$. (auth)

EDE SUBJECT CATEGORY: EDB-400201

<467>

<467>

O'Hare, F.A.G.; Johnson, G.K.

Lithium nitride (Li_3N): standard enthalpy of formation by solution calorimetry ERA-01:013733; EDB-76:048938; J. Chem. Thermodyn. (JCTDA) 7:13-20 (1975)

SUBJECT DESCRIPTORS: AMMONIUM CHLORIDES; CALORIMETRY; CHEMICAL REACTIONS; ENTHALPY; FORMATION HEAT: Q1; HYDROCHLORIC ACID; LITHIUM CHLORIDES; LITHIUM NITRIDES: T1; SOLUTION HEAT; WATER

ABSTRACT: The enthalpies of reaction of a carefully prepared and characterized specimen of lithium nitride, Li_3N , with H_2 and with $\text{HCl}(\text{aq})$ were found to be $-(581.62 \pm 1.42)$ kJ mol⁻¹ [$-(139.01 \pm 0.34)$ kcal/sub th/ mol⁻¹] and $-(803.50 \pm 1.26)$ kJ mol⁻¹ [$-(192.04 \pm 0.30)$ kcal/sub th/ mol⁻¹], respectively. Enthalpies of solution of LiCl and NH_4Cl were also measured. Combination of the measured results with auxiliary thermochemical data from the literature yielded values of $-(165.14 \pm 1.55)$ kJ mol⁻¹ [$-(39.47 \pm 0.37)$ kcal/sub th/ mol⁻¹] and $-(164.77 \pm 1.51)$ kJ mol⁻¹ [$-(39.38 \pm 0.36)$ kcal/sub th/ mol⁻¹], respectively, for the standard enthalpy of formation, $\Delta H_f^\circ(\text{Li}_3\text{N}, \text{c}, 298.15 \text{ K})$. A weighted mean value, $-(164.93 \pm 1.09)$ kJ mol⁻¹ [$-(39.42 \pm 0.26)$ kcal/sub th/ mol⁻¹], is recommended for $\Delta H_f^\circ(\text{Li}_3\text{N}, \text{c}, 298.15 \text{ K})$. This result differs by about 8 kcal/sub th/ mol⁻¹ from previous determinations. (auth)

EDB SUBJECT CATEGORY: EDB-360603

<468>

Oden, L.L.; Sanker, P.E.; Babitzke, H.R.; Sumner, D.H., Bureau of Mines, Albany, Oreg. (USA). Albany Metallurgy Research Center

In Phase relations in the alkaline earth fluoride-yttria systems (in three parts). 1. The CaF_2 - Y_2O_3 phase diagram ** EDB-77:120762; (Oct 1970) BM-RI--7437

SUBJECT DESCRIPTORS: CALCIUM FLUORIDES: T1; DIFFERENTIAL THERMAL ANALYSIS; ELECTRON MICROSCOPY ANALYSIS; PHASE STUDIES: Q1, Q2; YTTRIUM OXIDES: T2

ABSTRACT: The Bureau of Mines is developing melting data for binary combinations of the alkaline earth fluorides and yttria to facilitate the formulation of slag compositions. The present report deals with the CaF_2 - Y_2O_3 system for which solidus and liquidus temperatures and

<468> CONT.

solid solubilities were determined. The information is presented as the phase diagram. All specimen compositions were blended from high-purity materials and encapsulated in Mo or Pt20Rh prior to heat treatment. Liquidus temperatures were determined by ceramographic examination of heat-treated and quenched specimens. Solidus temperatures were determined by differential thermal analysis and by electron microprobe analysis of quenched specimens. Terminal solid solubilities were derived by parametric methods. The peritectic system was found to be pseudobinary due to the formation of yttrium oxyfluoride at temperatures near the liquidus. The peritectic point occurs at 7 mole pct Y_{23}O_3 and $1,438 \pm 5^\circ\text{C}$. The terminal solid solubility of Y_{23}O_3 in CaF_2 decreases from 7 mole pct at $1,438^\circ\text{C}$ to 1 mole pct at $1,100^\circ\text{C}$, and the solubility of CaF_2 in Y_{23}O_3 decreases from 9.5 to 7 mole pct over the same temperature interval.

EDB SUBJECT CATEGORY: EDB-400201;360202

<469>

Oden, L.L.; Sanker, P.E.; Babitzke, H.R.; Sumner, D.H., Bureau of Mines, Albany, Oreg. (USA). Albany Metallurgy Research Center

In Phase relations in the alkaline earth fluoride-yttria systems (in three parts). 2. The BaF_2 - Y_{23}O_3 phase diagram ** EDB-77:120763; (Oct 1970) BM-RI--7438

SUBJECT DESCRIPTORS: RARIUM FLUORIDES: T1; BINARY MIXTURES; EUTECTICS; PHASE STUDIES: Q1, Q2; THERMAL ANALYSIS; X-RAY DIFFRACTION; YTTRIUM OXIDES: T2

ABSTRACT: The Bureau of Mines is developing melting data for binary combinations of the alkaline earth fluorides and yttria to facilitate the formulation of slag composition. Liquidus and solidus temperatures and solid solubilities were determined for the BaF_2 - Y_{23}O_3 system, and the information is presented as the phase diagram. The phase diagram was constructed from data obtained by ceramography, x-ray diffraction, and thermal analysis of specimens encapsulated in small molybdenum ampoules. The data indicate a simple eutectic system with maximum terminal solid solubilities of less than 1 mole pct. The eutectic temperature is $1,343^\circ\text{C}$, and the composition of the eutectic is less than 1 mole pct Y_{23}O_3 .

EDB SUBJECT CATEGORY: EDB-400201;360202

<470>

<470>

Odoj, R.; Hilpert, K.; Nuernberg, H.W. (Kernforschungsanlage Juelich G.m.b.H. (F.R. Germany)); Foerthmann, R.; Naoumidis, A. (Kernforschungsanlage Juelich G.m.b.H. (F.R. Germany). Inst. fuer Reaktorwerkstoffe und Heisse Zellen)

Study of the retention of Ba in UO_2 nuclear fuel particles by ZrO_2 as getter

AIX-07:263502; ERA-02:016547; EDB-77:023622; J. Nucl. Mater. (JNUMA) 60:(2):216-222 (May 1976)

SUBJECT DESCRIPTORS: BARIUM; BARIUM OXIDES; COATED FUEL PARTICLES: Q1, T2; FISSION PRODUCTS; GETTERS: Q2; HTGR TYPE REACTORS: T1; MASS SPECTROSCOPY; URANIUM DIOXIDE; ZIRCONIUM OXIDES

ABSTRACT: A new concept to retain solid fission products in the coated particles of gas-cooled high-temperature reactors is the addition of refractory oxides to the fuel kernels as a getter. The suitability of this concept for the retention of BaO by ZrO_2 was determined. Therefore, the evaporation of $BaZrO_3$ was studied by high temperature mass spectrometry with a Knudsen cell made of Mo and the loss of Ba from model kernels heated for 100 h at different temperatures in a furnace was determined. From the results obtained with each of the two methods which were carefully discussed by theoretical consideration it follows independently that Ba can be retained by ZrO_2 but that Al_2O_3 is even more efficient for the retention of Ba than ZrO_2 . Moreover, it could be shown that BaO is the main species in the gaseous phase over $BaZrO_3$.

EDB SUBJECT CATEGORY: EDB-210300
INIS CATEGORY: E33

<471>

Cifer, H.P. ** Hill, M.R. (comp.), Oak Ridge National Lab., Tenn. (USA)

Metallurgical characterization of prototype tubesheet forging
In Mechanical properties test data for structural materials.
Quarterly progress report for period ending April 30, 1977
** ERA-03:004358; EDB-78:004440; (Jun 1977) ORNL--5287

SUBJECT DESCRIPTORS: ANNEALING; CHROMIUM-MOLYBDENUM STEELS: M1; HIGH TEMPERATURE; LMFBR TYPE REACTORS; MECHANICAL PROPERTIES; REACTOR MATERIALS; TENSILE PROPERTIES: Q1; VERY HIGH TEMPERATURE

ABSTRACT: Room and elevated temperature tensile tests have been performed on the prototype lower tubesheet forging procured by GE from Cameron Iron Works (CIW). These preliminary tensile tests were performed on tubesheet specimens given

<471> CONT.

either a full anneal plus a 40 hour 727°C (1340°F) post weld heat treatment (PWHT) (designated heat treatment F), or a full anneal plus this PWHT plus a 1000 hour 510°C (950°F) thermal aging/embrittlement treatment (designated heat treatment Q). The results are given in Table 8.2.1 along with the room temperature results supplied by CIW for heat treatment P.

EDB SUBJECT CATEGORY: EDB-360103

<472>

Ofitserov, A.A.; Golubtsov, V.V.; Dubrovin, M.N. et al.
(Moskovskii Inst. Stali i Splavov (USSR))

Determination of surface tension of a metal with high vapor pressure AIX-07:270426; EDB-77:030766; Izv. Vyssh. Uchebn. Zaved., Tsvet. Metall. (IVUTA) (4):171-172 (1975)

SUBJECT DESCRIPTORS: EQUATIONS; LEAST SQUARE FIT; LIQUID METALS: T1; SURFACE TENSION: Q1; TELLURIUM; TEMPERATURE DEPENDENCE; VAPOR PRESSURE

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-360104

INIS CATEGORY: B22

<473>

Ohno, H.; Furukawa, K. (Molten Materials Laboratory, Japan Atomic Energy Research Institute, Tokai, Ibaraki, Japan)

Phase diagrams in the pseudobinary systems of $\text{Li}_{2\text{BeF}_4}$, alkali fluoroborates and alkali fluorides
AIX-08:310031; EDB-77:094499; J. Nucl. Mater. (JNMMA) 64:(1): 37-43 (Jan 1977)

ABSTRACT: The phase diagrams in the systems $\text{Li}_{2\text{BeF}_4}$ - NaBF_4 , $\text{Li}_{2\text{BeF}_4}$ - KBF_4 , $\text{Li}_{2\text{BeF}_4}$ - LiBF_4 , NaI - LiI , NaBF_4 - LiF and NaBF_4 - KF were examined. The systems $\text{Li}_{2\text{BeF}_4}$ - NaBF_4 , $\text{Li}_{2\text{BeF}_4}$ - KBF_4 and NaBF_4 - LiF have a miscibility gap in the liquid phase. The much greater miscibility in the $\text{Li}_{2\text{BeF}_4}$ - NaBF_4 system compared to the $\text{Li}_{2\text{BeF}_4}$ - KBF_4 system can be attributed to the smaller difference in the radii of the cations. The solubility of LiF in molten NaBF_4 at 500°C is about 1 mole% and it is slightly greater than that in the KBF_4 - LiF system. The good similarity of the phase diagrams of the systems RF - $\text{R}'\text{BF}_4$ and RF - $\text{R}'\text{I}$ ($\text{R}, \text{R}' = \text{Li}, \text{Na}$ and K) is also explained by the fact that the EF_{sub}

<473>

<473> CONT.

$4f^{14}7s^2$ ion has nearly the same ionic radius as the $1f^{14}7s^2$ ion.

EDB SUBJECT CATEGORY: EDB-360602

INIS CATEGORY: B24

<474>

Ohse, R.W.; Babelot, J.P.; Kinsman, P.R. (Commission of the European Communities, Karlsruhe (Germany, F.R.). European Inst. for Transuranium Elements); Brumme, G.D. (Technische Hochschule Darmstadt (Germany, F.R.). Inst. fuer Angewandte Physik)

Vapor pressure studies over liquid uranium oxide and uranium plutonium oxide up to 5,000 K
AIX-08:289670; ERA-02:024848; EDB-77:049773; Ber. Bunsenges. Phys. Chem. (BBPCA) 80: (8):780-786 (Aug 1976)

SUBJECT DESCRIPTORS: ABSORPTION; FUNCTIONAL MODELS; LASER-RADIATION HEATING; LIQUIDS; LMFBR TYPE REACTORS: T1; MELTDOWN: M2, Q1; MIXTURES; NUCLEAR FUELS; OPTICAL PYROMETERS; PLUTONIUM; PLUTONIUM DIOXIDE: T4; PULSED IRRADIATION; SUBLIMATION HEAT; TEMPERATURE DEPENDENCE; THERMODYNAMICS: Q2; URANIUM DIOXIDE: T3; VAPOR PRESSURE: Q3, Q4; VAPORIZATION HEAT; VERY HIGH TEMPERATURE

ABSTRACT: The estimation of the energy release in a hypothetical fast breeder reactor core meltdown accident requires a detailed analysis of the disassembly process. Since the vapor pressure presents the ultimate shut-down mechanism, the required data are given by the equation of state of the irradiated fuel. The pressure over the ternary uranium-plutonium oxide was determined to $\log p(\text{atm}) = 7.966 - 28,137/T$, yielding a heat of evaporation of $\Delta H_{\text{sub}}(\text{evap}) = 128.7 \text{ kcal/mol}$, fully consistent with the heat of sublimation of $\Delta H_{\text{sub}}(\text{sub}) = 148.4 \text{ kcal/mol}$ below the melting point, and the heat of fusion of 19.4 kcal/mol . The corresponding equation for liquid UO_2 was determined as $\log p(\text{atm}) = 7.7 - 27,900/T$, with a heat of evaporation of $\Delta H_{\text{sub}}(\text{evap}) = 127.6 \text{ kcal/mol}$, again in consistency with the assessed data below the melting point of $\Delta H_{\text{sub}}(\text{sub}) = 144.1 \text{ kcal/mol}$, and the heat of fusion of 17.7 kcal/mol . For this a new high-energy laser technique, including fast temperature recording in the microsecond range, was developed. The first quantitative measurements on $(\text{U,Pu})\text{O}_2$ above 4,000 K were reported to the IAEA Symposium on Thermodynamics of Nuclear Materials in Vienna. The investigation of the total temperature range from 3,000 K to $T_{\text{sub}}(\text{c}) = 7,560 \text{ K}$ revealed two principle experimental difficulties, a liquid layer movement at temperatures below 4,000 K, and optical absorption due to increasing thermal

<474> CONT.

ionisation above 5,000 K. A new double intensity laser pulse technique is presented, which allows the extension of measurements to below 4,000 K. In addition a multi-wavelength pyrometer technique has been developed to eliminate, and at a later stage to determine, spectral emissivities above the melting point, during laser heating within pulse times of micro and nanoseconds.

EDB SUBJECT CATEGORY: EDB-220900;210500;360204

INIS CATEGORY: B23

<475>

Ohse, R.W.; Berrie, P.G. (Commission of the European Communities, Karlsruhe (F.R. Germany). European Inst. for Transuranium Elements); Bogensberger, H.G.; Fischer, E.A. (Kernforschungszentrum Karlsruhe (F.R. Germany). Inst. fuer Angewandte Systemanalyse)

Extension of vapour pressure measurements of nuclear fuels (U,Pu)O₂ and UO₂ to 7000 K for fast reactor safety analysis AIX-07:244140; ERA-02:006655; EDB-77:000340; J. Nucl. Mater. (JNUMA) 59:(2):112-124 (Feb 1976)

SUBJECT DESCRIPTORS: BINARY MIXTURES; FUSION
HEAT; LASERS; MELTDOWN; OPTICAL PYROMETERS; PLUTONIUM DIOXIDE:
T1; RADIATION HEATING; SOLID FUELS; SUBLIMATION HEAT; TEMPERATURE
DEPENDENCE; TEMPERATURE MEASUREMENT; ULTRAHIGH
TEMPERATURE; URANIUM DIOXIDE: T2; VAPOR PRESSURE:
Q1, Q2; VAPORIZATION HEAT; VERY HIGH TEMPERATURE

ABSTRACT: A new high-energy laser technique, including fast temperature recording in the microsecond range, was developed for measuring the vapour pressure of fast breeder uranium-plutonium oxide fuels up to 7000 K. In the ternary system, the pressure of uranium-plutonium oxide above its melting point was determined to be $\log p \text{ (atm)} = 7.966 - (28137/T)$, yielding a high-temperature heat of evaporation of 128.7 kcal/mol. Measurement of the UO₂ partial pressure over the solid gave $\log p \text{ (atm)} = 9.365 - (32436/T)$, resulting in a heat of sublimation of 148.4 kcal/mol. The difference of these heats yields the heat of fusion 19.7 kcal/mol, which is in good agreement with the literature value of 19.4 kcal/mol. In the binary UO₂ system, the pressure above the melting point was determined to be $\log p \text{ (atm)} = 7.7 - (27900/T)$, giving a heat of evaporation of 127.6 kcal/mol. An assessment of literature data for below the melting point yielded $\log p \text{ (atm)} = 8.846 - (31506/T)$, and a heat of sublimation of 144.1 kcal/mol. The resulting heat of fusion,

<475>

<475> CONT.

16.5 kcal/mol, is only slightly below the published value of
17.7 kcal/mol.

EDB SUBJECT CATEGORY: EDB-050700;360202
INIS CATEGORY: B23

<476>

Chse, R.W.; Berrie, P.G. (Commission of the European
Communities, Karlsruhe (Germany, F.R.). European Inst. for
Transuranium Elements); Brumme, G.D. (Technische Hochschule
Darmstadt (Germany, F.R.)) ** Blank, H.; Lindner, R. (eds.)

Advances in vapour pressure studies over liquid uranium
plutonium oxides up to 5000K In Plutonium 1975 and other
actinides ** AIX-08:310032;ERA-02:041845;EDB-77:094423;
(1976) North-Holland

SUBJECT DESCRIPTORS: EVAPORATION: Q1,Q2;LASER-RADIATION
HEATING;LIQUID CRYSTALS;PHASE TRANSFORMATIONS;PLUTONIUM
OXIDES: T2;PYROMETERS;ULTRAHIGH TEMPERATURE;URANIUM OXIDES:
T1;VAPOR PRESSURE: Q1,Q2;VERY HIGH TEMPERATURE

ABSTRACT: Vapour pressure measurements over liquid
($U_{sub}(0.80)Pu_{sub}(0.20)O_{sub}(1.95)$) between 4000 K and 5000 K
have been extended below 4000 K. In previous studies a radial
displacement of liquid and subsequent crater formation,
caused by the recoil force of the vapour jet during laser
heating, prevented meaningful measurements below 4000 K. This
effect could be eliminated by restricting the liquid layer
thickness to the minimum value at the final steady state
temperature, by shortening the rise time of the temperature,
using a new double intensity laser pulse heating technique
with a high initial density, and by focal spot enlargement.
The new series of measurements, applying a high-speed
multi-wavelength pyrometer, waveform analyzing and data
processing system, confirmed the previously presented
equation: $\log p(\text{atm}) = 7.966 - 28.137/T$. and heat of
evaporation $\Delta H_{sub}(\text{evap}) = 128.7 \text{ kcal/mol}$ below 4000 K.

EDB SUBJECT CATEGORY: EDB-360204
INIS CATEGORY: B24

<477>

Ohse, R.W.; Berrie, P.G.; Bogensberger, H.G.; Fischer, E.A.,
Kernforschungszentrum Karlsruhe (Germany, F.R.). Inst. fuer
Neutronenphysik und Reaktortechnik; Kernforschungszentrum
Karlsruhe (Germany, F.R.). Projekt Schneller Brueter

In Vapour pressure measurements to 7,000 K and equation of state
of oxide fuels for fast reactor safety analysis. I.
Extension of vapour pressure measurements of nuclear fuels
(U,PU) $\text{O}_{2\text{S}}$ and $\text{UO}_{2\text{S}}$ to 7,000 K **
AIX-07:278368; ERA-02:025985; EDB-77:053597; (May 1976)
KFK--2272

SUBJECT DESCRIPTORS: BINARY MIXTURES; FBR TYPE REACTORS; FUSION
HEAT; MELTDOWN; NUCLEAR FUELS; PLUTONIUM DIOXIDE: T1; REACTOR
SAFETY; SUBLIMATION HEAT; ULTRAHIGH TEMPERATURE; URANIUM
DIOXIDE: T2; VAPOR PRESSURE: Q1, Q2; VAPORIZATION HEAT

ABSTRACT: A new high-energy laser technique, including fast
temperature recording in the microsecond range, was developed
for measuring the vapour pressure of fast breeder
uranium-plutonium oxide fuels up to 7,000 K. In the ternary
system, the pressure of uranium-plutonium oxide above its
melting point was determined to be $\log p \text{ (atm)} = 7.966 -$
 $(28,137/T)$, yielding a high-temperature heat of evaporation
of 128.7 kcal/mol. Measurement of the $\text{UO}_{2\text{S}}$ partial
pressure over the solid gave $\log p \text{ (atm)} = 9.365 -$
 $(32,436/T)$, resulting in a heat of sublimation of 148.4
kcal/mol. The difference of these heats yields the heat of
fusion of 19.7 kcal/mol, which is in good agreement with the
literature value of 19.4 kcal/mol. In the binary $\text{UO}_{2\text{S}}$
system, the pressure above the melting point was determined
to be $\log p \text{ (atm)} = 7.7 - (27,900/T)$, giving a heat of
evaporation of 127.6 kcal/mol. An assessment of literature
data for below the melting point yielded $\log p \text{ (atm)} = 8.846$
 $- 31,506/T$, and a heat of sublimation of 144.1 kcal/mol. The
resulting heat of fusion of 16.5 kcal/mol, is only slightly
below the published value of 17.7 kcal/mol.

EDB SUBJECT CATEGORY: EDB-050700; 360204
INIS CATEGORY: E23

<478>

Ohse, R.W.; Berrie, P.G.; Kinsman, P.R.; Brumme, G.D.,
Commission of the European Communities, Karlsruhe (Germany,
F.R.). European Inst. for Transuranium Elements

In Vapour pressure studies over liquid uranium plutonium oxides
up to 5,000 K ** AIX-07:278369; EDB-77:050519; (1976)
AED-Conf--76-087-016

SUBJECT DESCRIPTORS: DIAGRAMS; LASER RADIATION; LIQUIDS; MELTING

<478>

<478> CONT.

PCINTS;PLUTONIUM OXIDES: T1;PRESSURE DEPENDENCE;PULSE
TECHNIQUES;TEMPERATURE DEPENDENCE;TEMPERATURE
MEASUREMENT;TERNARY ALLOY SYSTEMS;URANIUM OXIDES: T2;VAPOR
PRESSURE: Q1,Q2;VERY HIGH TEMPERATURE

ABSTRACT: The pressure over the ternary uranium-plutonium oxide was determined, yielding a heat of evaporation fully consistent with the heat of sublimation below the melting point, and with the heat of fusion. The corresponding equation for liquid UO_2 was determined with a heat of evaporation again in consistency with the assessed data below the melting point and with the heat of fusion. For this, a new high-energy laser technique, including fast temperature recording in the microsecond range, was developed. The investigation of the total temperature range from 3,000 K to $T_{\text{sub}}/c = 7,560$ K revealed two principal experimental difficulties, a liquid layer movement at temperatures below 4,000 K, and optical absorption due to increasing thermal ionisation above 5,000 K. A new double intensity laser pulse technique is presented which allows the extension of measurements to below 4,000 K. In addition a multi-wavelength pyrometer technique has been developed in order to eliminate, and at a later stage to determine, spectral emissivities above the melting point during laser heating with pulse times of micro and nanoseconds.

EDB SUBJECT CATEGORY: EDB-360204;050700
INIS CATEGORY: E23

<479>

Ohse, R.W.; Capone, P. ** Commission of the European Communities, Karlsruhe (Germany, F.R.). European Inst. for Transuranium Elements

Evaporation behaviour of the ternary uranium plutonium carbides
In 5th international conference on plutonium and other actinides ** AIX-07:233148;EDF-76:087310; (1975)
AED-Conf--75-417-018

SUBJECT DESCRIPTORS: CHEMICAL COMPOSITION;EVAPORATION:
Q1,Q2;PLUTONIUM CARBIDES: T1;SUBLIMATION HEAT;TEMPERATURE
DEPENDENCE;URANIUM CARBIDES: T2;VAPOR PRESSURE;VERY HIGH
TEMPERATURE

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-360204
INIS CATEGORY: E23

<480>

Ohse, R.W.; Capone, F. (Commission of the European Communities, Karlsruhe (F.R. Germany). European Inst. for Transuranium Elements)

Evaporation behaviour of the ternary uranium plutonium carbides
In Reactor conference, Duesseldorf, 30.3.-2.4.1976. Section
2: Fuel elements, fuel cycle and isotope technique **
AIX-07:242008;EDB-76:092583; (1976) ZAED

SUBJECT DESCRIPTORS: COMPATIBILITY;HEAT TREATMENTS;HEATING;MASS
SPECTROSCOPY;MIXTURES;NUCLEAR FUELS;PHASE DIAGRAMS;PLUTONIUM
CARBIDES: T1;SOLID SOLUTIONS;SUBLIMATION HEAT:
Q1,Q2;TEMPERATURE DEPENDENCE;TUNGSTEN;URANIUM CARBIDES:
T2;VAPOR PRESSURE;VERY HIGH TEMPERATURE

ABSTRACT: Simultaneous measurement of total and partial
pressures; variation of plutonium pressure with $1/T$ in the
single phase region MC from $C/M = 1.014$ to 1.15 and two-phase
region $MC + M_{2/3}C_{3/3}$ from $C/M = 1.15$ to 1.30 from
 $2,000$ K to $2,500$ K and determination of the corresponding
sublimation heat from the measured values; pressure
composition diagram of the ternary (U,Pu) carbides giving the
phase boundaries between MC and $MC + M_{2/3}C_{3/3}$ phase
fields.

EDB SUBJECT CATEGORY: EDB-360204;050700
INIS CATEGORY: B23

<481>

Ohse, R.W.; Capone, F. (Commission of the European Communities, Karlsruhe (Germany, F.R.). European Inst. for Transuranium Elements) ** Blank, H.; Lindner, R. (eds.)

Evaporation behaviour of the ternary uranium plutonium carbides
In Plutonium 1975 and other actinides **
AIX-08:310033;ERA-02:041846;EDB-77:094424; (1976)
North-Holland

SUBJECT DESCRIPTORS: CHEMICAL
COMPOSITION;COMPATIBILITY;EVAPORATION:
Q1,Q2;IMPURITIES;KNUDSEN FLOW;MASS
SPECTROSCOPY;MICROSTRUCTURE;PHASE STUDIES;PLUTONIUM CARBIDES:
T1;SEGREGATION;SUBLIMATION;SUBLIMATION HEAT;TEMPERATURE
DEPENDENCE;URANIUM CARBIDES: T2;VAPOR PRESSURE: Q1,Q2;VERY
HIGH TEMPERATURE

ABSTRACT: The evaporation behaviour of uranium plutonium
carbides ($U_{0.80}Pu_{0.20}C_{1-x}$) was studied by a
combined application of mass spectrometry, using the uranium
isotope U-233, and the Knudsen effusion target collection
technique in the temperature range from $1500^{\circ}C$ to the

<481>

<481> CONT.

liquids temperature measured at 2458°C and the composition range from $\text{C/M} = 0.95$ to 1.4 . High temperature compatibility tests were made with W-cells, carburized Ta and TaC-liners up to 2500°C . The influence of oxygen and nitrogen impurities on vapour pressure, and composition changes in continued evaporation of the mixed carbides were investigated. The effects of plutonium depletion and segregation were studied.

EDB SUBJECT CATEGORY: EDB-360204;C50700
INIS CATEGORY: B24

<482>

Ohse, R.W.; Ciani, V.

Evaporation behavior and high-temperature thermal analysis of substoichiometric plutonium oxide in the composition range from $\text{O/Pu} = 1.51$ to 2.00 in high temperature technology **
EDE-77:120461; (1969) Butterworths

SUBJECT DESCRIPTORS: CHEMICAL PREPARATION;EVAPORATION:
Q1;PLUTONIUM DIOXIDE: T1;STOICHIOMETRY;THERMAL ANALYSIS;VAPOR
PRESSURE;VERY HIGH TEMPERATURE;X-RAY DIFFRACTION

ABSTRACT: None.

EDB SUBJECT CATEGORY: EDB-360204;C50700

<483>

Ohwada, K. (Japan Atomic Energy Research Inst., Tokai, Ibaraki.
Tokai Research Establishment)

IR spectroscopic study of dipotassium and dicesium uranyl
tetrabromides AIX-08:346258;EDB-78:028036; J. Inorg. Nucl.
Chem. (JINCA) 39: (9):1601-1605 (1977)

SUBJECT DESCRIPTORS: CESIUM BROMIDES;CHEMICAL BONDS: Q1;INFRARED
SPECTRA: Q1;POTASSIUM BROMIDES;URANYL COMPOUNDS:
T1;VALENCE;VIBRATIONAL STATES

ABSTRACT: The IR spectra of dipotassium and dicesium uranyl
tetrabromides ($\text{K}_2\text{UO}_2\text{Br}_4$, $\text{Cs}_2\text{UO}_2\text{Br}_4$) have been measured in the region from 4000 down
to 30 cm^{-1} . A normal coordinate analysis of the
(UO_2Br_4) $^{2-}$ ion apart from the K^+ and Cs^+
 Cs^+ ions has been made using Wilson's FG matrix method.
The force constants associated with the U-O and U-Br bonds
have been obtained on the basis of the two force models of
modified Urey-Bradley and valence force fields. An effort
has been directed towards the understanding of a negative

<483> CONT.

force constant of the U-O bond-bond interaction which plays an important role in the calculations of symmetric and asymmetric U-O vibrations, and detailed discussion has been made on it from the point of view of the mutual polarizability between the two U-O bonds.

EDB SUBJECT CATEGORY: EDB-400702
INIS CATEGORY: B13

<484>

Olson, C.G.; Piacentini, M.; Lynch, D.W. ** Koch, E.E. (ed.)

Optical properties of rare earth trifluorides in vacuum
ultraviolet radiation physics ** ERA-01:023970; EDB-76:066061;
(1974) Friedr. Vieweg and Sohn Verlagsgesellschaft mbH

SUBJECT DESCRIPTORS: CERIUM FLUORIDES: T2; DYSPROSIUM FLUORIDES:
T4; ENERGY-LEVEL TRANSITIONS; FAR ULTRAVIOLET
RADIATION; GADOLINIUM FLUORIDES: T5; LANTHANUM FLUORIDES:
T1; MONOCRYSTALS; PRASEODYMIUM FLUORIDES: T3; REFLECTIVITY:
Q1, Q2, Q3, Q4, Q5; SPECTRA

ABSTRACT: The reflectivity spectra of single crystals of the trifluorides of La, Ce, Pr, and Dy and of polycrystalline GdF_3 were measured in the 4 to 30 eV range using synchrotron radiation. The transmission of thin films of these salts on LiF substrates was measured. (GHT)

EDB SUBJECT CATEGORY: EDB-360204; 640302

<485>

Olson, J.J., California Univ., San Diego, La Jolla (USA). Dept.
of Physics

In KKR analysis of the Anderson-McMillan theory of electronic structure of liquid metals. Technical report **
ERA-02:010428; EDB-77:007024; (Apr 1976) AD-A--024386

SUBJECT DESCRIPTORS: BAND THEORY; DISPERSION RELATIONS; ELECTRONIC
STRUCTURE: Q1; LIQUID METALS: T1; SOLIDS

ABSTRACT: The method of Anderson and McMillan (AM) has recently been demonstrated to fail for resonant scatterers while nevertheless providing a useful band structure approximation for non-resonant scatterers. The authors explain this behavior by using the formalism of Korringa, Kohn, and Rostoker (KKR) to relate the AM method to well known approximations in the band theory of crystalline solids. In particular the authors show that the AM method for the

<485>

<485> CONT.

dispersion relation of a liquid metal is essentially an optimized single-orthogonalized-plane-wave approximation to the band structure of a crystal at the same density. (GRA)

EDB SUBJECT CATEGORY: EDB-360104

<486>

Oosterbroek, M.; van de Braak, H.P.; Gellings, P.J. (Technische Hogeschool Twente, Enschede (Netherlands)) ** Evans, R.; Greenwood, D.A. (eds.)

Improved model for solutions of nonmetals in liquid alkali metals. Calculation of enthalpy of solution and electrical resistivity In Liquid metals, 1976 **

AIX-08:320690;EDB-77:120825; (1977) Institute of Physics

SUBJECT DESCRIPTORS: ALKALI METALS: T1;ELECTRIC CONDUCTIVITY;LIQUID METALS;MATHEMATICAL MODELS;NONMETALS: T2;SOLUTION HEAT: Q2;SOLUTIONS;SOLVENT PROPERTIES: Q1

ABSTRACT: A new model is proposed for the calculation of the enthalpy of solution of electronegative elements and noble gases in liquid alkali metals. This is based on the difference between the internal energies of the pure metal and of the solution. Contributions to those energies arise from free-electron terms and the Madelung energy, expressed in terms of simple model potentials and partial structure factors. Screening is accounted for by the band-structure energy. Satisfactory agreement between experimental and calculated values is obtained using pure metal empty-core radii and crystal radii for the dissolved atoms in the model potentials and adjusting hard-sphere diameters in Percus-Yevick calculations for the structure factors. The model is able to explain the positive enthalpy of solution of the noble gases. Volume effects give an important contribution to corrections to the Madelung energy. The volume dependence of the electron-gas and band-structure terms are relatively small. The electrical resistivity is obtained from Ziman's formula using identical structure factors and form factors based on the model potentials as used in the enthalpy calculations. Reasonable values for the resistivity change on dissolving the nonmetal are obtained but the nearly complete lack of experimental data makes verification practically impossible at this moment.

EDB SUBJECT CATEGORY: EDB-400201;360104
INIS CATEGORY: B12

<487>

Osborne, D.W.; Flotow, H.E.

Half-life of ^{242}Pu from precise low-temperature heat capacity measurements

INS-76:020557; ERA-02:002060; EDB-76:083721; Phys. Rev., C (PRVCA) 14: (3): 1174-1178 (Sep 1976)

SUBJECT DESCRIPTORS: ALPHA DECAY: Q1; HALF-LIFE: LOW TEMPERATURE; MEDIUM TEMPERATURE; PLUTONIUM 242: T1; PLUTONIUM FLUORIDES: T3; PLUTONIUM OXIDES: T2; SPECIFIC HEAT: Q2, Q3; ULTRALOW TEMPERATURE; VERY LOW TEMPERATURE

ABSTRACT: Heat capacity measurements on three pure compounds, PuF_3 , PuF_4 , and PuO_2 , made from 99.91 at. % ^{242}Pu , have been used to evaluate the power produced by radioactive decay. The mean specific power was found to be (0.12099 ± 0.00024) mW per gram of plutonium in the samples. The contribution of nuclides other than ^{242}Pu to the power was calculated to be (5.15 ± 0.11) μW per gram of plutonium, and so the specific power of pure ^{242}Pu is (0.11594 ± 0.00026) mW g $^{-1}$. A correction for the (5.50×10^{-4}) % spontaneous fission branching reduces the specific power due to α decay to (0.11591 ± 0.00026) mW g $^{-1}$. Combining this value of the specific power with the total disintegration energy of (4.9823 ± 0.0012) MeV, we obtain (3.763 ± 0.009) x 10^5 years as the α half-life of pure ^{242}Pu . (AIP)

EDB SUBJECT CATEGORY: EDB-652022; 360104
INIS CATEGORY: A34; B22

<488>

Oyamada, R. (Nagasaki Univ. (Japan). Faculty of Education); Yoshida, S.

Raman spectra of solid ThCl_4 and fused ThCl_4 --KCl system AIX-07:266529; EDB-77:007484; Nippon Seirigaku Zasshi (NISEA) 38: (6): 1786 (Jun 1975)

SUBJECT DESCRIPTORS: ANIONS; CHEMICAL SHIFT; COMPLEXES; LASERS; MOLTEN SALTS; POTASSIUM CHLORIDES: T1; RAMAN EFFECT: Q1, Q2; SPECTRA; SPECTRAL SHIFT; THORIUM CHLORIDES: T2

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-400201
INIS CATEGORY: A13

<489>

<489>

Papatheodrou, G.N.

Raman spectroscopic studies of yttrium (III) chloride--alkali metal chloride melts and of $\text{Cs}_2\text{NaYCl}_6$ and YCl_3 solid compounds INS-77:008015; ERA-02:032799; EDB-77:074216; J. Chem. Phys. (JCPA) 66:(7):2893-2900 (1 Apr 1977)

SUBJECT DESCRIPTORS: ALKALI METAL COMPOUNDS: T2; BINARY MIXTURES: Q1, Q2, T3; CHLORIDES; EMISSION SPECTRA; MOLTEN SALTS; RAMAN EFFECT: Q1, Q2, Q3; YTTRIUM CHLORIDES: T1

ABSTRACT: The Raman spectra of molten YCl_3 --A Cl (A=Cs, K, Li) mixtures have been measured at different compositions and temperatures up to 890 degreeC. The Raman spectra of polycrystalline $\text{Cs}_2\text{NaYCl}_6$ and YCl_3 were also measured from 25 degreeC to temperatures above melting. The factor group analysis of crystalline $\text{Cs}_2\text{NaYCl}_6$ was used to identify the three Raman active modes (ν_{1g} , ν_{2g} , ν_{5g}) of the YCl_6^{3-} octahedra. For liquid mixtures rich in alkali halide, the predominant features of the spectra are characterized by a polarized and a depolarized band with frequencies near the ν_{1g} and ν_{5g} frequencies of the YCl_6^{3-} octahedron and thus indicates the existence of such species in the melt. In melts containing above 25% YCl_3 a new polarized band D appears in the spectra which shifts continuously and rapidly to higher energies with increasing YCl_3 concentration. The frequency shift is attributed to a continuous distortion mechanism of the YCl_6^{3-} octahedra by the neighboring yttrium ions. The continuous frequency shift of the D band and a comparison of the liquid and solid Raman spectra of yttrium chloride suggest the existence of lattice-type modes in these melts.

EDB SUBJECT CATEGORY: EDB-360000;360602

INIS CATEGORY: B20;B24

<490>

Partashnikova, M.Z.; Oboznenko, V.A.; Khrepinyuk, T.E.

Differential-spectrophotometric determination of niobium in its compounds EDB-76:054265; Zh. Anal. Khim. (ZAKHA) 31:(1):187-189 (Jan 1976)

SUBJECT DESCRIPTORS: ALKALI METAL COMPOUNDS; ALKALINE EARTH METAL COMPOUNDS; CHEMICAL ANALYSIS: Q2; NIOBATES: T2; NIOBIUM: T1; QUANTITATIVE CHEMICAL ANALYSIS; RARE EARTH COMPOUNDS; SPECTROPHOTOMETRY: Q1

ABSTRACT: A method has been developed of differential-spectrophotometric determination of niobium in

<490> CONT.

niobates of alkali, alkaline earth, and rare earth metals from absorbance of a peroxide complex of niobium in a medium of concentrated sulfuric acid.

EDE SUBJECT CATEGORY: EDB-400104

<491>

Fathak, N.L.; Sen, S.C. (Indian Inst. of Tech., Kanpur. Dept. of Physics)

Effect of luminescence centers on optical absorption, excitation and fluorescence spectra of CsCl:Pb

AIX-07:236142; EDB-76:054157; Indian J. Pure Appl. Phys. (IJOPA) 13: (5):288-291 (May 1975)

SUBJECT DESCRIPTORS: ABSORPTION; BAND THEORY; CESIUM CHLORIDES: T1; COLOR CENTERS; ELECTRONIC STRUCTURE; EMISSION SPECTRA; ENERGY-LEVEL TRANSITIONS; EXCITATION; FLUORESCENCE: Q1, Q2; LEAD CHLORIDES: T2; PHOSPHORS

ABSTRACT: The optical absorption, excitation and fluorescence spectra of CsCl containing lead of different concentrations have been studied. Absorption bands have been observed at approximately 5.55 and 4.40 eV at room temperature as well as at liquid nitrogen temperature. Four fluorescence bands have been observed at approximately 4.00, 3.50, 3.07 and 2.15 eV at liquid nitrogen temperature, being efficiently excited by their excitation bands at approximately 4.67, 4.40, 4.72 and 4.28 eV, respectively. The different excitation and fluorescence bands have been attributed to the different luminescence centers, i.e. Pb^{2+} -vacancy pairs (dipoles), trimers and pentamers.

EDE SUBJECT CATEGORY: EDB-360605

INIS CATEGORY: B12

<492>

Peacock, R.D. (King's Coll., London (UK). Dept. of Chemistry) **
 Dunitz, J.D. (ed.) (Eidgenossische Technische Hochschule, Zurich (Switzerland). Lab. fuer Organische Chemie);
 Hemmerich, P. (ed.) (Konstanz Univ. (F.R. Germany). Fachbereich Biologie); Holm, R.H. (ed.) (Massachusetts Inst. of Tech., Cambridge (USA). Dept. of Chemistry); Ibers, J.A. (ed.) (Northwestern Univ., Evanston, Ill. (USA). Dept. of Chemistry); Joergensen, C.K. (ed.) (Geneva Univ. (Switzerland). Dept. de Chimie Minerale et Analytique); Neillands, J.B. (ed.) (California Univ., Berkeley (USA). Dept. of Biochemistry); Reinen, D. (ed.) (Marburg Univ. (F.R. Germany). Fachbereich Chemie); Williams, R.J.P. (ed.) (Oxford Univ. (UK). Inorganic Chemistry Lab.)

<492>

<492> CONT.

Intensities of lanthanide f-f transitions In Rare earths **
 Structure and Bonding, AIX-07:257201;EDB-77:007478; (1975)
 Springer

SUBJECT DESCRIPTORS: COUPLING;DIPOLE MOMENTS;ELECTRIC
 DIPOLES;ELECTRONIC STRUCTURE;ENERGY-LEVEL TRANSITIONS: Q1;P
 STATES;MATHEMATICS;MATRICES;PRASEODYMIUM COMPOUNDS;QUADRUPOLE
 MOMENTS;RARE EARTH COMPOUNDS;RARE EARTHS:
 T1;REVIEWS;SPECTRA;TENSORS

ABSTRACT: Forced electric dipole transitions and the Judd-Cfelt
 theory, other multipole transitions, the success of the
 Judd-Cfelt theory, hypersensitive transitions, the
 exceptional behaviour of praseodymium(III), the $5D_{05}$ - $7F_{05}$ transitions of europium(III), and
 structural implications are treated.

EDB SUBJECT CATEGORY: EDB-400201
 INIS CATEGORY: B12

<493>

Permyakov, P.G.; Korshunov, B.G.; Krokhin, V.A. (Moskovskij
 Inst. Stali i Splavov (USSR); Gosudarstvennyj Inst. Redkikh
 Metallov, Moscow (USSR))

Lanthanum chloride interaction with magnesium oxide
 AIX-07:277861;EDB-77:031066; Izv. Vyssh. Uchebn. Zaved.,
 Tsvet. Metall. (IVUTA) (3):93-95 (1975)

SUBJECT DESCRIPTORS: CHEMICAL REACTIONS;DIFFERENTIAL THERMAL
 ANALYSIS;LANTHANUM CHLORIDES: T1;MAGNESIUM OXIDES: T2;PHASE
 DIAGRAMS: Q1,Q2;QUANTITY RATIO;SOLUBILITY

ABSTRACT: Chlorides of rare-earth elements, obtained as a result
 of chlorination of complex ores, enter into complex reactions
 with the oxygen-containing components of the charge and
 refractories. This work was aimed, in particular, at studying
 the interaction of components of a ternary system $LaCl_3$ +
 MgO reversible $LaOCl$ + $MgCl_2$. The study was based on
 differential-thermographic analysis. The melting-point curve
 of this ternary system is given. It is concluded that the
 lanthanum chloride formed as a result of chlorination of a
 raw material containing rare-earth elements may interact with
 MgO and, most probably, with other basic oxides forming part
 of the charge or refractories. The lanthanum hydroxychloride
 formed in the reaction is highly soluble in lanthanum and
 magnesium chlorides. A higher concentration of magnesium
 chloride in the melt enhances the solubility of $LaOCl$, which

<493> CCNT.

should be taken into account when selecting the composition of the melt for chlorination.

EDB SUBJECT CATEGORY: EDB-360602
INIS CATEGORY: B12

<494>

Permyakov, P.G.; Korshunov, B.G.; Krokhin, V.A. (Moskovskii Inst. Stali i Splavov (USSR); Gosudarstvennyi Inst. Redkikh Metallov, Moscow (USSR))

Interaction of neodymium- and yttrium chlorides with magnesium oxide AIX-07:265089; EDB-77:031426; Zh. Neorg. Khim. (ZNOKA) 20:(7):1965-1969 (Jul 1975)

SUBJECT DESCRIPTORS: CHEMICAL REACTIONS: Q1,Q2,Q3; DIFFERENTIAL THERMAL ANALYSIS; MAGNESIUM OXIDES: T1; NEODYMIUM CHLORIDES: T2; PHASE DIAGRAMS; X-RAY DIFFRACTION; YTTRIUM CHLORIDES: T3

ABSTRACT: By differential-thermographic and roentgenographic analysis interaction of neodymium and yttrium chlorides with magnesium oxide was studied. Diagrams of reversible system $\text{NdCl}_3 + \text{MgO} \rightleftharpoons \text{NdOCl} + \text{MgCl}_2$, $\text{YCl}_3 + \text{MgO} \rightleftharpoons \text{YCl}_2 + \text{MgCl}_2$ fusibility were plotted. Compound of the assumed composition $3\text{MgCl}_2 \cdot 3\text{x}2\text{NdOCl}$ is formed in the MgCl_2 - NdOCl system.

EDB SUBJECT CATEGORY: EDB-400201
INIS CATEGORY: B12

<495>

Permyakov, P.G.; Korshunov, B.G.; Krokhin, V.A. (Moskovskij Inst. Stali i Splavov (USSR); Gosudarstvennyj Inst. Redkikh Metallov, Moscow (USSR))

Solubility of rare earth and yttrium oxochlorides in molten salts - MCl_2 (M-Mg, Ca, Sr, Ba) AIX-08:298331; EDB-77:069031; Zh. Neorg. Khim. (ZNOKA) 20:(8):2184-2187 (Aug 1975)

SUBJECT DESCRIPTORS: CALCIUM COMPOUNDS: M1; DIFFERENTIAL THERMAL ANALYSIS; LANTHANUM COMPOUNDS: M2; MOLTEN SALTS; NIOBATES: M3; OXYGEN COMPOUNDS; PHASE DIAGRAMS: Q1,Q2,Q3; RARE EARTH COMPOUNDS; SATURATION; SOLUBILITY; STRONTIUM CHLORIDES

ABSTRACT: The results of phase equilibria studying in the $\text{CaO} \cdot 2\text{La}_2\text{O}_3$ - La_2O_3 system in the temperature range from 1.250 deg C to the melting point are summarized in this paper. A phase diagram of the system has been constructed from the results of X-ray and

<495>

<495> CONT.

crystal-optical analyses, which reveals its eutectic nature with a melting point of 1.345 ± 10 deg C at 36 mol.% $\text{LaNb}_{0.9}$ and 64 mol.% $\text{CaNb}_{2.6}$, as well as limited solid solutions based on the starting compounds with the following content of the dissolved components: 10 ± 2.5 mol.% $\text{LaNb}_{0.9}$ and 25 ± 2.5 mol.% $\text{CaNb}_{2.6}$.

EDF SUBJECT CATEGORY: EDB-400201

INIS CATEGORY: B12

<496>

Permyakov, P.G.; Korshunov, B.G.; Krokhin, V.A. (Moskovskij Inst. Stali i Splavov (USSR); Gosudarstvennyj Nauchno-Issledovatel'skij i Proektnyj Inst. Redkometallicheskoj Promyshlennosti, Moscow (USSR))

Interaction of thorium chloride with calcium oxide

AIX-08:295591; EDB-77:069323; Izv. Vyssh. Uchebn. Zaved., Tsvet. Metall. (IVUTA) (3):95-97 (1976)

SUBJECT DESCRIPTORS: CALCIUM OXIDES: T1; CHEMICAL REACTIONS: Q1, Q2; DIFFERENTIAL THERMAL ANALYSIS; EUTECTICS; FREE ENERGY; MOLTEN SALTS; PHASE DIAGRAMS; QUANTITY RATIO; THORIUM CHLORIDES: T2; VERY HIGH TEMPERATURE; X-RAY DIFFRACTION

ABSTRACT: None

EDF SUBJECT CATEGORY: EDB-400702

INIS CATEGORY: B16

<497>

Perov, P.A.; Nedyak, S.V.; Mal'tsev, A.A. (Moskovskii Gosudarstvennyi Univ. (USSR). Kafedra Fizicheskoi Khimii)

Investigation into absorption infrared spectra of vapors under $\text{ScEr}_{0.3}$, $\text{YCl}_{0.3}$, $\text{LaCl}_{0.3}$, $\text{GdCl}_{0.3}$, $\text{ErCl}_{0.3}$, $\text{LuCl}_{0.3}$ by means of isolation method in rare gas matrix AIX-07:265090; EDB-77:031416; Vestn. Mosk. Univ., Ser. II. Khim. (VMUKA) 16:(3):281-283 (May 1975)

SUBJECT DESCRIPTORS: ABSORPTION SPECTRA; COMPARATIVE EVALUATIONS; ERBIUM CHLORIDES: T5; FREQUENCY DEPENDENCE; GADOLINIUM CHLORIDES: T4; INFRARED SPECTRA: Q1, Q2, Q3, Q4, Q5, Q6; LANTHANUM CHLORIDES: T3; LUTETIUM: T6; RARE GASES; SCANDIUM BROMIDES: T1; VERY LOW TEMPERATURE; VIBRATIONAL STATES; YTTRIUM CHLORIDES: T2

ABSTRACT: Infra red spectra of $\text{ScBr}_{0.3}$, $\text{YCl}_{0.3}$, $\text{LaCl}_{0.3}$, $\text{GdCl}_{0.3}$, $\text{ErCl}_{0.3}$ and $\text{LuCl}_{0.3}$ molecules in the

<497> CONT.

inert gas matrix at 20-30⁰ K was studied. Comparison of the results with published data shows that frequencies ν_2 , ν_3 , ν_4 , of the Sc, Y, La, trihalogen series, are gradually decreasing from Sc-trihalogen to La-trihalogen. In the lanthanum trihalogen series (Gd....La) tendency was observed of the frequency increasing with the increasing of the atomic weight of the metal central atom. Estimation of the oscillation frequencies for the trihalogens of scandium, yttrium, lanthanum and some lanthanide trichlorides experimentally not investigated was made by comparative method.

EDB SUBJECT CATEGORY: EDB-400201
INIS CATEGORY: B12

<498>

Ferrin, A. (Rennes-1 Univ., 35 (France))

Preparational, structural and vibrational study of $M_2U_2O_5Cl_4$, $2H_2O$ (M = Rb, Cs). Detection of a tetranuclear anion $[(UO_2)_2]^{4+} [Cl_8]^{8-} (H_2O)_2$ ⁴⁻ AIX-08:337074; EDB-77:141188; J. Inorg. Nucl. Chem. (JINCA) 39:(7):1169-1172 (1977)

SUBJECT DESCRIPTORS: CESIUM CHLORIDES; CHEMICAL BONDS: Q1; CHEMICAL PREPARATION: Q1; INFRARED SPECTRA; MOLECULAR STRUCTURE; RAMAN SPECTRA; RUBIDIUM CHLORIDES; URANYL COMPLEXES: T1; X-RAY DIFFRACTION

ABSTRACT: The interaction between $M_2U_2O_5Cl_4$. nH_2O (M = Rb, Cs) and $UO_2(OH)_2$ leads to new complexes $M_2U_2O_5Cl_4$, $2H_2O$; however, structural determination by X-ray diffraction exhibits discrete tetranuclear anions $[(UO_2)_2]^{4+} [Cl_8]^{8-} (H_2O)_2$ ⁴⁻: uranyl groups are bonded together in their equatorial plane by O/O or O/Cl double bridges in which oxygen atom is tricoordinated with U-O (bridge) distances lying between 2.20 and 2.29 Å. Vibrational spectra are recorded and interpreted and hydrogen bonds discussed.

EDB SUBJECT CATEGORY: EDB-400702
INIS CATEGORY: B13

<499>

<499>

Pervov, V.S. (AN SSSR, Moscow. Inst. Obshchej i Neorganicheskoy Khimii)

Energies of valence states of atoms in halides
AIX-08:336995;EDB-78:009992; Koord. Khim. (KOKHD)
2:(9):1164-1171 (Sep 1976)

SUBJECT DESCRIPTORS: ATOMIC MODELS;BERYLLIUM FLUORIDES:
T1;BINDING ENERGY: Q1,Q2,Q3,Q4,Q5,Q6,Q7,Q8;BORON FLUORIDES:
T2;ELECTRONIC STRUCTURE;FORMATION HEAT;GASES;MOLYBDENUM
CHLORIDES: T3;NIOBIUM CHLORIDES: T4;STRONTIUM FLUORIDES:
T5;TANTALUM CHLORIDES: T6;TUNGSTEN FLUORIDES:
T7;VALENCE;VANADIUM CHLORIDES: T8

ABSTRACT: Variation in energy of valence states (E) in alternate-valence series involving the elements of the groups 5-7 of the periodic system has been first considered on the basis of review of experimental data on the formation heat of gaseous halides (Be, Sr, B, W fluorides and Mo, V, Ta, Nb chlorides). The dependences are compared with calculation data for E. The principal reasons of non-monotonous variation of E in alternate-valence series of multivalence metals are discussed.

EDB SUBJECT CATEGORY: EDB-400201
INIS CATEGORY: B12

<500>

Pervov, V.S.; Nikolaev, N.C. (AN SSSR, Moscow. Inst. Obshchej i Neorganicheskoy Khimii)

Progress in fluorine calorimetry AIX-08:314936;EDB-77:101149;
Usp. Khim. (USKHA) 45:(4):640-660 (Apr 1976)

SUBJECT DESCRIPTORS: BERYLLIUM;CADMIUM;CALORIMETRY: Q1;CHEMICAL REACTIONS: Q2;FLUORINATION: T1;FLUORINE: T2;FORMATION HEAT;MEASURING METHODS;REVIEWS;TRANSITION ELEMENTS;URANIUM

ABSTRACT: The development of a new area of thermochemistry (the calorimetry of fluorination reactions) is reviewed. The high reactivity of fluorine offers some important advantages over alternative reagents, but at the same time it introduces some experimental difficulties. We have attempted to generalise the practical problems of fluorine calorimetry by systematically examining a number of specific thermochemical investigations by the most widely adopted modification of the method: fluorine bomb calorimetry. Current developments and

<500> CONT.

possible new applications are reviewed, and the basic experimental procedures and apparatus are described.

EDB SUBJECT CATEGORY: EDB-400201

INIS CATEGORY: E12

<501>

Peterson, J.R., Tennessee Univ., Knoxville (USA). Dept. of Chemistry

In Physical--chemical studies of transuranium elements.

Progress report, April 1, 1976--March 31, 1977 **

INS-77:006516; ERA-02:027159; EDB-77:056434; (1977)

ORO--4447-048

SUBJECT DESCRIPTORS: BERKELIUM: T6; BERKELIUM CHLORIDES: T1; CALIFORNIUM BROMIDES: T5; CALIFORNIUM CHLORIDES: T2; CHEMICAL PROPERTIES; EINSTEINIUM BROMIDES: T4; EINSTEINIUM CHLORIDES: T3; PHYSICAL PROPERTIES; RESEARCH PROGRAMS: Q7; STRUCTURAL CHEMICAL ANALYSIS: Q1, Q2, Q3, Q4, Q5, Q6; TRANSURANIUM ELEMENTS: T7

ABSTRACT: Major advances in our continuing program to determine, interpret, and correlate the basic chemical and physical properties of the transuranium elements are summarized for the period April 1, 1976, through March 31, 1977. Implementation of data reduction programs and acquisition of a CRT time-sharing graphics terminal/stand-alone computer have advanced significantly the handling capabilities of single-beam spectral data obtained by our microscope spectrophotometer. EsCl_3 , EsBr_3 , and EsI_3 have been well characterized spectroscopically, and limited X-ray diffraction data have been obtained from EsBr_3 and EsI_3 . The reduction of mixed Es-lanthanide trihalides has produced what might be Es(II) . Dimorphism in BkCl_3 , CfCl_3 , and BkBr_3 has been studied spectrophotometrically and the results confirmed by X-ray analysis. Our solution microcalorimeter was improved by reducing the system-generated electrical noise and developing a novel sample container. The operating sensitivity was determined to be within the desired 0.1 percent precision requirement. The necessary hardware was obtained and software development was initiated for the capability to acquire, store, and analyze the heat of solution data automatically. A very sensitive apparatus (SQUID) for the determination of magnetic susceptibility has been constructed and is being evaluated. Our device should greatly facilitate the study of

<501>

<501> CONT.

microgram-sized samples of transuranium elements and compounds.

EDE SUBJECT CATEGORY: EDB-400702

INIS CATEGORY: B13

<502>

Pfann, J. (Ústav Jaderneho Vyzkumu, Rez (Czechoslovakia))

New description of liquid metal heat transfer in closed conduits

AIX-08:332735; EDB-78:022544; Nucl. Eng. Des. (NEDEA)

41: (2):149-163 (Apr 1977)

SUBJECT DESCRIPTORS: BOUNDARY CONDITIONS; CIRCULAR CONFIGURATION; CONTINUITY EQUATIONS; ERRORS; FRICTION; HEAT TRANSFER: Q1; INTEGRALS; LAMINAR FLOW; LIQUID FLOW; LIQUID METALS: T1; NUSSELT NUMBER; PRANDTL NUMBER; ROUGHNESS; THERMAL CONDUCTIVITY; TUBES; TURBULENT FLOW; VELOCITY; VISCOSITY

ABSTRACT: Heat transfer with liquid metals in conduits with fully developed flow is analyzed. Four typical regions are distinguished for which different equations hold. The influence of the Prandtl number and of the relative roughness are investigated. Physical conclusions are illustrated for circular tubes and computed results compared with liquid metal heat transfer formulae currently in use. A new formula for the friction factor is proposed.

EDE SUBJECT CATEGORY: EDB-420400

INIS CATEGORY: E11

<503>

Fialoux, A.; Lode, M.

In Contribution to the establishment of the phase diagram for the U--O--C system by means of x-ray diffractometry at high temperatures and under controlled pressure **

ERA-03:001162; EDB-78:009578; J. Nucl. Mater. (JNUMA)

56::221-228 (1975) ORNL-tr--4327

SUBJECT DESCRIPTORS: GRAPHITE; PHASE DIAGRAMS: Q1, Q2; REDUCTION; URANIUM CARBIDES: M2; URANIUM DIOXIDE; URANIUM OXIDES: M1; VERY HIGH TEMPERATURE; X-RAY DIFFRACTION

ABSTRACT: During the progressive reduction of uranium dioxide by graphite under known pressures of carbon monoxide, we have measured the lattice parameters of the compounds UO_{2-x} (fcc), $\alpha-U_2O_3$ (tetragonal) and UO_3 (fcc), and have evaluated the compositions of these nonstoichiometric phases near 1770°C. By this means,

<503> CONT.

and for the first time thanks to these direct measurements at such high temperatures, it has been possible to establish with precision the range of the main univariant phase fields [UO_2 , C, $\alpha\text{-UC}_2$, CO] and [UO_2 , $\alpha\text{-UC}_2$, 'UC', CO]. The part of the U-C-O diagram thus mapped out completes the earlier isotherm for 1700°C determined by Henry et al. 48 refs.

EDB SUBJECT CATEGORY: EDB-360202;C50700

<504>

Flakhctnik, V.N.; Steba, V.K. (Dnepropetrovskij Inst. Inzhenerov Zheleznodorozhnogo Transporta (Ukrainian SSR))

Influence of salt background on kinetics of BF_4 -alkali hydrolysis AIX-08:339958;EDB-78:017360; Izv. Vyssh. Uchebn. Zaved., Khim. Khim. Tekhnol. (IVUKA) 20:(2):306 (1977)

SUBJECT DESCRIPTORS: ACTIVATION ENERGY;ANIONS;BORON FLUORIDES: T1;CHEMICAL REACTION KINETICS: Q2,Q3;CORRELATIONS;ENTROPY;HYDROLYSIS: Q1;MEDIUM TEMPERATURE;QUANTITY RATIO;SODIUM CHLORIDES: T2;SODIUM PERCHLORATES: T3

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-400201

INIS CATEGORY: B12

<505>

Podafa, E.P.; Dubovoi, P.G.

Thermodynamic analysis of melting point diagrams of systems consisting of alkali metal chlorides, thallium chloride and beryllium chloride AIX-07:277868;EDB-77:031091; Ukr. Khim. Zh. (UKZHA) 42:(4):430-432 (Apr 1976)

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-360602;400201

INIS CATEGORY: B12

<506>

<506>

Podafa, B.P.; Dubovoi, P.G.; Barchuk, V.T. (AN Ukrainskoj SSR,
Kiev, Inst. Obshchej i Neorganicheskoy Khimii)

Meltability diagram of RbCl-BeCl₂ system

AIX-08:286145; EDB-77:044215; Zh. Neorg. Khim. (ZNOKA)
20:(8):2289-2290 (Aug 1975)

SUBJECT DESCRIPTORS: BERYLLIUM CHLORIDES: T1; BINARY
MIXTURES; CHEMICAL ANALYSIS; CHEMICAL COMPOSITION; DIFFERENTIAL
THERMAL ANALYSIS; EUTECTICS; MELTING POINTS; MOLTEN SALTS; PHASE
DIAGRAMS: Q1, Q2; RUBIDIUM CHLORIDES: T2

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-400201
INIS CATEGORY: B12

<507>

Eclart, J.; Degobert, P.; Bloch, O.

Intensity-voltage curves of hydrogen diffusion electrodes in
eutectic LiCl--KCl EDB-76:064854; C. R. Hebd. Seances Acad.
Sci. (COREA) 255:(3):515-517 (16 Jul 1962)

SUBJECT DESCRIPTORS: ANODES; ELECTRICAL
PROPERTIES; ELECTROCHEMISTRY; ELECTROLYTES:
Q1; EUTECTICS; HYDROGEN FUEL CELLS: T1; LITHIUM
CHLORIDES; POTASSIUM CHLORIDES

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-300503

<508>

Politis, C., Kernforschungszentrum Karlsruhe (Germany, F.R.).
Inst. fuer Material- und Festkuerperforschung

In Thermal lattice dilatation of some transition metal compounds
** AIX-07:230587; EDB-76:087319; (Aug 1975) KFK--2168

SUBJECT DESCRIPTORS: DATA; DIAGRAMS; DILATOMETRY; GRUENEISEN
CONSTANT; HAFNIUM CARBIDES: T1; HAFNIUM NITRIDES: T2; LATTICE
PARAMETERS; TANTALUM CARBIDES: T3; TANTALUM NITRIDES:
T4; THERMAL EXPANSION: Q1, Q2, Q3, Q4, Q5, Q6, Q7, Q8, Q9, Q10; THORIUM
CARBIDES: T5; THORIUM NITRIDES: T6; URANIUM CARBIDES:
T7; URANIUM NITRIDES: T8; VERY HIGH TEMPERATURE; X-RAY
DIFFRACTOMETERS; ZIRCONIUM CARBIDES: T9; ZIRCONIUM NITRIDES: T10

ABSTRACT: The report deals with the thermal lattice dilatation
of cubic transition metal compounds. The thermal dilatation

<508> CONT.

is determined through the variation of the lattice constants. The measurements are carried out 'in situ' by use of a high-temperature X-ray diffractometer chamber. The evaluation relates to both the linear thermal expansion coefficient α and, for some compounds, the specific heat at constant volume $C_{sub}(V)$ and the Grueneisen constant γ . In general a higher thermal dilatation is observed for nitrides than for carbides with the compounds formed by the transition metals belonging to the IVA and VA groups. The influence exerted by vacancies and by the oxygen dissolved in the lattice on the thermal dilatation of carbonitrides of zirconium, hafnium and tantalum is explained by the more pronounced anharmonic character of atomic vibrations in the crystal lattice.

EDB SUBJECT CATEGORY: EDB-360204

INIS CATEGORY: B23

<509>

Politis, C. (Kernforschungszentrum Karlsruhe (F.R. Germany).
Inst. fuer Material- und Festkoerperforschung)

Phase analysis on simulated LWR fuel rods and its evaluation with the aid of constitution studies in the system uranium-zirconium-oxygen In Reactor conference, Duesseldorf, 30.3.-2.4.1976. Section 1: Reactor design and experiments
** AIX-07:240832; ERA-02:013001; EDB-77:017547; (1976) ZAED

SUBJECT DESCRIPTORS: BWR TYPE REACTORS: T2; CRYSTAL STRUCTURE; FUEL ELEMENTS: T, Q1, Q2; OXYGEN; PHASE DIAGRAMS; PHASE STUDIES; PHASE TRANSFORMATIONS; PWR TYPE REACTORS: T1; SIMULATION; TEMPERATURE DEPENDENCE; URANIUM; ZIRCALOY

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-210100; 210200

INIS CATEGORY: E23

<510>

Politis, C. (Kernforschungszentrum Karlsruhe (Germany, F.R.).
Inst. fuer Material- und Festkoerperforschung) ** Nuclear Energy Agency, 75 - Paris (France)

Phase analysis at simulated LWR fuel rods and their interpretation by constitution investigations in the uranium-zirconium-oxygen system In Specialists' meeting on the behaviour of water reactor fuel elements under accident conditions ** AIX-07:277870; ERA-02:026935; EDB-77:055249;
([nd]) INIS-mf--3224

<510>

<510> CONT.

SUBJECT DESCRIPTORS: BWR TYPE REACTORS; FUEL CANS; FUEL
 PELLETS; FUEL RODS; FUEL-CLADDING INTERACTIONS; MELTDOWN; OXYGEN:
 T1; PHASE DIAGRAMS; PHASE STUDIES: Q1, Q2, Q3; PWR TYPE
 REACTORS; SIMULATORS; URANIUM: T2; URANIUM DIOXIDE; VERY HIGH
 TEMPERATURE; ZIRCONIUM: T3; ZIRCONIUM OXIDES

ABSTRACT: None

EDE SUBJECT CATEGORY: EDB-360102; 210100; 050700
 INIS CATEGORY: B12

<511>

Poluehktov, N.S.; Mishchenko, V.T. (AN Ukrainskoi SSR, Odessa.
 Inst. Obshchei i Neorganicheskoi Khimii)

Use of oscillator strengths of 'supersensitive' ion transfers
 of lanthanides for calculation of some parameters determining
 the intensity of their absorption spectra

AIX-08:280676; EDB-77:031391; Teor. Eksp. Khim. (TEKHA)

10:(5):681-684 (Sep 1974)

SUBJECT DESCRIPTORS: ABSORPTION SPECTRA: Q1; COMPARATIVE
 EVALUATIONS; CORRELATIONS; DIAGRAMS; ELECTRIC
 FIELDS; ENERGY-LEVEL TRANSITIONS; IONS; OSCILLATOR
 STRENGTHS; RARE EARTH COMPOUNDS: T1

ABSTRACT: None

EDE SUBJECT CATEGORY: EDB-400201
 INIS CATEGORY: B12

<512>

Pclukhin, V.A.; Gel'chinskii, B.R.; Evseev, A.M.; Sryvalin,
 I.I.; Ukhov, V.F.; Vatolin, N.A.; Esin, O.A.

Using molecular dynamics method for the evaluation of Fe-Ni and
 Al-Zr melts structure properties AIX-09:349516; EDB-78:027556;
 Izv. Akad. Nauk SSSR, Met. (IZNMA) (3):35-37 (May 1976)

ABSTRACT: None

EDE SUBJECT CATEGORY: EDB-360103; 360102
 INIS CATEGORY: B22

<513>

Pong, R.; Donaghey, L.F.

Thermodynamic studies of gallium--indium liquid alloys by solid state electrochemistry with oxide electrolytes
 INS-76:019916; ERA-02:001031; EDB-76:087226; J. Chem. Eng. Data (JCEAA) 21: (3):370-374 (Jul 1976)

SUBJECT DESCRIPTORS: ELECTROLYTIC CELLS; ENTROPY; GALLIUM ALLOYS: T1; INDIUM ALLOYS: T2; LIQUID METALS; MIXING HEAT; THERMODYNAMIC PROPERTIES: Q1, Q2

ABSTRACT: The activity of gallium in Ga--In liquid alloys was studied in a high temperature galvanic cell using calcia stabilized zirconia as an oxide electrolyte. The measured activity data for gallium in the Ga-In melts were used to calculate the interaction coefficient in the liquid alloy, which has the form $\alpha_{\text{GaIn}} = \ln \gamma_{\text{Ga}} / (1 - x_{\text{Ga}})^2 = 0.2862 + 0.0352 x_{\text{Ga}} + (399.3 + 220.0 x_{\text{Ga}}) T^{-1}$. The derived expressions for the enthalpy and excess entropy of mixing were found to be given by $\Delta H^{\text{M}} = (791.4 + 218.6 x_{\text{Ga}} / (1 - x_{\text{Ga}}))$ and $\Delta S^{\text{xs}} = -(0.568 - 0.0350 x_{\text{Ga}}) x_{\text{Ga}} / (1 - x_{\text{Ga}})$. The enthalpy of mixing reaches a maximum of 226 cal/mol at a gallium mole fraction of 0.53 at x_{Ga} . Results are compared to thermodynamic data obtained by other methods. Experimental techniques and limitations are critically reviewed.

EDB SUBJECT CATEGORY: EDB-360104; 400400
 INIS CATEGORY: E22; B12

<514>

Popel, S.I.; Spiridonov, M.A.

Structure of In_{25}Bi melted compound
 AIX-09:349517; EDB-78:027523; Izv. Akad. Nauk SSSR, Met. (IZNMA) (3):47-50 (May 1976)

SUBJECT DESCRIPTORS: BISMUTH ALLOYS: T1; CRYSTAL STRUCTURE: Q1, Q2; DISTRIBUTION; ELECTRON DIFFRACTION; INDIUM ALLOYS: T2; INTERATOMIC DISTANCES: Q1, Q2; INTERMETALLIC COMPOUNDS; LIQUID METALS

ABSTRACT: Based on the study of the long-range maxima of the atom radial distribution function, interatomic distances have been revealed in a molten compound In_{25}Bi . Comparison of model curves describing electron diffraction activity with an experimental curve points to the existence of the following four structural elements in the melt: spherical regions with a statistical distribution of various types of atoms (20%), clusters of atoms of pure Indium and Bismuth (40 and 10%

<514>

<514> CONT.

respectively), and a chain quasi-structure of various types of atoms (30%).

EDB SUBJECT CATEGORY: EDB-360102

INIS CATEGORY: B22

<515>

Popov, B.N. (Skoplje Univ. (Yugoslavia) Tehnolosko-Metalurski Fakultet)

Electrochemical reduction of molybdenum oxychloride in molten lithium chloride - potassium chloride eutectic

AIX-08:339961;EDB-78:017436; Bull. Tokyo Inst. Technol. (Engl. Ed.) (BTITA) 41:(3):. 99-105 (1976)

SUBJECT DESCRIPTORS: CURRENT

DENSITY;ELECTRODEPOSITION;EUTECTICS;LITHIUM

CHLORIDES;MOLYBDENUM COMPOUNDS: T1;OXYCHLORIDES: T2;POTASSIUM CHLORIDES;POTENTIOMETRY;QUANTITY RATIO;REDUCTION: Q1,Q2

ABSTRACT: Chronopotentiometric study of the electrochemical reduction of MoO_2Cl_2 (I) in molten LiCl-KCl eutectic revealed one irreversible reduction step. Quarter-wave potential vs. a Pt(II)/Pt reference electrode was found to be -0.351 ± 0.003 V. Reduction of I is diffusion controlled with the diffusion coefficient estimated to be $1.008 \times 10^{-5} \text{ cm}^2 \text{ s}^{-1}$. The product of electrochemical reduction of I obtained by constant current electrolysis as a brownish-violet deposit was identified as MoO_2 by chemical analysis and X-ray powder method.

EDB SUBJECT CATEGORY: EDB-400400

INIS CATEGORY: B12

<516>

Pesypaiko, V.I.; Shter, G.E.; Vasina, N.A.; Gryzalova, E.S.; Trunin, A.S. (Vsesoyuznyj Zaochnyj Politekhnikheskij Inst., Moscow (USSR); Kujbyshevskij Meditsinskij Inst. (USSR); Kujbyshevskij Politekhnikheskij Inst. (USSR))

Practical application of conversion method in study of five-component mutual system consisting of nine salts NaF , Na_2MoO_4 , Na_2WO_4 , KF , K_2MoO_4 , K_2WO_4 , BaF_2 , BaMoO_4 , BaWO_4
 AIX-08:295364;EDB-77:056206; Dokl. Akad. Nauk SSSR (DANKA) 228:(3):613-616 (21 Jun 1976)

SUBJECT DESCRIPTORS: BARIUM COMPOUNDS: T7;BARIUM FLUORIDES: T1;BARIUM TUNGSTATES: T6;CHEMICAL REACTIONS;COMPLEXES;CONVERSION;DIFFERENTIAL THERMAL

<516> CONT.

ANALYSIS; MIXTURES; MOLYBDATES: T2; PHASE DIAGRAMS:
Q1, Q2, Q3, Q4, Q5, Q6, Q7, Q8, Q9, Q10; POTASSIUM COMPOUNDS:
T8; POTASSIUM FLUORIDES: T3; POTASSIUM TUNGSTATES: T9; SODIUM
FLUORIDES: T4; SODIUM TUNGSTATES: T10; TUNGSTATES: T5; X-RAY
SPECTROSCOPY

ABSTRACT: None

EDE SUBJECT CATEGORY: EDB-400201
INIS CATEGORY: B12

<517>

Posypaiko, V.I.; Stratilatov, B.V.; Pervikova, V.N.

Calculated method of eutectics of multicomponent systems on base
of algorithm and digital computers
AIX-07:277874; EDB-77:031093; Zh. Neorg. Khim. (ZNOKA)
21: (5):1298-1301 (May 1976)

SUBJECT DESCRIPTORS: ALGORITHMS; COMPUTER
CALCULATIONS; CRYSTALLIZATION; DIGITAL
COMPUTERS; EQUATIONS; EQUILIBRIUM; EUTECTICS; MIXTURES; PHASE
TRANSFORMATIONS: Q1, Q2, Q3; POTASSIUM CHLORIDES: T1; POTASSIUM
FLUORIDES: T2; POTASSIUM IODIDES: T3

ABSTRACT: None

EDE SUBJECT CATEGORY: EDB-360602; 400201
INIS CATEGORY: B12

<518>

Posypaiko, V.I.; Trunin, A.S.; Kosmynin, A.S.; Shter, G.E.
(Kujbyshevskii Politekhnikeskii Inst. (USSR); Kujbyshevskii
Meditsinskii Inst. (USSR))

Projection-thermographic method for study of ternary and ternary
mutual systems AIX-08:280681; EDB-77:031509; Dokl. Akad. Nauk
SSSR (DANKA) 228: (4):911-913 (1 Jun 1976)

SUBJECT DESCRIPTORS: BARIUM CHLORIDES: T1; BARIUM FLUORIDES:
T4; CRYSTALLIZATION; DIFFERENTIAL THERMAL
ANALYSIS; EUTECTICS; MEASURING METHODS; MIXTURES; PHASE
DIAGRAMS; PHASE STUDIES: Q1, Q2, Q3, Q4; SODIUM CHLORIDES:
T2; TUNGSTATES: T3

<518>

<519> CONT.

ABSTRACT: None

EDE SUBJECT CATEGORY: EDB-400201

INIS CATEGORY: B12

<519>

Pcsypaiko, V.I.; Trunin, A.S.; Miftakhov, T.T.; Gasanaliyev, A.M.; Garkushin, I.K. (Kujbyshevskij Politekhnikheskij Inst. (USSR))

Thermal analysis of the system of NaF, CaF₂, Na₂MoO₄, CaMoO₄ AIX-08:289314; EDB-77:050884; (UKZHAUkr. Khim. Zh.) 42:(7):687-691 (Jul 1976)

SUBJECT DESCRIPTORS: CALCIUM COMPOUNDS: T4; CALCIUM FLUORIDES: T1; DIFFERENTIAL THERMAL ANALYSIS; EUTECTICS; MELTING POINTS; MOLYBDATE: T2; PHASE DIAGRAMS: Q1, Q2, Q3, Q4, Q5; SODIUM COMPOUNDS: T5; SODIUM FLUORIDES: T3; SOLID SOLUTIONS

ABSTRACT: None

EDE SUBJECT CATEGORY: EDB-400201

INIS CATEGORY: B12

<520>

Posypajko, V.I.; Trunin, A.S.; Khitrova, L.M. (Vsesoyuznii Zaochnii Politekhnikheskii Inst., Moscow (USSR); Kujbyshevskii Politekhnikheskii Inst. (USSR); Voronezhskii Meditsinskii Inst. (USSR))

KF-KCl-K₂MoO₄ system AIX-07:277873; EDB-77:031083; Zh. Neorg. Khim. (ZNOKA) 21:(2):547-550 (Feb 1976)

SUBJECT DESCRIPTORS: CRYSTALLIZATION; DIFFERENTIAL THERMAL ANALYSIS; MOLYBDATE: T1; PHASE DIAGRAMS: Q1, Q2, Q3; POTASSIUM CHLORIDES: T2; POTASSIUM FLUORIDES: T3

ABSTRACT: None

EDE SUBJECT CATEGORY: EDB-360602

INIS CATEGORY: B12

<521>

Pculain, M.; Chanthanasinh, M.; Lucas, J.

New fluoride glasses EDB-78:004622; Mater. Res. Bull. (MRBUA)
12:(2):151-156 (Feb 1977)

SUBJECT DESCRIPTORS: BARIUM FLUORIDES: T3;CHEMICAL
COMPOSITION;CRYSTALLIZATION;GLASS: T4;MELTING POINTS;OPTICAL
PROPERTIES: Q1,Q2,Q3,Q4;SYNTHESIS;THERMODYNAMIC
PROPERTIES;THORIUM FLUORIDES: T2;ZIRCONIUM FLUORIDES: T1

ABSTRACT: New fluoride glasses have been obtained in the ZrF_4 - ThF_4 - BaF_2 systems. Large samples have been prepared in order to determine some physical properties. Optical transmission range lies from 0.22 to 7μ , the refractive index is about 1.53 and can be modulated by chemical composition. Glass temperature, crystallization and melting temperature have been measured.

EDB SUBJECT CATEGORY: EDB-360204

<522>

Poyarkov, A.M.; Lebedev, V.A.; Nichkov, I.F.; Rspopin, S.P.

Thermodynamic properties of liquid thorium-gallium alloys
EDB-76:042177; Izv. Akad. Nauk SSSR, Met. (IZNMA) (3):87-88
(May 1975)

SUBJECT DESCRIPTORS: ELECTROMOTIVE FORCE;GALLIUM ALLOYS: T1;HIGH
TEMPERATURE;INTERMETALLIC COMPOUNDS;LIQUID METALS;SOLID
SOLUTIONS;THERMODYNAMIC PROPERTIES: Q1,Q2;THORIUM ALLOYS: T2

ABSTRACT: The emf (electromotive force) method was used to study the thermodynamic properties of liquid alloys of thorium with gallium and of an intermetallic compound existing in equilibrium with them. The solubility of thorium in gallium was also determined. The formation of the intermetallic compound is accompanied by the release of a considerable amount of heat, and by a decrease in entropy and in the free energy of the thorium. This change of thermodynamic functions suggests an energetic interaction of thorium with gallium in the fused state.

EDB SUBJECT CATEGORY: EDB-360102
INIS CATEGORY: B22

<523>

<523>

Predel, E.; Berka, H. (Max-Planck-Institut fuer Metallforschung, Stuttgart (F.R. Germany). Inst. fuer Werkstoffwissenschaften; Stuttgart Univ. (TH) (F.R. Germany). Inst. fuer Metallkunde)

Thermodynamic investigation of liquid alloys in the system
Cd-In-Tl AIX-07:275118;EDB-77:018629; Z. Metallkd. (ZEMTA)
67:(5):296-302 (May 1976)

SUBJECT DESCRIPTORS: CADMIUM;CADMIUM ALLOYS: T1;ENTROPY;FREE
ENTHALPY;HIGH TEMPERATURE;INDIUM ALLOYS: T2;LIQUID
METALS;MIXING HEAT;QUANTITY RATIO;TERNARY ALLOY
SYSTEMS;THALLIUM ALLOYS: T3;THERMODYNAMIC ACTIVITY: Q1,Q2,Q3

ABSTRACT: The thermodynamic activities of Cd in liquid Cd-In-Tl alloys have been determined by the application of the isopiestic method. The entire field of concentrations has been covered at temperatures of 500, 550, 595, 650 and 700°C. The positive deviations of the activity isotherms from Raoult's law are indicating a prevailing tendency for demixing throughout the system. Nevertheless, a more detailed consideration of the partial and integral free excess enthalpies, of the enthalpies of mixing and of the excess entropies, as resulting from the fundamental experimental values leads to the conclusion that - similar to the conditions in the bordering systems Cd-In and Cd-Tl - there are also individual bonding influences in the ternary case: In spite of the general demixing tendency due to the differences of the atomic volumes of the component species this tendency is considerably reduced in the concentration area of about 75 at.-% Cd.

EDB SUBJECT CATEGORY: EDB-360102
INIS CATEGORY: B22

<524>

Predel, E.; Berka, H. (Max-Planck-Institut fuer Metallforschung, Stuttgart (F.R. Germany). Inst. fuer Werkstoffwissenschaften; Stuttgart Univ. (TH) (F.R. Germany). Inst. fuer Metallkunde)

Contribution to the problem of the thermodynamic properties and the structure of liquid Cd-In alloys
AIX-07:262718;EDB-77:018793; Z. Metallkd. (ZEMTA)
67:(3):198-204 (Mar 1976)

SUBJECT DESCRIPTORS: CADMIUM ALLOYS: T1;ENTROPY;HIGH
TEMPERATURE;INDIUM ALLOYS: T2;LIQUID METALS;MIXING
HEAT;THERMODYNAMIC PROPERTIES: Q1,Q2;VAPOR PRESSURE

ABSTRACT: In order to obtain a new set of thermodynamic data of liquid Cd-In alloys partial vapor pressure measurements by an isopiestic method have been carried out in the whole

<524> CONT.

concentration range at temperatures between 500 and 700⁰C. An analysis of the concentration dependences of the enthalpies and entropies of mixing resulting from the experimental data shows that there is a tendency towards compound formation in the melt near 75 at.-% Cd, where an intermetallic compound is occurring in the solid state. The volume fraction of the parts of the melt, where this tendency is prevailing is about 0.3 at 500⁰C and decreases to about 0.03 at 700⁰C.

EDB SUBJECT CATEGORY: EDB-360104
INIS CATEGORY: B22

<525>

Protopapas, P.; Parlee, N.A.D.

Theory for the derivation of a single hard-sphere potential describing the overall molecular interactions in binary liquid metal alloys as a function of temperature and composition EDB-76:048728; High Temp. Sci. (HITSA) 7:(4):259-287 (Dec 1975)

SUBJECT DESCRIPTORS: BIBLIOGRAPHIES; BINARY ALLOY SYSTEMS:
M1; LIQUID METALS; MATHEMATICAL MODELS; MOLECULAR STRUCTURE:
Q1; REVIEWS; SIMULATION

ABSTRACT: The hard sphere potential provides the simplest mathematical formalism of a kinetic theory for liquids. Numerous attempts have been made to develop kinetic theories based on more complex potential models, but their mathematics become almost intractable, and the necessary approximations and assumptions introduce errors. Furthermore, the necessary input data are extensive, involved, and seldom available. The situation worsens in the case of binary liquid mixtures. One way of improving the hard-sphere results is the implementation of the mathematical formalism for hard-sphere potential with hard spheres shrinking at higher temperatures, i.e., the implementation of the kinetic theory for hard-sphere potential with "effective" soft spheres. In previous papers, a corresponding state principle were developed and numerically tested for the calculation of temperature-dependent hard spheres for liquid metals. It is shown how a binary liquid metal alloy can be simulated by a hypothetical single-component hard-sphere liquid, the overall molecular interactions of which are described by the "effective" soft-sphere potential for pure liquid metals. The simulation and potential are tested on the calculation of the structure factor of 71 binary alloys of different components, at different compositions and temperatures. The theoretical vehicle is the Percus--Yevick theory for the

<525>

<525> CONT.

structure factor of pure liquids. Results are in excellent agreement with the Percus--Yevick theory for binary hard-sphere fluids and experimental data. 70 references. (DLC)

EDB SUBJECT CATEGORY: EDB-360104

<526>

Prusakov, V.N.; Ezhov, V.K.; Efremov, E.A.

Investigation of liquid-steam equilibrium in system with diluted solutions of metal fluorides in uranium hexafluoride
AIX-08:295482;EDB-77:056092; At. Energ. (USSR) (AENGA)
41: (2):98-101 (Aug 1976)

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-400105

INIS CATEGORY: B13

<527>

Pugh, C.E.; Robinson, D.N., Oak Ridge National Lab., Tenn. (USA)

In Some trends in constitutive equation model development for high-temperature behavior of fast-reactor structural alloys
** FRA-03:000677;INS-77:017890;EDB-78:008226; (1977)
CONF-770807--27

SUBJECT DESCRIPTORS: CHROMIUM-MOLYBDENUM STEELS: T3;LMFBR TYPE REACTORS: T1;MATHEMATICAL MODELS;MECHANICAL PROPERTIES: Q2,Q3;REACTOR MATERIALS: M2,Q1

ABSTRACT: The paper addresses some important features of the inelastic behavior of $2\frac{1}{4}\%$ Cr--1Mo steel and indicates a mathematical framework that is capable of representing these types of response. While the constitutive model discussed embraces capabilities beyond those of equations presently used in design analyses: their implementation into practicable analysis methods (such as finite-element programs) is more demanding. For example, in the case of slow time-dependent deformations, the equations governing accumulation of the inelastic strain components and the evolution of the tensorial state variable α are intimately coupled. A part of recommending any such model for use in design must be a quantitative assessment of the economic feasibility of implementation.

EDB SUBJECT CATEGORY: EDB-210500;360103

INIS CATEGORY: E35;B22

<528>

Pulham, R.J. (Nottingham Univ. (UK))

Recent developments in the chemistry of solutions of salts in liquid alkali metals AIX-08:312192; EDB-77:094032; Pure Appl. Chem. (PACHA) 49: (1):83-92 (1977)

SUBJECT DESCRIPTORS: ALKALI METAL COMPOUNDS; BIBLIOGRAPHIES; CESIUM: T1; CHEMICAL REACTIONS; ELECTRIC CONDUCTIVITY; ENTHALPY; LIQUID METALS; LITHIUM: T2; NONMETALS; PHASE DIAGRAMS: Q1, Q2, Q3, Q4, Q5; POTASSIUM: T3; REVIEWS; RUBIDIUM: T4; SODIUM: T5; SOLUBILITY; SOLUTIONS

ABSTRACT: Essential features of phase equilibria between alkali metal-salt-non-metal are summarized and narrowed to detailed consideration of dilute solutions of salts in metals. The use of resistivity techniques is illustrated to determine non-metal solubilities and hence derive partial molar quantities and anion solvation enthalpies. Dissolved non-metals increase the resistivity of a specific metal by a characteristic amount and this property is exploited in the study of solute-solute interactions. Examples are given of solute association, dissociation and equilibrium in metallic solution and the reactions are correlated with the corresponding solid state reaction enthalpies.

EDB SUBJECT CATEGORY: EDB-360102; 400201
INIS CATEGORY: B12

<529>

Pulham, R.J.; Adams, P.F.; Hubberstey, P.; Parry, G.; Thunder, A.E. ** Watson, J.S.; Wiffen, F.W.; Bishop, J.L.; Breeden, B.K., Oak Ridge National Lab., Tenn. (USA)

Reaction rates and electrical resistivities of the hydrogen isotopes with, and their solubilities in, liquid lithium. In Radiation effects and tritium technology for fusion reactors. Vol. IV ** INS-76:021788; ERA-02:004615; EDB-76:095984; (Mar 1976) CONF-750989--P4

SUBJECT DESCRIPTORS: DEUTERIUM; ELECTRIC CONDUCTIVITY; ENTHALPY; HIGH TEMPERATURE; LIQUID METALS; LITHIUM; LITHIUM DEUTERIDES: T2; LITHIUM HYDRIDES: T1; LITHIUM TRITIDES: T3; PHASE DIAGRAMS: Q1, Q2, Q3; PRESSURE DEPENDENCE; SOLUBILITY; SOLVATION; TRITIUM

ABSTRACT: The rate of reaction, k , of hydrogen and of deuterium with liquid lithium have been determined up to pressures of $20 \text{ kNm}^2 \text{sup } -2$ and at temperatures between 230 and $270 \text{sup } 0^\circ \text{C}$. The reaction is first order with an apparent activation energy of 52.8 and $55.2 \text{ kJmol}^2 \text{sup } -1$ for hydrogen and

<529>

<529> CONT.

deuterium, respectively. The deuterium isotope effect, $k_{\text{sub H}}/k_{\text{sub D}}$, decreases from 2.95 at 230 to 2.83 at 270[°]C. Tritium is predicted to react even more slowly than deuterium. The freezing point of lithium is depressed by 0.082 and 0.075[°]C, respectively, by dissolved hydride and deuteride giving eutectics at 0.016 mol percent H and 0.012 mol percent D in the metal-salt phase diagrams. The depression and eutectic concentration are expected to be less for tritium. The increase in the resistivity of liquid lithium caused by dissolved hydrogen isotopes is linear and relatively large, $5 \times 10^{-8} \Omega \text{m}$ (mol percent H or D)⁻¹. The solubility of lithium hydride and deuteride was determined from the marked change in resistivity on saturation. The liquidus of the metal-salt phase diagram rises steeply from the eutectic point to meet the two-immiscible liquid region. Tritium is expected to be less soluble than deuterium. The partial molar enthalpies of solution are 44.2 and 55.0 kJmol⁻¹ for hydrogen and deuterium, respectively. These values are used to calculate the solvation enthalpies of the isotope anions in the metal.

EDB SUBJECT CATEGORY: EDB-C60100;400201

INIS CATEGORY: R31;R12

<530>

Pyrina, V.K.; Prostakov, M.E. (Ural'skiy Gosudarstvennyy Univ., Sverdlovsk (USSR))

Interaction of potassium tetrafluoroborate with fluorides of potassium, lanthanum, neodymium in melt EDB-76:036686; Zh. Neorg. Khim. (ZNOKA) 20:(4):1140-1142 (Apr 1975)

SUBJECT DESCRIPTORS: CHEMICAL REACTIONS; FLUOROBORATES: T2; MOLTEN SALTS; PHASE DIAGRAMS: Q1, Q2, Q3, Q4, Q5; POTASSIUM COMPOUNDS: T1; THERMAL ANALYSIS; X-RAY DIFFRACTION

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-360602

INIS CATEGORY: B12

<531>

Rakitskii, A.N. (AN Ukrainskoj SSR, Kiev. Inst. Metallofiziki)

Investigation on the resistance of high-refractory materials to chromium melt AIX-08:321154; EDB-77:113764; Ogneupory (OGNEA) (12):47-51 (Dec 1975)

SUBJECT DESCRIPTORS: ALUMINIUM OXIDES: T8; ANNEALING; BERYLLIUM OXIDES: T2; CHEMICAL REACTIONS; CHROMIUM; CHROMIUM BASE

<531> CONT.

ALLOYS; CRUCIBLES: T9; MAGNESIUM OXIDES:
T7; MELTING; METALLOGRAPHY; MOLYBDENUM
COMPOUNDS; ELASTICITY; POROSITY; POWDERS; RARE EARTH
ADDITIONS; REFRACTORIES: T1; SILICIDES; STABILITY:
Q1, Q2, Q3, Q4, Q5, Q6, Q7, Q8, Q9; STRUCTURAL CHEMICAL
ANALYSIS; TABLES; THERMAL TESTING; TITANIUM BORIDES: T3; TUNGSTEN
COMPOUNDS; ZIRCONIUM BORIDES: T4; ZIRCONIUM CARBIDES:
T5; ZIRCONIUM OXIDES: T6

ABSTRACT: An investigation was made of the stability of refractory crucibles used in melting ERKh-type chromium (electrolytically refined in hydrogen atmosphere) and its alloys with rare-earth metals. As refractory materials for the crucibles powdered oxides (Al_2O_3 , ZrO_2 , MgO , BeO), carbides (TiC , ZrC), borides (TiB_2 , ZrB_2) and silicides (MoSi_2 , W_5Si_3) were used. Crucibles having a capacity up to 1 kg of liquid chromium were made from powdered refractory materials by the slip casting method and pressing technique. The porosity of the crucibles after calcining at 1850-2000 deg C was 4-8%. The melting process was carried out in a laboratory induction furnace in purified argon atmosphere. The metal in liquid state was kept in the crucible for 5 min. The suitability of the examined refractory materials for making crucibles and ingot molds, used in contact with molten chromium and its alloys, was determined from the degree of interaction of chromium melt with the refractory materials and from the mechanical (first of all, plastic) properties of the cast metal most sensitive to the presence of impurities in Cr. The degree of interaction was established by x-ray structural analysis and petrographic measurements of the refractory materials, as well as by chemical and metallographic analyses of the surface layer of the casting. Crucibles made of BeO , ZrC , Al_2O_3 and ThO_2 were found to be most inert towards non-alloyed molten chromium. Minimum contamination of Cr-rare-earth elements melts was observed in the case of crucibles made of ZrC .

EDB SUBJECT CATEGORY: EDB-360204
INIS CATEGORY: B23

<532>

Rakovshchik, G.M.; Linchevskij, P.V.

Nitrogen solubility in liquid nickel--niobium alloys
EDB-76:048704; Tekhnol. Legk. Splavov (TLSPD) (8):61-65 (1974)

SUBJECT DESCRIPTORS: DECASSING; ENTHALPY; ENTROPY; LIQUID
METALS; NICKEL BASE ALLOYS: T1; NIOBIUM ALLOYS: T2; NITROGEN:
T3; PRESSURE DEPENDENCE; QUANTITY RATIO; REACTION
KINETICS; SOLUBILITY: Q3; SOLVENT PROPERTIES: Q1, Q2; TEMPERATURE

<532>

<532> CONT.
DEPENDENCE

ABSTRACT: None

EDF SUBJECT CATEGORY: EDB-360104;360603
INIS CATEGORY: E22

<533>

Rapp, R.A., Ohio State Univ., Columbus (USA)

In Fundamental studies of metal fluorination reactions.
Progress report, May 1, 1976--April 31, 1977 **
ERA-02:030370;EDB-77:062487; (Jan 1977) COO--2671-2

SUBJECT DESCRIPTORS: CALCIUM
FLUORIDES; COPPER; DIFFUSION; FLUORINATION: Q1, Q2; METALS:
T2; NICKEL: T1; RESEARCH PROGRAMS; SOLUBILITY

ABSTRACT: The present research is being undertaken to study some fundamental thermodynamic and transport properties and detailed mechanisms pertinent to metal fluorination reactions. The program consists of determining the following properties using solid-state, CaF_2 -electrolyte galvanic cells: (A) solubility and diffusivity of fluorine in metals, specifically nickel, and (B) the nature of the point defects (type, concentration and mobility) in metal fluorides as a function of fluorine partial pressure, temperature and impurity concentration. The latter study encompasses the study of the electrical conductivity of metal fluorides under thermodynamically well-defined conditions. Another objective of the program is to design and construct a fluorine probe for the measure of fluorine chemical potentials in gaseous environments. The probe must be evaluated in oxygen-containing atmospheres which might limit its accuracy and lifetime. For several technical reasons, the specific experimentation has deviated somewhat from that originally proposed. For two reasons, a probable phase change in CaF_2 at a temperature around 1175°C and a thermal expansion coefficient which differs greatly from solid nickel, CaF_2 single crystal electrolytes could not yet be interfaced perfectly to solid nickel for electrochemical studies. However, these problems were avoided by the successful substitution of copper for nickel in the studies of solubility and diffusivity. Some problems also arose in the study of the electrical conductivity of NiF_2 , namely, only relatively impure NiF_2 was available, and NiF_2 exhibited poor mechanical stability as electrodes. Therefore, the proposed methods were used mostly for the study of B-PbF_2 , for which previous studies also

<533> CONT.

provided values for comparison. With the experience gained from the work reported here, future experimentation with nickel metal and NiF_2 should be successful.

EDB SUBJECT CATEGORY: EDB-400201

<534>

Reeder, P.L.; Alquist, L.J.; Lin, J.; Wright, J.F. (Battelle Pacific Northwest Labs., Richland, Wash. (USA))

Separation of alkaline earth from alkali metal fission products in the SOLAR on-line mass spectrometer

AIX-07:257637; ERA-02:010661; EDB-77:007656; Nucl. Instrum. Methods (NUIMA) 133: (3):501-502 (15 Mar 1976)

SUBJECT DESCRIPTORS: BARIUM FLUORIDES; BARIUM ISOTOPES: T1; CESIUM ISOTOPES: T2; COUNTING RATES; EFFICIENCY; FISSION PRODUCTS; ISOTOPE PRODUCTION; MASS SPECTROMETERS; MASS SPECTROSCOPY: Q1, Q2, Q3, Q4; MOLECULAR IONS; MULTI-ELEMENT SEPARATION; ON-LINE MEASUREMENT SYSTEMS; RUBIDIUM ISOTOPES: T3; STRONTIUM FLUORIDES; STRONTIUM ISOTOPES: T4; SURFACE IONIZATION

ABSTRACT: Fission product nuclides of Sr and Ba have been separated from their Rb and Cs isobars in an on-line spectrometer by the use of molecular fluoride ions of Ba and Sr formed on a hot Re surface.

EDB SUBJECT CATEGORY: EDB-400703

INIS CATEGORY: D13

<535>

Reetz, T.

In Measurement of the cell voltage of ThO_2 - Y_2O_3 -solid electrolyte cells in the range of mixed conduction

** ERA-02:000825; EDB-76:080725; Z. Phys. Chem. (Leipzig) (ZPCLA) 250: (3/4):205-216 (1972) ANL-Trans--1060

SUBJECT DESCRIPTORS: ELECTROLYTIC CELLS: T1; ELECTROMOTIVE FORCE: Q1; OXYGEN; PRESSURE DEPENDENCE; THORIUM OXIDES; YTTRIUM OXIDES

ABSTRACT: The cell voltage drop of galvanic chains with oxygen ion-conducting solid electrolytes by electron transfer can be estimated with a knowledge of either the parameters for electron conduction of the transfer numbers applicable to different oxygen pressures. For a given electrode system, it is possible to determine an optimum reference system in terms of the mean transfer number. Approximation equations are derived for the relationship between the transfer number and

<535>

<535> CONT.

mean transfer number of galvanic chains. The electrical conduction properties of the $\text{ThO}_2\text{--Y}_2\text{O}_3$ solid electrolyte are discussed on the basis of experimental results with the chains $\text{Pt/H}_2\text{O}(20\text{ atm/sat.})/\text{ThO}_2 + 20\text{ Mol- percent YO}_{1.5}/\text{air/Pt}$ (Chain 1) and $\text{Ni/Na,Na}_2\text{O(sat.)/ThO}_2 + 20\text{ Mol- percent YO}_{1.5}/\text{O}_2(\text{p'of } 1)/\text{Ag/Ni}$ (Chain 2) and literature data. The measured values of chain 2 at different oxygen pressures of the reference system furnished values for electron conduction parameters. 7 figures, 4 tables.

EDE SUBJECT CATEGORY: EDB-250902;400702

<536>

Reeves, F.; Chassay, R., National Aeronautics and Space Administration, Huntsville, Ala. (USA). George C. Marshall Space Flight Center

In Space Processing Applications Rocket project, SPAR 1. Final report ** INS-77:015213; ERA-02:057093; EDB-77:127846; (Dec 1976) N--77--16082

SUBJECT DESCRIPTORS: FABRICATION: Q2;SPACE

ABSTRACT: The experiment objectives, design/operational concepts, and final results of each of nine scientific experiments conducted during the first Space Processing Applications Rocket (SPAR) flight are summarized. The nine individual SPAR experiments, covering a wide and varied range of scientific materials processing objectives, were entitled: solidification of Pb-Sb eutectic, feasibility of producing closed-cell metal foams, characterization of rocket vibration environment by measurement of mixing of two liquids, uniform dispersions of crystallization processing, direct observation of solidification as a function of gravity levels, casting thorium dispersion-strengthened interfaces, contained polycrystalline solidification, and preparation of a special alloy for manufacturing of magnetic hard superconductor under zero-g environment.

EDB SUBJECT CATEGORY: EDB-360101;360201
INIS CATEGORY: B21;B23

<537>

Reil, K.C.; Cronenberg, A.W., Sandia Labs., Albuquerque, N.Mex.
(USA); EG and G Idaho, Inc., Idaho Falls (USA)

In Effective equation-of-state measurements on uranium dioxide
** NTS-78:000279; ERA-03:006804; INS-78:000240; EDB-78:008749;
(1977) SAND--77-1094C

SUBJECT DESCRIPTORS: EQUATIONS OF STATE; LMFBR TYPE REACTORS:
T1; MELTDOWN: M2, Q1; THERMODYNAMICS: Q2; URANIUM DIOXIDE:
T3; VAPOR PRESSURE: Q3

ABSTRACT: The safety analysis of Liquid Metal Fast Breeder Reactor (LMFBR) systems requires knowledge of the vapor pressure of reactor fuels up to 6000 K. A new technique was developed to obtain experimental vapor pressure data for reactor fuel materials at high temperatures and pressures. Fuel samples are pulse fission heated in the Annular Core Pulse Reactor (ACPR) at heating rates comparable to those anticipated during a severe LMFBR transient. Since the vapor of reactor fuels may contain several dissociated fuel species, volatile fission products, and/or fission gas, and the evolution of that pressure may be time or rate dependent, the terminology "effective equation-of-state (EEOS) measurement" was used to differentiate these results from the vapor pressure of pure materials. This technique can be easily applied to plutonium bearing and previously irradiated fuels. The EEOS technique was used to determine the vapor pressure of high purity uranium dioxide with an oxygen-to-metal ratio of 2.08. Energy depositions of up to 2720 J/g relative to room temperature were obtained, resulting in measured pressures in excess of 38 mpa.

EE SUBJECT CATEGORY: EDB-220900; 210500; 360204
INIS CATEGORY: C52; E35; B23

<538>

Reisfeld, R. (Hebrew Univ., Jerusalem (Israel). Dept. of Inorganic and Analytical Chemistry) ** Dunitz, J.D. (ed.) (Eidgenossische Technische Hochschule, Zurich (Switzerland). Lab. fuer Organische Chemie); Hemmerich, P. (ed.) (Konstanz Univ. (F.R. Germany). Fachbereich Biologie); Holm, R.H. (ed.) (Massachusetts Inst. of Tech., Cambridge (USA). Dept. of Chemistry); Ibers, J.A. (ed.) (Northwestern Univ., Evanston, Ill. (USA). Dept. of Chemistry); Joergensen, C.K. (ed.) (Geneva Univ. (Switzerland). Dept. de Chimie Minerale et Analytique); Neillands, J.B. (ed.) (California Univ., Berkeley (USA). Dept. of Biochemistry); Reinen, D. (ed.) (Marburg Univ. (F.R. Germany). Fachbereich Chemie); Williams, R.J.F. (ed.) (Oxford Univ. (UK). Inorganic Chemistry Lab.)

Radiative and non-radiative transitions of rare-earth ions in

<538>

<538> CCNT.

glasses In Rare Earths ** Structure and Bonding,
AIX-07:257202;EDB-77:007479; (1975) Springer

SUBJECT DESCRIPTORS: CATIONS;DATA;ELECTRIC MOMENTS;EMISSION
SPECTRA;ENERGY LEVELS;ENERGY-LEVEL TRANSITIONS:
Q1;FLUORESCENCE SPECTROSCOPY;GLASS;MAGNETIC
MOMENTS;QUADRUPOLES;RARE EARTH COMPOUNDS;RARE EARTHS:
T1;RELAXATION;REVIEWS;TEMPERATURE DEPENDENCE;VALENCE

ABSTRACT: Magnetic dipole and electric quadrupole transitions,
forced electric dipole transitions, intensity parameters of
trivalent rare earth ions in glasses, and nonradiative
relaxations are treated.

EDB SUBJECT CATEGORY: EDB-400201
INIS CATEGORY: B12

<539>

Reisfeld, R.; Lieblich, N.; Boehr, L. (Hebrew Univ., Jerusalem
(Israel)); Barnett, B. (Soreq Nuclear Research Center, Yavne,
Israel)

Energy transfer between $\text{Bi}^{3+} \rightarrow \text{Eu}^{3+}$, $\text{Bi}^{3+} \rightarrow \text{Sm}^{3+}$ and $\text{UO}_2^{2+} \rightarrow \text{Eu}^{3+}$
in oxide glasses AIX-07:253226;ERA-02:008296;EDB-77:004188;
J. Lumin. (JLUMA) 12-13::749-753 (Mar 1976)

SUBJECT DESCRIPTORS: EFFICIENCY;ENERGY TRANSFER:
Q1,Q2,Q3,Q4,Q5,Q6;EXCITATION;FLUORESCENCE;GLASS;LIFETIME;PROB
ABILITY;SPECTRA

ABSTRACT: Probabilities and efficiencies of energy transfer from
 Bi^{3+} to Eu^{3+} , Bi^{3+} to Sm^{3+} in borax
and germanate glasses, and from UO_2^{2+} to Eu^{3+}
in phosphate glass were measured. Enhancement of acceptor
fluorescence by two orders of magnitude was achieved as a
result of energy transfer. From the decrease of donor
fluorescence lifetime the energy transfer from UO_2^{2+}
to Eu^{3+} was found to be nonradiative.

EDB SUBJECT CATEGORY: EDB-640302
INIS CATEGORY: A12

<540>

Reshetnikova, L.F.; Shaimuradov, I.B.; Novoselova, A.V.
(Mskcvskij Gosudarstvennyj Univ. (USSR))

Investigation into reaction of rubidium and holmium fluorides
AIX-08:320706;EDB-77:114204; Izv. Akad. Nauk SSSR, Neorg.
Mater. (IVNMA) 12:(4):638-642 (Apr 1976)

SUBJECT DESCRIPTORS: CHEMICAL COMPOSITION;CHEMICAL REACTIONS:
Q1,Q2;DIFFERENTIAL THERMAL ANALYSIS;EUTECTICS;HIGH
TEMPERATURE;HOLMIUM FLUORIDES: T1;MELTING;PHASE DIAGRAMS:
Q1,Q2;RUBIDIUM FLUORIDES: T2;X-RAY DIFFRACTION

ABSTRACT: A study was made of the phase diagram of the system
 $\text{RbF}-\text{HoF}_{0.3}$. The investigation was carried out by
differential thermal and X-ray diffraction analyses. The
following double fluorides were formed in the system: $\text{Rb}_{0.3}\text{Ho}_{0.7}\text{F}_6$,
 $\text{Rb}_{0.2}\text{Ho}_{0.7}\text{F}_{10}$ and compounds with assumed
composition $\text{Rb}_{0.4}\text{Ho}_{0.6}\text{F}_{10}$, $\text{Rb}_{0.3}\text{Ho}_{0.7}\text{F}_{22}$. The compound
 $\text{Rb}_{0.4}\text{Ho}_{0.6}\text{F}_{10}$ was found only thermographically. There are two eutectic points: the
eutectic between rubidium fluoride and $\text{Rb}_{0.3}\text{Ho}_{0.7}\text{F}_6$ has
the composition 7.5 mol.% $\text{HoF}_{0.3}$ and melts at 745 deg C;
the eutectic between $\text{Rb}_{0.3}\text{Ho}_{0.7}\text{F}_6$ and $\text{Rb}_{0.2}\text{Ho}_{0.7}\text{F}_{10}$
contains 45% $\text{HoF}_{0.3}$ and melts at 850 deg C. Rubidium
hexafluoroholmate ($\text{Rb}_{0.3}\text{Ho}_{0.7}\text{F}_6$) is the only compound
of the system which melts in a congruent way ($t_{\text{sub}}(m)=1020$
deg C).

EDB SUBJECT CATEGORY: EDB-400201:360603
INIS CATEGORY: B12

<541>

Rhoades, S.C., Bureau of Mines, Washington, D.C. (USA)

In $\text{K}_{0.2}\text{Hf}_{0.6}\text{F}_{1.25}-\text{K}_{0.2}\text{Zr}_{0.6}\text{F}_{1.25}$ percent of HF
system at 40 deg C with other solubility curves from 25 deg C
to 70 deg C ** INS-77:011944;EDB-77:120764; (1973)
EM-RI--7785

SUBJECT DESCRIPTORS: DIAGRAMS;HAFNIUM FLUORIDES: T2;HYDROFLUORIC
ACID: T4;MEDIUM TEMPERATURE;PHASE STUDIES:
Q1,Q2,Q3,Q4;POTASSIUM FLUORIDES:
T1;RECRYSTALLIZATION;SOLUBILITY: Q1,Q2,Q3;SOLVENT PROPERTIES:
Q4;ZIRCONIUM FLUORIDES: T3

ABSTRACT: The Bureau of Mines investigated the complete ternary
system $\text{K}_{0.2}\text{Hf}_{0.6}\text{F}_{1.25}-\text{K}_{0.2}\text{Zr}_{0.6}\text{F}_{1.25}$
weight-percent HF at 40 deg C. In addition, solubility
curves have been established at 25 deg C, 30 deg C, 50 deg C,
60 deg C, and 70 deg C. These solubility data
implemented a study of hafnium enrichment in feed solutions

<541>

<541> CONT.

for liquid-liquid extraction processes. Selected $K_{2\text{HfF}_6\text{--}K_2\text{ZrF}_6}$ 1.25 weight-percent HF mixtures were equilibrated in a constant temperature bath. Compositions of the equilibrated solutions and wet residues were established from combined hafnium and zirconium oxide and total fluorine analyses. All solubility curves were essentially parallel on triangular coordinates. At 40°C, the solid phase of the $K_{2\text{HfF}_6\text{--}K_2\text{ZrF}_6}$ 1.25 percent HF system is a series of solid solution hydrates with $K_{2\text{HfF}_6}$, H_2O and $K_{2\text{ZrF}_6}$, $3\text{H}_2\text{O}$ segments existing at the respective extremes. Since this system contains $K_{2\text{HfF}_6}$, H_2O and $K_{2\text{ZrF}_6}$, $3\text{H}_2\text{O}$ segments, it is not ideal for simple separation of high-purity $K_{2\text{HfF}_6}$ from $K_{2\text{HfF}_6\text{--}K_2\text{ZrF}_6}$ mixtures by recrystallization. However, it would be useful for producing hafnium-enriched feed solutions from zircon concentrates for liquid-liquid extraction of hafnium from zirconium.

EDP SUBJECT CATEGORY: EDP-400201;400105
INIS CATEGORY: F12;B11

<542>

Rosa, C.J., Cincinnati Univ., Ohio (USA). Dept. of Materials Science and Metallurgical Engineering

In Basic mechanisms providing oxidation resistance in structural metals at high temperatures. II. The defect structure of nonstoichiometric rutile (TiO_2) and nonstoichiometric rutile containing 1 to 10 mole % Ce_2O_3 . Final report 21 Dec 1973-20 Dec 1974 **
ERA-02:010462;EDB-77:007113: (Aug 1975) AD-A024540

SUBJECT DESCRIPTORS: ELECTROLYTES;ELECTROLYTIC CELLS;NIOBIUM OXIDES: T3;OXIDATION: Q1;RUTILE: T2;THERMODYNAMIC PROPERTIES: Q2,Q3;THORIUM OXIDES;TITANIUM BASE ALLOYS: T1;VERY HIGH TEMPERATURE;YTTRIUM OXIDES

ABSTRACT: The overall research program in which this investigation is a part is a study of oxidation of titanium and titanium alloys in hopes of yielding information concerning possible alloy additions which may improve the oxidation resistance of titanium structural materials at temperatures in the range 800 to 1200°C. Previous investigations have shown marked improvement in the oxidation resistance of titanium by additions of several metals, of which columbium is one. Because the thermodynamic stability limits for nonstoichiometric rutile is in the range of oxygen chemical potentials which can be conveniently measured by electrochemical techniques, an EMF (electromotive force) cell

<542> CCNT.

using yttria doped thoria as a solid electrolyte was designed to determine the thermodynamic properties of nonstoichiometric rutile and rutile containing columbium. (GRA)

EDB SUBJECT CATEGORY: EDB-360105;360204

<543>

Roy, S.K. (Ranchi Univ. (India). Dept. of Chemistry)

Heats of formation of solid solutions of CsCl with CsBr

AIX-08:331987;EDB-78:022284; Indian J. Chem., Sect. A (IJCAD) 14:(12):1008-1009 (Dec 1976)

SUBJECT DESCRIPTORS: CESIUM BROMIDES: T1;CESIUM CHLORIDES: T2;CRYSTAL DEFECTS;FORMATION HEAT: Q1,Q2;SOLID SOLUTIONS

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-400201

INIS CATEGORY: B12

<544>

Rustamov, P.G.; Seidova, N.A.; Shakhbazov, M.G.

(Azerbajdzhanskij Gosudarstvennyj Pedagogicheskij Inst., Baku (USSR); AN Azerbajdzhanskoy SSR, Baku. Inst. Neorganicheskoy i Fizicheskoy Khimii)

Ga-Bi-Te system AIX-08:329911;EDB-78:009649; Zh. Neorg. Khim. (ZNOKA) 21:(3):764-768 (Mar 1976)

SUBJECT DESCRIPTORS: BISMUTH: T3;BISMUTH ALLOYS;BISMUTH TELLURIDES: T1;CRYSTALLIZATION;DIFFERENTIAL THERMAL ANALYSIS;EUTECTICS;GALLIUM: T4;GALLIUM ALLOYS;GALLIUM TELLURIDES: T2;MICROHARDNESS;PHASE DIAGRAMS: Q1,Q2,Q3,Q4

ABSTRACT: To elucidate the nature of interaction in the system Ga-Bi-Te, a study has been made of sections $\text{GaTe-Bi}_{\text{sub } 2}\text{Te}_{\text{sub } 3}$, $\text{Ga}_{\text{sub } 2}\text{Te}_{\text{sub } 3}\text{-Bi}$, GaTe-Bi and $\text{Bi}_{\text{sub } 2}\text{Te}_{\text{sub } 3}\text{-Ga}$. The alloys have been prepared by direct melting of the components or their alloys with subsequent homogenizing and annealing at 400 deg C. The study has been made by the methods of differential thermal, microstructural analysis and by microhardness measurements. On the basis of literature data and data obtained a projection of the liquidus surface of the phase diagram for the system Ga-Bi-Te has been constructed. In the ternary system there are 17 curves of monovariant equilibrium dividing the liquidus into 10 fields of primary crystallization of phases, 9 points of

<544>

<544> CONT.

non-variant equilibrium of which 4 points are triple eutectics and 5 points are triple peritectics.

EDB SUBJECT CATEGORY: EDB-360602
INIS CATEGORY: B12

<545>

Rustamov, P.G.; Seidova, N.A.; Shakhbazov, M.G. (Arzamasskiy Gosudarstvennyy Pedagogicheskiy Inst. (USSR); AN Azerbajdzhanskoj SSR, Baku. Inst. Neorganicheskoy i Fizicheskoy Khimii)

GaTe-Pi\$sub 2fTe\$sub 3f system AIX-08:329912;EDB-78:009650; Zh. Neorg. Khim. (ZNOKA) 21:(3):860-862 (Mar 1976)

SUBJECT DESCRIPTORS: BISMUTH ALLCYS;BISMUTH TELLURIDES:
T1:EUTECTICS;GALLIUM ALLCYS;GALLIUM TELLURIDES:
T2:MICROHARDNESS;MICROSTRUCTURE;PHASE DIAGRAMS: Q1,Q2;SOLID SOLUTIONS

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-360602
INIS CATEGORY: B12

<546>

Ruzinova, E.A.; Kozhemyakin, V.A.; Korshunov, B.G.; Pirkovskaya, N.V. ** Poletaev, I.F.; Rubajlova, K.M. (eds.), Gosudarstvennyy Nauchno-Issledovatel'skiy i Proektnyj Inst. Redkometallicheskoj Promyshlennosti, Moscow (USSR)

Practice of experiment planning in studying limited regions of multicomponent system in rare earth metals **
AIX-08:292458;EDB-77:074877; (1976) INIS-mf--3315

SUBJECT DESCRIPTORS: ALUMINIUM CHLORIDES:
T1:EQUATIONS;EXPERIMENT PLANNING: T;IRON CHLORIDES:
T2:MAGNESIUM CHLORIDES: T3;MELTING;MOLTEN SALTS;PHASE
DIAGRAMS: Q1,Q2,Q3,Q4,Q5,Q6;POTASSIUM CHLORIDES: T4;SODIUM
CHLORIDES: T5;VAPOR PRESSURE;ZIRCONIUM CHLORIDES: T6

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-360602
INIS CATEGORY: B12

<547>

Ryabov, E.N.; Sandler, R.A.; Nguen Ngok Kuan; Klyuchnikova, E.F.; Kozhina, I.I. (Leningradskij Gornyj Inst. (USSR))

MoCl₃-NaCl and MoCl₃-KCl binary systems

EDE-76:048907; Zh. Neorg. Khim. (ZNOKA) 20:(3):765-768 (Mar 1975)

SUBJECT DESCRIPTORS: BINARY MIXTURES; DIFFERENTIAL THERMAL ANALYSIS; MOLYBDENUM CHLORIDES: T3; PHASE DIAGRAMS: Q1, Q2, Q3; X-RAY DIFFRACTION

ABSTRACT: None

EDE SUBJECT CATEGORY: EDE-360602

INIS CATEGORY: B12

<548>

Ryabov, E.N.; Aleksandrovskii, S.V.; Sandler, R.A.; Zakharevich, A.A.; Naumchik, A.N.

ZrCl₄-TiCl₃-NaCl ternary system

AIX-08:283015; EDB-77:044276; Zh. Prikl. Khim. (ZPKHA) 49:(6):1378-1379 (Jun 1976)

SUBJECT DESCRIPTORS: EUTECTICS; MELTING POINTS; MOLTEN SALTS; PHASE DIAGRAMS: Q1, Q2, Q3; SODIUM CHLORIDES: T1; TITANIUM CHLORIDES: T2; ZIRCONIUM CHLORIDES: T3

ABSTRACT: None

EDE SUBJECT CATEGORY: EDE-400201

INIS CATEGORY: B12

<549>

Rykhail, R.M.; Zarechnyuk, O.S. (L'vovskij Gosudarstvennyj Univ. (Ukrainian SSR)) ** Ministerstvo Vysshego i Srednego Spetsial'nogo Obrazovaniya SSSR, Moscow; L'vovskij Gosudarstvennyj Univ. (Ukrainian SSR)

New representatives of CeMn₄Al₈ structural type in 2. all-union conference on crystallochemistry of intermetallic compounds ** AIX-07:275122; EDB-77:050192; (1974) INIS-mf--3191

SUBJECT DESCRIPTORS: ALUMINIUM BASE ALLOYS: T2; DIFFRACTION METHODS; INTERMETALLIC COMPOUNDS; LATTICE PARAMETERS: Q1, Q2, Q3; RARE EARTH ALLOYS: T1; RHENIUM ALLOYS: T3; TETRAGONAL LATTICES

<549>

<549> CONT.

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-360102

INIS CATEGORY: B22

<550>

Sabounji, M.; Rahman, A.

Application of molecular dynamics computations to the conformational ionic solution theory: Thermodynamics of binary molten salt mixtures INS-76:019963; ERA-02:001161; EDB-76:081827; J. Chem. Phys. (JCPSA) 65:(6):2393-2399 (15 Sep 1976)

SUBJECT DESCRIPTORS: BINARY MIXTURES; FREE ENERGY; MIXING; MOLTEN SALTS: T1; STATISTICAL MECHANICS; THERMODYNAMIC PROPERTIES: Q1

ABSTRACT: An evaluation of the coefficient $a(T,V)$ as defined in the equation for the excess Helmholtz free energy of mixing of binary solutions, $\Delta A_m / RT = a(T,V) \sum_i (\lambda_i - \lambda_j) X_i / X_j + \dots$, derived from the conformational ionic solution theory is presented. This coefficient is shown to be negative and proportional to a fluctuation energy. The sign agrees with experimental calorimetric data of binary molten salt mixtures. A molecular dynamics computation of $a(T,V)$ was made to compare theory and experiments. The calculated values are strongly dependent on the form of the repulsive contribution to the pair potentials (Tosi and Fumi model and Michielsen, Woerlee, Graaf, and Ketelaar model). Criteria for testing the repulsive part of proposed pair potentials are suggested as a result of this investigation. (AIP)

EDB SUBJECT CATEGORY: EDB-400201

INIS CATEGORY: B12

<551>

Sabounji, M.I.; Blander, M.

Electromotive force measurements in molten lithium--magnesium alloys ERA-01:023780; EDB-76:065775; J. Electrochem. Soc. (JESOA) 122:(12):1631-1634 (Dec 1975)

SUBJECT DESCRIPTORS: ELECTROMOTIVE FORCE: Q1, Q2; LIQUID METALS; LITHIUM ALLOYS: T1; MAGNESIUM ALLOYS: T2; PHASE STUDIES; THERMODYNAMIC PROPERTIES

ABSTRACT: Electromotive force measurements of molten lithium-magnesium alloys are presented. Results of the

<551> CONT.

measurements may be combined with data from the measured Li-Mg phase diagram to calculate the thermodynamic properties of solid Li-Mg alloys at the solidus. (MCG)

EDB SUBJECT CATEGORY: EDB-360104;400400

<552>

Sabcunqi, M.L.; Blander, M., Argonne National Lab., Ill. (USA)

In Theoretical concepts useful in the calculation or storage of phase diagrams of ionic systems **

ERA-02:041863;EDB-77:094490; (Jan 1977) CONF-770116--1

SUBJECT DESCRIPTORS: COMPUTER CALCULATIONS: Q2;MOLTEN SALTS: T1;PHASE DIAGRAMS: Q1,T2

ABSTRACT: Equations for the a priori calculation or for the storage of phase diagrams and thermodynamic properties of molten ternary ionic systems were derived from the conformal ionic solution theory, which rigorously takes into account the ionic nature of the melts. Thermodynamic functions for ternary ionic systems were derived for cases in which all the cations have the same charge and all the anions have the same charge. Equations for the excess free energy of mixing, ΔG^E , are obtained for reciprocal systems, e.g., $A^+B^-X^-$, $B^+X^-Y^-$, defined by the three components AY, BX, and BY, and for additive systems, e.g., $A^+B^+C^-$, $B^+C^-X^-$. From these equations and classical thermodynamic relations, one can calculate phase diagrams of the two classes of ternary systems a priori from data on the lower order systems. Comparisons of such calculations with experimental data were made for liquidus phase diagrams and miscibility gaps in a number of ternary systems; the agreement was generally very good. An empirical extension of the theory to a broader range of systems, in which a pair of cations or anions have different charges, led to a good representation of the phase diagrams. The success of these calculations indicates their usefulness in evaluating the self-consistency of measurements and in the extrapolation of data into unmeasured regions. The equations should also prove useful for the storage of phase diagrams. The accuracy of such stored information can be improved by treating the coefficients as somewhat adjustable parameters and by adding more terms.

EDB SUBJECT CATEGORY: EDB-360602

<553>

<553>

Safonov, V.V. .; Pes'kova, Zh.K.; Grigor'eva, N.M.; Ksenzenko,
V.I. (Moskovskij Inst. Tonkoj Khimicheskoy Tekhnologii (USSR))

BiCl_3 - GeCl_4 - TeCl_4 system
AIX-07:277889; EDB-77:031085; Zh. Neorg. Khim. (ZNOKA)
21: (2):588-589 (Feb 1976)

SUBJECT DESCRIPTORS: BISMUTH CHLORIDES: T1; DIFFERENTIAL THERMAL
ANALYSIS; EUTECTICS; GERMANIUM CHLORIDES: T2; MELTING
POINTS; PHASE DIAGRAMS: Q1, Q2, Q3; TELLURIUM CHLORIDES: T3

ABSTRACT: None

EDF SUBJECT CATEGORY: EDB-360602
INIS CATEGORY: B12

<554>

Safonov, V.V.; Burshina, V.V.; Ivnitskaya, R.B. (Moskovskij
Inst. Tonkoj Khimicheskoy Tekhnologii (USSR))

FeCl_3 - MoCl_5 - NbCl_5 - WCl_6 system
AIX-08:329913; EDB-78:010015; Zh. Neorg. Khim. (ZNOKA)
21: (3):863-865 (Mar 1976)

SUBJECT DESCRIPTORS: CHEMICAL REACTIONS; DIFFERENTIAL THERMAL
ANALYSIS; EUTECTICS; IRON CHLORIDES: T1; MOLYBDENUM CHLORIDES:
T2; NIOBIUM CHLORIDES: T3; PHASE DIAGRAMS: Q1, Q2, Q3, Q4; TUNGSTEN
CHLORIDES: T4

ABSTRACT: None

EDF SUBJECT CATEGORY: EDB-400201
INIS CATEGORY: B12

<555>

Safonov, V.V.; Burshina, V.V.; Ivnitskaya, R.B.; Osipov, N.V.;
Korshunov, B.G. (Institut Tonkoj Khimicheskoy Tekhnologii,
Moscow (USSR); Moskovskij Inst. Stali i Splavov (USSR))

Thermographic study of iron, molybdenum, tantalum and tungsten
high chlorides interaction AIX-08:339969; EDB-78:017378; Izv.
Vyssh. Uchebn. Zaved., Tsvet. Metall. (IVUTA) (1):77-79 (Jan
1976)

SUBJECT DESCRIPTORS: CHEMICAL REACTIONS:
Q1, Q2, Q3, Q4; DIFFERENTIAL THERMAL ANALYSIS; EUTECTICS; IRON
CHLORIDES: T1; MELTING POINTS; MOLYBDENUM CHLORIDES: T2; PHASE
DIAGRAMS; PHASE STUDIES; SOLID SOLUTIONS; TANTALUM CHLORIDES:
T3; TUNGSTEN CHLORIDES: T4

<555> CONT.

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-400201

INIS CATEGORY: B12

<556>

Safonov, V.V.; Kuzina, T.V.; Titov, L.G.; Korshunov, B.G.
(Moskovskij Inst. Tonkoj Khimicheskoy Tekhnologii (USSR))

Investigation into reactions of hexabromotellurates of
potassium, cesium and thallium(1) in melt
AIX-07:277890; EDB-77:031086; Zh. Neorg. Khim. (ZNOKA)
21: (2):586-587 (Feb 1976)

SUBJECT DESCRIPTORS: BROMINE COMPOUNDS: T1; CESIUM COMPOUNDS:
T2; CHEMICAL REACTIONS; DIFFERENTIAL THERMAL ANALYSIS; MOLTEN
SALTS; PHASE DIAGRAMS: Q1, Q2, Q3, Q4, Q5; POTASSIUM COMPOUNDS:
T3; SOLID SOLUTIONS; TELLURATES: T5; THALLIUM COMPOUNDS:
T4; X-RAY DIFFRACTION

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-360602

INIS CATEGORY: B12

<557>

Saito, Shinroku (Tokyo Inst. of Tech. (Japan). Research Lab. of
Engineering Materials) ** Saito, Shinroku; Somiya, Shigeyuki
(Tokyo Inst. of Tech. (Japan). Research Lab. of Engineering
Materials); Kotera, Yoshihide; Pask, J.A.; Batha, H.D. (eds.)

Recent development of ceramic basic science in Japan In
Equilibria and kinetics in modern ceramic processing **
EDB-76:053947; (1972) Tokyo Inst. of Tech.

SUBJECT DESCRIPTORS: ALUMINIUM OXIDES; BERYLLIUM OXIDES:
T1; CERAMICS; CHEMICAL PREPARATION; DOPED MATERIALS; INERT
ATMOSPHERE; MAGNESIUM OXIDES; POWDER METALLURGY:
Q1, Q2, Q3; PRESSURE DEPENDENCE; REVIEWS; SINTERING; SOL-GEL
PROCESS; URANIUM DIOXIDE: T2; ZIRCONIUM OXIDES: T3

ABSTRACT: Topics on the basic science on ceramics are discussed,
limiting to the recent works carried out in Japan on
densification and its concerns. The significant factors
contributing to densification are considered to be: 1) effect
of preparation of starting materials, 2) influence of
atmosphere, 3) special treatment of powder, and 4) sintering
mechanism under vacuum, atmosphere and high pressure. On the
preparation of specially modified starting powder materials
for the purpose of densification, there are two ways; one is

<557>

<557> CONT.

the hysteresis of preparation from mother salt, and the other is the effect of minor additions. The studies on the easily sinterable powder of UO_2 prepared by sol-gel process, the effects of Cl^+ and F^+ on MgO , very fast shrinkage at the initial stage of sintering of Al_2O_3 , ZrO_2 , and BeO derived from sulfates, the densification effect of Li_2O and B_2O_3 , the effect of valency of metallic ions using NiO as a sintering sample, and others are described. The study on the effect of vapor pressure on the sintering of MgO is reviewed. Findings were applied to the work of activation hot pressing of MgO . On the special treatment of powder, two reports are introduced. One was done with explosive, and the other is powder treatment by electro-discharge. Studies on sintering mechanism, and models for sintering under atmospheric pressure are reviewed.

EDE SUBJECT CATEGORY: EDB-360201

INIS CATEGORY: B23

<558>

Sakurai, T.; Takahashi, A. (Japan Atomic Energy Research Inst., Tokai, Ibaraki. Tokai Research Establishment)

Behavior of ruthenium in fluoride-volatility process. II.

Fluorination of RuO_2 by F_2

AIX-08:312303; ERA-02:047677; EDB-77:098250; J. Inorg. Nucl. Chem. (JINCA) 39:(3):427-429 (1977)

SUBJECT DESCRIPTORS: FLUORIDE VOLATILITY PROCESS:

T; FLUORINATION: Q1; FLUORINE; HIGH TEMPERATURE; INFRARED

SPECTRA; MOLECULAR STRUCTURE; OXYFLUORIDES; RUTHENIUM OXIDES:

T1; SPENT FUELS; TEMPERATURE DEPENDENCE

ABSTRACT: RuO_2 was fluorinated by F_2 in the temperature range 250 to 500°C. Mass spectrometry confirmed that the reaction product was RuOF_4 . Small amount of RuO_4 was produced as a by-product in the fluorination. Temperatures higher than 400°C were necessary to convert RuO_2 smoothly into RuOF_4 . IR absorption spectrum of RuOF_4 was obtained. The product is an unstable material and tends to decompose into RuF_4 , releasing oxygen. Discrepancies with earlier published literature were discussed.

EDE SUBJECT CATEGORY: EDB-050800

INIS CATEGORY: B16

<559>

Sanchez, J.P.; Friedt, J.M.; Shenoy, G.K.; Percheron, A.;
 Achard, J.C., Strasbourg-1 Univ., 67 (France). Centre de
 Recherches Nucleaires

In Electronic and magnetic properties of rare earth-Sn₃
 compounds from ¹¹⁹Sn Moessbauer spectroscopy **
 AIX-07:246634;EDB-77:018704; (1975) CRN-CNPA--75-27

SUBJECT DESCRIPTORS: HYPERFINE STRUCTURE; ISOMER SHIFT; MAGNETIC
 FIELDS; MAGNETIC PROPERTIES: Q2, Q3; MOESSBAUER EFFECT:
 Q1; ORIENTATION; QUADRUPOLES; RARE EARTH ALLOYS: M2; TIN 119:
 M1; TIN ALLOYS: M3

ABSTRACT: The electronic and magnetic properties of RE-Sn₃
 compounds (RE=La, Ce, Pr, Nd, Sm, Eu, Gd, Yb) have been
 investigated using the 23.8keV Moessbauer resonance of ¹¹⁹Sn.
 The isomer shifts and quadrupole interactions are nearly the same
 in all compounds. The transferred magnetic fields and their
 orientation with respect to the principal electric field gradient
 axis at various Sn sites in the magnetically ordered state of
 RE-Sn₃ (RE=Pr, Nd, Sm, Eu, Gd) have been utilized to get
 information about the magnetic structure. An evaluation of the
 transferred fields in Pr-Sn₃ and Nd-Sn₃ shows that the spin
 density at the Sn nucleus is nearly the same in both compounds.

EDB SUBJECT CATEGORY: EDB-360104
 INIS CATEGORY: A13

<560>

Sandenshaw, T.A. (Los Alamos Scientific Lab., N.Mex. (USA)) **
 Blank, H.; Lindner, R. (eds.)

Anomalous thermal property behaviour of uranium at low
 temperatures In Plutonium 1975 and other actinides **
 AIX-08:309909; ERA-02:041793; EDB-77:094098; (1976)
 North-Holland

SUBJECT DESCRIPTORS: BRILLUIN ZONES; CRYSTAL-PHASE
 TRANSFORMATIONS; DISLOCATIONS; ENTHALPY; ENTROPY; FERMI
 LEVEL; GRAIN SIZE; HARDNESS; IMPURITIES; LOW
 TEMPERATURE; POLYCRYSTALS; SPECIFIC HEAT: Q1, Q2; STACKING
 FAULTS; TEMPERATURE DEPENDENCE; THERMAL CONDUCTIVITY:
 Q2; URANIUM; URANIUM 235: T1; URANIUM 238: T2; VERY LOW
 TEMPERATURE

ABSTRACT: Low temperature heat capacity curves are presented for
 polycrystalline ²³⁵U and ²³⁸U metals in different
 microstructural states and of different purities. Thermal
 conductivity versus temperature curves are shown for
 polycrystalline ²³⁸U metal after undergoing varied

<560>

<560> CONT.

fabrication procedures such as casting, swaging, γ -phase extrusion and α -phase rolling. Purity and the microstructure existing at the time of thermal property measurements appear to determine whether α -phase uranium is modified reversibly or irreversibly, if at all, on cooling below 250-270 K. It is suggested that the cooling rate through the $\beta \rightarrow \alpha$ phase transformation temperature of uranium has a considerable influence on observed low temperature properties because of the resulting defect (dislocation and stacking fault) structure. It is speculated that the reproducible heat capacity behaviour observed between 36 K and 250 K in a specimen of ^{238}U is due to a type of defect ordering which is compatible with a martensitic transformation.

EED SUBJECT CATEGORY: EDB-360103

INIS CATEGORY: E22

<561>

Schaefer, H.; Giegling, D. (Muenster Univ. (F.R. Germany). Anorganisch-Chemisches Inst.)

Chemistry of elements niobium and tantalum. 87. Studies on the thermal formation and decomposition of the complexes $[\text{NbCl}_6]^{2+}$, $[\text{NbI}_6]^{3+}$, and $[\text{TaX}_6]^{(n)+}$ (X=Cl, Br, I)
AIX-07:262572; EDB-77:019361; Z. Anorg. Allg. Chem. (ZAACA) 420: (1):1-24 (1976)

SUBJECT DESCRIPTORS: CHEMICAL REACTION KINETICS; CRYSTAL STRUCTURE; DEBYE-SCHERRER METHOD; MASS SPECTROSCOPY; NIOBIUM BROMIDES: T1; NIOBIUM CHLORIDES: T2; NIOBIUM COMPLEXES: T3; NIOBIUM IODIDES: T4; PYROLYSIS: Q1, Q2, Q3, Q4, Q5, Q6, Q7, Q8; SOLVENT EXTRACTION; STABILITY; TANTALUM BROMIDES: T5; TANTALUM CHLORIDES: T6; TANTALUM COMPLEXES: T7; TANTALUM IODIDES: T8; THERMAL GRAVIMETRIC ANALYSIS; X-RAY DIFFRACTION

ABSTRACT: Thermal decompositions of the compounds NbCl_3 , NbBr_3 , NbI_3 , $[\text{NbCl}_6]\text{Cl}_2$, $[\text{NbI}_6]\text{I}_2$, $[\text{NbCl}_6]\text{I}_3$, $[\text{TaCl}_6]\text{Cl}_3$, $[\text{TaBr}_6]\text{Br}_3$, $[\text{TaI}_6]\text{I}_3$ and $[\text{TaI}_6]\text{I}_2$ have been investigated by means of the thermobalance. These observations have been completed by chemical analysis of the solid phases, X-ray photographs, extractions of the solid with ethanol and mass spectra. The formation of the crystal lattices containing Me_6 -complexes during the thermal decomposition of MeX_3 or Me_3X is kinetically difficult. The decomposition in the vacuum requires much higher temperatures

<561> CONT.

as expected from thermodynamics. During the decomposition of TaCl_3 occurs an X-ray amorphous phase, from which Ta -complexes can be extracted. The retardation in respect to the formation of the corresponding crystal lattice is explained by the assumption that the complexes within the amorphous phase have different oxidation numbers. This requires a redox mechanism connected with diffusion. The decomposition reaction leading from $[\text{TaBr}_{12}]$ to $[\text{TaBr}_{12}]$ proceeds in such a way, that a small part of the complexes is decomposed and acts as reducing agent for the remaining undestroyed complexes.

EDB SUBJECT CATEGORY: EDB-400201

INIS CATEGORY: B12

<562>

Schenk, H.J.; Bohres, E.W.; Schwochau, K. (Kernforschungsanlage Juelich G.m.b.H. (F.R. Germany). Inst. fuer Nuklearchemie)

Diffuse reflectance spectra of protactinium(IV), uranium(IV) and neptunium(IV) tetrachlorides and tetraformates

AIX-07:236211; EDB-76:092884; J. Inorg. Nucl. Chem. (JINCA) 37:(12):2491-2495 (Dec 1975)

SUBJECT DESCRIPTORS: CHLORIDES; COORDINATION NUMBER: T7; CRYSTAL FIELD; ELECTRON TRANSFER; ELECTRONIC STRUCTURE; ENERGY-LEVEL TRANSITIONS; EXCITED STATES; FORMATES; LIGANDS; NEPTUNIUM CHLORIDES: T4; NEPTUNIUM COMPLEXES: T1; PROTACTINIUM CHLORIDES: T6; PROTACTINIUM COMPLEXES: T2; REFLECTION: Q1, Q2, Q3, Q4, Q5, Q6, Q7; SPECTRA; URANIUM CHLORIDES: T5; URANIUM COMPLEXES: T3

ABSTRACT: Diffuse reflectance spectra of 8-coordinate AnCl_4 and $\text{An}(\text{HCOO})_4$ ($\text{An}: \text{Pa}, \text{U}, \text{Np}$) have been measured in the range 18,000 to 2200 Å at room temperature. The observed excited levels arise from $5f^q$, $5f^{q-1}6d$ and $5f^{q+1}6p$ electron configurations. The $5f^q$ spectra of corresponding chlorides and formates are quite similar, shifted towards higher wave numbers in the latter case. Owing to the actinide contraction the crystal-field splitting of the $5f^q$ levels decreases with increasing atomic number. The energy variation of the $5f^q \rightarrow 5f^{q-1}6d$ transitions as a function of q is more pronounced in the predominantly ionic tetraformates than in the more covalent tetrachlorides. The cubic splitting parameter of the $6d$ shell is evaluated to be $\Delta_{\text{cubic}}(6d) = 9 \text{ kK}$ and $\Delta_{\text{cubic}}(6d) = 15 \text{ kK}$ in PaCl_4 and $\text{Pa}(\text{HCOO})_4$, respectively. The electron transfer spectra indicate the optical electronegativities $X_{\text{opt}} =$

<562>

<562> CONT.

1.42 for protactinium(IV) and $X_{\text{sub}}(\text{opt}) = 3.13$ for the formate ion.

EDE SUBJECT CATEGORY: EDB-400702

INIS CATEGORY: B13

<563>

Schenke, M.; Broers, G.H.J.; Ketallaar, J.A.A.

Solubility of oxygen in molten carbonates EDB-76:086741; J. Electrochem. Soc. (JESOA) 113:(4):404 (Apr 1966)

SUBJECT DESCRIPTORS: CARBONATES;ELECTROLYTES:

Q3;HIGH-TEMPERATURE FUEL CELLS: T3;MOLTEN SALTS: T1;OXYGEN:

T2;SOLUBILITY: Q2;SOLUTION HEAT;SOLVENT PROPERTIES: Q1

ABSTRACT: None

EDE SUBJECT CATEGORY: EDB-300505

<564>

Schirmacher, W., Freie Univ. Berlin (Germany, F.R.). Fachbereich Physik

In Nuclear quadrupole relaxation and viscosity in liquid metals

** A1X-01:264025;EDB-77:040109; (1976) AED-Conf--76-087-010

SUBJECT DESCRIPTORS: BESSEL

FUNCTIONS;EQUATIONS;HYDRODYNAMICS;INELASTIC

SCATTERING;INTEGRALS;LIQUID METALS: T1;NEUTRON

DIFFRACTION;NUCLEAR MAGNETIC RESONANCE;NUCLEAR QUADRUPOLE

RESONANCE: Q1,T2;RELAXATION: Q2;SELF-DIFFUSION;TEMPERATURE

DEPENDENCE;VISCOSITY

ABSTRACT: It is shown that the nuclear quadrupole relaxation rate due to the molecular motions in liquid metals is related to the shear and bulk viscosity and hence to the absorption coefficient of ultrasound. Application of the 'extended liquid phonon' model of Ortoleva and Nelkin - which is the third of a series of continued-fraction-approximations for the van Hove neutron scattering function - gives a relation to the self diffusion constant. The predictions of the theory concerning the temperature dependence are compared with quadrupole relaxation measurements of Riegel et al. and

<565> CONT.

Kerlin et al. in liquid gallium. Agreement is found only with the data of Riegel et al.

EEB SUBJECT CATEGORY: EDB-656000

INIS CATEGORY: A13

<565>

Schmitz, F. (CEA Centre d'Etudes Nucleaires de Fontenay-aux-Roses, 92 (France))

Oxygen potential and vacancy structure of the mixed oxide
(UPu)O/sub 2-x/ AIX-07:239904;EDB-77:013204; J. Nucl. Mater.
(JNUMA) 58:(3):357-360 (Dec 1975)

SUBJECT DESCRIPTORS: CHEMICAL

COMPOSITION;IMPURITIES;OXYGEN;PLUTONIUM OXIDES: T1;QUANTITY
RATIO;REDOX POTENTIAL: Q1,Q2;URANIUM OXIDES: T2;VACANCIES:
Q1,Q2

ABSTRACT: The authors have analyzed reported $\Delta G_{\text{O}_2} = f(\text{O/M})$ values. For concentrations of oxygen vacancies, $V_{\text{O}} = x$ and $(\text{PuO}_2) = y - 2x$, $\log x/(y - 2x) = \alpha \log P_{\text{O}_2} + \log k$. The value $x/(y - 2x)$, rather than the oxygen to metal ratio (O/M), determines the variation of the mixed oxide as a function of deviation from stoichiometry up to the level or near approach to stoichiometry or phase limits; the impurities determine the extrinsic value of the property considered. The hypostoichiometric domain of the mixed oxide $(\text{U}_{1-y}\text{Pu}_y)\text{O}_{2-x}$ is characterized by a defect complex, $V_{\text{O}} + 2\text{Pu}$ atoms. The energy of formation of the complex is about 2.25 eV.

EEB SUBJECT CATEGORY: EDB-360202

INIS CATEGORY: B24

<566>

Schcenfeld, I.; Ungarish, M. ** Israel Atomic Energy Commission, Tel Aviv

Spectrochemical method for determination of zirconium in molten salts In Research laboratories annual report 1975 **
AIX-08:320548;EDB-77:114116; (Jun 1976) IA--1321

SUBJECT DESCRIPTORS: ABSORPTION SPECTROSCOPY: Q1;CHEMICAL

ANALYSIS: Q2;EMISSION SPECTROSCOPY: Q1;MOLTEN SALTS:

T2;QUANTITATIVE CHEMICAL ANALYSIS;ZIRCONIUM: T1;ZIRCONIUM
SILICATES

<566>

<566> CONT.

ABSTRACT: None

EDE SUBJECT CATEGORY: EDB-400104

INIS CATEGORY: B11

<567>

Schommers, W. ** Pintschovius, L., Kernforschungszentrum
Karlsruhe (Germany, F.R.). Inst. fuer Angewandte Kernphysik

Frequency spectrum and scattering law for liquid copper. A
molecular dynamics investigation In Progress report of the
Teilinstitut Nukleare Festkoerperphysik.
(1.6.1974--31.5.1975) ** AIX-07:253507; EDB-77:007047; (Nov
1975) KFK--2183

SUBJECT DESCRIPTORS: COPPER: T1; CORRELATION FUNCTIONS; LIQUID
METALS; MATHEMATICAL MODELS; MOLECULAR STRUCTURE: Q1; SCATTERING

ABSTRACT: None

EDE SUBJECT CATEGORY: EDB-360104

INIS CATEGORY: A13

<568>

Schouler, E.; Hammou, A.; Kleitz, M.

Complex impedance of electrochemical cells based on yttria doped
thoria INS-77:006416; EDB-77:055745; Mater. Res. Bull. (MRBUA)
11: (9):1137-1146 (Sep 1976)

SUBJECT DESCRIPTORS: AGING; CHEMICAL COMPOSITION; ELECTRIC
CONDUCTIVITY; ELECTRIC IMPEDANCE; ELECTRICAL PROPERTIES:
Q1, Q2, Q3; ELECTROCHEMICAL CELLS: T1; IONIC
CONDUCTIVITY; QUANTITY RATIO; TEMPERATURE DEPENDENCE; THORIUM
OXIDES: T2; YTTRIUM OXIDES: T3

ABSTRACT: In the frequency range 10^{-2} to $5 \cdot 10^5$ Hz
and the temperature interval 400 to 1200°C , the
impedance diagram of a symmetrical cell based on yttria-doped
thoria is composed of three semicircles. The influence of the
geometrical factor of the thoria samples indicates that the
two low-frequency semicircles characterize interface
processes. The electrical resistance of the electrolyte is
determined by the low frequency intercept of the high
frequency semicircle with the real axis. The electronic
conductivity of thoria based solid solutions deduced from the
variation of this point obeys the law $\sigma_{\pm} = kP_{\pm}^{1/4}$. Ionic conductivity measured as a function of
the oxide vacancy concentration and the relevant activation

<568> CONT.

energies are also reported. An aging process was observed on this material. 7 figures.

EDB SUBJECT CATEGORY: EDB-360204
INIS CATEGORY: B23

<569>

Seddon, J.W.

In Nuclear reactor containments **

AIX-08:296900; ERA-02:034217; EDB-77:073461; (15 Sep 1976)

SUBJECT DESCRIPTORS: CLOSURES; COMPOSITE MATERIALS; CONTAINMENT SHELLS: M2, Q1; DESIGN: Q2; LMFBR TYPE REACTORS: M1; PRESSURE VESSELS; PRESTRESSED CONCRETE; STEELS; THERMAL EXPANSION

ABSTRACT: The containment comprises a cavity-defining prestressed concrete structure and a composite concrete and steel top closure which slides horizontally relative to the structure. The closure has a recess complementary to the cavity to form a chamber for housing a primary vessel containing a nuclear reactor core and ancillary equipment. The chamber bounded by a metal membrane lining the concrete, and wherein the concrete is relieved to form a peripheral groove adjacent to the lining, the width of the groove extending across the junction of the closure with the structure.

EDB SUBJECT CATEGORY: EDB-210500
INIS CATEGORY: E35

<570>

Seleznev, B.I.; Vlasov, Yu.G. (Leningradskij Gosudarstvennyj Univ. (USSR))

Thermodynamics of solid solutions CsCl-RbCl and CsBr-RbBr at 25¹sup 0°C AIX-08:298343; EDB-77:069082; Zh. Fiz. Khim. (ZFKHA) 50:(6):1605-1606 (Jun 1976)

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-400201
INIS CATEGORY: B12

<571>

<571>

Seppelt, K. (Heidelberg Univ. (F.R. Germany). Inst. fuer
Anorganische Chemie)

Covalent compounds of uranium of the type $F_{x/5}U(OTeF_5)_{6-x}$ AIX-07:272453; EDB-77:026655; Chem. Ber. (CHBEA)
109: (3):1046-1052 (1976)

SUBJECT DESCRIPTORS: CHEMICAL PREPARATION: Q1,Q2; COVALENCE; MASS
SPECTRA; NUCLEAR MAGNETIC RESONANCE; TELLURIUM COMPOUNDS:
T2; TELLURIUM FLUORIDES; URANIUM COMPOUNDS: T1; URANIUM
FLUORIDES; URANIUM OXIDES; VOLATILITY

ABSTRACT: Pure $U(OTeF_5)_6$ is prepared. It is volatile
in spite of the molecular weight of 1,669 mass units and
seems to be the inorganic nonpolymeric molecule with the
highest molecular weight known. All compounds
 $F_xU(OTeF_5)_{6-x}$ have been detected by the
n.m.r. method.

EIE SUBJECT CATEGORY: EDB-400702
INIS CATEGORY: E12

<572>

Shajmuradov, I.B.; Reshetnikova, L.P.; Novoselova, A.V.
(Moskovskij Gosudarstvennyj Univ. (USSR))

CsF-HoF₃ system EDB-76:008910; Zh. Neorg. Khim. (ZNOKA)
20: (4):1077-1080 (Apr 1975)

SUBJECT DESCRIPTORS: CESIUM FLUORIDES: T1; CHEMICAL
REACTIONS; DIFFERENTIAL THERMAL ANALYSIS; HOLMIUM FLUORIDES:
T2; PHASE DIAGRAMS: Q1,Q2; X-RAY DIFFRACTION

ABSTRACT: None

EIE SUBJECT CATEGORY: EDB-360602
INIS CATEGORY: B12

<573>

Shaner, J.W., California Univ., Livermore (USA). Lawrence
Livermore Lab.

In Thermal expansion of metals over the entire liquid range **
INS-77:019140; NTS-78:000095; ERA-03:007095; EDB-78:004477; (25
Aug 1977) UCRL--80007

SUBJECT DESCRIPTORS: DENSITY; HIGH TEMPERATURE; LIQUID METALS:
T1; MEASURING METHODS: Q2; REVIEWS; THERMAL EXPANSION:
Q1,T2; VERY HIGH TEMPERATURE

<573> CONT.

ABSTRACT: This paper reviews the current state of the art for measuring liquid metal densities. Conventional high precision techniques for use below 2000K as well as new techniques for more extreme temperatures are addressed. Pertinent data, which have appeared since the last critical reviews, for elemental metals are discussed.

EDB SUBJECT CATEGORY: EDB-360104

INIS CATEGORY: B22

<574>

Shaner, J.W.; Gathers, G.R.; Minichino, C., California Univ., Livermore (USA). Lawrence Livermore Lab.

In New apparatus for thermophysical measurements above 2500 K **
INS-04:016966; ERA-01:023767; EDB-76:065458; (Mar 1976)
UCRL--77929

SUBJECT DESCRIPTORS: ENTHALPY; FUSION HEAT; LIQUID METALS; MEASURING INSTRUMENTS; MELTING; MELTING POINTS; MOLYBDENUM: M2; NIOBIUM: M1; PHYSICAL PROPERTIES: Q1, Q2, Q3, Q4; PRESSURE DEPENDENCE; TANTALUM: M3; TUNGSTEN: M4; VERY HIGH TEMPERATURE

ABSTRACT: An apparatus was developed to make simultaneous measurements of enthalpy, temperature, volume, and electrical resistivity on metals in the temperature range from 2500 to 8000^{sup} 0°K and at pressures up to 1 GPa. Cylindrical samples are resistively heated by a 10-50 μ s current pulse in a constant pressure inert gas environment. Data for the refractory transition metals Nb, Mo, Ta, and W are presented.

EDB SUBJECT CATEGORY: EDB-360104

INIS CATEGORY: B22

<575>

Shatinskii, V.F.; Goikhman, M.S.; Mendel'son, D.A.; Garlinskii, R.N. (AN Ukrainskoj SSR, Lvov. Fiziko-Mekhanicheskij Inst.)

Friction in metal melts AIX-08:295560; EDB-77:055396; Fiz.-Khim. Mekh. Mater. (PKMMA) 12:(4):91-96 (Jul 1976)

SUBJECT DESCRIPTORS: ADHESION; EUTECTICS; FRICTION: Q1, Q2, Q3, Q4; HIGH TEMPERATURE; IRON: T1; LEAD ALLOYS: T2; LIQUID METALS: T3; PRESSURE DEPENDENCE; TEMPERATURE DEPENDENCE; TIN ALLOYS: T4

<575>

<575> CONT.

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-360103

INIS CATEGORY: B15

<576>

Shejko, I.N.; Grechina, T.N.; Sidorenko, I.A. (AN Ukrainskoj SSR, Kiev. Inst. Obshchej i Neorganicheskoy Khimii)

Viscosity of melts of $KCl-KF-K_{2.2}ZrF_{6.6}$ ternary system
EDB-76:042421; Ukr. Khim. Zh. (UKZHA) 41:(2):206-208 (Feb 1975)SUBJECT DESCRIPTORS: MOLTEN SALTS; POTASSIUM CHLORIDES:
T1; POTASSIUM FLUORIDES: T2; VISCOSITY: Q1, Q2, Q3; ZIRCONIUM
FLUORIDES: T3

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-360602; 400201

INIS CATEGORY: B12

<577>

Sheth, A.; Leibowitz, L., Argonne National Lab., Ill. (USA)

In Equation of state and transport properties of uranium and
plutonium carbides in the liquid region **
INS-77:003552; ERA-02:017804; EDB-77:026214; (Sep 1975)
ANL-AFP--11

SUBJECT DESCRIPTORS: DATA

COMPILATION; DENSITY; EMISSIVITY; ENTHALPY; EQUATIONS OF
STATE; PLUTONIUM CARBIDES: T2; SPECIFIC HEAT; THERMAL
CONDUCTIVITY; THERMODYNAMIC PROPERTIES: Q1, Q2; URANIUM
CARBIDES: T1; VAPOR PRESSURE; VISCOSITYABSTRACT: By the use of available low-temperature data for
various thermophysical and transport properties for uranium
and plutonium carbides, values above the melting point were
estimated. Sets of recommended values have been prepared for
the compounds UC, PuC, and (U,Pu)C. The properties that have
been evaluated are density, heat capacity, enthalpy, vapor
pressure, thermal conductivity, viscosity, and emissivity.

EDB SUBJECT CATEGORY: EDB-360204; 050700

INIS CATEGORY: B23

<578>

Sheth, A.; Leibowitz, L., Argonne National Lab., Ill. (USA)

In Equation of state and transport properties of uranium and plutonium nitrides in the liquid region **

INS-77:002299; ERA-02:017805; EDB-77:026215; (Oct 1976)

ANI-AFP--12

SUBJECT DESCRIPTORS: DENSITY; ENTHALPY; EQUATIONS OF STATE: Q1, Q2; PLUTONIUM NITRIDES: T2; SPECIFIC HEAT; THERMAL CONDUCTIVITY; THERMODYNAMIC PROPERTIES: Q1, Q2; URANIUM NITRIDES: T1; VAPOR PRESSURE; VISCOSITY

ABSTRACT: By the use of available low-temperature data for various thermophysical and transport properties for uranium and plutonium nitrides, values above the melting point of density, heat capacity, enthalpy, vapor pressure, thermal conductivity, and viscosity were estimated. Sets of recommended values have been prepared for the compounds UN, PuN, and (U_{0.8}Pu_{0.2})N.

EDB SUBJECT CATEGORY: EDB-360204

INIS CATEGORY: B23

<579>

Sheth, A.; Leibowitz, L.; Winslow, G.

Densities of liquid carbides and nitrides of uranium and plutonium ERA-02:002460; INS-76:020668; EDB-76:090325; Trans. Am. Nucl. Soc. (TANSA) 23::130-131 (Jun 1976)

SUBJECT DESCRIPTORS: DENSITY: Q1, Q2, Q3, Q4; LIQUIDS; NUCLEAR FUELS; PLUTONIUM CARBIDES: T3; PLUTONIUM NITRIDES: T4; THERMAL EXPANSION; THERMODYNAMIC PROPERTIES; URANIUM CARBIDES: T1; URANIUM NITRIDES: T2

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-050700; 360204

INIS CATEGORY: B23

<580>

Shimizu, S.; Kase, K.; Midoriyama, S.; Murakami, T. (Kawasaki Heavy Industries Ltd., Kobe (Japan))

Properties of structural materials for sodium-cooled fast breeder reactor AIX-08:315087; EDB-77:120379; FAPIG (Tokyo) (FPIGA) (81):19-31 (May 1976)

SUBJECT DESCRIPTORS: CARBURIZATION; CHROMIUM-MOLYBDENUM STEELS: T1; CORROSION: Q1, Q2, Q3, Q4, Q5, Q6; CORROSIVE EFFECTS:

<580>

<580> CONT.

Q7,Q8;CRACKS;CREEP;DECARBONIZATION;FATIGUE;FRACTURE
 PROPERTIES;INCOLOY 800: T6;LIQUID METALS;LMFBR TYPE
 REACTORS;SODIUM: T7;SODIUM COOLED REACTORS;STAINLESS
 STEEL-304: T2;STAINLESS STEEL-316: T3;STAINLESS STEEL-321:
 T4;STAINLESS STEEL-347: T5;STEAM;TENSILE PROPERTIES;WATER: T8

ABSTRACT: For the selection of the structural materials for superheaters and reheaters of sodium-cooled fast breeder reactors, it is important to grasp the change of strength due to the complex change of material properties, such as the combination of surface corrosion, the decarbonization of 2-1/4 Cr-1Mo steel, the carburization of austenitic stainless steel, the structural change due to heating history, etc. in sodium environment. The stress corrosion cracking of austenitic materials in water must be studied also. The materials taken up in this paper are austenitic stainless steels such as SUS 304, SUS 316, SUS 321, and SUS 347, iron-based superalloy Incoloy 800, and ferritic alloy steel 2-1/4Cr-1Mo steel. The data on the above described properties of the materials are given. Also the tensile strength, creep rupture and fatigue characteristics of the parent materials in the amount of corrosion in sodium. The strength of ferritic alloy steel is lowered owing to the decarbonization in sodium, but the change of strength due to carburization was not observed. There is some possibility that the unstabilized steels such as SUS 304 and SUS 316 become sensitive to stress corrosion cracking, and the stabilized steels such as SUS 321 and SUS 347 become sensitive to it in long hour heating. The tensile strength of welded joints is almost same as that of parent materials, but the elongation decreases by about 10%.

EDF SUBJECT CATEGORY: EDB-360105;210500
 INIS CATEGORY: B22

<581>

Shohoji, N.; Katsura, M.; Sano, T. (Osaka Univ. (Japan))

Decomposition of the UN-CeN solid solution

AIX-07:254866;EDB-76:097712; J. Nucl. Mater. (JNUMA)
 60: (1):52-58 (Apr 1976)

SUBJECT DESCRIPTORS: CERIUM NITRIDES:

T1;DECOMPOSITION;EQUATIONS;FORMATION FREE ENERGY;LOW
 PRESSURE;MEDIUM VACUUM;PHASE TRANSFORMATIONS: Q1,Q2;SOLID
 SOLUTIONS;THERMAL ANALYSIS;URANIUM NITRIDES: T2;VERY HIGH
 TEMPERATURE

ABSTRACT: The decomposition temperature of CeN,
 Usub(0.5)Cesub(0.5)N and Usub(0.75)Cesub(0.25)N has been
 measured as a function of nitrogen pressure. Thermodynamic

<581> CONT.

analysis based on the reaction, in which the solid solution, $U_{sub}(1-x)Ce_{sub}(x)N$, decomposes into $U_{sub}(1-x)Ce_{sub}(x)(1)$ and $1/2 N_{sub} 2(g)$, results in a very simple relation. The influence of solubility of nitrogen in $U_{sub}(1-x)Ce_{sub}(x)(1)$ has been also taken into consideration. The present experimental data are in good agreement with the equation derived from the present thermodynamic analysis.

EDB SUBJECT CATEGORY: EDB-360602
INIS CATEGORY: B12

<582>

Shulakov, A.S.; Zimkina, T.M.; Fomichev, V.A.; Neshpor, V.S.

X-ray spectral study of the carbides and nitrides of group IV and V metals EDB-77:050532; Fiz. Tverd. Tela (FTVTA) 18: (3):793-796 (Mar 1976)

SUBJECT DESCRIPTORS: CARBIDES: T2;HAFNIUM;NIOBIUM;NIOBIUM CARBIDES;NIOBIUM NITRIDES;NITRIDES: T3;TANTALUM;TITANIUM;TRANSITION ELEMENT COMPOUNDS: T1;VALENCE;VANADIUM;X-RAY SPECTRA: Q1,Q2,Q3;ZIRCONIUM

ABSTRACT: The very long-wave x-ray emission bands of Group IV and V transition metals reflecting the distribution of the d valence state were studied in the corresponding carbides and nitrides and the results were compared with those obtained for the pure metals (Ti, V, Nb, Hf, Ta, and Zr). For each group the structure of the d-type bands of the metals in monotypic compounds was very similar and depended mainly on the electron concentration. Special attention was paid to the cases of NbC and NbN, and the groups of electron states associated with various spatial bonds were distinguished. Analysis of the band shapes showed that in Nb and Zr some of the metal d electrons came under the influence of the lighter elements (C or N).

EDB SUBJECT CATEGORY: EDB-360204

<583>

Shurmovskaya, N.A.; Egorova, E.M.; Kornacheva, G.M. (AN SSSR, Moscow. Inst. Elektrokhimii)

Work function of gallium AIX-08:317782;EDB-77:120280; Elektrokhiimiya (ELKKA) 12: (4):590-591 (Apr 1976)

SUBJECT DESCRIPTORS: ELECTRODES;ELECTROMOTIVE FORCE;GALLIUM: T1;LIQUID METALS;STAINLESS STEELS;TANTALUM;VACUUM SYSTEMS;WORK FUNCTIONS: Q1

<583>

<583> CONT.

ABSTRACT: None

EED SUBJECT CATEGORY: EDB-360104

INIS CATEGORY: B12

<584>

Silbert, M. (East Anglia Univ. (UK). School of Mathematics and Physics) ** Evans, R.; Greenwood, D.A. (eds.)

Pressure and entropy calculations for liquid Na, K and Rb along the melting line in liquid metals, 1976 **

AIX-08:319612;EDB-77:113423; (1977) Institute of Physics

SUBJECT DESCRIPTORS: ENTROPY;EQUATIONS OF STATE: Q1,Q2,Q3;FREE ENERGY;LIQUID METALS;MELTING POINTS;POTASSIUM: T3;PRESSURE DEPENDENCE;RUBIDIUM: T2;SODIUM: T1;VARIATIONAL METHODS

ABSTRACT: The Gibbs-Bogoliubov variational principle is used to derive an equation of state for liquid metals, following a procedure indicated by Watabe et al (J.Phys.F:Metal Phys.;4L:29(1974)). Results for liquid Na, K and Rb along the melting curve are presented. It is found that for liquid Na there is overall agreement with experimental results up to 80 kbar, but for liquid K and Rb only up to 25 to 30 kbar.

EED SUBJECT CATEGORY: EDB-360102

INIS CATEGORY: A13

<585>

Silverman, M.D.; Huntley, W.R.; Robertson, H.E., Oak Ridge National Lab., Tenn. (USA)

In Heat transfer measurements in a forced convection loop with two molten-fluoride salts: $\text{LiF--BeF}_2\text{--ThF}_4$ and eutectic $\text{NaBF}_4\text{--NaF}$ **

INS-77:000119;ERA-02:009794;EDB-77:006325; (Oct 1976)

ORNL/TM--5335

SUBJECT DESCRIPTORS: BERYLLIUM FLUORIDES;COOLANTS: Q1;HEAT TRANSFER;LITHIUM FLUORIDES;MOLTEN SALT FUELS: T3;MOLTEN SALT REACTORS: T1;MOLTEN SALTS: T2;REACTOR COOLING SYSTEMS;SODIUM FLUORIDES;THERMODYNAMIC PROPERTIES: Q2,Q3;THORIUM FLUORIDES;URANIUM FLUORIDES

ABSTRACT: Heat transfer coefficients were determined experimentally for two molten-fluoride salts [$\text{LiF--BeF}_2\text{--ThF}_4$ (72-16-12-0.3 mole %) and $\text{NaBF}_4\text{--NaF}$ (92-8 mole %)] proposed as the fuel salt and coolant salt, respectively, for molten-salt breeder reactors. Information was obtained over a wide range of variables, with

<585> CONT.

salt flowing through 12.7-mm-OD (0.5-in.) Hastelloy N tubing in a forced convection loop (FCL-2b). Satisfactory agreement with the empirical Sieder-Tate correlation was obtained in the fully developed turbulent region at Reynolds moduli above 15,000 and with a modified Hausen equation in the extended transition region (Re approx. 2100-15,000). Insufficient data were obtained in the laminar region to allow any conclusions to be drawn. These results indicate that the proposed salts behave as normal heat transfer fluids with an extended transition region.

EDB SUBJECT CATEGORY: EDB-210500;360204
INIS CATEGORY: E35;B23

<586>

Silvester, L.F.; Pitzer, K.S.

Thermodynamics of electrolytes. 8. High-temperature properties, including enthalpy and heat capacity, with application to sodium chloride
INS-78:000307;ERA-03:008849;EDB-78:009960; J. Phys. Chem. (JPCCHA) 81:(19):1822-1828 (22 Sep 1977)

SUBJECT DESCRIPTORS: ELECTROLYTES;ENTHALPY: Q1;HIGH TEMPERATURE;MEDIUM TEMPERATURE;REACTION KINETICS;SODIUM CHLORIDES: T1;SPECIFIC HEAT: Q1;THERMODYNAMIC ACTIVITY: Q1;THERMODYNAMICS

ABSTRACT: The ion-interaction type of equation which has proven so successful for activity and osmotic coefficients of both pure and mixed electrolytes near room temperature is extended for wide temperature ranges and for the enthalpy and heat capacity. An improved equation is obtained for the dielectric constant of water over a wide range of temperature and density; it is used to calculate the Debye-Hueckel parameters to 350[°]C for enthalpy and heat capacity as well as for osmotic and activity coefficients. These equations are applied to the extensive array of thermodynamic data for aqueous sodium chloride from 0 to 300[°]C. Good agreement is obtained and the equations thus provide a convenient analytical representation of the thermodynamic properties of NaCl(aq). Also the results indicate that at 300[°]C sodium chloride is still a strong electrolyte with little tendency toward the formation of ion pairs.

EDB SUBJECT CATEGORY: EDB-400201;400400
INIS CATEGORY: E12

<587>

<587>

Sinha, V.K. (Indian Inst. of Tech., Kanpur)

Terminal solid solubility of hydrogen in the zirconium-2.5 weight % niobium alloy AIX-07:270433; EDB-77:030672; J. Chem. Soc. (London), Faraday Trans., I (JCFTA) 72: (pt.1):134-142 (1976)

SUBJECT DESCRIPTORS: HIGH TEMPERATURE; HYDRIDATION; HYDROGEN: T3; NIOBIUM ALLOYS: T1; PHASE STUDIES; SOLID SOLUTIONS; SOLUBILITY: Q3; SOLVENT PROPERTIES: Q1, Q2; VERY HIGH TEMPERATURE; ZIRCONIUM; ZIRCONIUM BASE ALLOYS: T2; ZIRCONIUM HYDRIDES

ABSTRACT: The terminal solid solubility (TSS) of hydrogen in the Zr-2.5 wt.% Nb alloy has been determined by measuring the decomposition pressures of several Zr-2.5 wt.% Nb alloys containing hydrogen in the temperature range 425 to 900°C. The TSS boundaries $\alpha/(\alpha + \delta)$ and $\alpha/(\alpha + \beta)$ were determined from the points of inflection of the van't Hoff decomposition pressure against temperature plots. The TSS of hydrogen in Zr-2.5 wt.% Nb alloy is about 1 to 2.5 atom % more than that in unalloyed zirconium (the maximum difference being at the eutectoid temperature of the Zr/H₂ system). The decomposition pressure against temperature relation in the α and $(\alpha + \delta)$ phase regions and the temperature against concentration relation along the TSS boundary have also been derived. The mean relative partial molar enthalpy of hydrogen in the α -phase and the mean enthalpy of formation of the δ -hydride from the saturated α -phase in the alloy are - 24.6 kcal(mol H₂)⁻¹ and - 40.9 kcal(mol H₂)⁻¹ respectively.

EDE SUBJECT CATEGORY: EDB-360102

INIS CATEGORY: B22

<588>

Skrinichenko, T.M.; Klibanov, E.L.; Kashin, V.I.

Investigation into outgassing from molybdenum during melting
EDE-76:036209; Fiz. Khim. Obrab. Mater. (FKOMA) (1):40-43
(Jan 1975)

SUBJECT DESCRIPTORS: DEGASSING: Q1; DESORPTION; DIFFUSION; DROPLETS; ELECTRIC ARCS; EQUATIONS; FILMS; HYDRODYNAMICS; HYDROGEN; LIQUID METALS; MASS TRANSFER; MOLYBDENUM: T1; OXYGEN; REFINING; REYNOLDS NUMBER; VACUUM MELTING

<588> CONT.

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-360101

INIS CATEGORY: B21

<589>

Skripko, G.A.; Makhanek, A.G.; Gintoft, R.I.

Structure of $4f^{6s}$ and $4f^{8s}$ levels and Eu^{3+} and Tb^{3+} configurations in short-range ultra-violet
AIX-C8:280690;EDB-77:031506; Zh. Prikl. Spektrosk. (ZPSBA)
24: (6):968-971 (Jun 1976)

SUBJECT DESCRIPTORS: CALCIUM FLUORIDES;COMPARATIVE
EVALUATIONS;CRYSTALS;ELECTRONIC STRUCTURE: Q1,Q2;ENERGY
LEVELS;EUROPIUM COMPOUNDS:
T1;LASERS;LUMINESCENCE;SPECTRA;TERBIUM COMPOUNDS: T2

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-400201

INIS CATEGORY: B12

<590>

Slcan, S.A. ** Cooper, M.H. (ed.), American Nuclear Society,
Hinsdale, Ill.; American Inst. of Mining, Metallurgical, and
Petroleum Engineers, Inc., New York; Energy Research and
Development Administration, Washington, D.C. (USA); American
Nuclear Society, Pittsburgh, Pa. Pittsburgh Section

Reaction of potassium superoxide in excess potassium In
Proceedings of the international conference on liquid
technology in energy production **
INS-77:016019;ERA-02:059704;EDB-77:139579; (Nov 1976)
CONF-760503--P2

SUBJECT DESCRIPTORS: CHEMICAL REACTION KINETICS:
Q3;COCLANTS;EUTECTICS;EXPLOSIONS: Q1,Q2;LIQUID METALS;LMFBR
TYPE REACTORS: T1;POTASSIUM ALLOYS: T2;POTASSIUM OXIDES:
T3;SIMULATION;SODIUM ALLOYS;TEMPERATURE DEPENDENCE

ABSTRACT: Experiments are described in which liquid potassium
and potassium superoxide are brought into contact at
different temperatures. There is no obvious reaction below
390 K, a controlled reaction between 390 K and 450 K, and an
explosive reaction occurs when the reactants are mixed at
higher temperatures. If the reactants are mixed at a low
temperature and heated at 100 K min^{-1} , an explosion
occurs. When heated at 3 K min^{-1} , the potassium quietly
ingests the superoxide. The results are discussed in terms of

<590>

<590> CONT.

a thermal explosion model. The explosions occur in the absence of organic contaminants. The system is not critically sensitive to shock or agitation.

EDB SUBJECT CATEGORY: EDB-220900;210500;400201

INIS CATEGORY: C52;E35;B12

<591>

Smith, D.F.; Miller, W.C.; Taylor, A.R. Jr., Bureau of Mines, Tuscaloosa, Ala. (USA)

In High-temperature heat contents and entropies of zirconium fluocride and zirconium sulfate ** EDB-77:101119; (May 1961)
PM-RI--5964

SUBJECT DESCRIPTORS: ENTHALPY;ENTROPY;HIGH TEMPERATURE;MEDIUM TEMPERATURE;TEMPERATURE DEPENDENCE;THERMODYNAMIC PROPERTIES: Q1,Q2;VERY HIGH TEMPERATURE;ZIRCONIUM FLUORIDES: M1;ZIRCONIUM SULFATES: M2

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-400201

<592>

Smotrakov, V.G.; Popovskaya, N.P.; Protsenko, P.I.
(Rostovskij-na-Donu Gosudarstvennyj Univ. (USSR))

Viscosity of binary of nitrate systems melts
AIX-08:334371;EDB-77:140978; Zh. Fiz. Khim. (ZFKHA)
50: (12):3087-3090 (Dec 1976)

SUBJECT DESCRIPTORS: ALKALI METAL COMPOUNDS;BINARY MIXTURES;CADMIUM NITRATES: T1;CADMIUM NITRATES: T4;CALCIUM NITRATES: T2;MOLTEN SALTS;NITRATES;QUANTITY RATIO;SILVER NITRATES: T5;THALLIUM NITRATES: T3;VISCOSITY: Q1,Q2,Q3,Q4,Q5

ABSTRACT: The systematic studies results of the physical and chemical properties of the binary mixtures melts of univalent metals nitrates with calcium and cadmium nitrates have been generalized. It has been noted that the concentration dependence in such compounds becomes more complicated, which is manifested by an increase in the deviation from additivity with the increasing difference in the energy characteristics of cations. The somewhat greater deviations from additivity observed in the sequence of cadmium systems may be explained by the more significant structural changes taking place in melts which contain the cadmium ions. The relationship

<592> CONT.

between the deviation of viscosity from additivity and the values the accessive modal volume is discussed.

EDE SUBJECT CATEGORY: EDB-400201

INIS CATEGORY: B12

<593>

Sobolev, B.P.; Sejranyan, K.B. (AN SSSR, Moscow. Inst. Kristallografii; Erevanskii Gosudarstvennyi Univ. (USSR))

Phases of variable composition with $\text{LaF}_{\text{sub } 3}$ structure in $\text{MF}_{\text{sub } 2}$ -(Y, Ln) $\text{F}_{\text{sub } 3}$ systems. 1. $\text{SrF}_{\text{sub } 2}$ -(Y, Ln) $\text{F}_{\text{sub } 3}$ systems (thermal characteristics, formation of berthollides) AIX-07:265104; EDB-77:013488; Kristallografiya (KRISA) 20:(4):763-768 (Jul 1975)

SUBJECT DESCRIPTORS: DIFFERENTIAL THERMAL ANALYSIS; PHASE DIAGRAMS; PHASE STUDIES: Q1, Q2, Q3; PHASE TRANSFORMATIONS; RARE EARTH COMPCUNDS: T3; SOLID SOLUTIONS; STRONTIUM FLUORIDES: T1; X-RAY DIFFRACTION; YTTRIUM FLUORIDES: T2

ABSTRACT: Using thermal and X-ray diffraction analysis, portions of the phase diagrams have been studied of 13 systems of $\text{SrF}_{\text{sub } 2}$ -(Y, Ln) $\text{F}_{\text{sub } 3}$ in the region containing 60-100 mol. % (Y, Ln) $\text{F}_{\text{sub } 3}$. In all the systems the solid solutions $\text{Ln}_{\text{sub } 1-x}\text{Sr}_{\text{sub } x}\text{F}_{\text{sub } 3-x}$ are formed of the $\text{LaF}_{\text{sub } 3}$ -type structure. As a result of a morphotropic transition in the $\text{LnF}_{\text{sub } 3}$ -series these phases get isolated acquiring all characteristic features of berthollides. The observed relation of the berthollide phase with solid solutions based on actually existing tiscnite modifications ($\text{LaF}_{\text{sub } 3}$ -type) of rare earth trifluorides uniquely determines the relationship of berthollides to the type of solid solutions on the base of "imaginary" modifications of components. The berthollide phases studied refer to a new variety characterized by a congruent melting which is stipulated by the stabilization of the $\text{LaF}_{\text{sub } 3}$ -type structure with heterovalent isomcrphic substitutions. The coordinates of the maxima have been given in the phase diagrams of the tiscnite phases. In the systems of trifluorides of yttrium and rare earths (erbium through ytterbium) solid solutions have been discovered based on high-temperature modifications $\alpha\text{-LnF}_{\text{sub } 3}$ (structural type $\alpha\text{-UO}_{\text{sub } 3}$).

EDE SUBJECT CATEGORY: EDB-400201

INIS CATEGORY: B12

<594>

<594>

Sommers, J.A.; Westrum, E.F. Jr. (Michigan Univ., Ann Arbor (USA). Dept. of Chemistry)

Thermodynamics of the lanthanide halides. II. Heat capacities and Schottky anomalies of SmCl_3 , EuCl_3 and GdCl_3 from 50 to 350 K AIX-08:305623; EDB-77:107385; J. Chem. Thermodyn. (JCTDA) 9:(1):1-26 (Jan 1977)

SUBJECT DESCRIPTORS: CALORIMETRY; ENTROPY; EUROPTIUM CHLORIDES: T1; GADOLINIUM CHLORIDES: T2; LOW TEMPERATURE; MEDIUM TEMPERATURE; SAMARIUM CHLORIDES: T3; SPECIFIC HEAT: Q1, Q2, Q3; TEMPERATURE DEPENDENCE; ULTRALOW TEMPERATURE; VERY LOW TEMPERATURE

ABSTRACT: The heat capacities of SmCl_3 , EuCl_3 , and GdCl_3 have been measured from 5 to 350 K by adiabatic calorimetry. For SmCl_3 and EuCl_3 , the calculated Schottky heat capacities may be compared with the difference in heat capacity between LaCl_3 and the paramagnetic members, as before. For GdCl_3 which lacks a Schottky contribution, the difference between the heat capacity of LaCl_3 and GdCl_3 may be discussed in terms of spectroscopic data on lattice vibration. Comparison between the measured entropies and the results of estimation schemes is presented. The values of $(S^\circ(298.15 \text{ K}) - S^\circ(0))$ for SmCl_3 , EuCl_3 , and GdCl_3 are 35.88, 34.43 and 36.19 cal/(mol K) respectively.

EDB SUBJECT CATEGORY: EDB-400201
INIS CATEGORY: E12

<595>

Sproule, R.T.

Zirconia and yttria-thoria ceramics find new uses as solid electrolytes EDB-77:043916; Mater. Eng. (MAENB) 79:(1):48-49 (1974)

SUBJECT DESCRIPTORS: CERAMICS; ELECTRIC CONDUCTIVITY; ELECTRICAL PROPERTIES; ELECTROLYTES: Q4; HIGH TEMPERATURE; HIGH-TEMPERATURE FUEL CELLS: T4; MASS TRANSFER; OXYGEN IONS; THORIUM OXIDES: T3; USES: Q1, Q2, Q3; YTTRIUM OXIDES: T2; ZIRCONIUM OXIDES: T1

ABSTRACT: Zirconia and yttria-thoria are electrical insulators at low temperatures, but become electrical conductors at elevated temperatures. Their high-temperature conductivity can be controlled by varying the type and amount of stabilizer such as calcia, yttria, or magnesia. O^{2-} ions will pass through the zirconia lattice: it is impervious to all

<595> CONT.

other gases. This property is used in devices for determining O in gases, liquids, and metals. It also makes it useful in fuel cells.

EDB SUBJECT CATEGORY: EDB-360204;300503

<596>

Steffens, F.; Reinen, D. (Marburg Univ. (F.R. Germany).
Fachbereich Chemie)

Preparation and structure of transition metal fluorides
 $UE_{sup}(II)M_{sup}(IV)F_{sub 6}[UE_{sup}(II) : Cr-Zn; M_{sup}(IV) : Hf, Zr]$ AIX-07:274884; EDB-77:031503; Z. Naturforsch., b
 (ZENEA) 31:(6):894-896 (Jun 1976)

SUBJECT DESCRIPTORS: CHEMICAL BONDS; CHEMICAL
 PREPARATION; COMPARATIVE EVALUATIONS; CRYSTAL STRUCTURE:
 Q1, Q2; HAFNIUM FLUORIDES: T1; JAHN-TELLER EFFECT; LATTICE
 PARAMETERS; TRANSITION ELEMENT COMPOUNDS; X-RAY
 DIFFRACTION; ZIRCONIUM FLUORIDES: T2

ABSTRACT: Compounds $UE_{sup}(II)M_{sup}(IV)F_{sub 6}[UE : Mn-Ni, Zn]$
 crystallize in the $LiSbF_{sub 6}$ - or in the ordered $ReO_{sub 3}$ -
 structure. The influence of the Jahn-Teller ions $Cr_{sup} 2+$
 and $Cu_{sup} 2+$ on these structure types is studied.

EDE SUBJECT CATEGORY: EDB-400201

INIS CATEGORY: B12

<597>

Steiger, R.P.; Cater, E.D.

Mass spectrometric determination of dissociation energies of
 high-temperature molecules: ScS, YS, LaS, ZrS, UO
 INS-76:014749; ERA-01:020217; EDB-76:054358; High Temp. Sci.
 (HITSA) 7:(4):288-300 (Dec 1975)

SUBJECT DESCRIPTORS: DISSOCIATION ENERGY:

Q1, Q2, Q3, Q4, Q5; GASES; LANTHANUM SULFIDES: T3; MASS
 SPECTROSCOPY; MOLECULES; SCANDIUM SULFIDES: T1; URANIUM OXIDES:
 T5; YTTRIUM SULFIDES: T2; ZIRCONIUM SULFIDES: T4

ABSTRACT: Gaseous isomolecular equilibria of types $MS + N = NS + M$
 and $MO + N = NO + M$, where M and N are variously the
 elements Sc, Y, La, Zr, and U, were studied mass
 spectrometrically, over appropriate mixed solid phases heated
 in a tungsten Knudsen cell. Dissociation energies were
 derived, using as reference values $D_{sub 0}^{sup 0}(US) = 124.0 \pm 2.3$,
 $D_{sub 0}^{sup 0}(YS) = 125.4 \pm 2.5$, and $D_{sub 0}^{sup 0}(YO) = 170.0 \pm 3.0$ kcal/mole.
 "Preferred values"

<597>

<597> CONT.

obtained by considering the results of the present study and literature values are ScS, 113.4 $\pm$ 3.0; YS, 125.4 $\pm$ 2.5; LaS, 136.0 $\pm$ 3.0; ZrS, 136.6 $\pm$ 4.0; and UO, 181.0 $\pm$ 4.0 kcal/mole. Determination of ground state degeneracies of US and UO by calculating the electronic entropies of the molecules was attempted. 4 tables, 33 references.

EDB SUBJECT CATEGORY: EDB-400201;400702;360204

INIS CATEGORY: P12;P13;P23

<598>

Steiner, P.; Hoechst, H.; Huefner, S. (Universitaet des Saarlandes, Saarbruecken (Germany, F.R.))

XPS valence bands of La, Ce and Gd and their aluminium alloys
AIX-08:319619;EDB-77:113442; J. Phys., F (London) (JPFMA)
7:(5):L145-L149 (May 1977)

SUBJECT DESCRIPTORS: ALUMINIUM ALLOYS: T1;BAND
THEORY;CERIUM;CERIUM ALLOYS: T2;ELECTRONIC STRUCTURE:
Q1,Q2,Q3,Q4;FERMI LEVEL;GADOLINIUM;GADOLINIUM ALLOYS:
T3;LANTHANUM;LANTHANUM ALLOYS: T4;PHOTOELECTRON
SPECTROSCOPY;SPECTRA;VALENCE;X RADIATION

ABSTRACT: The XPS valence band spectra of La, Ce and Gd metal and of their aluminium alloys show the 5d band in La metal at 0.8 eV below the Fermi energy with a width of about 1.5 eV; in Ce the 4f band and 5d band are at 0.5 eV and 1.7 eV below the Fermi energy with widths of 0.7 eV and 1.5 eV respectively. The Gd 5d band is very similar to the La 5d band, while the 4f level is well localized at 8.0 eV below the Fermi energy.

EDB SUBJECT CATEGORY: EDB-360102

INIS CATEGORY: A13

<599>

Stepanov, V.G.; Kozhevnikov, V.F. (Moskovskij Aviatsionnyj Inst. (USSR))

Density and enthalpy of liquid alkaline metals at pressures up to 1500 bar AIX-09:349543;EDB-78:027597; Inzh.-Fiz. Zh.
(INFZA) 30:(5):855-862 (May 1976)

SUBJECT DESCRIPTORS: CESIUM;DENSITY: Q1;ENTHALPY;FREE
ENERGY;HIGH PRESSURE;HIGH TEMPERATURE;LIQUID METALS:
T1;LITHIUM;MEDIUM TEMPERATURE;POTASSIUM;PRESSURE
DEPENDENCE;RUBIDIUM;SODIUM;TEMPERATURE DEPENDENCE;VERY HIGH
PRESSURE

<599> CONT.

ABSTRACT: On the basis of the similarity principle justified theoretically for liquid alkali metals, generalized expressions for thermophysical characteristics have been derived. From data for rubidium density and enthalpy of liquid lithium, sodium, potassium are calculated at pressures up to 1500 bars and those of liquid cesium up to 1000 bars.

EDB SUBJECT CATEGORY: EDB-360104

INIS CATEGORY: E22

<600>

Stephens, F.B.; Gutmacher, R.G.; Ernst, K.; Harrar, J.E.,
California Univ., Livermore (USA). Lawrence Livermore Lab.

In Methods for the accountability of uranium dioxide **
INS-76:014263; ERA-01:018796; EDB-76:057861; (Jun 1975)
PB--241763

SUBJECT DESCRIPTORS: ACCOUNTING; CHEMICAL
ANALYSIS; DISSOLUTION; IMPURITIES; MASS SPECTROSCOPY; NUCLEAR
MATERIALS MANAGEMENT: Q1; SAMPLING; URANIUM DIOXIDE: T1

ABSTRACT: This report deals with methods for the analysis of uranium dioxide and contains sections describing: (1) sampling and sample handling, (2) dissolution of samples, (3) the determination of impurities in the matrix material, (4) isotopic analysis by surface ionization mass spectrometry, (5) chemical analysis, and (6) error analysis. Each section deals in sufficient detail with the subject matter to allow an experienced analyst to implement the procedures in a reasonably well-equipped analytical laboratory. (GRA)

EDB SUBJECT CATEGORY: EDB-055001

INIS CATEGORY: F42

<601>

Stievenart, M.; Bouffloux, P.; Egleme, M.; Fabry, J.P.; Lamotte, H. (Societe Belge pour l'Industrie Nucleaire, Brussels) **
Jaeger, T.A. (comp.)

Analysis of LMFBR explosion model experiments by means of the SUBPOUM-II code in Structural mechanics in reactor technology. Vol. 2 **
AIX-08:287060; ERA-02:024800; EDB-77:049734; (1975)
North-Holland

SUBJECT DESCRIPTORS: BOUNDARY CONDITIONS; COMPUTER
CALCULATIONS; DEFORMATION; INCOMPRESSIBLE FLOW; REACTOR CORE
DISRUPTION: T2, Q1; S CODES; SIMULATION: Q2; SNR-1 REACTOR:
T1; TWO-DIMENSIONAL CALCULATIONS

<601>

<601> CONT.

ABSTRACT: Experiments have been carried out to simulate in small scale vessels the occurrence of an Hypothetical Core Disruptive Accident (HCDA) for the SNR-300 reactor. In parallel with the experimental programme, the (2D) computer code SURBOUM-II has been developed. The code computes the two-dimensional fluid flow within the system in case of a core explosion. The fluid is assumed incompressible and the deformations of the concrete shell(s) and vessel(s) are calculated by means of the thin shell theory. Perforated dip plates which are included in the model are treated as a particular type of boundary conditions involving empirical pressure drop versus flow relationships. The first part of the interpretation work was the determination of the pressure-volume relationship of the slow burning charge used to simulate the HCDA. This has been achieved by an original trial and error method built in the code which fits best the experimental impulse time records obtained in bare charge experiments fired in overstrong vessels. Other experiments carried out in overstrong vessels and involving a perforated dip plate above the core to damp out the fluid impact on the roof were calculated and the comparison of the theoretical and experimental impulse time curves was satisfactory. Further experiments, with and without the perforated dip plate, carried out in yielding vessels were also interpreted. For those experiments the scope of the work was the comparison of calculated and measured deformation of the vessel. The agreement obtained is satisfactory, through the code seems to overestimate slightly the final deformation.

EEB SUBJECT CATEGORY: EDB-220900;210500
 INIS CATEGORY: C52

<602>

Storonkin, A.V.; Vasil'kova, I.V.; Efimov, A.I.; Baruzdina, V.A.

Thermographic investigation of the $\text{PbCl}_{\text{sub } 2\text{f}}\text{-PbBr}_{\text{sub } 2\text{f}}\text{-PbI}_{\text{sub } 2\text{f}}$ and $\text{PbCl}_{\text{sub } 2\text{f}}\text{-PbBr}_{\text{sub } 2\text{f}}\text{-PbF}_{\text{sub } 2\text{f}}$ systems
 AIX-08:317792;EDB-77:133394; Vestn. Leningr. Univ., Fiz.,
 Khim. (VLUFB) (16):137-139 (Aug 1975)

SUBJECT DESCRIPTORS: DIFFERENTIAL THERMAL
 ANALYSIS;EUTECTICS;LEAD BROMIDES: T1;LEAD CHLORIDES: T2;LEAD
 FLUORIDES: T4;LEAD IODIDES: T3;PHASE DIAGRAMS:
 Q1,Q2,Q3,Q4;SOLID SOLUTIONS

ABSTRACT: None

EEB SUBJECT CATEGORY: EDB-400201
 INIS CATEGORY: B12

<603>

Storozhenko, T.P.; Khanaev, E.I.; Afanas'ev, Yu.A. (Kubanskiy Gosudarstvennyy Univ., Krasnodar (USSR))

Formation enthalpy of gadolinium, terbium and dysprosium trifluorides AIX-08:292491; EDB-77:069064; Zh. Fiz. Khim. (ZFKHA) 50:(4):1060 (Apr 1976)

SUBJECT DESCRIPTORS: CALORIMETRY; DYSPROSIUM FLUORIDES: T1; FORMATION HEAT: Q1, Q2, Q3; GADOLINIUM FLUORIDES: T2; TERBIUM FLUORIDES: T3

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-400201

INIS CATEGORY: P12

<604>

Streicher, M.A.

Effect of composition and structure on crevice, intergranular, and stress corrosion of some wrought Ni--Cr--Mo alloys
INS-76:016231; EDB-76:053892; Corrosion (CORRA) 32:(3):79-93 (Mar 1976)

SUBJECT DESCRIPTORS: CHEMICAL COMPOSITION; CORROSIVE EFFECTS: Q2, Q3; HASTELLOYS: T1; INORGANIC ACIDS: T2; INTERGRANULAR CORROSION: Q1; INTERMETALLIC COMPOUNDS; MAGNESIUM CHLORIDES: T3; MICROSTRUCTURE; PRECIPITATION HARDENING; STRESS CORROSION: Q1; VERY HIGH TEMPERATURE

ABSTRACT: The relationship of composition and microstructure to the occurrence of localized corrosion in Hastelloy alloys C, C-276, and C-4 was investigated. One hour exposures of these alloys in the range of 1200 to 2200°F may result in the formation of a molybdenum-rich intermetallic compound and, in alloys with more than about 0.004 percent C, a molybdenum-rich carbide. The boiling ferric sulfate-50 percent H₂SO₄ test readily detects the presence of both of these precipitates which cause rapid intergranular attack in this solution. The molybdenum-rich M₆C carbide precipitate impairs resistance to intergranular, crevice, and stress corrosion. In contrast, the molybdenum-rich intermetallic compound, Mu-phase, impairs resistance only to intergranular attack, primarily in oxidizing acids. When the carbon content is progressively increased above 0.004 percent, formation of the molybdenum-rich carbide precipitate at grain boundaries by one hour heat treatments results first in loss of resistance to stress corrosion cracking (SCC) in the boiling 45 percent MgCl₂ test. Further increases in carbon content (0.05 percent) result in loss of resistance to intergranular attack

<604>

<604> CONT.

in acids, to crevice corrosion in 10 percent ferric chloride at 50⁰C and also to SCC in this solution. All SCC is of the intergranular type. Formation of the molybdenum-rich carbide precipitate is prevented in alloy C-4 by keeping the carbon content low and by addition of titanium. Formation of the intermetallic compound is minimized by adjustment of the alloy content. In the heat investigated, titanium stabilization was effective in preventing intergranular attack in reducing acids and SCC in the $MgCl_2$ test. However, metallographic examination and intergranular attack in the ferric sulfate test on material heated 1 hour at 1600⁰F indicated that a compound was still being formed at grain boundaries.

EDB SUBJECT CATEGORY: EDB-360105

INIS CATEGORY: B15

<605>

Stremousov, V.I.; Rubtsov, A.S.

Thermophysical properties of the ternary liquid-metal system tin--lead--cadmium ERA-02:001027; EDB-76:081560; High Temp. (USSR) (Engl. Transl.) (HITEA) 13: (6):1201-1203 (Nov 1975)

SUBJECT DESCRIPTORS: CADMIUM ALLOYS: T1; COMPRESSIBILITY: Q1, Q2, Q3; EUTECTICS; LEAD ALLOYS: T3; MEDIUM TEMPERATURE; SPECIFIC HEAT: Q1, Q2, Q3; TIN ALLOYS: T2; ULTRASONIC WAVES

ABSTRACT: Data on the velocity of ultrasound and adiabatic compressibility are used to calculate specific heats and isothermal compressibility of Pb-Sn-Cd alloys. (AIP) (AIP)

EDB SUBJECT CATEGORY: EDB-360104

<606>

Stremousov, V.I.; Rubtsov, A.S. (Volgogradskij Pedagogicheskij Inst. (USSR))

Thermodynamic properties of the ternary Sn-Pb-Cd liquid metal system AIX-08:324282; EDB-77:132712; Teplofiz. Vys. Temp. (TVYTA) 13: (6):1298-1300 (Nov 1975)

SUBJECT DESCRIPTORS: CADMIUM ALLOYS: T1; CHEMICAL COMPOSITION; COMPRESSIBILITY; EQUATIONS; LEAD ALLOYS: T2; LIQUID METALS; MEASURING METHODS; SPECIFIC HEAT; TEMPERATURE DEPENDENCE; TERNARY ALLOY SYSTEMS; THERMODYNAMIC PROPERTIES: Q1, Q2, Q3; TIN ALLOYS: T3; ULTRASONIC WAVES

<606> CONT.

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-360104

INIS CATEGORY: B22

<607>

Stuve, J.M., Bureau of Mines, Reno, Nev. (USA). Reno Metallurgy Research Center

In Heat of formation of europium sesquioxide and europium trichloride ** EDB-77:107188; (Oct 1964) BM-RI--6640

SUBJECT DESCRIPTORS: CALORIMETRY;EUROPIUM CHLORIDES: T2;EUROPIUM OXIDES: T1;FORMATION HEAT: Q1,Q2;MEDIUM TEMPERATURE

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-360204

<608>

Stuve, J.M., Bureau of Mines, Reno, Nev. (USA). Reno Metallurgy Research Center

In Heats of formation of holmium and terbium trichlorides ** EDE-77:120761; (Nov 1967) BM-RI--7046

SUBJECT DESCRIPTORS: FORMATION HEAT: Q1,Q2;HOLMIUM CHLORIDES: T1;SOLUTION HEAT;TERBIUM CHLORIDES: T2

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-400201

<609>

Suzuki, K. (Tohoku Univ., Sendai (Japan). Research Inst. for Iron, Steel and Other Metals)

Structural study of liquids with strong short-range correlation in the atomic distribution AIX-08:333464;EDB-78:022010; Ber. Bunsenges. Phys. Chem. (BBPCA) 80:(8):689-694 (Aug 1976)

SUBJECT DESCRIPTORS: AMORPHOUS STATE;ATOMIC MODELS;BOND LENGTHS;CARBON SULFIDES: T3;COMPARATIVE EVALUATIONS;CORRELATIONS;CRYSTAL STRUCTURE;DISTRIBUTION;INTERACTION RANGE;LIQUID METALS;MOLECULAR STRUCTURE;NEUTRON DIFFRACTION: Q1,Q2,Q3;PALLADIUM SILICIDES: T2;SELENIUM: T1;SOLIDS;SPECTRA;STRUCTURAL MODELS;TIME-OF-FLIGHT METHOD

<609>

<609> CONT.

ABSTRACT: Structure factors of liquids and amorphous solids having a relatively high degree of ordering in their short-range structures have been measured over a wide range of scattering vector by means of the T-O-F neutron diffraction using epithermal pulsed neutrons generated by a linear electron accelerator. It has been shown in the case of CS_2 liquid that the size and shape of a molecule existing in the liquid are determined from the behaviour of the structure factor in the range of high scattering vector, and that the structure factor at low scattering vectors then informs on inter-molecular orientational and center-center correlations in the liquid state. Moreover, based on highly-resolved radial distribution functions, a free rotating chain model has been discussed for chain molecules contained in liquid Se and a splitting of the nearest neighbour Pd-Pd and Pd-Si correlation has been clearly found in amorphous $\text{Pd}_{0.8}\text{Si}_{0.2}$ alloy.

EDB SUBJECT CATEGORY: EDB-360602

INIS CATEGORY: A13

<610>

Svechnikov, V.N.; Kobzenko, G.F.; Ivanchenko, V.G. (AN Ukrainskoj SSR, Kiev. Inst. Metallofiziki)

Gd, Tb, Dy, Ho, Er solubility in chromium

AIX-09:349545; EDB-78:027614; Metallofizika (MFIZA) (59):77-83 (1975)

SUBJECT DESCRIPTORS: CHROMIUM: T1; DOPED

MATERIALS; INTERSTITIALS; MICROHARDNESS; MICROSTRUCTURE; PHASE TRANSFORMATIONS; RARE EARTHS: T2; SOLID SOLUTIONS; SOLUBILITY: Q1, Q2; TRANSITION TEMPERATURE; VERY HIGH TEMPERATURE

ABSTRACT: Using metallographic, differential, thermal and X-ray crystal analysis and microhardness methods, the authors determined the limiting solubility in chromium of five rare-earth metals of the yttrium sub-group at the temperatures of monotectic transformation and at 1700deg, 1600deg, 1400deg and 900degC. The influence of the quantity of alloying additive on the microhardness of the chrome-based solid solution and on the microstructure and hardness of the alloys was investigated. The temperatures of monotectic and eutectic transformation were determined. The changes in the physical properties of the solid solution as a function of

<611> CONT.

composition are explained on the basis of notions of phase equilibria in multi-component systems.

EDB SUBJECT CATEGORY: EDB-360104

INIS CATEGORY: B22

<611>

Tagawa, H. (Japan Atomic Energy Research Inst., Tokai, Ibaraki.
Tokai Research Establishment)

Nonstoichiometry in cubic uranium sesquinitride In U.S.-Japan
joint seminar on defects and diffusion in solids **

AIX-08:320731;EDB-77:114344; (1976) Japan Society for
Promotion of Science

SUBJECT DESCRIPTORS: CHEMICAL COMPOSITION; CRYSTAL DEFECTS:
C1; CRYSTAL-PHASE TRANSFORMATIONS; CUBIC LATTICES; DENSITY; HIGH
TEMPERATURE; LATTICE PARAMETERS; NEUTRON DIFFRACTION; PHASE
DIAGRAMS; PRESSURE DEPENDENCE; THERMODYNAMIC PROPERTIES; URANIUM
NITRIDES: T1; VERY HIGH TEMPERATURE; X-RAY DIFFRACTION

ABSTRACT: Of four phases in the U-N system, the cubic
sesquinitride $\alpha\text{-U}_{2/3}\text{N}_{3/2}$ exhibits a wide
nonstoichiometric range of composition at high temperatures.
The defect characteristic of $\alpha\text{-U}_{2/3}\text{N}_{3/2}$ has
been examined from the standpoint of crystal structure and
thermodynamic properties. Two separate linear relationships
between lattice parameter and nitrogen content are obtained:
one is characteristic of $\text{U}_{2/3}\text{N}_{3/2}$ and the other of
 $\text{UN}_{1/2}$. Using the density data, it is deduced that the
nonstoichiometry in the $\text{U}_{2/3}\text{N}_{3/2}$ region is caused by
the interstitial nitrogen atoms, and the nonstoichiometry in
 $\text{UN}_{1/2}$ is due to the nitrogen vacancies. X-ray and neutron
diffraction analyses also show that the excess nitrogen atoms
fill the vacant interstitial positions without the structure
change. Thermodynamic data suggest that the excess nitrogen
is dissolved in $\text{U}_{2/3}\text{N}_{3/2}$ with the wide homogeneous
range of composition; the results are consistent with those
obtained from the measurements for lattice parameter and
density, and from the structure analysis.

EDB SUBJECT CATEGORY: EDB-400702;360202

INIS CATEGORY: B12

<612>

<612>

Tagawa, H.; Fujino, T.; Tateno, J., Japan Atomic Energy Research
Inst., Tokyo

In Phase relations, crystal structures and physical properties
of nuclear fuels **

AIX-08:281013; ERA-02:032849; EDB-77:074802; (Jul 1975)

JAERI-M--6180

SUBJECT DESCRIPTORS: CRYSTAL STRUCTURE; ELECTRICAL
PROPERTIES; MAGNETIC PROPERTIES; MELTING POINTS; PHASE
DIAGRAMS; PHYSICAL PROPERTIES:
Q1, Q2, Q3, Q4, Q5, Q6, Q7, Q8, Q9, Q10, Q11, Q12, Q13, Q14, Q15; PLUTONIUM
CARBIDES: M1; PLUTONIUM NITRIDES: M2; PLUTONIUM OXIDES:
M3; PLUTONIUM PHOSPHIDES: M4; PLUTONIUM SULFIDES:
M5; REVIEWS; SELF-DIFFUSION; THERMAL EXPANSION; THORIUM CARBIDES:
M6; THORIUM NITRIDES: M7; THORIUM OXIDES: M8; THORIUM
PHOSPHIDES: M9; THORIUM SULFIDES: M10; URANIUM CARBIDES:
M11; URANIUM NITRIDES: M12; URANIUM OXIDES: M13; URANIUM
PHOSPHIDES: M14; URANIUM SULFIDES: M15

ABSTRACT: Phase relations, crystal structures and physical
properties of the compounds for nuclear fuels are presented,
including melting point, thermal expansion, diffusion and
magnetic and electric properties. Emphasis is on oxides,
carbides and nitrides of thorium, uranium and plutonium.

EDB SUBJECT CATEGORY: EDB-360204; 360202; 220300

INIS CATEGORY: B23

<613>

Takahashi, Yoichi; Kamimoto, Masayuki (Tokyo Univ. (Japan)).
Faculty of Engineering)

Thermal properties of nuclear fuel ceramics EDB-76:054019;
Seramikku (SERAA) 10: (5): 304-310 (May 1975)

SUBJECT DESCRIPTORS: CERAMICS; NUCLEAR FUELS; PLUTONIUM OXIDES:
T1; SOLID SOLUTIONS; SPECIFIC HEAT; TEMPERATURE
DEPENDENCE; THERMAL CONDUCTIVITY; THERMODYNAMIC PROPERTIES:
Q1, Q2, Q3, Q4; THORIUM OXIDES: T2; URANIUM CARBIDES: T3; URANIUM
OXIDES: T4

ABSTRACT: The contribution of electronic structure to thermal
properties was studied by examining the heat capacity and
heat conductivity of ceramics as nuclear fuel material. The
heat capacity of NaCl type uranium compounds at low
temperature and the relationship between state density and
the coefficient of electronic heat capacity are illustrated.
Further, the heat capacity of UC, $U_{2/3}C_{1/3}$ and FuC at
high temperature is shown. The deviation of the heat capacity
of UC and $U_{2/3}C_{1/3}$ from the ordinary temperature

<613> CONT.

dependence of heat capacity has been explained as dependent on the formation of lattice defects in crystals, whereas the deviation of that of PuC has to be examined still further. The heat conductivity of UO_2 is small. The heat conductivity of NaCl type uranium compounds is in an intermediate range between metallic uranium and UO_2 . It is markedly low at low temperature, but increases with temperature rise, and shows large value as compared with UO_2 at high temperature. Therefore, the NaCl type uranium compounds are promising as nuclear fuel ceramics for future. The heat conductivity of UO_2 is reviewed on the basis of phonon concept. Attention is paid to the localized 5 f electrons of UP and US. As a result, the behavior of the heat conductivity at transition point differs considerably from that of UN. The research on solid solutions such as $(\text{U,Pu})\text{O}_2$ and $\text{U}(\text{P,S})$ are reviewed. Finally, the method of measuring thermal property by laser flash is briefly described.

EDB SUBJECT CATEGORY: EDB-360204
INIS CATEGORY: B23

<614>

Talbot, J.B.; Smith, F.J.; Land, J.F.; Barton, P. (Oak Ridge National Lab., Tenn. (USA))

Tritium sorption in lithium--bismuth and lithium--aluminum alloys AIX-08:295850; ERA-02:027007; EDB-77:055610; J. Less-Common Met. (JCOMA) 50:(1):23-28 (Nov 1976)

ABSTRACT: The sorption of tritium by molten lithium-bismuth, Li-Bi (15 at.% lithium), and solid equiatomic lithium-aluminum, Li-Al, was investigated to evaluate the potential application of both materials in controlled thermonuclear reactors. The solubility of tritium in molten Li-Bi is less than 0.1 ppb at 500 - 700 $^{\circ}\text{C}$ and tritium partial pressures of 10^{-1} - 10^{-3} Torr. Therefore, extraction of tritium from molten Li_2BeF_4 salt with Li-Bi is not practical. The solubility of hydrogen in solid Li-Al (50 - 50 at.%) at 500 $^{\circ}\text{C}$ follows Sieverts' Law; the Sieverts' constant was measured to be $1.9 (+0.1) \times 10^4 \text{ Torr}^{1/2} / \text{atomic fraction}$. Tritium sorption in Li-Al ranged from 0.01 to 7 ppm at 400 - 600 $^{\circ}\text{C}$ at respective tritium partial pressures of 0.14 - 0.52 Torr.

EDB SUBJECT CATEGORY: EDB-360105;700209
INIS CATEGORY: B22

<615>

<615>

Tamaki, S. (Niigata Univ. (Japan). Dept. of Physics)

Application of the hard-sphere model to thermodynamic properties of liquid 3d transition metals AIX-07:251112;EDB-76:097500; J. Phys., F (London) (JPFMA) 6:(4):L89-L92 (Apr 1976)

SUBJECT DESCRIPTORS: ANALYTICAL SOLUTION;HARD-SPHERE
MODEL;LIQUID METALS;PERCUS-YEVICK EQUATION;SPECIFIC
HEAT;THERMODYNAMIC PROPERTIES: Q1;TRANSITION ELEMENTS:
T1;VANADIUM;VAPORIZATION HEAT

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-360104

INIS CATEGORY: B22

<616>

Tamura, S.; Yokokawa, T. (Hokkaido Univ., Sapporo (Japan).
Faculty of Science)

Enthalpy of vitreous beryllium fluoride
AIX-08:280698;EDB-77:094784; Bull. Chem. Soc. Jpn. (BCSJA)
48:(9):2542-2547 (Sep 1975)

SUBJECT DESCRIPTORS: BERYLLIUM FLUORIDES:
M1;CALORIMETRY;ENTHALPY;GLASS;HIGH TEMPERATURE;LIQUIDS;PHASE
TRANSFORMATIONS;SPECIFIC HEAT;TEMPERATURE
DEPENDENCE;THERMODYNAMIC PROPERTIES: Q1;VISCOSITY

ABSTRACT: The enthalpy increments of beryllium fluoride were measured by the drop method in the glassy and liquid states from 450 to 914 K. The other thermodynamic functions were also calculated. The estimated configurational entropy of the supercooled liquid showed its temperature dependence to be small compared with those of nearly all other glass-forming liquids. The theory of Adam and Gibbs was applied to explain both the present enthalpy data and the Arrhenius viscosity behaviour over an unusually wide temperature range. The similarity between beryllium fluoride and silicon dioxide was also shown with respect to the heat capacities of the glassy and supercooled liquid states.

EDB SUBJECT CATEGORY: EDB-400201

INIS CATEGORY: B12

<617>

Taylor, A.R. Jr.; Gardner, T.E., Bureau of Mines, Tuscaloosa, Ala. (USA). Tuscaloosa Metallurgy Research Lab.

In Some thermal properties of beryllium fluoride from 8\$sup 0\$ to 1,200\$sup 0\$ K ** EDB-77:101124; (Dec 1964) BM-RI--6664

SUBJECT DESCRIPTORS: BERYLLIUM FLUORIDES: T1;GLASS;HIGH TEMPERATURE;LOW TEMPERATURE;MEDIUM TEMPERATURE;SPECIFIC HEAT: Q1;ULTRALOW TEMPERATURE;VERY HIGH TEMPERATURE;VERY LOW TEMPERATURE

ABSTRACT: None

EDE SUBJECT CATEGORY: EDE-400201

<618>

Tetenbaum, M.

Total pressures of uranium- and plutonium-bearing species above the U--Pu--O system

ERA-02:002461;INS-76:020669;EDB-76:090326; Trans. Am. Nucl. Soc. (TANSA) 23::131-132 (Jun 1976)

SUBJECT DESCRIPTORS: NUCLEAR FUELS;PLUTONIUM DIOXIDE: T2;PRESSURE DEPENDENCE;SUBLIMATION;URANIUM DIOXIDE: T1;VAPOR PRESSURE: Q1,Q2;VERY HIGH TEMPERATURE

ABSTRACT: None

EDE SUBJECT CATEGORY: EDB-050700;360204
INIS CATEGORY: B23

<619>

Thayer, W.L.; Robbins, J.L. (California Univ., Livermore (USA). Lawrence Livermore Lab.) ** Blank, H.; Lindner, R. (eds.)

Enthalpy and heat capacity of liquid plutonium and uranium In Plutonium 1975 and other actinides **

AIX-(8:309932;ERA-02:041806;EDB-77:094171; (1976)
North-Holland

SUBJECT DESCRIPTORS: ENTHALPY: Q1,Q2;FILMS;FISSION;HIGH TEMPERATURE;LIQUID METALS;NEUTRON REACTIONS;PLUTONIUM: T1;RADIATION HEATING;RADICMETRIC ANALYSIS;SPECIFIC HEAT: Q1,Q2;SUPER KUKLA REACTOR;TEMPERATURE MEASUREMENT;URANIUM: T2

ABSTRACT: The enthalpy and effective heat capacity of delta-plutonium and or alloy have been measured at temperatures high in the liquid region. The Super Kukla prompt-burst reactor was used to heat thin wafers of these

<619>

<619> CONTI.

metals by neutron-induced fission. The temperature increase in the samples was measured with thermocouples and the number of fissions in the sample determined radiochemically. From these data, heat capacity and enthalpy were calculated. The heat capacity of oralloy was calculated as 159 J/kg.^{0°C} up to 2200^{0°C}. A constant value of 180 J/Kg.^{0°C} was computed for the effective heat capacity of liquid delta-plutonium up to 2200^{0°C}. Because molten plutonium and uranium would chemically attack even the most refractory crucible materials, it would be very difficult to obtain such data by conventional laboratory calorimetry; it is believed that the use of neutron heating in these investigations is unique.

EDB SUBJECT CATEGORY: EDB-360104
INIS CATEGORY: B22

<620>

Thayer, W.L.; Robbins, J.L. (California Univ., Livermore (USA). Lawrence Livermore Lab.) ** Commission of the European Communities, Karlsruhe (Germany, F.R.). European Inst. for Transuranium Elements

Enthalpy and heat capacity of liquid plutonium and uranium In 5th international conference on plutonium and other actinides. Book of abstracts **
AIX-07:233077;EDB-76:087038; (1975) AED-Conf-417-007

SUBJECT DESCRIPTORS: ENTHALPY: Q1,Q2;FISSION RATIO:LIQUID METALS;PLUTONIUM-DELTA: T1;SPECIFIC HEAT: Q1,Q2;URANIUM: T2;VERY HIGH TEMPERATURE

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-360104
INIS CATEGORY: B22

<621>

Thirriot, C.; Lacas, L. (Toulouse-3 Univ., 31 (France)); Amoros, D. (Centre National de la Recherche Scientifique, 31 - Toulouse (France))

Experimental study of the temperature dependence on energy exchanges in liquid metals In Comptes rendus du 95 **
AIX-07:279004;EDB-77:026115; (1975) Bibliotheque Nationale

SUBJECT DESCRIPTORS: ENERGY TRANSFER: Q1;LIQUID METALS: T1;TEMPERATURE DEPENDENCE;VISCOSITY

ABSTRACT: Because of their simple structure, liquid metals are

<621> CONT.

ideal for fundamental research on the liquid state. The very wide temperature region between the adjacent solid-vapour states is a great advantage in studying the influence of temperature on parameters characteristics of energy transfer. Following a bibliographical survey aimed at establishing the possibilities of theoretical prediction the results of experiments carried out at the Toulouse Institute of Fluid Mechanics, concerning measurements of the viscosity, specific heat and diffusivity parameters for several liquid metals and an alloy, are presented in detail. The experimental results are compared with theoretical predictions. By using the law of the corresponding states it is possible to give graphs relating dimensionless parameters.

EDB SUBJECT CATEGORY: EDB-360104

INIS CATEGORY: E11

<622>

Tippelskirch, H. v., Karlsruhe Univ. (TH) (Germany, F.R.). Inst. fuer Physikalische Chemie und Elektrochemie

In Viscosities of cesium vapor to 1,620 K and of liquid gallium to 1,800 K ** AIX-07:268015;EDB-77:036624; (1976)
AED-CONF--76-087-008

SUBJECT DESCRIPTORS: CESIUM: T2;GALLIUM: T1;LIQUID METALS;TEMPERATURE DEPENDENCE;VAPORS;VERY HIGH TEMPERATURE;VISCOSITY: Q1,Q2

ABSTRACT: The viscosity of cesium at 1,620 K and 40 bar has been determined to $41 \times 10^{\text{sup}} -6 \text{ (Pa} \times \text{s)}$ by the oscillating cup method. The saturated vapor density at 1,580 K could be derived from the viscosity measurements. The viscosity of liquid gallium has been determined from 370 K to 1,800 K. The experimental results have been compared with calculations based on the Enskog hard-sphere transport theory for dense fluids.

EDB SUBJECT CATEGORY: EDB-360103

INIS CATEGORY: B22

<623>

Tiptsova-Yakovleva, V.G.; Kopnina, O.I.; Lopatina, E.A. (Moskovskij Inst. Stali i Splavov (USSR))

Spectrophotometric determination of rare earths and yttrium in gallium base alloys AIX-08:342659;EDB-78:017233; Izv. Vyssh. Uchebn. Zaved., Khim. Khim. Tekhnol. (IVUKA) 20:(3):465 (1977)

SUBJECT DESCRIPTORS: CHEMICAL ANALYSIS: Q1;GALLIUM BASE ALLOYS:

<623>

<623> CONT.

T1;QUANTITATIVE CHEMICAL ANALYSIS;RARE EARTHS:

T2;SPECTROPHOTOMETRY: Q2,Q3;YTTRIUM: T3

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-400104

INIS CATEGORY: B11

<624>

Tishura, T.A.; Budarina, A.N. (AN Ukrainskoj SSR, Kiev. Inst. Obshchey i Neorganicheskoy Khimii)

Degree of thermal dissociation of $M\text{CaCl}_3$ (M-K,Rb,Cs) complexes during melting AIX-08:286199;EDB-77:094806; Ukr. Khim. Zh. (UKZHA) 42:(3):280-282 (Mar 1976)

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-400201

INIS CATEGORY: B12

<625>

Toft Soerensen, O., Danish Atomic Energy Commission, Risoe. Research Establishment; Lund Univ. (Sweden)

In Rare earth and actinide oxides. Thermodynamic and electron microscopy studies ** AIX-07:272766;EDB-77:055674; (Oct 1975) RISO--331

ABSTRACT: Partial molar thermodynamic quantities for oxygen in substoichiometric cerium oxides (CeO_{2-x}), plutonium oxides (PuO_{2-x}) and mixed uranium--plutonium oxides ($(\text{U}, \text{Pu})\text{O}_{2-x}$) were determined by thermogravimetric analysis in atmospheres of controlled oxygen pressures (CO_2/CO mixtures) in the temperature range 900 to 1450 deg C. Detailed analysis of the data obtained showed that the nonstoichiometric phase ranges for the three oxide-systems can be divided into several subregions each consisting of an apparent nonstoichiometric single phase. The finer details of the thermodynamic data, however, suggest that some of these subregions can be further split into ordered intermediate phases with compositions following the series $M_n\text{O}_{2n-2}$. Supplementary high-temperature X-ray diffraction studies were made on CeO_{2-x} at temperatures up to 855 deg C. Between 790 and 855 deg C, a new phase of low symmetry was obtained. Indexing the powder pattern for this phase showed it to be isostructural with Pr_6O_{11} and with a monoclinic unit cell with $a=6.781 \pm 0.006$ A, $b=11.893 \pm 0.009$ A, $c=15.823 \pm 0.015$ A and $\beta=125.04 \pm 0.04$ deg. The Ce_6O_{11} phase observed in the X-ray studies

<625> CONT.

corresponds to one of the intermediate phases inferred from the thermodynamic data. Supplementary high-resolution electron microscopy studies were also conducted on reduced single crystals of CeO_2 . On some particles reduced by beam heating, a lamellae structure was observed and a model involving crystallographic shearing is proposed to explain this observation. In other beam-heated particles diffraction patterns were observed corresponding to the monoclinic superstructure found in the high-temperature X-ray studies. Finally patterns on particles reduced by a heat treatment in vacuum under well defined conditions showed that twinning can also take place in this oxide system.

EDB SUBJECT CATEGORY: EDB-360202;360204
INIS CATEGORY: B23

<626>

Toft Scerensen, O. (Danish Atomic Energy Commission, Risoe. Research Establishment)

Studies of non-stoichiometric oxides by thermoanalytical methods
AIX-07:247911;EDB-76:081891; Thermochim. Acta (THACA)
15:(2):227-237 (May 1976)

SUBJECT DESCRIPTORS: CERIUM OXIDES: T1;CHEMICAL COMPOSITION;CONTROLLED ATMOSPHERES;CRYSTAL LATTICES;HIGH TEMPERATURE;OXYGEN;PHASE STUDIES;PLUTONIUM OXIDES: T2;STOICHIOMETRY: Q1,Q2,Q3;TEMPERATURE DEPENDENCE;THERMAL GRAVIMETRIC ANALYSIS;THERMODYNAMIC PROPERTIES;URANIUM OXIDES: T3;VACANCIES;VAPOR PRESSURE;VERY HIGH TEMPERATURE

ABSTRACT: Non-stoichiometric oxides can be divided into metal-deficient and oxygen-deficient oxides, and a few examples of each type are given in order to show the materials of interest in this field. After a short review of the methods used in the study of these oxides, an account is given of the methods used in and the results obtained from a recent thermogravimetric investigation in atmospheres of controlled oxygen pressure on cerium oxides, plutonium oxides and mixed uranium/plutonium oxides, which are all of special interest in research on fuels for nuclear power reactors.

EDB SUBJECT CATEGORY: EDB-400702
INIS CATEGORY: B11

<627>

<627>

Tominaga, T.; Sakai, T.; Mimura, T. (Tokyo Univ. (Japan).
Faculty of Science)

A Moessbauer spectroscopic study of intercalation compounds of
graphite with ferric chloride and aluminum chloride
EDF-76:036719; Chem. Lett. (Tokyo) (CMLTA) (8):853-858 (Aug
1974)

SUBJECT DESCRIPTORS: ALUMINIUM CHLORIDES:
T1;CLATHRATES;GRAPHITE: T2;IRON CHLORIDES: T3;ISOMER
SHIFT;MOESSBAUER EFFECT;QUADRUPOLES;SPIN;STRUCTURAL CHEMICAL
ANALYSIS: Q1,Q2,Q3

ABSTRACT: The Mossbauer spectra of graphite- FeCl_3 and
graphite- FeCl_3 - AlCl_3 compounds with varying
 $\text{FeCl}_3/\text{AlCl}_3$ ratio were measured at 80°K and
298°K. In the spectra of the latter compounds with
 $\text{FeCl}_3/\text{AlCl}_3$ ratio of approximately one and less
were observed two quadrupole doublets indicating the presence
of high-spin Fe(II). (auth)

EDF SUBJECT CATEGORY: EDB-360603
INIS CATEGORY: A13

<628>

Toropov, A.N.; Sokhovich, E.V.; Rubinshtein, O.V.; Ermakova,
T.M. (Leningradskij Tekhnologicheskij Inst. (USSR))

Refractory materials resistance to molten barium alloy
AIX-08:340277;EDB-78:016989; Ogneupory (OGNPA) (7):64-68 (Jul
1975)

SUBJECT DESCRIPTORS: ARGON;BARIUM ALLOYS: T1;BARIUM
COMPOUNDS;CALCIUM COMPOUNDS: T2;CHEMICAL REACTIONS;CORROSION:
Q2,Q3,Q4,Q5;CORROSIVE EFFECTS: Q1;CORUNDUM: T3;LIQUID
METALS;MATERIALS TESTING;MICROHARDNESS;MICROSTRUCTURE;PHASE
STUDIES;SPINELS: T4;TEMPERATURE DEPENDENCE;ZIRCONATES: T5

ABSTRACT: An investigation was made of the resistance to barium
fusion of dense sintered refractors materials - corundum,
magnesium clay spinel, magnesium oxide, and calcium and
barium zirconates. The tests were made by the crucible method
at 1,600°C in an argon atmosphere. The refractory
materials resist well to barium fusion. The presence of
barium oxide reduces the resistance of corundum and spinel as
a result of the formation of barium monoaluminate in
equilibrium with the eutectic that fuses at 1,480°C.
Calcium zirconate in the presence of BaO becomes barium
zirconate, which causes cracking of the samples. The most
promising materials resistant to the fusion of barium and its

<628> CONT.

oxide are the solid refractories of pure magnesium oxide and barium zirconate.

EDB SUBJECT CATEGORY: EDB-360205

INIS CATEGORY: B23

<629>

Tcvmas'yan, I.K.; Krasnova, O.M.; Bukhalova, G.A.
(Rostovskij-na-Donu Pedagogicheskij Inst. (USSR))

Quarter mutual system of chlorides, bromides, iodides of sodium and rubidium EDB-76:036671; Zh. Neorg. Khim. (ZNOKA) 20:(1):254-256 (Jan 1975)

SUBJECT DESCRIPTORS: CHEMICAL REACTIONS; MELTING POINTS; MIXTURES; PHASE DIAGRAMS: Q1, Q2, Q3, Q4, Q5, Q6; SOLID SOLUTIONS; THERMAL ANALYSIS

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-360602

INIS CATEGORY: B12

<630>

Tovmas'yan, I.K.; Krasnova, O.M. (Rostovskij-na-Donu Pedagogicheskij Inst. (USSR))

Quaternary mutual system of chlorides, bromides, iodides of potassium and cesium EDB-76:048911; Zh. Neorg. Khim. (ZNOKA) 20:(4):1103-1105 (Apr 1975)

SUBJECT DESCRIPTORS: CESIUM BROMIDES: T2; CESIUM CHLORIDES: T1; CESIUM IODIDES: T3; CRYSTAL STRUCTURE; DIFFERENTIAL THERMAL ANALYSIS; PHASE DIAGRAMS: Q1, Q2, Q3, Q4, Q5, Q6; POTASSIUM BROMIDES: T5; POTASSIUM CHLORIDES: T4; POTASSIUM IODIDES: T6

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-360602

INIS CATEGORY: B12

<631>

<631>

Tovmas'yan, I.K.; Tsushba, T.M.

Sections of quaternary mutual system consisting of sodium, potassium and cesium chlorides and bromides EDB-76:036681; Zh. Neorg. Khim. (ZNOKA) 20:(2):513-515 (Feb 1975)

SUBJECT DESCRIPTORS: ALKALI METAL COMPOUNDS: T1; CESIUM BROMIDES; CESIUM CHLORIDES; CROSS SECTIONS; CRYSTALLIZATION; HALIDES: T2; PHASE DIAGRAMS: Q1, Q2; POTASSIUM BROMIDES; POTASSIUM CHLORIDES; SODIUM BROMIDES; SODIUM CHLORIDES; SOLID SOLUTIONS

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-360602

INIS CATEGORY: B12

<632>

Trebeljahr, S. (Zentralinstitut fuer Kernforschung, Rossendorf bei Dresden (German Democratic Republic)) ** Muenze, R. (ed.), Zentralinstitut fuer Kernforschung, Rossendorf bei Dresden (German Democratic Republic)

Spectrophotometric determination of uranium in spent fuels of WWER-2 reactors In Annual report, 1975 ** AIX-07:272349; EDB-77:054177; (Aug 1975) ZfK--294

SUBJECT DESCRIPTORS: BURNUP; QUANTITATIVE CHEMICAL ANALYSIS: Q2; SPECTROPHOTOMETRY: T; SPENT FUEL ELEMENTS: T2, Q1; SPENT FUELS; URANIUM; URANIUM COMPLEXES; WWER-2 REACTOR: T1

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-210200; 440105; 400104

INIS CATEGORY: B11

<633>

Tret'yakova, K.V.; Tyurin, V.I.; Nisel'son, L.A.

Systems of NbCl_5 - GaCl_3 and TaCl_5 - GaCl_3 AIX-07:277929; EDB-77:031094; Zh. Neorg. Khim. (ZNOKA) 21:(5):1310-1314 (May 1976)

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-360602; 400201

INIS CATEGORY: B12

<634>

Tretyakov, Yu.D.; Kaul, A.R.; Makukhin, N.V.

Electrochemical study of high-temperature stability of compounds between the rare earths and copper oxide

INS-76:013209;EDB-76:042345; J. Solid State Chem. (JSSCB) 17:(1-2):183-189 (Feb 1976)

SUBJECT DESCRIPTORS: COPPER OXIDES: T1;CRYSTAL STRUCTURE: Q1,Q2,Q3;ELECTROCHEMISTRY;ELECTROLYTES;FORMATION FREE ENERGY;OXIDES: T3;RARE EARTH COMPOUNDS: T2;THERMODYNAMIC PROPERTIES;VERY HIGH TEMPERATURE;X-RAY DIFFRACTION

ABSTRACT: Equilibrium conditions for formation of the compounds $\text{CuLn}_{20}\text{O}_{40}$ ($\text{Ln} = \text{La}, \text{Nd}, \text{Sm}, \text{Eu}, \text{Gd}$) and $\text{CuR}_{20}\text{O}_{50}$ ($\text{R} = \text{Tb}, \text{Dy}, \text{Er}, \text{Yb}, \text{Y}, \text{In}$) were studied in galvanic cells with solid electrolyte $\text{ZrO}_{20}(\text{Y}_{20}\text{O}_{30})$ in the temperature range 950 to 1150°C. The results, together with results of a study of the equilibrium $\text{CuO}-\text{Cu}_{20}\text{O}$, were used to calculate the ΔG^0 of formation of the double oxides from CuO and the $\text{Ln}_{20}\text{O}_{30}$ ($\text{In}_{20}\text{O}_{30}$; $\text{Y}_{20}\text{O}_{30}$) as listed above. A decrease was found in the stability of the compounds $\text{CuLn}_{20}\text{O}_{40}$ relative to the initial oxides in the sequence $\text{La}-\text{Gd}$ and an increase in the stability of $\text{Cu}_{20}\text{Ln}_{20}\text{O}_{50}$ in the sequence $\text{Tb}-\text{Yb}$. The results are discussed on the basis of special features of the crystalline structure of the compounds examined.

EDB SUBJECT CATEGORY: EDB-360202
INIS CATEGORY: B23

<635>

Trojanov, S.I.; Marek, G.S.; Tsirel'nikov, V.I. (Moskovskij Gosudarstvennyj Univ. (USSR))

Thermographic determination of formation heats of lower zirconium chlorides and bromides EDB-76:036718; Izv. Sib. Otd. Akad. Nauk SSSR, Ser. Khim. Nauk (IZSKA) (9):111-114 (Jul 1974)

SUBJECT DESCRIPTORS: CHEMICAL PREPARATION;DIFFERENTIAL THERMAL ANALYSIS;FORMATION HEAT: Q1,Q2;ZIRCONIUM BROMIDES: T1;ZIRCONIUM CHLORIDES: T2

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-360603
INIS CATEGORY: B12

<636>

<636>

Trunin, A.S.; Bukhalova, G.A.; Petrova, D.G.; Garkushin, I.K.
(Kujbyshevskij Politekhnikheskij Inst. (USSR))

Thermal analysis of NaF, NaCl, NaMoO₄ system
AIX-08:298365;EDB-77:069128; Zh. Neorg. Khim. (ZNOKA)
21:(9):2506-2510 (Sep 1976)

SUBJECT DESCRIPTORS: CHEMICAL
COMPOSITION;CRYSTALLIZATION;DIFFERENTIAL THERMAL ANALYSIS:
Q1,Q2,Q3;EUTECTICS;MELTING POINTS;MIXTURES;MOLTEN
SALTS;MOLYBDATES: T1;PHASE DIAGRAMS;SODIUM CHLORIDES:
T2;SODIUM FLUORIDES: T3

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-400201
INIS CATEGORY: B12

<637>

Trunin, A.S.; Garkushin, I.K.; Vasil'chenko, L.M.

System NaF₂WO₄-NaCl-CaWO₄-CaCl₂
AIX-08:337037;EDB-77:140954; Zh. Neorg. Khim. (ZNOKA)
22:(2):495-498 (Feb 1977)

SUBJECT DESCRIPTORS: CALCIUM CHLORIDES: T1;CALCIUM TUNGSTATES:
T2;CRYSTALLIZATION: Q1,Q2,Q3,Q4;EUTECTICS;MELTING
POINTS;PHASE DIAGRAMS;SODIUM CHLORIDES: T3;SODIUM TUNGSTATES:
T4

ABSTRACT: Using the methods of thermal analysis (DTA, VPM), the authors have investigated the liquidus of the triple mutual system Na, Ca long Cl, WO₄. The crystallization fields of the components and the compounds of congruent melting of composition NaClxNa₂WO₄ have been separated. The system Na, Ca long Cl, WO₄ is singular and is characterized by the presence of three triple eutectics: E₁=494°C (degenerated eutectic), E₂=656°C, E₃=627°C.

EDB SUBJECT CATEGORY: EDB-400201
INIS CATEGORY: B12

<638>

Trunin, A.S.; Shter, G.E.; Kosmynin, A.S. (Kujbyshevskij
Sel'skohozyajstvennyj Inst. (USSR); Kujbyshevskij
Meditinskij Inst. (USSR))

NaMoO_4 -- BaMoO_4 -- NaF -- BaF system EDB-76:042600; Zh.
Neorg. Khim. (ZNOKA) 20:(6):1647-1651 (Jun 1975)

SUBJECT DESCRIPTORS: BARIUM COMPOUNDS: T5; BARIUM FLUORIDES:
T1; CHEMICAL COMPOSITION; CHEMICAL
REACTIONS; CRYSTALLIZATION; DIFFERENTIAL THERMAL
ANALYSIS; EUTECTICS; MELTING; MOLYBDATE: T2; PHASE DIAGRAMS:
Q1, Q2, Q3, Q4, Q5; SODIUM COMPOUNDS: T4; SODIUM FLUORIDES:
T3; SOLID SOLUTIONS; X-RAY DIFFRACTION

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-400201

INIS CATEGORY: B12

<639>

Trunov, V.K. (Moskovskii Gosudarstvennyi Univ. (USSR))

Double potassium rare earth tungstates
AIX-07:265110; EDB-77:013490; Zh. Neorg. Khim. (ZNOKA)
20:(7):1974-1978 (Jul 1975)

SUBJECT DESCRIPTORS: COMPARATIVE EVALUATIONS; DIFFERENTIAL
THERMAL ANALYSIS; ERBIUM COMPOUNDS: T5; PHASE DIAGRAMS:
Q1, Q2, Q3, Q4, Q5; POTASSIUM TUNGSTATES: T1; PROMETHIUM COMPOUNDS:
T3; RARE EARTH COMPOUNDS; TERBIUM COMPOUNDS: T4; TUNGSTATES:
T2; X-RAY DIFFRACTION

ABSTRACT: By roentgenophase and differential-thermal analysis
phase diagrams of the potassium tungstates-tungstates r.e.e.
(for Pm, Tb, Er) system was studied. On the basis of
published and this study data the question of changing
properties of double potassium tungstate in the rare earth
elements series is discussed.

EDB SUBJECT CATEGORY: EDB-400201

INIS CATEGORY: B12

<640>

<640>

Tsushba, T.M.; Tovmas'yan, I.B.; Krasnova, O.M.
(Rostovskij-na-Donu Pedagogicheskij Inst. (USSR))

Quaternary mutual system of chlorides, bromides, iodides of
rubidium and cesium EDB-76:048912; Zh. Neorg. Khim. (ZNCKA)
20:(5):1376-1377 (May 1975)

SUBJECT DESCRIPTORS: CESIUM BROMIDES: T1;CESIUM CHLORIDES:
T2;CESIUM IODIDES: T3;DIFFERENTIAL THERMAL ANALYSIS;PHASE
DIAGRAMS: Q1,Q2,Q3,Q4,Q5,Q6;RUBIDIUM BROMIDES: T6;RUBIDIUM
CHLORIDES: T4;RUBIDIUM IODIDES: T5;SOLID SOLUTIONS;X-RAY
DIFFRACTION

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-360602
INIS CATEGORY: B12

<641>

Tsvigunov, A.; Polunina, G.P.; Kovba, L.M.; Impolitova, I.A. **
AN SSSR, Sverdlovsk. Inst. Khimii; Vostochnyj
Nauchno-Issledovatel'skij i Proektnyj Inst. Ogneupornoj
Promyshlennosti, Sverdlovsk (USSR); Ural'skij
Nauchno-Issledovatel'skij Khimicheskij Inst., Sverdlovsk
(USSR)

Phases of high pressure in UO_2 - UO_3 - PbO system In
Summaries of reports of the all-union meeting on solid body
chemistry. 3--5 June 1975. III. Synthesis, phase
equilibrium and use of solid oxide compounds **
AIX-08:289418;EDB-77:074781; (1975) INIS-mf--3246

SUBJECT DESCRIPTORS: ALLOY SYSTEMS;CRYSTAL STRUCTURE;HIGH
PRESSURE;HIGH TEMPERATURE;LATTICE PARAMETERS;LEAD OXIDES:
T1;PHASE STUDIES: Q1,Q2,Q3;PHASE TRANSFORMATIONS;URANIUM
DIOXIDE: T3;URANIUM TRIOXIDE: T2

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-360202
INIS CATEGORY: B13

<642>

Twidwell, L.G.

Thermodynamic properties of sulphur in liquid antimony
 EDB-77:107253; Inst. Min. Metall., Trans., Sect. C (TMEMA)
 85::C52-C54 (Mar 1976)

SUBJECT DESCRIPTORS: ANTIMONY: T3; HIGH TEMPERATURE; LIQUID
 METALS; PHASE STUDIES; SOLVENT PROPERTIES: Q3; SULFUR:
 T1; THERMODYNAMIC PROPERTIES: Q1

ABSTRACT: The thermodynamic activity of sulphur in liquid
 antimony was determined over the temperature range
 640--750^oC. This study is a part of a series of
 investigations to determine the thermodynamic properties of
 sulphur in liquid metallic solutions.

EDB SUBJECT CATEGORY: EDB-360603

<643>

Tyagunov, G.V.; Baum, B.A.; Kushnir, M.N.; Khasin, G.A.;
 Ermanovich, N.A.

Influence of the technology of the alloy fabrication on its
 properties in a liquid state EDB-76:036326; Fiz. Khim. Obrab.
 Mater. (FKOMA) (1):48-50 (Jan 1975)

SUBJECT DESCRIPTORS: CORRELATIONS; CRYSTALLIZATION; ELECTRIC
 CONDUCTIVITY; ELONGATION; HYSTERESIS; LIQUID METALS; MECHANICAL
 PROPERTIES: Q1; MELTING POINTS; PHYSICAL PROPERTIES:
 Q1; PLASTICITY; SMELTING; STEELS: T1; TEMPERATURE
 DEPENDENCE; THERMAL CONDUCTIVITY; VACUUM
 MELTING; VISCOSITY; YIELD STRENGTH

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-360103; 360104

INIS CATEGORY: B21

<644>

Ugajin, M.

Thermodynamic activity of carbon in molybdenum-containing
 uranium carbide EDB-77:005578; J. Nucl. Sci. Technol. (Tokyo)
 (JNSTA) 12:(6):381-384 (Jun 1975)

SUBJECT DESCRIPTORS: CARBON; COMPATIBILITY: Q1, Q2, Q3, Q4; FAST
 REACTORS; FREE ENTHALPY; FUEL-CLADDING INTERACTIONS; MOLYBDENUM:
 T3; PHASE DIAGRAMS; PHASE STUDIES; PLUTONIUM CARBIDES:
 T2; STAINLESS STEELS: T4; THERMODYNAMIC PROPERTIES; URANIUM
 CARBIDES: T1

<644>

<644> CONT.

ABSTRACT: The carbides UC and (U, Pu)C are promising fuel materials for fast reactors. The problem involved in their utilization is that the stainless steel used in the cladding tends to pick up carbon from the fuel, and this carburization of the steel may eventually undermine the intrinsic utility of these compounds. Calculations predict that molybdenum-stabilized carbide fuel is thermodynamically compatible with austenitic stainless steel cladding. 8 refs.

EDB SUBJECT CATEGORY: EDB-050700;360105

<645>

Ugajin, M., Japan Atomic Energy Research Inst., Tokyo

In Phase equilibrium study on system

uranium-plutonium-tungsten-carbon **

AIX-08:315114;EDB-77:113695; (Nov 1976) JAERI-M--6804

SUBJECT DESCRIPTORS: FORMATION FREE ENERGY;METALLURGICAL EFFECTS: Q3;PHASE DIAGRAMS;PHASE STUDIES;PHASE TRANSFORMATIONS: Q1,Q2;PLUTONIUM CARBIDES: T2;QUATERNARY ALLOY SYSTEMS;SEGREGATION;SOLUTION HEAT;TERNARY ALLOY SYSTEMS;THERMAL EQUILIBRIUM;TUNGSTEN: T3;TUNGSTEN ADDITIONS;URANIUM CARBIDES: T1

ABSTRACT: Metallurgical properties of the U-Pu-W-C system have been studied with emphasis on phases and reactions. Free energy of compound formation, carbon activity and U/Pu segregation in the W-doped carbide fuel are estimated using phase diagram data. The results indicate that tungsten metal is useful as a thermochemical stabilizer of the carbide fuel. Tungsten has high temperature stability in contact with uranium carbide and mixed uranium-plutonium carbide.

EDB SUBJECT CATEGORY: EDB-360202;050700

INIS CATEGORY: B23

<646>

Ugajin, M.; Ate, J.; Kurihara, M.

Phase behavior and thermodynamics of U--Mo--C system

EDB-77:013203; J. Nucl. Sci. Technol. (Tokyo) (JNSTA) 12:(9):560-566 (Sep 1975)

SUBJECT DESCRIPTORS: CARBIDES: M5;CARBON;CARBON COMPOUNDS: M3;CHEMICAL REACTION KINETICS;CHEMICAL REACTIONS;CRYSTAL STRUCTURE;EQUATIONS;EQUILIBRIUM;FORMATION FREE ENERGY: Q4,Q5;MOLYBDENUM;MOLYBDENUM ALLOYS: M1;PHASE DIAGRAMS;PHASE STUDIES: Q1,Q2,Q3;PHASE TRANSFORMATIONS;THERMODYNAMIC PROPERTIES: Q4,Q5;THERMODYNAMICS;URANIUM;URANIUM ALLOYS:

<646> CONT.

M2;URANIUM CARBIDES: M4

ABSTRACT: The U-Mo-C system has been studied to supplement the hitherto available phase-diagram data and to deduce therefrom the thermodynamic property of $\text{UMoC}_{\text{sub } 1\% \text{ sub } 7\%}$ and $\text{UMoC}_{\text{sub } 2\%}$. It is shown that a peritectic four-phase in equilibrium reaction takes place: $\text{UC} + \text{Mo} \rightarrow \text{UMoC}_{\text{sub } 1\% \text{ sub } 7\%} + \text{liquid}$, at $1,830 \pm 30^\circ\text{C}$. The phase relations between orthorhombic $\text{UMoC}_{\text{sub } 2\%}$ and monoclinic $\text{UMoC}_{\text{sub } 1\% \text{ sub } 7\%}$ may be defined by the in equilibrium reaction $\text{UMoC}_{\text{sub } 1\% \text{ sub } 7\%} + \text{UMoC}_{\text{sub } 2\%} + \text{liquid}$, which will occur at some temperature between $2,200^\circ\text{C}$ and $2,350^\circ\text{C}$, the reported melting point of $\text{UMoC}_{\text{sub } 2\%}$. Liquidus projection for the U-Mo-C system is presented on the basis of these results and from data found in literature. 26 refs.

EDB SUBJECT CATEGORY: EDB-360202;050700

<647>

Ugajin, M.; Abe, J.; Kurihara, M. (Japan Atomic Energy Research Inst., Tokai, Ibaraki. Tokai Research Establishment)

Phase behavior and thermodynamics of U--Mo--C system

AIX-07:262737;EDB-77:018944; J. Nucl. Sci. Technol. (Tokyo) (JNSTA) 12:(9):560-566 (Sep 1975)

SUBJECT DESCRIPTORS: ENTROPY;FORMATION FREE ENERGY;MOLYBDENUM CARBIDES: T1;ORTHORHOMBIC LATTICES;PHASE DIAGRAMS;PHASE STUDIES: Q1,Q2;THERMODYNAMICS;URANIUM CARBIDES: T2;VERY HIGH TEMPERATURE

ABSTRACT: The U-Mo-C system has been studied to supplement the hitherto available phase-diagram data and to deduce therefrom the thermodynamic property of $\text{UMoC}_{\text{sub } (1.7)}$ and $\text{UMoC}_{\text{sub } 2\%}$. It is shown that a peritectic four-phase reaction takes place: $\text{UC} + \text{Mo} \rightarrow \text{UMoC}_{\text{sub } (1.7)} + \text{liquid}$, at $1,830 \pm 30^\circ\text{C}$. The phase relations between orthorhombic $\text{UMoC}_{\text{sub } 2\%}$ and monoclinic $\text{UMoC}_{\text{sub } (1.7)}$ may be defined by $\text{UMoC}_{\text{sub } (1.7)} + \text{UMoC}_{\text{sub } 2\%} + \text{liquid}$, which will occur at some temperature between $2,200^\circ\text{C}$ and $2,350^\circ\text{C}$, the reported melting point of $\text{UMoC}_{\text{sub } 2\%}$. Liquidus projection for the U-Mo-C system is presented on the basis of these results and from data found in literature. The free energy equations for the formation of $\text{UMoC}_{\text{sub } (1.7)}$ and $\text{UMoC}_{\text{sub } 2\%}$ are respectively estimated to be $\Delta G_{\text{sub } (f)}^\circ(\text{UMoC}_{\text{sub } (1.7)}) = -40,350 - 0.69T \text{ cal/mol}$ and

<647>

<648> CONT.

$\Delta G_{\text{sub}}(f)_{\text{sup } 0}(\text{UMoC}_{\text{sub } 2}) = -46,360 - 0.69T \text{ cal/mol,}$
 both in the temperature range 298--2,100 K.

EDB SUBJECT CATEGORY: EDB-360202;360204

INIS CATEGORY: B23

<648>

Ugajin, M.; Abe, J.; Suzuki, Y.; Takahashi, I.; Kurihara, M.
 (Japan Atomic Energy Research Inst., Tokai, Ibaraki. Tokai
 Research Establishment)

Phase equilibria in system U--Pu--W--C

AIX-08:289684; ERA-02:025072; EDB-77:055683; J. Nucl. Sci.
 Technol. (Tokyo) (JNSTA) 13:(1):36-39 (Jan 1976)

SUBJECT DESCRIPTORS: COMPATIBILITY; EQUILIBRIUM; PHASE DIAGRAMS:
 Q1, Q2, Q3; PHASE STUDIES; PLUTONIUM CARBIDES: T1; QUATERNARY
 ALLOY SYSTEMS; TUNGSTEN CARBIDES: T2; URANIUM CARBIDES: T3; VERY
 HIGH TEMPERATURE

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-360202;050700

INIS CATEGORY: B23

<649>

Ugajin, M.; Takahashi, I.

In Process for preparing uranium carbonitride **

AIX-07:259850; ERA-02:012466; EDB-77:016966; (5 Sep 1969)

SUBJECT DESCRIPTORS:

CONTAINERS; EMBRITTLEMENT; GRAPHITE; MOLYBDENUM; NITROGEN; NUCLEAR
 FUELS; PHASE DIAGRAMS; PRODUCTION: Q1, Q2; SOLID
 SOLUTIONS; TUNGSTEN; URANIUM; URANIUM CARBIDES: M1; URANIUM
 COMPOUNDS; URANIUM NITRIDES: M2

ABSTRACT: In the conventional production of uranium carbonitride
 $\text{UC}_{\text{sub}(x)}\text{N}_{\text{sub}(y)}$ ($x+y=1$, x not equal 0, y not equal 0) as a
 reactor fuel, the partial pressure of $\text{N}_{\text{sub } 2}$ must be
 strictly controlled in order to prevent the formation of
 $\text{UC}_{\text{sub } 2}$ which carburizes the stainless steel-made container
 to make it fragile. Herein provided is a new process for
 preparing $\text{UC}_{\text{sub}(x)}\text{N}_{\text{sub}(y)}$ in which the partial pressure of
 $\text{N}_{\text{sub } 2}$ may be arbitrary. The process comprises fusing
 uranium carbide or a mixture of metallic uranium and graphite
 (the atomic ratio U:C=1:0.76 to 0.94) in the presence of
 metallic W or Mo and $\text{N}_{\text{sub } 2}$ gas (0.02 to 1.2 atm). The W or
 Mo reacts with the formed $\text{UC}_{\text{sub } 2}$ to convert it to harmless
 $\text{UWC}_{\text{sub } 2}$ or $\text{UMoC}_{\text{sub } 2}$. The minimum quantity of W or Mo to

<649> CONT.

be added can be predicted by referring to a U-C-W (or Mo)-N quaternary phase diagram (given in the specification), so far as the quantities of U and C and the partial pressure of N_2 are known. In one example, metallic uranium (1 mole), graphite (0.89 mole) and metallic W (0.07) were fused in an atmosphere of N_2 +Ar (total pressure 0.7 atm, N_2 0.45 atm) in a vacuum arc furnace (15 to 20 V, 90 to 250A) for about 2 minutes. This was repeated 7 times to yield 93.4% by mole of $UC_{0.82}N_{0.18}$ together with 6.6% by mole of $UWC_{0.82}N_{0.18}$.

EDB SUBJECT CATEGORY: EDB-050700;360201
INIS CATEGORY: B23

<650>

Ugajin, Mitsuhiro (Japan Atomic Energy Research Inst., Tokai, Ibaraki. Tckai Research Establishment)

Thermodynamic activity of carbon in molybdenum-containing uranium carbide AIX-07:267905;EDB-77:007636; J. Nucl. Sci. Technol. (Tokyo) (JNSTA) 12:(6):381-384 (Jun 1975)

SUBJECT DESCRIPTORS: CARBON: T1;CHEMICAL REACTION KINETICS: Q1,Q2;COMPATIBILITY;CRYSTAL-PHASE TRANSFORMATIONS;FREE ENERGY;FUEL-CLADDING INTERACTIONS;MOLYBDENUM CARBIDES;STAINLESS STEELS;TERNARY ALLOY SYSTEMS;URANIUM CARBIDES: T2

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-400702
INIS CATEGORY: B16

<651>

Ukanwa, A.O.

In Diffusion in liquid metal systems. Final report **
ERA-02:010434;INS-77:001229;EDB-77:013073; (Jun 1975)
N--76-10302

SUBJECT DESCRIPTORS: DATA COMPILATION;DENSITY;DIFFUSION: Q2,Q3;ELECTRIC CONDUCTIVITY;GALLIUM ALLOYS: A3;HEAT TRANSFER;INDIUM ALLCYS: A2;LIQUID METALS: A1;PHYSICAL PROPERTIES;REVIEWS: Q1;SAFETY;SPECIFIC HEAT;TABLES;THERMAL CONDUCTIVITY

ABSTRACT: Physical properties of twenty liquid metals are reported; some of the data on such liquid metal properties as density, electrical resistivity, thermal conductivity, and heat capacity are summarized in graphical form. Data on

<651>

<651> CONT.

laboratory handling and safety procedure are summarized for each metal; heat-transfer-correlations for liquid metals under various conditions of laminar and turbulent flow are included. Where sufficient data were available, temperature equations of properties were obtained by the method of least-squares fit. All values of properties given are valid in the given liquid phase ranges only. Additional tabular data on some 40 metals are reported in the appendix. Included is a brief description of experiments that were performed to investigate diffusion in liquid indium-gallium systems.
(Author) (GRA)

EDE SUBJECT CATEGORY: EDB-360104

INIS CATEGORY: B22

<652>

Ullmann, H.; Lang, H.J.; Nebelung, C. (Zentralinstitut fuer Kernforschung, Rossendorf bei Dresden (German Democratic Republic)) ** Muenze, R. (ed.)

Investigations on chemical properties of thorium oxide solid electrolyte ceramics In Annual report, 1976 **
AIX-08:324308;EDB-78:009583; (Jul 1976) ZfK--312

SUBJECT DESCRIPTORS: CERAMICS;CHEMICAL PROPERTIES:
Q1;ELECTROCHEMICAL CORROSION;ELECTROLYTES;ELECTROLYTIC
CELLS;ELECTROMOTIVE FORCE;IMPURITIES;SODIUM;THORIUM OXIDES: T1

ABSTRACT: None

EDE SUBJECT CATEGORY: EDB-360204;400201

INIS CATEGORY: B23

<653>

Uzuki, K., Tohoku Univ., Sendai (Japan). Research Inst. for Iron, Steel and Other Metals

In Structural study of liquids with strong short-range correlation in the atomic distribution **
AIX-07:264041;EDB-77:040110; (1976) AED-Conf--76-097-011

SUBJECT DESCRIPTORS: AMORPHOUS STATE: T1;ATOMIC MODELS;BOND LENGTHS;COMPARATIVE EVALUATIONS;CORRELATIONS;CRYSTAL STRUCTURE: Q1,Q2;DIAGRAMS;DISTRIBUTION;INTERACTION RANGE;LIQUID METALS;LIQUIDS: T2;MOLECULAR STRUCTURE;NEUTRON DIFFRACTION;PALLADIUM ALLOYS;QUANTITY RATIO;SELENIUM;SILICON ALLOYS;TIME-OF-FLIGHT METHOD;X-RAY DIFFRACTION

ABSTRACT: Structure factors of liquids and amorphous solids having a relatively high degree of ordering in their

<653> CONT.

short-range structures have been measured over a wide range of scattering vectors by means of the T-O-F neutron diffraction using epithermal pulsed neutrons generated by an electron linear accelerator. It has been shown in the case of liquid CS₂ that the size and shape of a molecule existing in the liquid phase are determined from the behaviour of the structure factor in the range of high scattering vectors, and that the structure factor in the region of low scattering vectors informs on inter-molecular orientational and center-center correlations in the liquid state. Moreover, based on highly resolved radial distribution functions, a free rotating chain model has been discussed for chain molecules contained in liquid Se, and a splitting of the nearest neighbour Pd-Pd and Pd-Si correlation has been clearly found in the amorphous Pd_{0.8} - Si_{0.2} alloy.

EDB SUBJECT CATEGORY: EDB-656000
INIS CATEGORY: A13

<654>

Vahed, A.; Kay, D.A.R.

Deoxidation thermodynamics of cerium and lanthanum in liquid iron EDB-76:087291; Metall. Trans., B (MTTBC) 6B:(1):285-287 (Mar 1975)

SUBJECT DESCRIPTORS: CERIUM OXIDES: T2; FREE ENERGY; IRON
ALLOYS; LANTHANUM OXIDES: T1; LIQUID METALS; PHASE
DIAGRAMS; REDUCTION: Q1, Q2; THERMODYNAMICS; VERY HIGH TEMPERATURE

ABSTRACT: The deoxidation thermodynamics of the Fe-Ce-O and Fe-La-O systems are estimated by combining the free energies of solution of cerium and lanthanum in iron, obtained from the respective phase diagrams, with the standard free energies of formation of cerium and lanthanum oxides given in the literature. Values for the deoxidation constants $(w/o \text{ Ce})^2 \cdot (w/o \text{ O})^3$ and $(w/o \text{ La})^2 \cdot (w/o \text{ O})^3$ of about 1×10^{-20} are predicted at steelmaking temperatures. These predicted values are much lower than those obtained experimentally. It is suggested that refractory interactions, and sampling and analytical difficulties, in both laboratory and industrial determinations of these deoxidation constants, lead to erroneously high results.

EDB SUBJECT CATEGORY: EDB-360202; 360105

<655>

<655>

Vaidya, V.N.; Prasad, R.; Venugopal, V.; Iyer, P.N.; Sood, D.D.;
Ramaniah, M.V., Bhabha Atomic Research Centre, Bombay (India)

In Molten salt chemistry : Part IV - Solubility behaviour of
PuF₃ in fluoride salts of interest in molten salt
reactor technology ** AIX-08:289419;EDB-77:075232; (1976)
BARC--854

ABSTRACT: Solubility of PuF₃ in eight ternary mixtures
containing LiF, BeF₂ and ThF₄ and in four binary
mixtures containing LiF and ThF₄ has been studied in
the temperature range 510-787 deg C. From the observed
solubility data, an attempt has been made to correlate the
solubility of PuF₃ at a particular temperature as a
function of composition of salt melts. Activity coefficients
of PuF₃ in various fluoride mixtures using liquid
PuF₃ as standard state, partial molal heat of mixing
and excess partial molal entropy of mixing liquid PuF₃
have also been calculated from the solubility data. These
thermodynamic data indicate that PuF₃ interacts with
the melts, the interaction being higher for less ordered
melts. It is concluded that PuF₃ is sufficiently
soluble and stable in fluoride melts of interest, making it a
suitable fuel for molten salt reactors of breeder or
converter type.

EDB SUBJECT CATEGORY: EDB-400702

INIS CATEGORY: B13

<656>

Van Uitert, L.G.; Guggenheim, H.J.; O'Bryan, H.M.; Warner, A.W.
Jr.; Brownlow, D.; Bernstein, J.L.; Pasteur, G.A.; Johnson,
L.F.

Physical properties of thorium fluoride

INS-76:023005;EDB-76:097986; Mater. Res. Bull. (MRBUA)

11: (6):669-671 (Jun 1976)

SUBJECT DESCRIPTORS: CRYSTAL STRUCTURE;DIELECTRIC
PROPERTIES;HARDNESS;MONOCRYSTALS;OPTICAL PROPERTIES;PHYSICAL
PROPERTIES: Q1;THERMAL EXPANSION;THORIUM FLUORIDES: T1

ABSTRACT: Thorium fluoride has many properties that make it of
interest for infrared windows. It is transparent to about
eleven microns, is unaffected by moisture, has a moderate
hardness, and suffers little dimensional change upon heating.

EDB SUBJECT CATEGORY: EDB-400702

INIS CATEGORY: B13

<657>

Varfolomeev, M.B.; Sotnikova, M.N.; Chibirova, F.Kh.; Shpinel, V.S. (Moskovskij Inst. Tonkoj Khimicheskoy Tekhnologii (USSR))

Some structure differences of double oxides of indium and gallium with antimony(5) EDB-76:048832; Zh. Neorg. Khim. (ZNOKA) 20:(5):1163-1166 (May 1975)

SUBJECT DESCRIPTORS: ANTIMONY OXIDES: T1;CRYSTAL STRUCTURE: Q1,Q2,Q3;CRYSTAL-PHASE TRANSFORMATIONS;MOESSBAUER EFFECT;SPECTRA;X-RAY DIFFRACTION

ABSTRACT: None

EDE SUBJECT CATEGORY: EDB-360202
INIS CATEGORY: B12

<658>

Vasil'eva, I.A.; Sukhushina, I.S.; Granovskaya, Zh.V.; Balabaeva, R.F.; Maicrova, A.P. (Moskovskij Gosudarstvennyj Univ. (USSR))

Thermodynamic properties of NbO , $\text{Fe}_{0.95}\text{O}$, CoO , NiO , Cu_{2}O and CuO oxides as comparative electrodes in electromotive forces method AIX-08:317802;EDB-77:120476; Zh. Fiz. Khim. (ZFKHA) 49:(9):2169-2174 (Sep 1975)

SUBJECT DESCRIPTORS: COMPARATIVE EVALUATIONS;ELECTRODES;ELECTROLYTES;ELECTROMOTIVE FORCE;NIOBIUM;NIOBIUM OXIDES: T1;THERMODYNAMIC PROPERTIES: Q1;THORIUM OXIDES;VERY HIGH TEMPERATURE

ABSTRACT: On the basis of critical analysis and statistical treatment of literature data on thermodynamic properties of oxides FeO , NiO and CoO , the values of $\Delta H_{\text{O}_{2}}^{\text{O}}$, $\Delta S_{\text{O}_{2}}^{\text{O}}$ and the dependences $\Delta G_{\text{O}_{2}}^{\text{O}}(T)$ with corresponding errors have been calculated in the range 1000-1400 K. For oxides NbO , Cu_{2}O and CuO , for which the literature data are not reliable enough, the above thermodynamic functions are determined experimentally by the solid electrolyte e.m.f. method. A mixture $\text{Nb}+\text{NbO}$ has been used as a reference electrode with a thorium dioxide base solid electrolyte to determine the thermodynamic properties of NbO . A convenient method of combination of the dependences $\Delta G_{\text{O}_{2}}^{\text{O}}(T) \pm \sigma(T)$ for different electrodes is proposed

<658>

<659> CONT.

which can be used in the course of treatment of experimental data obtained by the e.m.f. method.

EDB SUBJECT CATEGORY: EDB-360204

INIS CATEGORY: B12

<659>

Vasil'kcva, I.V.; Efimov, A.I.; Ryazantseva, L.V.

Interaction in AgCl-CsCl, CsCl-KCl systems and liquidus surface of AgCl-CsCl-KCl ternary system AIX-08:295418;EDB-77:056191; Vestn. Leningr. Univ., Fiz., Khim. (VLUFB) 10:(2):144-147 (May 1976)

SUBJECT DESCRIPTORS: CESIUM CHLORIDES: T1;CHEMICAL COMPOSITION;CRYSTALLIZATION;DIFFERENTIAL THERMAL ANALYSIS;EUTECTICS;MELTING POINTS;MOLTEN SALTS;PHASE DIAGRAMS;PHASE STUDIES: Q1,Q2,Q3;POTASSIUM CHLORIDES: T2;SILVER CHLORIDES: T3;X-RAY DIFFRACTION

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-400201

INIS CATEGORY: B12

<660>

Vasudeva Rao, P.R.; Patil, S.K., Bhabha Atomic Research Centre, Bombay (India)

In Spectrophotometric method for the determination of neptunium and plutonium in process solutions **
AIX-07:274807;ERA-02:027122;EDB-77:056017; (1976) BARC--842

SUBJECT DESCRIPTORS:

ARSENazo;BIBLIOGRAPHIES;CALIBRATION;NEPTUNIUM: T1;NEPTUNIUM COMPLEXES;PLUTONIUM: T2;PLUTONIUM COMPLEXES;SOLUTIONS;SOLVENT EXTRACTION;SPECTROPHOTOMETRY: Q1,Q2;TTA;ZIRCONIUM

ABSTRACT: A method is described for the determination of neptunium and plutonium in process solutions. This involves the separation of these elements followed by their spectrophotometric determinations as Arsenazo III complexes. Neptunium(IV) and plutonium(IV) are separated using TTA extraction method, and the separated Np(IV) and Pu(IV) are then determined as their Arsenazo III complexes in 5 M HNO₃. A few solutions from the reprocessing plant at Trombay are analyzed for plutonium using this method and the results were compared with those obtained by the radiometric method.

<660> CONT.

An attempt was made to use Arsenazo III to determine uranium in the plant solutions.

EDB SUBJECT CATEGORY: EDB-400104;050800
INIS CATEGORY: B11

<661>

Vatolin, N.A.; Ukhov, V.P.; Pastukhov, Eh.A.; Gel'chinskiy, B.R.; Esin, O.A.

Investigation of particles interaction in liquid metals and alloys EDB-76:036470; Izv. Akad. Nauk SSSR, Met. (IZNMA) (1):45-49 (Jan 1975)

SUBJECT DESCRIPTORS: BOLTZMANN EQUATION;CORRELATION FUNCTIONS;DISTRIBUTION FUNCTIONS;INTERATOMIC DISTANCES;IRON ALLOYS: T1;LIQUID METALS;NICKEL ALLOYS: T2;PARTICLE INTERACTIONS;PERCUS-YEVICK EQUATION;SILICON ALLOYS: T3;STRUCTURAL CHEMICAL ANALYSIS;THERMODYNAMIC MODEL;THERMODYNAMIC PROPERTIES: Q1,Q2,Q3;X-RAY DIFFRACTION

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-360104
INIS CATEGORY: B22

<662>

Veal, B.W.; Lam, D.J.; Hoekstra, H.R.; Diamond, H.; Carnall, W.T., Argonne National Lab., Ill. (USA)

In XPS studies of actinide materials **
ERA-02:030285;INS-77:008094;EDB-77:069590; ([nd])
CONF-760956--4

SUBJECT DESCRIPTORS: ACTINIDES: M14;AMERICIUM OXIDES: M11;BAND THEORY;BERKELIUM OXIDES: M13;CALIFORNIUM OXIDES: M8;CRYSTAL FIELD;CURIUM OXIDES: M12;ELECTRONIC STRUCTURE: Q1,Q2,Q3,Q4,Q5,Q6,Q7,Q8,Q9,Q10,Q11,Q12,Q13,Q14;EMISSION SPECTRA;ENERGY LEVELS;NEPTUNIUM OXIDES: M5;PHOTOEMISSION;PLUTONIUM OXIDES: M10;REVIEWS;THORIUM: M2;THORIUM OXIDES: M1;TRANSURANIUM ELEMENTS: M4;URANIUM: M6;URANIUM COMPOUNDS: M7;URANIUM OXIDES: M9;URANIUM-ALPHA: M3;X-RAY SPECTRA

ABSTRACT: Applications of x-ray photoemission spectroscopy (XPS) to the study of actinide materials are reviewed. Examples discussed here include the band structures of thorium and uranium metal, the multiplet structure associated with the 5f electron states in oxides of the transuranium elements, the test for temporal configurations in NpO_2^{2+} , crystal field

<662>

<662> CONT.

splitting of the $U 6p_{3/2}$ level in a series of uranyl compounds, mixed oxidation states in $Cf_{7f_{12}}$, and a test for the participation of 5f electrons in bonding in a series of uranium compounds.

EEL SUBJECT CATEGORY: EDB-360106;360206

INIS CATEGORY: B25

<663>

Veleckis, E.; Maroni, V.A. ** Watson, J.S.; Wiffen, F.W.; Bishop, J.L.; Breeden, B.K. (eds.), Oak Ridge National Lab., Tenn. (USA)

Thermodynamic properties of solutions of hydrogen isotopes in metals and alloys of interest to fusion reactor technology In Radiation effects and tritium technology for fusion reactors
** ERA-02:022938;INS-77:005755;EDB-77:043997; (Mar 1976)
CONF-750989--P3

SUBJECT DESCRIPTORS: ALUMINIUM HYDRIDES: T4;DEUTERIUM: T5;HYDROGEN: T6;ISOTOPE EFFECTS: Q5;LIQUID METALS:LITHIUM: T3;LITHIUM DEUTERIDES: T2;LITHIUM HYDRIDES: T1;SOLUBILITY: Q2,Q3;SOLVENT PROPERTIES: Q1,Q4,Q6;THERMONUCLEAR REACTOR MATERIALS

ABSTRACT: Recent ANL work on the solution thermodynamics of systems composed of a hydrogen isotope and either pure lithium or a lithium-containing alloy has been directed toward (1) the sources and magnitude of isotope effects in the systems Li--LiH and Li--LiD, (2) the solubility of LiD in liquid lithium (200 to 700 $^{\circ}$ C), and (3) the solubility and decomposition pressure of hydrogen in the ternary Li--Al--H system. The H/D isotope effect appears to originate primarily in the gas phase and within the two-phase regions of the phase diagram, its range of values is 1.43 ± 0.05 . At 200 $^{\circ}$ C, the solubility of LiD in lithium is approximately 250 appM D, thus essentially ruling out hydrogen isotope separation methods based on cold-trapping concepts. Two isotherms (at 550 and 599 $^{\circ}$ C) have been obtained for solutions of hydrogen in the β -phase of the Li--Al system (48 to 56 mol % Li). They indicate that, at low hydrogen concentrations (<1 atom % H), the dissociation pressures are much higher ($\times 100$) than those over the lithium metal. Within the miscibility gap, which extends from 6 to 33.3 atom % D, the plateau pressures, P , may be represented by $\ln P$ (torr) = $20.46 - 11,600/T$.

EEL SUBJECT CATEGORY: EDB-360603;700209;400203

INIS CATEGORY: B24;A14;B12

<664>

Vinogradov, E.E.; Kirilenko, I.A.; Krasilov, Yu.I.; Solokha, A.F. (AN SSSR, Moscow. Inst. Obshchej i Neorganicheskoy Khimii)

Structure of Nd^{3+} terms in nitrate glasses

AIX-08:283412; EDB-77:036910; Izv. Akad. Nauk SSSR, Neorg. Mater. (IVNMA) 11:(8):1501-1505 (Aug 1975)

SUBJECT DESCRIPTORS: CATIONS; ENERGY LEVELS:

Q1, Q2, Q3; GLASS; LUMINESCENCE; MAGNESIUM NITRATES: T1; NEODYMIUM NITRATES: T2; SODIUM NITRATES: T3; SPECTRA; STARK EFFECT

ABSTRACT: The structure of the neodymium energy levels in nitrate glasses and crystals has been studied. In these matrices an activator has a highly-symmetrical identical surroundings conditioned by the existence of weakly distorted neodymium-nitrate complexes. Great amounts of water and hydroxyl present in the matrix cause deactivation of the neodymium excited state.

EDB SUBJECT CATEGORY: EDB-360204

INIS CATEGORY: B24

<665>

Vinogradov, E.E.; Kirilenko, I.A. (AN SSSR, Moscow. Inst. Obshchej i Neorganicheskoy Khimii)

Constitution equilibrium diagram of the $\text{LiNO}_3\text{-LiIO}_3$, $\text{NaNO}_3\text{-NaIO}_3$, $\text{KNO}_3\text{-KIO}_3$ system

AIX-08:289374; EDB-77:050852; (ZNOKAZh. Neorg. Khim.) 21:(4):1157-1159 (Apr 1976)

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-400201

INIS CATEGORY: P12

<666>

Vinogradov, E.E.; Kirilenko, I.A.

Glass formation in nitrate systems containing rare earth ions

AIX-08:289375; EDB-77:050853; (ZNOKAZh. Neorg. Khim.) 21:(4):1130-1132 (Apr 1976)

SUBJECT DESCRIPTORS: DIFFERENTIAL THERMAL

ANALYSIS; GLASS; MAGNESIUM NITRATES: T1; PHASE DIAGRAMS:

Q1, Q2, Q3; RARE EARTH COMPOUNDS: T2; SODIUM NITRATES: T3; VITRIFICATION

<666>

<666> CONT.

ABSTRACT: None

EDE SUBJECT CATEGORY: EDB-400201

INIS CATEGORY: B12

<667>

Vinogradov, E.E.; Kirilenko, I.A. (AN SSSR, Moscow. Inst.
Obshchej i Neorganicheskoy Khimii)

Phase diagram of $\text{NaNO}_3\text{--Nd}(\text{NO}_3)_3$ system
AIX-08:295420; EDB-77:056239; Zh. Neorg. Khim. (ZNOKA)
21: (7): 1986-1987 (Jul 1976)

SUBJECT DESCRIPTORS: CHEMICAL PREPARATION; HIGH
TEMPERATURE; NEODYMIUM NITRATES: T1; PHASE DIAGRAMS:
Q1, Q2; QUANTITY RATIO; SODIUM NITRATES: T2

ABSTRACT: None

EDE SUBJECT CATEGORY: EDB-400201

INIS CATEGORY: B12

<668>

Vinogradov, E.E.; Kirilenko, I.A. (AN SSSR, Moscow. Inst.
Obshchej i Neorganicheskoy Khimii)

Phase diagrams of the $\text{Mg}(\text{NO}_3)_2\text{--Nd}(\text{NO}_3)_3$
system AIX-08:337041; EDB-77:140956; Zh. Neorg. Khim. (ZNOKA)
22: (2): 583-585 (Feb 1977)

SUBJECT DESCRIPTORS: EUTECTICS; MAGNESIUM NITRATES: T1; MELTING
POINTS; NEODYMIUM NITRATES: T2; PHASE DIAGRAMS: Q1, Q2

ABSTRACT: None

EDE SUBJECT CATEGORY: EDB-400201

INIS CATEGORY: B12

<669>

Visser, R.G.; Schiavi, W.F.; Berard, M.F.

Interdiffusion in the $\text{CaF}_2\text{--YF}_3$ system
ERA-01:016754; EDB-76:042456; J. Am. Ceram. Soc. (JACTA)
58: (9-10): 438-441 (1975)

SUBJECT DESCRIPTORS: CALCIUM FLUORIDES: T1; DEFECTS; DIFFUSION:
Q1, Q2; ELECTRON PROBES; SOLUBILITY; TEMPERATURE DEPENDENCE; VERY
HIGH TEMPERATURE; YTTRIUM FLUORIDES: T2

<669> CONT.

ABSTRACT: An electron microprobe interdiffusion study was made in single crystals between 1200 and 1331⁰C over the solid-solution compositional range 0 to 20 mol percent YF₃ in CaF₂. At constant temperature, the value $\tilde{D}$, the interdiffusion coefficient, increased approximately exponentially with increasing YF₃ content. As YF₃ content increased, the temperature dependence of $\tilde{D}$ decreased. Extrapolation of $\tilde{D}$ to 0 percent YF₃ combined with consideration of the defect formation mechanism at infinite dilution allowed estimation of the impurity diffusion coefficient for Y in essentially pure CaF₂.

EDB SUBJECT CATEGORY: EDB-360603

<670>

Viting, L.M.; Gorbovskaya, G.P. (Moskovskii Gosudarstvennyi Univ. (USSR))

Estimate of cryoscopic calculations accuracy from fusibility diagrams AIX-08:280706; EDB-77:031069; Izv. Akad. Nauk SSSR, Neorg. Mater. (IVNMA) 11:(7):1237-1240 (Jul 1975)

ABSTRACT: The melting points of some lead and zinc salts, that can be used as solvents for ferrites in systems: PbMoO₄-MgFe₂O₄, Zn₂V₂O₇-NiFe₂O₄, Pb₃(VO₄)₂-MgFe₂O₄, have been calculated in accordance with the hypothetical mechanism of the solvent dissociation. The accuracy of cryoscopic calculations based on melting point curves is evaluated. Cryoscopic calculations permit to determine the solvent activity with the accuracy of $\pm 0.3\%$ and the heat of its fusion, with the accuracy of $\pm 3\%$. The comparison of the calculated and experimental values of the entropy of melting, as well as calculated and experimental values of the cryoscopic constant elucidates the mechanism of dissociation of both the dissolved compound and the solvent.

EDB SUBJECT CATEGORY: EDB-360602

INIS CATEGORY: B12

<671>

Vivchar, O.I.; Zarechnyuk, O.S. (L'vovskij Gosudarstvennyj Univ. (Ukrainian SSR)) ** Ministerstvo Vysshego i Srednego Spetsial'nogo Obrazovaniya SSSR, Moscow; L'vovskij Gosudarstvennyj Univ. (Ukrainian SSR)

Compounds with the structure of ThMn₁₂ type in the R-Fe-Al systems In 2. all-union conference on crystallochemistry of intermetallic compounds ** AIX-07:275147; EDB-77:050198; (1974) INIS-mf--3191

<671>

<671> CONT.

SUBJECT DESCRIPTORS: ALKALI METALS: M4; ALUMINIUM ALLOYS:
M1; INTERMETALLIC COMPOUNDS; IRON ALLOYS: M2; LATTICE
PARAMETERS: Q1, Q2, Q3, Q4; RARE EARTH ALLOYS: M3; TERNARY ALLOY
SYSTEMS; TETRAGONAL LATTICES; X-RAY DIFFRACTION

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-360102
INIS CATEGORY: B22

<672>

Volkov, S.V. (AN Ukrainskoj SSR, Kiev. Inst. Obshchej i
Neorganicheskoy Khimii)

Isorhodate complexes of 3d-metal ions in molten salts
AIX-08:286214; EDB-77:044310; Koord. Khim. (KOKHD)
2: (8):1106-1109 (Aug 1976)

SUBJECT DESCRIPTORS: CHEMICAL BONDS; ELECTRONIC STRUCTURE; MOLTEN
SALTS; RHODIUM COMPLEXES: T2; SPECTROSCOPY; THERMODYNAMICS:
Q1, Q2; THIOCYANATES; VANADIUM COMPLEXES: T1

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-400201
INIS CATEGORY: B12

<673>

Volkov, S.V.; Buryak, N.I. (AN Ukrainskoj SSR, Kiev. Inst.
Obshchej i Neorganicheskoy Khimii)

Effect of second sphere cations on formation of nickel (2)
chloride complexes in molten salts
AIX-08:339996; ERA-03:012887; EDB-78:022276; Zh. Neorg. Khim.
(ZNCKA) 22: (1):150-154 (Jan 1977)

SUBJECT DESCRIPTORS: ABSORPTION SPECTRA; ALKALI METAL COMPLEXES:
T2; CESIUM CHLORIDES; CESIUM NITRATES; CHEMICAL
PREPARATION; ELECTRONIC STRUCTURE: Q1, Q2; EQUILIBRIUM; MOLTEN
SALTS; NICKEL COMPLEXES: T1

ABSTRACT: The effect of the outer-sphere cations of alkali
metals (A-Li^+ , Na^+ , K^+ , Rb^+ or Cs^+) on complexing of Ni(2) in melts of $\text{ANO}_{3\text{H}}$ and ACl has been studied. The effect of these cations on the
equilibrium octahedral-tetrahedral nickel (2) complexes in
melts of $\text{ANO}_{3\text{H}}$ and ACl has been elucidated. It has been
established that octahedral complex $[\text{NiCl}_6]^{4-}$ is
formed with an outer-sphere cation Li^+ , tetrahedral
complex $[\text{NiCl}_4]^{2-}$ with Rb^+ and Cs^+ ,

<673> CONT.

and equilibrium $[\text{NiCl}_6]^{4-}$ is observed in the case of Na^+ and K^+ . The regularities in a change of spectroscopic parameters of the complex depending on an outer-sphere cation have been investigated.

EDB SUBJECT CATEGORY: EDB-400201

INIS CATEGORY: B12

<674>

Vclkov, S.V.; Evtushenko, N.P.; Yatsimirskii, K.B. (AN Ukrainskoj SSR, Kiev. Inst. Fizicheskij Khimii; AN Ukrainskoj SSR, Kiev. Inst. Obshchej i Neorganicheskij Khimii)

Spectra of combination scattering and structure of 3d-metals chloride complexes in molten salts
AIX-08:286215; EDB-77:044239; Teor. Eksp. Khim. (TEKHA) 12:(1):111-115 (Jan 1976)

SUBJECT DESCRIPTORS: CESIUM COMPLEXES: T1; CHLORIDES; CRYSTAL STRUCTURE: Q1; ELECTRON SPECTRA; MOLTEN SALTS; SCATTERING; SPECTRA; VALENCE; VIBRATIONAL STATES

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-400201

INIS CATEGORY: B12

<675>

Volkovich, A.V.; Potapenko, O.G.; Semin, A.N.; Nichkov, I.F. (Moskovskij Khimiko-Tekhnologicheskij Inst. (USSR). Novomoskovskij Filial; Ural'skij Politekhnikeskij Inst., Sverdlovsk (USSR))

Anodic behavior of cadmium and cadmium-zinc alloys in eutectic melt of potassium- and lithium chlorides
AIX-07:272465; EDB-77:026035; Izv. Vyssh. Uchebn. Zaved., Tsvet. Metall. (IVUTA) (4):153-155 (1975)

SUBJECT DESCRIPTORS: ANODES; CADMIUM: T1; CADMIUM ALLOYS: T2; DISSOLUTION; ELECTROCHEMISTRY: Q1, Q2, Q3; EUTECTICS; LITHIUM CHLORIDES; MOLTEN SALTS; POLARIZATION; POTASSIUM CHLORIDES; TEMPERATURE DEPENDENCE; ZINC ALLOYS: T3

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-360100; 400400

INIS CATEGORY: B12

<676>

<676>

Vollrath, D., Kernforschungszentrum Karlsruhe (Germany, F.R.).
Inst. fuer Material- und Festkoerperforschung

In Parameter for optimizing the synthesis of (U,Pu)C **
INS-76:022946; ERA-02:007615; EDB-77:003012; (1976)
CRNL-tr--4179

SUBJECT DESCRIPTORS: ACTIVATION ENERGY; BINARY MIXTURES; CHEMICAL
REACTION KINETICS:
Q3, Q4; GRAPHITE; GRINDING; LOSSES; OPTIMIZATION; PHASE
STUDIES; PLUTONIUM CARBIDES: T2; PLUTONIUM DIOXIDE:
T4; SYNTHESIS: Q1, Q2; THERMAL GRAVIMETRIC ANALYSIS; URANIUM
CARBIDES: T1; URANIUM DIOXIDE: T3

ABSTRACT: Experiments were carried out on the carbothermal
reaction of mixed (U,Pu)O₂ with graphite. The reaction
was followed with an electronic thermobalance, and the phases
were analyzed roentgenographically. Results showed that for
the mixed powder, the reactions of the U and Pu phases are
independent of each other, whereas in powders ground for 36
h, U_{0.8}Pu_{0.2}O₂ formed first,
followed by (U,Pu) (O_x/C_{1-x}), (U,Pu)₂C₃,
and finally U_{0.8}Pu_{0.2}C. Nucleus
formation occurs in the 36-h-milled UO₂--PuO₂--
graphite mixture during the milling process. Pu losses
can be minimized by lower temperatures and by formation of
mixed carbide. (DLC)

EDF SUBJECT CATEGORY: EDB-360201; 050700
INIS CATEGORY: B23

<677>

Voloshin, V.A.; Krimnus, M.G. (Donetskij Fiziko-Tekhnicheskij
Inst. (Ukrainian SSR))

Pressure effect on spectral properties of CeF₃:Pr³⁺
AIX-08:286216; EDB-77:044216; Ukr. Fiz. Zh. (Russ. Ed.)
(UFIZA) 20:(8):1266-1272 (Aug 1975)

SUBJECT DESCRIPTORS: ABSORPTION SPECTRA: Q1, Q2; CATIONS; CERIUM
FLUORIDES: T1; COUPLING CONSTANTS; ELECTROMAGNETIC
INTERACTIONS; INTERATOMIC DISTANCES; L-S COUPLING; LOW
TEMPERATURE; MONOCRYSTALS; PRASEODYMIUM; PRASEODYMIUM COMPOUNDS:
T2; PRESSURE DEPENDENCE; ULTRALOW TEMPERATURE; VERY HIGH PRESSURE

ABSTRACT: The absorption spectrum of a single crystal of CeF₃:Pr³⁺
has been studied in the range of 400 to 700 nm
at the atmospheric pressure and temperatures of 78 and 4.2
deg K, as well as at the pressure of 25 kbar and a
temperature of 78 deg K. Under pressure a slight change in
the Stark splitting has been observed together with a more

<677> CONT.

significant shift of the j-levels into the long-wavelength region of the spectrum (approximately by 15 cm^{-1}). Calculated from the experimental data are the parameters of the electrostatic and spin-orbital interactions F_2 and $\zeta(4f)$, as well as the nephelauxetic factor $\beta = F_2(K)/F_2(fr)$, where $F_2(K)$ and $F_2(fr)$ are the Slater parameters for Pr^{3+} in the crystal and free ion of Pr (4), respectively. Also observed under pressure is an increase in the parameter of bond $b^{(1/2)}[(1-\beta)/2]^{(1/2)}$. It is assumed that the shorter interatomic distances in the case of higher pressure result in the covalence contribution into the chemical bond becoming more substantial.

EDB SUBJECT CATEGORY: EDB-400201
INIS CATEGORY: B12

<678>

von Amberger, H.D.; Rosenbauer, G.G. (Erlangen-Nuernberg Univ., Erlangen (Germany, F.R.). Inst. fuer Anorganische Chemie); Fischer, R.D.

Electronic structure of highly symmetric lanthanoid and actinoid compounds. VI. The electronic Raman spectrum of $\text{Cs}_2\text{NaCe(III)Cl}_6$ AIX-08:294154; EDB-77:068761; Mol. Phys. (MOPHA) 32:(5):1291-1298 (Nov 1976)

SUBJECT DESCRIPTORS: CERIUM CHLORIDES: T1; CESIUM CHLORIDES: T2; CRYSTAL FIELD; ELECTRONIC STRUCTURE: Q1, Q2, Q3; GROUND STATES; LOW TEMPERATURE; MAGNETIC SUSCEPTIBILITY; RAMAN SPECTRA; SODIUM CHLORIDES: T3; TEMPERATURE DEPENDENCE

ABSTRACT: The low-temperature Raman spectrum of the elpasolite system $\text{Cs}_2\text{NaCe(III)Cl}_6$ exhibits five new sharp lines (compared with the room-temperature spectrum) which are assigned to crystal-field (=CF) Raman transitions between the CF ground state $\Gamma_7(\frac{5}{2})$ and the partially split 'octahedral' CF states of the free ion terms $\frac{5}{2}F_5(\frac{5}{2})$ and $\frac{5}{2}F_7(\frac{7}{2})$. A crystal-field calculation yields: $\zeta(4f) = 635 \text{ cm}^{-1}$, $B_4 = 248 \text{ cm}^{-1}$ and $B_6 = 2,4 \text{ cm}^{-1}$. The temperature dependence of the paramagnetic susceptibility calculated on the basis of these parameters is in good agreement with earlier experimental results.

EDB SUBJECT CATEGORY: EDB-360602
INIS CATEGORY: A12

<679>

<679>

von Goldbeck, O. ** Kubaschewski, O. (ed.)

Phase diagrams In Thorium: physico-chemical properties of its compounds and alloys ** EDB-76:036261; (1975) International Atomic Energy Agency

SUBJECT DESCRIPTORS: BINARY ALLOY SYSTEMS; HIGH TEMPERATURE; MEDIUM TEMPERATURE; PHASE DIAGRAMS: Q1; REVIEWS; THORIUM ALLOYS: T1; THORIUM CARBIDES; VERY HIGH TEMPERATURE

ABSTRACT: The phase diagrams of binary alloy systems of thorium with different elements in various temperature regions from 400⁰C to 3000⁰C are given. Altogether 44 binary systems are reviewed.

EDE SUBJECT CATEGORY: EDB-360102

INIS CATEGORY: B22

<680>

von Goldbeck, O. (Technische Hochschule Aachen (Germany, F.R.). Lehrstuhl fuer Metallurgie der Kernbrennstoffe und Theoretische Huettenkunde)

Phase diagrams In Zirconium: Physico-chemical properties of its compounds and alloys ** AIX-07:277762; EDB-77:026064; (1976) IAEA

SUBJECT DESCRIPTORS: BINARY ALLOY SYSTEMS; PHASE DIAGRAMS: Q1; REVIEWS; SILVER ALLOYS; SOLUBILITY; ZIRCONIUM ALLOYS: T1

ABSTRACT: The phase diagrams of binary alloys of Zr with 64 different elements are discussed and the diagrams given. For a few of the alloys, a complete phase diagram was not available. The elements are: Ag, Al, As, Au, B, Be, Bi, C, Cd, Ce, Co, Cr, Cu, Dy, Er, Fe, Ga, Gd, Ge, H, Hf, Hg, Ho, In, Ir, K, La, Li, Mg, Mn, Mo, N, Na, Nb, Nd, Ni, O, Os, P, Pd, Pt, Pu, rare earths (solubility in α -Zr and β -Zr), Re, Rh, Ru, S, Sb, Sc, Se, Si, Sm, Sn, Ta, Tb, Tc, Te, Th, Ti, U, V, W, Y, Zu.

EDE SUBJECT CATEGORY: EDB-360102

INIS CATEGORY: B12

<681>

von Tippelskirch, H. (Karlsruhe Univ. (TH) (Germany, F.R.).
Inst. fuer Physikalische Chemie und Elektrochemie)

Viscosities of cesium vapor to 1620 K and of liquid gallium to
1800 K AIX-08:334614; EDB-77:132682; Ber. Bunsenges. Phys.
Chem. (BBPCA) 80: (8):726-729 (Aug 1976)

SUBJECT DESCRIPTORS: CESIUM: T3; DENSITY; GALLIUM: T1; LIQUID
METALS; MEASURING METHODS; TABLES; TEMPERATURE DEPENDENCE; VAPOR
PRESSURE; VAPORS; VERY HIGH TEMPERATURE; VISCOSITY: Q1, Q3

ABSTRACT: The viscosity of cesium at 1620 K and 40 bar has been
determined to $41 \times 10^{\text{sup}} -6$ (Pa.s) by the oscillating cup
method. The saturated vapor density at 1580 K could be
derived from the viscosity measurements. The viscosity of
liquid gallium has been determined from 370 K to 1800 K. The
experimental results have been compared with calculations
based on the Enskog hard-sphere transport theory for dense
fluids.

EDB SUBJECT CATEGORY: EDB-360104
INIS CATEGORY: B22

<682>

Vorobei, M.P.; Desyatnik, V.N.; Kurbatov, N.N.; Skiba, O.V.

Interaction of plutonium trichloride in melts of ternary systems
AIX-08:334441; EDB-77:133673; Radiokhimiya (RADKA)
18: (1):119-122 (Jan 1976)

ABSTRACT: The systems $\text{KCl-UCl}_{\text{sub } 3} - \text{PuCl}_{\text{sub } 3}$, $\text{NaCl-ThCl}_{\text{sub } 4} - \text{PuCl}_{\text{sub } 3}$, $\text{KCl-ThCl}_{\text{sub } 4} - \text{PuCl}_{\text{sub } 3}$ have been studied
by differential-thermal analysis. Meltability diagrams have
been plotted and invariant points of the systems have been
determined as well as low-temperature regions of the
compositions which can be recommended for using as
electrolytes in the process of producing or refining
plutonium by electrochemical method.

EDB SUBJECT CATEGORY: EDB-400702
INIS CATEGORY: B13

<683>

<683>

Waber, J.T. (Commission of the European Communities, Karlsruhe
(F.R. Germany). European Inst. for Transuranium Elements) **
Mueller, W.; Lindner, R. (eds.)

Electronic structure of actinides and related metals In
Transplutonium 1975 **
AIX-07:270441; ERA-02:020483; EDB-77:036581; (1976)
North-Holland

SUBJECT DESCRIPTORS: AMERICIUM: T6; BAND THEORY; BRILLOUIN
ZONES; CERIUM; CURIUM: T7; EIGENVALUES; ELECTRONIC STRUCTURE:
Q1, Q2, Q3, Q4, Q5, Q6, Q7; MATRIX ELEMENTS; NEPTUNIUM: T4; OCCUPATION
NUMBER; PLUTONIUM: T5; PROTACTINIUM: T2; SLATER METHOD; THORIUM:
T1; TITANIUM; URANIUM: T3; WAVE FUNCTIONS

ABSTRACT: In order to elucidate the role of 5f-electrons in the
light actinide elements, a series of Confined Atom
calculations are reported. These emphasize the sensitivity of
the 5f electrons to the volume available for the Pu atoms
i.e. the electronic configurations which minimize the total
energy vary with the sphere radius $R_{sub}(s)$. The problems and
significance of finding which states are occupied in the
Brillouin zone, are reviewed to illustrate a point made by
Freeman at other conferences. Since relativistic effects are
of considerable importance in discussing both light and heavy
actinides, the structure of Dirac-Slater wave functions are
briefly commented on.

EDB SUBJECT CATEGORY: EDB-360102
INIS CATEGORY: B22

<684>

Walch, P.F.; Ellis, D.E.

Effects of secondary ligands on the electronic structure of
uranyls INS-76:020005; ERA-02:001215; EDB-76:081893; J. Chem.
Phys. (JCPA) 65: (6):2387-2392 (15 Sep 1976)

SUBJECT DESCRIPTORS: CRYSTAL FIELD; ELECTRONIC STRUCTURE:
Q1; LIGANDS; MOLECULAR ORBITAL METHOD; URANIUM OXIDES; URANYL
COMPLEXES: T1

ABSTRACT: The effects of secondary ligands on the electronic
structure of uranyl, UO_4^{2+} , are investigated in a
perturbing point ion model, making use of the relativistic
Dirac-Slater molecular orbital (MO) approach. Variation of
'crystal-field splittings' with primary U-O bond length is
explored; relaxation of the uranyl MO's is taken into account
by self-consistent iterations. The theoretical splittings are
found to agree rather well with the x-ray photoemission
spectroscopy (XPS) data of Veal et al. Comparison is made

<684> CONT.

with optical data and reasons are given for the poor agreement. (AIP)

EDB SUBJECT CATEGORY: EDB-400702
INIS CATEGORY: B13

<685>

Wang, W.L.; Singh, R.N.

Equilibrium index for fission gas bubbles
ERA-02:056709; INS-77:015992; EDB-77:131685; Trans. Am. Nucl. Soc. (TANSA) 26::377-378 (Jun 1977)

SUBJECT DESCRIPTORS: BUBBLES: Q1; COALESCENCE; FISSION
PRODUCTS; GASES; SOLID FUELS: T1; URANIUM DIOXIDE

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-220300
INIS CATEGORY: E23

<686>

Waseda, Y. (Tohoku Univ., Sendai (Japan). Research Inst. of Mineral Dressing and Metallurgy) ** Evans, R.; Greenwood, D.A. (eds.)

Structure of liquid transition metals and their alloys In Liquid metals, 1976 ** AIX-08:319649; EDB-77:113431; (1977) Institute of Physics

SUBJECT DESCRIPTORS: ATOMIC NUMBER; BIBLIOGRAPHIES; ELECTRONIC STRUCTURE: Q1; INTERATOMIC DISTANCES; INTERATOMIC FORCES; LIQUID METALS; RARE EARTH ALLOYS; RARE EARTHS; REVIEWS; TRANSITION ELEMENT ALLOYS: T1; TRANSITION ELEMENTS; VARIATIONS

ABSTRACT: An attempt is made in this review to cover the characteristics of the structure of liquid transition metals, including the lanthanide elements, and alloys with a transition metal as a constituent. One difference in the structure of liquid transition metals from that of normal metals is that, as the atomic number increases from Ti to Ni or from Ce to Yb, the oscillations in the structure factor increase in amplitude. This gives qualitative support for the suggestion that the partial overlap between atoms which have a nearly empty 3d-shell or 4f-shell, such as Ti or Ce, is larger than those having a nearly filled 3d-shell or 4f-shell, such as Ni or Yb. Similar behaviour is found in the structure of liquid Al-Ti and Al-Fe alloys. Results are

<686>

<686> CONT.

also given for liquid transition metal-metalloid systems,
such as Ni-P alloys.

EDE SUBJECT CATEGORY: EDB-360102

INIS CATEGORY: A13

<687>

Waseda, Y.; Tamaki, S.

Structures of 3d-transition metals in the liquid state

EDB-76:042168; Philos. Mag. (PHMAA) 32:(2):273-281 (1975)

SUBJECT DESCRIPTORS: CHROMIUM: T3; COBALT: T6; COORDINATION
NUMBER; FOURIER ANALYSIS; INTERATOMIC DISTANCES; IRON: T5; LIQUID
METALS; MANGANESE: T4; MELTING POINTS; MOLECULAR STRUCTURE:
Q1, Q2, Q3, Q4, Q5, Q6, Q7; NICKEL: T7; TITANIUM: T1; TRANSITION
ELEMENTS; VANADIUM: T2; X-RAY DIFFRACTION

ABSTRACT: The structures of 3d-transition metals (Ti, V, Cr, Mn,
Fe, Co, Ni, and Cu) in the liquid state have been
systematically examined by x-ray diffraction at temperatures
above their mp. In terms of the usual Fourier analysis, the
atomic radial distribution functions were obtained from which
interatomic distances and coordination numbers were estimated.

EDE SUBJECT CATEGORY: EDB-360102

<688>

Waseda, Y.; Tamaki, S. (Tohoku Univ., Sendai (Japan). Research
Inst. of Mineral Dressing and Metallurgy)

Structure of rare earth metals in the liquid state

AIX-08:338942; EDB-78:021630; Philos. Mag. (PHMAA) 36:(1):1-8
(Jul 1977)

SUBJECT DESCRIPTORS: CERIUM: T1; COORDINATION NUMBER; DISTRIBUTION
FUNCTIONS; ELECTRONIC STRUCTURE:
Q1, Q2, Q3, Q4, Q5, Q6, Q7, Q8, Q9; EUROPIUM: T2; FOURIER
ANALYSIS; GADOLINIUM: T3; INTERATOMIC DISTANCES; LANTHANUM:
T4; LIQUID METALS: T5; PRASEODYMIUM: T6; SCANDIUM: T7; TERBIUM:
T8; X-RAY DIFFRACTION; YTTERBIUM: T9

ABSTRACT: The structure of liquid rare earth metals (Sc, La, Ce,
Pr, Eu, Gd, Tb and Yb) have been systematically examined by
X-ray diffraction at temperatures above their melting points.
The structure factors of liquid La, Ce, Pr, Eu and Yb were
temperature insensitive in the presently available
experimental temperature range. In terms of the usual
Fourier analysis, the atomic radial distribution functions

<688> CONT.

were obtained from which interatomic distances and coordination numbers were estimated.

EDB SUBJECT CATEGORY: EDB-360102

INIS CATEGORY: A13

<689>

Waseda, Y.; Tamaki, S. (Tohoku Univ., Sendai (Japan). Research Inst. of Mineral Dressing and Metallurgy).

Structure of liquid Ce-Ni alloys AIX-08:329099;EDB-78:021633; J. Phys., F (London) (JPFMA) 7:(6):L151-L153 (Jun 1977)

SUBJECT DESCRIPTORS: CERIUM ALLOYS: T1;CHARGE TRANSPORT;CRYSTAL STRUCTURE: Q1,Q2;HARD-CORE POTENTIAL;LIQUID METALS;NICKEL ALLOYS: T2;X-RAY DIFFRACTION

ABSTRACT: The structures of liquid Ce-Ni alloys with 25 and 75at% Ni have been investigated by x ray diffraction. The results suggest that the charge transfer effect makes a major contribution to the effective hard core diameter of Ce-Ce pairs in these alloys.

EDB SUBJECT CATEGORY: EDB-360102

INIS CATEGORY: A13

<690>

Waterbury, G.R. (comp.), Los Alamos Scientific Lab., N.Mex. (USA)

In Analytical methods for fissionable materials in the nuclear fuel cycle. Progress report, July 1, 1975--September 30, 1976 ** INS-77:008150;ERA-02:032889;EDB-77:082774; (Dec 1976) LA--6577-PF

SUBJECT DESCRIPTORS: AUTOMATION;CHEMICAL ANALYSIS: Q2,Q3,Q4;DISSOLUTION;FISSIONABLE MATERIALS: T2;FUEL CYCLE;HALF-LIFE: Q1;ISOTOPE DILUTION;MASS SPECTROSCOPY: Q5;PLUTONIUM: T3;PLUTONIUM 239: T1;PLUTONIUM OXIDES;SAMELE PREPARATION: Q1;SEPARATION PROCESSES;URANIUM: T4;URANIUM 235: T5

ABSTRACT: Progress continued on development of dissolution techniques for difficult-to-dissolve nuclear materials, development of methods and automated instruments for determinations of plutonium and uranium, preparation of plutonium-containing materials for the Safeguards Analytical Laboratory Evaluation (SALE) program, analysis of SALE uranium materials, and measurement of plutonium isotope half-lives. Gas-solid reactions at elevated temperatures using reactive gases such as chlorine continue to show

<690>

<690> CONT.

promise for separating uranium from refractory materials. An extensive study of nonaqueous solvents for the dissolution of refractory materials is in progress. An extraction-separation procedure, highly specific for microgram amounts of uranium, has been developed, and its adaptation to the Los Alamos Scientific Laboratory (LASL) automated spectrophotometer is being evaluated. Development of an electrometric analysis method for plutonium is nearing completion, and design of an automated instrument using the method has been started. Batches of plutonium oxide and mixed uranium-plutonium, intended for issue as Secondary Reference and Calibration Test Materials, are being recharacterized for assay and isotopic contents. The half-life of ^{239}Pu has been determined by isotope-dilution mass-spectrometric measurement of ^{235}U grow-in as a function of time.

EDB SUBJECT CATEGORY: EDB-400100;055001

INIS CATEGORY: P11;F41

<691>

Wedemeyer, H.; Guenther, E., Kernforschungszentrum Karlsruhe (Germany, F.R.). Inst. fuer Material- und Festkoerperforschung; Kernforschungszentrum Karlsruhe (Germany, F.R.). Projekt Schneller Brueter

In Fabrication of uranium-plutonium mixed carbide **

AIX-07:255197;ERA-02:020576;EDB-77:068633; (Dec 1975)
KPK--2238

SUBJECT DESCRIPTORS: CHEMICAL COMPOSITION;CHEMICAL PREPARATION;DENSITY;FABRICATION: Q1,Q2;GRINDING;LOSSES;MICROSTRUCTURE;MIXTURES;PHASE DIAGRAMS;PLUTONIUM;PLUTONIUM CARBIDES: T1;PRESSURE DEPENDENCE;PRODUCTION;REDUCTION;SINTERING;SOLID SOLUTIONS;TEMPERATURE DEPENDENCE;TIME DEPENDENCE;URANIUM CARBIDES: T2;X-RAY DIFFRACTION

ABSTRACT: In the present report the carbide fuel fabrication-line of the Institut fuer Material- und Festkoerperforschung (IMF) is outlined. The carbide fabrication by carbothermic reduction as well of original produced urania-/plutonia-powders as of sintered oxide waste is described. In special, test-results of the stoichiometry-behaviour and the reduction of plutonium-losses are given. With this, the reduction of losses is possible by an early step of formation of mixed-crystals within the carbide fabrication process. With the present results it

<691> CONT.

seems to be possible, to depress the total material-loss to values lower than 1 wt.% if the batch-size is large enough.

EDB SUBJECT CATEGORY: EDB-360201;050700

INIS CATEGORY: B23

<692>

Weeks, R.W. ** Hill, M.R. (comp.), Oak Ridge National Lab., Tenn. (USA)

Argonne National Laboratory In Mechanical properties test data for structural materials. Quarterly progress report for period ending April 30, 1977 ** ERA-03:004353;EDB-78:004435; (Jun 1977) CRNL--5287

SUBJECT DESCRIPTORS: CORROSION;CORROSIVE EFFECTS;DEFORMATION;FATIGUE: Q1,Q2;FRACTURE PROPERTIES: Q1,Q2;INCOLOY 800: T3;LMFBR TYPE REACTORS;REACTOR MATERIALS;SODIUM;STAINLESS STEEL-16-8-2: T2;STAINLESS STEEL-304: T1;STAINLESS STEEL-316: T4;TENSILE PROPERTIES: Q3,Q4

ABSTRACT: Results obtained on the fatigue behavior of Type 16-8-2 stainless steel weld metal, additional analysis of notch effects on the cyclic life of Type 304 stainless steel, and results from an experimental determination of bilinear stress-strain parameters for Incoloy 800 are described. An analysis of true stress-true strain tensile data obtained on titanium-modified Type 316 stainless steel exposed to flowing sodium is presented, and additional results on the effects of sodium environment on the fatigue behavior of Type 304 stainless steel are reported.

EDB SUBJECT CATEGORY: EDB-360103;210500

<693>

Wenzl, H.; Sorajic, V.; Bischof, B., Kernforschungsanlage Juelich G.m.b.H. (Germany, F.R.). Inst. fuer Festkoerperforschung

In Investigation of the solubility and diffusion of Fe atoms in Cu at high temperature using molten salt electrochemistry ** AIX-08:334389;EDB-78:016794; (Jan 1977) Juel--1384

SUBJECT DESCRIPTORS: COPPER ALLOYS: T1;DIFFUSION: Q2;ELECTROCHEMISTRY;HIGH TEMPERATURE;IRON: T2;IRON ALLOYS;MOLTEN SALTS;PERMEABILITY: Q1;SOLUBILITY;VERY HIGH TEMPERATURE

ABSTRACT: The electrochemical cell $\text{CuFe}_{\text{sub}}(n)/\text{KF}, \text{LiF}, \text{FeF}_{\text{sub}}$

<693>

<693> CONT.

2% (molten solution)/Fe was used between 800 and 1,000°C to produce CuFe_n alloys of various copper rich compositions n by electrochemically controlled diffusion. From measurements of cell voltage and current we determined composition, bulk diffusion coefficient D, and atomic solubility limit x₀ of Fe in Cu. The numerical values at the temperature of 950°C are $D = 0.9 \times 10^{-9} \text{ cm}^2/\text{sec}$, $x_0 = 1.2 \text{ at\%}$.

EED SUBJECT CATEGORY: EDB-360104

INIS CATEGORY: B12

<694>

Wichmann, U.; Berndt, U.; Keller, C. (Kernforschungszentrum Karlsruhe (Germany, F.R.). Inst. fuer Radiochemie)

Phase equilibria and thermodynamic data for the ternary UO₂-UO₃-REO_{1.5} (Ho, Tm, Lu) systems

AIX-08:346212;EDB-78:027694; Rev. Chim. Miner. (RVCMA)

14:(2):105-118 (1977)

ABSTRACT: Investigations of the phase equilibria in the oxygen-rich part of the ternary systems U-RE-O with the heavy lanthanides holmium, thulium and lutetium by using X-ray and EMF methods have shown that the following single phase regions exist in all systems: a fluorite phase (U_{1-y}RE_y)(U_{2+x}), a rhombohedral phase and a C-type phase. The phase width in the three systems for the different single phases differs, however, especially for the rhombohedral phase. No solubility of REO_{1.5} in U₃₋₈ could be detected whereas REO_{1.5} dissolves small quantities of UO_{2+x} the amount being the higher, the smaller the value of x is. The stoichiometric members of the fluorite phase, (RE_{0.5}U_{0.5})O₂, are thermodynamically stable only at high temperatures for RE=Ho, Tm, Lu. A detailed description of the phase diagrams of these systems is given together with thermodynamic data for the free solution enthalpy of oxygen in selected compounds of these systems, obtained by using a specially developed high-temperature galvanic cell.

EED SUBJECT CATEGORY: EDB-360202;400702

INIS CATEGORY: B12

<695>

Williams, J.M. (Argonne National Lab., IL); Dowling, N.;
Gunde, R.; Hadzi, D.; Orel, B.

X-ray diffraction and infrared: Raman spectroscopic study of
the hydrogen dinitrate ion, $(\text{O} \text{sub } 2 \text{NO-H-ONO} \text{sub } 2) \text{sup -}$,
in cesium hydrogen dinitrate

ERA-01:016813; INS-76:014750; EDB-76:049146; J. Am. Chem. Soc.
(JACSA) 98:(6):1581-1582 (17 Mar 1976)

SUBJECT DESCRIPTORS: BOND ANGLE; BOND LENGTHS; CESIUM NITRATES:
T3; CRYSTAL GROWTH; DEUTERIUM: T4; HYDROGEN COMPOUNDS:
T1; INFRARED SPECTRA; ISOTOPE EFFECTS: Q4; PATTERSON
METHOD; RAMAN SPECTRA; STRUCTURAL CHEMICAL ANALYSIS:
Q1, Q3; X-RAY DIFFRACTION

ABSTRACT: A structural chemical analysis of $\text{CsH}(\text{NO} \text{sub } 3) \text{sub } 2$
was performed using the Patterson map and the least
squares technique. The Cs ion coordination sphere was found
to consist of ten oxygen atoms from surrounding nitrate
groups; and the $\text{H}(\text{NO} \text{sub } 3) \text{sub } 2 \text{sup -}$ ion was found to
have four oxygen atoms from two nitrate groups arranged in
tetrahedral fashion. Bond angles and lengths were reported,
and deuterium isotope effects were discussed. (DDA)

EDE SUBJECT CATEGORY: EDB-400201; 400202
INIS CATEGORY: B12

<696>

Wit, H.P.; Niesen, L. (Rijksuniversiteit Groningen (Netherlands))

Accurate nuclear parameters derived from the Moessbauer spectrum
of thulium metal AIX-07:276505; EDB-77:026148; Hyperfine
Interact. (HYIND) 1:(6):501-504 (Aug 1976)

SUBJECT DESCRIPTORS: ALUMINIUM; ERBIUM 169: T2; ERBIUM
IONS; HYPERFINE STRUCTURE: Q1; ION IMPLANTATION; MOESSBAUER
EFFECT: Q2; SPECTRA; THULIUM: T1; ULTRALOW TEMPERATURE

ABSTRACT: Accurate values of hyperfine interaction parameters
have been obtained for thulium metal from Moessbauer spectra,
using an implanted source of $\text{sup } 169 \text{Er}$ in Al and Moire
velocity calibration. A magnetic moment ratio
 $\mu^*/\mu = -2.223(13)$ has been measured along with a ground
state magnetic dipole coupling strength $\mu^*H = 33.26(9)$ cm/sec
and an excited state quadrupole coupling strength $e \text{sup } 2qQ^*/h = 1933(12)$ MHz.

EDB SUBJECT CATEGORY: EDB-360104
INIS CATEGORY: A13

<697>

<697>

Woolfrey, J.L. (Australian Atomic Energy Commission Research Establishment, Lucas Heights. Materials Div.); Wright, S.J. (New South Wales Univ., Kensington (Australia). Dept. of Ceramic Engineering)

Measurement of interfacial energies in the UO_2 - Gd_2O_3 system AIX-08:327849; EDB-77:140670; J. Aust. Ceram. Soc. (JAUCA) 12:(2):21-26 (Nov 1976)

SUBJECT DESCRIPTORS: GADOLINIUM OXIDES: T2; GRAIN BOUNDARIES: Q1, Q2; INTERFACES; PHASE STUDIES; POROSITY; SURFACE TENSION: Q1, Q2; SURFACES; THERMAL EQUILIBRIUM; URANIUM DIOXIDE: T1

ABSTRACT: The multiphase equilibration method is being used to investigate the interfacial energies in the UO_2 - Gd_2O_3 system as part of a program to determine the effect of Gd_2O_3 additions on the surface energy of UO_2 . The methods used for determining surface and interfacial energies are reviewed, the experimental techniques used in the present investigation are described and some results are presented. Specimens of UO_2 -5 weight per cent Gd_2O_3 were equilibrated at 1700°C for two weeks in hydrogen. High purity nickel (99.99%) was used to investigate the solid-liquid equilibrium. The results indicated that the presence of gadolinia reduced the ratios of grain boundary to solid surface energy and grain boundary to solid-liquid interfacial energy for UO_2 by approximately 13 and 47 per cent respectively. It also increased the grain boundary-pore 'dihedral' angle of UO_2 from 90.6 to 100.1 degrees.

EDB SUBJECT CATEGORY: EDB-360202
INIS CATEGORY: B23

<698>

Wurzinger, H.; Mueller, K. (Technische Hochschule, Graz (Austria))

Determination of trace impurities of titanium and zirconium in molybdenum and tungsten and their oxygen compounds by X-ray fluorescence spectrometry AIX-08:331932; EDB-78:022212; Z. Anal. Chem. (ZACFA) 284:(2):101-105 (Apr 1977)

ABSTRACT: A method for the determination of trace impurities of titanium and zirconium in metallic molybdenum and tungsten or in their basic products MoO_3 , WO_3 and H_2WO_4 is presented. The method is based on coprecipitation of the trace impurities titanium and zirconium with cobalt hydroxide as carrier from molybdenum and tungsten matrices. The hydroxide precipitates are collected on filter paper and measured by XRF after drying.

<698> CONT.

That way it is possible to get linear calibration curves up to $1 \times 10^{\text{sup } -4} \text{ g}$; the limits of detection are determined to be $3.4 \times 10^{\text{sup } -8} \text{ g}$ for titanium and $5.0 \times 10^{\text{sup } -7} \text{ g}$ for zirconium. About 45 min are necessary for one analysis, but several samples can be analyzed simultaneously.

EDB SUBJECT CATEGORY: EDB-400103

INIS CATEGORY: B11

<699>

Yagi, E.; Maeda, M.; Miyajima, K.; Kagami, H., Japan Atomic Energy Research Inst., Tokyo

In Technical experiences in an engineering-scale facility for uranium fluorination studies **

AIX-C8:300964; ERA-02:041098; EDB-77:092370; (Mar 1976)

JAERI-M--6488

SUBJECT DESCRIPTORS: FLOWSHEETS; FLUIDIZED BED; FLUORIDE VOLATILITY PROCESS; FLUORINATION: Q1; FLUORINE; FUEL CYCLE; OFF-GAS SYSTEMS; OPERATION; PERFORMANCE; PILOT PLANTS; POWER REACTORS; URANIUM: T1; URANIUM DIOXIDE; URANIUM HEXAFLUORIDE

ABSTRACT: In the research program of power reactor fuel reprocessing by fluoride volatility process, an engineering-scale facility for uranium fluorination has been operated batchwise and semi-continuously since 1969 in Fuel Reprocessing Laboratory, JAERI, to establish the techniques of handling kilogram quantities of fluorine and uranium hexafluoride and to obtain engineering data on the uranium fluidized-bed fluorination. The experiences gained with the facility are described, giving some evaluations and recommendations for the equipment design, its performance and operation of the system.

EDB SUBJECT CATEGORY: EDB-050800

INIS CATEGORY: B16

<700>

Yagi, E.; Saito, S.; Horiuchi, M., Japan Atomic Energy Research Inst., Tokyo

In Description of an engineering-scale facility for uranium fluorination studies **

AIX-C8:300965; ERA-02:041097; EDB-77:092369; (Mar 1976)

JAERI-M--6487

SUBJECT DESCRIPTORS: DIAGRAMS; FLUIDIZED BED; FLUORIDE VOLATILITY PROCESS; FLUORINATION: Q2; FLUORINE; FUEL CYCLE; MATERIALS

<700>

<700> CONT.

HANDLING; OFF-GAS SYSTEMS; OXIDATION; PILOT PLANTS; POWER
 REACTORS; SAFETY; URANIUM: T2; URANIUM DIOXIDE; URANIUM
 HEXAFLUORIDE

ABSTRACT: In the research program of power reactor fuel reprocessing by fluoride volatility process, the engineering facility was constructed to establish the techniques of handling kilogram quantities of fluorine and uranium hexafluoride and to obtain engineering data on the uranium fluidized-bed oxidation and fluorination. This facility is designed for a capacity of 5 kg per batch. Descriptions on the facility and equipment are given, including design philosophy, safety and its analysis.

EDB SUBJECT CATEGORY: EDB-050800

INIS CATEGORY: B16

<701>

Yenco, R.M.; Maroni, V.A.

Solubility of lithium oxide in liquid lithium

INS-77:015238; ERA-02:057131; EDB-77:132589; Trans. Am. Nucl.
 Soc. (TANSA) 26::166-167 (Jun 1977)

SUBJECT DESCRIPTORS: HIGH TEMPERATURE; LIQUID METALS; LITHIUM:
 M1; LITHIUM OXIDES: M2; SOLUBILITY: Q2; SOLVENT PROPERTIES: Q1

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-360104

INIS CATEGORY: P22

<702>

Yonco, R.M.; Veleckis, E.; Maroni, V.A. (Argonne National Lab.,
 Ill. (USA))

Solubility of nitrogen in liquid lithium and thermal

decomposition of solid Li_3N EDB-76:057072; J. Nucl.
 Mater. (JNUMA) 57:(3):317-324 (Sep 1975)

SUBJECT DESCRIPTORS: FORMATION FREE ENERGY; HIGH
 TEMPERATURE; LIQUID METALS; LITHIUM: T1; LITHIUM NITRIDES:
 T3; MELTING POINTS; NITROGEN: T2; PYROLYSIS: Q3; SOLUBILITY:
 Q2; SOLVENT PROPERTIES: Q1; TEMPERATURE
 DEPENDENCE; THERMONUCLEAR REACTOR MATERIALS; VERY HIGH
 TEMPERATURE

ABSTRACT: The solubility of nitrogen in liquid lithium was
 determined from 195 to 441°C by direct sampling and
 equilibrium nitrogen pressure over solid Li_3N were

<702> CONT.

measured at eight temperatures between 660 and 778⁰C. The solubility data may be represented by $\log_{10} S = 3.323 - 2107/T^{\circ}K$, where S is in mol%Li₃N and T is in K. From a thermodynamic analysis of the combined solubility and decomposition data, the standard free energy of formation of solid Li₃N was estimated to be $\Delta G^{\circ} = 33.2 \times 10^3 - 3T$ (kcal/mol). For dilute solutions of Li₃N in lithium, the Sieverts' law constant was determined. The system Li₃N appears to conform to a simple eutectic diagram in which the eutectic point occurs at approximately 0.05 mol%Li₃N and 180.3⁰C. The melting point of Li₃N was found to be (813 $\pm$ 1)⁰C. Implications of the results of this study regarding the compatibility of liquid lithium with the structural materials of interest to fusion reactors are discussed.

EDE SUBJECT CATEGORY: EDB-700209;360603
INIS CATEGORY: B12

<703>

Yoshida, S.; Kuroda, T. (Tokyo Inst. of Tech. (Japan). Research Lab. of Nuclear Reactor); Oyamada, Ryoza

Preparation of thorium metal by electrolysis of fused salt ThCl₄-LiCl+NaCl+KCl(eut) system
AIX-07:262603;EDB-77:018565; Bull. Tokyo Inst. Technol. (Engl. Ed.) (BTITA) (127):105-112 (Mar 1975)

SUBJECT DESCRIPTORS: CHEMICAL PREPARATION;ELECTRIC CONDUCTIVITY;ELECTROCHEMISTRY;ELECTROLYSIS;ELECTROMETALLURGY; ELECTROMOTIVE FORCE;ELECTROREFINING: Q1;LITHIUM CHLORIDES;MOLTEN SALTS;NUCLEAR FUELS;PHASE DIAGRAMS;POTASSIUM CHLORIDES;SODIUM CHLORIDES;TEMPERATURE DEPENDENCE;THORIUM: T1;THORIUM CHLORIDES

ABSTRACT: Although uranium is mainly used as nuclear fuel, thorium attracts attention because it is more abundant than uranium. At present, thorium fuel is mainly in the form of ceramics for HWRs and HTGRs, and fluoride for MSFRs. In the future, thorium will be used as an alloying element. The electrolytic production of metallic thorium is most promising for industrial purpose, hence, the authors chose thorium chloride as a component of electrolyte, because they had succeeded to get high purity recrystalline ThCl₄ free from hygroscopic property. Basic data concerning the electrolytic production of high purity metallic thorium were already reported by the authors. The present study is concerned with the conditions of the electrolysis of fused salt ThCl₄-LiCl+NaCl+KCl (eut) system. ThCl₄ and the eutectic mixture of LiCl+NaCl+KCl were prepared, and their fused salt mixture was made. The measurements of

<703>

<703> CONT.

liquidus, specific electric conductance, and electromotive force were carried out, then the electrolysis of the fused salt was made. The optimum conditions for the electrolysis of the fused salt were estimated from the basic measurements as follows: ThCl_4 concentration in electrolyte 5--60 wt.%, specific conductance of the electrolyte containing 10--30 wt. % ThCl_4 at 550--650°C is not inferior to other systems, lower e.m.f. than other systems is advantageous. Therefore, ThCl_4 concentration 30 wt.%, electrolysis temperature 550--600°C, and initial cathode current density as high as possible, were found as most optimum conditions.

EDB SUBJECT CATEGORY: EDB-360101
INIS CATEGORY: B12

<704>

Yoshizawa, S.; Takehara, Z.; Ito, Y.; Morimoto, S.

Lithium alloys--chlorine secondary battery. I. Basic studies on a lithium alloys--chlorine secondary battery using a molten salt as the electrolyte EDB-77:112750; Denki Kagaku (DNKKA) 39:(10):834-839 (1971)

SUBJECT DESCRIPTORS: AUTOMOBILES:

T2; CARBON; ELECTROLYTES; ELECTROMOTIVE FORCE; ENERGY; FEASIBILITY STUDIES: Q1; HIGH TEMPERATURE; LEAD ALLOYS; LITHIUM ALLOYS; LITHIUM CHLORIDES; LITHIUM-CHLORINE BATTERIES: M1, Q2; MOLTEN SALTS; POTASSIUM CHLORIDES; TIN ALLOYS; ZINC ALLOYS

ABSTRACT: The feasibility as an automobile battery of a $\text{Li}(\text{Zn}, \text{Sn}, \text{or Pb})/\text{LiCl--KCl/C--Cl}_2$ battery is examined. The emf was somewhat less than that of a conventional Li/Cl_2 battery. Operating temperature was 450°C, and the output energy density of the anode was about 130 Wh/kg. (RWR)

EDB SUBJECT CATEGORY: EDB-250901

<705>

Young, W.H., East Anglia Univ. (UK). School of Mathematics and Physics

In Electronic states densities from structure and thermodynamic data for liquid transition metals **
AIX-07:271683; EDB-77:055413; (1976) AED-Conf--76-087-015

SUBJECT DESCRIPTORS: COMPARATIVE EVALUATIONS; DATA; ELECTRONIC STRUCTURE: Q1; ELECTRONS; ENTROPY; LIQUID METALS; SOLIDS; THERMODYNAMIC PROPERTIES: Q1; THERMODYNAMICS; TRANSITION ELEMENTS: T1; VALENCE

<705> CONT.

ABSTRACT: The entropies of many liquid metals can be written in the form $S = S_{\text{gas}} + S_{\text{pack}} + S_{\text{elec}}$, where S_{gas} is the ideal gas expression, S_{pack} arises from the finite sizes of the ions (described by effective hard spheres) and S_{elec} is the valence-electron contribution $\frac{1}{3} \pi^{1/2} N(E_F) k_B T$. The shape of the observed structure factor tells us whether the above formalism is applicable in any given case; if it is, we thereby deduce the effective hard sphere diameter and thence S_{pack} . Measured thermodynamic data allow us to obtain S and S_{gas} and so S_{elec} and thus $N(E_F)$ may be inferred in principle. In practice, for non-transition metals S_{elec} is typically 1% of S , and so it is difficult to extract meaningful results from the experimental data. However, for transition metals (for which $N(E_F)$ and the melting temperature are both much higher) S_{elec} is typically 20% of S and useful information on $N(E_F)$ may be obtained. Such data are presented and related to the additional relevant experimental and theoretical information available to us.

EEB SUBJECT CATEGORY: EDB-360104

INIS CATEGORY: A13

<706>

Zakharov, D.M.

Eutectic point of gallium-indium system

AIX-08:321120;EDB-77:113371; Zh. Fiz. Khim. (ZFKHA)

49:(9):2208-2210 (Sep 1975)

SUBJECT DESCRIPTORS: BINARY ALLOY SYSTEMS;COMPARATIVE EVALUATIONS;EUTECTICS;GALLIUM ALLOYS: T1;INDIUM ALLOYS: T2;LIQUID METALS;MELTING;PHASE DIAGRAMS: Q1,Q2;THERMODYNAMICS

ABSTRACT: On the base of thermodynamic analysis of the gallium-indium alloy system its eutectic concentration has been determined: 13.8 ± 0.4 at.% of In. The computed value agrees with the experimental value determined by a chemical analysis of liquid indium-gallium alloy formed in the process of a contact melting of Ga and In at the eutectic temperature.

EEB SUBJECT CATEGORY: EDB-360102

INIS CATEGORY: B22

<707>

<707>

Zavalishin, N.I.; Mal'tsev, A.A. (Moskovskij Gosudarstvennyj Univ. (USSR). Kafedra Fizicheskoy Khimii)

IR spectra of the NbCl_5 , TaCl_5 , NbBr_5 and NbOCl_3 vapors AIX-08:283089; EDB-77:069073; Vestn. Mosk. Univ., Ser. II. Khim. (VMUKA) 17:(3):314-317 (May 1976)

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-400201

INIS CATEGORY: B12

<708>

Zhul'kov, V.I. (Vsesoyuznyj Nauchno-Issledovatel'skij Inst. Sinteticheskogo Kauchuka, Leningrad (USSR))

Investigation into properties of liquidus curves of binary salt system melts where chemical compounds formation. 2. Systems of mono-monovalent and bi-bivalent cations with common anion AIX-08:340003; EDB-78:017351; Zh. Fiz. Khim. (ZFKHA) 51:(1):11-15 (Jan 1977)

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-400201

INIS CATEGORY: B12

<709>

Zimina, T.D.; Zhukova, L.D. (Irkutskij Gosudarstvennyj Univ. (USSR))

Cesium, calcium and barium chlorides ternary system AIX-08:286228; EDB-77:044209; Izv. Vyssh. Uchebn. Zaved., Khim. Khim. Tekhnol. (IVUKA) 18:(8):1307-1309 (1975)

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-400201

INIS CATEGORY: B12

<710>

In High density thoria-silica-metal (III) oxide fibers ** AIX-07:233183; EDB-76:060445; (18 Dec 1974)

SUBJECT DESCRIPTORS: ALUMINIUM OXIDES: T1; BAKING; BORON OXIDES: T4; CHEMICAL COMPOSITION; CHEMICAL PREPARATION: Q1, Q2, Q3, Q4; DENSITY; DISPERSIONS; EXTRUSION; FIBERS; FLEXIBILITY; PROTECTIVE CLOTHING; SHIELDING MATERIALS; SILICON OXIDES: T2; THORIUM OXIDES: T3; VISCOSITY

<710> CONT.

ABSTRACT: Transparent refractory fibers, at least 50% thoria and additionally containing silica and metal(III) oxides, particularly Al_2O_3 and B_2O_3 or Cr_2O_3 are made by shaping and dehydratively gelling, particularly by extruding in air, viscous aqueous thoria solutions or sols containing colloidal silica with boric acid-stabilized aluminum acetate, or additionally chromium acetate or colloidal Cr_2O_3 , and heating the resulting gelled fibers in a controlled manner to decompose and volatilize undesired constituents and convert fibers to refractory fibers which are useful to form, for example, refractory fabrics, or as reinforcement for composites. The fabrics are heat resistant. A special application is X-ray protective clothing.

EEB SUBJECT CATEGORY: EDB-360204

INIS CATEGORY: B24

<711>

Standard methods for chemical, mass spectrometric, and spectrochemical analysis of nuclear-grade uranium dioxide powders and pellets INS-76:013238;EDB-76:042495; Annu. Book ASTM Stand. (ASTSA) (45):175-213 (1975)

SUBJECT DESCRIPTORS: ALUMINIUM: T11;ANTIMONY: T31;ARSENIC: T12;BARIUM: T14;BERYLLIUM: T15;BISMUTH: T16;BORON: T13;CADMIUM: T18;CALCIUM: T17;CARBON: T4;CHEMICAL ANALYSIS: Q1;CHLORINE: T5;CHROMIUM: T20;COBALT: T19;COPPER: T21;EMISSION SPECTROSCOPY: Q8,Q9,Q11,Q12,Q13,Q14,Q15,Q16,Q17,Q18,Q19,Q20,Q21,Q22,Q23,Q24,Q25,Q26,Q27,Q28,Q29,Q30,Q31,Q32,Q33,Q34,Q35,Q36,Q37,Q40,Q41,Q2;FLUORINE: T6;FUEL PELLETS;GRAVIMETRIC ANALYSIS: Q2,Q3;INDIUM: T23;ION SELECTIVE ELECTRODE ANALYSIS: Q5,Q6;IRON: T22;LEAD: T30;MAGNESIUM: T24;MANGANESE: T25;MASS SPECTROSCOPY: Q38;MERCURY: T40;MOISTURE;MOLYBDENUM: T26;NICKEL: T28;NITROGEN: T7;OXYGEN: T3;PHOSPHORUS: T29;POTASSIUM: T42;POWDERS;QUANTITATIVE CHEMICAL ANALYSIS: Q4;RARE EARTHS: T9;SILICON: T32;SILVER: T8;SODIUM: T27;SPECTROPHOTOMETRY: Q10;SULFUR: T10;THORIUM: T41;TIN: T33;TITANIUM: T34;TITRATION: Q7,Q2;TRACE AMOUNTS;URANIUM: T2;URANIUM DIOXIDE: T1;URANIUM ISOTOPES: T38;VANADIUM: T35;VOLTAMETRY;ZINC: T36;ZIRCONIUM: T37

ABSTRACT: Chemical methods are described for the determination of U by FeSO_4 reduction in H_3PO_4 and titration with $\text{Cr}_2\text{O}_7^{2-}$, atomic ratio by ignition (gravimetric) impurity correction method, total C by direct combustion-thermal conductivity method, N by the Kjeldahl method, surface area of powders by the N-absorption method, and total gas (other than H_2O) in pellets by a vacuum outgassing and heat release method. Electrochemical

<711>

<711> CONT.

methods for the determination of moisture, Cl^{sup} -, F^{sup} -, and O:U atomic ratio are described. The isotopic U content is determined by a mass spectrometric method, and S is determined by a distillation spectrophotometric method. Various spectrochemical methods are included for the following trace elements present in the UO_2 volatile F^{sup} -, Ag, rare earths, and 26 other elements whose concentration ranges and recommended analytical spectral lines are tabulated. A spark-source mass spectrographic method is described for the determination of impurities in the atom-ppm range. All methods are described in detail including procedure, detection limits, and interferences when applicable. (BLM)

EDB SUBJECT CATEGORY: EDB-400102;400104;050700

INIS CATEGORY: B11

<712>

Standard methods for chemical, mass spectrometric, and spectrochemical analysis of nuclear-grade plutonium dioxide powders and pellets INS-76:013239;EDB-76:042496; Annu. Book ASTM Stand. (ASTSA) (45):214-250 (1975)

SUBJECT DESCRIPTORS: ALUMINIUM: T8;AMPEROMETRY: Q45;ANTIMONY: T32;ARSENIC: T10;BARIUM: T13;BERYLLIUM: T14;BISMUTH: T15;BORON: T12;CADMIUM: T17;CALCIUM: T16;CARBON: T3;CHEMICAL ANALYSIS: Q1;CHLORINE: T5;CHROMIUM: T20;COBALT: T18;COFFER: T19;EMISSION SPECTROSCOPY: Q7,Q8,Q9,Q10,Q11,Q12,Q13,Q14,Q15,Q16,Q17,Q18,Q19,Q20,Q21,Q22,Q23,Q24,Q25,Q26,Q27,Q28,Q29,Q30,Q31,Q32,Q33,Q34,Q35,Q36,Q37,Q3,Q39,Q40,Q41,Q42;FLUORINE: T4;FUEL PELLETS;GALLIUM: T22;GOLD: T11;INDIUM: T23;ION SELECTIVE ELECTRODE ANALYSIS: Q4,Q5;IRON: T21;LEAD: T30;MAGNESIUM: T25;MANGANESE: T26;MASS SPECTROSCOPY: Q44;MERCURY: T41;MOISTURE;MOLYBDENUM: T27;NICKEL: T28;NITROGEN: T2;PALLADIUM: T31;PHOSPHORUS: T29;PLUTONIUM: T45;PLUTONIUM DIOXIDE: T1;PLUTONIUM ISOTOPES: T44;POTASSIUM: T24;POWDERS;QUANTITATIVE CHEMICAL ANALYSIS: Q3;RARE EARTHS: T7;SILICON: T33;SILVER: T9;SPECTROPHOTOMETRY: Q2,Q4,Q5,Q6;STRONTIUM: T35;SULFUR: T6;THORIUM: T42;TIN: T34;TITANIUM: T36;TITRATION: Q4,Q45;TRACE AMOUNTS;TUNGSTEN: T38;VANADIUM: T37;VOLTAMETRY: Q45;ZINC: T39;ZIRCONIUM: T40

ABSTRACT: Various methods are described in detail as to procedure, interferences, and limits of detection or applicability for the determination of Pu and various impurities in PuO_2 powders and pellets. Methods for the determination of Pu are controlled-potential coulometry, ceric sulfate titration, and amperometric titration with $\text{Fe}^{\text{sup}} 2+$. Spectrophotometric methods are given for N and S, and a direct combustion-thermal conductivity method for the determination of total C is described. A pyrohydrolysis

<712> CONT.

method is described for the determination of μg amounts of $\text{Cl}^{\text{sup}}-\$$ and $\text{F}^{\text{sup}}-\$$. A mass spectrometric determination of the Pu isotopic content of the oxide is included, and general chemical methods are included for the determination of moisture in both powders and pellets and for total gas--exclusive of moisture--in pellets. Spectrochemical methods are described for the determination of rare earths, 36 trace elements whose recommended analytical spectral lines and concentration ranges are tabulated, and impurities occurring in the atom-ppB range. (BLM)

EDB SUBJECT CATEGORY: EDB-400102;400104;050700
INIS CATEGORY: B11

<713>

Standard methods for chemical, mass spectrometric, and spectrochemical analysis of nuclear-grade mixed oxides (U,Pu) $\text{O}_{\text{sub } 2}$ INS-76:013240;EDB-76:042497; Annu. Book ASTM Stand. (ASTSA) (45):251-292 (1975)

SUBJECT DESCRIPTORS: ALUMINIUM: T16;AMPEROMETRY: Q4;BARIUM: T18;BERYLLIUM: T19;BISMUTH: T20;BORON: T17;CADMIUM: T22;CALCIUM: T21;CARBON: T8;CHEMICAL ANALYSIS: Q1,Q2;CHLORINE: T9;CHROMIUM: T24;COBALT: T23;COPPER: T25;EMISSION SPECTROSCOPY: Q14,Q15,Q16,Q17,Q18,Q19,Q20,Q21,Q22,Q23,Q24,Q25,Q26,Q27,Q28,Q29,Q30,Q31,Q32,Q33,Q34,Q35,Q36,Q37,Q38,Q39,Q40,Q42,Q43;FLUORINE: T10;FUEL PELLETS;ION SELECTIVE ELECTRODE ANALYSIS: Q9,Q10;IRON: T26;LEAD: T33;MAGNESIUM: T27;MANGANESE: T28;MASS SPECTROSCOPY: Q5,Q6;MERCURY: T42;MOLYBDENUM: T29;NICKEL: T31;NITROGEN: T11;PHOSPHORUS: T32;PLUTONIUM: T4;PLUTONIUM DIOXIDE: T2;PLUTONIUM ISOTOPES: T6;POTASSIUM: T40;POWDERS;QUANTITATIVE CHEMICAL ANALYSIS: Q8;RARE EARTHS: T14;SILICON: T34;SILVER: T15;SODIUM: T30;SPECTROPHOTOMETRY: Q9,Q10,Q11,Q12,Q13;SULFUR: T12;THORIUM: T43;TIN: T35;TITANIUM: T36;TITRATION: Q4,Q9;TRACE AMOUNTS;TUNGSTEN: T13;URANIUM: T3;URANIUM DIOXIDE: T1;URANIUM ISOTOPES: T5;VANADIUM: T37;VOLTAMETRY: Q3,Q4;ZINC: T38;ZIRCONIUM: T39

ABSTRACT: Methods are described for the determination of both Pu and U in mg quantities and for trace impurities in mixed oxides (U,Pu) $\text{O}_{\text{sub } 2}$. Controlled-potential coulometry methods are described for both U and Pu, and an amperometric titration method using $\text{Fe}^{\text{sup } 2+}$ is included for Pu. The isotopic content of the oxide is determined by a mass spectrometric method. Methods are described for the determination of moisture and of total gas--other than moisture--in mixed oxide pellets. Direct combustion-thermal conductivity and pyrohydrolysis methods are included for the determination of C and $\text{Cl}^{\text{sup}}-\$$ and $\text{F}^{\text{sup}}-\$$, respectively. Spectrophotometric methods are described for the

<713>

<713> CONT.

determination of N, S, and W. Trace amounts of rare earths and 25 other elements whose recommended analytical spectral lines and concentration ranges are tabulated are determined by spectroscopic methods. (BLM)

EDB SUBJECT CATEGORY: EDB-400102;400104;050700

INIS CATEGORY: B11

<714>

Standard methods for chemical, mass spectrometric, spectrochemical, nuclear, and radiochemical analysis of nuclear-grade plutonium metal INS-76:013241;EDB-76:042498; Annu. Book ASTM Stand. (ASTSA) (45):356-366 (1975)

SUBJECT DESCRIPTORS: ALUMINIUM: T16;AMERICIUM 241: T10;ANTIMONY: T39;ARSENIC: T18;BARIUM: T21;BERYLLIUM: T22;BISMUTH: T23;BORON: T20;CADMIUM: T25;CALCIUM: T24;CARBON: T2;CHEMICAL ANALYSIS: Q1;CHLORINE: T6;CHROMIUM: T28;COBALT: T26;COPPER: T27;EMISSION SPECTROSCOPY: Q5,Q11,Q12,Q13,Q14,Q15,Q16,Q17,Q18,Q19,Q20,Q21,Q22,Q23,Q24,Q26,Q27,Q28,Q30,Q31,Q32,Q33,Q34,Q35,Q36,Q37,Q38,Q39,Q40,Q41,Q42,Q43,Q44,Q45,Q46,Q47;FLUORINE: T7;GALLIUM: T30;GAMMA SPECTROSCOPY: Q3,Q4;GOLD: T19;INDIUM: T31;IRON: T5;LEAD: T38;LITHIUM: T15;MAGNESIUM: T33;MANGANESE: T34;MERCURY: T47;MOLYBDENUM: T35;NICKEL: T36;NIOBIUM: T13;NITROGEN: T8;PHOSPHORUS: T37;PLUTONIUM: T1;POTASSIUM: T32;RADIOMETRIC ANALYSIS: Q10;RARE EARTHS: T11;SILICON: T40;SILVER: T17;SODIUM: T14;SPECTROPHOTOMETRY: Q2,Q3,Q4,Q5,Q6,Q7,Q8,Q9;STRONTIUM: T42;SULFUR: T9;THORIUM: T4;TIN: T41;TITANIUM: T43;TRACE AMOUNTS;TUNGSTEN: T12;URANIUM: T3;VANADIUM: T44;ZINC: T45;ZIRCONIUM: T46

ABSTRACT: Methods are described for the analysis of matrix materials and various impurities in Pu metal. Controlled-potential coulometry, amperometric titration with $\text{Fe}^{\text{sup } 2+}$, or titration with ceric sulfate methods are described for the determination of Pu. A mass spectrometric method is given for determination of the isotopic content of the metal. Determination of C by direct combustion-thermal conductivity is described. Methods for the spectrophotometric determination of U with arsenazo I, Th with thorin, Fe with 1,10-phenanthroline or α , α' -dipyridyl, and $\text{Cl}^{\text{sup } -}$, $\text{F}^{\text{sup } -}$, N, and S following separation by distillation are described. If the Pu metal does not contain significant amounts of other radioactive fission products or high specific-activity γ emitters, ^{241}Am may be determined by γ counting. Determination of γ -emitting fission products, U and Th, is determined by γ spectroscopy. Spectrochemical methods for the

<714> CONT.

determination of rare earths, W, Nb, Ta, and about 40 other trace elements are included. (BLM)

EDB SUBJECT CATEGORY: EDB-400102;400103;400104;050700
INIS CATEGORY: B11

<715>

Standard method of test for atom percent fission in uranium and plutonium fuel (mass spectrometric method)

INS-76:014736;EDB-76:049037; Annu. Book ASTM Stand. (ASTSA) (45):739-744 (1975)

SUBJECT DESCRIPTORS: BURNUP: Q3;FUEL ELEMENTS: T3;MASS SPECTROSCOPY: Q4,Q5,Q6,Q7,Q8,Q9,Q10;PLUTONIUM;PLUTONIUM 239: T6;PLUTONIUM 240: T7;PLUTONIUM 241: T8;PLUTONIUM 242: T9;URANIUM;URANIUM 235: T4;URANIUM 236: T5;URANIUM 238: T10

ABSTRACT: A method is described for the determination of the heavy element atom percent fission in irradiated U fuel with initial Pu content of from 0 to 50 percent from the isotopic content of the fuel before and after irradiation. The method is restricted to samples where accurate preirradiation isotopic analysis can be obtained, and both analyses should be performed on the same instrument under similar conditions. Atomic ratios of the isotopes ^{235}U , ^{236}U , ^{239}Pu , ^{240}Pu , ^{241}Pu , and ^{242}Pu to ^{238}U are determined by mass spectrometry. The atom percent fission attributed to fission of ^{235}U , ^{238}U , and ^{241}Pu is separately calculated and then summed to obtain the total heavy element atom percent fission. The calculations for the determination are explained in detail. (BLM)

EDB SUBJECT CATEGORY: EDB-400104;C50800
INIS CATEGORY: B11;B16

<716>

Standard method of test for atom percent fission in uranium and plutonium fuel (neodymium-148 method)

INS-76:014738;EDB-76:049039; Annu. Book ASTM Stand. (ASTSA) (45):806-814 (1975)

SUBJECT DESCRIPTORS: BURNUP: Q4;CHEMICAL ANALYSIS;FISSION PRODUCTS;ION EXCHANGE;ISOTOPE DILUTION;MASS SPECTROSCOPY: Q3;NEODYMIUM 148: T3;NUCLEAR FUELS: T4;PLUTONIUM;URANIUM

ABSTRACT: Measurement of ^{148}Nd by mass spectroscopy in irradiated U fuel with initial Pu content of 0 to 50 percent is described as a method for determination of burnup. A blended spike of ^{150}Nd , ^{233}U , and ^{242}Pu is

<716>

<716> CONT.

added to the sample, and then the Nd, U, and Pu fractions are separated from each other by ion exchange. The decontaminated Nd is mounted on a Re filament for mass analysis. The atom percent fission is directly proportional to the $\frac{^{148}\text{Nd}}{\text{fuel}}$ ratio in the irradiated fuel. (BLM)

EDB SUBJECT CATEGORY: EDB-400104;050800

INIS CATEGORY: B11;B16

<717>

CODATA recommended key values for thermodynamics, 1976. Report of the CODATA Task Group on key values for thermodynamics, 1976 AIX-08:339840;EDB-78:017278; J. Chem. Thermodyn. (JCTDA) 9: (8):705-706 (Aug 1977)

SUBJECT DESCRIPTORS: BERYLLIUM: T1;BERYLLIUM OXIDES: T2;DATA;ENTHALPY: Q1,Q2,Q3,Q4,Q5;ENTROPY: Q1,Q2,Q3,Q4,Q5;FORMATION HEAT: Q1,Q2,Q3,Q4,Q5;INTERNATIONAL ORGANIZATIONS;RECOMMENDATIONS;THORIUM: T3;THORIUM OXIDES: T4;URANIUM: T5

ABSTRACT: None

EDB SUBJECT CATEGORY: EDB-400201

INIS CATEGORY: B12

<718>

Carnegie Institution of Washington, D.C. (USA). Dept. of Terrestrial Magnetism

In Optical spectroscopy of Np^{3+} in CaF_2 . Final report ** INS-77:011429;ERA-02:043798;EDB-77:101128; (1976) COO--3245-5

SUBJECT DESCRIPTORS: ABSORPTION SPECTRA: Q1;CALCIUM FLUORIDES: T1;DOPED MATERIALS;EMISSION SPECTRA: Q1;FLUORESCENCE;NEPTUNIUM 237: T

ABSTRACT: An analysis was made of the optical absorption and fluorescence spectra of green-colored Np^{3+} : CaF_2 specimens. The absorption spectra were attributed to Np^{3+} substituents at Ca^{2+} [000] sites with three distinct modes of charge compensation and correspondingly lowered local symmetries: (1) C_{3v} , associated with an F^{3+} compensator at a $[\text{F}^{3+}_{1/2}\text{F}^{3+}_{1/2}\text{F}^{3+}_{1/2}]$ -type site, (2) C_{3v} , associated with the replacement of a nn F^{3+} by an O^{2+} , (3) C_{4v} , associated with an F^{3+} compensator at a $[\text{O}^{2+}_{1/2}\text{F}^{3+}_{1/2}\text{F}^{3+}_{1/2}]$ -type site. A two-parameter fit to the positions of the multiplet centers yielded for these

<718> CONT.

sites, the following $F_{2\text{S}}$ and zeta values (in $\text{cm}^{\text{sup}}-1$): (1) 214.5 and 2032, (2) 212.7 and 2001, (3) 219.1 and 2083. "Crystal-field" parameters, derived from fits to the multiplet splittings, were calculated by molecular orbital techniques. Fluorescence, observed in the infrared, was assigned solely to the C_{4v} -symmetry site.

EDB SUBJECT CATEGORY: EDB-400201;400702

INIS CATEGORY: E12;B13

<719>

Combustion Engineering, Inc., Windsor, Conn. (USA)

In Fuel-clad interaction. C-E thermo-structural fuel evaluation methods **

INS-76:017364;ERA-01:023241;EDB-76:064254; (Apr 1976)

CENPD--179

SUBJECT DESCRIPTORS: CRACKS;FUEL CANS;FUEL PELLETS;FUEL-CLADDING

INTERACTIONS: M2,Q1;MATHEMATICAL MODELS: Q2;PWR TYPE

REACTORS: T1;STRAINS;STRESSES;URANIUM DIOXIDE;ZIRCALOY

ABSTRACT: Thermal and irradiation induced distortions experienced during the service life of a fuel rod lead to pellet-clad gap closure and local clad stresses and strains. In addition, presence of fission products can be detrimental to the ability of the clad to sustain such stresses and strains. A thermo-structural fuel evaluation model is described which has been developed to analyze the response of Zircaloy clad uranium dioxide fuel rods during pellet-clad interaction (PCI). The approach taken has been to utilize several analyses consisting of (1) an axisymmetric, cylindrical, pellet average model of history dependent behavior of the fuel rod and (2) pellet distortion and cracking models. Stresses and strains are to be compared with failure criteria to define conditions for which incipient fuel rod failure would exist. A description of the mathematical and behavioral models, failure criteria, and results for typical fuel rod geometries and power histories is presented.

EDB SUBJECT CATEGORY: EDB-210200

INIS CATEGORY: E32

<720>

<720>

Electrochimica Corp., Menlo Park, Calif. (USA)

In Research on a high-energy nonaqueous battery system.

Quarterly progress report No. 1, 1 Dec 1965--1 Mar 1966 **

EDB-76:047724; (Mar 1966) AD--484014

SUBJECT DESCRIPTORS: ELECTRODES: Q1; LITHIUM-COPPER CHLORIDE
BATTERIES: T1

ABSTRACT: None

EDE SUBJECT CATEGORY: EDB-250901;400201

<721>

Florida Univ., Gainesville (USA). Dept. of Chemistry

In Electrochemical methods for monitoring purity of molten
salts. Semiannual progress report, July--December 1975 **

INS-76:014764; ERA-01:020254; EDB-76:054434; (1975) ORO--4947-1

SUBJECT DESCRIPTORS: CHEMICAL ANALYSIS: Q1, Q2; ELECTROCHEMISTRY:
Q1, Q2; ELECTRODES; EUTECTICS; LITHIUM CHLORIDES: T1; MOLTEN
SALTS; PLATINUM; POTASSIUM CHLORIDES: T2; QUALITATIVE CHEMICAL
ANALYSIS; RESEARCH PROGRAMSABSTRACT: The tentative determination of visible impurities in
the LiCl-KCl eutectic salt as Fe, silica, and nitrides was
reported. Moisture was removed by bubbling HCl through the
melt. Of the three electrodes tested Pt had the least
residual current and was the most stable. Some etching of the
Pt electrode was observed in the preliminary stages of the
experiment. (JSR)

EDE SUBJECT CATEGORY: EDB-400400;360603

INIS CATEGORY: B24

<722>

General Atomic Co., San Diego, Calif. (USA)

In Thorium utilization program. Quarterly progress report for
the period ending May 31, 1977 **

ERA-03:006028; EDB-78:001097; (Jun 1977) GA-A--14441

SUBJECT DESCRIPTORS: ADSORPTION; CENTRIFUGATION; COATED FUEL
PARTICLES: T8; COMBUSTION PROPERTIES; CONTAINMENT
SYSTEMS; CONTROL ELEMENTS; CONVEYORS; CRUSHING: Q2; DESIGN:
Q7; DISSOLVERS; DUSTS; FLOWSHEETS: Q4; FLUIDIZED-BED COMBUSTION:
Q8, Q2; FLUIDIZED-BED COMBUSTORS: T9; FUEL CYCLE: Q1; FUEL
REPROCESSING PLANTS: T5; GAS FLOW; GRAPHITE; HEAD END
PROCESSES; HIGH TYPE REACTORS:

<722> CONT.

T1; IODINE; MAINTENANCE; MANAGEMENT; MATERIAL BALANCE; MATERIALS
HANDLING; NITROGEN OXIDES; NON-PROLIFERATION TREATY: T; OFF-GAS
SYSTEMS: T4; OPERATION; PERFORMANCE TESTING: Q9; PILOT PLANTS:
Q5; QUANTITY RATIO; RESEARCH PROGRAMS: Q3; RUTHENIUM
FLUORIDES; SEALS; SEPARATION PROCESSES; SILICON
CARBIDES; SOLIDS; SOLVENT EXTRACTION; SPENT FUEL CASKS: T7; SPENT
FUEL ELEMENTS: T2; SULFUR DIOXIDE; THOREX PROCESS:
T3; TRITIUM; URANYL NITRATES; VENTILATION; VOLATILITY

ABSTRACT: Results of work performed under the National HTGR Fuel
Recycle Program (also known as the Thorium Utilization
Program) at General Atomic Company are presented. Results of
work on this program prior to June 1974 were included in a
quarterly series on the HTGR Base Program. The work reported
includes the development of unit processes and equipment for
reprocessing of High-Temperature Gas-Cooled Reactor (HTGR)
fuel, the design and development of an integrated pilot line
to demonstrate the head end of HTGR reprocessing using
unirradiated fuel materials, and design work in support of
Hot Engineering Tests (HET). Work is also described on
trade-off studies concerning the required design of
facilities and equipment for the large-scale recycle of HTGR
fuels in order to guide the development activities for HTGR
fuel recycle.

EDB SUBJECT CATEGORY: EDB-C50800

<723>

Institute of Nuclear Physics, Krakow (Poland)

Nuclear spectroscopy. Laboratory of chemistry and radiochemistry
In Progress report 1971. Vol. 1 ** EDB-76:037142; (Nov
1972) INIS-mf--1946

SUBJECT DESCRIPTORS: CARRIER-FREE ISOTOPES:

T1; CHEMISTRY; COMPLEXES; ELEMENTS; EXTRACTION

CHROMATOGRAPHY; METALS; RADIOCHEMISTRY: T, Q1; RARE

EARTHS; SOLVENT EXTRACTION; SYNTHESIS; TRANSPLUTONIUM ELEMENTS

ABSTRACT: NONE

EDB SUBJECT CATEGORY: EDB-400702

INIS CATEGORY: B13

<724>

<724>

Japan Atomic Energy Research Inst., Tokyo

In Adaptation of the continuous cold trap system of
fluidized-bed to the fluoride volatility process **
AIX-08:300953;ERA-02:041078;EDB-77:092326; (Feb 1976)
JAERI-M--6405

SUBJECT DESCRIPTORS: COLD TRAPS:

Q1;DECONTAMINATION;EFFICIENCY;FLUIDIZED BED;FLUORIDE
VOLATILITY PROCESS: T1,Q2,Q3;PERFORMANCE;PLUTONIUM FLUORIDES:
T2;PURIFICATION;RADIOLYSIS;URANIUM HEXAFLUORIDE: T3;VAPOR
CONDENSATION

ABSTRACT: A continuous cold trap system consisting of fluidized
condenser and stripper has been evaluated with a view to
adapt it to the Fluoride Volatility Process in establishing
the continuous purification process without radiation
decomposition of PuF_6 . Its feasibility is shown by the
test with UF_6 -air. Necessary conditions for the cold
trap, and performance of the 2-in. dia fluidized bed cold
trap system are presented, and also a model of mist formation
in the condenser.

EDB SUBJECT CATEGORY: EDB-050400;050800
INIS CATEGORY: E16

<725>

Japan Atomic Energy Research Inst., Tokyo

In Fluorination process studies of plutonium dioxide by
fluid-bed ** AIX-08:300956;ERA-02:041095;EDB-77:092367; (Feb
1976) JAERI-M--6392

SUBJECT DESCRIPTORS: CONTROL;FLUIDIZED BED;FLUORIDE VOLATILITY
PROCESS;FLUORINATION: Q2,Q3;NEUTRON MONITORS;PERFORMANCE
TESTING;PLUTONIUM: T1;PLUTONIUM DIOXIDE: T2;POWDERS;RECOVERY:
Q1;SAMPLERS;SAMPLING;SYSTEMS ANALYSIS;URANIUM DIOXIDE: T3

ABSTRACT: As the first stage towards establishing the F_2
two-step fluorination process for stable recovery of
plutonium, the fluorination process has been studied with
 PuO_2 and UO_2 powder in the 2" fluid-bed.
Fluorination process characteristics and related problems
were clarified. This report describes outlines of
experimental system and its design philosophy, overall
operating test with UO_2 , fluorination studies of

<726> CONT.

PuO₂, analysis with a neutron process monitor and powder sampling.

EDE SUBJECT CATEGORY: EDB-050800

INIS CATEGORY: B16

<726>

Japan Atomic Energy Research Inst., Tokyo

In Analysis of uranium carbides ** AIX-08:300779;EDB-77:094584;
(Mar 1976) JAERI--1242

ABSTRACT: The Committee on the Analytical Chemistry of Nuclear Fuels and Reactor Materials, JAERI, undertook the analytical problems on the analysis of uranium carbides during the period November, 1971 to September, 1973. Nine laboratories participated in the cooperative analyses of three uranium carbide samples. (Samples K, M, N; two uranium monocarbide and one uranium dicarbide) for total carbon, free carbon, oxygen and nitrogen, and useful information was obtained. The analyzed samples were also useful, and the preparation and the characteristics were described together with the analytical results.

EDE SUBJECT CATEGORY: EDB-400100

INIS CATEGORY: B11

<727>

Mallory (P.R.) and Co., Inc., Burlington, Mass. (USA)

In Research on electrochemical parameters for nonaqueous batteries: task I. Quarterly progress report No. 3 **
EDB-76:059426; (Jul 1970) AD--874598

SUBJECT DESCRIPTORS: BATTERY SEPARATORS;COMPARATIVE EVALUATIONS;COPPER CHLORIDES;ELECTROCHEMISTRY: Q1;ELECTRODEPOSITION;ELECTROLYTES;LITHIUM;MEDIUM TEMPERATURE;METAL-NONMETAL BATTERIES: M1;ORGANIC NITROGEN COMPOUNDS;POLARIZATION;SOLUBILITY;STABILITY;TEMPERATURE DEPENDENCE

ABSTRACT: None

EDE SUBJECT CATEGORY: EDB-250901

<728>

<728>

Massachusetts Inst. of Tech., Cambridge (USA)

In Basic research in crystalline and noncrystalline ceramic systems. Annual report, May 1, 1975--April 1, 1976 **
 INS-08:019472; ERA-02:001048; EDB-76:081613; (1976)
 COO--2390-10

ABSTRACT: Activities in research programs on ceramics are reported in sections on electric conductivity and dielectric properties, microstructure and properties, ion transport and diffusion, defect interactions and grain boundary phenomena, and future developments. (JRD)

EDB SUBJECT CATEGORY: EDB-360200
 INIS CATEGORY: B23

<729>

Oak Ridge National Lab., Tenn. (USA)

Molten-salt systems In Chemistry Division annual progress report for period ending November 1, 1975 **
 NSA-33:28433; ERA-01:013800; EDB-76:036889; (Feb 1976)
 ORNL--5111

SUBJECT DESCRIPTORS: BENCH-SCALE EXPERIMENTS; CHROMIUM FLUORIDES: T9; ELECTRODES: T10; FABRICATION: Q10; MOLTEN SALT REACTORS; MOLTEN SALTS: T11; PHASE DIAGRAMS: Q6, Q7; RESEARCH PROGRAMS: Q11; SODIUM FLUORIDES: T8; STABILITY: Q8, Q9

ABSTRACT: Progress is reported on studies concerned with the molten-salt reactor program. Estimates of the deposition of Te on Hastelloy N surfaces in a molten salt breeder reactor (MSBR) were calculated. Experiments were conducted to evaluate the extent of Te-induced grain boundary cracking that occurs when Hastelloy N or its various modified alloys are exposed to Te vapor. Results of preliminary studies on hydrogen reduction equilibrium of UF_4 dissolved in molten fluoride mixtures and studies on the effects of oxygen on the equilibrium of UF_3 -- UF_4 molten fluoride solutions contained in graphite are reported. In studies on the chemistry of Na fluoroborate, the following compounds were prepared and characterized: H_3BO_3 , BF_3 , $2\text{H}_2\text{O}$, $\text{HBF}_2(\text{OH})_2$, NaBF_3OH , $\text{Na}_2\text{B}_2\text{F}_6$, and $\text{Na}_3\text{B}_3\text{F}_6$. Progress in studies on the thermodynamics of corrosion product fluorides in molten-salt reactor coolants is given. In experiments on the chemistry of fluorination and fuel reconstitution, a detailed investigation of the reaction $\text{UF}_5(\text{d}) + \text{H}_2(\text{g}) \rightarrow \text{UF}_4(\text{d}) + \text{HF}(\text{g})$ was carried out. A bench-scale test assembly

<729> CONT.

incorporating a small packed-bed electrode of glassy carbon spheres was constructed and was tested on the LiCl--KCl eutectic system. (JGB)

EDB SUBJECT CATEGORY: EDB-400201;400702

<730>

Oak Ridge National Lab., Tenn. (USA)

Chemistry of transuranium elements In Chemistry Division annual progress report for period ending November 1, 1975 **
NSA-33:28596;ERA-01:013852;EDB-76:037139; (Feb 1976)
CRNL--5111

SUBJECT DESCRIPTORS: CHEMISTRY;MOLECULAR STRUCTURE: Q10;RESEARCH PROGRAMS: Q18;TRANSURANIUM ELEMENTS: T18

ABSTRACT: The half-wave amalgamation potential of No (element 102) was determined. Studies on the preparation and properties of Cf are reported; previously reported results on Es were confirmed. The heat of solution of Bk was measured. Spectroscopic and x-ray diffraction studies of Es-253 bromides are reported. Salicylate complexes with Sm(III), Am(III), Th(IV), and U(IV)O₂ were prepared; crystal data are given. Results of studies on the effect of secondary bonds on the strength of O double bond Np double bond O bonds are reported. The M series x rays of Np were measured. Anomalous spin-orbit splitting of the 4f level in the actinide series is reported. Progress is reported in studies on the Cf oxide system for oxygen pressures up to 760 mm and temperatures between 25 and 1000°C, as well as structural parameters of transplutonium oxysulfates and oxysulfides and actinide and lanthanide hydroxides. The electron paramagnetic resonance spectrum of ²⁵³Es²⁺ was observed in the cubic monocrystal hosts BaF₂ and SrCl₂. (JGB)

EDB SUBJECT CATEGORY: EDB-400702;640302

<731>

Oak Ridge National Lab., Tenn. (USA)

In Summary of U.S. LMFBR programs on high temperature structural design and associated materials testing **
ERA-02:024426;EDB-77:047626; (Oct 1976) ERDA--76/146

SUBJECT DESCRIPTORS: DESIGN;LMFBR TYPE REACTORS: T1;MATERIALS TESTING;REACTOR COMPONENTS: T2,Q1;REACTOR MATERIALS: T3,Q1;RESEARCH PROGRAMS: Q2,Q3;REVIEWS

<731>

<731> CONT.

ABSTRACT: This document was prepared at the request of the Division of Reactor Development and Demonstration (DRDD), U.S. Energy Research and Development Administration. Four general areas of research and development are included: high-temperature structural design; irradiation effects--mechanical properties of structural materials; sodium environmental effects--influence of sodium on mechanical properties; and general material qualification.

EDB SUBJECT CATEGORY: EDB-210500

AUTHOR INDEX

*** A ***

- Abdel-Aziz, A.H.K. 1
- Abe, J. 646, 647, 648
- Abramov, L.A. 426
- Achard, J.C. 559
- Ackermann, R.J. 2
- Adachi, H. 259
- Adair, H.L. 18
- Adams, P.F. 529
- Afanas'ev, Y.A. 306
- Afanas'ev, Y.A. (Kubanskiy Gosudarstvennyy Univ., Krasnodar (USSR)) 40
- Afanas'ev, Yu.A. 3, 39
- Afanas'ev, Yu.A. (Kubanskiy Gosudarstvennyy Univ., Krasnodar (USSR)) 603
- Afinogenov, Yu.P. 4
- Ahmad, U.M. 5
- Alabyshev, A.F. 6
- Albright, D.I. 211, 212
- Alcock, C.B. 7
- Aleksandrovskii, S.V. 8, 548
- Alexander, C.A. 9
- Alikeev, B.D. 185
- Alire, R.M. 10
- Allen, C.B. 276
- Allen, C.W. 11
- Allen, G.C. 12
- Almazova, N.G. 339
- Alquist, L.J. 534
- Amangel'diev, B.R. 340
- Amberger, H.D. 13, 14
- Amirova, S.A. 356
- Amoros, D. (Centre National de la Recherche Scientifique, 31 - Toulou 621
- AN Azerbajdzhanskoj SSR, Baku. Inst. Neorganicheskoy i Fizicheskoy Kh 544, 545
- AN Ukrainskoj SSR, Kiev. Inst. Obshchej i Neorganicheskoy Khimii) 674
- Andrew, J.F. 15
- Andrews, L. 16, 17
- Angelini, P. 18
- Angerer, G. (Kernforschungszentrum Karlsruhe (Germany, F.R.). Inst. fu 19
- Antonov, B.D. (AN SSSR, Sverdlovsk. Inst. Ehlektrokhimii) 20
- Arabei, B.G. 215
- Ardashnikova, E.I. 21
- Arkharov, V.I. 313, 463
- Arncl'dov, M.N. 22
- Arrance, F.C. 23
- Arsent'ev, P.P. 24
- Asher, R.C. 25

AUTHOR INDEX

Astakhova, I.S. (Sibirskii
Metallurgicheskii Inst.,
Novokuznetsk (USSR) 361

Averin, V.V. 172

*** B ***

Babayan, G.G. 26

Babelot, J.F. 474

Babikova, T.F. 27

Babitzke, H.R. 468, 469

Baer, Y. 28

Baernighausen, H. 29, 378

Bagai, R.K. 30

Balabaeva, R.F. 658

Balashov, D.B. 31

Balej, J. (Ceskoslovenska Akademie
Ved, Prague) 32

Balourdet, Marcel 33

Bannella, R. 75

Barbanel, Y.A. 331

Barbanel', Yu.A. 34

Barbier, M.J. (Ecole Nationale
Superieure d'Electrochimie et
d'Electr 68

Barchuk, V.T. (AN Ukrainskoj SSR,
Kiev, Inst. Obshchej i Neorganiches
506

Barinova, M.V. (AN SSSR,
Vladivostok. Inst. Khimii) 427

Barker, M.G. 35

Barnett, B. (Soreq Nuclear Research
Center, Yavne, Israel) 539

Barton, P. (Oak Ridge National Lab.
, Tenn. (USA)) 614

Baruzdina, V.A. 602

Bas'kova, N.A. 36

Bassani, F. 186

Bates, J.B. 102

Batha, H.D. (eds.) 557

Batsanova, L.R. 21

Baum, B.A. 643

Bauman, H.F. 37, 38

Bautista, R.G. 355

Bayanov, A.P. 3, 39, 40

Becker, R. 41

Beetham, S.A. (UKAEA Research
Group, Harwell. Atomic Energy
Research 25

Begun, G.M. 203, 204

Behl, W.K. 42

Belov, B.F. 313

Belov, S.F. 43

Belyi, M.U. 44

Bennion, D.N. 246

Berard, M.F. 669

Berdieva, R.N. 256

Beresneva, T.I. 45

Berezhinsky, L.I. 46

Berezhnaya, V.T. 87, 88

Berezin, B.Y. 47

Berka, H. (Max-Planck-Institut fuer
Metallforschung, Stuttgart (F.R.G.)
523, 524

AUTHOR INDEX

- Berndt, K. 48
- Berndt, U. 49, 50, 694
- Berndt, U. (Kernforschungszentrum
Karlsruhe (F.R. Germany). Inst. fue
303
- Bernstein, E.R. 51, 52, 53
- Bernstein, J.L. 656
- Berrie, P.G. 477, 478
- Berrie, P.G. (Commission of the
European Communities, Karlsruhe (F.
R. 475
- Berrie, P.G. (Commission of the
European Communities, Karlsruhe
(Germ 476
- Besmann, T.M. 54, 55
- Bhise, V.S. 56
- Bikina, G.V. 430
- Bischof, B. 693
- Bishop, E.C. 57
- Bishop, J.L. 529, 663
- Bixon, M. (Tel Aviv Univ. (Israel).
Dept. of Chemistry) 428
- Blackburn, P.E. 281
- Blander, M. 58, 551, 552
- Blank, H. 9, 83, 110, 133, 209,
227, 398, 414, 476, 481, 560, 619
- Blasse, G. (Rijksuniversiteit
Utrecht (Netherlands). Fysisch Lab.
) 59
- Bloch, O. 507
- Blomquist, R. 367
- Bloxom, S.R. 236
- Blucher, R.L. 116
- Bober, M. 60
- Bode, D.D. 61
- Boehm, L. (Hebrew Univ., Jerusalem
(Israel)) 539
- Boerio, J. 466
- Bogatov, Yu.Eh. 153
- Bogdanov, Yu.V. 22
- Bogensberger, H.G. 475, 477
- Bohet, J. 407
- Bohres, E.W. 562
- Boivineau, J.C. 62
- Bonilla, C.F. 56
- Boom, R. 425
- Borgstedt, H.U. 63, 64
- Borsese, A. 65
- Borzenkova, M.P. 21
- Borzone, G. 65
- Bouffieux, P. 601
- Bougon, R. 66, 409
- Boundry, J.C. 67
- Bourgeois, M. 67
- Bouteillon, J. 68
- Bowen, H.K. 401
- Bowersox, D.P. 69
- Brandberg, S. 70
- Braunstein, J. 71
- Breeden, B.K. 529
- Breeden, B.K. (eds.) 663
- Breitung, W. 60, 72, 73, 74
- Bresesti, A.M. 75

AUTHOR INDEX

- Bresemi, M. 75
- Breyer, N.N. 211, 212
- Broadhead, K.G. 76
- Broers, G.H.J. 77, 563
- Bron, J. (Natal Univ., Durban
(South Africa). Dept. of Chemistry)
78
- Bronstein, H.R. 71
- Broutman, L.J. 212
- Brown, C.H. Jr. 79
- Brown, C.L. 85
- Brown, D. 80, 81, 82
- Browning, P. 83, 84
- Brownlow, D. 656
- Brumme, G.D. 478
- Brumme, G.D. (Technische Hochschule
Darmstadt (Germany, F.R.). Inst.
474
- Brumme, G.D. (Technische Hochschule
Darmstadt (Germany, F.R.)) 476
- Bryan, G.H. 85
- Budarina, A.N. (AN Ukrainskoj SSR,
Kiev. Inst. Obshchej i Neorganiche
624
- Bukhalova, G.A. 86, 87, 88, 89,
90, 636
- Bukhalova, G.A. (Rostovskij-na-Donu
Pedagogicheskij Inst. (USSR)) 629
- Bulatova, V.F. 91, 92
- Buraya, I.T. 93
- Burch, W.D. 94
- Burlakova, V.M. 86, 88
- Burlakova, V.M. (Rostovskij-na-Donu
Inzhenerno-Stroitel'nyj Inst. (US
87
- Burmeister, R.A. 95
- Burris, L. 58, 405
- Burshina, V.V. 554, 555
- Buryak, N.I. (AN Ukrainskoj SSR,
Kiev. Inst. Obshchej i
Neorganichesk 673
- Buryakov, A.V. (Rostovskij-na-Donu
Inzhenerno-Stroitel'nyj Inst. (USS
90
- Burylev, B.P. 96, 97, 98
- Buryleva, E.B. 99
- Buschow, K.H.J. 392
- Busol, F.I. 108
- Butler, C.C. 319
- Butler, J.N. 119
- Butskii, E.V. 100
- Bykhovskij, A.I. 101
- Bystrova, O.N. 136
- *** C ***
- Cafasso, F.A. 58, 405
- Cairns, E.J. 187
- Calabretta, A. 65
- Calais, D. 134
- Calaway, W.F. 404, 405
- Cantor, S. 71
- Cape, T.W. 102
- Capelli, R. 65

AUTHOR INDEX

- Capone, F. 479
- Capone, F. (Commission of the European Communities, Karlsruhe (F. R. G. 480
- Capone, F. (Commission of the European Communities, Karlsruhe (German 481
- Carnall, W.T. 353, 662
- Carroll, K. 281
- Castanet, R. (Centre National de la Recherche Scientifique, 13 - Mars 290, 291
- Cater, E.D. 597
- Cha, Y.S. (Argonne National Lab., IL) 103
- Chackraburttty, D.M. 278
- Chadha, A. 278
- Chandola, L.C. 104
- Chanthanasinh, M. 521
- Chaplygin, G.V. (AN SSSR, Moscow. Inst. Kristallografii) 105
- Charpin, P. 66, 409
- Chassay, R. 536
- Cheburkov, V.I. 106
- Chekhovskoi, V.Y. 47
- Chelyabinskij Politekhnikheskij Inst. (USSR)) 201
- Chereau, P. 107
- Chernogorenko, V.E. 321
- Chernov, V.S. 108
- Cherpillo, V.P. 394
- Chervinskii, Y.F. (Ural'skij Politekhnikheskij Inst., Sverdlovsk (USS 142
- Chervinskii, Yu.F. 141
- Chervonnyi, A.D. 253
- Chibirova, F.Kh. 657
- Chikalla, T.D. 171
- Childs, W.J. 353
- Chilton, G.R. 109, 110
- Chilton, J.E. 38
- Chilton, J.E. Jr. 111
- Chizhov, N.N. (Permskij Politekhnikheskij Inst. (USSR)) 356
- Cho, K. (Korea Univ., Kodaira, Tokyo (Japan). Faculty of Engineering) 112
- Chudinov, E.G. 113
- Chupin, A.N. 114
- Chuprikov, G.E. 115
- Chuvaev, V.F. (Moskovskij Gosudarstvennyj Univ. (USSR)) 21
- Ciani, V. 482
- Clark, R.B. 9
- Clark, R.P. 116, 117
- Clinton, J.E. Jr. 118
- Cogley, D.R. 119
- Collins, J.L. 120
- Coninck, R. de 121
- Conner, W.J. 118
- Cook, G.M. 118
- Cooper, A.R. 401
- Cooper, M.H. (ed.) 56, 103, 590
- Corbett, J.D. 378

AUTHOR INDEX

- Costello, T.J. 103
- Courths, R. 123
- Courths, R. (Freie Univ. Berlin (F. R. Germany). Inst. fuer Atom- und F 122
- Cox, D.L. 124
- Crawford, J.H. Jr. 410
- Crofts, J.A. 12
- Cronenberg, A.W. 537
- Cronenberg, A.W. (New Mexico Univ., Albuquerque (USA)) 124
- Cronenberg, A.W. (New Mexico Univ., Albuquerque (USA). Dept. of Nucle 225
- Cronenberg, A.W. (New Mexico Univ., Albuquerque (USA)) 354
- Cunningham, F.T. (Argonne National Lab., Ill. (USA)) 102
- Curtiss, L.A. 58
- Cuypers, M. 75
- *** D ***
- D'Adamo, D. 75
- Dairova, A.M. 340
- Dally, J.W. 212
- Dalvi, A.G.I. 125
- Danan, J. (CEA Centre d'Etudes Nucleaires de Fontenay-aux-Roses, 92 (F 126
- Danil'tseva, N.A. 129
- Dantseva, V.A. (Voronezhskij Gosudarstvennyj Univ. (USSR)) 4
- Dao, N.Q. 127
- Davies, B.S.J. 128
- Davies, D. 25
- Davis, S.A. 175
- Davranov, M. 256
- Davydov, A.D. 129
- De Batist, R. 132
- de Beer, R. 130
- de Bievre, P. 131
- de Boer, F.R. 425
- De Coninck, R. 132
- de Franco, M. 133
- de Franco, M. (CEA Centre d'Etudes Nucleaires de Fontenay-aux-Roses, 107
- De Keroulas, F. 134
- De Paoli, G. 80, 81
- Dean, G. 107
- Dean, J.A. (ed.) 319
- Dedjurnin, A.J. 209
- Degobert, P. 507
- Dem'yanovich, M.A. 135, 311
- Demidov, A.I. 430
- Demidov, A.I. (Leningradskij Politekhnikeskij Inst. (USSR)) 91
- den Hartog, H.W. (Rijksuniversiteit Groningen (Netherlands)) 130
- Denisova, L.A. 45
- Denisova, N.D. 136
- Deodhar, C.S. 125
- Deodhar, S.S. 137
- Dergacheva, M.B. 138, 339

AUTHOR INDEX

- Dernov-Pegarev, V.F. 396
 Desmoulin, J.F. 66, 409
 Desre, Pierre 33
 Desyatnik, V.N. 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 682
 DeVan, J.H. 309
 DeVries, L.E. 274, 275
 Dewally, D. 149
 Diamond, H. 662
 Dobrotin, R.E. (Belorusskij Tekhnologicheskij Inst., Minsk) 196
 Dode, M. 503
 Donaghey, L.F. 513
 dos Santos, A.M. 150
 Dowling, N. 695
 Downey, J.R. Jr. 276
 Dozhdeva, N.M. 426
 Drobacheva, T.I. 151
 Drotning, W.E. 152
 Dubinin, B.V. 139, 140
 Dubovoi, P.G. 505, 506
 Dubrovin, M.N. et al. (Moskovskii Inst. Stali i Splavov (USSR)) 472
 Dudareva, A.G. 153
 Dudchik, G.P. 391
 Dudnik, E.M. 215
 Dumoulin, E. 154
 Dunaeva, T.I. 255
 Dunaeva, T.I. (Rostovskij-na-Donu Filial Zaochnogo Inst. Sovetskoi To 254
 Dunitz, J.D. (ed.) (Eidgenoessische Technische Hochschule, Zurich (Swi 280, 492, 538
 Dutchak, Ya.I. 155
 Duyckaerts, G. (Liege Univ. (Belgium)) 360, 385, 386, 407, 408
 Dybkov, V.I. 156
 Dzeranova, K.B. 288
 Dzhuraev, T.D. (AN Tadzhikskoi SSR, Dushanbe. Inst. Khimii) 446
 Dzyubo, L.N. 264

 *** E ***
 Eagles, D.M. 157
 Eckert, E.R.G. 158
 Eder, O.J. 159
 Efimov, A.I. 602, 659
 Efremov, E.A. 526
 Efremov, V.A. 160, 444
 Egerev, O.I. 288
 Egleme, M. 601
 Egorov, R.F. 161
 Egorova, E.M. 583
 Ehlert, G.V. 162
 Ehpov, D.G. 427
 Elliott, F. 375
 Elliott, G.R.B. 163
 Ellis, D.E. 223, 684

AUTHOR INDEX

- Elovskikh, E.M. 164
- Epifanova, K.I. 115
- Eremenko, V.H. 156
- Eren, A. 165
- Erevanskii Gosudarstvennyi Univ. (USSR)) 593
- Ermakova, T.M. (Leningradskij Tekhnologicheskij Inst. (USSR)) 628
- Ermanovich, N.A. 643
- Ernst, K. 600
- Ershov, G.S. 198
- Esin, O.A. 512, 661
- Esin, Yu.O. 166, 201
- Evans, R. 167, 224, 247, 249, 486, 584, 686
- Evdckimov, A.A. 168
- Evers, J. 224
- Evseev, A.M. 512
- Evtushenko, N.P. 674
- Ewart, F.T. 169
- Ezhov, V.K. 170, 526
- Ezure, H. 453
- *** F ***
- Fabry, J.P. 601
- Fahey, J.A. (Ercnx Community Coll., New York (US)) 171
- Falta, G. 214
- Fassel, V.A. 319
- Fedorchenko, V.I. 172
- Fedorov, N.F. 173
- Fedorov, N.F. (AN SSSR, Leningrad. Inst. Khimii Silikatov) 206
- Fedorov, P.I. 174
- Fedorova, N.V. 45
- Fee, D.C. 175
- Feldman, M.J. 94
- Felmlee, T.L. 95
- Ferris, L.M. 176
- Ferro, R. 65
- Fes'kova, Zh.K. 553
- Filippov, E.S. (Moskovskij Inst. Stali i Splavov (USSR)) 177
- Filippov, S.I. (Moskovskij Inst. Stali i Splavov (USSR)) 372
- Fillipov, A.F. 100
- Finch, A. 178
- Fink, J.K. 367
- Fischer, D.F. 179
- Fischer, E.A. 477
- Fischer, E.A. (Kernforschungszentrum Karlsruhe (F.R. Germany). Inst. f 180
- Fischer, E.A. (Kernforschungszentrum Karlsruhe (F.R. Germany). Inst. 475
- Fischer, R.D. 13, 14, 678
- Fisher, S.B. (Central Electricity Generating Board, Berkeley (UK). Ber 181
- Fitzner, K. 182
- Fleury, P.A. 183
- Flotow, H.E. 184, 383, 384, 487

AUTHOR INDEX

- Foerthmann, R. 470
 Pcmichev, V.A. 582
 Pcmichev, V.V. 185
 Forney, J.J. (Ecole Polytechnique Federale, Lusanne, Switzerland) 186
 Foster, M.S. 187, 282
 Fournier, J.M. (CEA Centre d'Etudes Nucleaires de Grenoble, 38 (France) 188
 Fred, M. 353
 Freeman, A.J. (Northwestern Univ., Evanston, Ill. (USA). Dept. of Phys 189
 Frees, G. 63, 64
 Frenchko, V.S. (L'vovskij Gosudarstvennyj Univ. (Ukrainian SSR)) 155
 Freyland, W. 190
 Fried, S.M. 184
 Friederich, E. 191
 Friedt, J.M. 559
 Froberg, M.G. 420
 Frolov, Yu.A. 3, 39
 Fromm, E. 192, 193
 Fronczek, F.F. 194
 Fuger, J. (Liege Univ. (Belgium)) 82
 Fujino, T. 612
 Fukushima, Y. 195
 Furukawa, K. (Molten Materials Laboratory, Japan Atomic Energy Research 473
 Furuya, H. 445
 Gabelnick, S.D. 218
 Galkin, N.P. 308
 Gamanovich, N.M. 196
 Ganchenko, E.N. 40
 Ganin, G.V. (AN SSSR, Sverdlovsk. Inst. Fiziki Metallov) 161
 Gaprindashvili, V.N. (AN SSSR, Moscow. Inst. Obshchej i Neorganichesk 434
 Garanina, I.A. 99
 Garashina, L.S. 197
 Gardner, T.E. 617
 Garkushin, I.K. 637
 Garkushin, I.K. (Kujbyshevskij Politekhnikeskij Inst. (USSR)) 519, 636
 Garlinskii, R.N. (AN Ukrainskoj SSR, Lvov. Fiziko-Mekhanicheskij Inst 575
 Garner, P.L. 310
 Gasanaliyev, A.M. 519
 Gates, P.N. 178
 Gatesoupe, J.P. (CEA Centre d'Etudes Nucleaires de Fontenay-aux-Roses 133
 Gathers, G.R. 574
 Gava, S.A. 329
 Gavrilin, I.V. 198
 Gawel, W. 199
 Gazizov, R.K. 311
 Gee, R. 200
 Gel'chinskii, B.R. 512

AUTHOR INDEX

- Gel'chinskij, B.R. 661
- Gel'd, P.V. 201
- Gel'd, P.V. (Ural'skij Politekhnikheskij Inst., Sverdlovsk (USSR)) 166
- Gellings, P.J. (Technische Hogeschool Twente, Enschede (Netherlands)) 486
- Gerasimenko, L.N. 430
- Gerdanian, P. 107
- Gershunina, V.Ya. (Krasnodarskij Politekhnikheskij Inst. (USSR)) 96
- Giacchetti, G. 202
- Giegling, D. (Muenster Univ. (F.R. Germany). Anorganisch-Chemisches I 561
- Gijs, A. (Centre d'Etude de l'Energie Nucleaire, Mol (Belgium)) 121, 132
- Gilbert, B. 204
- Gilbert, B. (Univ. of Tennessee, Knoxville) 203
- Gintoft, B.I. 589
- Girifalco, L.A. 205
- Giuriato, A. 75
- Gladneva, A.F. (Moskovskij Inst. Tonkoj Khimicheskoy Tekhnologii (USSR) 43
- Glushkova, V.E. 206
- Glybin, V.P. 196
- Gnatovskaya, V.N. 46
- Gcikhman, M.S. 575
- Goldsmith, H.J. 116
- Golub, A.M. 207
- Golubin, M.A. 208
- Golubtsov, V.V. 472
- Gomozov, L.I. 209
- Goodman, L.S. 353
- Goodwin, G.M. 309
- Gopala Rao, R.V. 210
- Gorbovskaya, G.P. (Moskovskii Gosudarstvennyi Univ. (USSR)) 670
- Gordon, P. 211, 212
- Gorecki, T. (Wyzsza Szkola Inzynierska, Opole (Poland). Lab. of Physic 213
- Gorokhov, L.N. (AN SSSR, Moscow. Inst. Vysokikh Temperatur 216
- Goroncek, K. 214
- Goryachev, Yu.M. 215
- Gosudarstvennyi Inst. Redkikh Metallov, Moscow (USSR)) 494
- Gosudarstvennyj Inst. Redkikh Metallov, Moscow (USSR)) 493, 495
- Gosudarstvennyj Nauchno-Issledovatel'skij i Proektnyj Inst. Redkometa 496
- Gotkis, I.S. 216
- Goudy, A.J. 217
- Grankina, Z.A. (AN SSSR, Novosibirsk. Inst. Neorganicheskoy Khimii) 305
- Granovskaya, Zh.V. 658
- Graves, A.D. 263
- Grechina, T.N. 576
- Green, D.W. 218, 219, 220
- Greenwood, D.A. (eds.) 247, 249, 486, 584, 686

AUTHOR INDEX

Griffiths, A.J. (Central
Electricity Generating Board,
Berkeley (UK). 12

Grigor'eva, N.M. 553

Grigoryan, V.A. 100

Grishchenko, V.F. 221

Grishin, I.A. 393

Groenier, W.S. 94

Gruenberg, H. 222

Gryzalova, E.S. 516

Gubanov, V.A. (AN SSSR, Sverdlovsk.
Inst. Khimii) 223

Guenther, E. 691

Guentherodt, H.-J. 224

Guggenheim, H.J. 656

Gunde, R. 695

Gunnerson, F.S. 225

Gusarov, A.V. 216

Gutmacher, R.G. 600

*** H ***

Hadari, Z. 226

Hadari, Z. (Israel Atomic Energy
Commission, Beersheba. Nuclear
Resea 428

Hadzi, D. 695

Haines, H.R. 227, 228, 414, 415

Hall, D.G. 229

Hall, R.O.A. 414, 415

Halstead, G.W. 194

Hammou, A. 568

Harder, J.M. 230

Hardy, R.J. (Univ. of Nebraska,
Lincoln) 231

Harrar, J.E. 600

Hashimoto, T. 232

Hastings, J.C. 400

Hatakeyama, F. 445

Hatterer, A. (Ecole Supérieure de
Chimie, 68 - Mulhouse (France). Lab
154

Hauser, E. 224

Hausman, J. 233

Heady, H.H. 76

Heatherly, D. 71

Heatherly, D.E. 234

Heestand, R.L. 309

Heiss, A. 235

Hemmerich, P. (ed.) (Konstanz Univ.
(F.R. Germany). Fachbereich Biolo
280, 492, 538

Henley, E.J. (ed.) 448

Herranz, J. 236

Heuer, A.H. (eds.) 401

Hightower, J.R. Jr. 79, 237

Hildenbrand, D.L. 238, 239, 240,
241

Hill, H.H. (Los Alamos Scientific
Lab., N.Mex. (USA)) 318

Hill, M.R. (comp.) 57, 471, 692

Hilpert, K. 470

Hindman, J.C. 353

Hirata, K. 242

AUTHOR INDEX

- Hiskes, R. 95
- Hoch, M. 243, 244
- Hoechst, H. 598
- Hoekstra, H.R. 384, 662
- Hoekstra, H.R. (Argonne National Lab., Ill. (USA)) 383
- Hoerz, G. (Max-Planck-Institut fuer Metallforschung, Stuttgart (Germany)) 192
- Holleck, H. (Kernforschungszentrum Karlsruhe (Germany, F.R.)). Inst. fu 245
- Holm, R.H. (ed.) (Massachusetts Inst. of Tech., Cambridge (USA)). Dept 280, 492, 538
- Holsinger, R.W. 118
- Homs, M.H. (Technische Hogeschool Delft (Netherlands)) 130
- Hooper, A.J. (Nottingham Univ. (UK)) 35
- Hopenfeld, J. 103
- Hoppe, R. (Giessen Univ. (F.R. Germany)). Inst. fuer Anorganische und 419
- Hori, T. (Osaka Univ., Suita (Japan). Faculty of Engineering) 259
- Horiuchi, M. 700
- Horspool, J.M. 169
- Hseuh, L. 246
- Hsu, C.C. 58
- Hubberstey, P. 529
- Hubberstey, P. (Nottingham Univ. (UK)) 247
- Huefner, S. (Freie Univ. Berlin (F.R. Germany)). Inst. fuer Atom- und 123
- Huefner, S. (Universitaet des Saarlandes, Saarbruecken (Germany, F.R.)) 598
- Huijben, M.J. 248, 249, 250
- Hulet, E.K. (California Univ., Livermore (USA)). Lawrence Livermore La 61
- Huntley, W.R. 585
- Hurtgen, C. (UKAEA Research Group, Harwell. Atomic Energy Research Es 82
- *** I ***
- Ibers, J.A. (ed.) (Northwestern Univ., Evanston, Ill. (USA)). Dept. of 280, 492, 538
- Ichise, Mitsunojo 314
- Ignat'eva, L.N. (Krasnodarskij Politekhnikeskij Inst. (USSR)) 99
- Ikeda, Y. 251
- Ikhenov, D.A. 31
- Iksanov, B.A. 252
- Il'in, V.K. 253
- Il'iasov, I.I. 254, 255, 256, 257, 258, 377
- Il'iasov, I.I. (Zaochnyj Inst. Sovetskij Torgovli, Rostov-na-Donu. (US 268
- Imangaziev, E.K. (AN Kazakhskoj SSR, Alma-Ata. Inst. Organicheskogo K 138
- Imoto, S. 259
- Impolitova, I.A. 641
- Inaba, H. 260, 261, 262
- Inman, D. 263

AUTHOR INDEX

- Inyushkina, T.L. 264, 265
- Inzenhofer, A. (Kraftwerk Union A. G., Erlangen (Germany, F.R.). Abt. B 266, 267
- Isaacs, H. 309
- Iskandarov, K.I. 256, 268
- Iskandarov, K.I. (Samarkandskij Meditsinskij Inst. (USSR)) 257
- Ito, Y. 269, 704
- Ivanchenko, V.G. (AN Ukrainskoj SSR, Kiev. Inst. Metallofiziki) 610
- Ivanov, I.I. (Leningradskij Gornyj Inst. (USSR)) 8
- Ivanov, O.S. (AN SSSR, Moscow. Inst. Metallurgii) 209
- Ivanov, P.M. 135
- Ivanova, I.A. 270
- Ivanova, M.M. 185
- Ivanovskii, M.N. 22
- Ivanovskij Khimiko-Tekhnologicheskij Inst. (USSR) 216
- Ivnitskaya, B.B. 555
- Ivnitskaya, B.B. (Moskovskij Inst. Tonkoj Khimicheskij Tekhnologii (U 554
- Iyer, F.N. 655
- Iyer, B.H. 271
- *** J ***
- Jacob, K.T. 7
- Jacob, K.T. (Toronto Univ., Ontario (Canada). Dept. of Metallurgy and 182
- Jacobs, H. 272
- Jacucci, G. 273
- Jaeger, T.A. (comp.) 19, 601
- James, G. (UKAEA Reactor Group, Sellafield) 169
- James, S.D. 274, 275
- Janz, G.J. 276, 277
- Jayadevan, N.C. 278
- Jefferson, J.H. 279
- Jehn, H. 192
- Joergensen, C.K. (ed.) (Geneva Univ. (Switzerland). Dept. de Chimie M 280, 492, 538
- Joergensen, C.K. (Geneva Univ. (Switzerland). Dept. de Chimie Minerale 280
- Johnson, C.E. 175, 281, 282
- Johnson, G.K. 467
- Johnson, I. 175
- Johnson, L.F. 656
- Joshi, B.D. (Bhabha Atomic Research Centre, Bombay (India). Radiochem 125
- *** K ***
- Kabankova, N.N. 283
- Kadochnikov, V.A. 284
- Kagami, H. 699
- Kakihana, H. 285
- Kaldis, E. 224
- Kalina-Khorokhorova, L.A. 286

AUTHOR INDEX

- Kalinin, S.N. (Rostovskij-na-Donu Inzhenerno-Stroitel'nyj Inst. (USSR) 286
- Kalinyuk, N.N. (AN Ukrainskoj SSR, Kiev. Inst. Elektrosvariki) 287
- Kaloev, N.I. 288
- Kamegashira, N. 448
- Kamenetskii, M.V. 6
- Kamimoto, Masayuki (Tokyo Univ. (Japan). Faculty of Engineering) 613
- Kamkin, A.N. 129
- Kanashvili, N.V. 289, 434
- Kang, T. 290, 291
- Kanno, M. (Tokyo Univ. (Japan). Faculty of Engineering) 292, 293
- Kapoor, M.L. (Technische Univ. Berlin (F.R. Germany). Inst. fuer Meta 420
- Karapet'yants, M.Kh. 36
- Karapet'yants, M.Kh. (Moskovskij Khimiko-Tekhnologicheskij Inst. (USSR) 364
- Karasev, N.M. 294, 295, 296
- Karo, A.M. 231
- Karow, H.U. 60, 297
- Karpenko, I.F. 343
- Kartsev, V.E. 298, 299, 300
- Kase, K. 580
- Kashin, V.I. 588
- Kasimov, G.G. (Ural'skij Politekhnikeskij Inst., Sverdlovsk (USSR)) 164
- Kastenberg, W.E. (California Univ., Los Angeles (USA). Dept. of Energ 456
- Kats, S.A. 47
- Katsura, M. 581
- Katyshev, S.F. 141, 142
- Kaul, A.R. 326, 634
- Kay, D.A.R. 654
- Kazanjian, A.R. 301
- Kazanskii, K.A. 302
- Keeler, J.B. 236
- Kehiaian, H.V. 291
- Keler, E.h.k. 206
- Keller, C. 49, 50, 303
- Keller, C. (Kernforschungszentrum Karlsruhe (Germany, F.R.). Inst. fu 399, 694
- Kellogg, L.S. 416
- Kent, R.A. 15, 304
- Ketallaar, J.A.A. 563
- Ketov, A.N. (Permskij Farmatsevticheskij Inst. (USSR)) 45
- Khajmenov, A.P. (AN SSSR, Sverdlovsk. Inst. Elektrokhimii) 208
- Khaldoyanidi, K.A. 305
- Khalimonova, I.N. 46
- Khanaev, E.I. 306, 603
- Khasin, G.A. 643
- Khitrova, L.M. (Vsesoyuznii Zaochnii Politekhnikeskii Inst., Moscow 520
- Khomyakov, V.I. 307, 308
- Khorlbekk, V. 402
- Khrepinyuk, T.E. 490

AUTHOR INDEX

- Khromov, A.D. 357
- Kikindai, T. (Ecole Centrale des Arts et Manufactures, 92 - Chatenay- 424
- Kim, C.W. 366
- King, R.T. 309
- Kinsman, P.R. 180, 478
- Kinsman, P.R. (Commission of the European Communities, Karlsruhe (Ger 474
- Kirbiyik, M. 310
- Kirchheim, R. (Max-Planck-Institut fuer Metallforschung, Stuttgart (F 193
- Kirihara, T. 451
- Kirilenko, I.A. 664, 666
- Kirilenko, I.A. (AN SSSR, Moscow. Inst. Obshchej i Neorganicheskoy Kh 665, 667, 668
- Kirillov, A.I. 329
- Kiriolovich, A.P. 311
- Kirkham, I.A. 109
- Kirkham, I.A. (UKAEA Reactor Group, Windscale. Reactor Development La 110
- Kirshah, M.B. (Alexandria Univ. (Egypt). Faculty of Engineering) 1
- Kiselev, Y.M. 312
- Kisun'ko, V.Z. 313
- Kitamura, T. 314
- Klein, J.A. 79
- Klein, M.L. 273
- Kleitz, M. 568
- Klevtsov, P.V. 338
- Kleykamp, H. 315
- Kleykamp, H. (Kernforschungszentrum Karlsruhe (Germany, F.R.). Inst. f 316
- Klibanov, E.L. 588
- Klinkova, L.A. 317
- Klym, N.M. 155
- Klyuchnikova, E.F. 547
- Kmetko, E.A. 318
- Knidiri, M. (Ecole Centrale des Arts et Manufactures, 92 - Chatenay-M. 127
- Kniseley, R.N. 319
- Knoll, W. 359
- Kobayashi, K. 232
- Kober, V.I. 320
- Kobzenko, G.P. 321, 610
- Koch, E.E. (ed.) 382, 484
- Koch, G. (Kernforschungszentrum Karlsruhe (Germany, F.R.). Inst. fuer 322
- Koelling, D.D. 323
- Koelling, D.D. (Argonne National Lab., Ill. (USA)) 189
- Koizumi, M. (Power Reactor and Nuclear Fuel Development Corp., Tokai, 445
- Kojima, T. (Government Industrial Research Inst., Takamatsu (Japan)) 314
- Kolin, V.V. (Radievyj Inst., Leningrad (USSR)) 34
- Kolster, B.H. 324
- Komarevtseva, N.I. 325
- Komm, T.Z. 326

AUTHOR INDEX

- Komshilova, G.M. 93
- Kononenko, V.I. 327, 328
- Konstanchuk, I.G. 338
- Kopnina, O.I. 623
- Korenev, Y.M. (Moskovskij Gosudarstvennyj Univ. (USSR)) 296
- Korenev, Yu.M. 330
- Korenev, Yu.M. (Moskovskij Gosudarstvennyj Univ. (USSR)) 294, 295
- Kornacheva, G.M. (AN SSSR, Moscow. Inst. Elektrokhimii) 583
- Kornelli, M.E.. (AN Ukrainskoj SSR, Odessa. Inst. Obshchej i Neorgani 329
- Kornilova, V.T. (Ural'skij Politehnicheskij Inst., Sverdlovsk (USSR) 265
- Koroleva, G.N. 329
- Korshunov, B.G. 299, 300, 493, 494, 495, 496, 546
- Korshunov, B.G. (Gosudarstvennyj Inst. Redkikh Metallov, Moscow (USSR) 298
- Korshunov, B.G. (Institut Tonkoj Khimicheskoy Tekhnologii, Moscow (US 555
- Korshunov, B.G. (Moskovskij Inst. Tonkoj Khimicheskoy Tekhnologii (US 556
- Korshunov, I.A. (Gor'kovskij Gosudarstvennyj Univ. (USSR)) 351
- Kosmynin, A.S. 518
- Kosmynin, A.S. (Kujbyshevskij Sel'skohozyajstvennyj Inst. (USSR) 638
- Kosorukov, A.A. 330
- Kotera, Yoshihide 557
- Kotlin, V.P. 34, 331
- Kouhsen, C. 332, 333, 334
- Kovalev, F.V. 298, 299, 300
- Kovalevskii, A.V. 335
- Kovba, L.M. 325, 641
- Kovenskaya, B.A. 215
- Kozachenko, E.G. 336, 337
- Kozeeva, L.P. 338
- Kozhemyakin, V.A. 546
- Kozhevnikov, V.P. (Moskovskij Aviatsionnyj Inst. (USSR)) 599
- Kozhina, I.I. (Leningradskij Gornyj Inst. (USSR)) 547
- Kozin, L.F. 138, 339, 340
- Kozin, V.F. 341
- Kozina, L.N. 342
- Kozlov, V.B. 343
- Kozlova, O.B. (Vsesoyuznyj Inst. Ogneuporov, Leningrad (USSR)) 344
- Krascella, N.L. 345
- Krasilov, Yu.I. 664
- Krasnii, Yu.P. 346
- Krasnoperova, A.P. (Khar'kovskij Gosudarstvennyj Univ. (Ukrainian SSR) 347
- Krasnova, I.A. (Moskovskij Inst. Stali i Splavov (USSR)) 100
- Krasnova, O.M. 629
- Krasnova, O.M. (Rostovskij-na-Donu Pedagogicheskij Inst. (USSR)) 630, 640
- Krasulin, Yu.L. 348

AUTHOR INDEX

- Krenev, V.A. (AN SSSR, Moscow. Inst. Obshchej i Neorganicheskoy Khimi 253
- Krimnus, M.G. (Donetskij Fiziko-Tekhnicheskij Inst. (Ukrainian SSR)) 677
- Krivanev, M. 349
- Krohn, B.J. 350
- Krokhin, V.A. (Moskovskij Inst. Stali i Splavov (USSR) 494
- Krokhin, V.A. (Moskovskij Inst. Stali i Splavov (USSR) 493, 495, 496
- Krttil, J. 349
- Kruger, O.L. (Battelle Columbus Labs., Ohio (USA)) 9
- Krylov, E.I. 114, 164
- Kryukova, A.I. 351
- Ksenzenko, V.I. (Moskovskij Inst. Tonkoj Khimicheskoy Tekhnologii (US 336, 337, 553
- Kubaschewski, O. (ed.) 679
- Kucera, G.H. 58, 352
- Kuenzi, H.U. (Easel Univ. (Switzerland)) 224
- Kugel, R. 353
- Kujbyshevskii Meditsinskii Inst. (USSR)) 518
- Kujbyshevskii Politekhnikeskii Inst. (USSR) 520
- Kujbyshevskij Meditsinskij Inst. (USSR) 516
- Kujbyshevskij Meditsinskij Inst. (USSR)) 638
- Kujbyshevskij Politekhnikeskij Inst. (USSR)) 516
- Kulagin, N.M. 361
- Kuliasha, M.A. 354
- Kulova, L.K. (Gruzinskij Politekhnikeskij Inst., Tbilisi (USSR)) 288
- Kumaravadivel, R. (Bristol Univ. (UK). H.H. Wills Physics Lab.) 167
- Kunsch, B. 159
- Kuntz, L.K. (Ames Lab., IA) 355
- Kurbatov, N.N. 140, 143, 148, 682
- Kurihara, M. 646
- Kurihara, M. (Japan Atomic Energy Research Inst., Tokai, Ibaraki. Tok 647, 648
- Kuroda, T. (Tokyo Inst. of Tech. (Japan). Research Lab. of Nuclear Re 703
- Kushnir, M.N. 643
- Kushnirenko, I.Ya. 44
- Kutsenko, S.A. 356
- Kuz'micheva, E.U. 325
- Kuz'min, N.M. 357
- Kuzina, T.V. 556
- Kuznetsov, E.V. 161
- Kuzyakov, Yu.Ya. (Moskovskii Gosudarstvennyi Univ. (USSR)) 283
- Kyrs, M. 358
- *** L ***
- L'vov, S.N. 321
- Lacas, L. (Toulouse-3 Univ., 31 (France)) 621
- Lacroix, R. (Geneva Univ. (Switzerland)) 432

AUTHOR INDEX

- Lam, D.J. 662
- Lamotte, H. (Societe Belge pour l'Industrie Nucleaire, Brussels) 601
- Lamparter, P. 359
- Land, J.F. 614
- Landresse, G. 360
- Lang, H.J. 652
- Laptev, D.M. 361
- Larin, A.A. 362
- Lau, K.P. 363
- Lavrinovich, Yu.G. 311
- Lazarev, V.M. 364
- Le Bouhellec, J. 67
- Le Roy, J. 365
- Lebedev, V.A. 284, 335, 522
- Lee, H.J. 366
- Lee, J.A. 415
- Lee, J.A. (UKAEA Research Group, Harwell Atomic Energy Research Establishment) 414
- Leibowitz, L. 220, 367, 577, 578, 579
- Leibowitz, L. (Argonne National Lab., Ill. (USA)) 179
- Lemmer, R.H. (Rand Afrikaans Univ., Johannesburg (South Africa)) 368
- Lengyel, T. (Magyar Tudomanyos Akademia Izotop Intezete, Budapest) 369, 370, 371
- Lentz, A. 41
- Leonov, A.M. (Gosudarstvennyj Nauchno-Issledovatel'skij i Proektnyj I 458
- Lesnaya, M.I. 321
- Levin, Y.B. 372
- Levina, N.A. 337
- Levinskii, Yu.V. 373
- Lezzoli, L. 75
- Li, K.C. 374
- Liang, W.W. 375
- Liao, K.C. 11
- Liebllich, N. 539
- Lierde, W. van 121
- Lin, J. 534
- Linchevskij, B.V. 532
- Lindemer, T.B. 55
- Lindemer, T.B. (Oak Ridge National Lab., Tenn. (USA)) 54
- Lindgard, P. 376
- Lindner, R. (eds.) 9, 83, 110, 133, 209, 227, 303, 360, 398, 414, 476, 481, 560, 619, 683
- Lippincott, E.P. 416
- Lisinskii, V.E. (Leningradskii Politekhnikheskij Inst. (USSR)) 6
- Lisitsa, M.P. 46
- Litvinov, Yu.G. 255, 268, 377
- Litvinov, Yu.G. (Rostovskij-na-Donu Filial Zashchitnogo Inst. Sovetskij 258
- Litvinova, G.N. 87, 88
- Lityagov, V.Ya. 153
- Lochner, U. 378
- Lojchenko, V.Ya. (AN Ukrainskoj SSR, Kiev. Inst. Obshchej i Neorganicheskij 402

AUTHOR INDEX

- Lomovtsev, V.I. 327
 Lopatina, E.A. (Moskovskij Inst. Stali i Splavov (USSR)) 623
 Lorenz, R.A. 120
 Lovell, G.H.E. 379
 Lucas, J. 521
 Lucovsky, G. (ed.) 183
 Lufkin, D.A. 380
 Lundin, C.E. (ed.) 11, 65, 425
 Lutsenko, V.G. (AN Ukrainskoj SSR, Kiev. Inst. Obshchej i Neorganicheskoj 221
 Lyall, A.E. 381
 Lynch, D.W. 382, 484
 Lynchak, K.A. 321
 Lyon, W.G. 383, 384
 Lyon, W.S. (ccmp.) 423
 Lyons, K.B. 183
 Lysy, R. 360, 385, 386
 Lyzlov, Y.N. 457
- *** M ***
- MacDonald, P.E. 387
 Machado, I.J. (Bhabha Atomic Research Centre, Bombay (India). Spectro 104
 Machavariani, Z.I. (Rostovskij-na-Donu Inzhenerno-Stroitel'nyj Inst. 394
 Maeda, M. 699
 Maier, D. 49
 Maiorova, A.P. (Moskovskij Gosudarstvennyj Univ. (USSR)) 658
 Maiorova, E.A. 388
 Maiti, H.S. 389
 Majer, A.A. 36
 Makarov, L.I. (Leningradskij Gosudarstvennyj Univ. (USSR)) 390
 Makhanek, A.G. 589
 Makhmadmurodov, A. 391
 Maksyukova, E.V. (AN SSSR, Moscow. Inst. Obshchej i Neorganicheskoj K 435
 Makukhin, N.V. 634
 Mal, H.H. van 392
 Mal'kevich, N.V. 393
 Mal'tsev, A.A. (Moskovskij Gosudarstvennyj Univ. (USSR). Kafedra Fizi 497
 Mal'tsev, A.A. (Moskovskij Gosudarstvennyj Univ. (USSR). Kafedra Fizi 707
 Mal'tsev, V.T. 394
 Malek, Z. 358
 Malinauskas, A.P. 120
 Malinin, G.V. 395
 Malinin, S.D. 396
 Malm, J.G. 66, 184, 353
 Malmejac, Yves 33
 Malova, N.S. 174
 Malyshev, V.V. (Gosudarstvennyj Komitet po Ispol'zovaniyu Atomnoj Ehne 397
 Malyutina, T.M. 115
 Mamantov, G. 203

AUTHOR INDEX

- Mamantov, G. (Univ. of Tennessee, Knoxville) 204
 Manes, L. 398
 Mangelmann, M. 399
 Manning, S.R. 120
 Mar, R.W. 400
 Marchant, D.D. 401
 Marcon, J.P. 134
 Mardon, P.G. 227, 228
 Marek, G.S. 635
 Marenkova, I.N. 264
 Markov, B.F. 402
 Markov, B.G. 403
 Maroni, V.A. 102, 404, 405, 663, 701
 Maroni, V.A. (Argonne National Lab., Ill. (USA)) 702
 Marr, J.J. 58
 Martinot, L. 360, 407, 408
 Martinot, L. (Liege Univ. (Belgium)) 406
 Martynov, S.Z. 252
 Masson, J.P. 409
 Matsui, T. 260
 Matsumoto, G. (Nagoya Univ. (Japan). Faculty of Engineering) 251
 Matsumoto, W.Y. 416
 Matsunaga, M. 269
 Matsuura, S. 453
 Matthews, G.E. Jr. 410
 Mauri, R.E. 38
 Mayer, A.I. 364
 McClelland, J.F. 411
 McCloud, G.H. 187
 McClure, D.S. (Princeton Univ., N. J. (USA). Dept. of Chemistry) 412
 McCurdy, H.C. 309
 McDiarmid, R. 413
 McElroy, D.L. 415
 McElroy, D.L. (Oak Ridge National Lab., Tenn. (USA)) 414
 McElroy, W.N. 416
 McLachlan, A.D. 417
 McNeese, L.E. 176, 237
 McQuillan, A.D. (Birmingham Univ. (UK). Dept. of Physical Metallurgy a 418
 Mehlhorn, B. 419
 Mehrotra, G.M. 420
 Mel'nikov, Yu.T. 139
 Mel'nikova, O.V. 173
 Mendel'son, D.A. 575
 Meredith, G.R. 51, 52, 53
 Merrill, L. 421
 Meyer, A. 422
 Meyer, A.S. 423
 Michaels, R.G. 381
 Michaille, P. 424
 Midoriyama, S. 580
 Miedema, A.R. (Philips Research Labs., Eindhoven (The Netherlands)) 392

AUTHOR INDEX

- Miedema, A.R. (Philips Research Labs., Eindhoven, Netherlands) 425
- Miftakhov, T.T. 519
- Mikhailichenko, A.I. 426
- Mikhajlov, M.A. 427
- Miller, A.E. 11
- Miller, R.I. 103
- Miller, W.C. 591
- Mimura, T. (Tokyo Univ. (Japan). Faculty of Science) 627
- Minaev, Y.A. 252
- Minichino, C. 574
- Mintz, M.H. 226, 428
- Miguel, P. 67
- Mironov, V.L. 98, 99
- Mironov, V.L. (Krasnodarskij Politekhnikeskij Inst. (USSR)) 97
- Mirsoyanova, N.N. 89
- Mirza, Z.I. 246
- Mishchenko, V.T. (AN Ukrainskoi SSR, Odessa. Inst. Obshchei i Neorgan. 511
- Mishra, S.L. 447
- Miyajima, K. 699
- Mkrtchyan, R.F. 26
- Molodkin, A.K. (Universitet Druzhy Narodov, Moscow (USSR)) 153
- Montgomery, E.L. 429
- Morachevskii, A.G. 6, 91, 430
- Morachevskii, A.G. (Leningradskij Politekhnikeskij Inst. (USSR)) 92, 388
- Moravec, J. 431
- Moravec, J. (Ustav Jaderneho Vyzkumu, Rez (Czechoslovakia)) 349
- Moreau, J.M. 365
- Moret, J.M. 432
- Morimoto, S. 704
- Moriya, K. 242
- Morozov, A.I. 434, 435
- Morozov, A.I. (AN SSSR, Moscow. Inst. Obshchej i Neorganicheskoi Khim 289
- Morozov, A.I. (AN SSSR, Moscow. Inst. Obshchej i Neorganicheskoi Khimii 433
- Morozov, I.S. 434
- Morozov, V.A. 22
- Moser, Z. 436
- Moshnerova, I. (Statni Vyzkumny Ustav pro Stavbu Stroj, Bechovice (Cz 437
- Moskovskij Gosudarstvennyj Univ. (USSR)) 312
- Moskovskij Inst. Stali i Splavov (USSR)) 298, 555
- Moskvitina, E.N. 283
- Mueller-Buschbaum, H. 438
- Mueller, K. (Technische Hochschule, Graz (Austria)) 698
- Mueller, W. 303, 360, 683
- Muonze, R. (ed.) 632, 652
- Mukhina, I.Yu. 439
- Muller, W. 407
- Munir, Z.A. 440
- Murakami, T. (Kawasaki Heavy Industries Ltd., Kobe (Japan)) 580

AUTHOR INDEX

Murch, G.E. 442

Murch, G.E. (Flinders Univ. of South Australia, Bedford Park. School o 441

Murr, L.E. 5

Murthy, T.K.S. 447

*** N ***

Naegele, J. (Commission of the European Communities, Karlsruhe (Germa 398

Nafikova, S.Kh. 443, 444

Nagai, S. 445

Nagibina, I.E. 446

Nair, V.R. 447

Naito, K. 448

Naito, K. (Nagoya Univ. (Japan). Faculty of Engineering) 260, 261, 262, 459

Nakagawa, I. (Government Industrial Research Inst., Nagoya (Japan)) 449, 450, 451

Nakajima, T. 452

Nakanishi, K. 452

Naoumidis, A. 334

Naoumidis, A. (Kernforschungsanlage Juelich G.m.b.H. (F.R. Germany). 470

Natanzon, Ya.V. 156

Natarajan, P.R. (eds.) 271

Natsume, H. 453

Naumchik, A.N. 8, 548

Nazare, S. 454

Nebelung, C. (Zentralinstitut fuer Kernforschung, Rossendorf bei Dres 652

Nedyak, S.V. 497

Neilands, J.B. (ed.) (California Univ., Berkeley (USA). Dept. of Bioc 280, 492, 538

Nelson, A.G. 246

Neshpor, V.S. 582

Nguen Ngok Kuan 547

Nichkov, I.E. (Ural'skij Politekhnikheskij Inst., Sverdlovsk (USSR)) 320

Nichkov, I.P. 284, 335, 522

Nichkov, I.P. (Moskovskij Khimiko-Tekhnologicheskij Inst. (USSR). Nov 675

Nicholson, E.L. Jr. 455

Nickel, H. (Kernforschungsanlage Juelich G.m.b.H. (F.R. Germany). Ins 334

Nickerson, G.M. 456

Niesen, L. (Rijksuniversiteit Groningen (Netherlands)) 696

Nigmatova, R.Sh. 340

Nikolaev, N.C. (AN SSSR, Moscow. Inst. Obshchej i Neorganicheskij Khi 500

Nisel'son, L.A. 457, 458, 633

Nobile, A.A. 263

Noda, Y. 459

Novichkov, V.Kh. 461, 462

Novikov, G.I. (Belorusskij Tekhnologicheskij Inst., Minsk) 93

Novikov, I.I. 460

Novikov, N.I. 461, 462

AUTHOR INDEX

- Novikov, V.M. 302
- Novokhatskii, I.A. 463
- Novokhatskij, I.A. 313
- Novoselov, G.F. 135
- Novoselova, A.V. 21, 330, 444
- Novoselova, A.V. (Moskovskij Gosudarstvennyj Univ. (USSR). Kafedra Ne 443
- Novoselova, A.V. (Moskovskij Gosudarstvennyj Univ. (USSR)) 540, 572
- Nowotny, H. 464
- Nuernberg, H.W.
(Kernforschungsanlage Juelich G.m.b.H. (F.R. Germany) 470
- *** O ***
- O'Bryan, H.M. 656
- O'Donnell, T.A. 465
- O'Hare, P.A.G. 466, 467
- Oboznenko, V.A. 490
- Oden, L.L. 468, 469
- Odinov, B.V. 113
- Odoj, B. 470
- Offer, H.P. 471
- Ofitserov, A.A. 472
- Ohno, H. 473
- Ohse, R.W. 474, 475, 476, 477, 478, 479, 480, 481, 482
- Ohse, R.W. (Commission of the European Communities, Karlsruhe (F.R.G. 180
- Ohwada, K. (Japan Atomic Energy Research Inst., Tokai, Ibaraki. Tokai 483
- Okashita, H. 453
- Olson, C.G. 382, 484
- Olson, J.J. 485
- Ondracek, G. 454
- Ono, S. 261
- Oosterbroek, M. 486
- Orel, B. 695
- Orshich, J. 381
- Osborne, D.W. 184, 383, 384, 487
- Osborne, M.F. 120
- Osipov, N.V. 555
- Overend, J. 350
- Oyamada, R. (Nagasaki Univ. (Japan). Faculty of Education) 488
- Oyamada, Ryoza 703
- *** P ***
- Paccard, D. 365
- Paez, O.A. 380
- Panova, T.I. 206
- Paoletti Gualandi, M. 75
- Papatheodorou, G.N. 58, 489
- Parlee, N.A.D. 525
- Parry, G. 529
- Partashnikova, M.Z. 490
- Parthe, E. (Geneva Univ. (Switzerland). Lab. Crystallographie aux Ray 365

AUTHOR INDEX

- Pashchenko, A.V. (Ukrainskaya Sel'skokhozyajstvennaya Akademiya, Kiev) 101
- Pask, J.A. 557
- Pasteur, G.A. 656
- Pastukhov, Eh.A. 661
- Pathak, N.L. 491
- Patil, S.K. 660
- Pavlov, Y.A. (Moskovskij Inst. Stali i Splavov (USSR)) 252
- Peacock, R.D. (King's Coll., London (UK). Dept. of Chemistry) 492
- Peake, S.J. (Royal Holloway Coll., London (UK)) 178
- Pelipas, V.P. 357
- Percheron, A. 559
- Perepelitsa, A.P. 207
- Permyakov, P.G. 493, 494, 495, 496
- Peroni, P. 75
- Perov, P.A. 497
- Perrin, A. (Rennes-1 Univ., 35 (France)) 498
- Perrot, P. (Lille-1 Univ., 59 - Villeneuve-d'Ascq (France)) 149
- Person, W.B. 350
- Pervikova, V.N. 517
- Pervov, V.S. 500
- Pervov, V.S. (AN SSSR, Moscow. Inst. Obshchej i Neorganicheskoi Khimii) 499
- Peterson, J.R. 501
- Petrov, K.I. (Moskovskij Inst. Tonkoj Khimicheskoi Tekhnologii (USSR)) 185
- Petrov, V.A. 6
- Petrova, D.G. 636
- Petrova, G.K. 174
- Petrova, M.A. 270
- Petrushevskii, M.S. 201
- Petukhova, L.P. 265
- Pfann, J. (Ustav Jaderneho Vyzkumu, Rez (Czechoslovakia)) 502
- Phillips, B.A. 83, 84
- Piacentini, M. 484
- Pialoux, A. 503
- Pichlmayer, F. 214
- Pilkuhn, M.H. (ed.) 186
- Pintschovius, L. 567
- Pirkovskaya, N.V. 546
- Pitzer, K.S. 586
- Plakhotnik, V.N. 504
- Plizga, M.J. 23
- Podafa, B.P. 505, 506
- Podkolzina, I.G. 270
- Polart, J. 507
- Poletaev, I.F. 461, 462, 546
- Politis, C. 508
- Politis, C. (Kernforschungszentrum Karlsruhe (F.R. Germany). Inst. fue 509
- Politis, C. (Kernforschungszentrum Karlsruhe (Germany, F.R.). Inst. fu 510
- Polley, M.V. 128
- Poluehktov, N.S. 329, 511

AUTHOR INDEX

- Polukhin, V.A. 512
- Polunina, G.P. 641
- Polyachenok, O.G. 93
- Polyachenok, O.G. (Belorusskij Tekhnologicheskij Inst., Minsk) 391
- Pong, R. 513
- Popel, S.I. 514
- Popel', P.F. (Kievskij Gosudarstvennyj Univ. (Ukrainian SSR)) 207
- Popov, B.N. (Skoplje Univ. (Yugoslavia) Tehnoloskc-Metalurski Fakultet 515
- Popovskaya, N.P. 592
- Posypaiko, V.I. 516, 517, 518, 519
- Posypajko, V.I. 520
- Potapenko, O.G. 675
- Potter, P.E. 83, 84, 228
- Potter, P.E. (UKAEA Research Group, Harwell. Atomic Energy Research E 227
- Poulain, M. 521
- Poyarkov, A.M. 522
- Prasad, R. 655
- Predel, B. 523, 524
- Pressl, G. 159
- Prokopenko, V.K. (AN Ukrainskoj SSR, Donetsk. Fiziko-Tekhnicheskij In 463
- Proshina, O.F. 364
- Proskurin, V.E. 460
- Prostakov, M.E. (Ural'skij Gosudarstvennyj Univ., Sverdlovsk (USSR)) 530
- Protopapas, P. 525
- Protsenko, P.I. (Rostovskij-na-Donu Gosudarstvennyj Univ. (USSR)) 592
- Prusakov, V.N. 135, 526
- Pugh, C.E. 527
- Pulham, R.J. 529
- Pulham, R.J. (Nottingham Univ. (UK)) 528
- Pyrina, V.K. 530
- *** Q ***
- Quattropiani, A. 186
- *** R ***
- Rahman, A. 550
- Rakitskii, A.N. (AN Ukrainskoj SSR, Kiev. Inst. Metallofiziki) 531
- Rakovshchik, G.M. 532
- Ramaniah, M.V. 655
- Rand, M.H. 84
- Rand, M.H. (UKAEA Research Group, Harwell. Atomic Energy Research Est 83
- Randall, C.H. (Melbourne Univ., Parkville (Australia)) 465
- Rapp, R.A. 533
- Raspopin, S.P. 139, 141, 142, 144, 145, 146, 147, 284, 335
- Raspopin, S.P. (Ural'skij Politekhnikeskij Inst., Sverdlovsk (USSR). 143
- Rauh, E.G. 2

AUTHOR INDEX

- Ray, W.E. 57
- Raymond, K.N. (California Univ., Berkeley (USA). Dept. of Chemistry) 194
- Reeder, P.L. 534
- Reedy, G.T. 218, 219, 220
- Reetz, T. 535
- Reeves, F. 536
- Reeves, R.D. 277
- Refling, J.G. 310
- Reil, K.O. 537
- Reinen, D. (ed.) (Marburg Univ. (F.R. Germany). Fachbereich Chemie) 280, 492, 538
- Reinen, D. (Marburg Univ. (F.R. Germany). Fachbereich Chemie) 596
- Reinhardt, F.W. 117
- Reisfeld, R. 539
- Reisfeld, R. (Hebrew Univ., Jerusalem (Israel). Dept. of Inorganic and 538
- Remanick, A.H. 380
- Renard, Eh.V. 135
- Renner, T.A. 58
- Reshetnikova, L.P. 443, 444, 540, 572
- Revyakin, A.V. 342
- Reynolds, A.E. (Virginia Univ., Charlottesville (USA). Dept. of Nucle 310
- Reznik, E.M. 185
- Rhoades, S.C. 541
- Robbins, J.L. (California Univ., Livermore (USA). Lawrence Livermore 9, 619, 620
- Robertson, H.E. 585
- Robinson, D.N. 527
- Romanova, V.P. 342
- Rosa, C.J. 542
- Rosen, A. 223
- Rosenbauer, G.G. 13
- Rosenbauer, G.G. (Erlangen-Nuernberg Univ., Erlangen (Germany, F.R.). 14, 678
- Roth, S.R. 236
- Roy, S.K. (Ranchi Univ. (India). Dept. of Chemistry) 543
- Rozanov, A.N. 106
- Rspopin, S.P. 522
- Rubajlova, K.M. (eds.) 461, 462, 546
- Ruhinshtein, O.V. 628
- Rubtsov, A.S. 605
- Rubtsov, A.S. (Volgogradskij Pedagogicheskij Inst. (USSR)) 606
- Rustamov, P.G. 544, 545
- Ruzinova, E.A. 546
- Ryabov, E.N. 547
- Ryabov, Eh.N. 548
- Ryan, J.L. 85
- Ryazantseva, L.V. 659
- Rybkina, R.D. 461, 462
- Rykhail, R.M. 549
- Ryss, G.M. 201

AUTHOR INDEX

*** S ***

- Saboungi, M. 550
- Saboungi, M.I. 58, 352, 551, 552
- Safonov, V.V. 553
- Safonov, V.V. 336, 337, 554, 555, 556
- Safronov, E.K. 136
- Saito, S. 700
- Saito, Shinroku 557
- Saito, Shinroku (Tokyo Inst. of Tech. (Japan). Research Lab. of Engine 557
- Sakai, T. 627
- Sakurai, T. 558
- Saltykova, V.A. (Leningradskij Tekhnologicheskij Inst. (USSR)) 173
- Samokhin, G.A. 115
- Sanchez, J.P. 559
- Sandenshaw, T.A. (Los Alamos Scientific Lab., N.Mex. (USA)) 560
- Sandler, R.A. 8, 547, 548
- Sanker, P.E. 468, 469
- Sano, T. (Osaka Univ. (Japan)) 581
- Sara, V. 431
- Sari, C. 202
- Sauteron, J. 67
- Savva, V.I. (Zaochnyj Inst. Sovetskoy Torgovli, Rostov-na-Donu (USSR) 377
- Sawodny, W. (Ulm Univ. (F.R. Germany). Abt. Anorganische Chemie) 41
- Schaefer, H. 561
- Schenk, H.J. 562
- Schenke, M. 563
- Schiavi, W.F. 669
- Schirmacher, W. 564
- Schmitz, F. (CEA Centre d'Etudes Nucleaires de Fontenay-aux-Roses, 92 565
- Schneider, H. 64
- Schneider, H. (Kernforschungszentrum Karlsruhe (Germany, F.R.). Inst. 63
- Schoenfeld, I. 566
- Schommers, W. 567
- Schouler, E. 568
- Schretzmann, K. (Kernforschungszentrum Karlsruhe (F.R. Germany). Inst. 60
- Schulz, B. (Kernforschungszentrum Karlsruhe (Germany, F.R.). Inst. fu 454
- Schwochau, K. (Kernforschungsanlage Juelich G.m.b.H. (F.R. Germany). 562
- Seddon, J.W. 569
- Seidova, N.A. 544, 545
- Seifert, K. (Vienna Univ. (Austria). Inst. fuer Physikalische Chemie) 464
- Seiger, H.N. 381
- Sejranyan, K.B. (AN SSSR, Moscow. Inst. Kristallografii 593
- Seleznev, B.L. 570
- Selman, J.R. (ed.) 269

AUTHOR INDEX

- Sementsova, D.V. (Rostovskij-na-Donu Inzhenerno-Stroitel'nyj Inst. (U 86
- Semin, A.N. 675
- Sen, D. (Jadavpur Univ., Calcutta (India)) 210
- Sen, S.C. (Indian Inst. of Tech., Kanpur. Dept. of Physics) 491
- Seppelt, K. (Heidelberg Univ. (F.R. Germany). Inst. fuer Anorganische 571
- Serov, A.V. 135
- Sevast'yanov, V.G. 312
- Severyanina, E.N. 215
- Shaimuradov, I.B. 540
- Shajmuradov, I.B. 572
- Shakhbazov, M.G. (Arzamasskij Gosudarstvennyj Pedagogicheskij Inst. (545
- Shakhbazov, M.G. (Azerbajdzhanskij Gosudarstvennyj Pedagogicheskij In 544
- Shaner, J.W. 573, 574
- Shatinskii, V.F. 575
- Shaw, M. 380
- Shchetinina, G.P. 427
- Shejko, I.N. 576
- Sheka, I.A. (AN Ukrainskoj SSR, Kiev. Inst. Obshchej i Neorganichesk 341
- Shelton, R.A.J. 200
- Shenoy, G.K. 559
- Sheth, A. 577, 578, 579
- Shevchenko, V.B. 135
- Shimizu, H. 262
- Shimizu, S. 580
- Shinn, W.A. 175
- Shirokovskij, V.P. 161
- Shishkov, Yu.D. 307, 308
- Shmidt, V.S. 135
- Shohoji, N. 581
- Shpinel, V.S. (Moskovskij Inst. Tonkoj Khimicheskoy Tekhnologii (US 657
- Shter, G.E. 516, 638
- Shter, G.E. (Kujbyshevskii Politehnicheskii Inst. (USSR) 518
- Shulakov, A.S. 582
- Shurmovskaya, N.A. 583
- Shvets, V.T. (Odesskij Gosudarstvennyj Univ. (Ukrainian SSR)) 346
- Sidorenko, I.A. (AN Ukrainskoj SSR, Kiev. Inst. Obshchej i Neorganich 576
- Sidorov, L.N. 294, 295, 296
- Silbert, M. (East Anglia Univ. (UK). School of Mathematics and Physics 584
- Silverman, M.D. 585
- Silvester, L.F. 586
- Simonova, M.V. 286
- Singh, R.N. 685
- Sinha, V.K. (Indian Inst. of Tech., Kanpur) 587
- Sittig, L. 191
- Skiba, O.V. 135, 311, 682
- Skoropad, T.S. (Rostovskij-na-Donu Inzhenerno-Stroitel'nyj Inst. (USS 151

AUTHOR INDEX

- Skrinichenko, T.M. 588
- Skripko, G.A. 589
- Skyrme, G. 128
- Sloan, S.A. 590
- Slobodyanik, N.S. 207
- Slovyanskikh, V.K. (AN SSSR, Moscow. Inst. Obshchej i Neorganicheskij Khim 312
- Smith, D.F. 591
- Smith, F.J. 614
- Smotrakov, V.G. 592
- Sobolev, B.P. 593
- Sokhovich, E.V. 628
- Sokolova, T.D. 458
- Sokolovskii, Yu.S. 27
- Solokha, A.F. (AN SSSR, Moscow. Inst. Obshchej i Neorganicheskij Khim 664
- Solov'ev, S.I. 457
- Somiya, Shigeyuki (Tokyo Inst. of Tech. (Japan). Research Lab. of Eng 557
- Scmmers, J.A. 594
- Sood, D.D. 655
- Sorajic, V. 693
- Sotnikova, M.N. 657
- Scurkova, L. 431
- Spiegl, A.W. (Erlangen-Nuernberg Univ., Erlangen (F.R. Germany). Inst 13
- Spiridonov, M.A. 514
- Spiridonova, T.N. (AN SSSR, Moscow. Inst. Metallurgii) 460
- Spitsyn, V.I. (AN SSSR, Moscow. Inst. Obshchej i Neorganicheskij Khim 312
- Sproule, R.T. 595
- Sroelova, M.M. 27
- Sryvalin, I.T. 512
- Sryvalin, I.T. (Krasnodarskij Politehnicheskij Inst. (USSR)) 98
- Staahl, G.E. 175
- Steba, V.K. (Dnepropetrovskij Inst. Inzhenerov Zheleznodorozhnogo Tra 504
- Steeb, S. 359
- Steffens, F. 596
- Steiger, R.P. 597
- Steiner, P. 598
- Steinleitner, G. 190
- Stepanov, V.G. 599
- Stephens, F.B. 600
- Stevens, K.W.H. (Nottingham Univ. (UK)) 279
- Stievenart, M. 601
- Storonkin, A.V. 602
- Storozhenko, T.P. 603
- Storozhenko, T.P. (Kubanskij Gosudarstvennyj Univ., Krasnodar (USSR)) 306
- Stott, M.J. 422
- Stratilatov, B.V. 517
- Streicher, M.A. 604
- Stremousov, V.I. 605, 606
- Stroganov, A.I. (Ural'skij Politehnicheskij Inst., Sverdlovsk (USSR) 201

AUTHOR INDEX

Stuttgart Univ. (TH) (F.R.
Germany). Inst. fuer Metallkunde)
523, 524

Stuve, J.M. 607, 608

Subbarao, E.C. 389

Sukhman, A.L. 327

Sukhushina, I.S. 658

Sumner, D.H. 468, 469

Suponitskii, Yu.L. 364

Sus, F. 349

Suvorov, S.A. (Leningradskij
Tekhnologicheskij Inst. (USSR))
344

Suzuki, K. (Tohoku Univ., Sendai
(Japan). Research Inst. for Iron,
St 195

Suzuki, K. (Tohoku Univ., Sendai
(Japan). Research Inst. for Iron,
Ste 609

Suzuki, Y. 648

Svechnikov, V.N. 610

Synnott, J.C. 119

*** T ***

Tagawa, H. 612

Tagawa, H. (Japan Atomic Energy
Research Inst., Tokai, Ibaraki.
Tokai 611

Takahashi, A. (Japan Atomic Energy
Research Inst., Tokai, Ibaraki. To
558

Takahashi, I. 648, 649

Takahashi, Ycichi 613

Takehara, Z. 704

Talbot, J.B. 614

Tamaki, M. 251

Tamaki, S. 687

Tamaki, S. (Niigata Univ. (Japan).
Dept. of Physics) 615

Tamaki, S. (Tohoku Univ., Sendai
(Japan). Research Inst. of Mineral
D 688, 689

Tamura, S. 616

Tanamas, R. 50

Tarasov, G.G. 46

Tateno, J. 612

Taylor, A.R. Jr. 591, 617

Taylor, R. (Paris-11 Univ., 91 -
Orsay (France). Centre de Calcul)
273

Taylor, R.G. (UKAEA Research Group,
Harwell. Atomic Energy Research E
169

Ter-Arakelyan, K.A. (Erevanskij
Gosudarstvennyj Univ. (USSR)) 26

Terpilowski, J. (Akademia Medyczna,
Wroclaw (Poland). Instytut Chemii
199

Tetenbaum, M. 618

Thayer, W.L. 619, 620

Thirriot, C. 621

Thompson, J.K. 85

Thompson, L.B. (eds.) 387

Thorn, R.J. (Argonne National Lab.,
Ill. (USA)) 442

Thunder, A.E. 529

Tikhonova, A.I. 357

Tinelli, G. 154

AUTHOR INDEX

- Tippelskirch, H. v. 622
 Tiptsova-Yakovleva, V.G. 623
 Tishura, T.A. 624
 Titov, L.G. 556
 Titov, S.G. 209
 Toft Soerensen, O. 625
 Toft Soerensen, O. (Danish Atomic Energy Commission, Risoe. Research E 626
 Tolmachev, Yu.M. (Radievyj Inst., Leningrad (USSR)) 395
 Tominaga, T. 627
 Tomkins, R.P.T. 276
 Topilin, V.V. 100
 Toropov, A.N. 628
 Tovmas'yan, I.B. 640
 Tovmas'yan, I.K. 629, 630, 631
 Trekeljahr, S. (Zentralinstitut fuer Kernforschung, Rossendorf bei Dre 632
 Tret'yakov, Yu.D. 326
 Tret'yakova, K.V. 633
 Tretyakov, Yu.D. 634
 Trifonov, K.I. 146, 147
 Trifonov, K.I. (Ural'skij Politekhnikheskij Inst., Sverdlovsk (USSR)). 145
 Troyanov, S.I. 635
 Trunin, A.S. 518, 519, 520, 636, 637, 638
 Trunin, A.S. (Vsesoyuznyj Zashchnyj Politekhnikheskij Inst., Moscow (U 516
 Trunov, V.K. (Moskovskii Gosudarstvennyi Univ. (USSR)) 639
 Trunov, V.K. (Moskovskij Gosudarstvennyj Univ. (USSR)) 160
 Trunov, V.K. (Moskovskij Gosudarstvennyj Univ. (USSR). Kafedra Neorga 168
 Tsarenko, A.F. 135
 Tsemekhman, L.Sh. 98
 Tsirel'nikov, V.I. (Moskovskij Gosudarstvennyj Univ. (USSR)) 635
 Tsushba, T.M. 631, 640
 Tsvigunov, A. 641
 Turcotte, R.P. 171
 Twidwell, L.G. 642
 Tyagunov, G.V. 643
 Tyurin, V.I. 633
 *** U ***
 Ugajin, M. 644, 645, 646, 647, 648, 649
 Ugajin, Mitsuhiro (Japan Atomic Energy Research Inst., Tokai, Ibaraki. 650
 Ukanwa, A.O. 651
 Ukhov, V.P. 512, 661
 Ukshe, E.A. 317
 Ullmann, H. 652
 Ulseth, J.A. 416
 Umezawa, H. 453
 Ungarish, M. 566
 Ural'skij Politekhnikheskij Inst., Sverdlovsk (USSR)) 675

AUTHOR INDEX

Uzuki, K. 653

*** V ***

Vahed, A. 654

Vaidya, V.N. 655

Vakhobov, A.V. 446

Valishev, M.G. 166

Vallet, C.E. 71, 234

van Audenhove, J. (Commission of
the European Communities, Geel
(Belg 131

Van Ballegoy, H.J.J. 77

van de Braak, H.P. 486

van der Lugt, W. (Rijksinstituut
voor Zuivering van Afvalwater,
Voorb 248van der Lugt, W. (Rijksuniversiteit
Groningen (Netherlands)) 249, 250

van Ormondt, E. 130

van Tets, A. (Atomic Energy Board,
Pelindaba, Pretoria (South Africa)
379

Van Tuyl, H.H. 85

Van Uiter, L.G. 656

Varfolomeev, M.B. 657

Vasil'chenko, L.M. 637

Vasil'eva, I.A. 658

Vasil'kova, I.V. 602, 659

Vasina, N.A. 516

Vasudeva Rao, P.R. 660

Vatolin, N.A. 512, 661

Veal, B.W. 662

Veerkamp, G.R. 440

Veleckis, E. 404, 405, 663, 702

Venugopal, V. 655

Vernardakis, T. (Cincinnati Univ.,
Ohio (USA). Dept. of Materials Sci
243, 244

Vesely, V. 358

Viljoen, J. du P 368

Vinogradov, B.G. (Moskovskij Inst.
Stali i Splavov (USSR)) 24Vinogradov, E.E. 664, 665, 666,
667, 668

Vishnyakov, Y.S. 197

Visser, R.G. 669

Viting, L.M. 670

Vivchar, O.I. 671

Vlasov, V.G. 362

Vlasov, Yu.G. (Leningradskij
Gosudarstvennyj Univ. (USSR)) 570Volchanskaya, V.V. (Zaochnyj Inst.
Sovetskoj Torgovli, Rostov-na-Donu
255

Volkov, S.V. 673, 674

Volkov, S.V. (AN Ukrainskoj SSR,
Kiev. Inst. Obshchej i
Neorganicheskoe 672

Volkovich, A.V. 673

Vollrath, D. 676

Voloshin, V.A. 677

von Amberger, H.D. 678

von Goldbeck, O. 679

von Goldbeck, O. (Technische
Hochschule Aachen (Germany, F.R.).
Lehrst 680

AUTHOR INDEX

- von Tippelskirch, H. (Karlsruhe Univ. (TH) (Germany, F.R.). Inst. fuer 681
- Vendra, B.L. 94
- Vorob'eva, G.V. (Moskovskij Inst. Tcnkoj Khimicheskoy Tekhnologii (US 174
- Vorobei, M.P. 148, 682
- Vorobey, M.P. 311
- Voronezhskii Meditsinskii Inst. (USSR)) 520
- *** W ***
- Waber, J.T. (Commission of the Eurcpean Communities, Karlsruhe (F. R. G 683
- Walch, P.F. 684
- Wang, W.L. 685
- Warke, W.R. 211
- Warner, A.W. Jr. 656
- Warrier, A.V.R. (Solid State Physics Lab., Delhi (India)) 30
- Waseda, Y. 687, 688, 689
- Waseda, Y. (Tohcku Univ., Sendai (Japan). Research Inst. of Mineral D 242
- Waseda, Y. (Tohcku Univ., Sendai (Japan). Research Inst. of Mineral Dr 686
- Watanabe, N. (Kycto Univ. (Japan). Faculty of Engineering) 452
- Waterbury, G.R. (comp.) 690
- Watson, J.S. 529, 663
- Waugh, A.B. 465
- Wedemeyer, H. 691
- Weeks, R.W. 692
- Wenzl, H. 693
- Westrum, E.F. Jr. (Michigan Univ., Ann Arbor (USA). Dept. of Chemistr 594
- Whittaker, B. 81
- Whittaker, B. (UKAEA Research Group, Harwell. Atomic Energy Research 80
- Wichmann, U. 694
- Wiffen, P.W. 529, 663
- Wild, J.F. 61
- Williams, C. 353
- Williams, J.M. (Argonne National Lab., IL) 695
- Williams, R.J.P. (ed.) (Oxford Univ. (UK). Inorganic Chemistry Lab.) 280, 492, 538
- Winslow, G. 579
- Wit, H.P. 696
- Woolfrey, J.L. (Australian Atomic Energy Commission Research Establish 697
- Wright, J.F. (Battelle Pacific Northwest Labs., Richland, Wash. (USA) 534
- Wright, S.J. (New South Wales Univ., Kensington (Australia). Dept. of 697
- Wurzinger, H. 698
- *** Y ***
- Yagi, E. 699, 700
- Yagub'yan, E.S. 90

AUTHOR INDEX

- Yagub'yan, E.S. (Rostovskij-na-Donu Inzhenerno-Stroitel'nyj Inst. (US 89
- Yamagami, S. 292, 293
- Yamamoto, M. 232
- Yato, Y. (Tokyo Inst. of Tech. (Japan). Research Lab. of Nuclear Reac 285
- Yatsenko, S.F. 328
- Yatsimirskii, K.B. (AN Ukrainskoj SSR, Kiev. Inst. Fizicheskoi Khimii 674
- Yokokawa, T. (Hokkaido Univ., Sapporo (Japan). Faculty of Science) 616
- Yonco, R.M. 404, 405, 701, 702
- Yoshida, S. 488, 703
- Yoshizawa, S. 269, 704
- Young, W.H. 705
- Young, W.H. (East Anglia Univ. (UK). School of Mathematics and Physics 230
- Young, W.H. (Queen's Univ., Kingston, Ontario (Canada). Dept. of Phys 422
- Zarechnyuk, O.S. (L'vovskij Gosudarstvennyj Univ. (Ukrainian SSR)) 549, 671
- Zavalishin, N.I. 707
- Zhukova, L.D. (Irkutskij Gosudarstvennyj Univ. (USSR)) 709
- Zhul'kov, V.I. (Vsesoyuznyj Nauchno-Issledovatel'skij Inst. Sinteticheskoe 708
- Zimina, T.D. 709
- Zimkina, T.M. 582
- Zocher, R.W. 15, 304
- Zuercher, C. 28

*** Z ***

Zador, S. (Toronto Univ., Ontario (Canada). Dept. of Metallurgy and M 7

Zakharevich, A.A. 548

Zakharov, D.M. 706

Zakulski, W. 436

Zaporozhets, E.G. 90

SUBJECT DESCRIPTORS INDEX

*** A ***

ABSORPTION 152, 345, 474, 491

ABSORPTION HEAT 101

ABSORPTION SPECTRA 13, 31, 44, 46,
81, 165, 219, 350, 385, 386, 413,
497, 673ABSORPTION SPECTRA: Q1 34, 208,
283, 331, 353, 511, 718

ABSORPTION SPECTRA: Q1,Q2 30, 677

ABSORPTION SPECTRA: Q1,Q2,Q3 12

ABSORPTION SPECTRA: Q1,Q2,Q3,Q4 53

ABSORPTION SPECTRA: Q1,Q2,Q3,Q4,Q5,
Q6,Q7,Q8,Q9,Q10,Q11,Q12 382

ABSORPTION SPECTROSCOPY 12, 122

ABSORPTION SPECTROSCOPY: Q1 319,
566ABSORPTION SPECTROSCOPY: Q1,Q2
123, 431

ABSORPTION: Q4 217

ACCOUNTING 600

ACCOUNTING: Q2 131, 349

ACCURACY 22, 104, 152, 349

ACTINIDE ALLOYS: T1 189

ACTINIDE ALLOYS: T3 323

ACTINIDE COMPLEXES 13

ACTINIDE COMPOUNDS: T2 2, 323

ACTINIDES: M1 188

ACTINIDES: M14 662

ACTINIDES: T1 318, 323

ACTINIDES: T2 189

ACTINIUM 188

ACTIVATION ENERGY 24, 106, 137,
141, 146, 198, 312, 313, 328,
341, 395, 436, 504, 676

ADDITIVES 111

ADHESION 575

ADHESION: Q1,Q2,Q3 327

ADSORPTION 94, 108, 722

AFFINITY 342

AFTER-HEAT REMOVAL: Q2 367

AGING 568

ALGORITHMS 517

ALKALI METAL COMPLEXES: T2 673

ALKALI METAL COMPOUNDS 490, 528,
592ALKALI METAL COMPOUNDS: T1 20,
317, 631

ALKALI METAL COMPOUNDS: T2 489

ALKALI METALS 63, 269

ALKALI METALS: M4 671

ALKALI METALS: T1 192, 486

ALKALI METALS: T2 161

ALKALI METALS: T3 64, 343

ALKALINE EARTH METAL COMPOUNDS 490

ALKALINE EARTH METALS: T2 192

ALLOTROPY 206, 361

ALLOY SYSTEMS 641

ALLOY-VM-1: T1 129

ALLOYS 99

ALPHA DECAY: Q1 487

ALPHA SOURCES 301

ALUMINATES 23, 111, 357

SUBJECT DESCRIPTORS INDEX

ALUMINATES: T1 326	AMERICIUM COMPOUNDS: M2 61
ALUMINATES: T3 77	AMERICIUM FLUORIDES 61, 307, 308
ALUMINIUM 104, 111, 211, 440, 696	AMERICIUM OXIDES 61, 303
ALUMINIUM ALLCYS 212, 269	AMERICIUM OXIDES: M11 662
ALUMINIUM ALLCYS: M1 671	AMERICIUM 241: T10 714
ALUMINIUM ALLCYS: T1 201, 598	AMERICIUM 243: T1 61
ALUMINIUM BASE ALLCYS: T1 48	AMERICIUM: T1 308
ALUMINIUM BASE ALLOYS: T2 549	AMERICIUM: T2 307
ALUMINIUM CHLORIDES 111, 118	AMERICIUM: T6 683
ALUMINIUM CHLORIDES: T1 434, 546, 627	AMMONIA 70
ALUMINIUM FLUORIDES 440	AMMONIUM CHLORIDES 467
ALUMINIUM FLUORIDES: T1 26, 203, 204	AMMONIUM COMPOUNDS: T1 127
ALUMINIUM HYDRIDES: T4 663	AMMONIUM COMPOUNDS: T6 427
ALUMINIUM NITRIDES 23	AMMONIUM COMPOUNDS: T8 66
ALUMINIUM OXIDES 71, 246, 557	AMORPHOUS STATE 609
ALUMINIUM OXIDES: T1 104, 244, 710	AMORPHOUS STATE: T1 653
ALUMINIUM OXIDES: T2 203	AMORPHOUS STATE: T3,Q1 229
ALUMINIUM OXIDES: T3 243	AMPEROMETRY: Q4 713
ALUMINIUM OXIDES: T4 303	AMPEROMETRY: Q45 712
ALUMINIUM OXIDES: T5 372	ANALYTICAL SOLUTION 437, 615
ALUMINIUM OXIDES: T8 531	ANGULAR MOMENTUM 230
ALUMINIUM: T1 156, 159, 167, 327	ANIONS 30, 44, 314, 424, 488, 504
ALUMINIUM: T11 711	ANISOTROPY 122, 197, 363
ALUMINIUM: T16 713, 714	ANL 58
ALUMINIUM: T8 712	ANNEALING 471, 531
AMERICIUM 188	ANNIHILATION 48
AMERICIUM CARBONATES 61	ANODES 450, 507, 675
AMERICIUM CHLORIDES 34	ANODES: Q1 38, 269, 274
	ANODES: Q1,Q2 275

SUBJECT DESCRIPTORS INDEX

ANOMALOUS DIMENSION 183

ANTIFERROMAGNETISM 368

ANTIMONY 211, 328

ANTIMONY ALLCYS: M1 359

ANTIMONY ALLCYS: T3 190

ANTIMONY OXIDES: T1 657

ANTIMONY: T16 104

ANTIMONY: T3 642

ANTIMONY: T31 711

ANTIMONY: T32 712

ANTIMONY: T39 714

AQUEOUS SOLUTIONS 70

ARGON 16, 17, 628

ARSENATO 660

ARSENIC 211

ARSENIC COMPOUNDS 321

ARSENIC: T10 712

ARSENIC: T12 711

ARSENIC: T18 714

ARSENIC: T2 321

ARSENIDES: T1 321

ATCM TRANSPORT 159

ATCM TRANSPORT: Q1 401

ATOMIC MODELS 499, 609, 653

ATOMIC NUMBER 426, 686

AUTOMATION 690

AUTOMOBILES: T2 704

AUTORADIOLYSIS: Q1 301

*** B ***

B CODES 75

B CODES: Q3 272

BAKING 710

BAND THEORY 52, 157, 189, 259,
323, 411, 485, 491, 598, 662, 683

BARIUM 470

BARIUM ALLOYS: T1 628

BARIUM BROMIDES: T1 240

BARIUM CHLORIDES: T1 518

BARIUM CHLORIDES: T2 240

BARIUM CHLORIDES: T4 86

BARIUM CHLORIDES: T5 139

BARIUM COMPOUNDS 628

BARIUM COMPOUNDS: T5 638

BARIUM COMPOUNDS: T7 516

BARIUM FLUORIDES 95, 534

BARIUM FLUORIDES: T1 419, 469,
516, 638

BARIUM FLUORIDES: T2 462

BARIUM FLUORIDES: T3 521

BARIUM FLUORIDES: T4 518

BARIUM IODIDES: T3 241

BARIUM ISOTOPES: T1 534

BARIUM NITRATES 42

BARIUM OXIDES 95, 202, 470

BARIUM TUNGSTATES: T6 516

BARIUM: T1 224

BARIUM: T13 712

SUEJECT DESCRIPTORS INDEX

BARIUM: T14	711	BINARY ALLOY SYSTEMS	1, 48, 91, 137, 190, 201, 205, 284, 290, 291, 320, 321, 335, 339, 388, 403, 439, 446, 679, 680, 706
BARIUM: T18	713	BINARY ALLOY SYSTEMS: M1	525
BARIUM: T21	714	BINARY MIXTURES	73, 74, 97, 98, 139, 145, 146, 151, 160, 266, 267, 270, 288, 303, 370, 435, 443, 469, 475, 477, 506, 547, 550, 592, 676
BARIUM: T3	202	BINARY MIXTURES: Q1, Q2, T3	489
BATTERY SEPARATORS	727	BINDING ENERGY	113, 188, 209, 213, 398
BATTERY SEPARATORS: T2, Q1	23	BINDING ENERGY: Q1, Q2	186
BCC LATTICES	177, 198, 318	BINDING ENERGY: Q1, Q2, Q3, Q4, Q5, Q6, Q7, Q8	499
BENCH-SCALE EXPERIMENTS	729	BISMUTH	79, 237, 328
BERKELIUM	188	BISMUTH ALLOYS	420, 544, 545
BERKELIUM CHLORIDES: T1	501	BISMUTH ALLOYS: M2	359
BERKELIUM OXIDES: M13	662	BISMUTH ALLOYS: T1	514
BERKELIUM: T6	501	BISMUTH ALLOYS: T2	1
BERYLLIUM	500	BISMUTH BASE ALLOYS	187
BERYLLIUM CHLORIDES: T1	506	BISMUTH BROMIDES: T1	288
BERYLLIUM FLUORIDES	79, 176, 585	BISMUTH CHLORIDES: T1	553
BERYLLIUM FLUORIDES: M1	616	BISMUTH IODIDES: T2	288
BERYLLIUM FLUORIDES: T1	21, 499, 617	BISMUTH OXIDES	420
BERYLLIUM OXIDES: T1	557	BISMUTH TELLURIDES: T1	544, 545
BERYLLIUM OXIDES: T2	531, 717	BISMUTH: T1	1
BERYLLIUM: T1	717	BISMUTH: T15	712
BERYLLIUM: T14	712	BISMUTH: T16	711
BERYLLIUM: T15	711	BISMUTH: T2	420
BERYLLIUM: T19	713	BISMUTH: T20	713
BERYLLIUM: T22	714	BISMUTH: T23	714
BESSEL FUNCTIONS	564		
BIBLIOGRAPHIES	276, 412, 448, 525, 528, 660, 686		
BIBLIOGRAPHIES: Q1	158		
BIMETALS	156		

SUBJECT DESCRIPTORS INDEX

BISMUTH: T3 544
 BISMUTH: T4 104
 BOLTZMANN EQUATION 661
 BOND ANGLE 325, 695
 BOND LENGTHS 12, 66, 325, 609, 653, 695
 BOR-60 REACTOR 311
 BOR-60 REACTOR: T1 135
 BORANES 194
 BORATES 44, 246
 BORIDES: T2 215
 BORON 108
 BORON ALLOYS 274, 275
 BORON CARBIDES: T1 440
 BORON COMPOUNDS: A4 176
 BORON FLUORIDES: A6 176
 BORON FLUORIDES: T1 504
 BORON FLUORIDES: T2 499
 BORON NITRIDES 71
 BORON OXIDES 95
 BORON OXIDES: T3 183
 BORON OXIDES: T4 710
 BORON OXIDES: T7 303
 BORON: T12 712
 BORON: T13 711
 BORON: T17 713
 BORON: T20 714
 BORON: T3 104
 BOUNDARY CONDITIONS 273, 310, 401, 437, 502, 601
 BREEDER REACTORS 79, 236
 BREEDING BLANKETS 175
 BRILLOUIN ZONES 161, 560, 683
 BROMIDES 111
 BROMIDES: M1 17
 BROMIDES: T3 20
 BROMINE COMPOUNDS: T1 556
 BUBBLES: Q1 685
 BUILDING MATERIALS: T1 212
 BURNABLE POISONS 50
 BURNUP 75, 281, 315, 316, 349, 416, 453, 632
 BURNUP: Q3 715
 BURNUP: Q4 716
 BURNUP: T2, Q1 214
 BWR TYPE REACTORS 455, 510
 BWR TYPE REACTORS: T2 509

 *** C ***
 CADMIUM 211, 212, 417, 500, 523
 CADMIUM ALLOYS: T1 340, 523, 524, 605, 606
 CADMIUM ALLOYS: T2 403, 675
 CADMIUM CHLORIDES: T1 199, 257
 CADMIUM COMPLEXES: T1 314
 CADMIUM IODIDES: T2 257
 CADMIUM NITRATES: T1 592
 CADMIUM NITRATES: T4 592
 CADMIUM SULFIDES: T2 186

SUBJECT DESCRIPTORS INDEX

CADMIUM: T 210	CALCIUM OXIDES: M1 372
CADMIUM: T1 198, 403, 675	CALCIUM OXIDES: T1 496
CADMIUM: T17 712	CALCIUM OXIDES: T2 389
CADMIUM: T18 711	CALCIUM OXIDES: T3 352
CADMIUM: T22 713	CALCIUM TUNGSTATES: T2 637
CADMIUM: T25 714	CALCIUM: T16 712
CADMIUM: T5 104	CALCIUM: T17 711
CALCINATION 114, 164	CALCIUM: T21 713
CALCITE: T1 396	CALCIUM: T24 714
CALCIUM 380	CALCULATION METHODS 231, 328
CALCIUM BROMIDES: T3 240	CALIBRATION 660
CALCIUM CARBONATES 70	CALIBRATION: T1 64
CALCIUM CHLORIDES 452	CALIFORNIUM 188
CALCIUM CHLORIDES: T1 117, 146, 637	CALIFORNIUM BROMIDES: T5 501
CALCIUM CHLORIDES: T2 352	CALIFORNIUM CHLORIDES: T2 501
CALCIUM CHLORIDES: T3 139	CALIFORNIUM OXIDES: M8 662
CALCIUM COMPOUNDS 23	CALORIMETERS 179
CALCIUM COMPOUNDS: M1 495	CALORIMETRY 107, 290, 306, 355, 383, 467, 594, 603, 607, 616
CALCIUM COMPOUNDS: T2 628	CALORIMETRY: Q1 500
CALCIUM COMPOUNDS: T3 117	CAPILLARY FLOW: Q5 101
CALCIUM COMPOUNDS: T4 116, 519	CARBIDES: M5 646
CALCIUM FLUORIDES 307, 308, 533, 589	CARBIDES: T2 582
CALCIUM FLUORIDES: T1 363, 468, 519, 669, 718	CARBON 63, 64, 83, 324, 442, 644, 646, 704
CALCIUM FLUORIDES: T2 26	CARBON ADDITIONS 313
CALCIUM IODIDES: T1 241	CARBON BLACK 381
CALCIUM NITRATES: T1 254	CARBON COMPOUNDS: M3 646
CALCIUM NITRATES: T2 592	CARBON DIOXIDE 70
	CARBON MONOXIDE: T2 304

SUBJECT DESCRIPTORS INDEX

CARBON NITRIDES 332	CERIUM ALLOYS: T2 598
CARBON NITRIDES: T3 333	CERIUM CHLORIDES: T1 678
CARBON STEELS: M2 311	CERIUM COMPLEXES 424
CARBON SULFIDES: T3 609	CERIUM FLUORIDES 424
CARBON TETRACHLORIDE: T6 301	CERIUM FLUORIDES: T1 312, 461, 677
CARBON: T1 650	CERIUM FLUORIDES: T2 382, 484
CARBON: T2 714	CERIUM HYDROXIDES 424
CARBON: T3 235, 712	CERIUM NITRATES: T2 164
CARBON: T4 711	CERIUM NITRIDES: T1 28, 581
CARBON: T8 713	CERIUM OXIDES 281
CARBONATES 71, 563	CERIUM OXIDES: T1 626
CARBONATES: T2 77	CERIUM OXIDES: T2 654
CARBOXYLIC ACID ESTERS 38, 111, 118, 381	CERIUM OXIDES: T5 191
CARBURIZATION 580	CERIUM: T1 125, 230, 355, 688
CARRIER-FREE ISOTOPES: T1 723	CESIUM 120, 175, 343, 599
CARRIERS 411, 447	CESIUM ALLOYS: T1 190
CAST IRON 25	CESIUM ALLOYS: T2 249, 339
CATHODES 92, 111	CESIUM ALLOYS: T3 138
CATHODES: Q1 38, 381	CESIUM BROMIDES 483, 631
CATIONS 68, 123, 127, 130, 261, 314, 341, 386, 423, 445, 450, 538, 664, 677	CESIUM BROMIDES: M3 17
CAVITATION: Q2 103	CESIUM BROMIDES: T1 268, 543, 640
CAVITIES 115	CESIUM BROMIDES: T2 630
CENTRIFUGATION 722	CESIUM CARBONATES: T2 120
CERAMICS 64, 226, 244, 557, 595, 613, 652	CESIUM CHLORIDES 386, 498, 631, 673
CERAMICS: T3 25	CESIUM CHLORIDES: M3 16
CERIUM 28, 598, 683	CESIUM CHLORIDES: T1 6, 13, 14, 44, 46, 96, 98, 178, 196, 402, 433, 435, 491, 630, 659
CERIUM ALLOYS: T1 39, 689	CESIUM CHLORIDES: T2 87, 88, 265, 408, 543, 640, 678

SUBJECT DESCRIPTORS INDEX

- CESIUM CHLORIDES: T4 91
 CESIUM COMPLEXES: T1 674
 CESIUM COMPOUNDS 34
 CESIUM COMPOUNDS: T2 556
 CESIUM COMPOUNDS: T4 66
 CESIUM FLUORIDES 307
 CESIUM FLUORIDES: T1 41, 572
 CESIUM FLUORIDES: T2 127
 CESIUM HYDROXIDES: T3 120
 CESIUM IODIDES: T2 178
 CESIUM IODIDES: T3 630, 640
 CESIUM ISOTOPES: T2 534
 CESIUM NITRATES 673
 CESIUM NITRATES: T1 377, 466
 CESIUM NITRATES: T2 254
 CESIUM NITRATES: T3 695
 CESIUM OXIDES 281
 CESIUM TUNGSTATES: T1 151
 CESIUM URANATES: M4 120
 CESIUM 134 75
 CESIUM 137 75, 214
 CESIUM: T1 22, 248, 249, 250, 460, 528
 CESIUM: T2 346, 622
 CESIUM: T3 681
 CHAPMAN-ENSKOG THEORY 124
 CHARGE CARRIERS 165
 CHARGE DENSITY 223
 CHARGE EXCHANGE: Q1,Q2,Q3,Q4 218
 CHARGE TRANSPORT 689
 CHEMICAL ACTIVATION: Q1 63
 CHEMICAL ANALYSIS 27, 45, 319, 506, 600, 716
 CHEMICAL ANALYSIS: Q1 104, 203, 423, 623, 711, 712, 714
 CHEMICAL ANALYSIS: Q1,Q2 713, 721
 CHEMICAL ANALYSIS: Q15,Q16 125
 CHEMICAL ANALYSIS: Q2 447, 490, 566
 CHEMICAL ANALYSIS: Q2,Q3,Q4 690
 CHEMICAL BONDS 16, 17, 24, 127, 223, 259, 280, 596, 672
 CHEMICAL BONDS: Q1 483, 498
 CHEMICAL BONDS: Q1,Q2,Q3 398
 CHEMICAL COMPOSITION 3, 4, 9, 24, 64, 73, 74, 107, 115, 116, 133, 146, 147, 169, 190, 234, 247, 276, 278, 286, 290, 296, 316, 321, 330, 359, 371, 378, 379, 389, 398, 414, 420, 433, 441, 443, 456, 479, 481, 506, 521, 540, 565, 568, 604, 606, 611, 626, 636, 638, 659, 691, 710
 CHEMICAL COMPOSITION: Q1 216
 CHEMICAL COMPOSITION: Q1,Q2 93, 109, 110
 CHEMICAL COMPOSITION: Q1,Q2,Q3,Q4 418
 CHEMICAL PREPARATION 41, 81, 105, 206, 267, 304, 325, 332, 399, 419, 482, 557, 596, 635, 667, 673, 691, 703
 CHEMICAL PREPARATION: Q1 18, 305, 498
 CHEMICAL PREPARATION: Q1,Q2 194, 266, 278, 330, 409, 571
 CHEMICAL PREPARATION: Q1,Q2,Q3 80

SUBJECT DESCRIPTORS INDEX

- CHEMICAL PREPARATION: Q1,Q2,Q3,Q4,
Q5,Q6,Q7,Q8 82
- CHEMICAL PREPARATION: Q1,Q2,Q3 114
- CHEMICAL PREPARATION: Q1,Q2,Q3,Q4,
Q5,Q6,Q7,Q8 191
- CHEMICAL PREPARATION: Q1,Q2,Q3 207
- CHEMICAL PREPARATION: Q1,Q2,Q3,Q4,
Q5,Q6,Q7,Q8,Q9 314
- CHEMICAL PREPARATION: Q1,Q2,Q3,Q4,
Q5 333
- CHEMICAL PREPARATION: Q1,Q2,Q3,Q4
338, 710
- CHEMICAL PREPARATION: Q2 55
- CHEMICAL PROPERTIES 282, 501
- CHEMICAL PROPERTIES: Q1 305, 426,
652
- CHEMICAL PROPERTIES: Q1,Q2,Q3 2,
360
- CHEMICAL PROPERTIES: Q1,Q2,Q3,Q4,
Q5,Q6,Q7,Q8,Q9 66
- CHEMICAL PROPERTIES: Q5 311
- CHEMICAL RADIATION EFFECTS: Q1,Q2
169
- CHEMICAL REACTION KINETICS 16, 17,
45, 64, 66, 91, 247, 265, 320,
333, 356, 360, 395, 424, 445,
561, 646
- CHEMICAL REACTION KINETICS: Q1,Q2
27
- CHEMICAL REACTION KINETICS: Q1 68
- CHEMICAL REACTION KINETICS: Q1,Q2,
Q3 156
- CHEMICAL REACTION KINETICS: Q1,Q2
295
- CHEMICAL REACTION KINETICS: Q1,Q2,
Q3,Q4 341
- CHEMICAL REACTION KINETICS: Q1 385
- CHEMICAL REACTION KINETICS: Q1,Q2,
Q3,Q4,Q5,Q6 427
- CHEMICAL REACTION KINETICS: Q1,Q2
650
- CHEMICAL REACTION KINETICS: Q2,Q3
106
- CHEMICAL REACTION KINETICS: Q2,Q3,
Q4,Q5 332
- CHEMICAL REACTION KINETICS: Q2,Q3
504
- CHEMICAL REACTION KINETICS: Q3 590
- CHEMICAL REACTION KINETICS: Q3,Q4
676
- CHEMICAL REACTION KINETICS: Q9 367
- CHEMICAL REACTION YIELD 356
- CHEMICAL REACTION YIELD: Q2 54,
301
- CHEMICAL REACTION YIELD: Q4 120
- CHEMICAL REACTIONS 105, 106, 118,
124, 174, 194, 235, 247, 257,
314, 354, 380, 385, 423, 427,
467, 493, 516, 528, 530, 531,
554, 556, 572, 628, 629, 638, 646
- CHEMICAL REACTIONS: Q1 81
- CHEMICAL REACTIONS: Q1,Q2 206,
496, 540
- CHEMICAL REACTIONS: Q1,Q2,Q3 35,
113, 120, 281, 286, 494
- CHEMICAL REACTIONS: Q1,Q2,Q3,Q4 26
- CHEMICAL REACTIONS: Q1,Q2,Q3,Q4,Q5
91, 139
- CHEMICAL REACTIONS: Q1,Q2,Q3,Q4
555
- CHEMICAL REACTIONS: Q1,Q3 400
- CHEMICAL REACTIONS: Q2 304, 500
- CHEMICAL REACTIONS: Q2,Q3,Q4,Q5,Q6,
Q7 202

SUBJECT DESCRIPTORS INDEX

CHEMICAL REACTIONS: Q4	80	CHROMIUM COMPOUNDS	176
CHEMICAL SHIFT	488	CHROMIUM COMPOUNDS: M1	122
CHEMICAL SHIFT: Q1	363	CHROMIUM COMPOUNDS: T1	123
CHEMICAL STATE: Q1,Q2,Q3	40	CHROMIUM FLUORIDES: T9	729
CHEMILUMINESCENCE	241	CHROMIUM OXIDES: T1	35
CHEMISORPTION	135	CHROMIUM-MOLYBDENUM STEELS: M1	471
CHEMISTRY	438, 723, 730	CHROMIUM-MOLYBDENUM STEELS: T1	580
CHEMISTRY: Q1,Q8	176	CHROMIUM-MOLYBDENUM STEELS: T3	527
CHLORATES	111	CHROMIUM: T1	610
CHLORIDES	34, 380, 426, 447, 465, 489, 562, 674	CHROMIUM: T20	711, 712
CHLORIDES: M1	16	CHROMIUM: T24	713
CHLORINATION	81	CHROMIUM: T28	714
CHLORINATION: Q1	356	CHROMIUM: T3	687
CHLORINE	433	CHROMIUM: T7	104, 115
CHLORINE COMPOUNDS	111	CIRCULAR CONFIGURATION	502
CHLORINE 35: T1	31	CLATHRATES	627
CHLORINE: T5	711, 712	CLOSURES	569
CHLORINE: T6	714	CLUSTER MODEL	198
CHLORINE: T9	713	COALESCENCE	685
CHROMATES: T2	35, 375	COATED FUEL PARTICLES: Q1,T2	315, 470
CHROMATES: T4	117	COATED FUEL PARTICLES: T1	334
CHROMATES: T5	116	COATED FUEL PARTICLES: T8	722
CHROMIUM	422, 531	COATINGS	25, 315
CHROMIUM ADDITIONS	313, 324	COBALT	422
CHROMIUM ALLOYS: T1	24	COBALT ALLOYS: T2	11
CHROMIUM ALLOYS: T3	172	COBALT CARBIDES: T3	165
CHROMIUM BASE ALLOYS	531	COBALT CHLORIDES: T2	98, 200
CHROMIUM CHLORIDES: T1	200	COBALT COMPLEXES: T7	314

SUBJECT DESCRIPTORS INDEX

- COBALT FLUORIDES 38
- COBALT: T18 712
- COBALT: T19 711
- COBALT: T23 713
- COBALT: T26 714
- COBALT: T6 104, 687
- COLD TRAPS: Q1 724
- COLOR CENTERS 491
- COLOR: Q1,Q2,Q3,Q4,Q5,Q6,Q7,Q8 191
- COMBUSTION PROPERTIES 722
- COMFACTS 440
- COMPARATIVE EVALUATIONS 29, 69, 73, 147, 190, 196, 233, 280, 357, 359, 397, 409, 418, 453, 460, 497, 511, 589, 596, 609, 639, 653, 658, 705, 706, 727
- COMPARATIVE EVALUATIONS: Q2 309
- COMPATIBILITY 25, 71, 311, 480, 481, 648, 650
- COMPATIBILITY: Q1,Q2,Q3,Q4 644
- COMPATIBILITY: Q1,Q2,Q3,Q4,Q5,Q6, Q7,Q8,Q9,Q10 115
- COMPATIBILITY: Q5 311
- COMPLEXES 27, 113, 142, 170, 268, 488, 516, 723
- COMPOSITE MATERIALS 569
- COMPRESSIBILITY 460, 606
- COMPRESSIBILITY: Q1,Q2,Q3 605
- COMPRESSIBILITY: Q1,Q2,Q3,Q4,Q5,Q6 403
- COMPTON EFFECT 48
- COMPTON SPECTROMETERS 48
- COMPUTER CALCULATIONS 19, 73, 98, 138, 259, 517, 601
- COMPUTER CALCULATIONS: Q2 552
- COMPUTER CODES 431
- COMPUTER CODES: T3 272
- CONTAINERS 118, 649
- CONTAINERS: Q1 111
- CONTAINMENT SHELLS: M2,Q1 569
- CONTAINMENT SYSTEMS 722
- CONTINUITY EQUATIONS 502
- CONTROL 725
- CONTROL ELEMENTS 722
- CONTROL SYSTEMS 94
- CONTROLLED ATMOSPHERES 164, 626
- CONVERSION 233, 516
- CONVEYORS 722
- COOLANTS 25, 590
- COOLANTS: Q1 585
- COOLING 22
- COORDINATION NUMBER 29, 138, 197, 419, 687, 688
- COORDINATION NUMBER: T7 562
- COPPER 533
- COPPER ALLOYS 211
- COPPER ALLOYS: T1 182, 290, 291, 693
- COPPER CHLORIDES 727
- COPPER CHLORIDES: T1 186
- COPPER COMPOUNDS: T2 314
- COPPER FLUORIDES 38

SUBJECT DESCRIPTORS INDEX

COPPER OXIDES: T1 634
 COPPER: T1 417, 567
 COPPER: T19 712
 COPPER: T21 711
 COPPER: T25 713
 COPPER: T27 714
 COPPER: T8 104
 COPRECIPITATION 447
 CORE CATCHERS 367
 CORIOLIS FORCE 285
 CORIUM: T1 454
 CORRELATION FUNCTIONS 210, 567, 661
 CORRELATIONS 75, 112, 189, 355, 504, 511, 609, 643, 653
 CORROSION 35, 64, 169, 309, 315, 692
 CORROSION INHIBITORS 25
 CORROSION PRODUCTS 324
 CORROSION PROTECTION 25
 CORROSION RESISTANCE: Q2, Q3, Q4 311
 CORROSION: Q1 324
 CORROSION: Q1, Q2 128
 CORROSION: Q1, Q2, Q3, Q4, Q5, Q6 580
 CORROSION: Q1, Q3, Q4, Q5, Q6, Q7, Q8, Q9, Q10, Q11, Q12, Q13, Q14, Q15, Q16, Q17, Q18, Q19, Q20, Q21, Q22, Q23 439
 CORROSION: Q2, Q3, Q4, Q5 628
 CORROSION: Q3 25
 CORROSIVE EFFECTS 276, 692
 CORROSIVE EFFECTS: Q1 25, 628
 CORROSIVE EFFECTS: Q10, Q11 335
 CORROSIVE EFFECTS: Q2 324, 439
 CORROSIVE EFFECTS: Q2, Q3 604
 CORROSIVE EFFECTS: Q3 128
 CORROSIVE EFFECTS: Q7, Q8 580
 CORUNDUM: T2 317
 CORUNDUM: T3 628
 COULOMB FIELD 189
 COUNTING RATES 534
 COUPLING 230, 492
 COUPLING CONSTANTS 677
 COVALENCE 280, 398, 571
 CRACKS 211, 400, 580, 719
 CREEP 580
 CRITICAL PRESSURE 180
 CRITICAL PRESSURE: Q2 56
 CRITICAL TEMPERATURE 180, 376, 442
 CRITICAL TEMPERATURE: Q2 56
 CRITICALITY 455
 CROSS SECTIONS 4, 48, 631
 CRUCIBLES: T9 531
 CRUSHING: Q2 722
 CRYSTAL DEFECTS 11, 267, 389, 441, 448, 543
 CRYSTAL DEFECTS: Q1 611
 CRYSTAL FIELD 13, 14, 161, 562, 662, 678, 684
 CRYSTAL GROWTH 108, 695
 CRYSTAL GROWTH: Q1, Q2, Q3, Q4, Q5, Q6 95

SUBJECT DESCRIPTORS INDEX

- CRYSTAL LATTICES 30, 46, 209, 228, 231, 411, 459, 626
- CRYSTAL MODELS 133, 442
- CRYSTAL STRUCTURE 114, 185, 217, 326, 338, 357, 359, 392, 398, 421, 422, 448, 509, 561, 609, 612, 630, 641, 646, 656
- CRYSTAL STRUCTURE: Q1 197, 378, 425, 674
- CRYSTAL STRUCTURE: Q1,Q2 29, 194, 278, 362, 438, 514, 596, 653, 689
- CRYSTAL STRUCTURE: Q1,Q2,Q3 173, 207, 634, 657
- CRYSTAL STRUCTURE: Q1,Q2,Q3,Q4 177
- CRYSTAL STRUCTURE: Q4 363
- CRYSTAL-PHASE TRANSFORMATIONS 560, 611, 650, 657
- CRYSTAL-PHASE TRANSFORMATIONS: Q2 31
- CRYSTAL-PHASE TRANSFORMATIONS: Q1 108
- CRYSTAL-PHASE TRANSFORMATIONS: Q1, Q2 421
- CRYSTALLIZATION 86, 108, 115, 117, 147, 162, 173, 254, 257, 270, 351, 434, 457, 517, 518, 520, 521, 544, 631, 636, 638, 643, 659
- CRYSTALLIZATION: Q1,Q2,Q3,Q4 637
- CRYSTALLOGRAPHY 20, 379
- CRYSTALS 13, 14, 26, 51, 52, 178, 296, 317, 330, 374, 589
- CUBIC LATTICES 379, 409, 611
- CURIE-WEISS LAW 279
- CURIUM 188
- CURIUM COMPLEXES: T1 34
- CURIUM OXIDES: M12 662
- CURIUM: T7 683
- CURRENT DENSITY 430, 450, 451, 515
- CYANIDES 314
- CZECHOSLOVAKIA 358
- CZECHOSLOVAKIA: T1 349
- *** D ***
- D STATES 28, 161
- DATA 12, 73, 79, 103, 141, 280, 283, 453, 454, 508, 538, 705, 717
- DATA COMPILATION 213, 577, 651
- DATA COMPILATION: Q1,Q2 276
- DATA COMPILATION: Q5 281
- DATA: Q1,Q2,Q3 192
- DE HAAS-VAN ALPHEN EFFECT 189
- DEBYE TEMPERATURE 126, 243, 414, 415
- DEBYE-SCHERRER METHOD 561
- DECARBONIZATION 580
- DECLADDING 135
- DECOMPOSITION 399, 581
- DECOMPOSITION: Q2 176, 345
- DECONTAMINATION 724
- DEFECTS 669
- DEFORMATION 175, 601, 692
- DEGASSING 311, 532
- DEGASSING: Q1 588
- DENDRITES 108
- DENSITY 9, 21, 24, 141, 173, 205, 226, 267, 277, 381, 397, 403,

SUBJECT DESCRIPTORS INDEX

DENSITY 422, 440, 460, 573, 577,
578, 611, 651, 681, 691, 710

DENSITY: Q1 599

DENSITY: Q1,Q2 142

DENSITY: Q1,Q2,Q3,Q4 1, 177, 579

DENSITY: Q1,Q2,Q3,Q4,Q5 112

DENSITY: Q6 176

DEPOSITION 324

DESIGN 23, 37, 94, 118, 119, 204,
343, 731

DESIGN: Q2 569

DESIGN: Q7 722

DESORPTION 588

DESTRUCTIVE TESTING 453

DEUTERIUM 529

DEUTERIUM: T4 695

DEUTERIUM: T5 663

DIAGRAMS 3, 63, 64, 73, 98, 101,
190, 359, 402, 478, 508, 511,
541, 653, 700

DIELECTRIC PROPERTIES 656

DIFFERENTIAL THERMAL ANALYSIS 21,
86, 87, 88, 117, 139, 143, 160,
162, 168, 270, 288, 317, 336,
361, 443, 468, 493, 494, 495,
496, 506, 516, 518, 519, 520,
540, 544, 547, 553, 554, 555,
556, 572, 593, 602, 630, 635,
638, 639, 640, 659, 666

DIFFERENTIAL THERMAL ANALYSIS: Q1,
Q2 4, 149, 305

DIFFERENTIAL THERMAL ANALYSIS: Q1,
Q2,Q3 636

DIFFRACTION METHODS 549

DIFFRACTION MODELS 346

DIFFUSION 101, 108, 128, 137, 156,
159, 175, 263, 267, 277, 292,
315, 448, 533, 588

DIFFUSION: Q1 10, 408

DIFFUSION: Q1,Q2 441, 669

DIFFUSION: Q2 401, 693

DIFFUSION: Q2,Q3 651

DIGITAL COMPUTERS 517

DILATOMETRY 508

DILUTION 6, 407

DIMERS 17

DIPOLE MOMENTS 492

DIPOLES: Q1 410

DIRAC APPROXIMATION 223

DISLOCATIONS 560

DISPERSION HARDENING 5

DISPERSION RELATIONS 485

DISPERSIONS 710

DISPERSIONS: Q1 452

DISSOCIATION 124, 295, 354, 395

DISSOCIATION ENERGY 240, 241

DISSOCIATION ENERGY: Q1,Q2,Q3,Q4,Q5
239, 597

DISSOLUTION 64, 306, 317, 347,
349, 450, 600, 675, 690

DISSOLUTION: Q1 455

DISSOLUTION: Q2,Q3 156

DISSOLUTION: Q4 94

DISSOLVERS 722

DISTILLATION 18

DISTRIBUTION 514, 609, 653

SUBJECT DESCRIPTORS INDEX

DISTRIBUTION FUNCTIONS 195, 248,
307, 308, 369, 370, 661, 688

DISTRIBUTION FUNCTIONS: Q1 426

DMSO 119, 369, 381

DOPED MATERIALS 30, 130, 193, 261,
363, 366, 389, 432, 557, 610, 718

DRAG 19

DROPLETS 588

DUSTS 722

DYSPROSIUM 399, 418

DYSPROSIUM ALLCYS: T2 399

DYSPROSIUM CHLORIDES: T1 29, 112

DYSPROSIUM COMPOUNDS: T6 82

DYSPROSIUM FLUORIDES: T1 603

DYSPROSIUM FLUORIDES: T2 21

DYSPROSIUM FLUORIDES: T4 484

DYSPROSIUM FLUORIDES: T8 382

DYSPROSIUM HYDRIDES: T3 418

DYSPROSIUM OXIDES: T5 427

DYSPROSIUM: T2 125

*** E ***

EER-2 REACTOR: T1 416

EFFICIENCY 269, 534, 539, 724

EFFICIENCY: Q2 430

EIGENVALUES 683

EINSTEINIUM 188

EINSTEINIUM FLUORIDES: T4 501

EINSTEINIUM CHLORIDES: T3 501

ELASTICITY: Q1 368

ELECTRIC ARCS 588

ELECTRIC BATTERIES 40, 96

ELECTRIC BATTERIES: T1 42, 274,
275, 277, 282, 380

ELECTRIC BATTERIES: T2 246

ELECTRIC CONDUCTIVITY 23, 164,
247, 248, 260, 263, 277, 321,
346, 381, 448, 486, 528, 529,
568, 595, 643, 651, 703

ELECTRIC CONDUCTIVITY: Q1 302

ELECTRIC CONDUCTIVITY: Q1,Q2 389

ELECTRIC CONDUCTIVITY: Q1,Q2,Q3,Q4,
Q5 112

ELECTRIC CONDUCTIVITY: Q1,Q2,Q3
193, 224

ELECTRIC CONDUCTIVITY: Q1,Q2,Q3,Q4,
Q5 461

ELECTRIC CONDUCTIVITY: Q1,Q3,Q4,Q5,
Q6,Q7,Q8 191

ELECTRIC CURRENTS 38

ELECTRIC DIPOLES 492

ELECTRIC FIELDS 511

ELECTRIC IMPEDANCE 568

ELECTRIC MOMENTS 538

ELECTRIC POTENTIAL 38, 269, 335,
402

ELECTRIC POTENTIAL: Q1 187

ELECTRIC-POWERED VEHICLES 23

ELECTRICAL PROPERTIES 215, 398,
507, 595, 612

ELECTRICAL PROPERTIES: Q1,Q2,Q3
568

ELECTROCHEMICAL CELLS 282

SUBJECT DESCRIPTORS INDEX

- ELECTROCHEMICAL CELLS: T1 568
 ELECTROCHEMICAL CORROSION 652
 ELECTROCHEMICAL CORROSION: Q1,Q2,
 Q3,Q4,Q5,Q6,Q7,Q8,Q9 335
 ELECTROCHEMICAL MACHINING: Q1 129
 ELECTROCHEMISTRY 74, 176, 275,
 380, 439, 507, 634, 693, 703
 ELECTROCHEMISTRY: Q1 111, 221,
 263, 293, 450, 727
 ELECTROCHEMISTRY: Q1,Q2 721
 ELECTROCHEMISTRY: Q1,Q2,Q3 675
 ELECTROCHEMISTRY: Q2 277, 451
 ELECTROCHEMISTRY: T 63
 ELECTRODEPOSITION 111, 515, 727
 ELECTRODEPOSITION: Q1 430
 ELECTRODES 68, 193, 234, 263, 274,
 583, 658, 721
 ELECTRODES: Q1 380, 720
 ELECTRODES: T10 729
 ELECTROLYSIS 91, 221, 263, 450,
 451, 703
 ELECTROLYSIS: Q1,Q2 234
 ELECTROLYTES 38, 64, 91, 129, 193,
 542, 586, 634, 652, 658, 704, 727
 ELECTROLYTES: Q1 42, 77, 116, 277,
 282, 380, 381, 507
 ELECTROLYTES: Q1,Q2 246
 ELECTROLYTES: Q2,Q3 263
 ELECTROLYTES: Q3 234, 389, 563
 ELECTROLYTES: Q4 595
 ELECTROLYTIC CELLS 32, 63, 68,
 200, 291, 513, 542, 652
 ELECTROLYTIC CELLS: T1 535
 ELECTROLYTIC CELLS: T2 430
 ELECTROMAGNETIC INTERACTIONS 677
 ELECTROMETALLURGY 703
 ELECTROMETALLURGY: T1,Q2 76
 ELECTROMOTIVE FORCE 6, 40, 50, 63,
 64, 96, 163, 182, 193, 200, 221,
 291, 293, 314, 339, 372, 388,
 406, 407, 420, 436, 522, 583,
 652, 658, 703, 704
 ELECTROMOTIVE FORCE: Q1 535
 ELECTROMOTIVE FORCE: Q1,Q2 32, 551
 ELECTRON BEAMS 132
 ELECTRON DENSITY 161, 346
 ELECTRON DIFFRACTION 514
 ELECTRON EXCHANGE 186
 ELECTRON MICROPROBE ANALYSIS 169,
 468
 ELECTRON PROBES 440, 669
 ELECTRON SPECTRA 674
 ELECTRON SPECTROSCOPY 113
 ELECTRON SPIN RESONANCE 409, 432
 ELECTRON TRANSFER 193, 412, 562
 ELECTRONEGATIVITY: Q1 429
 ELECTRONIC SPECIFIC HEAT 126, 209,
 243
 ELECTRONIC STRUCTURE 30, 53, 61,
 81, 164, 165, 190, 220, 259, 320,
 398, 412, 413, 417, 491, 492,
 499, 562, 672
 ELECTRONIC STRUCTURE: Q1 28, 157,
 188, 205, 280, 485, 684, 686, 705
 ELECTRONIC STRUCTURE: Q1,Q2 13,
 122, 161, 189, 223, 411, 460,
 589, 673
 ELECTRONIC STRUCTURE: Q1,Q2,Q3 14,

SUBJECT DESCRIPTORS INDEX

ELECTRONIC STRUCTURE: Q1,Q2,Q3 48

ELECTRONIC STRUCTURE: Q1,Q2,Q3,Q4,
Q5,Q6 215ELECTRONIC STRUCTURE: Q1,Q2,Q3,Q4,
Q5,Q6,Q7,Q8,Q9 230ELECTRONIC STRUCTURE: Q1,Q2,Q3,Q4,
Q5 249ELECTRONIC STRUCTURE: Q1,Q2,Q3,Q4,
Q5,Q6 323ELECTRONIC STRUCTURE: Q1,Q2,Q3,Q4
598ELECTRONIC STRUCTURE: Q1,Q2,Q3,Q4,
Q5,Q6,Q7,Q8,Q9,Q10,Q11,Q12,Q13,Q14
662

ELECTRONIC STRUCTURE: Q1,Q2,Q3 678

ELECTRONIC STRUCTURE: Q1,Q2,Q3,Q4,
Q5,Q6,Q7 683ELECTRONIC STRUCTURE: Q1,Q2,Q3,Q4,
Q5,Q6,Q7,Q8,Q9 688

ELECTRONS 48, 157, 205, 318, 705

ELECTROREFINING: Q1 703

ELEMENTS 723

ELONGATION 643

EMBRITTLEMENT 649

EMBRITTLEMENT: Q1 211, 212

EMBRITTLEMENT: Q2 25

EMISSION SPECTRA 44, 132, 409,
489, 491, 538, 662

EMISSION SPECTRA: Q1 374, 718

EMISSION SPECTROSCOPY: Q1 319, 566

EMISSION SPECTROSCOPY: Q1,Q2,Q3,Q4,
Q5,Q6,Q7,Q8,Q9,Q10,Q11,Q12,Q13,Q14
125EMISSION SPECTROSCOPY: Q14,Q15,Q16,
Q17,Q18,Q19,Q20,Q21,Q22,Q23,Q24,
Q25,Q26,Q27,Q28,Q29,Q30,Q31,Q32,
Q33,Q34,Q35,Q36,Q37,Q38,Q39,Q40,
Q42,Q43 713EMISSION SPECTROSCOPY: Q2,Q3,Q4,Q5,
Q6,Q7,Q8,Q9,Q10,Q11,Q12,Q13,Q14,
Q15,Q16,Q17,Q18,Q19,Q20,Q21 104EMISSION SPECTROSCOPY: Q5,Q11,Q12,
Q13,Q14,Q15,Q16,Q17,Q18,Q19,Q20,
Q21,Q22,Q23,Q24,Q25,Q26,Q27,Q28,
Q30,Q31,Q32,Q33,Q34,Q35,Q36,Q37,
Q38,Q39,Q4 714EMISSION SPECTROSCOPY: Q7,Q8,Q9,
Q10,Q11,Q12,Q13,Q14,Q15,Q16,Q17,
Q18,Q19,Q20,Q21,Q22,Q23,Q24,Q25,
Q26,Q27,Q28,Q29,Q30,Q31,Q32,Q33,
Q34,Q35,Q36, 712EMISSION SPECTROSCOPY: Q8,Q9,Q11,
Q12,Q13,Q14,Q15,Q16,Q17,Q18,Q19,
Q20,Q21,Q22,Q23,Q24,Q25,Q26,Q27,
Q28,Q29,Q30,Q31,Q32,Q33,Q34,Q35,
Q36,Q37,Q40 711

EMISSIVITY 121, 577

EMISSIVITY: Q1 132

EMISSIVITY: Q1,Q2,Q3,Q4,Q5 47

ENDOR: Q2 130

ENERGY 704

ENERGY GAP 398

ENERGY LEVELS 13, 14, 220, 223,
538, 589, 662

ENERGY LEVELS: Q1,Q2,Q3 664

ENERGY SPECTRA 161, 412

ENERGY TRANSFER: Q1 621

ENERGY TRANSFER: Q1,Q2,Q3,Q4,Q5,Q6
539

ENERGY-LEVEL DENSITY: Q1 422

ENERGY-LEVEL TRANSITIONS 16, 17,
413, 432, 484, 491, 511, 562

SUBJECT DESCRIPTORS INDEX

ENERGY-LEVEL TRANSITIONS: Q1 492,
538

ENGINEERING 94

ENRICHED URANIUM 226

ENTHALPY 73, 82, 101, 124, 133,
180, 199, 222, 243, 244, 251,
261, 262, 264, 281, 288, 297,
310, 316, 334, 340, 343, 347,
354, 367, 372, 386, 391, 410,
442, 467, 528, 529, 532, 560,
574, 577, 578, 591, 599, 616

ENTHALPY: Q1 184, 355, 586

ENTHALPY: Q1,Q2 179, 201, 383,
619, 620

ENTHALPY: Q1,Q2,Q3 166

ENTHALPY: Q1,Q2,Q3,Q4,Q5 717

ENTHALPY: Q2 78

ENTRAINMENT 135

ENTROPY 3, 9, 24, 36, 118, 133,
149, 238, 261, 262, 281, 286,
310, 332, 333, 343, 347, 367,
372, 386, 391, 406, 414, 422,
441, 442, 504, 513, 523, 524,
532, 560, 584, 591, 594, 647, 705

ENTROPY: Q1 184, 260, 466

ENTROPY: Q1,Q2 291, 383

ENTROPY: Q1,Q2,Q3 138

ENTROPY: Q1,Q2,Q3,Q4,Q5 717

ENTROPY: Q1,Q2,Q3,Q4,Q5,Q6,Q7,Q8,Q9
230

ENTROPY: Q2 78

EPITAXY 95

EQUATIONS 3, 6, 22, 36, 93, 96,
97, 98, 108, 190, 242, 284, 391,
420, 472, 517, 546, 564, 581,
588, 606, 646

EQUATIONS OF STATE 60, 136, 310,
397, 422, 537, 577

EQUATIONS OF STATE: Q1 180

EQUATIONS OF STATE: Q1,Q2 578

EQUATIONS OF STATE: Q1,Q2,Q3 584

EQUILIBRIUM 1, 4, 40, 54, 63, 64,
73, 87, 88, 109, 176, 238, 245,
314, 342, 385, 394, 420, 445,
446, 449, 450, 451, 517, 646,
648, 673

EQUILIBRIUM: Q1,Q2 457

EQUILIBRIUM: Q1,Q2,Q3,Q4 92

ERBIUM CHLORIDES: T5 497

ERBIUM COMPOUNDS: T2 123

ERBIUM COMPOUNDS: T5 639

ERBIUM COMPOUNDS: T6 196

ERBIUM COMPOUNDS: T8 82

ERBIUM FLUORIDES: T1 270

ERBIUM FLUORIDES: T10 382

ERBIUM IONS 696

ERBIUM 169: T2 696

ERBIUM: T3 125

ERRORS 163, 502

ETHANOL 347

ETHERS 111, 118

EUROPIUM 399

EUROPIUM ALLOYS: T3 399

EUROPIUM CHLORIDES: M1 361

EUROPIUM CHLORIDES: T1 594

EUROPIUM CHLORIDES: T2 607

EUROPIUM CHLORIDES: T3 14

EUROPIUM COMPOUNDS: T1 589

EUROPIUM FLUORIDES 307

SUEJECT DESCRIPTORS INDEX

- EUROPIUM FLUCHRIDES: T2 419
- EUROPIUM FLUCHRIDES: T4 241
- EUROPIUM IONS: T1 432
- EUROPIUM NITRATES: T1 165
- EUROPIUM OXIDES: T1 607
- EUROPIUM OXIDES: T3 50
- EUROPIUM OXIDES: T4 239
- EUROPIUM 154 75
- EUROPIUM: T2 224, 230, 688
- EUROPIUM: T4 125
- EUTECTICS 40, 68, 86, 89, 91, 92,
108, 116, 117, 139, 151, 177,
196, 199, 233, 254, 256, 267,
270, 274, 275, 286, 292, 293,
300, 307, 308, 314, 317, 351,
369, 370, 371, 377, 406, 407,
408, 430, 449, 452, 469, 496,
506, 507, 515, 517, 518, 519,
540, 544, 545, 548, 553, 554,
555, 575, 590, 602, 605, 636,
637, 638, 659, 668, 675, 706, 721
- EUTECTICS: T1 276
- EVAPORATION 54, 55, 281, 297, 391,
445, 448
- EVAPORATION: Q1 73, 482
- EVAPORATION: Q1,Q2 251, 476, 479,
481
- EXCHANGE INTERACTIONS 376
- EXCITATION 491, 539
- EXCITATION: Q1,Q2,Q3,Q4,Q5,Q6 183
- EXCITED STATES 280, 353, 374, 411,
562
- EXCITONS 411
- EXCURSIONS: Q1 60, 72
- EXPERIMENT PLANNING: T 546
- EXPLOSIONS: Q1,Q2 590
- EXTRACTION APPARATUSES 237
- EXTRACTION APPARATUSES: T1 236
- EXTRACTION APPARATUSES: T2 79
- EXTRACTION CHROMATOGRAPHY 723
- EXTRUSION 710
- *** F ***
- F CODES 310
- F STATES 28, 189, 280, 318, 323,
492
- FABRICATION 38, 111, 118, 381
- FABRICATION: Q1 226
- FABRICATION: Q1,Q2 691
- FABRICATION: Q10 729
- FABRICATION: Q2 536
- FAILURES 211
- FAR ULTRAVIOLET RADIATION 382, 484
- FAST NEUTRONS 225, 316
- FAST REACTORS 175, 644
- FATIGUE 580
- FATIGUE: Q1,Q2 692
- FBR TYPE REACTORS 477
- FBR TYPE REACTORS: T1 60
- FBR TYPE REACTORS: T3 135
- FCC LATTICES 177, 198, 411, 442
- FEASIBILITY STUDIES 85, 246
- FEASIBILITY STUDIES: Q1 704
- FERMI LEVEL 48, 52, 161, 189, 280,

SUBJECT DESCRIPTORS INDEX

FERMI LEVEL 346, 422, 560, 598
 FERMIUM 188
 FERROMAGNETISM 157
 FFTF REACTOR 416
 FIBERS 710
 FIELD THEORIES 376
 FILMS 25, 357, 588, 619
 FINE STRUCTURE: Q1,Q2,Q3 432
 FINITE DIFFERENCE METHOD 310
 FISSILE MATERIALS: T2 131
 FISSION 619
 FISSION PRODUCT RELEASE: Q2 315
 FISSION PRODUCTS 69, 75, 120, 169,
 175, 225, 226, 315, 453, 470,
 534, 685, 716
 FISSION PRODUCTS: Q1 202
 FISSION PRODUCTS: T1 307
 FISSION PRODUCTS: T3 281
 FISSION RATIO 620
 FISSION YIELD 214
 FISSIONABLE MATERIALS: T2 690
 FLASH HEATING 366
 FLEXIBILITY 710
 FLEXURAL STRENGTH 267
 FLOW BLOCKAGE: Q2 19
 FLOWSHEETS 699
 FLOWSHEETS: Q4 722
 FLUID FLOW 103, 158
 FLUIDIZED BED 158, 699, 700, 724,
 725
 FLUIDIZED-BED COMBUSTION: Q8,Q2
 722
 FLUIDIZED-BED COMBUSTORS: T9 722
 FLUORESCENCE 539, 718
 FLUORESCENCE SPECTROSCOPY 233, 538
 FLUORESCENCE: Q1,Q2 233, 491
 FLUORIDE VOLATILITY PROCESS 69,
 70, 135, 222, 311, 358, 699, 700,
 725
 FLUORIDE VOLATILITY PROCESS: T 558
 FLUORIDE VOLATILITY PROCESS: T1,Q2,
 Q3 724
 FLUORIDES 41, 329, 380, 427
 FLUORIDES: M2 16
 FLUORIDES: T1 463
 FLUORINATION 222, 237
 FLUORINATION: Q1 558, 699
 FLUORINATION: Q1,Q2 533
 FLUORINATION: Q2 700
 FLUORINATION: Q2,Q3 725
 FLUORINATION: T1 500
 FLUORINE 558, 699, 700
 FLUORINE COMPOUNDS: A5 176
 FLUORINE COMPOUNDS: T6 66
 FLUORINE 19 21, 463
 FLUORINE 19: T2 363
 FLUORINE: T10 713
 FLUORINE: T2 500
 FLUORINE: T4 712
 FLUORINE: T6 711
 FLUORINE: T7 714

SUEJECT DESCRIPTORS INDEX

FLUOROBORATES: T2 530
 FOCUSING 208, 331
 FORMATES 562
 FORMATION FREE ENERGY 36, 182, 199, 291, 428, 581, 634, 645, 647, 702
 FORMATION FREE ENERGY: Q1 184
 FORMATION FREE ENERGY: Q1,Q2 375
 FORMATION FREE ENERGY: Q4,Q5 646
 FORMATION FREE ENTHALPY 74, 83, 133
 FORMATION FREE ENTHALPY: Q1 407
 FORMATION FREE ENTHALPY: Q1,Q2 291, 326
 FORMATION HEAT 9, 36, 137, 240, 241, 332, 333, 341, 406, 409, 427, 428, 429, 499, 500
 FORMATION HEAT: Q1 65, 306, 425, 466, 467
 FORMATION HEAT: Q1,Q2 105, 238, 290, 543, 607, 608, 635
 FORMATION HEAT: Q1,Q2,Q3 166, 603
 FORMATION HEAT: Q1,Q2,Q3,Q4 137, 178
 FORMATION HEAT: Q1,Q2,Q3,Q4,Q5 245, 717
 FORMATION HEAT: Q1,Q3 206
 FORMATION HEAT: Q9,Q10 82
 FOURIER ANALYSIS 687, 688
 FOURIER TRANSFORMATION 159, 210
 FRACTURE PROPERTIES 211, 440, 580
 FRACTURE PROPERTIES: Q1,Q2 692
 FRACTURES 211, 212
 FREE ENERGY 5, 24, 54, 118, 134, 167, 180, 187, 206, 314, 342, 400, 496, 550, 584, 599, 650, 654
 FREE ENERGY: Q1,Q2,Q3 402
 FREE ENTHALPY 3, 110, 149, 279, 399, 406, 523, 644
 FREE ENTHALPY: Q1,Q2 383
 FREE ENTHALPY: Q2 78
 FREQUENCY DEPENDENCE 497
 FREQUENCY MEASUREMENT 185
 FREQUENCY RANGE 185
 FRICTION 502
 FRICTION: Q1,Q2,Q3,Q4 575
 FUEL ASSEMBLIES 75
 FUEL CANS 19, 169, 510, 719
 FUEL CANS: T1 175
 FUEL CELLS: T1 246
 FUEL CYCLE 232, 237, 690, 699, 700
 FUEL CYCLE: Q1 722
 FUEL CYCLE: Q2,Q3 455
 FUEL ELEMENTS: Q1 75
 FUEL ELEMENTS: T,Q1,Q2 509
 FUEL ELEMENTS: T1 214
 FUEL ELEMENTS: T3 715
 FUEL FABRICATION PLANTS 455
 FUEL FABRICATION PLANTS: M2 70
 FUEL PELLETS 226, 510, 711, 712, 713, 719
 FUEL PINS 72, 169, 175, 316, 416
 FUEL PINS: T2 456
 FUEL REPROCESSING PLANTS 455

SUBJECT DESCRIPTORS INDEX

FUEL REPROCESSING PLANTS: T1 131
 FUEL REPROCESSING PLANTS: T2 94
 FUEL REPROCESSING PLANTS: T5 722
 FUEL RODS 510
 FUEL SCANNING 131, 349
 FUEL-CLADDING INTERACTIONS 19,
 510, 644, 650
 FUEL-CLADDING INTERACTIONS: M2,Q1
 719
 FUEL-CLADDING INTERACTIONS: Q1,Q2
 169
 FUEL-COOLANT INTERACTIONS: T,Q1
 272
 FUNCTIONAL MODELS 474
 FURANS 194
 FURNACES 204, 208, 331
 FUSION HEAT 243, 272, 286, 475,
 477, 574

*** G ***

GADOLINIUM 399, 598
 GADOLINIUM ALLOYS: T3 598
 GADOLINIUM ALLOYS: T4 399
 GADOLINIUM BORIDES: T4 215
 GADOLINIUM CHLORIDES: T2 112, 594
 GADOLINIUM CHLORIDES: T4 497
 GADOLINIUM COMPOUNDS: T5 82
 GADOLINIUM FLUORIDES: T2 41, 603
 GADOLINIUM FLUORIDES: T5 484
 GADOLINIUM FLUORIDES: T6 382
 GADOLINIUM HYDRIDES: T1 418

GADOLINIUM IONS 130, 410
 GADOLINIUM OXIDES 366
 GADOLINIUM OXIDES: T2 266, 697
 GADOLINIUM OXIDES: T4 427
 GADOLINIUM: T1 370
 GADOLINIUM: T3 230, 688
 GADOLINIUM: T5 125
 GALLIUM 4, 328, 369
 GALLIUM ALLOYS 544, 545
 GALLIUM ALLOYS: A3 651
 GALLIUM ALLOYS: T1 513, 522, 706
 GALLIUM BASE ALLOYS: T1 623
 GALLIUM CHLORIDES: T1 4
 GALLIUM OXIDES: T2 105
 GALLIUM TELLURIDES: T2 544, 545
 GALLIUM: T1 583, 622, 681
 GALLIUM: T10 104
 GALLIUM: T2 327
 GALLIUM: T22 712
 GALLIUM: T30 714
 GALLIUM: T4 544
 GAMMA FUEL SCANNING 75
 GAMMA FUEL SCANNING: T,Q1,Q2 453
 GAMMA RADIATION 61
 GAMMA SPECTRA 453
 GAMMA SPECTROSCOPY 214
 GAMMA SPECTROSCOPY: Q3,Q4 714
 GAS FLOW 722
 GAS FUELED REACTORS: M1 345

SUBJECT DESCRIPTORS INDEX

GASES 2, 78, 180, 192, 281, 452,
499, 597, 685

*** H ***

GERMANIUM 108

H CODES 456

GERMANIUM CHLORIDES: T2 553

HAFNIUM 582

GERMANIUM OXIDES: T1 394

HAFNIUM CARBIDES: T1 508

GETTERING 64

HAFNIUM CHLORIDES: T1 136, 283

GETTERS: Q2 470

HAFNIUM FLUORIDES: T1 330, 596

GLASS 21, 183, 233, 538, 539, 616,
617, 664, 666

HAFNIUM FLUORIDES: T2 541

GLASS: T4 521

HAFNIUM NITRIDES: T2 508

GLOVES: T10 301

HALF-LIFE 487

GOLD ALLOYS: T2 190

HALF-LIFE: Q1 690

GOLD ALLOYS: T6 417

HALIDES 29, 99, 429

GOLD: T11 712

HALIDES: T2 631

GOLD: T19 714

HALL EFFECT 215

GOLD: T3 417

HAMILTONIANS 130, 229

GONIOMETERS 250

HARD-CORE POTENTIAL 689

GRAIN BOUNDARIES 5, 211, 456

HARD-SPHERE MODEL 615

GRAIN BOUNDARIES: Q1, Q2 697

HARDNESS 360, 656

GRAIN GROWTH 267

HARDNESS: Q1, Q3, Q4, Q5, Q6 191

GRAIN SIZE 440, 560

HARTREE-FOCK METHOD 223

GRAPHITE 111, 235, 327, 381, 503,
649, 676, 722

HASTELLOYS: T1 604

GRAPHITE: T2 627

HAZARDS 276

GRAVIMETRIC ANALYSIS: Q2, Q3 711

HEAD END PROCESSES 722

GREEN FUNCTION 161

HEAT STORAGE: T 276

GRINDING 676, 691

HEAT TRANSFER 272, 585, 651

GROUND STATES 230, 678

HEAT TRANSFER: Q1 437, 502

GRUENEISEN CONSTANT 231, 508

HEAT TRANSFER: T1 158

HEAT TREATMENTS 480

HEATING 22, 480

SUBJECT DESCRIPTORS INDEX

HEISENBERG MODEL 368
 HELIUM 345
 HIGH PRESSURE 599, 641
 HIGH TEMPERATURE 1, 6, 10, 22, 23,
 26, 27, 34, 35, 40, 66, 68, 79,
 92, 96, 106, 132, 154, 175, 179,
 182, 210, 224, 242, 243, 256,
 261, 262, 268, 269, 274, 275,
 284, 290, 292, 293, 324, 327,
 339, 340, 360, 385, 386, 388,
 391, 396, 399, 404, 405, 415,
 434, 463, 471, 522, 523, 524,
 529, 540, 558, 573, 575, 586,
 587, 591, 595, 599, 611, 616,
 617, 619, 626, 641, 642, 667,
 679, 693, 701, 702, 704
 HIGH-TEMPERATURE FUEL CELLS: M2
 263
 HIGH-TEMPERATURE FUEL CELLS: T1
 71, 77
 HIGH-TEMPERATURE FUEL CELLS: T3
 389, 563
 HIGH-TEMPERATURE FUEL CELLS: T4
 595
 HIGHLY ENRICHED URANIUM 214
 HOLES 186, 389
 HOLMIUM BORIDES: T6 215
 HOLMIUM CHLORIDES: T1 378, 608
 HOLMIUM FLUORIDES: T1 540
 HOLMIUM FLUORIDES: T2 572
 HOLMIUM FLUORIDES: T9 382
 HOLMIUM: T6 125
 HOT PRESSING 9, 440
 HTGR TYPE REACTORS 334
 HTGR TYPE REACTORS: T1 315, 470,
 722
 HYDRIDATION 587
 HYDRIDES 392, 429
 HYDRIODIC ACID 170
 HYDROCHLORIC ACID 467
 HYDRODYNAMICS 564, 588
 HYDRODYNAMICS: Q2 310
 HYDROFLUORIC ACID 306
 HYDROFLUORIC ACID: T4 541
 HYDROGEN 45, 55, 164, 176, 428,
 588
 HYDROGEN ADDITIONS 418
 HYDROGEN COMPOUNDS: T1 695
 HYDROGEN FUEL CELLS: T1 507
 HYDROGEN: T1 10, 287
 HYDROGEN: T2 22, 301
 HYDROGEN: T3 587
 HYDROGEN: T4 217, 392
 HYDROGEN: T6 663
 HYDROLYSIS: Q1 504
 HYDROLYSIS: Q4, Q5 176
 HYGROSCOPICITY 21
 HYPERFINE STRUCTURE 61, 559
 HYPERFINE STRUCTURE: Q1 130, 696
 HYSTERESIS 643
 *** I ***
 IMPURITIES 44, 46, 104, 125, 211,
 481, 560, 565, 600, 652
 IMPURITIES: Q4 404
 INCOLOY 800: T3 692

SUBJECT DESCRIPTORS INDEX

- INCOLOY 800: T6 580
- INCOMPRESSIBLE FLOW 601
- INDIUM 4, 105, 211, 212
- INDIUM ALLOYS: A2 651
- INDIUM ALLOYS: T1 388
- INDIUM ALLOYS: T2 177, 290, 291, 513, 514, 523, 524, 706
- INDIUM ALLOYS: T3 182, 328
- INDIUM CHLORIDES: T1 93, 174
- INDIUM CHLORIDES: T2 4, 341
- INDIUM OXIDES 182
- INDIUM OXIDES: T1 105
- INDIUM: T1 177, 341
- INDIUM: T11 104
- INDIUM: T2 182, 328
- INDIUM: T23 711, 712
- INDIUM: T3 327
- INDIUM: T31 714
- INELASTIC SCATTERING 564
- INERT ATMOSPHERE 557
- INFRARED SPECTRA 12, 21, 66, 80, 81, 127, 185, 218, 220, 278, 353, 409, 427, 431, 498, 558, 695
- INFRARED SPECTRA: Q1 219, 350, 483
- INFRARED SPECTRA: Q1,Q2 102, 305, 357
- INFRARED SPECTRA: Q1,Q2,Q3,Q4,Q5, Q6,Q7 41
- INFRARED SPECTRA: Q1,Q2,Q3,Q4,Q5,Q6 497
- INFRARED SPECTROMETERS: T 208
- INORGANIC ACIDS: T2 604
- INORGANIC COMPOUNDS 23
- INTEGRALS 502, 564
- INTERACTION RANGE 609, 653
- INTERATOMIC DISTANCES 197, 249, 419, 661, 677, 686, 687, 688
- INTERATOMIC DISTANCES: Q1,Q2 514
- INTERATOMIC FORCES 167, 224, 249, 686
- INTERATOMIC FORCES: Q1 248, 273
- INTERFACES 5, 108, 234, 697
- INTERGRANULAR CORROSION: Q1 604
- INTERMEDIATE STRUCTURE 16, 17
- INTERMETALLIC COMPOUNDS 137, 189, 190, 275, 359, 392, 399, 446, 514, 522, 549, 604, 671
- INTERMOLECULAR FORCES 273
- INTERNAL FRICTION 19
- INTERNATIONAL ORGANIZATIONS 717
- INTERSTITIALS 133, 193, 209, 410, 459, 610
- INTERSTITIALS: Q1,Q2 74
- INVENTORIES 349
- IODIDES 314
- IODIDES: M2 17
- IODIDES: T2 20
- IODINE 722
- IODINE: T3 94
- IODOX PROCESS: T 94
- ION DENSITY 20, 167
- ION EXCHANGE 358, 716
- ION IMPLANTATION 696

SUBJECT DESCRIPTORS INDEX

ION MOBILITY 246
 ION SELECTIVE ELECTRODE ANALYSIS: Q5, Q6 711
 ION SELECTIVE ELECTRODE ANALYSIS: Q4, Q5 712
 ION SELECTIVE ELECTRODE ANALYSIS: Q9, Q10 713
 ION SPECTROSCOPY 412
 IONIC CONDUCTIVITY 23, 246, 389, 568
 IONIC CRYSTALS 197
 IONIZATION 280, 445
 IONIZATION POTENTIAL: Q1, Q2, Q3, Q4, Q5 239
 IONS 46, 167, 273, 408, 412, 511
 IRON 422
 IRON ALLOYS 654, 693
 IRON ALLOYS: M2 671
 IRON ALLOYS: T1 661
 IRON ALLOYS: T2 172
 IRON BASE ALLOYS: T2 24
 IRON CHLORIDES: T1 97, 554, 555
 IRON CHLORIDES: T2 96, 433, 435, 546
 IRON CHLORIDES: T3 627
 IRON COMPLEXES: T8 314
 IRON OXIDES: M2 372
 IRON: T1 128, 313, 575
 IRON: T2 244, 693
 IRON: T21 712
 IRON: T22 711
 IRON: T26 713
 IRON: T5 687, 714
 IRON: T9 104
 IRRADIATION 169, 316, 416
 IRRADIATION: Q2 232
 ISOMER SHIFT 61, 559, 627
 ISOTHERMAL PROCESSES 97, 101, 340, 351
 ISOTOPE DILUTION 131, 690, 716
 ISOTOPE EFFECTS 220, 285, 405
 ISOTOPE EFFECTS: Q1 78
 ISOTOPE EFFECTS: Q2, Q3 353
 ISOTOPE EFFECTS: Q3 219
 ISOTOPE EFFECTS: Q4 695
 ISOTOPE EFFECTS: Q5 663
 ISOTOPE ENRICHED MATERIALS 374
 ISOTOPE PRODUCTION 534
 ISOTOPE RATIO 214, 232, 349, 453
 ISOTOPE SEPARATION 285
 ISOTOPIC EXCHANGE: T1 68

 *** J ***
 JAHN-TELLER EFFECT 53, 596
 JPDR REACTOR: T1 453

 *** K ***
 KERNFORSCHUNGSZENTRUM KARLSRUHE 60
 KINETIC EQUATIONS 101, 401
 KNUDSEN FLOW 120, 399, 481

SUBJECT DESCRIPTORS INDEX

*** L ***

L-S COUPLING 677

LABELLED COMPOUNDS 347

LAGRANGIAN FUNCTION 310

LAMBDA POINT 414

LAMINAR FLOW 502

LANTHANUM 598

LANTHANUM ALICYS: T1 3

LANTHANUM ALICYS: T4 598

LANTHANUM CHLORIDES 34, 130

LANTHANUM CHLORIDES: T1 493

LANTHANUM CHLORIDES: T3 112, 497

LANTHANUM COMPOUNDS 326

LANTHANUM COMPOUNDS: M2 495

LANTHANUM COMPOUNDS: T3 82

LANTHANUM COMPOUNDS: T7 196

LANTHANUM FLUORIDES 307

LANTHANUM FLUORIDES: T1 382, 484

LANTHANUM FLUORIDES: T2 461

LANTHANUM FLUORIDES: T3 41

LANTHANUM FLUORIDES: T4 26

LANTHANUM NITRATES: T1 164

LANTHANUM OXIDES: M1 49

LANTHANUM OXIDES: T1 438, 654

LANTHANUM OXIDES: T6 183

LANTHANUM SULFIDES: T3 597

LANTHANUM TUNGSTATES: T2 338

LANTHANUM 139: T2 130

LANTHANUM: T1 130

LANTHANUM: T4 230, 688

LANTHANUM: T7 125

LASER RADIATION 478

LASER-RADIATION HEATING 474, 476

LASERS 60, 475, 488, 589

LATTICE PARAMETERS 83, 168, 173,
194, 197, 207, 209, 242, 266,
303, 330, 332, 333, 334, 338,
379, 414, 419, 443, 508, 596,
611, 641

LATTICE PARAMETERS: Q1 81

LATTICE PARAMETERS: Q1,Q2,Q3 549

LATTICE PARAMETERS: Q1,Q2,Q3,Q4
671

LATTICE VIBRATIONS 41, 126

LAVES PHASES 137

LAWRENCIUM 188

LCAO METHOD 280

LEAD 42, 211, 212, 328, 346

LEAD ALLOYS 420, 704

LEAD ALLOYS: T2 436, 575, 606

LEAD ALLOYS: T3 605

LEAD ALLOYS: T4 1

LEAD ALLOYS: T6 403

LEAD BROMIDES: T1 602

LEAD CHLORIDES: T1 143

LEAD CHLORIDES: T2 402, 491, 602

LEAD COMPLEXES: T4 314

LEAD COMPOUNDS: M4 168

LEAD FLUORIDES 463

SUBJECT DESCRIPTORS INDEX

LEAD FLUORIDES: T3 26, 363, 419
 LEAD FLUORIDES: T4 602
 LEAD IODIDES: T3 602
 LEAD OXIDES 420
 LEAD OXIDES: T1 641
 LEAD OXIDES: T2 394
 LEAD: T1 25, 420
 LEAD: T15 104
 LEAD: T2 243
 LEAD: T3 1, 244
 LEAD: T30 711, 712
 LEAD: T33 713
 LEAD: T38 714
 LEAD: T4 403
 LEAST SQUARE FIT 98, 472
 LENSES 208, 331
 LEVITATION 355
 LIFETIME 233, 539
 LIGANDS 130, 194, 412, 562, 684
 LIGHT SCATTERING: Q1,Q2,Q3,Q4,Q5,Q6
 183
 LINE WIDTHS 61
 LIQUID CRYSTALS 476
 LIQUID FLOW 128, 502
 LIQUID FUELS 225
 LIQUID METAL COOLED REACTORS 25
 LIQUID METALS 1, 3, 10, 22, 24,
 25, 35, 39, 47, 63, 64, 79, 92,
 100, 105, 106, 128, 154, 156,
 158, 159, 167, 172, 177, 182,
 190, 201, 210, 224, 243, 244,
 247, 248, 249, 250, 273, 284,
 LIQUID METALS 287, 290, 291, 313,
 318, 324, 327, 328, 339, 340,
 342, 343, 355, 359, 388, 403,
 405, 420, 422, 425, 436, 446,
 460, 486, 513, 514, 522, 523,
 524, 525, 528, 529, 532, 551,
 567, 574, 580, 583, 584, 588,
 590, 606, 609, 615, 619, 620,
 622, 628, 642, 643, 653, 654,
 661, 663, 681, 686, 687, 689,
 701, 702, 705, 706
 LIQUID METALS: A1 651
 LIQUID METALS: T1 138, 152, 346,
 437, 472, 485, 502, 564, 573,
 599, 621
 LIQUID METALS: T2 198, 211, 212,
 229
 LIQUID METALS: T3 166, 575
 LIQUID METALS: T5 688
 LIQUID METALS: T9 230
 LIQUIDS 180, 225, 335, 457, 474,
 478, 579, 616
 LIQUIDS: T2 152, 653
 LITHIUM 16, 17, 23, 37, 38, 111,
 118, 281, 282, 343, 374, 380,
 381, 404, 529, 599, 727
 LITHIUM ALLOYS 269, 704
 LITHIUM ALLOYS: T1 91, 551
 LITHIUM BASE ALLOYS 274, 275
 LITHIUM BROMIDES: T2 268
 LITHIUM CHLORIDES 23, 68, 111,
 118, 234, 275, 282, 292, 293,
 314, 385, 386, 407, 430, 452,
 467, 507, 515, 675, 703, 704
 LITHIUM CHLORIDES: M10 335
 LITHIUM CHLORIDES: M2 406
 LITHIUM CHLORIDES: T1 40, 86, 142,
 721
 LITHIUM CHLORIDES: T2 91, 116,

SUBJECT DESCRIPTORS INDEX

LITHIUM CHLORIDES: T2 143, 196, 274
 LITHIUM CHLORIDES: T3 92, 341, 408, 435
 LITHIUM CHLORIDES: T6 89
 LITHIUM COMPOUNDS 111, 168, 194, 246
 LITHIUM COMPOUNDS: T4 77
 LITHIUM DEUTERIDES: T1 404
 LITHIUM DEUTERIDES: T2 529, 663
 LITHIUM DEUTERIDES: T3 405
 LITHIUM FLUORIDES 79, 176, 246, 314, 430, 452, 585
 LITHIUM FLUORIDES: T1 30, 89
 LITHIUM FLUORIDES: T2 270, 330
 LITHIUM FLUORIDES: T3 461, 462
 LITHIUM HALIDES: T2 282
 LITHIUM HYDRIDES: T1 374, 529, 663
 LITHIUM HYDRIDES: T2 404, 405
 LITHIUM IODIDES 282
 LITHIUM NITRATES: T1 195
 LITHIUM NITRATES: T2 351, 377
 LITHIUM NITRIDES 452
 LITHIUM NITRIDES: T1 467
 LITHIUM NITRIDES: T3 702
 LITHIUM NITRIDES: T5 404
 LITHIUM OXIDES 452
 LITHIUM OXIDES: M2 701
 LITHIUM TELLURIDES: A2 176
 LITHIUM TELLURIDES: T2 92
 LITHIUM TRITIDES: T3 404, 529
 LITHIUM TRITIDES: T4 405
 LITHIUM-CHLORINE BATTERIES: M1, Q2 269, 704
 LITHIUM-COPPER CHLORIDE BATTERIES: T1 720
 LITHIUM-SULFUR BATTERIES: T1 23
 LITHIUM-SULFUR BATTERIES: T2 275
 LITHIUM-SULFUR BATTERIES: T3 234
 LITHIUM: M1 106, 701
 LITHIUM: T1 161, 247, 273, 405, 430, 452, 702
 LITHIUM: T15 714
 LITHIUM: T2 10, 48, 528
 LITHIUM: T3 346, 663
 LITHIUM: T4 244, 404
 LMFBR TYPE REACTORS 57, 124, 354, 416, 471, 580, 692
 LMFBR TYPE REACTORS: M1 310, 569
 LMFBR TYPE REACTORS: T1 19, 56, 72, 94, 103, 272, 297, 309, 367, 456, 474, 527, 537, 590, 731
 LMFBR TYPE REACTORS: T3 455
 LMFBR TYPE REACTORS: T4 281
 LOSS OF FLOW: M2, Q1 19
 LOSSES 676, 691
 LOW PRESSURE 10, 22, 106, 581
 LOW TEMPERATURE 14, 31, 34, 126, 184, 261, 383, 384, 410, 414, 432, 487, 560, 594, 617, 677, 678
 LUMINESCENCE 44, 589, 664
 LUMINESCENCE: Q1, Q2 412
 LUTETIUM HYDRIDES: T4 418
 LUTETIUM OXIDES: T2 303

SUBJECT DESCRIPTORS INDEX

LUTETIUM: T6 497

LUTETIUM: T8 125

*** M ***

MACHINING 9

MAGNESIUM 48, 380, 400, 440

MAGNESIUM ALLCYS: T1 439

MAGNESIUM ALLCYS: T2 3, 39, 551

MAGNESIUM BASE ALLOYS: T3 48

MAGNESIUM CHLORIDES: T1 145

MAGNESIUM CHLORIDES: T2 139

MAGNESIUM CHLORIDES: T3 546, 604

MAGNESIUM COMPOUNDS 23

MAGNESIUM NITRATES: T1 664, 666,
668

MAGNESIUM NITRIDES: M3 400

MAGNESIUM OXIDES 23, 557

MAGNESIUM OXIDES: T1 242, 265,
352, 494

MAGNESIUM OXIDES: T2 493

MAGNESIUM OXIDES: T7 531

MAGNESIUM: T 210

MAGNESIUM: T12 104

MAGNESIUM: T24 711

MAGNESIUM: T25 712

MAGNESIUM: T27 713

MAGNESIUM: T33 714

MAGNETIC DIPOLES 123

MAGNETIC FIELDS 122, 130, 287, 559

MAGNETIC MATERIALS 95

MAGNETIC MOMENTS 538

MAGNETIC PROPERTIES 53, 137, 398,
612

MAGNETIC PROPERTIES: Q1 368

MAGNETIC PROPERTIES: Q1,Q2 189

MAGNETIC PROPERTIES: Q2,Q3 559

MAGNETIC SUSCEPTIBILITY 279, 305,
678MAGNETIC SUSCEPTIBILITY: Q1,Q2,Q3
190

MAGNETO-OPTICAL EFFECTS 122, 123

MAGNETOHYDRODYNAMICS 158

MAINTENANCE 722

MANAGEMENT 722

MANGANESE 422

MANGANESE ADDITIONS 313

MANGANESE CHLORIDES 6

MANGANESE CHLORIDES: T4 200

MANGANESE COMPLEXES: T9 314

MANGANESE: T25 711

MANGANESE: T26 712

MANGANESE: T28 713

MANGANESE: T34 714

MANGANESE: T4 687

MASS SPECTRA 295, 571

MASS SPECTRA: Q1 216

MASS SPECTROMETERS 534

MASS SPECTROSCOPY 120, 131, 214,
301, 349, 453, 470, 480, 481,
561, 597, 600

SUBJECT DESCRIPTORS INDEX

MASS SPECTROSCOPY: Q1,Q2,Q3,Q4 534
 MASS SPECTROSCOPY: Q3 716
 MASS SPECTROSCOPY: Q38 711
 MASS SPECTROSCOPY: Q4,Q5,Q6,Q7,Q8,
 Q9,Q10 715
 MASS SPECTROSCOPY: Q44 712
 MASS SPECTROSCOPY: Q5 690
 MASS SPECTROSCOPY: Q5,Q6 713
 MASS TRANSFER 25, 71, 103, 128,
 158, 193, 234, 237, 263, 277,
 356, 401, 588, 595
 MASS TRANSFER: Q1 236
 MASS TRANSFER: Q2 324
 MASS TRANSFER: Q2,Q3 79
 MATERIAL BALANCE 131, 722
 MATERIALS 38, 71
 MATERIALS HANDLING 700, 722
 MATERIALS TESTING 628, 731
 MATERIALS: Q2 23
 MATHEMATICAL MODELS 11, 167, 273,
 310, 367, 415, 425, 441, 486,
 525, 527, 567
 MATHEMATICAL MODELS: Q2 719
 MATHEMATICS 492
 MATRICES 492
 MATRIX ELEMENTS 16, 17, 683
 MATRIX MATERIALS 5
 MEAN FREE PATH 459
 MEASURING INSTRUMENTS 94, 574
 MEASURING METHODS 163, 500, 518,
 606, 681
 MEASURING METHODS: Q2 214, 573
 MEASURING METHODS: Q3 152
 MECHANICAL PROPERTIES 440, 471
 MECHANICAL PROPERTIES: Q1 209, 643
 MECHANICAL PROPERTIES: Q1,Q2,Q3 57
 MECHANICAL PROPERTIES: Q2,Q3 527
 MECHANICAL PROPERTIES: Q3,Q4 309
 MEDIUM PRESSURE 302
 MEDIUM TEMPERATURE 31, 35, 66,
 179, 184, 248, 261, 262, 284,
 306, 329, 347, 383, 384, 414,
 415, 432, 459, 466, 487, 504,
 541, 586, 591, 594, 599, 605,
 607, 617, 679, 727
 MEDIUM VACUUM 10, 106, 581
 MELTDOWN 72, 297, 475, 477, 510
 MELTDOWN: M2,Q1 474, 537
 MELTDOWN: T,Q2 454
 MELTING 19, 86, 115, 141, 224,
 300, 336, 372, 531, 540, 546,
 574, 638, 706
 MELTING POINTS 112, 116, 177, 188,
 198, 207, 213, 224, 256, 268,
 276, 282, 286, 288, 320, 338,
 399, 435, 454, 460, 478, 506,
 519, 521, 548, 553, 555, 574,
 584, 612, 629, 636, 637, 643,
 659, 668, 687, 702
 MELTING POINTS: Q1 18, 266, 267,
 321, 361
 MELTING POINTS: Q1,Q2 318
 MELTING POINTS: Q1,Q2,Q3 173
 MELTING POINTS: Q1,Q3,Q4,Q5,Q6,Q8
 191
 MELTING: Q1,Q2 421
 MELTING: Q1,Q2,Q3,Q4,Q5 213
 MENDELEVIVM 188

SUBJECT DESCRIPTORS INDEX

- MERCURY 211, 236
 MERCURY ALLOYS: T1 32, 339
 MERCURY ALLOYS: T2 138, 340
 MERCURY CHLORIDES: T2 31
 MERCURY: T40 711
 MERCURY: T41 712
 MERCURY: T42 713
 MERCURY: T47 714
 MERCURY: T5 101, 244
 METAL-METAL BATTERIES: M1 187
 METAL-METAL OXIDE BATTERIES 420
 METAL-NONMETAL BATTERIES: M1 37, 38, 111, 118, 119, 727
 METAL-NONMETAL BATTERIES: M3 263
 METAL-NONMETAL BATTERIES: T1 381
 METALLOGRAPHY 115, 129, 333, 531
 METALLOGRAPHY: Q2, Q3, Q4, Q5 332
 METALLURGICAL EFFECTS: Q1 25
 METALLURGICAL EFFECTS: Q2 211, 212
 METALLURGICAL EFFECTS: Q3 645
 METALLURGY 332
 METALS 190, 205, 723
 METALS: T1 108, 211
 METALS: T2 533
 METALS: T3 192
 METALS: T5 213
 METHANOL 347
 MICROHARDNESS 115, 209, 440, 544, 545, 610, 628
 MICROSCOPY 270
 MICROSTRUCTURE 481, 545, 604, 610, 628, 691
 MICROSTRUCTURE: Q1 440
 MICROSTRUCTURE: Q1, Q2 359, 463
 MICROWAVE RADIATION 130
 MIGRATION 401
 MISCH METAL 461
 MISCH METAL: T1 462
 MIXED SOLVENTS 347
 MIXING 550
 MIXING HEAT 205, 513, 523, 524
 MIXING HEAT: Q1, Q2, Q3 138
 MIXING RATIO 60
 MIXTURES 50, 60, 89, 125, 133, 165, 202, 292, 293, 299, 351, 474, 480, 516, 517, 518, 629, 636, 691
 MOCKUP 103
 MODULATION 411
 MOESSBAUER EFFECT 627, 657
 MOESSBAUER EFFECT: Q1 61, 559
 MOESSBAUER EFFECT: Q2 696
 MOISTURE 711, 712
 MOLECULAR IONS 2, 534
 MOLECULAR MODELS 78
 MOLECULAR ORBITAL METHOD 30, 684
 MOLECULAR STRUCTURE 13, 30, 80, 81, 102, 127, 194, 285, 498, 558, 609, 653
 MOLECULAR STRUCTURE: Q1 159, 350, 525, 567
 MOLECULAR STRUCTURE: Q1, Q2, Q3 51

SUBJECT DESCRIPTORS INDEX

MOLECULAR STRUCTURE: Q1,Q2,Q3,Q4,
Q5,Q6 195

MOLECULAR STRUCTURE: Q1,Q2,Q3,Q4
259

MOLECULAR STRUCTURE: Q1,Q2,Q3,Q4,
Q5,Q6,Q7 687

MOLECULAR STRUCTURE: Q10 730

MOLECULE-MOLECULE COLLISIONS 124,
354

MOLECULES 93, 412, 597

MOLTEN SALT FUELS: T1 237

MOLTEN SALT FUELS: T3 585

MOLTEN SALT REACTORS 236, 237, 729

MOLTEN SALT REACTORS: A1 176

MOLTEN SALT REACTORS: T1 79, 585

MOLTEN SALTS 20, 23, 27, 40, 42,
77, 79, 89, 91, 92, 93, 96, 97,
98, 102, 112, 142, 143, 145, 146,
147, 160, 195, 196, 221, 233,
244, 256, 265, 269, 275, 288,
292, 293, 299, 305, 307, 308,
314, 317, 351, 356, 360, 369,
370, 372, 375, 385, 386, 394,
402, 407, 408, 430, 433, 435,
449, 450, 451, 452, 461, 462,
488, 489, 495, 496, 506, 530,
546, 548, 556, 576, 592, 636,
659, 672, 673, 674, 675, 693,
703, 704, 721

MOLTEN SALTS: A8 176

MOLTEN SALTS: M1 163, 263

MOLTEN SALTS: T1 208, 331, 423,
550, 552, 563

MOLTEN SALTS: T11 729

MOLTEN SALTS: T2 71, 264, 276,
277, 371, 566, 585

MOLTEN SALTS: T3 204

MOLYEDATES: T1 102, 207, 520, 636

MOLYEDATES: T2 516, 519, 638

MOLYBDENUM 214, 316, 646, 649

MOLYBDENUM ALLOYS 202

MOLYBDENUM ALLOYS: M1 646

MOLYBDENUM ALLOYS: T1 100

MOLYBDENUM ALLOYS: T6 115

MOLYBDENUM CARBIDES 650

MOLYBDENUM CARBIDES: T1 647

MOLYBDENUM CHLORIDES: T2 554, 555

MOLYBDENUM CHLORIDES: T3 499, 547

MOLYBDENUM COMPLEXES 305

MOLYBDENUM COMPOUNDS 531

MOLYBDENUM COMPOUNDS: T1 515

MOLYBDENUM FLUORIDES: T1 51, 52,
397

MOLYBDENUM OXIDES 202, 214, 281

MOLYBDENUM OXIDES: T1 305, 316

MOLYBDENUM OXIDES: T7 191

MOLYBDENUM: M1 47

MOLYBDENUM: M2 574

MOLYBDENUM: T1 588

MOLYBDENUM: T10 115

MOLYBDENUM: T13 104

MOLYBDENUM: T2 25, 156, 202

MOLYBDENUM: T26 711

MOLYBDENUM: T27 712

MOLYBDENUM: T29 713

MOLYBDENUM: T3 644

MOLYBDENUM: T35 714

SUBJECT DESCRIPTORS INDEX

MONITORING: Q1,Q2,Q3 64
 MONOCRYSTALS 29, 30, 165, 363,
 419, 466, 484, 656, 677
 MONTE CARLO METHOD 229, 441, 442
 MOTION 19
 MUFFIN-TIN POTENTIAL 161
 MULTI-ELEMENT SEPARATION 534
 MULTICHARGED IONS 130, 261

*** N ***

N-TYPE CONDUCTORS 321
 NATURAL URANIUM 226
 NEODYMIUM 321
 NEODYMIUM ALICYS 321, 462
 NEODYMIUM BORIDES: T3 215
 NEODYMIUM CHLORIDES: T1 457
 NEODYMIUM CHLORIDES: T2 40, 434,
 494
 NEODYMIUM CHLORIDES: T4 112
 NEODYMIUM COMPOUNDS 237, 326
 NEODYMIUM COMPOUNDS: T1 357
 NEODYMIUM COMPOUNDS: T4 82
 NEODYMIUM FLUORIDES: T1 443
 NEODYMIUM FLUORIDES: T4 382, 461
 NEODYMIUM NITRATES: M1 233
 NEODYMIUM NITRATES: T1 667
 NEODYMIUM NITRATES: T2 664, 668
 NEODYMIUM NITRATES: T4 164
 NEODYMIUM OXIDES: T2 427

NEODYMIUM 148: T3 716
 NEODYMIUM: T2 233
 NEODYMIUM: T5 230
 NEODYMIUM: T9 125
 NEPTUNIUM 188, 428
 NEPTUNIUM CHLORIDES 406
 NEPTUNIUM CHLORIDES: T4 562
 NEPTUNIUM CHLORIDES: T5 360
 NEPTUNIUM COMPLEXES 660
 NEPTUNIUM COMPLEXES: T1 562
 NEPTUNIUM COMPOUNDS: T1 82
 NEPTUNIUM HYDRIDES: T1 428
 NEPTUNIUM OXIDES 385
 NEPTUNIUM OXIDES: M5 662
 NEPTUNIUM OXIDES: T9 303
 NEPTUNIUM 237: T 718
 NEPTUNIUM: T1 213, 360, 385, 386,
 660
 NEPTUNIUM: T4 683
 NEUTRON DIFFRACTION 195, 273, 359,
 379, 564, 611, 653
 NEUTRON DIFFRACTION: Q1,Q2,Q3 609
 NEUTRON FLUENCE 232, 416
 NEUTRON FLUX 169
 NEUTRON FLUX: Q1 416
 NEUTRON MONITORS 725
 NEUTRON REACTIONS 619
 NEUTRON SPECTRA 416
 NICKEL 275, 422
 NICKEL ADDITIONS 313

SUBJECT DESCRIPTORS INDEX

NICKEL ALLCYS: M1 5
NICKEL ALLCYS: T2 661, 689
NICKEL ALLCYS: T3 24
NICKEL ALLCYS: T4 172
NICKEL BASE ALLOYS: T1 532
NICKEL BASE ALLCYS: T2 100
NICKEL CHLORIDES: T3 200
NICKEL COMPLEXES: T1 673
NICKEL COMPLEXES: T6 314
NICKEL COMPOUNDS 68
NICKEL FLUORIDES 381
NICKEL OXIDES: T1 149
NICKEL 65 68
NICKEL: M4 311
NICKEL: T1 533
NICKEL: T14 104
NICKEL: T2 128
NICKEL: T28 711, 712
NICKEL: T31 713
NICKEL: T36 714
NICKEL: T7 687
NIOBATES: M3 495
NIOBATES: T 164
NIOBATES: T1 173
NIOBATES: T2 490
NIOBATES: T3 206
NIOBIUM 582, 658
NIOBIUM ADDITIONS 324
NIOBIUM ALLOYS: T1 587
NIOBIUM ALLOYS: T2 532
NIOBIUM ALLOYS: T3 100
NIOBIUM BROMIDES: T1 561
NIOBIUM CARBIDES 582
NIOBIUM CHLORIDES: T2 561
NIOBIUM CHLORIDES: T3 554
NIOBIUM CHLORIDES: T4 499
NIOBIUM COMPLEXES: T3 561
NIOBIUM FLUORIDES: T1 170, 216, 300
NIOBIUM IODIDES: T4 561
NIOBIUM IONS 261
NIOBIUM NITRIDES 582
NIOBIUM OXIDES 342
NIOBIUM OXIDES: T1 658
NIOBIUM OXIDES: T2 206
NIOBIUM OXIDES: T3 542
NIOBIUM OXIDES: T4 183, 191
NIOBIUM: M1 574
NIOBIUM: M2 47
NIOBIUM: T1 342, 490
NIOBIUM: T13 714
NIOBIUM: T8 115
NITRATES 426, 592
NITRATES: T2 113
NITRATES: T6 195
NITRIC ACID 113
NITRIC OXIDE: M4 218
NITRIDATION: Q1 400

SUBJECT DESCRIPTORS INDEX

NITRIDES 429
 NITRIDES: T3 582
 NITROGEN 373, 452, 649
 NITROGEN DIOXIDE: M3 218
 NITROGEN OXIDES 722
 NITROGEN OXIDES: T6 94
 NITROGEN 14 219
 NITROGEN 15: T3 219
 NITROGEN: T11 713
 NITROGEN: T2 702, 712
 NITROGEN: T3 532
 NITROGEN: T5 100
 NITROGEN: T7 711
 NITROGEN: T8 714
 NITROSO COMPOUNDS: T7 66
 NOBELIUM 188
 NON-PROLIFERATION TREATY: T 722
 NONDESTRUCTIVE ANALYSIS 453
 NONMETALS 190, 528
 NONMETALS: T2 486
 NUCLEAR FUELS 179, 214, 226, 272,
 308, 310, 349, 401, 431, 474,
 477, 480, 579, 613, 618, 649, 703
 NUCLEAR FUELS: Q1 345
 NUCLEAR FUELS: T1 73, 202
 NUCLEAR FUELS: T3 448
 NUCLEAR FUELS: T4 716
 NUCLEAR FUELS: T5, Q4 281
 NUCLEAR MAGNETIC RESONANCE 21, 66,
 463, 564, 571
 NUCLEAR MAGNETIC RESONANCE: Q2 363
 NUCLEAR MATERIALS MANAGEMENT: T2, Q1
 349
 NUCLEAR MATERIALS MANAGEMENT: Q1
 600
 NUCLEAR QUADRUPOLE RESONANCE: Q1
 31
 NUCLEAR QUADRUPOLE RESONANCE: Q1, T2
 564
 NUSSELT NUMBER 502
 *** O ***
 OCCUPATION NUMBER 683
 OFF-GAS SYSTEMS 94, 699, 700
 OFF-GAS SYSTEMS: T4 722
 OFF-PEAK ENERGY STORAGE: T2 269
 OILS: T5 301
 ON-LINE MEASUREMENT SYSTEMS 534
 ONE-DIMENSIONAL CALCULATIONS 229
 OPERATION 699, 722
 OPTICAL PROPERTIES 398, 656
 OPTICAL PROPERTIES: Q1, Q2, Q3, Q4
 521
 OPTICAL PYROMETERS 474, 475
 OPTIMIZATION 343, 676
 ORBIT STABILITY 320
 ORDER PARAMETERS 229, 414
 ORDER-DISORDER TRANSFORMATIONS
 224, 260, 463
 ORE PROCESSING 200
 ORGANIC BORON COMPOUNDS: T2 194

SUBJECT DESCRIPTORS INDEX

ORGANIC ION EXCHANGERS: T7	301	OXYGEN: T2	563
ORGANIC NITROGEN COMPOUNDS	727	OXYGEN: T3	711
ORGANIC SOLVENTS	111, 371	OXYGEN: T4	182
ORGANIC SOLVENTS: T9	301	OXYIODIDES	82
ORGANIC SULFUR COMPOUNDS	246		
ORGANOMETALLIC COMPOUNDS	111	*** P ***	
ORIENTATION	559		
ORTHORHOMBIC LATTICES	419, 432, 647	PAIRING ENERGY	159
OSCILLATOR STRENGTHS	511	PALLADIUM	422
OXIDATION	202, 342, 360, 700	PALLADIUM ALLOYS	202, 653
OXIDATION: Q1	424, 465, 542	PALLADIUM ALLOYS: T5	417
OXIDES	25, 64, 183, 206, 307, 399, 427, 429	PALLADIUM IODIDES: T5	94
OXIDES: T2	463	PALLADIUM SILICIDES: T2	609
OXIDES: T3	2, 634	PALLADIUM: T31	712
OXONIUM IONS: T	409	PARTIAL MOLAL VOLUME	302
OXYBROMIDES	82	PARTICLE INTERACTIONS	661
OXYCHLORIDES	82, 196, 221, 356	PARTITION FUNCTIONS	78
OXYCHLORIDES: T2	515	PARTITION FUNCTIONS: Q1	285
OXYFLUORIDES	558	PATTERSON METHOD	695
OXYGEN	35, 54, 74, 107, 110, 115, 175, 389, 452, 509, 535, 565, 588, 626	PERCHLORATES	246, 380
OXYGEN ADDITIONS	193	PERCUS-YEVICK EQUATION	210, 615, 661
OXYGEN COMPOUNDS	176, 185, 495	PERFORMANCE	38, 275, 315, 381, 699, 724
OXYGEN IONS	595	PERFORMANCE TESTING	416, 725
OXYGEN 16	220	PERFORMANCE TESTING: Q9	722
OXYGEN 18	220	PERFORMANCE: Q1	118
OXYGEN: T1	63, 510	PERMEABILITY: Q1	693
OXYGEN: T2	64, 235, 342, 401, 441,	PERMEABILITY: Q2	10
		PERMEABILITY: Q2, Q3, Q4	408

SUBJECT DESCRIPTORS INDEX

PERTURBATION THEORY 167, 279

PETROLEUM PRODUCTS 426

PH VALUE 329

PHASE DIAGRAMS 1, 4, 24, 26, 39,
54, 83, 108, 149, 164, 175, 209,
226, 269, 274, 290, 296, 320,
332, 333, 404, 421, 438, 440,
446, 461, 480, 494, 496, 509,
510, 518, 555, 593, 611, 612,
636, 637, 644, 645, 646, 647,
649, 654, 659, 691, 703

PHASE DIAGRAMS: Q1 62, 107, 321,
373, 376, 679, 680

PHASE DIAGRAMS: Q1,Q2 162, 174,
266, 267, 270, 288, 443, 448,
493, 503, 506, 540, 545, 572,
631, 667, 668, 706

PHASE DIAGRAMS: Q1,Q2,Q3 21, 50,
143, 151, 222, 254, 268, 317,
351, 377, 394, 434, 495, 520,
529, 547, 548, 553, 648, 666

PHASE DIAGRAMS: Q1,Q2,Q3,Q4 86,
90, 117, 228, 257, 336, 435, 544,
554, 602

PHASE DIAGRAMS: Q1,Q2,Q3,Q4,Q5 45

PHASE DIAGRAMS: Q1,Q2,Q3,Q4,Q5,Q6
89

PHASE DIAGRAMS: Q1,Q2,Q3,Q4,Q5
139, 147

PHASE DIAGRAMS: Q1,Q2,Q3,Q4,Q5,Q6,
Q7 160

PHASE DIAGRAMS: Q1,Q2,Q3,Q4,Q5,Q6,
Q7,Q8 196

PHASE DIAGRAMS: Q1,Q2,Q3,Q4,Q5 245

PHASE DIAGRAMS: Q1,Q2,Q3,Q4,Q5,Q6
256, 258

PHASE DIAGRAMS: Q1,Q2,Q3,Q4,Q5
299, 300

PHASE DIAGRAMS: Q1,Q2,Q3,Q4,Q5,Q6,
Q7,Q8,Q9,Q10 516

PHASE DIAGRAMS: Q1,Q2,Q3,Q4,Q5
519, 528, 530

PHASE DIAGRAMS: Q1,Q2,Q3,Q4,Q5,Q6
546

PHASE DIAGRAMS: Q1,Q2,Q3,Q4,Q5 556

PHASE DIAGRAMS: Q1,Q2,Q3,Q4,Q5,Q6
629, 630

PHASE DIAGRAMS: Q1,Q2,Q3,Q4,Q5
638, 639

PHASE DIAGRAMS: Q1,Q2,Q3,Q4,Q5,Q6
640

PHASE DIAGRAMS: Q1,Q3,Q4 168

PHASE DIAGRAMS: Q1,T2 552

PHASE DIAGRAMS: Q2,Q3 55

PHASE DIAGRAMS: Q2,Q3,Q4,Q5 116

PHASE DIAGRAMS: Q2,Q3,Q4,Q5,Q6 77

PHASE DIAGRAMS: Q2,Q4 177

PHASE DIAGRAMS: Q6,Q7 729

PHASE STABILITY 209

PHASE STUDIES 110, 120, 134, 182,
247, 335, 340, 359, 362, 393,
394, 400, 481, 509, 551, 555,
587, 626, 628, 642, 644, 645,
648, 676, 697

PHASE STUDIES: Q1 325, 428

PHASE STUDIES: Q1,Q2 136, 199,
468, 469, 647

PHASE STUDIES: Q1,Q2,Q3 49, 87,
88, 235, 510, 593, 641, 646, 659

PHASE STUDIES: Q1,Q2,Q3,Q4 518,
541

PHASE STUDIES: Q1,Q2,Q3,Q4,Q5,Q6,
Q7,Q8,Q9,Q10 303

PHASE STUDIES: Q2 282

PHASE TRANSFORMATIONS 24, 199,
261, 262, 361, 363, 438, 448,

SUBJECT DESCRIPTORS INDEX

- PHASE TRANSFORMATIONS 457, 459, 476, 509, 593, 610, 616, 641, 646
 PHASE TRANSFORMATIONS: Q1 18, 313
 PHASE TRANSFORMATIONS: Q1,Q2 11, 581, 645
 PHASE TRANSFORMATIONS: Q1,Q2,Q3 517
 PHONONS 459
 PHOSPHORS 491
 PHOSPHORUS 211
 PHOSPHORUS FLUORIDES 381
 PHOSPHORUS: T29 711, 712
 PHOSPHORUS: T32 713
 PHOSPHORUS: T37 714
 PHOTOELECTRON SPECTROSCOPY 28, 223, 280, 598
 PHOTOELECTRON SPECTROSCOPY: Q1,Q2,Q3,Q4,Q5,Q6 417
 PHOTOEMISSION 28, 662
 PHOTOLYSIS 16, 17
 PHYSICAL PROPERTIES 23, 215, 282, 501, 651
 PHYSICAL PROPERTIES: Q1 209, 226, 454, 643, 656
 PHYSICAL PROPERTIES: Q1,Q2 141, 409
 PHYSICAL PROPERTIES: Q1,Q2,Q3 114
 PHYSICAL PROPERTIES: Q1,Q2,Q3,Q4 574
 PHYSICAL PROPERTIES: Q1,Q2,Q3,Q4,Q5,Q6,Q7,Q8,Q9,Q10,Q11,Q12,Q13,Q14,Q15 612
 PHYSICAL PROPERTIES: Q3,Q4,Q5,Q6,Q7,Q8 367
 PILOT PLANTS 94, 699, 700
 PILOT PLANTS: Q5 722
 PLASTICITY 115, 531, 643
 PLATINUM 422, 721
 PLATINUM ALLOYS: T1 399
 PLEXIGLAS: T3 301
 PLUTONIUM 54, 75, 107, 131, 188, 349, 474, 691, 715, 716
 PLUTONIUM BROMIDES: T9 82
 PLUTONIUM CARBIDES 456
 PLUTONIUM CARBIDES: M1 612
 PLUTONIUM CARBIDES: T1 84, 414, 415, 479, 480, 481, 648, 691
 PLUTONIUM CARBIDES: T2 54, 55, 83, 134, 179, 577, 644, 645, 676
 PLUTONIUM CARBIDES: T3 579
 PLUTONIUM CARBIDES: T4 367
 PLUTONIUM CARBIDES: T7 281
 PLUTONIUM CHLORIDES: M3 406
 PLUTONIUM CHLORIDES: T1 147
 PLUTONIUM CHLORIDES: T4 360
 PLUTONIUM COMPLEXES 660
 PLUTONIUM COMPOUNDS 222
 PLUTONIUM COMPOUNDS: T2 82
 PLUTONIUM DIOXIDE 220, 416, 431
 PLUTONIUM DIOXIDE: T1 15, 54, 55, 74, 109, 110, 184, 281, 304, 455, 475, 477, 482, 712
 PLUTONIUM DIOXIDE: T10 303
 PLUTONIUM DIOXIDE: T15 125
 PLUTONIUM DIOXIDE: T2 60, 73, 618, 713, 725
 PLUTONIUM DIOXIDE: T3 175

SUBJECT DESCRIPTORS INDEX

PLUTONIUM DIOXIDE: T4 94, 474, 676	PLUTONIUM 239: T1 690
PLUTONIUM DIOXIDE: T6 367	PLUTONIUM 239: T2 353
PLUTONIUM FLUORIDES 308	PLUTONIUM 239: T6 715
PLUTONIUM FLUORIDES: T1 222, 353	PLUTONIUM 240: T7 715
PLUTONIUM FLUORIDES: T2 724	PLUTONIUM 241: T8 715
PLUTONIUM FLUORIDES: T3 487	PLUTONIUM 242 184
PLUTONIUM HYDRIDES: T2 222	PLUTONIUM 242: T1 487
PLUTONIUM IODIDES: T10 82	PLUTONIUM 242: T3 353
PLUTONIUM ISOTOPE: T44 712	PLUTONIUM 242: T9 715
PLUTONIUM ISOTOPE: T6 713	PLUTONIUM 243 61
PLUTONIUM NITRATES: T1 85	PLUTONIUM 244 184
PLUTONIUM NITRIDES: M2 612	PLUTONIUM-DELTA: T1 620
PLUTONIUM NITRIDES: T1 9	PLUTONIUM-EPSILON 318
PLUTONIUM NITRIDES: T2 578	PLUTONIUM: T1 619, 714, 725
PLUTONIUM NITRIDES: T4 579	PLUTONIUM: T2 213, 318, 360, 660
PLUTONIUM OXIDES 301, 316, 690	PLUTONIUM: T3 83, 690
PLUTONIUM OXIDES: M1 220	PLUTONIUM: T4 713
PLUTONIUM OXIDES: M10 662	PLUTONIUM: T45 712
PLUTONIUM OXIDES: M3 612	PLUTONIUM: T5 683
PLUTONIUM OXIDES: T1 62, 107, 133, 169, 398, 478, 565, 613	POISSON EQUATION 161
PLUTONIUM OXIDES: T2 476, 487, 626	POISSON RATIO 209
PLUTONIUM OXIDES: T3 55, 72, 222	POLARIZATION 38, 92, 193, 410, 675, 727
PLUTONIUM OXIDES: T7 202	POLAROGRAPHY 236
PLUTONIUM PHOSPHIDES: M4 612	POLYCRYSTALS 560
PLUTONIUM SULFIDES: M5 612	POLYETHYLENES 118
PLUTONIUM 238 301	POLYETHYLENES: T8 301
PLUTONIUM 238: T2 15	POLYMERIZATION 196
PLUTONIUM 239 301	POROSITY 115, 459, 531, 697

SUBJECT DESCRIPTORS INDEX

POTASSIUM 16, 17, 92, 108, 161,
343, 599

POTASSIUM ALICYS: T2 590

POTASSIUM BROMIDES 129, 483, 631

POTASSIUM BROMIDES: T1 90

POTASSIUM BROMIDES: T5 630

POTASSIUM CARBONATES 396

POTASSIUM CHLORIDES 6, 23, 27, 68,
234, 275, 293, 314, 385, 396,
406, 407, 449, 450, 451, 452,
507, 515, 631, 675, 703, 704

POTASSIUM CHLORIDES: M11 335

POTASSIUM CHLORIDES: T1 292, 299,
488, 517, 576

POTASSIUM CHLORIDES: T2 90, 117,
174, 231, 300, 432, 457, 520,
659, 721

POTASSIUM CHLORIDES: T3 40, 91,
116, 274

POTASSIUM CHLORIDES: T4 92, 196,
546, 630

POTASSIUM COMPOUNDS: T1 530

POTASSIUM COMPOUNDS: T2 66

POTASSIUM COMPOUNDS: T3 556

POTASSIUM COMPOUNDS: T6 77

POTASSIUM COMPOUNDS: T8 516

POTASSIUM FLUORIDES 314, 381, 452

POTASSIUM FLUORIDES: T1 295, 541

POTASSIUM FLUORIDES: T2 296, 299,
305, 517, 576

POTASSIUM FLUORIDES: T3 21, 90,
127, 300, 516, 520

POTASSIUM FLUORIDES: T6 41

POTASSIUM IODIDES 282

POTASSIUM IODIDES: T3 432, 517

POTASSIUM IODIDES: T4 90

POTASSIUM IODIDES: T6 630

POTASSIUM NITRATES 233

POTASSIUM NITRATES: T1 234

POTASSIUM NITRATES: T2 195

POTASSIUM NITRATES: T3 254, 351

POTASSIUM OXIDES: T3 590

POTASSIUM TUNGSTATES: M3 338

POTASSIUM TUNGSTATES: T1 639

POTASSIUM TUNGSTATES: T9 516

POTASSIUM: T2 167, 250

POTASSIUM: T24 712

POTASSIUM: T3 528, 584

POTASSIUM: T32 714

POTASSIUM: T40 713

POTASSIUM: T42 711

POTASSIUM: T5 249

POTASSIUM: T6 244

POTENTIAL ENERGY 273

POTENTIOMETRY 291, 293, 329, 515

POTENTIOMETRY: Q1 292

POWDER METALLURGY 226

POWDER METALLURGY: Q1, Q2, Q3 557

POWDERS 9, 431, 531, 711, 712,
713, 725

POWER REACTORS 699, 700

PRANDTL NUMBER 502

PRASEODYMIUM 677

SUBJECT DESCRIPTORS INDEX

- PRASEODYMIUM ALLCYS 462
 PRASEODYMIUM CHLORIDES: T5 112
 PRASEODYMIUM COMPOUNDS 168, 326, 492
 PRASEODYMIUM COMPOUNDS: T2 677
 PRASEODYMIUM FLUORIDES: T2 312
 PRASEODYMIUM FLUORIDES: T3 382, 484
 PRASEODYMIUM FLUORIDES: T5 461
 PRASEODYMIUM NITRATES: T2 165
 PRASEODYMIUM NITRATES: T3 164
 PRASEODYMIUM OXIDES: M2 395
 PRASEODYMIUM OXIDES: T1 427
 PRASEODYMIUM: T6 230, 688
 PRECIPITATION HARDENING 604
 PREDICTION EQUATIONS 231
 PRESSURE DEPENDENCE 9, 31, 213, 389, 405, 478, 529, 532, 535, 557, 574, 575, 584, 599, 611, 618, 677, 691
 PRESSURE DROP 103
 PRESSURE MEASUREMENT 3
 PRESSURE VESSELS 569
 PRESTRESSED CONCRETE 569
 PRIMARY COOLANT CIRCUITS 324
 PROBABILITY 539
 PRODUCTION 691
 PRODUCTION: Q1 232
 PRODUCTION: Q1,Q2 649
 PRODUCTION: Q1,Q2,Q3,Q4,Q5 334
 PRODUCTIVITY 129
 PROMETHIUM COMPOUNDS: T3 639
 PROMETHIUM OXIDES 18
 PROMETHIUM 147 18
 PROMETHIUM: T1 18
 PROTACTINIUM 188, 237
 PROTACTINIUM BROMIDES: T1 80
 PROTACTINIUM CARBIDES: T4 80
 PROTACTINIUM CHLORIDES: T2 80
 PROTACTINIUM CHLORIDES: T6 562
 PROTACTINIUM COMPLEXES: T2 562
 PROTACTINIUM FLUORIDES: T1 465
 PROTACTINIUM IODIDES: T3 80
 PROTACTINIUM: T2 683
 PROTECTIVE CLOTHING 710
 PROTECTIVE COATINGS 400
 PULSE TECHNIQUES 363, 478
 PULSED IRRADIATION 474
 PUREX PROCESS 69
 PURIFICATION 135, 724
 PURIFICATION: Q1 232
 PVC: T4 301
 PWR TYPE REACTORS 455, 510
 PWR TYPE REACTORS: T1 509, 719
 PYROLYSIS 66, 85, 168, 399
 PYROLYSIS: Q1,Q2 395
 PYROLYSIS: Q1,Q2,Q3 312
 PYROLYSIS: Q1,Q2,Q3,Q4,Q5,Q6 393
 PYROLYSIS: Q1,Q2,Q3,Q4,Q5,Q6,Q7,Q8 561

SUBJECT DESCRIPTORS INDEX

PYROLYSIS: Q3 702

*** R ***

PYROLYTIC CARBON 315

PYROMETALLURGY 69

PYROMETERS 179, 476

*** Q ***

QUADRUPOLE MOMENTS 492

QUADRUPOLES 130, 538, 559, 627

QUALITATIVE CHEMICAL ANALYSIS 721

QUANTITATIVE CHEMICAL ANALYSIS
104, 125, 292, 332, 333, 349,
431, 490, 566, 623QUANTITATIVE CHEMICAL ANALYSIS: Q2
632QUANTITATIVE CHEMICAL ANALYSIS: Q4
711QUANTITATIVE CHEMICAL ANALYSIS: Q3
712QUANTITATIVE CHEMICAL ANALYSIS: Q8
713QUANTITY RATIO 6, 22, 24, 27, 91,
96, 97, 98, 108, 129, 131, 138,
141, 142, 145, 146, 156, 166,
177, 190, 193, 201, 268, 299,
316, 321, 329, 336, 339, 340,
341, 349, 359, 369, 371, 389,
434, 450, 457, 461, 462, 463,
493, 496, 504, 515, 523, 532,
565, 568, 592, 653, 667, 722

QUANTUM MECHANICS 78

QUARTZ 18

QUATERNARY ALLOY SYSTEMS 332, 333,
645, 648

RADIATION HEATING 475, 619

RADIATION PROTECTION 135

RADICALS 195, 385

RADIOACTIVE WASTE DISPOSAL: M, Q1
311RADIOACTIVE WASTE PROCESSING: T
358

RADIOACTIVE WASTES: M5 311

RADIOACTIVE WASTES: T1 301

RADIOACTIVITY 135

RADIOCHEMISTRY: T, Q1 723

RADIOISOTOPE HEAT SOURCES 304

RADIOISOTOPE HEAT SOURCES: T, Q2 15

RADIOISOTOPES 76

RADIOLYSIS 85, 724

RADIOLYSIS: Q3, Q4, Q5, Q6, Q7, Q8, Q9,
Q10 301

RADIOMETRIC ANALYSIS 75, 619

RADIOMETRIC ANALYSIS: Q10 714

RADIUM CHLORIDES: T3 196

RAMAN EFFECT 14, 41

RAMAN EFFECT: Q1, Q2 488

RAMAN EFFECT: Q1, Q2, Q3 489

RAMAN SPECTRA 51, 66, 80, 127,
409, 498, 678, 695

RAMAN SPECTRA: Q1, Q2 13, 102

RAMAN SPECTRA: Q1, Q2, Q3 204

RAMAN SPECTRA: Q2 203

RARE EARTH ADDITIONS 531

SUBJECT DESCRIPTORS INDEX

RARE EARTH ALLOYS 686
 RARE EARTH ALLOYS: M2 559
 RARE EARTH ALLOYS: M3 671
 RARE EARTH ALLOYS: T 320
 RARE EARTH ALLOYS: T1 11, 65, 137, 205, 217, 376, 392, 549
 RARE EARTH COMPLEXES 13, 14
 RARE EARTH COMPLEXES: T1 329
 RARE EARTH COMPOUNDS 29, 123, 280, 308, 427, 490, 492, 495, 538, 639
 RARE EARTH COMPOUNDS: A4 338
 RARE EARTH COMPOUNDS: M1 168, 197, 279
 RARE EARTH COMPOUNDS: M2 122
 RARE EARTH COMPOUNDS: T1 2, 99, 114, 185, 206, 215, 391, 511
 RARE EARTH COMPOUNDS: T2 173, 207, 326, 357, 438, 447, 634, 666
 RARE EARTH COMPOUNDS: T3 113, 593
 RARE EARTH OXIDES: M2 49
 RARE EARTHS 122, 183, 237, 686, 723
 RARE EARTHS: M1 426
 RARE EARTHS: T1 115, 157, 280, 319, 412, 429, 492, 538
 RARE EARTHS: T11 714
 RARE EARTHS: T14 713
 RARE EARTHS: T2 610, 623
 RARE EARTHS: T3 213
 RARE EARTHS: T7 712
 RARE EARTHS: T9 711
 RARE GASES 497
 RARE GASES: T1 225
 REACTION HEAT 38, 118, 395
 REACTION KINETICS 3, 6, 32, 96, 221, 284, 339, 388, 406, 532, 586
 REACTION KINETICS: Q1, Q2, Q3, Q4, Q5, Q6 372
 REACTOR ACCIDENTS 19
 REACTOR ACCIDENTS: T, Q1 297
 REACTOR COMPONENTS 25
 REACTOR COMPONENTS: T2, Q1 731
 REACTOR COOLING SYSTEMS 585
 REACTOR COOLING SYSTEMS: M2, Q1 103
 REACTOR COOLING SYSTEMS: Q1 56
 REACTOR CORE DISRUPTION 124, 354
 REACTOR CORE DISRUPTION: M2, Q1 310, 367
 REACTOR CORE DISRUPTION: T2, Q1 601
 REACTOR CORES 310
 REACTOR CORES: T2 454
 REACTOR MATERIALS 57, 104, 454, 471, 692
 REACTOR MATERIALS: M10, Q1 367
 REACTOR MATERIALS: M2, Q1 527
 REACTOR MATERIALS: T2, Q1 309
 REACTOR MATERIALS: T3, Q1 731
 REACTOR OPERATION 453
 REACTOR SAFETY 60, 367, 477
 RECOMMENDATIONS 717
 RECOVERY 69
 RECOVERY: Q1 725
 RECOVERY: Q1, Q2 70

SUBJECT DESCRIPTORS INDEX

- RECRYSTALLIZATION 541
 REDOX POTENTIAL 293
 REDOX POTENTIAL: Q1 386, 449, 451
 REDOX POTENTIAL: Q1,Q2 565
 REDOX REACTIONS: Q4,Q5,Q6 360
 REDUCING AGENTS 304
 REDUCTION 18, 45, 332, 333, 360, 503, 691
 REDUCTION: Q1 54, 55, 304, 465
 REDUCTION: Q1,Q2 515, 654
 REDUCTION: Q1,Q2,Q3,Q4,Q5,Q6 399
 REDUCTION: Q3 176
 REFINING 588
 REFLECTION: Q1,Q2,Q3,Q4,Q5,Q6,Q7 562
 REFLECTIVITY: Q1,Q2,Q3,Q4,Q5 484
 REFRACTION 270
 REFRACTIVITY: Q1,Q2,Q3 173
 REFRACTORIES: T1 531
 RELAXATION 346, 410, 538
 RELAXATION: Q2 564
 RELIABILITY 94
 REMOVAL 404
 REMOVAL: Q3 94
 REPROCESSING 70
 REPROCESSING: M1 311
 REPROCESSING: Q1 69, 237, 358
 REPROCESSING: Q2 236
 REPROCESSING: Q4 135
 REPROCESSING: Q7 94
 REPROCESSING: T3,Q1 79
 RESEARCH PROGRAMS 58, 237, 533, 721
 RESEARCH PROGRAMS: Q1 71
 RESEARCH PROGRAMS: Q10 367
 RESEARCH PROGRAMS: Q11 729
 RESEARCH PROGRAMS: Q18 730
 RESEARCH PROGRAMS: Q2 94
 RESEARCH PROGRAMS: Q2,Q3 731
 RESEARCH PROGRAMS: Q3 722
 RESEARCH PROGRAMS: Q7 501
 RESOLUTION 152
 REVIEWS 65, 189, 263, 266, 280, 395, 412, 492, 500, 525, 528, 538, 557, 573, 612, 662, 679, 680, 686, 731
 REVIEWS: Q1 651
 REYNOLDS NUMBER 588
 RHENIUM ALLOYS: T3 549
 RHENIUM FLUORIDES: T1 53
 RHODIUM ALLOYS 202
 RHODIUM COMPLEXES: T2 672
 ROTATIONAL STATES 220
 ROUGHNESS 502
 RUBIDIUM 343, 599
 RUBIDIUM BROMIDES: T3 268
 RUBIDIUM BROMIDES: T6 640
 RUBIDIUM CHLORIDES 498
 RUBIDIUM CHLORIDES: T1 286
 RUBIDIUM CHLORIDES: T2 506
 RUBIDIUM CHLORIDES: T3 178, 231,

SUBJECT DESCRIPTORS INDEX

RUBIDIUM CHLORIDES: T3 257
 RUBIDIUM CHLORIDES: T4 408, 435, 640
 RUBIDIUM CHLORIDES: T5 196
 RUBIDIUM COMPOUNDS: T3 66
 RUBIDIUM COMPOUNDS: T4 127
 RUBIDIUM FLUORIDES: T2 443, 540
 RUBIDIUM IODIDES: T4 178, 257
 RUBIDIUM IODIDES: T5 640
 RUBIDIUM ISOTOPES: T3 534
 RUBIDIUM NITRATES: T3 195, 377
 RUBIDIUM TUNGSTATES: T3 151
 RUBIDIUM: T2 460, 584
 RUBIDIUM: T4 528
 RUDERMAN-KITTEL COUPLING 368
 RUTHENIUM ALLIGYS 202
 RUTHENIUM FLUORIDES 722
 RUTHENIUM OXIDES: T1 558
 RUTHENIUM: M5 47
 RUTILE: T2 542

*** S ***

S CODES 19, 54, 601
 S STATES 161
 SAFEGUARDS 349
 SAFEGUARDS: Q1 131
 SAFETY 276, 651, 700
 SALTS 102
 SAMARIUM 306, 399

SAMARIUM ALLOYS: T5 399
 SAMARIUM CHLORIDES: T3 594
 SAMARIUM COMPOUNDS 326
 SAMARIUM FLUORIDES: T1 306
 SAMARIUM FLUORIDES: T5 382
 SAMARIUM OXIDES: T1 303
 SAMARIUM OXIDES: T3 427
 SAMARIUM OXIDES: T5 241
 SAMARIUM: T10 125
 SAMPLE PREPARATION 104, 417
 SAMPLE PREPARATION: Q1 690
 SAMPLERS 725
 SAMPLING 600, 725
 SAPPHIRE 16, 17
 SATURATION 22, 97, 495
 SBR-5 REACTOR: T2 135
 SCALING LAWS 249, 376
 SCANDIUM BROMIDES: T1 497
 SCANDIUM FLUORIDES: T5 41
 SCANDIUM HYDROXIDES: M1 393
 SCANDIUM OXIDES: T6 303
 SCANDIUM SULFATES: M2 393
 SCANDIUM SULFIDES: T1 597
 SCANDIUM: T7 688
 SCATTERING 159, 567, 674
 SCHOTTKY EFFECT 414
 SCRUBBING: Q6 94
 SEALS 111, 722
 SECONDARY COOLANT CIRCUITS 324

SUBJECT DESCRIPTORS INDEX

SEGREGATION 332, 333, 481, 645

SEGREGATION: Q1,Q2 84

SEGREGATION: Q1,Q2,Q3,Q4 83

SELECTION RULES 186

SELENIUM 653

SELENIUM: T1 609

SELF-DIFFUSION 318, 564, 612

SELF-DIFFUSION: Q1,Q2,Q3 198

SELF-DIFFUSION: Q1,Q2,Q3,Q4 328

SELNI REACTOR: T1 75

SENSITIVITY 104, 319

SEPARATION PROCESSES 237, 404,
690, 722

SEPARATION PROCESSES: Q1 308

SEPARATION PROCESSES: Q1,Q2 307

SHEAR PROPERTIES 11

SHIELDING MATERIALS 710

SILICIDES 531

SILICON ADDITIONS 313

SILICON ALLOYS 653

SILICON ALLOYS: T3 661

SILICON CARBIDES 315, 722

SILICON FLUORIDES 307, 308

SILICON OXIDES 463

SILICON OXIDES: M3 372

SILICON OXIDES: T2 710

SILICON: T17 104

SILICON: T32 711

SILICON: T33 712

SILICON: T34 713

SILICON: T40 714

SILVER 42, 111, 328

SILVER ALLOYS 680

SILVER BASE ALLOYS: T4 417

SILVER CHLORIDES 111, 118

SILVER CHLORIDES: T3 659

SILVER COMPOUNDS 111

SILVER COMPOUNDS: T3 207

SILVER NITRATES: T2 234

SILVER NITRATES: T4 195

SILVER NITRATES: T5 592

SILVER: T15 713

SILVER: T17 714

SILVER: T2 104, 417

SILVER: T8 711

SILVER: T9 712

SIMULATION 19, 103, 509, 525, 590

SIMULATION: Q2 601

SIMULATORS 510

SINTERING 226, 266, 267, 557, 691

SLAGS 420

SLATER METHOD 161, 223, 683

SLOW NEUTRONS 159

SMALL ANGLE SCATTERING 250

SMELTING 643

SNR-1 REACTOR: T1 601

SODIUM 16, 17, 32, 103, 108, 151,
161, 187, 250, 272, 310, 343,
599, 652, 692

SUEJECT DESCRIPTORS INDEX

SODIUM ALLOYS 187, 590
 SODIUM ALLOYS: T2 32, 388
 SODIUM ALLOYS: T3 249
 SODIUM BROMIDES 631
 SODIUM CARBONATES 129, 396
 SODIUM CHLORIDES 27, 129, 187,
 406, 407, 452, 631, 703
 SODIUM CHLORIDES: T1 87, 88, 141,
 231, 449, 548, 586
 SODIUM CHLORIDES: T2 14, 86, 286,
 439, 504, 518, 636
 SODIUM CHLORIDES: T3 434, 637, 678
 SODIUM CHLORIDES: T4 300
 SODIUM CHLORIDES: T5 89, 546
 SODIUM COMPOUNDS 34, 35, 176
 SODIUM COMPOUNDS: T1 66, 375
 SODIUM COMPOUNDS: T4 638
 SODIUM COMPOUNDS: T5 77, 519
 SODIUM COOLED REACTORS 324, 580
 SODIUM FLUORIDES 307, 308, 314,
 363, 585
 SODIUM FLUORIDES: T2 204
 SODIUM FLUORIDES: T3 89, 299, 519,
 636, 638
 SODIUM FLUORIDES: T4 516
 SODIUM FLUORIDES: T5 299, 300
 SODIUM FLUORIDES: T8 729
 SODIUM HYDROXIDES 129
 SODIUM IODIDES: T1 154
 SODIUM NITRATES 42, 129, 233
 SODIUM NITRATES: T2 667
 SODIUM NITRATES: T3 664, 666
 SODIUM NITRATES: T5 195
 SODIUM OXIDES 35, 246
 SODIUM OXIDES: T1 383, 384
 SODIUM PERCHLORATES: T3 504
 SODIUM PHOSPHATES 381
 SODIUM TUNGSTATES: T10 516
 SODIUM TUNGSTATES: T4 637
 SODIUM: T1 243, 584
 SODIUM: T14 714
 SODIUM: T2 56, 154, 247, 324
 SODIUM: T27 711
 SODIUM: T3 35, 128, 167
 SODIUM: T30 713
 SODIUM: T4 249, 346
 SODIUM: T5 528
 SODIUM: T7 244, 580
 SODIUM: T9 367
 SOL-GEL PROCESS 557
 SOLDERING 211
 SOLID CLUSTERS 463
 SOLID FUELS 60, 332, 333, 475
 SOLID FUELS: T1 685
 SOLID SOLUTIONS 12, 44, 54, 83,
 86, 108, 174, 205, 209, 251, 266,
 267, 268, 288, 332, 333, 334,
 336, 379, 418, 480, 519, 522,
 543, 545, 555, 556, 581, 587,
 593, 602, 610, 613, 629, 631,
 638, 640, 649, 691
 SOLID WASTES 301
 SOLIDIFICATION 19

SUEJECT DESCRIPTORS INDEX

- SOLIDS 2, 29, 180, 193, 243, 395,
438, 485, 609, 705, 722
- SOLIDS: T1 229
- SOLUBILITY 21, 25, 36, 85, 101,
106, 108, 111, 128, 129, 156,
202, 245, 247, 284, 303, 305,
315, 327, 351, 380, 418, 439,
440, 452, 455, 493, 495, 528,
529, 533, 669, 680, 693, 727
- SOLUBILITY: Q1 10, 154, 225, 264,
265, 329, 347, 352, 396, 462
- SOLUBILITY: Q1,Q2 287, 379, 610
- SOLUBILITY: Q1,Q2,Q3 541
- SOLUBILITY: Q1,Q2,Q3,Q4 182, 405
- SOLUBILITY: Q1,Q2,Q3,Q4,Q5,Q6,Q7,
Q8,Q9,Q10 115
- SOLUBILITY: Q1,Q2,Q3,Q5 404
- SOLUBILITY: Q2 22, 176, 321, 342,
563, 701, 702
- SOLUBILITY: Q2,Q3 663
- SOLUBILITY: Q2,Q3,Q4,Q5,Q6 119
- SOLUBILITY: Q3 532, 587
- SOLUBILITY: Q4 392
- SOLUBILITY: Q5 94, 100
- SOLUTION HEAT 182, 418, 467, 563,
608, 645
- SOLUTION HEAT: Q1 107, 466
- SOLUTION HEAT: Q2 486
- SOLUTIONS 6, 63, 102, 154, 247,
385, 396, 406, 407, 486, 528, 660
- SOLVATION 247, 371, 529
- SOLVENT EXTRACTION 125, 135, 561,
660, 722, 723
- SOLVENT EXTRACTION: Q1 369, 370,
371, 426
- SOLVENT PROPERTIES 106, 321
- SOLVENT PROPERTIES: Q1 22, 156,
342, 486, 563, 701, 702
- SOLVENT PROPERTIES: Q1,Q2 247,
532, 587
- SOLVENT PROPERTIES: Q1,Q2,Q3 392
- SOLVENT PROPERTIES: Q1,Q2,Q3,Q4
100, 172
- SOLVENT PROPERTIES: Q1,Q4,Q6 663
- SOLVENT PROPERTIES: Q2 154, 225,
264, 265, 371
- SOLVENT PROPERTIES: Q2,Q3 352, 462
- SOLVENT PROPERTIES: Q3 642
- SOLVENT PROPERTIES: Q4 541
- SOLVENTS 95, 247, 277
- SORPTIVE PROPERTIES: Q1,Q2,Q3 217
- SOUND WAVES 460
- SPACE 536
- SPACE GROUPS 419
- SPATIAL DISTRIBUTION 316
- SPECIFIC HEAT 9, 123, 179, 206,
231, 243, 297, 366, 367, 384,
454, 456, 459, 577, 578, 606,
613, 615, 616, 651
- SPECIFIC HEAT: Q1 126, 184, 261,
262, 355, 414, 415, 586, 617
- SPECIFIC HEAT: Q1,Q2 291, 383,
560, 619, 620
- SPECIFIC HEAT: Q1,Q2,Q3 594, 605
- SPECIFIC HEAT: Q1,Q2,Q3,Q4,Q5,Q6,
Q7,Q8,Q9,Q10 244
- SPECIFIC HEAT: Q1,Q2,Q3,Q4,Q5,Q6
403
- SPECIFIC HEAT: Q2 78

SUBJECT DESCRIPTORS INDEX

- SPECIFIC HEAT: Q2,Q3 487
- SPECTRA 14, 130, 280, 359, 463, 484, 488, 492, 539, 562, 589, 598, 609, 657, 664, 674, 696
- SPECTRAL DENSITY: Q2,Q3 229
- SPECTRAL SHIFT 219, 285, 363, 488
- SPECTROPHOTOMETERS: T 208, 331
- SPECTROPHOTOMETRY 27, 360, 385, 386, 424
- SPECTROPHOTOMETRY: Q1 447, 490
- SPECTROPHOTOMETRY: Q1,Q2 660
- SPECTROPHOTOMETRY: Q10 711
- SPECTROPHOTOMETRY: Q2,Q3 623
- SPECTROPHOTOMETRY: Q2,Q3,Q4,Q5,Q6, Q7,Q8,Q9 714
- SPECTROPHOTOMETRY: Q2,Q4,Q5,Q6 712
- SPECTROPHOTOMETRY: Q9,Q10,Q11,Q12, Q13 713
- SPECTROPHOTOMETRY: T 632
- SPECTROSCOPY 2, 204, 232, 398, 672
- SPECTROSCOPY: Q1 44
- SPENT FUEL CASKS: T7 722
- SPENT FUEL ELEMENTS 75, 311
- SPENT FUEL ELEMENTS: T2 453, 722
- SPENT FUEL ELEMENTS: T2,Q1 632
- SPENT FUEL ELEMENTS: T4 135
- SPENT FUELS 79, 131, 453, 558, 632
- SPENT FUELS: M2 236
- SPENT FUELS: Q1,Q2,Q3 135
- SPENT FUELS: T1 69, 358
- SPENT FUELS: T7,Q1 94
- SPIN 130, 368, 627
- SPIN WAVES 368
- SPIN-LATTICE RELAXATION: Q3 363
- SPINELS: T4 628
- SQUARE-WELL POTENTIAL 210, 442
- STABILITY 16, 175, 312, 313, 314, 329, 332, 334, 439, 561, 727
- STABILITY: Q1,Q2,Q3,Q4 164
- STABILITY: Q1,Q2,Q3,Q4,Q5 333
- STABILITY: Q1,Q2,Q3,Q4,Q5,Q6,Q7,Q8, Q9 531
- STABILITY: Q8,Q9 729
- STACKING FAULTS 560
- STAINLESS STEEL-16-8-2: T2 692
- STAINLESS STEEL-304: T1 692
- STAINLESS STEEL-304: T2 580
- STAINLESS STEEL-304: T3 309
- STAINLESS STEEL-304: T7 367
- STAINLESS STEEL-316: T1 324
- STAINLESS STEEL-316: T3 580
- STAINLESS STEEL-316: T4 309, 692
- STAINLESS STEEL-316: T8 367
- STAINLESS STEEL-321: T4 580
- STAINLESS STEEL-347: T5 580
- STAINLESS STEELS 583, 650
- STAINLESS STEELS: M3 311
- STAINLESS STEELS: T4 644
- STARCH 381
- STARK EFFECT 664
- STATISTICAL MECHANICS 550

SUBJECT DESCRIPTORS INDEX

STATISTICAL MECHANICS: Q1 279

STEAM 580

STEELS 211, 212, 454, 569

STEELS: T1 643

STCICHIOMETRY 2, 281, 401, 448,
482

STCICHIOMETRY: Q1,Q2 316

STCICHIOMETRY: Q1,Q2,Q3 626

STCRAGE LIFE 119

STRAIN HARDENING 209

STRAINS 719

STRESS CORROSION: Q1 604

STRESSES 212, 719

STRONTIUM ALLCYS: T1 446

STRONTIUM BROMIDES: T4 240

STRONTIUM CHLORIDES 264, 495

STRONTIUM CHLORIDES: T2 97

STRONTIUM CHLORIDES: T3 06, 07,
88, 286

STRONTIUM CHLORIDES: T4 89, 139

STRONTIUM FLUORIDES 307, 534

STRONTIUM FLUORIDES: T1 410, 593

STRONTIUM FLUORIDES: T2 89

STRONTIUM FLUORIDES: T4 419

STRONTIUM FLUORIDES: T5 499

STRONTIUM IODIDES: T2 241

STRONTIUM ISOTOPES: T4 534

STRONTIUM NITRATES: T1 347, 351

STRONTIUM 89 347

STRONTIUM: T35 712

STRONTIUM: T42 714

STRUCTURAL CHEMICAL ANALYSIS 20,
35, 80, 81, 102, 127, 195, 278,
330, 359, 531, 661STRUCTURAL CHEMICAL ANALYSIS: Q2
61STRUCTURAL CHEMICAL ANALYSIS: Q1,
Q2,Q3,Q4,Q5,Q6,Q7,Q8,Q9 66STRUCTURAL CHEMICAL ANALYSIS: Q1
325STRUCTURAL CHEMICAL ANALYSIS: Q1,
Q2,Q3,Q4,Q5 419STRUCTURAL CHEMICAL ANALYSIS: Q1,
Q2,Q3,Q4,Q5,Q6 501STRUCTURAL CHEMICAL ANALYSIS: Q1,
Q2,Q3 627STRUCTURAL CHEMICAL ANALYSIS: Q1,Q3
695

STRUCTURAL MODELS 157, 609

SUBLIMATION 93, 391, 481, 618

SUBLIMATION HEAT 188, 216, 474,
475, 477, 479, 481

SUBLIMATION HEAT: Q1,Q2 480

SUBLIMATION: Q1,Q2 296

SUBSTRATES 327

SULFATES 429

SULFIDES 429

SULFOXIDES 426

SULFUR 162

SULFUR DIOXIDE 722

SULFUR: T1 642

SULFUR: T10 711

SULFUR: T12 713

SULFUR: T6 712

SUEJECT DESCRIPTORS INDEX

SULFUR: T9 714	TANTALUM CHLORIDES: T3 555
SUPER KUKLA REACTOR 619	TANTALUM CHLORIDES: T6 499, 561
SURFACE IONIZATION 534	TANTALUM COMPLEXES: T7 561
SURFACE PROPERTIES 193, 211	TANTALUM FLUORIDES: T2 170
SURFACE PROPERTIES: Q1,Q2 5	TANTALUM FLUORIDES: T4 299
SURFACE TENSION 108, 141	TANTALUM IODIDES: T8 561
SURFACE TENSION: Q1 472	TANTALUM NITRIDES: T4 508
SURFACE TENSION: Q1,Q2 142, 697	TANTALUM OXIDES: T2 183
SURFACE TENSION: Q1,Q2,Q3 167	TANTALUM: M3 574
SURFACE TENSION: Q1,Q2,Q3,Q4 1	TANTALUM: T1 450
SURFACES 129, 697	TANTALUM: T2 451
SYMMETRY 13, 14, 285	TBP 369, 426
SYNERGISM 370, 371	TELLURATES 44
SYNTHESIS 521, 723	TELLURATES: T5 556
SYNTHESIS: Q1,Q2 676	TELLURIDES 176
SYNTHESIS: Q1,Q2,Q3,Q4,Q5,Q6,Q7,Q8, Q9 66	TELLURIDES: T1 92
SYNTHESIS: Q2 400	TELLURIUM 162, 472
SYSTEMS ANALYSIS 725	TELLURIUM CHLORIDES: T1 336
	TELLURIUM CHLORIDES: T3 553
	TELLURIUM COMPOUNDS 176
	TELLURIUM COMPOUNDS: T1 221
	TELLURIUM COMPOUNDS: T2 571
	TELLURIUM FLUORIDES 571
	TELLURIUM SULFIDES: T1 162
	TEMPERATURE DEPENDENCE 1, 3, 5, 9, 31, 38, 45, 46, 49, 63, 97, 99, 100, 101, 108, 110, 114, 122, 123, 126, 132, 133, 136, 141, 142, 145, 146, 168, 170, 177, 182, 190, 193, 195, 206, 224, 233, 242, 243, 244, 247, 251, 262, 264, 265, 272, 284, 287, 296, 313, 327, 334, 335, 340,

*** T ***

TABLES 2, 161, 185, 280, 284, 339,
531, 651, 681

TANTALUM 193, 582, 583

TANTALUM ALLOYS: T5 115

TANTALUM BROMIDES: T5 561

TANTALUM CARBIDES: T3 508

TANTALUM CHLORIDES 449, 450

TANTALUM CHLORIDES: T1 451

SUBJECT DESCRIPTORS INDEX

- TEMPERATURE DEPENDENCE 341, 347,
363, 366, 372, 389, 391, 400,
403, 410, 414, 415, 418, 432,
433, 434, 442, 445, 449, 451,
463, 472, 474, 475, 478, 479,
480, 481, 509, 532, 538, 558,
560, 564, 568, 575, 590, 591,
594, 599, 606, 613, 616, 621,
622, 626, 628, 643, 669, 675,
678, 681, 691, 702, 703, 727
- TEMPERATURE EFFECTS 66, 345, 401,
437
- TEMPERATURE GRADIENTS 202, 401
- TEMPERATURE MEASUREMENT 132, 475,
478, 619
- TENSILE PROPERTIES 580
- TENSILE PROPERTIES: Q1 471
- TENSILE PROPERTIES: Q3,Q4 692
- TENSORS 492
- TERBIUM BORIDES: T5 215
- TERBIUM CHLORIDES: T2 608
- TERBIUM COMPLEXES 371
- TERBIUM COMPOUNDS 168
- TERBIUM COMPOUNDS: T2 589
- TERBIUM COMPOUNDS: T4 639
- TERBIUM COMPOUNDS: T7 82
- TERBIUM FLUORIDES: T3 312, 603
- TERBIUM FLUORIDES: T7 382
- TERBIUM HYDRIDES: T2 418
- TERBIUM OXIDES: T1 183
- TERBIUM 160 369, 371
- TERBIUM: T1 369, 371
- TERBIUM: T11 125
- TERBIUM: T7 230
- TERBIUM: T8 688
- TERNARY ALLOY SYSTEMS 162, 245,
340, 436, 478, 523, 606, 645,
650, 671
- TESTING 380
- TETRAGONAL LATTICES 330, 379, 549,
671
- TEXTURE 177
- THALLIUM 328
- THALLIUM ALLOYS: T3 523
- THALLIUM BROMIDES: T2 411
- THALLIUM CHLORIDES: T1 411
- THALLIUM CHLORIDES: T2 93
- THALLIUM COMPLEXES: T3 314
- THALLIUM COMPOUNDS: T1 278
- THALLIUM COMPOUNDS: T4 556
- THALLIUM NITRATES: T3 592
- THERMAL ANALYSIS 22, 26, 45, 89,
90, 278, 299, 300, 435, 469, 482,
530, 581, 629
- THERMAL BATTERIES 117
- THERMAL BATTERIES: T1 116
- THERMAL CONDUCTIVITY 9, 121, 226,
272, 346, 366, 367, 454, 456,
502, 577, 578, 613, 643, 651
- THERMAL CONDUCTIVITY: Q1 132, 354,
459
- THERMAL CONDUCTIVITY: Q1,Q2 124
- THERMAL CONDUCTIVITY: Q2 560
- THERMAL DIFFUSION 401
- THERMAL DIFFUSIVITY 9, 121, 366
- THERMAL DIFFUSIVITY: Q1 132, 459
- THERMAL EQUILIBRIUM 645, 697

SUEJECT DESCRIPTORS INDEX

THERMAL EXPANSION 9, 231, 266,
310, 456, 569, 579, 612, 656

THERMAL EXPANSION: Q1 15

THERMAL EXPANSION: Q1,Q2,Q3 242

THERMAL EXPANSION: Q1,Q2,Q3,Q4,Q5,
Q6 403

THERMAL EXPANSION: Q1,Q2,Q3,Q4,Q5,
Q6,Q7,Q8,Q9,Q10 508

THERMAL EXPANSION: Q1,T2 573

THERMAL EXPANSION: T3,Q1,Q2 152

THERMAL GRAVIMETRIC ANALYSIS 207,
312, 393, 427, 561, 626, 676

THERMAL GRAVIMETRIC ANALYSIS: Q1,Q2
149

THERMAL POWER PLANTS: T1 343

THERMAL TESTING 531

THERMOCHEMICAL PROCESSES 238

THERMODYNAMIC ACTIVITY 291, 465

THERMODYNAMIC ACTIVITY: Q1 586

THERMODYNAMIC ACTIVITY: Q1,Q2,Q3
523

THERMODYNAMIC CYCLES: Q1 343

THERMODYNAMIC MODEL 100, 661

THERMODYNAMIC PROPERTIES 50, 56,
118, 137, 158, 238, 287, 303,
398, 454, 521, 551, 579, 611,
626, 634, 644

THERMODYNAMIC PROPERTIES: Q1 6, 9,
121, 163, 302, 366, 446, 550,
615, 616, 642, 658, 705

THERMODYNAMIC PROPERTIES: Q1,Q2,Q3
2

THERMODYNAMIC PROPERTIES: Q1,Q2 3,
7, 32

THERMODYNAMIC PROPERTIES: Q1,Q2,Q3
36

THERMODYNAMIC PROPERTIES: Q1,Q2
39, 96, 133

THERMODYNAMIC PROPERTIES: Q1,Q2,Q3,
Q4 200

THERMODYNAMIC PROPERTIES: Q1,Q2
220

THERMODYNAMIC PROPERTIES: Q1,Q2,Q3
231

THERMODYNAMIC PROPERTIES: Q1,Q2,Q3,
Q4 240

THERMODYNAMIC PROPERTIES: Q1,Q2,Q3,
Q4,Q5 241, 243

THERMODYNAMIC PROPERTIES: Q1,Q2
284, 339

THERMODYNAMIC PROPERTIES: Q1,Q2,Q3
340

THERMODYNAMIC PROPERTIES: Q1,Q2
384, 388

THERMODYNAMIC PROPERTIES: Q1,Q2,Q3
397, 406

THERMODYNAMIC PROPERTIES: Q1,Q2
420, 433

THERMODYNAMIC PROPERTIES: Q1,Q2,Q3
436

THERMODYNAMIC PROPERTIES: Q1,Q2
513, 522, 524, 577, 578, 591

THERMODYNAMIC PROPERTIES: Q1,Q2,Q3
606

THERMODYNAMIC PROPERTIES: Q1,Q2,Q3,
Q4 613

THERMODYNAMIC PROPERTIES: Q1,Q2,Q3
661

THERMODYNAMIC PROPERTIES: Q2 71,
297

THERMODYNAMIC PROPERTIES: Q2,Q3,Q1
175

THERMODYNAMIC PROPERTIES: Q2,Q3
274, 542, 585

SUBJECT DESCRIPTORS INDEX

THERMODYNAMIC PROPERTIES: Q3 343,
448THERMODYNAMIC PROPERTIES: Q4,Q5
646THERMODYNAMIC PROPERTIES: Q6,Q7,Q1,
Q2,Q3 281THERMODYNAMICS 36, 55, 63, 89,
110, 147, 222, 284, 304, 312,
372, 397, 586, 646, 647, 654,
705, 706

THERMODYNAMICS: Q1 442

THERMODYNAMICS: Q1,Q2 134, 672

THERMODYNAMICS: Q2 474, 537

THERMOELECTRIC GENERATORS 15

THERMOELECTRIC PROPERTIES 346

THERMONUCLEAR REACTOR COOLING
SYSTEMS 10THERMONUCLEAR REACTOR MATERIALS
405, 663, 702

THERMONUCLEAR REACTORS 10

THICKNESS 400

THIOCYANATES 426, 672

THOREX PROCESS: T3 722

THORIUM 18, 188, 259, 349

THORIUM ALLOYS: T1 679

THORIUM ALLOYS: T2 392, 522

THORIUM CARBIDES 679

THORIUM CARBIDES: M6 612

THORIUM CARBIDES: T1 126, 245, 259

THORIUM CARBIDES: T5 508

THORIUM CHLORIDES 703

THORIUM CHLORIDES: T1 407

THORIUM CHLORIDES: T2 488, 496

THORIUM CHLORIDES: T3 143

THORIUM FLUORIDES 79, 176, 585

THORIUM FLUORIDES: T1 656

THORIUM FLUORIDES: T2 521

THORIUM HYDRIDES: T4 363

THORIUM IONS 261

THORIUM NITRIDES: M7 612

THORIUM NITRIDES: T2 259

THORIUM NITRIDES: T6 508

THORIUM OXIDES 23, 63, 64, 214,
535, 542, 658

THORIUM OXIDES: M1 662

THORIUM OXIDES: M2 5

THORIUM OXIDES: M8 612

THORIUM OXIDES: T1 193, 223, 239,
389, 431, 652THORIUM OXIDES: T2 232, 242, 398,
568, 613

THORIUM OXIDES: T3 595, 710

THORIUM OXIDES: T4 717

THORIUM OXIDES: T5 183

THORIUM OXIDES: T8 303

THORIUM PHOSPHIDES: M9 612

THORIUM PHOSPHIDES: T3 259

THORIUM SULFIDES: M10 612

THORIUM SULFIDES: T4 259

THORIUM 234 447

THORIUM: M2 662

THORIUM: T1 447, 683, 703

THORIUM: T3 717

SUBJECT DESCRIPTORS INDEX

THORIUM: T4 714
 THORIUM: T41 711
 THORIUM: T42 712
 THORIUM: T43 713
 THULIUM FLUORIDES: T11 382
 THULIUM: T1 696
 THULIUM: T12 125
 TIME DEPENDENCE 63, 64, 193, 292, 400, 437, 691
 TIME-OF-FLIGHT METHOD 609, 653
 TIN 211, 212
 TIN ALLOYS 269, 704
 TIN ALLOYS: M3 559
 TIN ALLOYS: T1 284
 TIN ALLOYS: T2 166, 605
 TIN ALLOYS: T3 436, 606
 TIN ALLOYS: T4 177, 328, 575
 TIN ALLOYS: T5 91, 403
 TIN 119: M1 559
 TIN: T1 328
 TIN: T18 104
 TIN: T3 177, 403
 TIN: T33 711
 TIN: T34 712
 TIN: T35 713
 TIN: T41 714
 TITANIUM 25, 422, 582, 683
 TITANIUM BASE ALLOYS: T1 542
 TITANIUM BORIDES: T3 531
 TITANIUM CHLORIDES: T2 548
 TITANIUM COMPOUNDS: T3 114, 173
 TITANIUM OXIDES: T1 191, 264
 TITANIUM OXIDES: T2 267
 TITANIUM OXIDES: T3 239
 TITANIUM OXIDES: T6 372
 TITANIUM: M4 47
 TITANIUM: T1 687
 TITANIUM: T19 104
 TITANIUM: T34 711
 TITANIUM: T36 712, 713
 TITANIUM: T4 327
 TITANIUM: T43 714
 TITANIUM: T8 244
 TITRATION 349
 TITRATION: Q4, Q45 712
 TITRATION: Q4, Q9 713
 TITRATION: Q7, Q2 711
 TOA 125
 TOA: T1 113
 TOLUENE 111, 118
 TRACE AMOUNTS 104, 125, 711, 712, 713, 714
 TRACER TECHNIQUES 68, 347, 369, 371, 441, 447
 TRACER TECHNIQUES: Q1 76
 TRANSIENTS 60
 TRANSIENTS: Q1, Q2 456
 TRANSITION ELEMENT ALLOYS 11
 TRANSITION ELEMENT ALLOYS: T1 425,

SUBJECT DESCRIPTORS INDEX

TRANSITION ELEMENT ALLOYS: T1 686	TUNGSTEN 480, 649
TRANSITION ELEMENT ALLOYS: T3 392	TUNGSTEN ADDITIONS 645
TRANSITION ELEMENT COMPOUNDS 596	TUNGSTEN ALLOYS: T4 100, 115
TRANSITION ELEMENT COMPOUNDS: T1 582	TUNGSTEN CARBIDES: T2 648
TRANSITION ELEMENTS 183, 500, 686, 687	TUNGSTEN CHLORIDES: T4 554, 555
TRANSITION ELEMENTS: T1 422, 615, 705	TUNGSTEN COMPOUNDS 531
TRANSITION ELEMENTS: T2 412	TUNGSTEN FLUORIDES: T2 51, 52, 397
TRANSITION TEMPERATURE 261, 262, 313, 361, 610	TUNGSTEN FLUORIDES: T7 499
TRANSMISSION 249	TUNGSTEN OXIDES 202
TRANSPLUTONIUM ELEMENTS 723	TUNGSTEN OXIDES: T6 191
TRANSPORT: Q1 85	TUNGSTEN: M4 574
TRANSURANIUM ELEMENTS 453	TUNGSTEN: T12 714
TRANSURANIUM ELEMENTS: M4 662	TUNGSTEN: T13 713
TRANSURANIUM ELEMENTS: T18 730	TUNGSTEN: T3 156, 198, 645
TRANSURANIUM ELEMENTS: T7 501	TUNGSTEN: T38 712
TRAPPING: Q8 281	TUNGSTEN: T5 202
TRICLINIC LATTICES 81	TUNGSTEN: T9 115
TRITIUM 176, 529, 722	TURBULENT FLOW 502
TRITIUM IONS: T8 281	TWO-DIMENSIONAL CALCULATIONS 601
TTA 369, 660	TWO-PHASE FLOW 343
TUBES 437, 502	
TUNGSTATES: M1 338	*** U ***
TUNGSTATES: M3 168	ULTRAHIGH TEMPERATURE 72, 73, 297, 475, 476, 477
TUNGSTATES: T2 102, 114, 151, 185, 639	ULTRALOW TEMPERATURE 126, 130, 383, 384, 414, 432, 487, 594, 617, 677, 696
TUNGSTATES: T3 518	ULTRASONIC WAVES 605, 606
TUNGSTATES: T5 516	ULTRASONIC WAVES: Q1, Q2, Q3, Q4, Q5, Q6 403

SUBJECT DESCRIPTORS INDEX

ULTRAVIOLET RADIATION 345
 ULTRAVIOLET SPECTRA: Q1 413
 ULTRAVIOLET SPECTRA: Q1,Q2,Q3 16, 17
 ULTRAVIOLET SPECTRA: Q2 345
 URANATES: T2 278
 URANIUM 131, 162, 175, 188, 214, 237, 349, 373, 453, 500, 509, 560, 632, 646, 649, 715, 716
 URANIUM ALLOYS: M2 646
 URANIUM ALLOYS: T2 284
 URANIUM ALLOYS: T5 323
 URANIUM CARBIDES 315, 456
 URANIUM CARBIDES: M1 132, 649
 URANIUM CARBIDES: M11 612
 URANIUM CARBIDES: M2 503
 URANIUM CARBIDES: M4 646
 URANIUM CARBIDES: T1 83, 121, 124, 134, 179, 209, 251, 333, 379, 442, 577, 579, 644, 645, 676
 URANIUM CARBIDES: T2 84, 332, 334, 362, 448, 479, 480, 481, 647, 650, 691
 URANIUM CARBIDES: T3 367, 613, 648
 URANIUM CARBIDES: T4 323
 URANIUM CARBIDES: T6 281
 URANIUM CARBIDES: T7 508
 URANIUM CHLORIDES 194, 292, 293
 URANIUM CHLORIDES: M1 406
 URANIUM CHLORIDES: T1 81, 139
 URANIUM CHLORIDES: T2 13, 141, 142, 145, 146, 199
 URANIUM CHLORIDES: T5 562
 URANIUM CHLORIDES: T6 360
 URANIUM COMPLEXES 81, 632
 URANIUM COMPLEXES: T1 194
 URANIUM COMPLEXES: T3 562
 URANIUM COMPOUNDS 408, 424, 649
 URANIUM COMPOUNDS: A7 176
 URANIUM COMPOUNDS: M7 662
 URANIUM COMPOUNDS: T1 571
 URANIUM COMPOUNDS: T2 409
 URANIUM COMPOUNDS: T5 66
 URANIUM DIOXIDE 135, 214, 220, 235, 315, 349, 416, 454, 455, 470, 503, 510, 685, 699, 700, 719
 URANIUM DIOXIDE: M1 120, 218, 366
 URANIUM DIOXIDE: T1 12, 50, 180, 226, 266, 267, 401, 441, 445, 600, 618, 697, 711, 713
 URANIUM DIOXIDE: T16 125
 URANIUM DIOXIDE: T2 72, 74, 109, 110, 175, 223, 225, 281, 297, 431, 475, 477, 557
 URANIUM DIOXIDE: T3 60, 73, 474, 537, 641, 676, 725
 URANIUM DIOXIDE: T4 243
 URANIUM DIOXIDE: T5 367
 URANIUM DIOXIDE: T9 244
 URANIUM FLUORIDES 307, 571, 585
 URANIUM FLUORIDES: T1 350, 409
 URANIUM FLUORIDES: T2 238
 URANIUM FLUORIDES: T9 66
 URANIUM HEXAFLUORIDE 70, 699, 700
 URANIUM HEXAFLUORIDE: M2 345

SUBJECT DESCRIPTORS INDEX

URANIUM HEXAFLUORIDE: T1 285, 302, 413	URANIUM OXIDES: T8 191
URANIUM HEXAFLUORIDE: T2 78	URANIUM PHOSPHIDES: M14 612
URANIUM HEXAFLUORIDE: T3 51, 52, 397, 724	URANIUM SULFIDES: M15 612
URANIUM HEXAFLUORIDE: T6 323	URANIUM SULFIDES: T2 162, 379
URANIUM ISOTOPES: T38 711	URANIUM TETRAFLUORIDE: A3 176
URANIUM ISOTOPES: T5 713	URANIUM TETRAFLUORIDE: T1 238
URANIUM NITRIDES: M12 612	URANIUM TRIOXIDE 220, 278
URANIUM NITRIDES: M2 400, 649	URANIUM TRIOXIDE: T2 50, 641
URANIUM NITRIDES: T1 219, 354, 368, 373, 578, 611	URANIUM 233: T1 232
URANIUM NITRIDES: T2 251, 333, 579, 581	URANIUM 235 285
URANIUM NITRIDES: T3 332, 334	URANIUM 235: T1 78, 560
URANIUM NITRIDES: T8 508	URANIUM 235: T4 715
URANIUM OXIDES 281, 571, 684	URANIUM 235: T5 690
URANIUM OXIDES U3O8 278	URANIUM 236: T5 715
URANIUM OXIDES U3O8: T1 262	URANIUM 238 78, 285
URANIUM OXIDES U3O8: T2 12	URANIUM 238: T10 715
URANIUM OXIDES: M1 325, 395, 503	URANIUM 238: T2 560
URANIUM OXIDES: M13 612	URANIUM-ALPHA: M3 662
URANIUM OXIDES: M2 218, 220	URANIUM: M1 70, 400
URANIUM OXIDES: M9 662	URANIUM: M6 662
URANIUM OXIDES: T1 260, 261, 362, 448, 459, 476	URANIUM: T1 235, 293, 408, 424, 699
URANIUM OXIDES: T2 30, 133, 149, 169, 316, 383, 384, 478, 565	URANIUM: T2 510, 619, 620, 700, 711
URANIUM OXIDES: T3 12, 398, 626	URANIUM: T3 360, 683, 713, 714
URANIUM OXIDES: T4 72, 613	URANIUM: T4 83, 213, 690
URANIUM OXIDES: T5 239, 597	URANIUM: T5 717
URANIUM OXIDES: T6 202	URANYL CHLORIDES: T2 27
	URANYL COMPLEXES: T1 498, 684
	URANYL COMPOUNDS 30

SUBJECT DESCRIPTORS INDEX

URANYL COMPOUNDS: T1 483
 URANYL FLUORIDES: T5 127
 URANYL NITRATES 722
 USES: Q1,Q2,Q3 595
 USSR 328

*** V ***

VACANCIES 133, 213, 389, 626
 VACANCIES: Q1,Q2 74, 565
 VACUUM MELTING 588, 643
 VACUUM SYSTEMS 583
 VALENCE 28, 29, 48, 50, 61, 161, 205, 224, 279, 423, 438, 450, 451, 483, 499, 538, 582, 598, 674, 705
 VALENCE: Q7 176
 VANADATES: T3 317
 VANADIUM 63, 422, 582, 615
 VANADIUM ALLOYS: T1 166
 VANADIUM CHLORIDES: T8 499
 VANADIUM COMPLEXES: T1 672
 VANADIUM OXIDES 63
 VANADIUM OXIDES: M4 372
 VANADIUM OXIDES: T1 356
 VANADIUM OXIDES: T3 191, 394
 VANADIUM OXIDES: T5 303
 VANADIUM: M3 47
 VANADIUM: T10 244
 VANADIUM: T2 687
 VANADIUM: T20 104

VANADIUM: T3 115, 193
 VANADIUM: T35 711
 VANADIUM: T37 712, 713
 VANADIUM: T44 714
 VANADIUM: T5 243
 VAPOR CONDENSATION 724
 VAPOR PRESSURE 3, 39, 120, 136, 180, 251, 312, 340, 367, 397, 399, 433, 472, 479, 480, 482, 524, 546, 577, 578, 626, 681
 VAPOR PRESSURE: Q1 99
 VAPOR PRESSURE: Q1,Q2 84, 93, 97, 98, 170, 295, 391, 475, 476, 477, 478, 481, 618
 VAPOR PRESSURE: Q1,Q2,Q3,Q4 83
 VAPOR PRESSURE: Q2 56
 VAPOR PRESSURE: Q2,Q3 60, 73
 VAPOR PRESSURE: Q2,Q3,Q4 72
 VAPOR PRESSURE: Q3 537
 VAPOR PRESSURE: Q3,Q4 474
 VAPORIZATION HEAT 9, 474, 475, 477, 615
 VAPORIZATION HEAT: Q1 445
 VAPORIZATION HEAT: Q1,Q2 295
 VAPORS 2, 176, 202, 216, 297, 354, 433, 457, 622, 681
 VAPORS: T2 124
 VARIATIONAL METHODS 584
 VARIATIONS 107, 441, 686
 VEGARD LAW 205, 334
 VELOCITY 460, 502
 VENTILATION 722

SUBJECT DESCRIPTORS INDEX

VERY HIGH PRESSURE 31, 213, 325,
363, 421, 599, 677

VERY HIGH TEMPERATURE 1, 5, 9, 10,
18, 39, 49, 50, 54, 60, 62, 72,
74, 83, 84, 97, 100, 105, 107,
109, 110, 114, 120, 121, 132,
141, 159, 162, 164, 166, 175,
179, 182, 201, 204, 208, 224,
242, 243, 244, 251, 281, 284,
287, 290, 291, 297, 302, 304,
325, 327, 331, 334, 342, 345,
352, 355, 375, 379, 399, 400,
405, 420, 440, 441, 445, 461,
462, 463, 471, 474, 475, 476,
478, 479, 480, 481, 482, 496,
503, 508, 510, 542, 573, 574,
581, 587, 591, 604, 610, 611,
617, 618, 620, 622, 626, 634,
647, 648, 654, 658, 669, 679,
681, 693, 702

VERY LOW TEMPERATURE 126, 184,
218, 383, 414, 432, 487, 497,
560, 594, 617

VIBRATIONAL STATES 41, 102, 219,
220, 283, 285, 350, 353, 483,
497, 674

VIBRATIONAL STATES: Q1 46

VIBRATIONAL STATES: Q1,Q2 185

VIBRATIONAL STATES: Q1,Q2,Q3 52

VIBRATIONAL STATES: Q1,Q2,Q3,Q4,Q5
127

VICKERS HARDNESS 209

VISCOSITY 19, 141, 159, 277, 313,
327, 328, 367, 381, 454, 502,
564, 577, 578, 616, 621, 643, 710

VISCOSITY: Q1,Q2 142, 145, 146,
622

VISCOSITY: Q1,Q2,Q3 24, 576

VISCOSITY: Q1,Q2,Q3,Q4 346

VISCOSITY: Q1,Q2,Q3,Q4,Q5 112, 592

VISCOSITY: Q1,Q3 681

VISIBLE RADIATION 47

VITRIFICATION 666

VOLATILITY 457, 571, 722

VOLOXIDATION PROCESS 94

VOLTAMETRY 42, 314, 711

VOLTAMETRY: Q3,Q4 713

VOLTAMETRY: Q45 712

VOLUME 1, 217

*** W ***

WALLS 437

WATER 236, 467

WATER COOLED REACTORS: T2 455

WATER VAPOR 105

WATER: T4 341

WATER: T8 580

WAVE FUNCTIONS 157, 161, 189, 683

WAVELENGTHS 374

WEAR 115

WETTABILITY: Q1,Q2,Q3,Q4 101

WETTABILITY: Q4 327

WIGNER-SEITZ METHOD 161

WIRES 380

WORK FUNCTIONS: Q1 583

WVER-2 REACTOR: T1 632

*** X ***

X RADIATION 223, 598

X-RAY DIFFRACTION 21, 35, 45, 62,

SUBJECT DESCRIPTORS INDEX

X-RAY DIFFRACTION 66, 80, 117,
132, 143, 160, 162, 164, 168,
173, 194, 207, 242, 248, 249,
262, 270, 278, 312, 317, 325,
334, 378, 379, 393, 394, 409,
419, 421, 427, 443, 469, 482,
494, 496, 498, 503, 530, 540,
547, 556, 561, 572, 593, 596,
611, 634, 638, 639, 640, 653,
657, 659, 661, 671, 687, 688,
689, 691, 695

X-RAY DIFFRACTION: Q1,Q2 250, 305

X-RAY DIFFRACTION: Q1,Q2,Q3 20

X-RAY DIFFRACTOMETERS 508

X-RAY SPECTRA 28, 662

X-RAY SPECTRA: Q1,Q2,Q3 582

X-RAY SPECTROSCOPY 26, 113, 516

XYLENES 125

*** Y ***

YIELD STRENGTH 209, 643

YOUNG MODULUS 209

YTTERBIUM 399

YTTERBIUM ALLOYS: T6 399

YTTERBIUM CHLORIDES: T2 29

YTTERBIUM FLUORIDES: T12 382

YTTERBIUM FLUORIDES: T4 41

YTTERBIUM: T13 125

YTTERBIUM: T3 224

YTTERBIUM: T8 230

YTTERBIUM: T9 688

YTTRIUM ALLOYS: T2 201

YTTRIUM BROMIDES: T2 391

YTTRIUM CHLORIDES: M5 393

YTTRIUM CHLORIDES: T1 489

YTTRIUM CHLORIDES: T2 497

YTTRIUM CHLORIDES: T3 494

YTTRIUM COMPOUNDS: T8 196

YTTRIUM FLUORIDES: T2 593, 669

YTTRIUM FLUORIDES: T7 41

YTTRIUM HYDROXIDES: M3 393

YTTRIUM NITRATES: M6 393

YTTRIUM OXIDES 63, 64, 535, 542

YTTRIUM OXIDES: M3 49

YTTRIUM OXIDES: T2 193, 468, 469,
595

YTTRIUM OXIDES: T3 242, 303, 568

YTTRIUM SULFATES: M4 393

YTTRIUM SULFIDES: T2 597

YTTRIUM: T14 125

YTTRIUM: T3 623

*** Z ***

ZEEMAN EFFECT 122, 123

ZEEMAN EFFECT: Q1,Q2,Q3 165

ZEOLITES 94

ZINC 211, 212, 417

ZINC ALLOYS 269, 704

ZINC ALLOYS: T1 436

ZINC ALLOYS: T3 340, 675

ZINC BASE ALLOYS: M9 335

ZINC CHLORIDES: T3 402

SUBJECT DESCRIPTORS INDEX

ZINC COMPLEXES: T5 314

ZINC: T21 104

ZINC: T36 711

ZINC: T38 713

ZINC: T39 712

ZINC: T45 714

ZIRCALLOY 454, 509, 719

ZIRCONATES 23

ZIRCONATES: T5 628

ZIRCONIUM 422, 582, 587, 660

ZIRCONIUM ALLOYS: T1 680

ZIRCONIUM ALLOYS: T2 7

ZIRCONIUM BASE ALLOYS: T2 587

ZIRCONIUM BORIDES: T4 531

ZIRCONIUM BROMIDES: T1 635

ZIRCONIUM CARBIDES: T4 332, 333,
334

ZIRCONIUM CARBIDES: T5 531

ZIRCONIUM CARBIDES: T9 508

ZIRCONIUM CHLORIDES: T1 27, 45

ZIRCONIUM CHLORIDES: T2 136, 635

ZIRCONIUM CHLORIDES: T3 548

ZIRCONIUM CHLORIDES: T6 546

ZIRCONIUM COMPOUNDS: T1 7, 7

ZIRCONIUM FLUORIDES 176

ZIRCONIUM FLUORIDES: M1 591

ZIRCONIUM FLUORIDES: T1 296, 521

ZIRCONIUM FLUORIDES: T2 295, 596

ZIRCONIUM FLUORIDES: T3 541, 576

ZIRCONIUM FLUORIDES: T5 419

ZIRCONIUM HYDRIDES 587

ZIRCONIUM NITRIDES: T10 508

ZIRCONIUM NITRIDES: T5 332, 333,
334

ZIRCONIUM OXIDES 470, 510

ZIRCONIUM OXIDES: T1 595

ZIRCONIUM OXIDES: T2 191, 239

ZIRCONIUM OXIDES: T3 557

ZIRCONIUM OXIDES: T6 531

ZIRCONIUM SILICATES 566

ZIRCONIUM SULFATES: M2 591

ZIRCONIUM SULFIDES: T4 597

ZIRCONIUM: T1 566

ZIRCONIUM: T2 115, 287

ZIRCONIUM: T3 510

ZIRCONIUM: T37 711

ZIRCONIUM: T39 713

ZIRCONIUM: T4 202

ZIRCONIUM: T40 712

ZIRCONIUM: T46 714

THIS PAGE
WAS INTENTIONALLY
LEFT BLANK

EDB SUBJECT CATEGORY INDEX

EDB-050400	724	EDB-300503	77, 507
EDB-050700	12, 54, 74, 169, 175, 176, 202, 220, 225, 226, 232, 266, 267, 281, 334, 366, 387, 445, 448, 475, 477, 579, 618, 644, 649	EDB-300505	563
EDB-050800	67, 69, 94, 135, 150, 236, 237, 307, 322, 455, 558, 699, 700, 722, 725	EDB-360000	489
EDB-050900	85	EDB-360100	192, 320, 675
EDB-052001	70, 301, 358	EDB-360101	26, 76, 115, 129, 290, 536, 588, 703
EDB-052002	311	EDB-360102	11, 33, 48, 100, 103, 137, 155, 157, 159, 177, 205, 210, 229, 235, 273, 313, 318, 323, 328, 359, 365, 376, 417, 425, 436, 510, 514, 522, 523, 528, 549, 584, 587, 598, 671, 679, 680, 683, 686, 687, 688, 689, 706
EDB-055001	131, 349, 600	EDB-360103	57, 212, 471, 512, 560, 575, 622, 643, 692
EDB-060100	529	EDB-360104	1, 3, 10, 18, 22, 24, 32, 39, 47, 65, 101, 138, 152, 154, 156, 161, 166, 167, 172, 182, 188, 189, 190, 198, 201, 213, 217, 224, 230, 243, 245, 248, 249, 250, 284, 287, 291, 327, 339, 340, 342, 346, 355, 373, 388, 392, 399, 403, 422, 446, 452, 460, 472, 485, 513, 524, 525, 532, 551, 559, 567, 573, 574, 583, 599, 605, 606, 610, 615, 619, 620, 621, 651, 661, 681, 693, 696, 701, 705
EDB-070300	15, 304	EDB-360105	25, 128, 181, 211, 324, 335, 400, 405, 439, 542, 580, 604, 614
EDB-152001	396	EDB-360106	662
EDB-200102	343	EDB-360200	728
EDB-210100	453, 509	EDB-360201	55, 95, 191, 333, 379, 557, 676, 691
EDB-210200	75, 632, 719	EDB-360202	28, 49, 50, 62, 84, 110, 121, 134, 171, 180, 223, 227, 228, 251, 260, 261, 303, 317, 362, 394, 398, 421, 440, 463, 503, 565, 625, 634, 641, 645, 646, 647, 648, 654, 657, 694, 697
EDB-210300	315, 470	EDB-360203	209, 348
EDB-210500	56, 79, 103, 309, 416, 527, 569, 585, 731		
EDB-220100	345		
EDB-220300	73, 214, 316, 685		
EDB-220900	19, 60, 72, 272, 297, 310, 367, 454, 456, 474, 537, 590, 601		
EDB-250600	276		
EDB-250901	37, 118, 119, 187, 282, 704, 720, 727		
EDB-250902	111, 277, 535		
EDB-250903	23, 38, 42, 116, 246, 269, 274, 275, 380, 381		
EDB-300501	71		

EDB SUBJECT CATEGORY INDEX

EDB-360204	5, 9, 83, 124, 126, 132, 133, 149, 179, 183, 184, 206, 215, 242, 252, 259, 262, 332, 344, 354, 368, 372, 389, 401, 414, 415, 441, 442, 459, 476, 478, 479, 480, 481, 482, 484, 508, 521, 531, 568, 577, 578, 582, 595, 607, 612, 613, 652, 658, 664, 710	EDB-400201	402, 404, 419, 426, 427, 429, 434, 435, 438, 443, 444, 449, 457, 458, 466, 468, 469, 486, 488, 492, 494, 495, 497, 499, 500, 504, 506, 511, 516, 518, 519, 533, 538, 540, 541, 543, 548, 550, 554, 555, 561, 570, 586, 589, 591, 592, 593, 594, 596, 597, 602, 603, 608, 616, 617, 624, 636, 637, 638, 639, 659, 665, 666, 667, 668, 672, 673, 674, 677, 695, 707, 708, 709, 717, 718, 729
EDB-360205	628		
EDB-360601	114		
EDB-360602	4, 21, 31, 45, 87, 88, 90, 162, 165, 174, 195, 197, 256, 258, 288, 330, 336, 337, 464, 473, 493, 505, 517, 520, 530, 544, 545, 546, 547, 552, 553, 556, 572, 576, 581, 609, 629, 630, 631, 633, 640, 670, 678	EDB-400203	68
EDB-360603	30, 36, 98, 130, 136, 141, 173, 200, 279, 321, 352, 357, 384, 390, 410, 411, 418, 428, 433, 461, 462, 467, 627, 635, 642, 663, 669	EDB-400400	64, 221, 234, 263, 293, 420, 430, 450, 451, 515, 721
EDB-360605	491	EDB-400700	271
EDB-400100	58, 690, 726	EDB-400702	2, 13, 27, 34, 59, 66, 78, 80, 81, 82, 107, 120, 127, 139, 140, 142, 143, 144, 145, 146, 147, 194, 199, 218, 222, 238, 278, 285, 292, 302, 325, 350, 360, 363, 383, 385, 386, 395, 406, 407, 408, 409, 424, 465, 483, 496, 498, 501, 562, 571, 611, 626, 650, 655, 656, 682, 684, 723, 730
EDB-400102	109, 711, 712, 713, 714	EDB-400703	534
EDB-400103	698	EDB-420400	158, 437, 502
EDB-400104	104, 125, 319, 423, 431, 447, 490, 566, 623, 660, 715, 716	EDB-440300	208, 331, 364
EDB-400105	308, 369, 370, 371, 526	EDB-640302	52, 219, 353, 382, 412, 413, 432, 539
EDB-400201	6, 7, 8, 14, 16, 17, 20, 29, 35, 40, 41, 43, 44, 46, 51, 53, 61, 63, 66, 89, 91, 92, 93, 96, 97, 99, 102, 105, 106, 112, 113, 117, 122, 123, 148, 151, 153, 160, 163, 164, 168, 170, 178, 185, 186, 193, 196, 203, 204, 207, 216, 231, 233, 239, 240, 241, 244, 247, 253, 254, 255, 257, 264, 265, 268, 270, 280, 283, 286, 289, 294, 295, 296, 298, 299, 300, 305, 306, 312, 314, 326, 329, 338, 341, 347, 351, 356, 361, 374, 375, 377, 378, 391, 393, 397,	EDB-652022	487
		EDB-656000	564, 653
		EDB-700209	702
		050700	9, 55, 70, 83, 109, 110, 132, 133, 180, 209, 262, 333, 345, 401, 455, 478, 480, 481, 482, 503, 510, 577, 645, 646, 648, 676, 691, 711, 712, 713, 714
		050800	79, 131, 222, 311, 358, 660, 715, 716, 724
		055001	690

EDB SUBJECT CATEGORY INDEX

060200	404	360602	77, 116, 282, 489
080201	392	360603	10, 52, 97, 225, 375, 532, 540, 702, 721
080800	418	400103	150, 271, 453, 714
210100	387, 510	400104	66, 203, 325, 374, 417, 632, 711, 712, 713, 714
210200	387, 509	400105	426, 430, 541
210500	19, 57, 60, 72, 176, 236, 237, 272, 297, 310, 367, 456, 474, 537, 580, 590, 601, 692	400201	2, 13, 22, 37, 52, 82, 90, 119, 120, 136, 140, 142, 147, 154, 156, 173, 176, 206, 218, 258, 357, 363, 364, 372, 390, 395, 396, 400, 505, 517, 528, 529, 576, 590, 633, 652, 720
220300	25, 169, 202, 612	400202	78, 353, 695
250902	234	400203	663
250903	111, 263	400301	113
300503	71, 246, 263, 389, 595	400400	277, 423, 513, 551, 586, 675
300505	71, 263	400600	301
320303	200	400702	12, 51, 52, 53, 54, 72, 148, 169, 176, 239, 244, 259, 323, 362, 373, 397, 442, 535, 597, 694, 718, 729
360100	58	400703	18
360102	1, 245, 274, 512	420400	103
360103	309, 527	440105	632
360104	5, 7, 56, 115, 181, 225, 244, 252, 290, 321, 367, 420, 436, 454, 462, 406, 487, 643	640302	374, 484, 730
360105	106, 287, 644, 654	700201	10
360200	281	700209	404, 405, 614, 663
360201	54, 226, 266, 304, 334, 400, 536, 649		
360202	73, 74, 83, 175, 266, 267, 323, 325, 348, 448, 468, 469, 475, 611, 612		
360203	387		
360204	2, 30, 72, 152, 191, 209, 218, 220, 239, 243, 244, 267, 305, 366, 367, 387, 445, 474, 477, 537, 542, 579, 585, 597, 618, 625, 647		
360206	169, 232, 662		

TITLE INDEX

AB

- Behavior of the AB-type compounds at high pressures and high temperatures* (421)
- absorption
- Optical absorption spectrum of UO_2 doped LiF * (30)
- Absorption spectra of curium (3) chloride complexes* (34)
- Optical absorption Zeeman spectroscopy of rare earth orthochromites. II. Theory* (122)
- Optical absorption Zeeman spectroscopy of rare earth orthochromites. I. ErCrO_3 * (123)
- Study of the compounds $\text{M}_3\text{UO}_2\text{F}_5$ ($\text{M} = \text{K}, \text{Rb}, \text{Cs}, \text{NH}_4$) by i.r. absorption and Raman diffusion spectrophotometry. III. Effect of cation on vibration frequency of the ion UO_2^{2+} * (127)
- Zeeman effect and satellite carrier structure in absorption spectra of pure and doped praseodymium and europium double nitrates* (165)
- High-temperature unit to the IKS-21 spectrometer for the detection of molten salts absorption spectra* (208)
- Absorption spectrum of HfCl molecule* (283)
- Flame emission and atomic absorption spectrometry. III. Elements and matrices* (319)
- Use of the SF-8 detection spectrophotometer for measuring absorption spectra of molten salts* (331)
- Quantitative study of the chemical equilibria in solution in molten LiCl-KCl (70-30 mol.%) by absorption spectrophotometry. Application to neptunium* (385)
- Determination of the $\text{Np(IV)} \rightarrow \text{Np(III)}$ normal potential in fused LiCl-CsCl (55-45%) by absorption spectrophotometry* (386)
- Hydrogen absorption of rare-earth (3d) transition intermetallic compounds* (392)
- Effect of luminescence centers on optical absorption, excitation and fluorescence spectra of CsCl:Pb * (491)

TITLE INDEX

absorption

Investigation into absorption infrared spectra of vapors under ScBr_3 , YCl_3 , LaCl_3 , GdCl_3 , ErCl_3 , LuCl_3 by means of isolation method in rare gas matrix* (497)

Use of oscillator strengths of 'supersensitive' ion transfers of lanthanides for calculation of some parameters determining the intensity of their absorption spectra* (511)

Abstracts

11. european congress on molecular spectroscopy. Abstracts* (44)

11. european congress on molecular spectroscopy. Abstracts* (46)

5th international conference on plutonium and other actinides.

Book of abstracts* (109)

Abstracts of papers and communications submitted to the 5.

all-union heat and mass transfer conference* (437)

5th international conference on plutonium and other actinides.

Book of abstracts* (620)

accident

Specialists' meeting on the behaviour of water reactor fuel elements under accident conditions* (510)

accountability

Methods for the accountability of uranium dioxide* (600)

accounting

Experience of the Central Control Laboratory (CCL) in accounting for and controlling nuclear material in Czechoslovakia* (349)

accuracy

Estimate of cryoscopic calculations accuracy from fusibility diagrams* (670)

Accurate

Accurate procedure to safeguard the fissile material content of input and output solutions of reprocessing plants* (131)

Accurate nuclear parameters derived from the Moessbauer spectrum of thulium metal* (696)

actinide

Comparison of the thermodynamic properties and high temperature chemical behavior of lanthanide and actinide oxides* (2)

Ternary UO_2 -- UO_3 -- EuO system and investigation of Eu(II) --actinide(IV)--perovskites* (50)

Preparation and some properties of actinide(III) and lanthanide(III) oxide halides, and the enthalpies of formation of $\text{PuBr}_3(\text{s})$ and $\text{PuI}_3(\text{s})$ * (82)

Bonding and the electronic structure of the actinide metals* (188)

Actinide metallocarbaborane complex; synthesis and x-ray structure determination of the bis[η^5 -

5 β -(3)-1,2-dicarbollyl]dichlorouranium(IV) dianion* (194)

Band structure studies of actinide systems* (323)

Electronic properties of actinide oxides* (398)

Some thermodynamic properties of dilute solutions of actinide chlorides in (Li-K)Cl and in (Na-K)Cl eutectics* (406)

Rare earth and actinide oxides. Thermodynamic and electron microscopy studies* (625)

XPS studies of actinide materials* (662)

actinides

Plutonium 1975 and other actinides* (9)

Plutonium 1975 and other actinides* (83)

TITLE INDEX

actinides

- 5th international conference on plutonium and other actinides* (84)
- 5th international conference on plutonium and other actinides. Book of abstracts* (109)
- Plutonium 1975 and other actinides* (110)
- Plutonium 1975 and other actinides* (133)
- Formation of actinides in irradiated HTGR fuel elements* (150)
- Electronic structure and properties of the actinides* (189)
- Plutonium 1975 and other actinides* (209)
- Plutonium 1975 and other actinides* (227)
- 5th international conference on plutonium and other actinides* (228)
- Plutonium 1975 and other actinides* (398)
- Plutonium 1975 and other actinides* (414)
- 5th international conference on plutonium and other actinides* (415)
- Plutonium 1975 and other actinides* (476)
- 5th international conference on plutonium and other actinides* (479)
- Plutonium 1975 and other actinides* (481)
- Plutonium 1975 and other actinides* (560)
- Plutonium 1975 and other actinides* (619)
- 5th international conference on plutonium and other actinides. Book of abstracts* (620)
- Electronic structure of actinides and related metals* (683)

actinoid

- Electronic structure of highly symmetric lanthanoid and actinoid compounds. VI. The electronic Raman spectrum of CsCe(III)Cl_6 * (678)

actinoids

- Electronic structure of highly symmetric complexes of lanthanoids and actinoids. Pt. 3. 'Electronic' Raman spectrum of CsU(IV)Cl_6 * (13)
- Electronic structure of highly symmetric complexes of lanthanoids and actinoids. Pt. 4. The electronic Raman spectrum of CsNaEu(III)Cl_6 * (14)

activities

- Characterisation of the chemical activities of non-metals dissolved in liquid alkali metals* (63)
- Characterisation of the chemical activities of non-metals dissolved in liquid alkali metals* (64)
- Some considerations in establishing thermodynamic activities from high temperature EMF measurements* (163)
- Partial pressures, activities and evaporation heat in the KF-ZrF_4 system* (295)
- Activities of lead and bismuth in their molten alloys* (420)

activity

- Calculation of thermodynamic activity and liquidus lines of components in binary systems of plutonium trichloride with alkali and alkaline earth metal chlorides* (147)
- Solubility and activity of oxygen in liquid indium and copper-indium alloys* (182)
- Study of the behavior of high activity waste produced during the regeneration of fast reactor fuel elements by the gaseous fluoride method* (311)

TITLE INDEX

activity

- Effect of basicity on ferrous oxide activity in complex oxide systems* (372)
- Calculation of thermodynamic activity of components in strontium liquid alloys* (446)
- Thermodynamic activity of carbon in molybdenum-containing uranium carbide* (644)
- Thermodynamic activity of carbon in molybdenum-containing uranium carbide* (650)

Adaptation

- Adaptation of the continuous cold trap system of fluidized-bed to the fluoride volatility process* (724)

additions

- Influence of small additions to the preparation and the properties of UO_2 ceramics. Part A. $\text{UO}_2/\text{Gd}_2\text{O}_3$ * (266)
- Influence of small additions on the preparation and the properties of UO_2 ceramics. Part B. The $\text{UO}_2\text{-TiO}_2$ system* (267)
- Effect of various additions on the temperature of structural transformation in liquid iron* (313)

Adhesion

- Adhesion properties of melted Al, Ga, In on solid titanium* (327)

Advanced

- Westinghouse Advanced Reactors Division* (57)
- Proceedings of the symposium and workshop on advanced battery research and design, March 22--24, 1976* (269)
- Preparation and thermochemical stability of uranium-zirconium-carbonitrides. A contribution to nuclear fuels for advanced high-temperature reactors* (333)

Advances

- Advances on thermochemistry of rare earths alloys* (65)
- Advances in nuclear science and technology. Vol. 9* (448)
- Advances in vapour pressure studies over liquid uranium plutonium oxides up to 5000K* (476)

AgCl

- Interaction in AgCl-CsCl, CsCl-KCl systems and liquidus surface of AgCl-CsCl-KCl ternary system* (659)

Agency

- Second specialist meeting on sodium/fuel interaction in fast reactors, Ispra, 21-23 November 1973. Sponsored by the Committee on the Safety of Nuclear Installations, OECD Nuclear Energy Agency, Paris, France* (272)

agitated

- Measurement of mass transfer coefficients in a mechanically agitated, nondispersing contactor operating with a molten mixture of $\text{LiF-BF}_3\text{-ThF}_4$ and molten bismuth* (79)

Al

- Gases and carbon in metals (thermodynamics, kinetics, and properties). I. Alkali metals, alkaline earth metals, light metals (Li, Na, K, Rb, Cs; Ca, Sr, Ba; Be, Mg, Al)* (192)
- Raman spectra of Al_2O_3 solutions in molten cryolite and other aluminum fluoride containing melts* (203)
- Adhesion properties of melted Al, Ga, In on solid titanium* (327)
- Wetting of certain refractories of the system $\text{MgO-Al}_2\text{O}_3\text{-ZrO}_2$ with metallic melts* (344)

TITLE INDEX

Al

Using molecular dynamics method for the evaluation of Fe-Ni and Al-Zn melts structure properties* (512)
 New representatives of $\text{CeMn}_{12}\text{Al}_8$ structural type* (549)
 Compounds with the structure of ThMn_{12} type in the R--Fe--Al systems* (671)

AlCl

Thermal investigation of NdCl_3 - AlCl_3 - KCl and NdCl_3 - AlCl_3 - CsCl system* (289)
 NdCl_3 - AlCl_3 - NaCl system* (434)

AlF

Study of chemical interaction in Pr_3AlF_6 - LaF_3 - CaF_2 system* (26)
 Density, viscosity, electric conductivity and meltability of Na_3AlF_6 - La_2O_3 and Na_3AlF_6 - Sm_2O_3 systems* (43)

algorithm

Calculated method of eutectics of multicomponent systems on base of algorithm and digital computers* (517)

alkali

Thermodynamic properties of diluted solutions of MnCl_2 in alkali metal chlorides* (6)
 X-ray diffraction study into molten bromides and iodides of alkali metals* (20)
 Temperature effect on vibrational spectra of the impurity molecular ions in alkali halides* (46)
 Characterisation of the chemical activities of non-metals dissolved in liquid alkali metals* (63)
 Characterisation of the chemical activities of non-metals dissolved in liquid alkali metals* (64)
 Structural study and properties of the alkali metal, nitrosyl, and ammonium hepta- and octafluorouranates(VI)* (66)
 Determination of vibrational contribution into excessive mixing entropy of liquid amalgams. Alkali metal amalgams* (138)
 Calculation of thermodynamic activity and liquidus lines of components in binary systems of plutonium trichloride with alkali and alkaline earth metal chlorides* (147)
 Electron spectrum of light alkali metals* (161)
 Gases and carbon in metals (thermodynamics, kinetics, and properties). I. Alkali metals, alkaline earth metals, light metals (Li, Na, K, Rb, Cs; Ca, Sr, Ba; Be, Mg, Al)* (192)
 Study of lanthanum, yttrium and erbium oxychlorides in melts of alkali metal chlorides* (196)
 Aspects of the solution chemistry of liquid alkali metals as elucidated from electrical resistivity studies* (247)
 Structure factors of liquid alkali metals* (250)
 Solubility of magnesium oxide in alkali metal chloride melt* (265)
 Redox and chemical properties of U, Np and Pu in molten alkali chlorides* (360)
 Origin of excess entropy of alkali metals halides solid solutions* (390)
 Pressure and composition of vapors over alkali metal tetrachloroferrates in chlorine atmosphere* (433)

TITLE INDEX

alkali

- Phase diagrams in the pseudobinary systems of Li_2BeF_4 , alkali fluoroborates and alkali fluorides* (473)
- Improved model for solutions of nonmetals in liquid alkali metals. Calculation of enthalpy of solution and electrical resistivity* (486)
- Raman spectroscopic studies of yttrium (III) chloride--alkali metal chloride melts and of $\text{Cs}_2\text{NaYCl}_6$ and YCl_3 solid compounds* (489)
- Influence of salt background on kinetics of BF_4^{sup} -alkali hydrolysis* (504)
- Thermodynamic analysis of melting point diagrams of systems consisting of alkali metal chlorides, thallium chloride and beryllium chloride* (505)
- Recent developments in the chemistry of solutions of salts in liquid alkali metals* (528)
- Separation of alkaline earth from alkali metal fission products in the SOLAR on-line mass spectrometer* (534)

alkaline

- Saturation vapor pressure in iron chloride-alkaline earth metal chlorides systems* (97)
- Interaction of uranium trichloride with alkaline earth metal chlorides* (139)
- Calculation of thermodynamic activity and liquidus lines of components in binary systems of plutonium trichloride with alkali and alkaline earth metal chlorides* (147)
- Gases and carbon in metals (thermodynamics, kinetics, and properties). I. Alkali metals, alkaline earth metals, light metals (Li, Na, K, Rb, Cs; Ca, Sr, Ba; Be, Mg, Al)* (192)
- Titanium dioxide solubility in melts of lithium- and alkaline earth metal chlorides* (264)
- Thermal decomposition of intermetallic compounds of platinum with alkaline earth and lanthanide metals at high temperatures* (399)
- Phase relations in the alkaline earth fluoride-yttria systems (in three parts). 1. The CaF_2 -- Y_2O_3 phase diagram* (468)
- Phase relations in the alkaline earth fluoride-yttria systems (in three parts). 2. The BaF_2 -- Y_2O_3 phase diagram* (469)
- Separation of alkaline earth from alkali metal fission products in the SOLAR on-line mass spectrometer* (534)
- Density and enthalpy of liquid alkaline metals at pressures up to 1500 bar* (599)

allowance

- The calculation of the consequences of sodium/fuel interactions with allowance for temperature gradients in the sodium* (272)

alloy

- Electrochemical machining of the molybdenum VM-1 alloy* (129)
- Studies on the negative alloy electrode* (269)
- Terminal solid solubility of hydrogen in the zirconium-2.5 weight % niobium alloy* (587)
- Refractory materials resistance to molten barium alloy* (628)
- Influence of the technology of the alloy fabrication on its properties in a liquid state* (643)

TITLE INDEX

alloys

- Density and temperature dependence of the surface tension of molten bismuth, lead, and bismuth-lead alloys* (1)
- Thermodynamic properties of the liquid La--Mg alloys* (3)
- Zirconium: Physico-chemical properties of its compounds and alloys* (7)
- Standard formation enthalpy of chloride alloys of $ZrCl_4$ --KCl (NaCl) systems* (8)
- Thermoelectric measurements in liquid alloys and electron transport heat calculations in the case of almost-free electrons* (33)
- Studies of the electronic structure of simple metals and their alloys with the aid of Compton scattering and positron annihilation using lithium, magnesium, lithium-magnesium and aluminium-zinc as examples* (48)
- Advances on thermochemistry of rare earths alloys* (65)
- Interaction of the lithium-tin alloys with the LiCl-KCl-CsCl melted eutectic mixture* (91)
- Equilibrium between Li-K-Te alloys and LiCl-KCl molten eutectic mixture* (92)
- Temperature dependence of vapor pressure of rare earth halides and their alloys* (99)
- Solubility of nitrogen in multicomponent nickel base alloys* (100)
- Formation enthalpy of liquid tin-vanadium alloys* (166)
- Atomic diffusion in alloys exposed to liquid sodium* (181)
- Solubility and activity of oxygen in liquid indium and copper-indium alloys* (182)
- Magnetic susceptibility and electronic structure in metallic and non-metallic liquid cesium-gold and cesium-antimony alloys* (190)
- Criteria for solid solution formation in NFE alloys* (205)
- Structure factor of liquid caesium and sodium-caesium alloys* (249)
- Spreading of iron-nickel alloys on ZrC and nickel-cobalt alloys on ZrN* (252)
- Structure and anodic discharge behavior of lithium-boron alloys in the LiCl-KCl eutectic melt.* (274)
- Structure of lithium--boron alloys and their performance as potential anodes for high power molten salt batteries. Progress report, 1 Aug 1974--30 Jun 1975* (275)
- Thermodynamic properties of the U-Sn-alloys riched with tin* (284)
- Interaction of atoms of rare earths-metal alloys* (320)
- Phase diagram and properties of Nd--As alloys* (321)
- Corrosion of liquid alloys of zinc with yttrium and with rare earth elements in LiCl--KCl melts* (335)
- Thermodynamic properties of Cd-Zn-Hg liquid alloys* (340)
- Structure of molten Bi--Sb alloys by means of neutron diffraction* (359)
- Theory of rare-earth alloys* (376)
- Speed of ultrasound and the thermophysical properties of the liquid metals Sn, Pb, Cd and of their binary alloys Pb:Sn and Pb:Cd* (403)
- Photoelectron spectroscopic studies of some transition metals and alloys* (417)

TITLE INDEX

alloys

- Activities of lead and bismuth in their molten alloys* (420)
 Heat of formation of solid and liquid alloys of transition metals* (425)
 Some regularities of the corrosion behavior of binary magnesium alloys* (439)
 Calculation of thermodynamic activity of components in strontium liquid alloys* (446)
 Zirconium: Physico-chemical properties of its compounds and alloys* (464)
 Thermodynamic studies of gallium--indium liquid alloys by solid state electrochemistry with oxide electrolytes* (513)
 Thermodynamic properties of liquid thorium-gallium alloys* (522)
 Thermodynamic investigation of liquid alloys in the system Cd-In-Tl* (523)
 Contribution to the problem of the thermodynamic properties and the structure of liquid Cd-In alloys* (524)
 Theory for the derivation of a single hard-sphere potential describing the overall molecular interactions in binary liquid metal alloys as a function of temperature and composition* (525)
 Some trends in constitutive equation model development for high-temperature behavior of fast-reactor structural alloys* (527)
 Nitrogen solubility in liquid nickel--niobium alloys* (532)
 Electromotive force measurements in molten lithium--magnesium alloys* (551)
 XPS valence bands of La, Ce and Gd and their aluminium alloys* (598)
 Effect of composition and structure on crevice, intergranular, and stress corrosion of some wrought Ni--Cr--Mo alloys* (604)
 Tritium sorption in lithium--bismuth and lithium--aluminum alloys* (614)
 Spectrophotometric determination of rare earths and yttrium in gallium base alloys* (623)
 Investigation of particles interaction in liquid metals and alloys* (661)
 Thermodynamic properties of solutions of hydrogen isotopes in metals and alloys of interest to fusion reactor technology* (663)
 Anodic behavior of cadmium and cadmium-zinc alloys in eutectic melt of potassium- and lithium chlorides* (675)
 Thorium: physico-chemical properties of its compounds and alloys* (679)
 Zirconium: Physico-chemical properties of its compounds and alloys* (680)
 Structure of liquid transition metals and their alloys* (686)
 Structure of liquid Ce-Ni alloys* (689)
 Lithium alloys--chlorine secondary battery. I. Basic studies on a lithium alloys--chlorine secondary battery using a molten salt as the electrolyte* (704)

almost

- Thermodiffusion measurements in liquid alloys and electron transport heat calculations in the case of almost-free electrons* (33)

alternate

- Report of task force on alternate structural materials for Liquid Metal Fast Breeder Reactors* (309)

TITLE INDEX

- aluminate Phase equilibria in Li--Na--K carbonate/aluminate systems* (77)
- aluminates Investigation of formation thermodynamic of rare earth aluminates with perovskite structure* (326)
Investigation of some rare earth aluminates by the method of infrared spectroscopy* (357)
- aluminium Studies of the electronic structure of simple metals and their alloys with the aid of Compton scattering and positron annihilation using lithium, magnesium, lithium-magnesium and aluminium-zinc as examples* (48)
Spectrographic determination of trace impurities in reactor grade aluminium* (104)
Molybdenum and tungsten interaction with liquid aluminium* (156)
Enthalpies of formation of melts of yttrium with aluminium* (201)
XPS valence bands of La, Ce and Gd and their aluminium alloys* (598)
- aluminum Structure factor of liquid aluminum and calculation of transport coefficients* (159)
Raman spectra of $\text{Al}_{2/3}\text{O}_{3/2}$ solutions in molten cryolite and other aluminum fluoride containing melts* (203)
Tritium sorption in lithium--bismuth and lithium--aluminum alloys* (614)
A Mossbauer spectroscopic study of intercalation compounds of graphite with ferric chloride and aluminum chloride* (627)
- amalgam Standard potential of sodium amalgam at 25°C* (32)
Cs-Hg amalgam thermodynamic properties* (339)
- amalgams Determination of vibrational contribution into excessive mixing entropy of liquid amalgams. Alkali metal amalgams* (138)
- ambient X-ray study of the plutonium--oxygen phase diagram at ambient temperatures up to 1100°C* (62)
- americium All-union conference on chemistry of transplutonium elements (americium, curium, berkelium, californium)* (34)
Mossbauer studies of americium compounds* (61)
Reaction of americium oxides with trivalent metal oxides* (303)
All-union conference on chemistry of transplutonium elements (americium, curium, berkelium, californium)* (307)
Investigation into americium extraction process from residues of irradiated nuclear fuel after its fluorination by SiF_4 -NaF and SiF_4 -CaF₂ melts of eutectic composition* (308)
- ammonium Structural study and properties of the alkali metal, nitrosyl, and ammonium hepta- and octafluorouranates(VI)* (66)
Thermogravimetric investigation of interaction of some rare earth oxides with ammonium bifluoride* (427)
- amorphous Structure and excitations of amorphous solids* (183)
- Amsterdam Spring meeting, 26 and 27 April 1976, Amsterdam* (250)

TITLE INDEX

analogous

Composition gradient during electrolysis in mixtures analogous to molten salt battery electrolytes* (234)

analysis

Thermal analysis of $\text{GaCl}_3 + 3\text{In} = \text{InCl} + \text{Ga}$ system* (4)

Investigation of fuel vapor pressure for fast breeder safety analysis* (60)

MATERO: Version 09. A handbook of materials properties for use in the analysis of light water reactor fuel rod behavior* (387)

Monte Carlo analysis of diffusion and thermodynamics in UO_2 * (441)

Extension of vapour pressure measurements of nuclear fuels (U, Pu) O_2 and UO_2 to 7000 K for fast reactor safety analysis* (475)

Vapour pressure measurements to 7,000 K and equation of state of oxide fuels for fast reactor safety analysis. I. Extension of vapour pressure measurements of nuclear fuels (U, Pu) O_2 and UO_2 to 7,000 K* (477)

Evaporation behavior and high-temperature thermal analysis of substoichiometric plutonium oxide in the composition range from $\text{O}/\text{Pu} = 1.51$ to 2.00 * (482)

KKR analysis of the Anderson-McMillan theory of electronic structure of liquid metals. Technical report* (485)

Thermodynamic analysis of melting point diagrams of systems consisting of alkali metal chlorides, thallium chloride and beryllium chloride* (505)

Phase analysis on simulated LWR fuel rods and its evaluation with the aid of constitution studies in the system uranium-zirconium-oxygen* (509)

Phase analysis at simulated LWR fuel rods and their interpretation by constitution investigations in the uranium-zirconium-oxygen system* (510)

Thermal analysis of the system of NaF , CaF_2 , Na_2MoO_4 , CaMoO_4 * (519)

Analysis of LMFBR explosion model experiments by means of the SURBUM-II code* (601)

Thermal analysis of NaF , NaCl , Na_2MoO_4 system* (636)

Standard methods for chemical, mass spectrometric, and spectrochemical analysis of nuclear-grade uranium dioxide powders and pellets* (711)

Standard methods for chemical, mass spectrometric, and spectrochemical analysis of nuclear-grade plutonium dioxide powders and pellets* (712)

Standard methods for chemical, mass spectrometric, and spectrochemical analysis of nuclear-grade mixed oxides (U, Pu) O_2 * (713)

Standard methods for chemical, mass spectrometric, spectrochemical, nuclear, and radiochemical analysis of nuclear-grade plutonium metal* (714)

Analysis of uranium carbides* (726)

Analytical

Analytical Chemistry Division annual progress report for period ending November 30, 1975* (423)

TITLE INDEX

Analytical

Analytical methods for fissionable materials in the nuclear fuel cycle. Progress report, July 1, 1975--September 30, 1976* (690)

Anderson

KKR analysis of the Anderson-McMillan theory of electronic structure of liquid metals. Technical report* (485)

anion

Preparational, structural and vibrational study of $M_{20}U_{50}Cl_4$, $2H_{20}(M = Rb, Cs)$. Detection of a tetranuclear anion $[(UO_2)_4O_{20}Cl_8(H_{20})_2]^{4-}$ (498)

Investigation into properties of liquidus curves of binary salt system melts where chemical compounds formation. 2. Systems of monc-monovalent and bi-bivalent cations with common anion* (708)

annihilation

Studies of the electronic structure of simple metals and their alloys with the aid of compton scattering and positron annihilation using lithium, magnesium, lithium-magnesium and aluminium-zinc as examples* (48)

Annual

Chemical Engineering Division physical inorganic chemistry. Annual report, July 1975--June 1976* (58)

Environmental sensitivity of structural metals: liquid metal embrittlement. Annual technical progress report No. 2, 25 Jun 1970--31 May 1971* (95)

Environmental sensitivity of structural metals: liquid metal embrittlement. Annual technical progress report No. 4, 1 Jun 1972--31 May 1973* (211)

Environmental sensitivity of structural metals: liquid metal embrittlement. Annual technical progress report No. 2, 25 Jun 1970--31 May 1971* (212)

Thermochemistry of gaseous compounds of metals. Annual summary report 5 Dec 74--4 Dec 75* (240)

Thermochemistry of gaseous compounds of metals. Annual summary report, 5 Dec 1975--4 Dec 1976* (241)

Radiochemistry Division annual progress report for 1973* (271)

Chemical Engineering Division fuels and materials chemistry.

Annual report, July 1975--June 1976* (281)

Chemical Engineering Division, reactor safety and physical property studies annual report, July 1974--June 1975* (367)

Eleventh annual seminar on theoretical physics, Johannesburg, 5-9 July 1976* (368)

Physical research on liquid-metal systems. Annual report, July 1975--June 1976* (405)

Analytical Chemistry Division annual progress report for period ending November 30, 1975* (423)

Research laboratories annual report 1975* (566)

Annual report, 1975* (632)

Annual report, 1976* (652)

Basic research in crystalline and noncrystalline ceramic systems. Annual report, May 1, 1975--April 1, 1976* (728)

Chemistry Division annual progress report for period ending November 1, 1975* (729)

Chemistry Division annual progress report for period ending November 1, 1975* (730)

TITLE INDEX

anode

Lithium anode limited-cycle battery investigation. Quarterly technical progress report No. 5* (37)

Lithium anode limited-cycle battery investigation. Quarterly technical progress report No. 3* (38)

anodes

Structure of lithium--boron alloys and their performance as potential anodes for high power molten salt batteries. Progress report, 1 Aug 1974--30 Jun 1975* (275)

anodic

Structure and anodic discharge behavior of lithium-boron alloys in the LiCl-KCl eutectic melt.* (274)

Electrochemistry of niobium and tantalum, (5). Anodic dissolution of tantalum and the average valence of tantalum ions in molten KCl * (450)

Anodic behavior of cadmium and cadmium-zinc alloys in eutectic melt of potassium- and lithium chlorides* (675)

anomalies

Lambda-type heat capacity anomalies in $\text{U}_{30}\text{O}_{8}$ * (262)

Thermodynamics of the lanthanide halides. II. Heat capacities and Schottky anomalies of SmCl_3 , EuCl_3 and GdCl_3 from 50 to 350 K* (594)

Anomalous

Anomalous quasielastic light scattering from metal oxide glasses* (183)

Anomalous melting of f electron metals (with attention to Pu)* (318)

Anomalous thermal property behaviour of uranium at low temperatures* (560)

anti

Partly filled shells constituting anti-bonding orbitals with higher ionization energy than their bonding counterparts* (280)

antimony

Magnetic susceptibility and electronic structure in metallic and non-metallic liquid cesium-gold and cesium-antimony alloys* (190)

Thermodynamic properties of sulphur in liquid antimony* (642)

Some structure differences of double oxides of indium and gallium with antimony(5)* (657)

apparatus

New apparatus for thermophysical measurements above 2500 K* (574)

applicability

Collection of gamma spectra data of irradiated light water moderated reactor spent fuel and study of the applicability of the methods for fuel identification. Final report for the period 1 February 1972--October 1975* (453)

Approximation

Approximation of tabulated thermodynamic data for purposes of optimization of two-phase liquid-metal power cycles* (343)

Apr

Thermochemistry of gaseous metal oxides. Final report, 1 Apr 1973--31 Dec 1975* (239)

Lithium--nickel fluoride secondary battery investigation. Quarterly technical progress report No. 1, 27 Apr--27 Jul 1966* (381)

TITLE INDEX

April

Mechanical properties test data for structural materials.
 Quarterly progress report for period ending April 30, 1977* (57)
 LMFER Fuel Reprocessing Program progress report for period April 1 to June 30, 1976* (94)
 Spring meeting, 26 and 27 April 1976, Amsterdam* (250)
 Electrochemical characterization of nonaqueous systems for secondary battery application. Quarterly report, February--April 1968* (380)
 Mechanical properties test data for structural materials.
 Quarterly progress report for period ending April 30, 1977* (471)
 Physical--chemical studies of transuranium elements. Progress report, April 1, 1976--March 31, 1977* (501)
 Fundamental studies of metal fluorination reactions. Progress report, May 1, 1976--April 31, 1977* (533)
 Mechanical properties test data for structural materials.
 Quarterly progress report for period ending April 30, 1977* (692)
 Basic research in crystalline and noncrystalline ceramic systems. Annual report, May 1, 1975--April 1, 1976* (728)

Ar

Identification of UN in Ar matrices* (219)

Argonne

Argonne National Laboratory* (692)

assemblies

Investigations on radioactive fission product correlations: gamma spectrometry measurements on spent fuel assemblies discharged from the Trino Vercellese reactor at the end of the 2nd irradiation cycle* (75)

assessment

Thermodynamic assessment of the conversion of plutonium dioxide to plutonium monocarbide* (54)
 Critical assessment of equation of state data for UO_2 * (180)

assessments

Preliminary assessments of carbide fuel pins during mild overpower transients in LMFBRs* (456)

Assignments

Assignments in the electronic spectrum of UF_6 * (413)

associated

Summary of U.S. LMFER programs on high temperature structural design and associated materials testing* (731)

atm

Thermophysical and electrophysical properties of uranium hexafluoride at temperatures of $(1-11) \times 10^3$ degree K and pressures of 0.1--100 atm* (302)

atmosphere

Stability of rare earth metaniobates in hydrogen atmosphere* (164)
 Pressure and composition of vapors over alkali metal tetrachloroferrates in chlorine atmosphere* (433)

atom

Standard method of test for atom percent fission in uranium and plutonium fuel (mass spectrometric method)* (715)
 Standard method of test for atom percent fission in uranium and plutonium fuel (neodymium-148 method)* (716)

TITLE INDEX

Atomic

Atomic structure of melts of complex system with eutectic transformations* (155)
 Atomic diffusion in alloys exposed to liquid sodium* (181)
 Flame emission and atomic absorption spectrometry. III. Elements and matrices* (319)
 Structural study of liquids with strong short-range correlation in the atomic distribution* (609)
 Structural study of liquids with strong short-range correlation in the atomic distribution* (653)

atoms

Interaction of atoms of rare earths-metal alloys* (320)
 Energies of valence states of atoms in halides* (499)
 Investigation of the solubility and diffusion of Fe atoms in Cu at high temperature using molten salt electrochemistry* (693)

attention

Anomalous melting of f electron metals (with attention to Pu)* (318)

attenuation

Thermal expansion of molten materials by the gamma attenuation technique* (152)

Aug

Structure of lithium--boron alloys and their performance as potential anodes for high power molten salt batteries. Progress report, 1 Aug 1974--30 Jun 1975* (275)

average

Electrochemistry of niobium and tantalum, (5). Anodic dissolution of tantalum and the average valence of tantalum ions in molten KCl* (450)

B

Influence of small additions on the preparation and the properties of UO_2 ceramics. Part B. The UO_2 - TiO_2 system* (267)

Ba

Gases and carbon in metals (thermodynamics, kinetics, and properties). I. Alkali metals, alkaline earth metals, light metals (Li, Na, K, Rb, Cs; Ca, Sr, Ba; Be, Mg, Al)* (192)
 Negative temperature coefficients of electrical resistivity: the divalent liquid metals Eu, Yb and Ba* (224)
 Study of the retention of Ba in UO_2 nuclear fuel particles by ZrO_2 as getter* (470)
 Solubility of rare earth and yttrium oxochlorides in molten salts - MCl_2 (M-Mg, Ca, Sr, Ba)* (495)

background

Influence of salt background on kinetics of BF_4^- - alkali hydrolysis* (504)

BaF

Misch metal solubility in melt of LiF - BaF_2 system* (462)
 Phase relations in the alkaline earth fluoride-yttria systems (in three parts). 2. The BaF_2 - Y_2O_3 phase diagram* (469)
 Practical application of conversion method in study of five-component mutual system consisting of nine salts NaF , Na_2MoO_4 , Na_2WO_4 , KF , K_2MoO_4 , K_2WO_4 , BaF_2 , BaMoO_4 , BaWO_4 * (516)

TITLE INDEX

BaF

NaMoO₄--BaMoO₄--NaF--BaF system* (638)

BaMoO

Practical application of conversion method in study of five-component mutual system consisting of nine salts NaF, Na₂MoO₄, Na₂WO₄, KF, K₂MoO₄, K₂WO₄, BaF₂, BaMoO₄, BaWO₄* (516)

band

Vibrational spectra of transition metal hexafluoride crystals. III. Exciton band structures of MoF₆, WF₆, and UF₆. Technical report* (52)

Band structure and chemical bond in ThX compounds* (259)

Band structure studies of actinide systems* (323)

bands

XPS valence bands of La, Ce and Gd and their aluminium alloys* (598)

bar

Density and enthalpy of liquid alkaline metals at pressures up to 1500 bar* (599)

barium

Electrode reactions in molten sodium nitrate--barium nitrate eutectic at 350 degrees C* (42)

Ternary systems from lithium, sodium, strontium and barium chlorides* (86)

Refractory materials resistance to molten barium alloy* (628)

Cesium, calcium and barium chlorides ternary system* (709)

base

Solubility of nitrogen in multicomponent nickel base alloys* (100)

Ceramic-base heat resisting constructional material* (348)

Calculated method of eutectics of multicomponent systems on base of algorithm and digital computers* (517)

Spectrophotometric determination of rare earths and yttrium in gallium base alloys* (623)

Basic

Basic problems of fast reactor fuel reprocessing* (135)

Properties of yttrium and scandium basic salts* (393)

Basic mechanisms providing oxidation resistance in structural metals at high temperatures. II. The defect structure of nonstoichiometric rutile (TiO₂) and nonstoichiometric rutile containing 1 to 10 mole % Nb₂O₆. Final report 21 Dec 1973-20 Dec 1974* (542)

Recent development of ceramic basic science in Japan* (557)

Lithium alloys--chlorine secondary battery. I. Basic studies on a lithium alloys--chlorine secondary battery using a molten salt as the electrolyte* (704)

Basic research in crystalline and noncrystalline ceramic systems. Annual report, May 1, 1975--April 1, 1976* (728)

basicity

Effect of basicity on ferrous oxide activity in complex oxide systems* (372)

batteries

Stable inorganic matrix materials for high temperature batteries. Final report No. 1, 2 June 1967--2 February 1968* (23)

TITLE INDEX

batteries

Composition of nonaqueous solutions of potential use in high energy density batteries. Final report, 15 Nov 1969--14 Nov 1972* (119)

Structure of lithium--boron alloys and their performance as potential anodes for high power molten salt batteries. Progress report, 1 Aug 1974--30 Jun 1975* (275)

Research on electrochemical parameters for nonaqueous batteries: task I. Quarterly progress report No. 3* (727)

battery

Lithium anode limited-cycle battery investigation. Quarterly technical progress report No. 5* (37)

Lithium anode limited-cycle battery investigation. Quarterly technical progress report No. 3* (38)

Lithium--silver chloride secondary battery investigation. Quarterly technical progress report No. 2* (111)

Lithium--silver chloride secondary battery investigation. Final report, December 1963--November 1964* (118)

Composition gradient during electrolysis in mixtures analogous to molten salt battery electrolytes* (234)

Proceedings of the symposium and workshop on advanced battery research and design, March 22--24, 1976* (269)

Electrochemical characterization of nonaqueous systems for secondary battery application. Quarterly report, February--April 1968* (380)

Lithium--nickel fluoride secondary battery investigation. Quarterly technical progress report No. 1, 27 Apr--27 Jul 1966* (381)

Lithium alloys--chlorine secondary battery. I. Basic studies on a lithium alloys--chlorine secondary battery using a molten salt as the electrolyte* (704)

Research on a high-energy nonaqueous battery system. Quarterly progress report No. 1, 1 Dec 1965--1 Mar 1966* (720)

BaWO

Practical application of conversion method in study of five-component mutual system consisting of nine salts NaF , Na_2MoO_4 , Na_2WO_4 , KF , K_2MoO_4 , K_2WO_4 , BaF_2 , BaMoO_4 , BaWO_4 * (516)

BaZrF

New hexafluorozirconates(IV): BaZrF_6 , PbZrF_6 , EuZrF_6 , SrZrF_6 * (419)

bearing

Total pressures of uranium- and plutonium-bearing species above the U--Pu--C system* (618)

BeCl

Meltability diagram of RbCl - BeCl_2 system* (506)

bed

Adaptation of the continuous cold trap system of fluidized-bed to the fluoride volatility process* (724)

Fluorination process studies of plutonium dioxide by fluid-bed* (725)

BeF

Study of KBeF_5 - DyF_3 cross-section of KF - BeF_2 - DyF_3 ternary system and some properties of this cross section glasses* (21)

TITLE INDEX

BeF

Measurement of mass transfer coefficients in a mechanically agitated, nondispersing contactor operating with a molten mixture of $\text{LiF--BeF}_{\text{sub } 2}\text{--ThF}_{\text{sub } 4}$ and molten bismuth* (79)
 Phase diagrams in the pseudobinary systems of $\text{Li}_{\text{sub } 2}\text{BeF}_{\text{sub } 4}$, alkali fluoroborates and alkali fluorides* (473)
 Heat transfer measurements in a forced convection loop with two molten-fluoride salts: $\text{LiF--BeF}_{\text{sub } 2}\text{--ThF}_{\text{sub } 2}\text{--UF}_{\text{sub } 4}$ and eutectic $\text{NaBF}_{\text{sub } 4}\text{--NaF}$ * (585)

behavior

Comparison of the thermodynamic properties and high temperature chemical behavior of lanthanide and actinide oxides* (2)
 Behavior of molybdenum in mixed-oxide fuel* (202)
 Electrochemical behavior of tellurium (4) in oxychloride melts* (221)
 Structure and anodic discharge behavior of lithium-boron alloys in the LiCl-KCl eutectic melt.* (274)
 Electrochemical behavior of U(III) and U(IV) in molten LiCl-KCl eutectic* (293)
 Study of the behavior of high activity waste produced during the regeneration of fast reactor fuel elements by the gaseous fluoride method* (311)
 Microprobe study of fission product behavior in high-burnup HTR fuels* (315)
 MATERO: Version 09. A handbook of materials properties for use in the analysis of light water reactor fuel rod behavior* (387)
 Solution behavior of hydrogen isotopes and other non-metallic elements in liquid lithium* (404)
 Behavior of the AB-type compounds at high pressures and high temperatures* (421)
 Some regularities of the corrosion behavior of binary magnesium alloys* (439)
 Vaporization behavior of $\text{UO}_{\text{sub } 2}$ * (445)
 Evaporation behavior and high-temperature thermal analysis of substoichiometric plutonium oxide in the composition range from $\text{O/Pu} = 1.51$ to 2.00 * (482)
 Some trends in constitutive equation model development for high-temperature behavior of fast-reactor structural alloys* (527)
 Behavior of ruthenium in fluoride-volatility process. II. Fluorination of $\text{RuO}_{\text{sub } 2}$ by $\text{F}_{\text{sub } 2}$ * (558)
 Phase behavior and thermodynamics of U--Mo--C system* (646)
 Phase behavior and thermodynamics of U--Mo--C system* (647)
 Anodic behavior of cadmium and cadmium-zinc alloys in eutectic melt of potassium- and lithium chlorides* (675)

behaviour

Evaporation behaviour of the ternary uranium plutonium carbides* (479)
 Evaporation behaviour of the ternary uranium plutonium carbides* (480)
 Evaporation behaviour of the ternary uranium plutonium carbides* (481)
 Specialists' meeting on the behaviour of water reactor fuel elements under accident conditions* (510)
 Anomalous thermal property behaviour of uranium at low temperatures* (560)

TITLE INDEX

behaviour

Molten salt chemistry : Part IV - Solubility behaviour of $\text{PuF}_{3\text{sub}}$ in fluoride salts of interest in molten salt reactor technology* (655)

berkelium

All-union conference on chemistry of transplutonium elements (americium, curium, berkelium, californium)* (34)

All-union conference on chemistry of transplutonium elements (americium, curium, berkelium, californium)* (307)

All-union conference on chemistry of transplutonium elements (americium, curium, berkelium, californium)* (308)

berthollides

Phases of variable composition with $\text{LaF}_{3\text{sub}}$ structure in $\text{MF}_{2\text{sub}}-(\text{Y,Ln})\text{F}_{3\text{sub}}$ systems. 1. $\text{SrF}_{2\text{sub}}-(\text{Y,Ln})\text{F}_{3\text{sub}}$ systems (thermal characteristics, formation of berthollides)* (593)

beryllium

Thermodynamic analysis of melting point diagrams of systems consisting of alkali metal chlorides, thallium chloride and beryllium chloride* (505)

Enthalpy of vitreous beryllium fluoride* (616)

Some thermal properties of beryllium fluoride from 8⁰ to 1,200⁰ K* (617)

BF

Influence of salt background on kinetics of $\text{BF}_{4\text{sup}}$ - alkali hydrolysis* (504)

Bi

Structure of molten Bi-Sb alloys by means of neutron diffraction* (359)

Structure of $\text{In}_{2\text{sub}}\text{Bi}$ melted compound* (514)

Energy transfer between $\text{Bi}_{3\text{sup}}^{3+}$ yields $\text{Eu}_{3\text{sup}}^{3+}$, $\text{Bi}_{3\text{sup}}^{3+}$ yields $\text{Sm}_{3\text{sup}}^{3+}$ and $\text{UO}_{2\text{sup}}^{2+}$ yields $\text{Eu}_{3\text{sup}}^{3+}$ in oxide glasses* (539)

Ga-Bi-Te system* (544)

GaTe-Bi₂Te₃ system* (545)

Investigation into properties of liquidus curves of binary salt system melts where chemical compounds formation. 2. Systems of monc-monovalent and bi-bivalent cations with common anion* (708)

bibliography

Heat transfer bibliography* (158)

EUTECTIC DATA: safety, hazards, corrosion, melting points, compositions, and bibliography* (276)

BiBr

Phase diagrams of $\text{BiBr}_{3\text{sub}}$ - $\text{BiI}_{3\text{sub}}$ system* (288)

BiCl

Phase diagrams of $\text{BiCl}_{3\text{sub}}$ - CuCl - $\text{TeCl}_{4\text{sub}}$ and CuCl - $\text{FeCl}_{3\text{sub}}$ - $\text{TeCl}_{4\text{sub}}$ systems* (337)

$\text{BiCl}_{3\text{sub}}$ - $\text{GeCl}_{4\text{sub}}$ - $\text{TeCl}_{4\text{sub}}$ system* (553)

biexciton

Structure of the biexciton molecule and optical transition selection rules in zincblende and wurtzite materials* (186)

bifluoride

Thermogravimetric investigation of interaction of some rare earth oxides with ammonium bifluoride* (427)

BiI

Phase diagrams of $\text{BiBr}_{3\text{sub}}$ - $\text{BiI}_{3\text{sub}}$ system* (288)

TITLE INDEX

binary

Viscosity of UCl_3 -- MgCl_2 binary system melts* (145)
 Calculation of thermodynamic activity and liquidus lines of components in binary systems of plutonium trichloride with alkali and alkaline earth metal chlorides* (147)
 Binary system of sodium, rubidium and cesium tungstates* (151)
 Binary molybdates of rare earths and silver (I)* (207)
 Thermodynamic study of the binary system copper-indium. I. Calorimetric study* (290)
 Thermodynamic study of the binary system copper--indium. II. Potentiometric study* (291)
 Studies on some binary solvent systems for the extraction of gadolinium* (370)
 Speed of ultrasound and the thermophysical properties of the liquid metals Sn, Pb, Cd and of their binary alloys Pb:Sn and Pb:Cd* (403)
 Some regularities of the corrosion behavior of binary magnesium alloys* (439)
 Theory for the derivation of a single hard-sphere potential describing the overall molecular interactions in binary liquid metal alloys as a function of temperature and composition* (525)
 MoCl_2 -- NaCl and MoCl_2 -- KCl binary systems* (547)
 Application of molecular dynamics computations to the conformal ionic solution theory: Thermodynamics of binary molten salt mixtures* (550)
 Viscosity of binary of nitrate systems melts* (592)
 Investigation into properties of liquidus curves of binary salt system melts where chemical compounds formation. 2. Systems of monc-monovalent and bi-bivalent cations with common anion* (708)

bis[n

Actinide metallocarborane complex; synthesis and x-ray structure determination of the bis[n^{sup}
 $5f$ -(3)-1,2-dicarbollyl]dichlorouranium(IV) dianion* (194)

bismuth

Density and temperature dependence of the surface tension of molten bismuth, lead, and bismuth-lead alloys* (1)
 Measurement of mass transfer coefficients in a mechanically agitated, nondispersing contactor operating with a molten mixture of LiF -- BeF_2 -- ThF_4 and molten bismuth* (79)
 Electrochemical studies of the sodium--bismuth system* (187)
 On interaction of tellurium tetrachloride with copper (1), mercury (2), bismuth and iron (3) chlorides* (336)
 Activities of lead and bismuth in their molten alloys* (420)
 Current yield during separation of lithium on a liquid bismuth cathode* (430)
 Tritium sorption in lithium--bismuth and lithium--aluminum alloys* (614)

bivalent

Investigation into properties of liquidus curves of binary salt system melts where chemical compounds formation. 2. Systems of monc-monovalent and bi-bivalent cations with common anion* (708)

body

Summaries of reports of the all-union meeting on solid body chemistry. 3--5 June 1975. III. Synthesis, phase equilibrium and use of solid oxide compounds* (114)

TITLE INDEX

body

Summaries of reports of the all-union meeting on solid body chemistry. 3--5 June 1975. III. Synthesis, phase equilibrium and use of solid oxide compounds* (641)

bond

Band structure and chemical bond in ThX compounds* (259)

Bonding

Bonding and the electronic structure of the actinide metals* (188)

Electron structure and bonding in ThO_2 and UO_2 * (223)

Partly filled shells constituting anti-bonding orbitals with higher ionization energy than their bonding counterparts* (280)

Structure and Bonding* (492)

Structure and Bonding* (538)

Book

5th international conference on plutonium and other actinides.

Book of abstracts* (109)

5th international conference on plutonium and other actinides.

Book of abstracts* (620)

boron

Structure and anodic discharge behavior of lithium-boron alloys in the LiCl-KCl eutectic melt.* (274)

Structure of lithium--boron alloys and their performance as potential anodes for high power molten salt batteries. Progress report, 1 Aug 1974--30 Jun 1975* (275)

Investigation of the parameters influencing the microstructure of hot-pressed boron carbide* (440)

Br

Optical spectra of the dibromide and diiodide ions in the matrix-isolated $\text{M}^{\text{sup}} + \text{Br}_2^{\text{sup}} -$ and $\text{M}^{\text{sup}} + \text{I}_2^{\text{sup}} -$ species* (17)

Chemistry of elements niobium and tantalum. 87. Studies on the thermal formation and decomposition of the complexes $[\text{NbCl}_{12}]^{\text{sup}} 2+$, $[\text{Nb}_6\text{I}_8]^{\text{sup}} 3+$, and $[\text{Ta}_6\text{X}_{12}]^{\text{sup}} (n+)$ ($\text{X}=\text{Cl}, \text{Br}, \text{I}$)* (561)

breeder

Investigation of fuel vapor pressure for fast breeder safety analysis* (60)

Reprocessing of fast breeder fuels in France* (67)

Report of task force on alternate structural materials for Liquid Metal Fast Breeder Reactors* (309)

Properties of structural materials for sodium-cooled fast breeder reactor* (580)

bromides

X-ray diffraction study into molten bromides and iodides of alkali metals* (20)

Pressure of saturated vapor of rare earth and yttrium bromides* (391)

Quaternary mutual system of chlorides, bromides, iodides of sodium and rubidium* (629)

Quaternary mutual system of chlorides, bromides, iodides of potassium and cesium* (630)

Sections of quaternary mutual system consisting of sodium, potassium and cesium chlorides and bromides* (631)

Thermographic determination of formation heats of lower zirconium chlorides and bromides* (635)

TITLE INDEX

bromides

Quaternary mutual system of chlorides, bromides, iodides of rubidium and cesium* (640)

bubbles

Equilibrium index for fission gas bubbles* (685)

burn

Determination of the burn up and differentiation of the fission yields of uranium-thorium-fuels* (214)

burnup

Microprobe study of fission product behavior in high-burnup HTR fuels* (315)

Fission rates, burnup, and neutron flux-fluence-spectra characterization for mixed oxide fuel experiments in the EBR-II* (416)

C

Standard potential of sodium amalgam at 25°C* (32)

Electrode reactions in molten sodium nitrate--barium nitrate eutectic at 350 degrees C* (42)

X-ray study of the plutonium--oxygen phase diagram at ambient temperatures up to 1100°C* (62)

Distribution coefficient for U and Pu in the system (UPu)₂C + (UPu)₃C* (134)

Solubility of sodium iodide in liquid sodium between 140 and 500°C* (154)

Thermodynamic study of the U-O-C system by measurement of the carbon monoxide pressures at equilibrium* (235)

Investigation of calcite solubility in K₂CO₃ and Na₂CO₃ solutions at temperatures of 200 to 350°C* (396)

Specific heat of PuC_(1-x) and Pu₂C₃ from 10 to 300 K* (414)

Specific heat of PuC_(1-x) and Pu₂C₃ from 10 to 300 K* (415)

Contribution to the establishment of the phase diagram for the U--C--C system by means of x-ray diffractometry at high temperatures and under controlled pressure* (503)

K₂HfF₆--K₂ZrF₆--1.25 percent of HF system at 40°C with other solubility curves from 25°C to 70°C* (541)

Thermodynamics of solid solutions CsCl-RbCl and CsBr-RbBr at 25°C* (570)

Phase behavior and thermodynamics of U--Mo--C system* (646)

Phase behavior and thermodynamics of U--Mo--C system* (647)

Phase equilibria in system U--Pu--W--C* (648)

Parameter for optimizing the synthesis of (U,Pu)₂C* (676)

Fuel-clad interaction. C-E thermo-structural fuel evaluation methods* (719)

Ca

Study of chemical interaction in Pb₃AlF₆--LaF₃--Ca₃AlF₆ system* (26)

Gases and carbon in metals (thermodynamics, kinetics, and properties). I. Alkali metals, alkaline earth metals, light metals (Li, Na, K, Rb, Cs; Ca, Sr, Ba; Be, Mg, Al)* (192)

LiNO₃--CsNO₃--Ca(NO₃)₂ and NaNO₃--CsNO₃--Ca(NO₃)₂ system* (255)

TITLE INDEX

- Ca**
Solubility of rare earth and yttrium oxochlorides in molten salts - MCl_2 (M-Mg, Ca, Sr, Ba)* (495)
- CaBr**
Phase diagrams of LiCl , CsCl , CaCl and LiBr , CsBr , CaBr ternary systems* (256)
- CaCl**
Phase diagram for the ternary system CaCl_2 -- KCl -- CaCrO_4 * (117)
Viscosity of molten CaCl_2 -- UCl_3 mixtures* (146)
Phase diagrams of LiCl , CsCl , CaCl and LiBr , CsBr , CaBr ternary systems* (256)
System Na_2WO_4 -- NaCl -- CaWO_4 -- CaCl_2 * (637)
- CaCrO**
Phase diagram of system LiCl -- KCl -- CaCrO_4 * (116)
Phase diagram for the ternary system CaCl_2 -- KCl -- CaCrO_4 * (117)
- cadmium**
Perturbation treatment of the Percus-Yevick equation and structures of liquid magnesium and cadmium* (210)
Ternary mutual system of chlorides and iodides of rubidium and cadmium* (257)
Thermophysical properties of the ternary liquid-metal system tin--lead--cadmium* (605)
Anodic behavior of cadmium and cadmium-zinc alloys in eutectic melt of potassium- and lithium chlorides* (675)
- Caesium**
Thermochemistry of polyhalides. Pt.1. Caesium and rubidium dichlorodates(I)* (178)
Static structure factor of liquid caesium* (248)
Structure factor of liquid caesium and sodium-caesium alloys* (249)
- CaF**
Investigation into americium extraction process from residues of irradiated nuclear fuel after its fluorination by SiF-NaF and SiF-CaF_2 melts of eutectic composition* (308)
Phase relations in the alkaline earth fluoride-yttria systems (in three parts). 1. The CaF_2 -- Y_2O_3 phase diagram* (468)
Thermal analysis of the system of NaF , CaF_2 , Na_2MoO_4 , CaMoO_4 * (519)
Interdiffusion in the CaF_2 -- YF_3 system* (669)
Optical spectroscopy of ^{237}Np in CaF_2 . Final report* (718)
- calcite**
Investigation of calcite solubility in K_2CO_3 and Na_2CO_3 solutions at temperatures of 200 to 350°C* (396)
- calcium**
Ternary system of potassium, cesium and calcium nitrates* (254)
Solubility of magnesium oxide in calcium oxide--calcium chloride mixtures* (352)
Interaction of thorium chloride with calcium oxide* (496)
Cesium, calcium and barium chlorides ternary system* (709)

TITLE INDEX

Calculated

Calculated method of eutectics of multicomponent systems on base of algorithm and digital computers* (517)

californium

All-union conference on chemistry of transplutonium elements (americium, curium, berkelium, californium)* (34)

All-union conference on chemistry of transplutonium elements (americium, curium, berkelium, californium)* (307)

All-union conference on chemistry of transplutonium elements (americium, curium, berkelium, californium)* (308)

calorimeter

High-temperature calorimeter for determination of reaction enthalpy* (364)

Calorimetric

Thermodynamic study of the binary system copper-indium. I.

Calorimetric study* (290)

calorimetry

Heat capacities and heat content of molten cerium by levitation calorimetry* (355)

Lithium nitride (Li_3N): standard enthalpy of formation by solution calorimetry* (467)

Progress in fluorine calorimetry* (500)

CaMoO

Thermal analysis of the system of NaF, CaF_2 , Na_2SiF_6 , CaMoO_4 * (519)

CaO

Electrical conduction in CaO-doped thoria electrolytes* (389)

capacities

Heat capacities and heat content of molten cerium by levitation calorimetry* (355)

Thermodynamics of the lanthanide halides. II. Heat capacities and Schottky anomalies of SmCl_3 , EuCl_3 and GdCl_3 from 50 to 350 K* (594)

carbide

Uranium carbide: thermal diffusivity, thermal conductivity and spectral emissivity at high temperatures* (121)

Theoretical prediction of the thermal conductivity of uranium carbide vapor* (124)

Enthalpy of uranium-plutonium carbide from 298K to the melting point* (179)

Investigation of the parameters influencing the microstructure of hot-pressed boron carbide* (440)

Preliminary assessments of carbide fuel pins during mild overpower transients in LMFBRs* (456)

Thermodynamic activity of carbon in molybdenum-containing uranium carbide* (644)

Thermodynamic activity of carbon in molybdenum-containing uranium carbide* (650)

Fabrication of uranium-plutonium mixed carbide* (691)

carbides

Physical and mechanical properties of some uranium-containing carbides* (209)

Evaporation behaviour of the ternary uranium plutonium carbides* (479)

TITLE INDEX

carbides

- Evaporation behaviour of the ternary uranium plutonium carbides* (480)
- Evaporation behaviour of the ternary uranium plutonium carbides* (481)
- Equation of state and transport properties of uranium and plutonium carbides in the liquid region* (577)
- Densities of liquid carbides and nitrides of uranium and plutonium* (579)
- X-ray spectral study of the carbides and nitrides of group IV and V metals* (582)
- Analysis of uranium carbides* (726)

carbon

- Segregation and vapour pressure studies on the uranium-plutonium-carbon system* (83)
- Segregation and vapour pressure studies in the uranium--plutonium--carbon system* (84)
- Gases and carbon in metals (thermodynamics, kinetics, and properties). I. Alkali metals, alkaline earth metals, light metals (Li, Na, K, Rb, Cs; Ca, Sr, Ba; Be, Mg, Al)* (192)
- Some constitutional studies on uranium-carbon- and plutonium-carbon-rhenium and technetium systems* (227)
- Some constitutional studies on uranium--carbon and plutonium--carbon--rhenium and technetium systems* (228)
- Thermodynamic study of the U-O-C system by measurement of the carbon monoxide pressures at equilibrium* (235)
- Structure of the thorium-(zirconium, niobium, ruthenium, rhodium)-carbon systems* (245)
- Vaporization of uranium-carbon-nitrogen system* (251)
- Thermodynamic activity of carbon in molybdenum-containing uranium carbide* (644)
- Phase equilibrium study on system uranium-plutonium-tungsten-carbon* (645)
- Thermodynamic activity of carbon in molybdenum-containing uranium carbide* (650)

carbonate

- Molten carbonate fuel cell research at ORNL* (71)
- Phase equilibria in Li--Na--K carbonate/aluminate systems* (77)

carbonates

- Thermodynamic properties of some rare earth carbonates and yttrium carbonates* (36)
- Solubility of oxygen in molten carbonates* (563)

carbonitride

- Process for preparing uranium carbonitride* (649)

carbonitrides

- Preparation and thermochemical stability of uranium-zirconium-carbonitrides.* (332)
- Preparation and thermochemical stability of uranium-zirconium-carbonitrides. A contribution to nuclear fuels for advanced high-temperature reactors* (333)
- Preparation and thermochemical stability of uranium-zirconium-carbonitrides* (334)

Carlo

- Monte Carlo analysis of diffusion and thermodynamics in UO_{2+x} * (441)

TITLE INDEX

carrier

Zeeman effect and satellite carrier structure in absorption spectra of pure and doped praseodymium and europium double nitrates* (165)

cathode

Current yield during separation of lithium on a liquid bismuth cathode* (430)

cation

Study of the compounds $M\text{UO}_2\text{F}_5$ ($M = \text{K, Rb, Cs, NH}_4$) by i.r. absorption and Raman diffusion spectrophotometry. III. Effect of cation on vibration frequency of the ion $\text{UO}_2\text{F}_5^{3-}$ * (127)

cations

Effect of second sphere cations on formation of nickel (2) chloride complexes in molten salts* (673)
Investigation into properties of liquidus curves of binary salt system melts where chemical compounds formation. 2. Systems of monc-monovalent and bi-bivalent cations with common anion* (708)

cavitation

Experimental investigation of cavitation inception and damage in a flowing sodium environment* (103)

CaWO

System $\text{Na}_2\text{WO}_4\text{-NaCl-CaWO}_4\text{-CaCl}_2$ * (637)

Cb

Basic mechanisms providing oxidation resistance in structural metals at high temperatures. II. The defect structure of nonstoichiometric rutile (TiO_2) and nonstoichiometric rutile containing 1 to 10 mole % Cb_2O_6 . Final report 21 Dec 1973-20 Dec 1974* (542)

CCL

Experience of the Central Control Laboratory (CCL) in accounting for and controlling nuclear material in Czechoslovakia* (349)

Cd

Thermodynamic properties of Cd-Zn-Hg liquid alloys* (340)
Speed of ultrasonic and the thermophysical properties of the liquid metals Sn, Pb, Cd and of their binary alloys Pb:Sn and Pb:Cd* (403)
Thermodynamic investigation of liquid alloys in the system Cd-In-Tl* (523)
Contribution to the problem of the thermodynamic properties and the structure of liquid Cd-In alloys* (524)
Thermodynamic properties of the ternary Sn-Pb-Cd liquid metal system* (606)

CdCl

Crycmetric studies of the system $\text{CdCl}_2\text{-UCl}_4$ * (199)

Ce

Investigation into thermodynamic properties of Ce-Mg melts* (39)
XPS valence bands of La, Ce and Gd and their aluminium alloys* (598)
Structure of liquid Ce-Ni alloys* (689)

CeF

Pressure effect on spectral properties of $\text{CeF}_3\text{:Pr}^{3+}$ * (677)

cell

Molten carbonate fuel cell research at ORNL* (71)

TITLE INDEX

cell

Knudsen cell--mass spectrometer studies of cesium--urania interactions* (120)
 Simple Raman cell and furnace usable at temperatures higher than 1000⁰ for corrosive melts* (204)
 Determination and correlation of mass transfer coefficients in a stirred cell* (236)
 Measurement of the cell voltage of ThO_2 -- Y_2O_3 --solid electrolyte cells in the range of mixed conduction* (535)

cells

Third international symposium on fuel cells* (77)
 Thermodynamic properties of Cr, Mn, Co and Ni dichlorides from emf measurements on cells with solid electrolytes* (200)
 Measurement of the cell voltage of ThO_2 -- Y_2O_3 --solid electrolyte cells in the range of mixed conduction* (535)
 Complex impedance of electrochemical cells based on yttria doped thoria* (568)

CeMn

New representatives of CeMn_4Al_8 structural type* (549)

CeN

Electronic structure of CeN studied by x-ray-photoemission spectroscopy* (28)
 Decomposition of the UN-CeN solid solution* (581)

centers

Effect of luminescence centers on optical absorption, excitation and fluorescence spectra of CsCl:Pb * (491)

Central

Experience of the Central Control Laboratory (CCL) in accounting for and controlling nuclear material in Czechoslovakia* (349)

ceramic

Uranium dioxide as a ceramic nuclear fuel* (226)
 Ceramic-base heat resisting constructional material* (348)
 High temperature chemistry of ceramic nuclear fuels with emphasis on nonstoichiometry* (448)
 Equilibria and kinetics in modern ceramic processing* (557)
 Basic research in crystalline and noncrystalline ceramic systems. Annual report, May 1, 1975--April 1, 1976* (728)

ceramics

Thermodynamic properties of solid and liquid metals and ceramics* (243)
 Specific heat of liquid metals and ceramics* (244)
 Influence of small additions to the preparation and the properties of UO_2 ceramics. Part A. $\text{UO}_2/\text{Gd}_2\text{O}_3$ * (266)
 Influence of small additions on the preparation and the properties of UO_2 ceramics. Part B. The UO_2 -- TiO_2 system* (267)
 Mass transport phenomena in ceramics* (401)
 Zirconia and yttria--thoria ceramics find new uses as solid electrolytes* (595)
 Thermal properties of nuclear fuel ceramics* (613)
 Investigations on chemical properties of thorium oxide solid electrolyte ceramics* (652)

TITLE INDEX

cerium

Heat capacities and heat content of molten cerium by levitation calorimetry* (355)
 Effect of fluoride ions on the kinetics of the oxidation of uranium(IV) by cerium(IV)* (424)
 Determination of thorium in rare earths chloride: coprecipitation with cerium (IV) iodate and spectrophotometric determination with thoron* (447)
 Deoxidation thermodynamics of cerium and lanthanum in liquid iron* (654)

cesium

Hydrogen solubility in liquid cesium* (22)
 Phase equilibria in the system of sodium chlorides, cesium chlorides and strontium chlorides* (87)
 Phase equilibria in the system of sodium chlorides, cesium chlorides and strontium chlorides* (88)
 Knudsen cell--mass spectrometer studies of cesium--urania interactions* (120)
 Binary system of sodium, rubidium and cesium tungstates* (151)
 Magnetic susceptibility and electronic structure in metallic and non-metallic liquid cesium-gold and cesium-antimony alloys* (190)
 Ternary system of potassium, cesium and calcium nitrates* (254)
 Phase diagram of the ternary system of lithium-, rubidium-, and cesium nitrates* (377)
 Interaction of rubidium and cesium fluorides with lutetium trifluorides* (444)
 Some structural characteristics of rubidium and cesium in the vicinity of their melting points* (460)
 Cesium nitrate. Standard enthalpies of solution and formation and the standard entropy at 298.15 K* (466)
 Investigation into reactions of hexabromotellurates of potassium, cesium and thallium(I) in melt* (556)
 Viscosities of cesium vapor to 1,620 K and of liquid gallium to 1,800 K* (622)
 Quaternary mutual system of chlorides, bromides, iodides of potassium and cesium* (630)
 Sections of quaternary mutual system consisting of sodium, potassium and cesium chlorides and bromides* (631)
 Quaternary mutual system of chlorides, bromides, iodides of rubidium and cesium* (640)
 Viscosities of cesium vapor to 1620 K and of liquid gallium to 1800 K* (681)
 X-ray diffraction and infrared: Raman spectroscopic study of the hydrogen dinitrate ion, $(\text{O} \text{ } \text{NO} \text{--} \text{H} \text{--} \text{ONO})^{\text{sup } -}$, in cesium hydrogen dinitrate* (695)
 Cesium, calcium and barium chlorides ternary system* (709)

Characterisation

Characterisation of the chemical activities of non-metals dissolved in liquid alkali metals* (63)
 Characterisation of the chemical activities of non-metals dissolved in liquid alkali metals* (64)

characterization

Preparation and characterization of higher uranates(VI) of thallium(I)* (278)

TITLE INDEX

characterization

- Rare earth metal--metal halide systems. 19. Structural characterization of the reduced holmium chloride $\text{Ho}_{0.5}\text{Cl}_{1.5}$ * (378)
- Electrochemical characterization of nonaqueous systems for secondary battery application. Quarterly report, February--April 1968* (380)
- Synthesis and characterization of a new uranium(V) compound: $\text{H}_{0.3}\text{O}_{0.3}\text{U}_{0.6}\text{F}_{0.6}$ * (409)
- Fission rates, burnup, and neutron flux-fluence-spectra characterization for mixed oxide fuel experiments in the EBR-II* (416)
- Metallurgical characterization of prototypic tubesheet forging* (471)

chemical

- Comparison of the thermodynamic properties and high temperature chemical behavior of lanthanide and actinide oxides* (2)
- Zirconium: Physico-chemical properties of its compounds and alloys* (7)
- Study of chemical interaction in $\text{Pb}_{0.3}\text{AlF}_{0.6}\text{-LaF}_{0.6}\text{-Ca}_{0.3}\text{AlF}_{0.6}$ system* (26)
- Chemical Engineering Division physical inorganic chemistry. Annual report, July 1975--June 1976* (58)
- Characterisation of the chemical activities of non-metals dissolved in liquid alkali metals* (63)
- Characterisation of the chemical activities of non-metals dissolved in liquid alkali metals* (64)
- Some chemical reactions of uranium hexachloride and the chlorination of uranium tetrachloride* (81)
- Chemical separation and spectrographic estimation of rare earths and yttrium in Eu_2O_3 and $(\text{U,Pu})_2\text{O}_3$ * (125)
- Chemical effects of composition changes in irradiated oxide fuel materials. II. Fission product segregation and chemical equilibria* (169)
- Band structure and chemical bond in ThX compounds* (259)
- Chemical Engineering Division fuels and materials chemistry. Annual report, July 1975--June 1976* (281)
- Chemical reprocessing of spent nuclear fuels* (322)
- Redox and chemical properties of U, Np and Pu in molten alkali chlorides* (360)
- Pulse NMR in solids: chemical shift, lead fluoride, and thorium hydride* (363)
- Chemical Engineering Division, reactor safety and physical property studies annual report, July 1974--June 1975* (367)
- Quantitative study of the chemical equilibria in solution in molten LiCl-KCl (70-30 mol.%) by absorption spectrophotometry. Application to neptunium* (385)
- Zirconium: Physico-chemical properties of its compounds and alloys* (464)
- Physical--chemical studies of transuranium elements. Progress report, April 1, 1976--March 31, 1977* (501)
- Investigations on chemical properties of thorium oxide solid electrolyte ceramics* (652)
- Thorium: physico-chemical properties of its compounds and alloys* (679)

TITLE INDEX

chemical

- Zirconium: Physico-chemical properties of its compounds and alloys* (680)
- Investigation into properties of liquidus curves of binary salt system melts where chemical compounds formation. 2. Systems of monc-monivalent and bi-bivalent cations with common anion* (708)
- Standard methods for chemical, mass spectrometric, and spectrochemical analysis of nuclear-grade uranium dioxide powders and pellets* (711)
- Standard methods for chemical, mass spectrometric, and spectrochemical analysis of nuclear-grade plutonium dioxide powders and pellets* (712)
- Standard methods for chemical, mass spectrometric, and spectrochemical analysis of nuclear-grade mixed oxides (U,Pu)O₂* (713)
- Standard methods for chemical, mass spectrometric, spectrochemical, nuclear, and radiochemical analysis of nuclear-grade plutonium metal* (714)

chemistry

- All-union conference on chemistry of transplutonium elements (americium, curium, berkelium, californium)* (34)
- Chemical Engineering Division physical inorganic chemistry. Annual report, July 1975--June 1976* (58)
- Summaries of reports of the all-union meeting on solid body chemistry. 3--5 June 1975. III. Synthesis, phase equilibrium and use of solid oxide compounds* (114)
- Chemistry* (176)
- Aspects of the solution chemistry of liquid alkali metals as elucidated from electrical resistivity studies* (247)
- Chemical Engineering Division fuels and materials chemistry. Annual report, July 1975--June 1976* (281)
- All-union conference on chemistry of transplutonium elements (americium, curium, berkelium, californium)* (307)
- All-union conference on chemistry of transplutonium elements (americium, curium, berkelium, californium)* (308)
- Analytical Chemistry Division annual progress report for period ending November 30, 1975* (423)
- Structural or solid state chemistry of rare earth sesquioxides from Goldschmidt to the present time* (438)
- High temperature chemistry of ceramic nuclear fuels with emphasis on nonstoichiometry* (448)
- Recent developments in the chemistry of solutions of salts in liquid alkali metals* (528)
- Chemistry of elements niobium and tantalum. 87. Studies on the thermal formation and decomposition of the complexes [Nb₆Cl₁₂]²⁺, [Nb₆I₈]³⁺, and [Ta₆X₁₂]⁽ⁿ⁺⁾ (X=Cl, Br, I)* (561)
- Summaries of reports of the all-union meeting on solid body chemistry. 3--5 June 1975. III. Synthesis, phase equilibrium and use of solid oxide compounds* (641)
- Molten salt chemistry: Part IV - Solubility behaviour of PuF₃ in fluoride salts of interest in molten salt reactor technology* (655)
- Nuclear spectroscopy. Laboratory of chemistry and radiochemistry* (723)

TITLE INDEX

Chemistry

Chemistry Division annual progress report for period ending November 1, 1975* (729)

Chemistry Division annual progress report for period ending November 1, 1975* (730)

chloride

Standard formation enthalpy of chloride alloys of $ZrCl_4$ -- KCl ($NaCl$) systems* (8)

Interaction of uranyl chloride with zirconium tetrachloride in the melt of sodium chloride and potassium chloride* (27)

Absorption spectra of curium (3) chloride complexes* (34)

Investigation into neodymium chloride state in equilibrium with metal in eutectic melt of potassium- and lithium chlorides* (40)

Saturation vapor pressure in iron chloride-alkaline earth metal chlorides systems* (97)

Lithium--silver chloride secondary battery investigation.

Quarterly technical progress report No. 2* (111)

Lithium--silver chloride secondary battery investigation. Final report, December 1963--November 1964* (118)

Density, surface tension, and viscosity of uranium

trichloride-sodium chloride melting* (141)

Density, surface tension and viscosity of melts of uranium

trichloride-lithium chloride* (142)

Interaction of lower indium chlorides with chloride and

chloroindates of potassium* (174)

Solubility of magnesium oxide in alkali metal chloride melt* (265)

Solubility of magnesium oxide in calcium oxide--calcium chloride mixtures* (352)

Rare earth metal--metal halide systems. 19. Structural characterization of the reduced holmium chloride $Ho_{0.5}Cl_{1.5}$ * (378)

Determination of thorium in rare earths chloride: coprecipitation with cerium (IV) iodate and spectrophotometric determination with thoron* (447)

Raman spectroscopic studies of yttrium (III) chloride--alkali metal chloride melts and of Cs_2NaYCl_6 and YCl_3 solid compounds* (489)

Lanthanum chloride interaction with magnesium oxide* (493)

Interaction of thorium chloride with calcium oxide* (496)

Thermodynamic analysis of melting point diagrams of systems consisting of alkali metal chlorides, thallium chloride and beryllium chloride* (505)

Electrochemical reduction of molybdenum oxychloride in molten lithium chloride - potassium chloride eutectic* (515)

Thermodynamics of electrolytes. 8. High-temperature properties, including enthalpy and heat capacity, with application to sodium chloride* (586)

A Moessbauer spectroscopic study of intercalation compounds of graphite with ferric chloride and aluminum chloride* (627)

Effect of second sphere cations on formation of nickel (2)

chloride complexes in molten salts* (673)

Spectra of combination scattering and structure of 3d-metals chloride complexes in molten salts* (674)

TITLE INDEX

chlorides

- Thermodynamic properties of diluted solutions of MnCl_2 in alkali metal chlorides* (6)
- Investigation into neodymium chloride state in equilibrium with metal in eutectic melt of potassium- and lithium chlorides* (40)
- Ternary systems from lithium, sodium, strontium and barium chlorides* (86)
- Phase equilibria in the system of sodium chlorides, cesium chlorides and strontium chlorides* (87)
- Phase equilibria in the system of sodium chlorides, cesium chlorides and strontium chlorides* (88)
- Quaternary mutual system from fluorides and chlorides of lithium, sodium and strontium* (89)
- Saturation vapor pressure in iron chloride-alkaline earth metal chlorides systems* (97)
- Interaction of uranium trichloride with alkaline earth metal chlorides* (139)
- Thermographic investigation into ternary systems containing sodium and potassium chlorides, uranium trichloride and thorium tetrafluoride* (140)
- Interaction of uranium- and thorium tetrachlorides with sodium- and potassium chlorides* (144)
- Calculation of thermodynamic activity and liquidus lines of components in binary systems of plutonium trichloride with alkali and alkaline earth metal chlorides* (147)
- Phase diagrams of ternary systems containing sodium and potassium chlorides, thorium tetrachloride and plutonium trichloride* (148)
- Interaction of lower indium chlorides with chloride and chloroindates of potassium* (174)
- Study of lanthanum, yttrium and erbium oxychlorides in melts of alkali metal chlorides* (196)
- Ternary mutual system of chlorides and iodides of rubidium and cadmium* (257)
- Titanium dioxide solubility in melts of lithium- and alkaline earth metal chlorides* (264)
- Reaction between sodium, rubidium and strontium chlorides* (286)
- On interaction of tellurium tetrachloride with copper (1), mercury (2), bismuth and iron (3) chlorides* (336)
- Redox and chemical properties of U, Np and Pu in molten alkali chlorides* (360)
- Some thermodynamic properties of dilute solutions of actinide chlorides in $(\text{Li-K})\text{Cl}$ and in $(\text{Na-K})\text{Cl}$ eutectics* (406)
- Diffusion of uranium species in several molten chlorides* (408)
- Electrode potentials of tantalum chlorides in a molten NaCl-KCl eutectic* (449)
- Electrochemistry of niobium and tantalum. VI. Electrode potentials of tantalum chlorides in molten KCl * (451)
- Interaction of neodymium- and yttrium chlorides with magnesium oxide* (494)
- Thermodynamic analysis of melting point diagrams of systems consisting of alkali metal chlorides, thallium chloride and beryllium chloride* (505)
- Thermographic study of iron, molybdenum, tantalum and tungsten high chlorides interaction* (555)

TITLE INDEX

chlorides

Quarter mutual system of chlorides, bromides, iodides of sodium and rubidium* (629)
 Quaternary mutual system of chlorides, bromides, iodides of potassium and cesium* (630)
 Sections of quaternary mutual system consisting of sodium, potassium and cesium chlorides and bromides* (631)
 Thermographic determination of formation heats of lower zirconium chlorides and bromides* (635)
 Quaternary mutual system of chlorides, bromides, iodides of rubidium and cesium* (640)
 Anodic behavior of cadmium and cadmium-zinc alloys in eutectic melt of potassium- and lithium chlorides* (675)
 Cesium, calcium and barium chlorides ternary system* (709)

chlorination

Some chemical reactions of uranium hexachloride and the chlorination of uranium tetrachloride* (81)
 Influence of intermediate products on chlorination kinetics of vanadium oxides in salt melt* (356)

chlorine

Pressure and composition of vapors over alkali metal tetrachloroferrates in chlorine atmosphere* (433)
 Lithium alloys--chlorine secondary battery. I. Basic studies on a lithium alloys--chlorine secondary battery using a molten salt as the electrolyte* (704)

chloroindates

Interaction of lower indium chlorides with chloride and chloroindates of potassium* (174)

chromium

Viscosity characteristics of the iron-chromium-nickel melts* (24)
 Reactions of liquid sodium with transition-metal oxides. VIII. The oxides of chromium-(III), -(IV), and -(VI) and disodium chromium(VI) tetraoxide* (35)
 Kinetic peculiarities of nitrogen dissolving in iron-chromium and iron-chromium-nickel melts* (172)
 Investigation on the resistance of high-refractory materials to chromium melt* (531)
 Gd, Tb, Dy, Ho, Er solubility in chromium* (610)

Chronopotentiometry

Chronopotentiometry of U(III) and U(IV) in molten LiCl-KCl eutectic* (292)

Cl

Electronic structure of highly symmetric complexes of lanthanoids and actinoids. Pt. 3. 'Electronic' Raman spectrum of CsU(IV)Cl_6 * (13)
 Electronic structure of highly symmetric complexes of lanthanoids and actinoids. Pt. 4. The electronic Raman spectrum of CsNaEu(III)Cl_6 * (14)
 Optical spectra of the difluoride, dichloride, and trichloride ions in the matrix-isolated $\text{M}^{\text{sup}} + \text{F}_2^{\text{sup}} -$, $\text{M}^{\text{sup}} + \text{Cl}_2^{\text{sup}} -$, and $\text{M}^{\text{sup}} + \text{Cl}_3^{\text{sup}} -$ species* (16)
 Rare earth metal--metal halide systems. 19. Structural characterization of the reduced holmium chloride $\text{Ho}_5\text{Cl}_{11}$ * (378)
 Some thermodynamic properties of dilute solutions of actinide chlorides in (Li-K)Cl and in (Na-K)Cl eutectics* (406)

TITLE INDEX

Cl

Thermodynamic properties of dilute solutions of ThCl_4 in $(\text{Li-K})\text{Cl}$ and (Na-K) eutectics* (407)
 Preparational, structural and vibrational study of $\text{M}_2\text{U}_5\text{Cl}_4$, $2\text{H}_2\text{O}$ ($\text{M} = \text{Rb}, \text{Cs}$). Detection of a tetranuclear anion $[(\text{UO}_2)_4\text{O}_8\text{Cl}_8(\text{H}_2\text{O})_2]^{4-}$ * (498)
 Chemistry of elements niobium and tantalum. 87. Studies on the thermal formation and decomposition of the complexes $[\text{NbCl}_{12}]^{2+}$, $[\text{NbI}_8]^{3+}$, and $[\text{TaX}_{12}]^{n+}$ ($\text{X} = \text{Cl}, \text{Br}, \text{I}$)* (561)
 Electronic structure of highly symmetric lanthanoid and actinoid compounds. VI. The electronic Raman spectrum of $\text{Cs}_2\text{NaCe(III)Cl}_6$ * (678)

clad

Fuel-clad interaction. C-E thermo-structural fuel evaluation methods* (719)

cladding

Computational simulation of cladding motion after melting including resolidification processes* (19)

closed

Study of mass transfer by liquid sodium in closed loop systems. Sodium technology development programme* (324)
 New description of liquid metal heat transfer in closed conduits* (502)

cluster

Thermodynamic properties of $(\text{U}, \text{Pu})\text{O}_{2+x}$ described by a cluster model* (133)

coatings

Synthesis of UN coatings on uranium* (400)

cobalt

Fault structures in rare earth--cobalt intermetallics* (11)
 Investigation into interaction of indium tribromide with dibromides of manganese, cobalt and nickel in melt* (153)
 Spreading of iron-nickel alloys on ZrC and nickel-cobalt alloys on ZrN* (252)

CoCl

Study of saturated vapor pressure above salt melts in CoCl_2 - MCl systems* (98)

CODATA

CODATA recommended key values for thermodynamics, 1976. Report of the CODATA Task Group on key values for thermodynamics, 1976* (717)

code

Analysis of LMFBR explosion model experiments by means of the SURECUM-II code* (601)

coefficient

Distribution coefficient for U and Pu in the system $(\text{UPu})\text{C} + (\text{UPu})_2\text{C}_3$ * (134)

coefficients

Measurement of mass transfer coefficients in a mechanically agitated, nondispersing contactor operating with a molten mixture of LiF--BeF_2 -- ThF_4 and molten bismuth* (79)
 Structure factor of liquid aluminum and calculation of transport coefficients* (159)

TITLE INDEX

coefficients

- Negative temperature coefficients of electrical resistivity: the divalent liquid metals Eu, Yb and Ba* (224)
- Determination and correlation of mass transfer coefficients in a stirred cell* (236)
- Diffusion coefficients in molten metals* (328)

cold

- Adaptation of the continuous cold trap system of fluidized-bed to the fluoride volatility process* (724)

Collection

- Collection of gamma spectra data of irradiated light water moderated reactor spent fuel and study of the applicability of the methods for fuel identification. Final report for the period 1 February 1972--October 1975* (453)

Committee

- Second specialist meeting on sodium/fuel interaction in fast reactors, Ispra, 21-23 November 1973. Sponsored by the Committee on the Safety of Nuclear Installations, OECD Nuclear Energy Agency, Paris, France* (272)

common

- Investigation into properties of liquidus curves of binary salt system melts where chemical compounds formation. 2. Systems of mono-monovalent and bi-bivalent cations with common anion* (708)

communications

- Abstracts of papers and communications submitted to the 5. all-union heat and mass transfer conference* (437)

Comparative

- Comparative study of systems of heavier rare-earth metals with hydrogen [Gd, Tb, Dy, Lu]* (418)
- Thermodynamic properties of NbO , $\text{Fe}_{0.95}\text{O}$, CoO , NiO , $\text{Cu}_{2.0}\text{O}$ and CuO oxides as comparative electrodes in electromotive forces method* (658)

compatibility

- Some observations on the compatibility of structural materials with molten lead* (25)
- Compatibility of molten rare earth metals with refractory materials* (115)

complex

- Atomic structure of melts of complex system with eutectic transformations* (155)
- Actinide metallocarbaborane complex; synthesis and x-ray structure determination of the bis[n^{sup} 5⁺- (3)-1,2-dicarbollyl]dichlorouranium(IV) dianion* (194)
- Comparison of the tendencies of complex formation of various metal ions in fused LiF-NaF-KF (eut.) and in fused LiCl-KCl (eut.)* (314)
- Effect of basicity on ferrous oxide activity in complex oxide systems* (372)
- Complex impedance of electrochemical cells based on yttria doped thoria* (568)

complexes

- Electronic structure of highly symmetric complexes of lanthanoids and actinoids. Pt. 3. 'Electronic' Raman spectrum of $\text{Cs}_{2.0}\text{U(IV)Cl}_{6.0}$ * (13)

TITLE INDEX

complexes

- Electronic structure of highly symmetric complexes of lanthanoids and actinoids. Pt. 4. The electronic Raman spectrum of $\text{Cs}^{2+}\text{NaEu}(\text{III})\text{Cl}_6$ * (14)
- Absorption spectra of curium (3) chloride complexes* (34)
- Chemistry of elements niobium and tantalum. 87. Studies on the thermal formation and decomposition of the complexes $[\text{NbCl}_{12}]^{2+}$, $[\text{Nb}_6\text{I}_8]^{3+}$, and $[\text{Ta}_6\text{X}_{12}]^{(n+)}$ ($\text{X}=\text{Cl}, \text{Br}, \text{I}$)* (561)
- Degree of thermal dissociation of MCaCl_3 ($\text{M}=\text{K}, \text{Rb}, \text{Cs}$) complexes during melting* (624)
- Isorhodate complexes of 3d-metal ions in molten salts* (672)
- Effect of second sphere cations on formation of nickel (2) chloride complexes in molten salts* (673)
- Spectra of combination scattering and structure of 3d-metals chloride complexes in molten salts* (674)

component

- Practical application of conversion method in study of five-component mutual system consisting of nine salts NaF , Na_2MoO_4 , Na_2WO_4 , KF , K_2MoO_4 , K_2WO_4 , BaF_2 , BaMoO_4 , BaWO_4 * (516)

components

- Calculation of thermodynamic activity and liquidus lines of components in binary systems of plutonium trichloride with alkali and alkaline earth metal chlorides* (147)
- Discrete variation of the structure and texture of liquid metals formed by components with discrete packing* (177)
- Calculation of thermodynamic activity of components in strontium liquid alloys* (446)

composition

- Tensimetric study of pressure and composition of saturated vapor in $\text{TlCl}-\text{InCl}_3$ system* (93)
- Composition of nonaqueous solutions of potential use in high energy density batteries. Final report, 15 Nov 1969--14 Nov 1972* (119)
- Chemical effects of composition changes in irradiated oxide fuel materials. II. Fission product segregation and chemical equilibria* (169)
- Vibration spectra of rare earth oxytungstates- Ln_2WO_6 composition* (185)
- Mass spectra and vapor composition of niobium pentafluoride* (216)
- Composition gradient during electrolysis in mixtures analogous to molten salt battery electrolytes* (234)
- Molecular vapor composition and mass spectra in $\text{KF}-\text{ZrF}_4$, $\text{CsF}-\text{ZrF}_4$ systems* (294)
- Investigation into americium extraction process from residues of irradiated nuclear fuel after its fluorination by SiF_4-NaF and $\text{SiF}_4-\text{CaF}_2$ melts of eutectic composition* (308)
- Solubility of $\text{Sr}(\text{NO}_3)_2$ in the $\text{LiNO}_3-\text{KNO}_3$ melt of variable composition* (351)
- Pressure and composition of vapors over alkali metal tetrachloroferrates in chlorine atmosphere* (433)
- Evaporation behavior and high-temperature thermal analysis of substoichiometric plutonium oxide in the composition range from $\text{O/Pu} = 1.51$ to 2.00 * (482)

TITLE INDEX

composition

Theory for the derivation of a single hard-sphere potential describing the overall molecular interactions in binary liquid metal alloys as a function of temperature and composition* (525)
 Phases of variable composition with $\text{LaF}_{\text{sub } 3}$ structure in $\text{MF}_{\text{sub } 2}$ -(Y, Ln) $\text{F}_{\text{sub } 3}$ systems. 1. $\text{SrF}_{\text{sub } 2}$ -(Y, Ln) $\text{F}_{\text{sub } 3}$ systems (thermal characteristics, formation of berthollides)* (593)

Effect of composition and structure on crevice, intergranular, and stress corrosion of some wrought Ni--Cr--Mo alloys* (604)

compositions

EUTECTIC DATA: safety, hazards, corrosion, melting points, compositions, and bibliography* (276)

compound

Interval of congruent and incongruent sublimation of $3\text{KF}\cdot\text{ZrF}_{\text{sub } 4}$ compound in the KF - $\text{ZrF}_{\text{sub } 4}$ system* (296)

Synthesis and characterization of a new uranium(V) compound: $\text{H}_{\text{sub } 3}\text{fO}_{\text{sup }} + \text{UF}_{\text{sub } 6}\text{O}_{\text{sup }} - \text{H}_2\text{O}$ * (409)

Structure of $\text{In}_{\text{sub } 2}\text{Bi}$ melted compound* (514)

compounds

Zirconium: Physico-chemical properties of its compounds and alloys* (7)

Mossbauer studies of americium compounds* (61)

Summaries of reports of the all-union meeting on solid body chemistry. 3--5 June 1975. III. Synthesis, phase equilibrium and use of solid oxide compounds* (114)

Study of the compounds $\text{M}_{\text{sub } 3}\text{UO}_{\text{sub } 2}\text{F}_{\text{sub } 5}$ ($\text{M} = \text{K}, \text{Rb}, \text{Cs}, \text{NH}_{\text{sub } 4}$) by i.r. absorption and Raman diffusion

spectrophotometry. III. Effect of cation on vibration frequency of the ion $\text{UO}_{\text{sub } 2}\text{F}_{\text{sub } 5}\text{O}_{\text{sup } 3-}$ * (127)

Some properties of hydrogen fluoride compounds with group 5 metal pentafluorides* (170)

Hydrogen solubility in rare earth intermetallic compounds* (217)

Thermochemistry of gaseous compounds of metals. Annual summary report 5 Dec 74--4 Dec 75* (240)

Thermochemistry of gaseous compounds of metals. Annual summary report, 5 Dec 1975--4 Dec 1976* (241)

Band structure and chemical bond in ThX compounds* (259)

Statistical mechanics of intermediate-valence rare earth compounds* (279)

$\text{R}_{\text{sub } 3}\text{T}_{\text{sub } 2}$ compounds ($\text{R} = \text{rare earth or Y}; \text{T} = \text{Rh}, \text{Pd}, \text{Pt}$) with the rhombohedral $\text{Er}_{\text{sub } 3}\text{Ni}_{\text{sub } 2}$ structure type* (365)

Hydrogen absorption of rare-earth (3d) transition intermetallic compounds* (392)

Thermal decomposition of intermetallic compounds of platinum with alkaline earth and lanthanide metals at high temperatures* (399)

Behavior of the AB-type compounds at high pressures and high temperatures* (421)

Tetrad effect in the extraction of the lanthanides with neutral oxygen-containing compounds* (426)

Zirconium: Physico-chemical properties of its compounds and alloys* (464)

Raman spectroscopic studies of yttrium (III) chloride--alkali metal chloride melts and of $\text{Cs}_{\text{sub } 2}\text{NaYCl}_{\text{sub } 6}$ and $\text{YCl}_{\text{sub } 3}$ solid compounds* (489)

TITLE INDEX

compounds

- Differential-spectrophotometric determination of niobium in its compounds* (490)
 Thermal lattice dilatation of some transition metal compounds* (508)
 2. all-union conference on crystallochemistry of intermetallic compounds* (549)
 Electronic and magnetic properties of rare earth-Sn₃ compounds from ¹¹⁹Sn Moesbauer spectroscopy* (559)
 Covalent compounds of uranium of the type $P_{x/5}U(O_2F_5)_{6-x}$ * (571)
 A Moesbauer spectroscopic study of intercalation compounds of graphite with ferric chloride and aluminum chloride* (627)
 Electrochemical study of high-temperature stability of compounds between the rare earths and copper oxide* (634)
 Summaries of reports of the all-union meeting on solid body chemistry. 3--5 June 1975. III. Synthesis, phase equilibrium and use of solid oxide compounds* (641)
 Compounds with the structure of ThMn₁₂ type in the R--Fe--Al systems* (671)
 Electronic structure of highly symmetric lanthanoid and actinoid compounds. VI. The electronic Raman spectrum of Cs₂NaCe(III)Cl₆* (678)
 Thorium: physico-chemical properties of its compounds and alloys* (679)
 Zirconium: Physico-chemical properties of its compounds and alloys* (680)
 Determination of trace impurities of titanium and zirconium in molybdenum and tungsten and their oxygen compounds by X-ray fluorescence spectrometry* (698)
 Investigation into properties of liquidus curves of binary salt system melts where chemical compounds formation. 2. Systems of monc-monovalent and bi-bivalent cations with common anion* (708)

Comptes

Comptes rendus du 95* (621)

compton

Studies of the electronic structure of simple metals and their alloys with the aid of compton scattering and positron annihilation using lithium, magnesium, lithium-magnesium and aluminium-zinc as examples* (48)

Computational

Computational simulation of cladding motion after melting including resolidification processes* (19)

computations

Application of molecular dynamics computations to the conformal ionic solution theory: Thermodynamics of binary molten salt mixtures* (550)

computers

Calculated method of eutectics of multicomponent systems on base of algorithm and digital computers* (517)

concentration

Interaction parameters in the Zn--Pb--Sn system at low zinc concentration* (436)

concepts

Theoretical concepts useful in the calculation or storage of phase diagrams of icnic systems* (552)

TITLE INDEX

conduction

Electrical conduction in CaO-doped thoria electrolytes* (389)
 Measurement of the cell voltage of ThO_2 - Y_2O_3 -solid electrolyte cells in the range of mixed conduction* (535)

conductivity

Density, viscosity, electric conductivity and meltability of NaAlF_6 - La_2O_3 and NaAlF_6 - Sm_2O_3 systems* (43)
 Study on conductivity, density, and viscosity of molten salt systems* (112)
 Uranium carbide: thermal diffusivity, thermal conductivity and spectral emissivity at high temperatures* (121)
 Theoretical prediction of the thermal conductivity of uranium carbide vapor* (124)
 Thermal diffusivity, thermal conductivity and spectral emissivity of uranium dicarbide at high temperatures* (132)
 Electronic conductivity and polarization effects of solid ThO_2 - Y_2O_3 electrolytes and of oxygen containing Va-metal electrodes* (193)
 Theoretical prediction of the thermal conductivity of uranium nitride vapor* (354)
 Thermal conductivity and diffusivity of $\text{U}_{40}\text{O}_{(9-y)}$ from 250 to 415 K* (459)
 Electric conductivity of melts of the LiF-misch metal fluoride system* (461)

conduits

New description of liquid metal heat transfer in closed conduits* (502)

conference

Proceedings of the 12th rare earth research conference. Vol. I* (11)
 All-union conference on chemistry of transplutonium elements (americium, curium, berkelium, californium)* (34)
 Proceedings of the international conference on liquid technology in energy production* (56)
 Proceedings of the 12th rare earth research conference. Vol. II* (65)
 5th international conference on plutonium and other actinides* (84)
 Proceedings of the international conference on liquid technology in energy production* (103)
 5th international conference on plutonium and other actinides. Book of abstracts* (109)
 5th international conference on plutonium and other actinides* (228)
 All-union conference on chemistry of transplutonium elements (americium, curium, berkelium, californium)* (307)
 All-union conference on chemistry of transplutonium elements (americium, curium, berkelium, californium)* (308)
 8th radiochemical conference* (347)
 8th radiochemical conference* (371)
 5th international conference on plutonium and other actinides* (415)

TITLE INDEX

conference

- Proceedings of the 12th rare earth research conference. Vol. II* (425)
 Abstracts of papers and communications submitted to the 5. all-union heat and mass transfer conference* (437)
 5th international conference on plutonium and other actinides* (479)
 Reactor conference, Duesseldorf, 30.3.-2.4.1976. Section 2: Fuel elements, fuel cycle and isotope technique* (480)
 Reactor conference, Duesseldorf, 30.3.-2.4.1976. Section 1: Reactor design and experiments* (509)
 2. all-union conference on crystallochemistry of intermetallic compounds* (549)
 Proceedings of the international conference on liquid technology in energy production* (590)
 5th international conference on plutonium and other actinides. Book of abstracts* (620)
 2. all-union conference on crystallochemistry of intermetallic compounds* (671)

configurations

- Structure of $4f^{sup 6}$ and $4f^{sup 8}$ levels and $Eu^{sup 3+}$ and $Tb^{sup 3+}$ configurations in short-range ultra-violet* (589)

conformal

- Application of molecular dynamics computations to the conformal ionic solution theory: Thermodynamics of binary molten salt mixtures* (550)

congress

11. european congress on molecular spectroscopy. Abstracts* (44)
 11. european congress on molecular spectroscopy. Abstracts* (46)

congruent

- Interval of congruent and incongruent sublimation of $3Kf \cdot ZrF_{sub 4}$ compound in the $KF-ZrF_{sub 4}$ system* (296)

constants

- Vibrational spectra and force constants of the elpasolites $Cs_{sub 2}KMF_{sub 6}$ ($M=Sc, Y, La, Gd, Yb$)* (41)

constituting

- Partly filled shells constituting anti-bonding orbitals with higher ionization energy than their bonding counterparts* (280)

constitution

- Phase analysis on simulated LWR fuel rods and its evaluation with the aid of constitution studies in the system uranium-zirconium-oxygen* (509)
 Phase analysis at simulated LWR fuel rods and their interpretation by constitution investigations in the uranium-zirconium-oxygen system* (510)
 Constitution equilibrium diagram of the $LiNO_{sub 3}-LiIO_{sub 3}$, $NaNO_{sub 3}-NaIO_{sub 3}$, $KNO_{sub 3}-KIO_{sub 3}$ system* (665)

constitutional

- Some constitutional studies on uranium-carbon- and plutonium-carbon-rhenium and technetium systems* (227)
 Some constitutional studies on uranium--carbon and plutonium--carbon--rhenium and technetium systems* (228)

constitutive

- Some trends in constitutive equation model development for high-temperature behavior of fast-reactor structural alloys* (527)

TITLE INDEX

constructional

Ceramic-base heat resisting constructional material* (348)

contactor

Measurement of mass transfer coefficients in a mechanically agitated, nondispersing contactor operating with a molten mixture of $\text{LiF--BeF}_2\text{--ThF}_4$ and molten bismuth* (79)

containments

Nuclear reactor containments* (569)

contaminated

Radiolytic gas generation in plutonium contaminated waste materials* (301)

contents

High-temperature heat contents and entropies of zirconium fluoride and zirconium sulfate* (591)

Control

Experience of the Central Control Laboratory (CCL) in accounting for and controlling nuclear material in Czechoslovakia* (349)

controlled

Contribution to the establishment of the phase diagram for the U--O--C system by means of x-ray diffractometry at high temperatures and under controlled pressure* (503)

controlling

Experience of the Central Control Laboratory (CCL) in accounting for and controlling nuclear material in Czechoslovakia* (349)

convection

Heat transfer measurements in a forced convection loop with two molten-fluoride salts: $\text{LiF--BeF}_2\text{--ThF}_2\text{--UF}_4$ and eutectic $\text{NaBF}_4\text{--NaF}$ * (585)

conversion

Thermodynamic assessment of the conversion of plutonium dioxide to plutonium monocarbide* (54)

Thermodynamics of the conversion of plutonium dioxide to plutonium monocarbide* (55)

Conversion of protactinium monocarbide to the penta- and tetra-halides* (80)

Practical application of conversion method in study of five-component mutual system consisting of nine salts NaF , Na_2MoO_4 , Na_2WO_4 , KF , K_2MoO_4 , K_2WO_4 , BaF_2 , BaMoO_4 , BaWO_4 * (516)

CoO

Thermodynamic properties of NbO , $\text{Fe}_{0.95}\text{O}$, CoO , NiO , Cu_2O and CuO oxides as comparative electrodes in electromotive forces method* (658)

cooled

Properties of structural materials for sodium-cooled fast breeder reactor* (580)

copper

Solubility and activity of oxygen in liquid indium and copper-indium alloys* (182)

Thermodynamic study of the binary system copper-indium. I. Calorimetric study* (290)

Thermodynamic study of the binary system copper--indium. II. Potentiometric study* (291)

On interaction of tellurium tetrachloride with copper (1), mercury (2), bismuth and iron (3) chlorides* (336)

TITLE INDEX

copper

Frequency spectrum and scattering law for liquid copper. A molecular dynamics investigation* (567)
 Electrochemical study of high-temperature stability of compounds between the rare earths and copper oxide* (634)

coprecipitation

Determination of thorium in rare earths chloride: coprecipitation with cerium (IV) iodate and spectrophotometric determination with thoron* (447)

corium

Properties of corium melts* (454)

correlation

Determination and correlation of mass transfer coefficients in a stirred cell* (236)
 Structural study of liquids with strong short-range correlation in the atomic distribution* (609)
 Structural study of liquids with strong short-range correlation in the atomic distribution* (653)

correlations

Investigations on radioactive fission product correlations: gamma spectrometry measurements on spent fuel assemblies discharged from the Trino Vercellese reactor at the end of the 2nd irradiation cycle* (75)

corrosion

EUTECTIC DATA: safety, hazards, corrosion, melting points, compositions, and bibliography* (276)
 Corrosion of liquid alloys of zinc with yttrium and with rare earth elements in LiCl--KCl melts* (335)
 Some regularities of the corrosion behavior of binary magnesium alloys* (439)
 Effect of composition and structure on crevice, intergranular, and stress corrosion of some wrought Ni--Cr--Mo alloys* (604)

corrosive

Simple Raman cell and furnace usable at temperatures higher than 1000\$sup 0\$ for corrosive melts* (204)

corundum

Dissolution of corundum in molten vanadates* (317)

counterparts

Partly filled shells constituting anti-bonding orbitals with higher ionization energy than their bonding counterparts* (280)

Covalent

Covalent compounds of uranium of the type F/sub x/ U(OTeF\$sub 5\$)/sub 6-x/* (571)

Cr

Thermodynamic properties of Cr, Mn, Co and Ni dichlorides from emf measurements on cells with solid electrolytes* (200)
 Preparation and structure of transition metal fluorides UEsup(II) Msup(IV) F\$sub 6\$[UEsup(II) : Cr-Zn; Msup(IV) : Hf, Zr]* (596)
 Effect of composition and structure on crevice, intergranular, and stress corrosion of some wrought Ni--Cr--Mo alloys* (604)

crevice

Effect of composition and structure on crevice, intergranular, and stress corrosion of some wrought Ni--Cr--Mo alloys* (604)

Criteria

Criteria for solid solution formation in NFE alloys* (205)

TITLE INDEX

critical

Experimental vapor pressure and critical point of sodium* (56)
 Critical assessment of equation of state data for UO_2 * (180)

CrO

Standard free energy of formation of liquid Na_2CrO_4 ,
 1100°K to 1220°K* (375)

cross

Study of KBeF_5 - DyF_3 cross-section of
 KF - BeF_2 - DyF_3 ternary system and some properties of
 this cross section glasses* (21)

cryolite

Raman spectra of Al_2O_3 solutions in molten cryolite
 and other aluminum fluoride containing melts* (203)

Cryometric

Cryometric studies of the system CdCl_2 - UCl_4 * (199)

cryoscopic

Estimate of cryoscopic calculations accuracy from fusibility
 diagrams* (670)

crystal

Mixed valence rare-earth halides and their unusual crystal
 structures* (29)

Phase relations, crystal structures and physical properties of
 nuclear fuels* (612)

crystalline

Basic research in crystalline and noncrystalline ceramic systems.
 Annual report, May 1, 1975--April 1, 1976* (728)

crystallochemistry

2. all-union conference on crystallochemistry of intermetallic
 compounds* (549)

2. all-union conference on crystallochemistry of intermetallic
 compounds* (671)

crystals

Vibrational spectra of transition metal hexafluoride crystals. I.
 Orthorhombic MoF_6 , WF_6 , UF_6 neat crystals. Technical report* (51)

Vibrational spectra of transition metal hexafluoride crystals.

III. Exciton band structures of MoF_6 , WF_6 , and UF_6 . Technical report* (52)

Interactions in inorganic molecular crystals - electronic spectra
 of ReF_6 pure and mixed crystals. Technical report, 1
 Jul--31 Dec 1975* (53)

Cs

Electronic structure of highly symmetric complexes of lanthanoids
 and actinoids. Pt. 3. 'Electronic' Raman spectrum of $\text{Cs}_2\text{U(IV)Cl}_6$ * (13)

Electronic structure of highly symmetric complexes of lanthanoids
 and actinoids. Pt. 4. The electronic Raman spectrum of $\text{Cs}_2\text{NaEu(III)Cl}_6$ * (14)

Vibrational spectra and force constants of the elpasolites Cs_2KMfF_6
 ($\text{M} = \text{Sc}, \text{Y}, \text{La}, \text{Gd}, \text{Yb}$)* (41)

Me_2ZrCl_6 - H_2 ($\text{Me} = \text{Na}, \text{K}, \text{Rb}, \text{Cs}$) system* (45)

Study of the compounds $\text{M}_3\text{UO}_2\text{F}_5$ ($\text{M} = \text{K}, \text{Rb}, \text{Cs},$
 NH_4) by i.r. absorption and Raman diffraction
 spectrophotometry. III. Effect of cation on vibration frequency
 of the ion $\text{UO}_2\text{F}_5^{3-}$ * (127)

TITLE INDEX

Cs

Cs--U--O phase diagram and its application to uranium--plutonium oxide fast reactor fuel pins* (175)
 Gases and carbon in metals (thermodynamics, kinetics, and properties). I. Alkali metals, alkaline earth metals, light metals (Li, Na, K, Rb, Cs; Ca, Sr, Ba; Be, Mg, Al)* (192)
 Cs-Hg amalgam thermodynamic properties* (339)
 Raman spectroscopic studies of yttrium (III) chloride--alkali metal chloride melts and of $\text{Cs}_2\text{NaYCl}_6$ and YCl_3 solid compounds* (489)
 Preparational, structural and vibrational study of $\text{M}_2\text{U}_2\text{O}_5\text{Cl}_4$, $2\text{H}_2\text{O}$ (M = Rb, Cs). Detection of a tetranuclear anion $[(\text{UO}_2)_4\text{O}_{12}\text{Cl}_8(\text{H}_2\text{O})_2]^{4-}$ * (498)
 Degree of thermal dissociation of M_3CaCl_3 (M = K, Rb, Cs) complexes during melting* (624)
 Electronic structure of highly symmetric lanthanoid and actinoid compounds. VI. The electronic Raman spectrum of $\text{Cs}_2\text{NaCe(III)Cl}_6$ * (678)

CsBr

Phase diagrams of LiCl, CsCl, CaCl and LiBr, CsBr, CaBr ternary systems* (256)
 LiBr, RbBr, CsBr ternary system* (268)
 Heats of formation of solid solutions of CsCl with CsBr* (543)
 Thermodynamics of solid solutions CsCl-RbCl and CsBr-RbBr at 25°C * (570)

CsCl

Interaction of the lithium-tin alloys with the LiCl-KCl-CsCl melted eutectic mixture* (91)
 Investigation into thermodynamic properties of FeCl_2 -CsCl system melts* (96)
 Phase diagrams of LiCl, CsCl, CaCl and LiBr, CsBr, CaBr ternary systems* (256)
 TlCl-RbCl-CsCl and NaI-TlI-CsI systems* (258)
 Thermal investigation of NdCl_3 - AlCl_3 -KCl and NdCl_3 - AlCl_3 -CsCl system* (289)
 Determination of the Np(IV)--Np(III) normal potential in fused LiCl-CsCl (55-45%) by absorption spectrophotometry* (386)
 Diagram of excess free energy of the PbCl_2 -CsCl- ZnCl_2 system* (402)
 Effect of luminescence centers on optical absorption, excitation and fluorescence spectra of CsCl:Pb * (491)
 Heats of formation of solid solutions of CsCl with CsBr* (543)
 Thermodynamics of solid solutions CsCl-RbCl and CsBr-RbBr at 25°C * (570)
 Interaction in AgCl-CsCl, CsCl-KCl systems and liquidus surface of AgCl-CsCl-KCl ternary system* (659)

CsF

Molecular vapor composition and mass spectra in KF-ZrF_4 , CsF-ZrF_4 systems* (294)
 CsF-HoF_3 system* (572)

CsI

TlCl-RbCl-CsCl and NaI-TlI-CsI systems* (258)

CsNO

LiNO_3 - CsNO_3 - $\text{Ca(NO}_3)_2$ and NaNO_3 - CsNO_3 - $\text{Ca(NO}_3)_2$ system* (255)

TITLE INDEX

- Csub Spectral properties of impurity molecules of the Csub (3v) symmetry groups in solid solutions of halide salts* (44)
- Cu Thermodynamic properties of NbO, Fe_{(0,95)O}, CoO, NiO, Cu₂O and CuO oxides as comparative electrodes in electromotive forces method* (658)
Investigation of the solubility and diffusion of Fe atoms in Cu at high temperature using molten salt electrochemistry* (693)
- cubic Nonstoichiometry in cubic uranium sesquinitride* (611)
- CuCl Phase diagrams of BiCl₃-CuCl-TeCl₄ and CuCl-FeCl₃-TeCl₄ systems* (337)
- CuO Thermodynamic properties of NbO, Fe_{(0,95)O}, CoO, NiO, Cu₂O and CuO oxides as comparative electrodes in electromotive forces method* (658)
- curium Absorption spectra of curium (3) chloride complexes* (34)
All-union conference on chemistry of transplutonium elements (americium, curium, berkelium, californium)* (307)
All-union conference on chemistry of transplutonium elements (americium, curium, berkelium, californium)* (308)
- Current Current yield during separation of lithium on a liquid bismuth cathode* (430)
- curves Vacancies and melting curves of metals at high pressure* (213)
Intensity-voltage curves of hydrogen diffusion electrodes in eutectic LiCl-KCl* (507)
K₂HfF₆-K₂ZrF₆-1.25 percent of HF system at 40°C with other solubility curves from 25°C to 70°C* (541)
Investigation into properties of liquidus curves of binary salt system melts where chemical compounds formation. 2. Systems of mono-monivalent and bi-bivalent cations with common anion* (708)
- cycles Approximation of tabulated thermodynamic data for purposes of optimization of two-phase liquid-metal power cycles* (343)
- Czechoslovakia Experience of the Central Control Laboratory (CCL) in accounting for and controlling nuclear material in Czechoslovakia* (349)
- damage Experimental investigation of cavitation inception and damage in a flowing sodium environment* (103)
- Daten Physik Daten/Physics Data* (192)
- Dec Interactions in inorganic molecular crystals - electronic spectra of ReF₆ pure and mixed crystals. Technical report, 1 Jul--31 Dec 1975* (53)
Thermochemistry of gaseous metal oxides. Final report, 1 Apr 1973--31 Dec 1975* (239)

TITLE INDEX

Dec

Thermochemistry of gaseous compounds of metals. Annual summary report 5 Dec 74--4 Dec 75* (240)

Thermocchemistry of gaseous compounds of metals. Annual summary report, 5 Dec 1975--4 Dec 1976* (241)

Basic mechanisms providing oxidation resistance in structural metals at high temperatures. II. The defect structure of nonstoichiometric rutile (TiO_{2-x}) and nonstoichiometric rutile containing 1 to 10 mole % $\text{Nb}_{2-x}\text{O}_{6-x}$. Final report 21 Dec 1973-20 Dec 1974* (542)

Research on a high-energy nonaqueous battery system. Quarterly progress report No. 1, 1 Dec 1965--1 Mar 1966* (720)

December

Lithium--silver chloride secondary battery investigation. Final report, December 1963--November 1964* (118)

Electrochemical methods for monitoring purity of molten salts. Semiannual progress report, July--December 1975* (721)

Decomposition

Decomposition, stoichiometry and structure of neptunium oxides* (171)

The spectral properties of uranium hexafluoride and its thermal decomposition products* (345)

Thermal decomposition of solid metal oxides* (395)

Thermal decomposition of intermetallic compounds of platinum with alkaline earth and lanthanide metals at high temperatures* (399)

Chemistry of elements niobium and tantalum. 87. Studies on the thermal formation and decomposition of the complexes $[\text{NbCl}_{12}]^{\text{sup } 2+}$, $[\text{NbI}_{12}]^{\text{sup } 3+}$, and $[\text{TaX}_{12}]^{\text{sup } (n+)}$ ($X=\text{Cl}, \text{Br}, \text{I}$)* (561)

Decomposition of the UN-CeN solid solution* (581)

Solubility of nitrogen in liquid lithium and thermal decomposition of solid Li_3N * (702)

defect

Calculation of the oxygen potential of $(\text{U}, \text{Pu})\text{O}_{2-x}$ with defect models and measured oxygen potential data in the temperature range from 1,000 to 1,700 K* (74)

Basic mechanisms providing oxidation resistance in structural metals at high temperatures. II. The defect structure of nonstoichiometric rutile (TiO_{2-x}) and nonstoichiometric rutile containing 1 to 10 mole % $\text{Nb}_{2-x}\text{O}_{6-x}$. Final report 21 Dec 1973-20 Dec 1974* (542)

defects

U.S.-Japan joint seminar on defects and diffusion in solids* (611)

definition

Fine definition of ir spectra from high temperature interactions of $\text{U} + \text{O}_{2-x}$. Part II. Final report, 1 July 1974--30 June 1975* (218)

Degree

Degree of thermal dissociation of MCl_3 ($\text{M}=\text{K}, \text{Rb}, \text{Cs}$) complexes during melting* (624)

degreeK

Thermophysical and electrophysical properties of uranium hexafluoride at temperatures of $(1-11) \times 10^3$ degreeK and pressures of 0.1--100 atm* (302)

TITLE INDEX

degrees

Electrode reactions in molten sodium nitrate--barium nitrate eutectic at 350 degrees C* (42)

delta

High-temperature microcalorimetric measurements of partial molar enthalpy of solution, $\Delta H(O_{2})$, in plutonium + oxygen* (107)

Densities

Densities of electronic states in liquid transition metals* (422)
Densities of liquid carbides and nitrides of uranium and plutonium* (579)
Electronic states densities from structure and thermodynamic data for liquid transition metals* (705)

Density

Density and temperature dependence of the surface tension of molten bismuth, lead, and bismuth-lead alloys* (1)
Density, viscosity, electric conductivity and meltability of $Na_{3}AlF_{6}$ - $La_{2}O_{3}$ and $Na_{3}AlF_{6}$ - $Sm_{2}O_{3}$ systems* (43)
Study on conductivity, density, and viscosity of molten salt systems* (112)
Composition of nonaqueous solutions of potential use in high energy density batteries. Final report, 15 Nov 1969--14 Nov 1972* (119)
Density, surface tension, and viscosity of uranium trichloride-sodium chloride melting* (141)
Density, surface tension and viscosity of melts of uranium trichloride-lithium chloride* (142)
Thermodynamic perturbation theory for the surface tension and ion density profile of a liquid metal* (167)
Spectral density functions for disordered systems* (229)
Density and viscosity of distillable products from $ZrCl_{4}$ and $HfCl_{4}$ interaction with $POCl_{3}$ * (458)
Density and enthalpy of liquid alkaline metals at pressures up to 1500 bar* (599)
High density thoria-silica-metal (III) oxide fibers* (710)

Deoxidation

Deoxidation thermodynamics of cerium and lanthanum in liquid iron* (654)

derivation

Theory for the derivation of a single hard-sphere potential describing the overall molecular interactions in binary liquid metal alloys as a function of temperature and composition* (525)

described

Thermodynamic properties of $(U,Pu)O_{(2+x)}$ described by a cluster model* (133)

describing

Theory for the derivation of a single hard-sphere potential describing the overall molecular interactions in binary liquid metal alloys as a function of temperature and composition* (525)

design

Proceedings of the symposium and workshop on advanced battery research and design, March 22--24, 1976* (269)
Reactor conference, Duesseldorf, 30.3.-2.4.1976. Section 1: Reactor design and experiments* (509)

TITLE INDEX

design

Summary of U.S. LMPER programs on high temperature structural design and associated materials testing* (731)

detection

High-temperature unit to the IKS-21 spectrometer for the detection of molten salts absorption spectra* (208)
 Use of the SF-8 detection spectrophotometer for measuring absorption spectra of molten salts* (331)
 Preparational, structural and vibrational study of $M_{20}U_{20}Cl_{40}$, $2H_{20}(M = Rb, Cs)$. Detection of a tetranuclear anion $[U_{20}O_{20}]_{40}Cl_{80}(H_{20}O_{20})_{20}]^{4-}$ * (498)

diagram

X-ray study of the plutonium--oxygen phase diagram at ambient temperatures up to $1100 \pm 0^\circ C$ * (62)
 Phase diagram of system $LiCl-KCl-CaCrO_4$ * (116)
 Phase diagram for the ternary system $CaCl_2-KCl-CaCrO_4$ * (117)
 Phase diagram of the $ThCl_4-LiCl-PbCl_2$ system* (143)
 Cs--U--O phase diagram and its application to uranium--plutonium oxide fast reactor fuel pins* (175)
 Phase diagram and properties of Nd--As alloys* (321)
 P--t diagram of the system uranium--nitrogen* (373)
 Phase diagram of the ternary system of lithium-, rubidium-, and cesium nitrates* (377)
 Diagram of excess free energy of the $PbCl_2-CsCl-ZnCl_2$ system* (402)
 Phase relations in the alkaline earth fluoride-yttria systems (in three parts). 1. The $CaF_2-Y_{20}O_{30}$ phase diagram* (468)
 Phase relations in the alkaline earth fluoride-yttria systems (in three parts). 2. The $BaF_2-Y_{20}O_{30}$ phase diagram* (469)
 Contribution to the establishment of the phase diagram for the U--O--C system by means of x-ray diffractometry at high temperatures and under controlled pressure* (503)
 Meltability diagram of $RbCl-BeCl_2$ system* (506)
 Constitution equilibrium diagram of the $LiNO_3-LiIO_3$, $NaNO_3-NaIO_3$, KNO_3-KIO_3 system* (665)
 Phase diagram of $NaNO_3-Nd(NO_3)_3$ system* (667)

diagrams

Phase diagrams of ternary systems containing sodium and potassium chlorides, thorium tetrachloride and plutonium trichloride* (148)
 Phase diagrams of $Li_{20}MoO_{40}-ZnMoO_4$, $K_{20}MoO_4$ and $K_{20}WO_4-ZnWO_4$ systems* (160)
 Calculation and discussion of state diagrams in the system Pu-F-O-H* (222)
 Phase diagrams of $LiCl, CsCl, CaCl$ and $LiBr, CsBr, CaBr$ ternary systems* (256)
 Phase diagrams of $BiBr_3-BiI_3$ system* (288)
 Phase diagrams of $BiCl_3-CuCl-TeCl_4$ and $CuCl-FeCl_3-TeCl_4$ systems* (337)
 Phase diagrams in the pseudobinary systems of $Li_{20}BeF_{40}$, alkali fluoroborates and alkali fluorides* (473)

TITLE INDEX

diagrams

Thermodynamic analysis of melting point diagrams of systems consisting of alkali metal chlorides, thallium chloride and beryllium chloride* (505)
 Theoretical concepts useful in the calculation or storage of phase diagrams of ionic systems* (552)
 Phase diagrams of the $\text{Mg}(\text{NO}_3)_2\text{-Nd}(\text{NO}_3)_3$ system* (668)
 Estimate of cryoscopic calculations accuracy from fusibility diagrams* (670)
 Phase diagrams* (679)
 Phase diagrams* (680)

dianion

Actinide metallocarborane complex; synthesis and x-ray structure determination of the bis[n⁵- (3)-1,2-dicarbollyl]dichlorouranium(IV) dianion* (194)

dibromide

Optical spectra of the dibromide and diiodide ions in the matrix-isolated $\text{M}^{\text{sup}} + \text{Br}_2^{\text{sup}} -$ and $\text{M}^{\text{sup}} + \text{I}_2^{\text{sup}} -$ species* (17)

dibromides

Investigation into interaction of indium tribromide with dibromides of manganese, cobalt and nickel in melt* (153)

dicarbide

Thermal diffusivity, thermal conductivity and spectral emissivity of uranium dicarbide at high temperatures* (132)

dicarbollyl]dichloro

Actinide metallocarborane complex; synthesis and x-ray structure determination of the bis[n⁵- (3)-1,2-dicarbollyl]dichlorouranium(IV) dianion* (194)

dicesium

IR spectroscopic study of dipotassium and dicesium uranyl tetrabromides* (483)

dichloride

Optical spectra of the difluoride, dichloride, and trichloride ions in the matrix-isolated $\text{M}^{\text{sup}} + \text{F}_2^{\text{sup}} -$, $\text{M}^{\text{sup}} + \text{Cl}_2^{\text{sup}} -$, and $\text{M}^{\text{sup}} + \text{Cl}_3^{\text{sup}} -$ species* (16)
 Melting point of europium dichloride* (361)

dichlorides

Thermodynamic properties of Cr, Mn, Co and Ni dichlorides from emf measurements on cells with solid electrolytes* (200)
 Kinetics and mechanism into preparation of samarium-, europium- and ytterbium dichlorides and their some properties* (253)

dichloroiodates

Thermochemistry of polyhalides. Pt.1. Caesium and rubidium dichloroiodates(I)* (178)

differences

Some structure differences of double oxides of indium and gallium with antimony(5)* (657)

Differential

Differential-spectrophotometric determination of niobium in its compounds* (490)

differentiation

Determination of the burn up and differentiation of the fission yields of uranium-thorium-fuels* (214)

TITLE INDEX

diffraction

- X-ray diffraction study into molten bromides and iodides of alkali metals* (20)
 High temperature thermal expansion of ThO_2 , MgO and Y_2O_3 by X-ray diffraction* (242)
 Structure of molten Bi--Sb alloys by means of neutron diffraction* (359)
 X-ray diffraction and infrared: Raman spectroscopic study of the hydrogen dinitrate ion, $(\text{O}_2\text{N}-\text{H}-\text{ONO})^+$, in cesium hydrogen dinitrate* (695)

diffractometry

- Contribution to the establishment of the phase diagram for the U--C--C system by means of x-ray diffractometry at high temperatures and under controlled pressure* (503)

Diffuse

- Diffuse reflectance spectra of protactinium(IV), uranium(IV) and neptunium(IV) tetrachlorides and tetraformates* (562)

diffusion

- Study of the compounds $\text{M}_3\text{UO}_2\text{F}_5$ ($\text{M} = \text{K}, \text{Rb}, \text{Cs}, \text{NH}_4$) by i.r. absorption and Raman diffusion spectrophotometry. III. Effect of cation on vibration frequency of the ion UO_2F_3^+ (127)
 Atomic diffusion in alloys exposed to liquid sodium* (181)
 Self-diffusion in liquid metals* (198)
 Diffusion coefficients in molten metals* (328)
 Diffusion of uranium species in several molten chlorides* (408)
 Monte Carlo analysis of diffusion and thermodynamics in UO_{2+x} * (441)
 Intensity-voltage curves of hydrogen diffusion electrodes in eutectic LiCl--KCl * (507)
 U.S.-Japan joint seminar on defects and diffusion in solids* (611)
 Diffusion in liquid metal systems. Final report* (651)
 Investigation of the solubility and diffusion of Fe atoms in Cu at high temperature using molten salt electrochemistry* (693)

diffusivity

- Uranium carbide: thermal diffusivity, thermal conductivity and spectral emissivity at high temperatures* (121)
 Thermal diffusivity, thermal conductivity and spectral emissivity of uranium dicarbide at high temperatures* (132)
 Thermal conductivity and diffusivity of U_4O_{9-y} from 250 to 415 K* (459)

difluoride

- Optical spectra of the difluoride, dichloride, and trichloride ions in the matrix-isolated $\text{M}^+ \text{F}_2^+$, $\text{M}^+ \text{Cl}_2^+$, and $\text{M}^+ \text{Cl}_3^+$ species* (16)

digital

- Calculated method of eutectics of multicomponent systems on base of algorithm and digital computers* (517)

diiodide

- Optical spectra of the dibromide and diiodide ions in the matrix-isolated $\text{M}^+ \text{Br}_2^+$ and $\text{M}^+ \text{I}_2^+$ species* (17)

TITLE INDEX

dilatation

Thermal lattice dilatation of some transition metal compounds* (508)

dilute

Some thermodynamic properties of dilute solutions of actinide chlorides in (Li-K)Cl and in (Na-K)Cl eutectics* (406)
Thermodynamic properties of dilute solutions of ThCl_4 in (Li-K)Cl and (Na-K) eutectics* (407)

diluted

Thermodynamic properties of diluted solutions of MnCl_2 in alkali metal chlorides* (6)
Thermodynamic properties of diluted sodium solutions in liquid indium* (388)
Investigation of liquid-steam equilibrium in system with diluted solutions of metal fluorides in uranium hexafluoride* (526)

dinitrate

X-ray diffraction and infrared: Raman spectroscopic study of the hydrogen dinitrate ion, $(\text{O}_2\text{NO-H-ONO}_2)^+$, in cesium hydrogen dinitrate* (695)

dioxide

Thermodynamic assessment of the conversion of plutonium dioxide to plutonium monocarbide* (54)
Thermodynamics of the conversion of plutonium dioxide to plutonium monocarbide* (55)
Uranium dioxide as a ceramic nuclear fuel* (226)
Titanium dioxide solubility in melts of lithium- and alkaline earth metal chlorides* (264)
Effective equation-of-state measurements on uranium dioxide* (537)
Methods for the accountability of uranium dioxide* (600)
Standard methods for chemical, mass spectrometric, and spectrochemical analysis of nuclear-grade uranium dioxide powders and pellets* (711)
Standard methods for chemical, mass spectrometric, and spectrochemical analysis of nuclear-grade plutonium dioxide powders and pellets* (712)
Fluorination process studies of plutonium dioxide by fluid-bed* (725)

dioxides

Determination of oxygen potentials of hyperstoichiometric U--Pu dioxides in the temperature range 1,500-1,850 K* (109)
Determination of oxygen potentials of hyperstoichiometric U-Pu dioxides in the temperature range 1500-1850 K* (110)

dipole

Ionic-thermoelectric study of the dipole relaxation and equilibrium in Gd-doped SrF_2 * (410)

dipotassium

IR spectroscopic study of dipotassium and dicesium uranyl tetrabromides* (483)

disassembly

Hydrodynamics of post-disassembly fuel expansions* (310)

discharge

Structure and anodic discharge behavior of lithium-boron alloys in the LiCl-KCl eutectic melt.* (274)

TITLE INDEX

discharged

Investigations on radioactive fission product correlations: gamma spectrometry measurements on spent fuel assemblies discharged from the Trino Vercellese reactor at the end of the 2nd irradiation cycle* (75)

discrete

Discrete variation of the structure and texture of liquid metals formed by components with discrete packing* (177)

disodium

Reactions of liquid sodium with transition-metal oxides. VIII. The oxides of chromium-(III), -(IV), and -(VI) and disodium chromium(VI) tetraoxide* (35)

disordered

Spectral density functions for disordered systems* (229)

Dispersion

Dispersion of metallic lithium in various molten salts* (452)

dissociation

Mass spectrometric determination of dissociation energies of high-temperature molecules: ScS, YS, LaS, ZrS, UO* (597)

Degree of thermal dissociation of $M\text{CaCl}_3$ (M-K, Rb, Cs) complexes during melting* (624)

Dissolution

Dissolution of corundum in molten vanadates* (317)

Investigation of solubility and thermodynamics of dissolution of salts in mixed solvents* (347)

Electrochemistry of niobium and tantalum, (5). Anodic dissolution of tantalum and the average valence of tantalum ions in molten KCl* (450)

PuO_2 dissolution problem for LWR plutonium recycle and LMFBR fuels: fabrication and reprocessing problems and their resolution* (455)

dissolved

Characterisation of the chemical activities of non-metals dissolved in liquid alkali metals* (64)

dissolving

Kinetic peculiarities of nitrogen dissolving in iron-chromium and iron-chromium-nickel melts* (172)

dissoved

Characterisation of the chemical activities of non-metals dissolved in liquid alkali metals* (63)

distillable

Density and viscosity of distillable products from ZrCl_4 and HfCl_4 interaction with POCl_3 * (458)

Distribution

Distribution coefficient for U and Pu in the system $(\text{UPu})\text{C} + (\text{UPu})_2\text{C}_3$ * (134)

Radial oxygen distribution in the fuel of irradiated mixed oxide fuel elements with different final stoichiometries* (316)

Structural study of liquids with strong short-range correlation in the atomic distribution* (609)

Structural study of liquids with strong short-range correlation in the atomic distribution* (653)

divalent

Negative temperature coefficients of electrical resistivity: the divalent liquid metals Eu, Yb and Ba* (224)

TITLE INDEX

Division

Westinghouse Advanced Reactors Division* (57)
 Chemical Engineering Division physical inorganic chemistry.
 Annual report, July 1975--June 1976* (58)
 Radiochemistry Division annual progress report for 1973* (271)
 Chemical Engineering Division fuels and materials chemistry.
 Annual report, July 1975--June 1976* (281)
 Chemical Engineering Division, reactor safety and physical
 property studies annual report, July 1974--June 1975* (367)
 Analytical Chemistry Division annual progress report for period
 ending November 30, 1975* (423)
 Chemistry Division annual progress report for period ending
 November 1, 1975* (729)
 Chemistry Division annual progress report for period ending
 November 1, 1975* (730)

doped

Optical absorption spectrum of UO_2 doped LiF * (30)
 Zeeman effect and satellite carrier structure in absorption
 spectra of pure and doped praseodymium and europium double
 nitrates* (165)
 Phase transition in doped UO_2 . I. Heat capacity
 measurement* (261)
 Electrical conduction in CaO -doped thoria electrolytes* (389)
 Ionic-thermocurrent study of the dipole relaxation and
 equilibrium in Gd -doped SrF_2 * (410)
 Complex impedance of electrochemical cells based on yttria doped
 thoria* (568)

double

Zeeman effect and satellite carrier structure in absorption
 spectra of pure and doped praseodymium and europium double
 nitrates* (165)
 Double tungstate $\text{K}_5\text{La}(\text{WO}_4)_4$ * (338)
 Double potassium rare earth tungstates* (639)
 Some structure differences of double oxides of indium and gallium
 with antimony(5)* (657)

du

Comptes rendus du 95* (621)

Duesseldorf

Reactor conference, Duesseldorf, 30.3.-2.4.1976. Section 2: Fuel
 elements, fuel cycle and isotope technique* (480)
 Reactor conference, Duesseldorf, 30.3.-2.4.1976. Section 1:
 Reactor design and experiments* (509)

Dy

Comparative study of systems of heavier rare-earth metals with
 hydrogen [Gd , Tb , Dy , Lu]* (418)
 Gd , Tb , Dy , Ho , Er solubility in chromium* (610)

DyF

Study of $\text{KBe}_{2/5}\text{F}_{5/3}\text{-DyF}_3$ cross-section of
 $\text{KF-Bef}_{2/3}\text{-DyF}_3$ ternary system and some properties of
 this cross section glasses* (21)

dynamical

Lattice-dynamical calculation of the thermodynamic properties of
 NaCl , KCl and RbCl * (231)

TITLE INDEX

dynamics

Using molecular dynamics method for the evaluation of Fe-Ni and Al-Zn melts structure properties* (512)
 Application of molecular dynamics computations to the conformal ionic solution theory: Thermodynamics of binary molten salt mixtures* (550)
 Frequency spectrum and scattering law for liquid copper. A molecular dynamics investigation* (567)

dysprosium

Formation enthalpy of gadolinium, terbium and dysprosium trifluorides* (603)

E

Fuel-clad interaction. C-E thermo-structural fuel evaluation methods* (719)

earth

Proceedings of the 12th rare earth research conference. Vol. I* (11)
 Mixed valence rare-earth halides and their unusual crystal structures* (29)
 Thermodynamic properties of some rare earth carbonates and yttrium carbonates* (36)
 Proceedings of the 12th rare earth research conference. Vol. II* (65)
 Saturation vapor pressure in iron chloride-alkaline earth metal chlorides systems* (97)
 Temperature dependence of vapor pressure of rare earth halides and their alloys* (99)
 Investigation into reactions between trioctylammonium nitrates and rare earth nitrates by X-ray electron spectroscopy* (113)
 Synthesis and properties of rare earth titanotungstates* (114)
 Compatibility of molten rare earth metals with refractory materials* (115)
 Optical absorption Zeeman spectroscopy of rare earth orthochromites. II. Theory* (122)
 Optical absorption Zeeman spectroscopy of rare earth orthochromites. I. ErCrO_3 * (123)
 Thermodynamics and kinetics of the formation of rare earth intermetallics* (137)
 Interaction of uranium trichloride with alkaline earth metal chlorides* (139)
 Calculation of thermodynamic activity and liquidus lines of components in binary systems of plutonium trichloride with alkali and alkaline earth metal chlorides* (147)
 Revised model for 5d-electrons in the heavy rare-earth metals* (157)
 Stability of rare earth metaniobates in hydrogen atmosphere* (164)
 Physicochemical investigation of rare earth titanoniobates, LnTiNbO_6 * (173)
 Vibration spectra of rare earth oxytungstates- Ln_2WO_6 composition* (185)
 Gases and carbon in metals (thermodynamics, kinetics, and properties). I. Alkali metals, alkaline earth metals, light metals (Li, Na, K, Rb, Cs; Ca, Sr, Ba; Be, Mg, Al)* (192)

TITLE INDEX

earth

- Thermodynamics of reactions of rare earth oxides with niobium oxide* (206)
- Electron structure of rare-earth metal tetraborides* (215)
- Hydrogen solubility in rare earth intermetallic compounds* (217)
- Entropies of liquid rare earth metals* (230)
- Titanium dioxide solubility in melts of lithium- and alkaline earth metal chlorides* (264)
- Statistical mechanics of intermediate-valence rare earth compounds* (279)
- Stability of rare earth tetrafluorides* (312)
- Rare earth elements* (319)
- Investigation of formation thermodynamic of rare earth aluminates with perovskite structure* (326)
- Corrosion of liquid alloys of zinc with yttrium and with rare earth elements in LiCl--KCl melts* (335)
- Investigation of some rare earth aluminates by the method of infrared spectroscopy* (357)
- $R\text{sub } 3T\text{sub } 2$ compounds (R = rare earth or Y; T = Rh, Pd, Pt) with the rhombohedral $\text{Er}\text{sub } 3\text{Ni}\text{sub } 2$ structure type* (365)
- Theory of rare-earth alloys* (376)
- Rare earth metal--metal halide systems. 19. Structural characterization of the reduced holmium chloride $\text{Ho}\text{sub } 5\text{Cl}\text{sub } 11$ * (378)
- Rare earth 4d spectra in rare earth trifluorides* (382)
- Pressure of saturated vapor of rare earth and yttrium bromides* (391)
- Hydrogen absorption of rare-earth (3d) transition intermetallic compounds* (392)
- Thermal decomposition of intermetallic compounds of platinum with alkaline earth and lanthanide metals at high temperatures* (399)
- Progress in the spectroscopy of rare earth and transition metal ions* (412)
- Comparative study of systems of heavier rare-earth metals with hydrogen [Gd, Tb, Dy, Lu]* (418)
- Proceedings of the 12th rare earth research conference. Vol. II* (425)
- Thermogravimetric investigation of interaction of some rare earth oxides with ammonium bifluoride* (427)
- Electronegativities of the rare-earth elements* (429)
- Structural or solid state chemistry of rare earth sesquioxides from Goldschmidt to the present time* (438)
- Rare earth metals* (461)
- Rare earth metals* (462)
- Phase relations in the alkaline earth fluoride-yttria systems (in three parts). 1. The $\text{CaF}\text{sub } 2$ -- $\text{Y}\text{sub } 2\text{O}\text{sub } 3$ phase diagram* (468)
- Phase relations in the alkaline earth fluoride-yttria systems (in three parts). 2. The $\text{BaF}\text{sub } 2$ -- $\text{Y}\text{sub } 2\text{O}\text{sub } 3$ phase diagram* (469)
- Optical properties of rare earth trifluorides* (484)
- Solubility of rare earth and yttrium oxochlorides in molten salts - $\text{MCl}\text{sub } 2$ (M-Mg, Ca, Sr, Ba)* (495)
- Separation of alkaline earth from alkali metal fission products in the SOLAR on-line mass spectrometer* (534)

TITLE INDEX

earth

Radiative and non-radiative transitions of rare-earth ions in glasses* (538)
 Rare earth metals* (546)
 Electronic and magnetic properties of rare earth-Sn₃ compounds from ¹¹⁹Sn Moessbauer spectroscopy* (559)
 Rare earth and actinide oxides. Thermodynamic and electron microscopy studies* (625)
 Double potassium rare earth tungstates* (639)
 Glass formation in nitrate systems containing rare earth ions* (666)
 Structure of rare earth metals in the liquid state* (688)

earths

Advances on thermochemistry of rare earths alloys* (65)
 Chemical separation and spectrographic estimation of rare earths and yttrium in PuO₂ and (U,Pu)O₂* (125)
 Binary molybdates of rare earths and silver (I)* (207)
 Measurement of fluorescence of rare earths in molten salts (Nd³⁺ in NaNO₃-KNO₃ eutectic)* (233)
 Rare earths* (280)
 Interaction of atoms of rare earths-metal alloys* (320)
 Determination of thorium in rare earths chloride: coprecipitation with cerium (IV) iodate and spectrophotometric determination with thoron* (447)
 Rare earths* (492)
 Rare Earths* (538)
 Spectrophotometric determination of rare earths and yttrium in gallium base alloys* (623)
 Electrochemical study of high-temperature stability of compounds between the rare earths and copper oxide* (634)

EER

Fission rates, burnup, and neutron flux-fluence-spectra characterization for mixed oxide fuel experiments in the EER-II* (416)

Effective

Effective equation-of-state measurements on uranium dioxide* (537)

elastic

Spin structure and elastic properties of uraniummononitride* (368)

electric

Density, viscosity, electric conductivity and meltability of Na₃AlF₆-La₂O₃ and Na₃AlF₆-6f-Sm₂O₃ systems* (43)
 Electric conductivity of melts of the LiF-misch metal fluoride system* (461)

electrical

Negative temperature coefficients of electrical resistivity: the divalent liquid metals Eu, Yb and Ba* (224)
 Aspects of the solution chemistry of liquid alkali metals as elucidated from electrical resistivity studies* (247)
 Electrical conduction in CaO-doped thoria electrolytes* (389)
 Improved model for solutions of nonmetals in liquid alkali metals. Calculation of enthalpy of solution and electrical resistivity* (486)

TITLE INDEX

electrical

Reaction rates and electrical resistivities of the hydrogen isotopes with, and their solubilities in, liquid lithium* (529)

Electrochemical

Electrochemical machining of the molybdenum VM-1 alloy* (129)

Electrochemical studies of the sodium--bismuth system* (187)

Electrochemical behavior of tellurium (4) in oxychloride melts* (221)

Electrochemical behavior of U(III) and U(IV) in molten LiCl-KCl eutectic* (293)

Electrochemical characterization of nonaqueous systems for secondary battery application. Quarterly report, February--April 1968* (380)

Electrochemical reduction of molybdenum oxychloride in molten lithium chloride - potassium chloride eutectic* (515)

Complex impedance of electrochemical cells based on yttria doped thorium* (568)

Electrochemical study of high-temperature stability of compounds between the rare earths and copper oxide* (634)

Electrochemical methods for monitoring purity of molten salts.

Semiannual progress report, July--December 1975* (721)

Research on electrochemical parameters for nonaqueous batteries: task I. Quarterly progress report No. 3* (727)

Electrochemistry

Electrochemistry of molten salts* (263)

Electrochemistry of niobium and tantalum, (5). Anodic dissolution of tantalum and the average valence of tantalum ions in molten KCl* (450)

Electrochemistry of niobium and tantalum. VI. Electrode potentials of tantalum chlorides in molten KCl* (451)

Thermodynamic studies of gallium--indium liquid alloys by solid state electrochemistry with oxide electrolytes* (513)

Investigation of the solubility and diffusion of Fe atoms in Cu at high temperature using molten salt electrochemistry* (693)

Electrode

Electrode reactions in molten sodium nitrate--barium nitrate eutectic at 350 degrees C* (42)

Studies on the negative alloy electrode* (269)

Electrode potentials of tantalum chlorides in a molten NaCl-KCl eutectic* (449)

Electrochemistry of niobium and tantalum. VI. Electrode potentials of tantalum chlorides in molten KCl* (451)

electrodes

Electronic conductivity and polarization effects of solid ThO_2 - Y_2O_3 electrolytes and of oxygen containing V-metal electrodes* (193)

Intensity-voltage curves of hydrogen diffusion electrodes in eutectic LiCl--KCl* (507)

Thermodynamic properties of NbO , $\text{Fe}_{0.95}\text{O}$, CoO , NiO , Cu_2O and CuO oxides as comparative electrodes in electromotive forces method* (658)

electrolysis

Composition gradient during electrolysis in mixtures analogous to molten salt battery electrolytes* (234)

Preparation of thorium metal by electrolysis of fused salt ThCl_4 - LiCl - NaCl - KCl (eut) system* (703)

TITLE INDEX

electrolyte

Nonaqueous electrolyte systems: ionic transport in nonaqueous media. Interim technical report No. 7* (246)

Measurement of the cell voltage of $\text{ThO}_2\text{--Y}_2\text{O}_3$ -solid electrolyte cells in the range of mixed conduction* (535)

Investigations on chemical properties of thorium oxide solid electrolyte ceramics* (652)

Lithium allcys--chlorine secondary battery. I. Basic studies on a lithium allcys--chlorine secondary battery using a molten salt as the electrolyte* (704)

electrolytes

Electronic conductivity and polarization effects of solid $\text{ThO}_2\text{--Y}_2\text{O}_3$ electrolytes and of oxygen containing Va-metal electrodes* (193)

Thermodynamic properties of Cr, Mn, Co and Ni dichlorides from emf measurements on cells with solid electrolytes* (200)

Composition gradient during electrolysis in mixtures analogous to molten salt battery electrolytes* (234)

Molten-salt electrolytes--transport properties* (277)

Phase equilibrium studies of lithium halide-containing electrolytes* (282)

Electrical conduction in CaO-doped thoria electrolytes* (389)

Thermodynamic studies of gallium--indium liquid alloys by solid state electrochemistry with oxide electrolytes* (513)

Thermodynamics of electrolytes. 8. High-temperature properties, including enthalpy and heat capacity, with application to sodium chloride* (586)

Zirconia and yttria--thoria ceramics find new uses as solid electrolytes* (595)

electrometallurgical

Radiotracer applications in electrometallurgical processing* (76)

Electromodulation

Electromodulation spectroscopy of sc and fcc phase TlCl and TlBr * (411)

Electromotive

Electromotive force measurements in molten lithium--magnesium allcys* (551)

Thermodynamic properties of NbO , $\text{Fe}_{0.95}\text{O}$, CoO , NiO , Cu_2O and CuO oxides as comparative electrodes in electromotive forces method* (658)

electron

Thermodiffusion measurements in liquid alloys and electron transport heat calculations in the case of almost-free electrons* (33)

Investigation into reactions between trioctylammonium nitrates and rare earth nitrates by X-ray electron spectroscopy* (113)

Electron spectrum of light alkali metals* (161)

Electron structure of rare-earth metal tetraborides* (215)

Electron structure and bonding in ThO_2 and UO_2 * (223)

Anomalous melting of f electron metals (with attention to Pu)* (318)

Electron contribution to the liquid metal viscosity* (346)

Rare earth and actinide oxides. Thermodynamic and electron microscopy studies* (625)

TITLE INDEX

Electronegativities

Electronegativities of the rare-earth elements* (429)

Electronic

Electronic structure of highly symmetric complexes of lanthanoids and actinoids. Pt. 3. 'Electronic' Raman spectrum of $\text{Cs}_{\text{sub}} 2\text{U(IV)Cl}_{\text{sub}} 6$ * (13)

Electronic structure of highly symmetric complexes of lanthanoids and actinoids. Pt. 4. The electronic Raman spectrum of $\text{Cs}_{\text{sub}} 2\text{NaEu(III)Cl}_{\text{sub}} 6$ * (14)

Electronic structure of CeN studied by x-ray-photoemission spectroscopy* (28)

Studies of the electronic structure of simple metals and their alloys with the aid of Compton scattering and positron annihilation using lithium, magnesium, lithium-magnesium and aluminium-zinc as examples* (48)

Interactions in inorganic molecular crystals - electronic spectra of $\text{ReF}_{\text{sub}} 6$ pure and mixed crystals. Technical report, 1 Jul--31 Dec 1975* (53)

Bonding and the electronic structure of the actinide metals* (188)

Electronic structure and properties of the actinides* (189)

Magnetic susceptibility and electronic structure in metallic and non-metallic liquid cesium-gold and cesium-antimony alloys* (190)

Electronic conductivity and polarization effects of solid $\text{ThO}_{\text{sub}} 2$ - $\text{Y}_{\text{sub}} 2\text{O}_{\text{sub}} 3$ electrolytes and of oxygen containing Va-metal electrodes* (193)

Electronic properties of actinide oxides* (398)

Assignments in the electronic spectrum of $\text{UF}_{\text{sub}} 6$ * (413)

Densities of electronic states in liquid transition metals* (422)

KKR analysis of the Anderson-McMillan theory of electronic structure of liquid metals. Technical report* (485)

Electronic and magnetic properties of rare earth- $\text{Sn}_{\text{sub}} 3$ compounds from $\text{sup } 119\text{Sn}$ Moessbauer spectroscopy* (559)

Electronic structure of highly symmetric lanthanoid and actinoid compounds. VI. The electronic Raman spectrum of $\text{Cs}_{\text{sub}} 2\text{NaCe(III)Cl}_{\text{sub}} 6$ * (678)

Electronic structure of actinides and related metals* (683)

Effects of secondary ligands on the electronic structure of uranyles* (684)

Electronic states densities from structure and thermodynamic data for liquid transition metals* (705)

electrons

Thermodiffusion measurements in liquid alloys and electron transport heat calculations in the case of almost-free electrons* (33)

Revised model for 5d-electrons in the heavy rare-earth metals* (157)

electrophysical

Thermophysical and electrophysical properties of uranium hexafluoride at temperatures of $(1--11) \times 10^{\text{sup } 3}$ degreeK and pressures of 0.1--100 atm* (302)

Eleventh

Eleventh annual seminar on theoretical physics, Johannesburg, 5-9 July 1976* (368)

TITLE INDEX

elpasolites

Vibrational spectra and force constants of the elpasolites Cs_2KMF_6 ($M=\text{Sc}, \text{Y}, \text{La}, \text{Gd}, \text{Yb}$)* (41)

elucidated

Aspects of the solution chemistry of liquid alkali metals as elucidated from electrical resistivity studies* (247)

embrittlement

Environmental sensitivity of structural metals: liquid metal embrittlement. Annual technical progress report No. 2, 25 Jun 1970--31 May 1971* (95)

Environmental sensitivity of structural metals: liquid metal embrittlement. Annual technical progress report No. 4, 1 Jun 1972--31 May 1973* (211)

Environmental sensitivity of structural metals: liquid metal embrittlement. Annual technical progress report No. 2, 25 Jun 1970--31 May 1971* (212)

EMF

Some considerations in establishing thermodynamic activities from high temperature EMF measurements* (163)

Thermodynamic properties of Cr, Mn, Co and Ni dichlorides from emf measurements on cells with solid electrolytes* (200)

emission

Vibrational fine structure in the uranium emission of Mg_3TeO_6 and Y_6WO_{12} * (59)

Raman and i.r.-emission studies of some tungstate- and molybdate-containing melts* (102)

Flame emission and atomic absorption spectrometry. III. Elements and matrices* (319)

emissivities

Spectral emissivities of molten refractory metals* (47)

emissivity

Uranium carbide: thermal diffusivity, thermal conductivity and spectral emissivity at high temperatures* (121)

Thermal diffusivity, thermal conductivity and spectral emissivity of uranium dicarbide at high temperatures* (132)

emphasis

High temperature chemistry of ceramic nuclear fuels with emphasis on nonstoichiometry* (448)

ENDOR

^{139}La ENDOR of $\text{LaCl}_3\text{:Gd}^{3+}$ * (130)

Energies

Energies of valence states of atoms in halides* (499)

Mass spectrometric determination of dissociation energies of high-temperature molecules: ScS , YS , LaS , ZrS , UO * (597)

Measurement of interfacial energies in the $\text{UO}_2\text{-Gd}_2\text{O}_3$ system* (697)

energy

Variation of the solid--solid--nickel--thoria interfacial free energy with temperature* (5)

At. Energy Rev.* (7)

Proceedings of the international conference on liquid technology in energy production* (56)

Proceedings of the international conference on liquid technology in energy production* (103)

TITLE INDEX

energy

- Composition of nonaqueous solutions of potential use in high energy density batteries. Final report, 15 Nov 1969--14 Nov 1972* (119)
- Heat capacity of PuO_2 from 12 to 350°K and of PuO_2 from 4 to 25°K. Entropy, enthalpy, and Gibbs energy of formation of PuO_2 at 298.15°K* (184)
- Second specialist meeting on sodium/fuel interaction in fast reactors, Ispra, 21-23 November 1973. Sponsored by the Committee on the Safety of Nuclear Installations, OECD Nuclear Energy Agency, Paris, France* (272)
- Partly filled shells constituting anti-bonding orbitals with higher ionization energy than their bonding counterparts* (280)
- Standard free energy of formation of liquid Na_2CrO_4 , 1100°K to 1220°K* (375)
- Diagram of excess free energy of the PbCl_2 - CsCl - ZnCl_2 system* (402)
- Energy programs: molten-salt reactor program* (423)
- At. Energy Rev.* (464)
- Energy transfer between Bi^{3+} yields Eu^{3+} , Bi^{3+} yields Sm^{3+} and UO_2^{2+} yields Eu^{3+} in oxide glasses* (539)
- Proceedings of the international conference on liquid technology in energy production* (590)
- Experimental study of the temperature dependence on energy exchanges in liquid metals* (621)
- Research on a high-energy nonaqueous battery system. Quarterly progress report No. 1, 1 Dec 1965--1 Mar 1966* (720)

Engineering

- Chemical Engineering Division physical inorganic chemistry. Annual report, July 1975--June 1976* (58)
- Chemical Engineering Division fuels and materials chemistry. Annual report, July 1975--June 1976* (281)
- Chemical Engineering Division, reactor safety and physical property studies annual report, July 1974--June 1975* (367)
- Technical experiences in an engineering-scale facility for uranium fluorination studies* (699)
- Description of an engineering-scale facility for uranium fluorination studies* (700)

enthalpies

- Preparation and some properties of actinide(III) and lanthanide(III) oxide halides, and the enthalpies of formation of PuBr_3 (s) and PuI_3 (s)* (82)
- Enthalpies of formation of melts of yttrium with aluminium* (201)
- Cesium nitrate. Standard enthalpies of solution and formation and the standard entropy at 298.15 K* (466)

enthalpy

- Standard formation enthalpy of chloride alloys of ZrCl_4 -- KCl (NaCl) systems* (8)
- High-temperature microcalorimetric measurements of partial molar enthalpy of solution, $\Delta H(\text{O}_2)$, in plutonium + oxygen* (107)
- Formation enthalpy of liquid tin-vanadium alloys* (166)
- Enthalpy of uranium-plutonium carbide from 298K to the melting point* (179)

TITLE INDEX

enthalpy

Heat capacity of PuO_2 from 12 to 350°K and of PuO_2 from 4 to 25°K. Entropy, enthalpy, and Gibbs energy of formation of PuO_2 at 298.15°K* (184)
 Thermodynamic state, specific heat, and enthalpy function of saturated UO_2 vapor between 3000 K and 5000 K* (297)
 Formation enthalpy of di- and trifluorides of samarium* (306)
 High-temperature calorimeter for determination of reaction enthalpy* (364)
 Lithium nitride (Li_3N): standard enthalpy of formation by solution calorimetry* (467)
 Improved model for solutions of nonmetals in liquid alkali metals. Calculation of enthalpy of solution and electrical resistivity* (486)
 Thermodynamics of electrolytes. 8. High-temperature properties, including enthalpy and heat capacity, with application to sodium chloride* (586)
 Density and enthalpy of liquid alkaline metals at pressures up to 1500 bar* (599)
 Formation enthalpy of gadolinium, terbium and dysprosium trifluorides* (603)
 Enthalpy of vitreous beryllium fluoride* (616)
 Enthalpy and heat capacity of liquid plutonium and uranium* (619)
 Enthalpy and heat capacity of liquid plutonium and uranium* (620)

entire

Thermal expansion of metals over the entire liquid range* (573)

Entropies

Entropies of liquid rare earth metals* (230)
 High-temperature heat contents and entropies of zirconium fluoride and zirconium sulfate* (591)

entropy

Determination of vibrational contribution into excessive mixing entropy of liquid amalgams. Alkali metal amalgams* (138)
 Heat capacity of PuO_2 from 12 to 350°K and of PuO_2 from 4 to 25°K. Entropy, enthalpy, and Gibbs energy of formation of PuO_2 at 298.15°K* (184)
 Entropy change for the lambda-type transition in nonstoichiometric UO_2 * (260)
 Origin of excess entropy of alkali metal halides solid solutions* (390)
 Cesium nitrate. Standard enthalpies of solution and formation and the standard entropy at 298.15 K* (466)
 Pressure and entropy calculations for liquid Na, K and Rb along the melting line* (584)

environment

Experimental investigation of cavitation inception and damage in a flowing sodium environment* (103)

Environmental

Environmental sensitivity of structural metals: liquid metal embrittlement. Annual technical progress report No. 2, 25 Jun 1970--31 May 1971* (95)
 Environmental sensitivity of structural metals: liquid metal embrittlement. Annual technical progress report No. 4, 1 Jun 1972--31 May 1973* (211)

TITLE INDEX

Environmental

Environmental sensitivity of structural metals: liquid metal embrittlement. Annual technical progress report No. 2, 25 Jun 1970--31 May 1971* (212)

equation

Critical assessment of equation of state data for UO_2 * (180)

Perturbation treatment of the Percus-Yevick equation and structures of liquid magnesium and cadmium* (210)

Vapour pressure measurements to 7,000 K and equation of state of oxide fuels for fast reactor safety analysis. I. Extension of vapour pressure measurements of nuclear fuels $(\text{U}, \text{Pu})\text{O}_2$ and UO_2 to 7,000 K* (477)

Some trends in constitutive equation model development for high-temperature behavior of fast-reactor structural alloys* (527)

Effective equation-of-state measurements on uranium dioxide* (537)

Equation of state and transport properties of uranium and plutonium carbides in the liquid region* (577)

Equation of state and transport properties of uranium and plutonium nitrides in the liquid region* (578)

equations

Thermodynamic similarity and universal equations of state of hexafluorides* (397)

equilibria

Phase equilibria in Li--Na--K carbonate/aluminate systems* (77)

Phase equilibria in the system of sodium chlorides, cesium chlorides and strontium chlorides* (87)

Phase equilibria in the system of sodium chlorides, cesium chlorides and strontium chlorides* (88)

Chemical effects of composition changes in irradiated oxide fuel materials. II. Fission product segregation and chemical equilibria* (169)

Quantitative study of the chemical equilibria in solution in molten LiCl-KCl (70-30 mol.%) by absorption spectrophotometry. Application to neptunium* (385)

Phase equilibria in the $\text{PbO-GeO}_2\text{-V}_2\text{O}_5$ ternary system* (394)

Equilibria and kinetics in modern ceramic processing* (557)

Phase equilibria in system U--Pu--W--C* (648)

Phase equilibria and thermodynamic data for the ternary $\text{UO}_2\text{-UO}_3\text{-REO}_{1.5}$ (Ho, Tm, Lu) systems* (694)

equilibrium

Investigation into neodymium chloride state in equilibrium with metal in eutectic melt of potassium- and lithium chlorides* (40)

Phase equilibrium in interlanthanide oxides systems* (49)

Study of the Ni-Ni (II) equilibrium exchange, by radiotracer techniques in fused LiCl-KCl eutectic* (68)

Calculation of vapor pressures of oxide fuels up to 5000 K in equilibrium and non-equilibrium evaporation* (72)

Calculation of vapor pressures of oxide fuels up to 5,000 K for equilibrium and nonequilibrium evaporation* (73)

Equilibrium between Li-K-Te alloys and LiCl-KCl molten eutectic mixture* (92)

TITLE INDEX

equilibrium

Summaries of reports of the all-union meeting on solid body chemistry. 3--5 June 1975. III. Synthesis, phase equilibrium and use of solid oxide compounds* (114)

Thermodynamic study of the U-O-C system by measurement of the carbon monoxide pressures at equilibrium* (235)

Phase equilibrium studies of lithium halide-containing electrolytes* (282)

Reduction of PuO_2 by CO and equilibrium partial pressures above plutonia* (304)

Ionic-thermocurrent study of the dipole relaxation and equilibrium in Gd-doped SrF_2 * (410)

Phase equilibrium in NdCl_3 -KCl system* (457)

Investigation of liquid-steam equilibrium in system with diluted solutions of metal fluorides in uranium hexafluoride* (526)

Summaries of reports of the all-union meeting on solid body chemistry. 3--5 June 1975. III. Synthesis, phase equilibrium and use of solid oxide compounds* (641)

Phase equilibrium study on system uranium-plutonium-tungsten-carbon* (645)

Constitution equilibrium diagram of the LiNO_3 - LiIO_3 , NaNCl_3 - NaIO_3 , KNO_3 - KIO_3 system* (665)

Equilibrium index for fission gas bubbles* (685)

Er

$\text{R}_{\text{sub } 3}\text{T}_{\text{sub } 2}$ compounds (R = rare earth or Y; T = Rh, Pd, Pt) with the rhombic Er $_{\text{sub } 3}\text{Ni}_{\text{sub } 2}$ structure type* (365)
Gd, Tb, Dy, Ho, Er solubility in chromium* (610)

erbiun

Study of lanthanum, yttrium and erbium oxychlorides in melts of alkali metal chlorides* (196)

ErCl

Investigation into absorption infrared spectra of vapors under ScBr_3 , YCl_3 , LaCl_3 , GdCl_3 , ErCl_3 , LuCl_3 by means of isolation method in rare gas matrix* (497)

ErCrO

Optical absorption Zeeman spectroscopy of rare earth orthochromites. I. ErCrO_3 * (123)

ErF

LiF-ErF_3 system* (270)

establishment

Contribution to the establishment of the phase diagram for the U--O--C system by means of x-ray diffractometry at high temperatures and under controlled pressure* (503)

Estimate

Estimate of cryoscopic calculations accuracy from fusibility diagrams* (670)

estimation

Chemical separation and spectrographic estimation of rare earths and yttrium in PuO_2 and $(\text{U,Pu})\text{O}_2$ * (125)

Eu

Ternary UO_2 -- UO_3 -- EuO system and investigation of Eu(II)--actinide(IV)--perovskites* (50)

Negative temperature coefficients of electrical resistivity: the divalent liquid metals Eu, Yb and Ba* (224)

TITLE INDEX

- Eu
Effect of temperature on the fine structure of Eu^{2+} in KCl and KI* (432)
Energy transfer between Bi^{3+} yields Eu^{3+} , Bi^{3+} yields Sm^{3+} and UO_2^{2+} yields Eu^{3+} in oxide glasses* (539)
Structure of $4f^{6+}$ and $4f^{8+}$ levels and Eu^{3+} and Tb^{3+} configurations in short-range ultra-violet* (589)
- EuCl
Thermodynamics of the lanthanide halides. II. Heat capacities and Schottky anomalies of SmCl_3 , EuCl_3 and GdCl_3 from 50 to 350 K* (594)
- EuO
Ternary UO_2 -- UO_3 -- EuO 1% 5% (EuO) system and investigation of Eu(II)--actinide(IV)--perovskites* (50)
- European
11. European congress on molecular spectroscopy. Abstracts* (44)
11. European congress on molecular spectroscopy. Abstracts* (46)
- europium
Zeeman effect and satellite carrier structure in absorption spectra of pure and doped praseodymium and europium double nitrates* (165)
Kinetics and mechanism into preparation of samarium-, europium- and ytterbium dichlorides and their some properties* (253)
Melting point of europium dichloride* (361)
Heat of formation of europium sesquioxide and europium trichloride* (607)
- eut
Comparison of the tendencies of complex formation of various metal ions in fused LiF-NaF-KF (eut.) and in fused LiCl-KCl (eut.)* (314)
Preparation of thorium metal by electrolysis of fused salt ThCl_4 - LiCl + NaCl + KCl (eut) system* (703)
- eutectic
Investigation into neodymium chloride state in equilibrium with metal in eutectic melt of potassium- and lithium chlorides* (40)
Electrode reactions in molten sodium nitrate--barium nitrate eutectic at 350 degrees C* (42)
Study of the Ni-Ni (II) equilibrium exchange, by radiotracer techniques in fused LiCl-KCl eutectic* (68)
Interaction of the lithium-tin alloys with the LiCl-KCl-CsCl melted eutectic mixture* (91)
Equilibrium between Li-K-Te alloys and LiCl-KCl molten eutectic mixture* (92)
Atomic structure of melts of complex system with eutectic transformations* (155)
Measurement of fluorescence of rare earths in molten salts (Nd^{3+} in NaNO_3 - KNO_3 eutectic)* (233)
Structure and anodic discharge behavior of lithium-boron alloys in the LiCl-KCl eutectic melt.* (274)
EUTECTIC DATA: safety, hazards, corrosion, melting points, compositions, and bibliography* (276)
Chronopotentiometry of U(III) and U(IV) in molten LiCl-KCl eutectic* (292)
Electrochemical behavior of U(III) and U(IV) in molten LiCl-KCl eutectic* (293)

TITLE INDEX

eutectic

- Investigation into americium extraction process from residues of irradiated nuclear fuel after its fluorination by $\text{SiF}_4\text{-NaF}$ and $\text{SiF}_4\text{-CaF}_2$ melts of eutectic composition* (308)
- Electrode potentials of tantalum chlorides in a molten NaCl-KCl eutectic* (449)
- Intensity-voltage curves of hydrogen diffusion electrodes in eutectic LiCl-KCl * (507)
- Electrochemical reduction of molybdenum oxychloride in molten lithium chloride - potassium chloride eutectic* (515)
- Heat transfer measurements in a forced convection loop with two molten-fluoride salts: LiF-BiF_3 , $\text{ThF}_4\text{-UF}_6$ and eutectic $\text{NaBF}_4\text{-NaF}$ * (585)
- Anodic behavior of cadmium and cadmium-zinc alloys in eutectic melt of potassium- and lithium chlorides* (675)
- Eutectic point of gallium-indium system* (706)

eutectics

- Some thermodynamic properties of dilute solutions of actinide chlorides in $(\text{Li-K})\text{Cl}$ and in $(\text{Na-K})\text{Cl}$ eutectics* (406)
- Thermodynamic properties of dilute solutions of ThCl_4 in $(\text{Li-K})\text{Cl}$ and (Na-K) eutectics* (407)
- Calculated method of eutectics of multicomponent systems on base of algorithm and digital computers* (517)

EuZrF

- New hexafluorozirconates(IV): BaZrF_6 , PbZrF_6 , EuZrF_6 , SrZrF_6 * (419)

evaluate

- Results of research to evaluate solid plutonium nitrate as a safe shipping form* (85)

evaporation

- Calculation of vapor pressures of oxide fuels up to 5000 K in equilibrium and non-equilibrium evaporation* (72)
- Calculation of vapor pressures of oxide fuels up to 5,000 K for equilibrium and nonequilibrium evaporation* (73)
- Partial pressures, activities and evaporation heat in the Kf-ZrF_4 system* (295)
- Evaporation behaviour of the ternary uranium plutonium carbides* (479)
- Evaporation behaviour of the ternary uranium plutonium carbides* (480)
- Evaporation behaviour of the ternary uranium plutonium carbides* (481)
- Evaporation behavior and high-temperature thermal analysis of substoichiometric plutonium oxide in the composition range from $\text{O/Pu} = 1.51$ to 2.00 * (482)

excess

- Origin of excess entropy of alkali metals halides solid solutions* (390)
- Diagram of excess free energy of the $\text{PbCl}_2\text{-CsCl-ZnCl}_2$ system* (402)
- Reaction of potassium superoxide in excess potassium* (590)

excessive

- Determination of vibrational contribution into excessive mixing entropy of liquid amalgams. Alkali metal amalgams* (138)

TITLE INDEX

exchanges

Experimental study of the temperature dependence on energy exchanges in liquid metals* (621)

excitation

Effect of luminescence centers on optical absorption, excitation and fluorescence spectra of CsCl:Pb* (491)

excitations

Structure and excitations of amorphous solids* (183)

Exciton

Vibrational spectra of transition metal hexafluoride crystals. III. Exciton band structures of MoF₆, WF₆, and UF₆. Technical report* (52)

expansion

Thermal expansion of ZrO_2 * (15)

Thermal expansion of molten materials by the gamma attenuation technique* (152)

High temperature thermal expansion of ThO₂, MgO and Y₂O₃ by X-ray diffraction* (242)

Thermal expansion of metals over the entire liquid range* (573)

expansions

Hydrodynamics of post-disassembly fuel expansions* (310)

Experience

Experience of the Central Control Laboratory (CCL) in accounting for and controlling nuclear material in Czechoslovakia* (349)

experiences

Technical experiences in an engineering-scale facility for uranium fluorination studies* (699)

explosion

Analysis of LMFBR explosion model experiments by means of the SUREDOM-II code* (601)

exposed

Atomic diffusion in alloys exposed to liquid sodium* (181)

Extension

Extension of vapour pressure measurements of nuclear fuels (U, Pu)O₂ and UO₂ to 7000 K for fast reactor safety analysis* (475)

Vapour pressure measurements to 7,000 K and equation of state of oxide fuels for fast reactor safety analysis. I. Extension of vapour pressure measurements of nuclear fuels (U, Pu)O₂ and UO₂ to 7,000 K* (477)

extraction

Investigation into americium extraction process from residues of irradiated nuclear fuel after its fluorination by SiF₄-NaF and SiF₄-CaF₂ melts of eutectic composition* (308)

Investigations on the solvent extraction from molten salts by tracer technique* (369)

Studies on some binary solvent systems for the extraction of gadolinium* (370)

Investigations of the solvent extraction from molten salts with the use of tracer technique* (371)

Tetrad effect in the extraction of the lanthanides with neutral oxygen-containing compounds* (426)

TITLE INDEX

F

Optical spectra of the difluoride, dichloride, and trichloride ions in the matrix-isolated $M^{\text{sup}} + F^{\text{sub}}_2^{\text{sup}} -$, $M^{\text{sup}} + Cl^{\text{sub}}_2^{\text{sup}} -$, and $M^{\text{sup}} + Cl^{\text{sub}}_3^{\text{sup}} -$ species* (16)

Study of $KBe^{\text{sub}}_2F^{\text{sub}}_5 - DyF^{\text{sub}}_3$ cross-section of $KF - BeF^{\text{sub}}_2 - DyF^{\text{sub}}_3$ ternary system and some properties of this cross section glasses* (21)

Study of the compounds $M^{\text{sub}}_3UO^{\text{sub}}_2F^{\text{sub}}_5$ ($M = K, Rb, Cs, NH^{\text{sub}}_4$) by i.r. absorption and Raman diffusion spectrophotometry. III. Effect of cation on vibration frequency of the ion $UO^{\text{sub}}_2F^{\text{sub}}_5^{\text{sup}}_3 -$ * (127)

Calculation and discussion of state diagrams in the system $Pu - F - O - H$ * (222)

Fusibility in $K^{\text{sub}}_2Nb(Ta)F^{\text{sub}}_7 - LiF - NaF$ systems* (298)

Anomalous melting of f electron metals (with attention to Pu)* (318)

Intensities of lanthanide f-f transitions* (492)

Behavior of ruthenium in fluoride-volatility process. II. Fluorination of RuO^{sub}_2 by F^{sub}_2 * (558)

Covalent compounds of uranium of the type $F^{\text{sub}}_x / U(O^{\text{sub}}_2F^{\text{sub}}_5)^{\text{sub}}_{6-x}$ * (571)

Phases of variable composition with LaF^{sub}_3 structure in $MF^{\text{sub}}_2 - (Y, Ln)F^{\text{sub}}_3$ systems. 1. $SrF^{\text{sub}}_2 - (Y, Ln)F^{\text{sub}}_3$ systems (thermal characteristics, formation of berthollides)* (593)

Preparation and structure of transition metal fluorides $UE^{\text{sup}}(II)M^{\text{sup}}(IV)F^{\text{sub}}_6 [UE^{\text{sup}}(II) : Cr - Zn; M^{\text{sup}}(IV) : Hf, Zr]$ * (596)

Fabrication

Fabrication and high temperature thermodynamic and transport properties of FuN * (9)

FuO^{sub}_2 dissolution problem for LWR plutonium recycle and LMFBR fuels: fabrication and reprocessing problems and their resolution* (455)

Influence of the technology of the alloy fabrication on its properties in a liquid state* (643)

Fabrication of uranium-plutonium mixed carbide* (691)

facility

Technical experiences in an engineering-scale facility for uranium fluorination studies* (699)

Description of an engineering-scale facility for uranium fluorination studies* (700)

Fault

Fault structures in rare earth--cobalt intermetallics* (11)

fcc

Electromodulation spectroscopy of sc and fcc phase $TlCl$ and $TlBr$ * (411)

Fe

Using molecular dynamics method for the evaluation of Fe-Ni and Al-Zn melts structure properties* (512)

Compounds with the structure of $ThMn^{\text{sub}}_{12}$ type in the R--Fe--Al systems* (671)

Investigation of the solubility and diffusion of Fe atoms in Cu at high temperature using molten salt electrochemistry* (693)

TITLE INDEX

February

Stable inorganic matrix materials for high temperature batteries. Final report No. 1, 2 June 1967--2 February 1968* (23)
 Molten-salt reactor program. Semiannual progress report for period ending February 29, 1976* (176)
 Molten-salt reactor program. Semiannual progress report for period ending February 29, 1976* (237)
 Electrochemical characterization of nonaqueous systems for secondary battery application. Quarterly report, February--April 1968* (380)
 Collection of gamma spectra data of irradiated light water moderated reactor spent fuel and study of the applicability of the methods for fuel identification. Final report for the period 1 February 1972--October 1975* (453)

FeCl

Investigation into thermodynamic properties of $\text{FeCl}_{\text{sub } 2}\text{-CsCl}$ system melts* (96)
 Phase diagrams of $\text{BiCl}_{\text{sub } 3}\text{-CuCl-TeCl}_{\text{sub } 4}$ and $\text{CuCl-FeCl}_{\text{sub } 3}\text{-TeCl}_{\text{sub } 4}$ systems* (337)
 $\text{FeCl}_{\text{sub } 3}\text{-MeCl}$ systems* (435)
 $\text{FeCl}_{\text{sub } 3}\text{-MoCl}_{\text{sub } 5}\text{-NbCl}_{\text{sub } 5}\text{-WCl}_{\text{sub } 6}$ system* (554)

ferric

A Mcesstauer spectroscopic study of intercalation compounds of graphite with ferric chloride and aluminum chloride* (627)

ferrous

Effect of basicity on ferrous oxide activity in complex oxide systems* (372)

Festkoerperphysik

Progress report of the Teilinstitut Nukleare Festkoerperphysik. (1.6.1974--31.5.1975)* (567)

Fesub

Thermodynamic properties of NbO , $\text{Fesub}(0,95)\text{O}$, CoO , NiO , $\text{Cu}_{\text{sub } 2}\text{O}$ and CuO oxides as comparative electrodes in electromotive forces method* (658)

fibers

High density thoria-silica-metal (III) oxide fibers* (710)

filled

Partly filled shells constituting anti-bonding orbitals with higher ionization energy than their bonding counterparts* (280)

Final

Stable inorganic matrix materials for high temperature batteries. Final report No. 1, 2 June 1967--2 February 1968*. (23)
 Lithium--silver chloride secondary battery investigation. Final report, December 1963--November 1964* (118)
 Composition of nonaqueous solutions of potential use in high energy density batteries. Final report, 15 Nov 1969--14 Nov 1972* (119)
 Fine definition of ir spectra from high temperature interactions of $\text{U} + \text{O}_{\text{sub } 2}$. Part II. Final report, 1 July 1974--30 June 1975* (218)
 Thermochemistry of gaseous metal oxides. Final report, 1 Apr 1973--31 Dec 1975* (239)
 Radial oxygen distribution in the fuel of irradiated mixed oxide fuel elements with different final stoichiometries* (316)

TITLE INDEX

Final

Collection of gamma spectra data of irradiated light water moderated reactor spent fuel and study of the applicability of the methods for fuel identification. Final report for the period 1 February 1972--October 1975* (453)

Space Processing Applications Rocket project, SPAR 1. Final report* (536)

Basic mechanisms providing oxidation resistance in structural metals at high temperatures. II. The defect structure of nonstoichiometric rutile (TiO_{2-x}) and nonstoichiometric rutile containing 1 to 10 mole % Nb_2O_5 . Final report 21 Dec 1973-20 Dec 1974* (542)

Diffusion in liquid metal systems. Final report* (651)

Optical spectroscopy of ^{237}Np in CaF_2 . Final report* (718)

find

Zirconia and yttria--thoria ceramics find new uses as solid electrolytes* (595)

fine

Vibrational fine structure in the uranium emission of MgTeO_6 and Y_6WO_{12} * (59)

Fine definition of ir spectra from high temperature interactions of $\text{U} + \text{O}_2$. Part II. Final report, 1 July 1974--30 June 1975* (218)

Effect of temperature on the fine structure of Eu^{2+} in KCl and KI * (432)

fissile

Accurate procedure to safeguard the fissile material content of input and output solutions of reprocessing plants* (131)

fission

Investigations on radioactive fission product correlations: gamma spectrometry measurements on spent fuel assemblies discharged from the Trino Vercellese reactor at the end of the 2nd irradiation cycle* (75)

Chemical effects of composition changes in irradiated oxide fuel materials. II. Fission product segregation and chemical equilibria* (169)

Determination of the burn up and differentiation of the fission yields of uranium-thorium-fuels* (214)

Use of fluoride melts for separation of some fission products and americium from mixtures of different fluorides being the simulators of residues from fast reactor fuel fluorination* (307)

Microprobe study of fission product behavior in high-burnup HTR fuels* (315)

Fission rates, burnup, and neutron flux-fluence-spectra characterization for mixed oxide fuel experiments in the EBR-II* (416)

Separation of alkaline earth from alkali metal fission products in the SOLAR on-line mass spectrometer* (534)

Equilibrium index for fission gas bubbles* (685)

Standard method of test for atom percent fission in uranium and plutonium fuel (mass spectrometric method)* (715)

Standard method of test for atom percent fission in uranium and plutonium fuel (neodymium-148 method)* (716)

TITLE INDEX

fissionable

Analytical methods for fissionable materials in the nuclear fuel cycle. Progress report, July 1, 1975--September 30, 1976* (690)

Flame

Flame emission and atomic absorption spectrometry. III. Elements and matrices* (319)

flow

Nonisothermal flow of liquid metals over the surface of solid ones* (101)

flowing

Experimental investigation of cavitation inception and damage in a flowing sodium environment* (103)

fluence

Fission rates, burnup, and neutron flux-fluence-spectra characterization for mixed oxide fuel experiments in the FRR-II* (416)

fluid

Fluorination process studies of plutonium dioxide by fluid-bed* (725)

fluidized

Adaptation of the continuous cold trap system of fluidized-bed to the fluoride volatility process* (724)

fluorescence

Measurement of fluorescence of rare earths in molten salts (Nd³⁺ in NaNO₃-KNO₃ eutectic)* (233)

Effect of luminescence centers on optical absorption, excitation and fluorescence spectra of CsCl:Pb* (491)

Determination of trace impurities of titanium and zirconium in molybdenum and tungsten and their oxygen compounds by X-ray fluorescence spectrometry* (698)

fluoride

Some properties of hydrogen fluoride compounds with group 5 metal pentafluorides* (170)

Raman spectra of Al₂O₃ solutions in molten cryolite and other aluminum fluoride containing melts* (203)

Use of fluoride melts for separation of some fission products and americium from mixtures of different fluorides being the simulators of residues from fast reactor fuel fluorination* (307)

Study of the behavior of high activity waste produced during the regeneration of fast reactor fuel elements by the gaseous fluoride method* (311)

Pulse NMR in solids: chemical shift, lead fluoride, and thorium hydride* (363)

Lithium--nickel fluoride secondary battery investigation. Quarterly technical progress report No. 1, 27 Apr--27 Jul 1966* (381)

Effect of fluoride ions on the kinetics of the oxidation of uranium(IV) by cerium(IV)* (424)

Electric conductivity of melts of the LiF-misch metal fluoride system* (461)

NMR study on structural microinhomogeneity of fluoride-oxide melts* (463)

Phase relations in the alkaline earth fluoride-yttria systems (in three parts). 1. The CaF₂-Y₂O₃ phase diagram* (468)

TITLE INDEX

fluoride

- Phase relations in the alkaline earth fluoride-yttria systems (in three parts). 2. The $\text{BaF}_2\text{--Y}_2\text{O}_3$ phase diagram* (469)
- New fluoride glasses* (521)
- Behavior of ruthenium in fluoride-volatility process. II.
- Fluorination of RuO_2 by F_2 * (558)
- Heat transfer measurements in a forced convection loop with two molten-fluoride salts: $\text{LiF--BeF}_2\text{--ThF}_2\text{--UF}_4$ and eutectic NaF--NaF * (585)
- High-temperature heat contents and entropies of zirconium fluoride and zirconium sulfate* (591)
- Enthalpy of vitreous beryllium fluoride* (616)
- Some thermal properties of beryllium fluoride from 8⁰ K to 1,200⁰ K* (617)
- Molten salt chemistry : Part IV - Solubility behaviour of PuF_3 in fluoride salts of interest in molten salt reactor technology* (655)
- Physical properties of thorium fluoride* (656)
- Adaptation of the continuous cold trap system of fluidized-bed to the fluoride volatility process* (724)

fluorides

- Quaternary mutual system from fluorides and chlorides of lithium, sodium and strontium* (89)
- Use of fluoride melts for separation of some fission products and americium from mixtures of different fluorides being the simulators of residues from fast reactor fuel fluorination* (307)
- Interaction of rubidium and cesium fluorides with lutetium trifluorides* (444)
- Reactivity of transition metal fluorides. IX. Protactinium pentafluoride and tetrafluoride* (465)
- Phase diagrams in the pseudobinary systems of Li_2BeF_4 , alkali fluoroborates and alkali fluorides* (473)
- Investigation of liquid-steam equilibrium in system with diluted solutions of metal fluorides in uranium hexafluoride* (526)
- Interaction of potassium tetrafluoroborate with fluorides of potassium, lanthanum, neodymium in melt* (530)
- Investigation into reaction of rubidium and holmium fluorides* (540)
- Preparation and structure of transition metal fluorides $\text{U}^{\text{II}}\text{M}^{\text{IV}}\text{F}_6$ [U^{II} : Cr-Zn; M^{IV} : Hf, Zr]* (596)

fluorination

- Use of fluoride melts for separation of some fission products and americium from mixtures of different fluorides being the simulators of residues from fast reactor fuel fluorination* (307)
- Investigation into americium extraction process from residues of irradiated nuclear fuel after its fluorination by $\text{SiF}_4\text{--NaF}$ and $\text{SiF}_4\text{--CaF}_2$ melts of eutectic composition* (308)
- Fundamental studies of metal fluorination reactions. Progress report, May 1, 1976--April 31, 1977* (533)
- Behavior of ruthenium in fluoride-volatility process. II.
- Fluorination of RuO_2 by F_2 * (558)
- Technical experiences in an engineering-scale facility for uranium fluorination studies* (699)

TITLE INDEX

fluorination

Description of an engineering-scale facility for uranium fluorination studies* (700)
 Fluorination process studies of plutonium dioxide by fluid-bed* (725)

fluorine

Interaction of lanthanide ions with fluorine-ions* (329)
 Progress in fluorine calorimetry* (500)

fluoroborates

Phase diagrams in the pseudobinary systems of Li_2BeF_4 , alkali fluoroborates and alkali fluorides* (473)

flux

Fission rates, burnup, and neutron flux-fluence-spectra characterization for mixed oxide fuel experiments in the EBR-II* (416)

force

Vibrational spectra and force constants of the elpasolites $\text{Cs}_2\text{KM}_6\text{F}_{24}$ (M=Sc, Y, La, Gd, Yb)* (41)
 Report of task force on alternate structural materials for Liquid Metal Fast Breeder Reactors* (309)
 Electromotive force measurements in molten lithium--magnesium alloys* (551)

forced

Heat transfer measurements in a forced convection loop with two molten-fluoride salts: $\text{LiF--BeF}_2\text{--ThF}_4\text{--UF}_4$ and eutectic $\text{NaBF}_4\text{--NaF}$ * (585)

forces

Thermodynamic properties of NbO , $\text{Fe}_{0.95}\text{O}$, CoO , NiO , Cu_2O and CuO oxides as comparative electrodes in electromotive forces method* (658)

forging

Metallurgical characterization of prototypic tubesheet forging* (471)

form

Results of research to evaluate solid plutonium nitrate as a safe shipping form* (85)

formation

Standard formation enthalpy of chloride alloys of $\text{ZrCl}_4\text{--KCl(NaCl)}$ systems* (8)
 Preparation and some properties of actinide(III) and lanthanide(III) oxide halides, and the enthalpies of formation of $\text{PuBr}_3(\text{s})$ and $\text{PuI}_3(\text{s})$ * (82)
 Experimental determination of formation heat of gaseous Ga_2O and In_2O * (105)
 Thermodynamics and kinetics of the formation of rare earth intermetallics* (137)
 Formation of actinides in irradiated HTGR fuel elements* (150)
 Formation enthalpy of liquid tin-vanadium alloys* (166)
 Heat capacity of $^{242}\text{PuO}_2$ from 12 to 350°K and of $^{244}\text{PuO}_2$ from 4 to 25°K. Entropy, enthalpy, and Gibbs energy of formation of PuO_2 at 298.15°K* (184)
 Enthalpies of formation of melts of yttrium with aluminium* (201)
 Criteria for solid solution formation in NFE alloys* (205)
 Formation enthalpy of di- and trifluorides of samarium* (306)

TITLE INDEX

formation

- Comparison of the tendencies of complex formation of various metal ions in fused LiF-NaF-KF (eut.) and in fused LiCl-KCl (eut.)* (314)
- Investigation of formation thermodynamic of rare earth aluminates with perovskite structure* (326)
- Study of kinetics of monovalent indium ions formation in the $\text{In}^0\text{-InCl}_3\text{-LiCl-H}_2\text{O}$ system* (341)
- Standard free energy of formation of liquid Na_2CrO_4 , 1100°K to 1220°K* (375)
- Heat of formation of solid and liquid alloys of transition metals* (425)
- Cesium nitrate. Standard enthalpies of solution and formation and the standard entropy at 298.15 K* (466)
- Lithium nitride (Li_3N): standard enthalpy of formation by solution calorimetry* (467)
- Heats of formation of solid solutions of CsCl with CsBr * (543)
- Chemistry of elements niobium and tantalum. 87. Studies on the thermal formation and decomposition of the complexes $[\text{Nb}_6\text{Cl}_{12}]^{2+}$, $[\text{Nb}_6\text{I}_8]^{3+}$, and $[\text{Ta}_6\text{X}_{12}]^{(n+)}$ ($\text{X}=\text{Cl}, \text{Br}, \text{I}$)* (561)
- Phases of variable composition with LaF_3 structure in $\text{MF}_2\text{-(Y, Ln)F}_3$ systems. 1. $\text{SrF}_2\text{-(Y, Ln)F}_3$ systems (thermal characteristics, formation of berthollides)* (593)
- Formation enthalpy of gadolinium, terbium and dysprosium trifluorides* (603)
- Heat of formation of europium sesquioxide and europium trichloride* (607)
- Heats of formation of holmium and terbium trichlorides* (608)
- Thermographic determination of formation heats of lower zirconium chlorides and bromides* (635)
- Glass formation in nitrate systems containing rare earth ions* (666)
- Effect of second sphere cations on formation of nickel (2) chloride complexes in molten salts* (673)
- Investigation into properties of liquidus curves of binary salt system melts where chemical compounds formation. 2. Systems of mono-monovalent and bi-bivalent cations with common anion* (708)

formed

- Discrete variation of the structure and texture of liquid metals formed by components with discrete packing* (177)

France

- Reprocessing of fast breeder fuels in France* (67)
- Second specialist meeting on sodium/fuel interaction in fast reactors, Ispra, 21-23 November 1973. Sponsored by the Committee on the Safety of Nuclear Installations, OECD Nuclear Energy Agency, Paris, France* (272)

frequency

- Study of the compounds $\text{M}_3\text{UO}_2\text{F}_5$ ($\text{M} = \text{K}, \text{Rb}, \text{Cs}, \text{NH}_4$) by i.r. absorption and Raman diffusion spectrophotometry. III. Effect of cation on vibration frequency of the ion $\text{UO}_2\text{F}_5^{3-}$ * (127)
- Frequency spectrum and scattering law for liquid copper. A molecular dynamics investigation* (567)

TITLE INDEX

Friction

Friction in metal melts* (575)

fuel

Investigation of fuel vapor pressure for fast breeder safety analysis* (60)

Molten carbonate fuel cell research at ORNL* (71)

Investigations on radioactive fission product correlations: gamma spectrometry measurements on spent fuel assemblies discharged from the Trino Vercellese reactor at the end of the 2nd irradiation cycle* (75)

Third international symposium on fuel cells* (77)

LMFBR Fuel Reprocessing Program progress report for period April 1 to June 30, 1976* (94)

Basic problems of fast reactor fuel reprocessing* (135)

Formation of actinides in irradiated HTGR fuel elements* (150)

Chemical effects of composition changes in irradiated oxide fuel materials. II. Fission product segregation and chemical equilibria* (169)

Cs--U--O phase diagram and its application to uranium--plutonium oxide fast reactor fuel pins* (175)

Behavior of molybdenum in mixed-oxide fuel* (202)

Uranium dioxide as a ceramic nuclear fuel* (226)

Fuel processing for molten-salt reactors* (237)

Second specialist meeting on sodium/fuel interaction in fast reactors, Ispra, 21-23 November 1973. Sponsored by the Committee on the Safety of Nuclear Installations, OECD Nuclear Energy Agency, Paris, France* (272)

Use of fluoride melts for separation of some fission products and americium from mixtures of different fluorides being the simulators of residues from fast reactor fuel fluorination* (307)

Investigation into americium extraction process from residues of irradiated nuclear fuel after its fluorination by SiF-NaF and SiF-CaF_2 melts of eutectic composition* (308)

Hydrodynamics of post-disassembly fuel expansions* (310)

Study of the behavior of high activity waste produced during the regeneration of fast reactor fuel elements by the gaseous fluoride method* (311)

Radial oxygen distribution in the fuel of irradiated mixed oxide fuel elements with different final stoichiometries* (316)

Twenty years of research in the processing of radioactive fuel and neutralization of radioactive waste at the Nuclear Research Institute (NRI)* (358)

Thermophysical properties of UO_2 fuel materials* (366)

MATERO: Version 09. A handbook of materials properties for use in the analysis of light water reactor fuel rod behavior* (387)

Fission rates, burnup, and neutron flux-fluence-spectra characterization for mixed oxide fuel experiments in the EBR-II* (416)

Collection of gamma spectra data of irradiated light water moderated reactor spent fuel and study of the applicability of the methods for fuel identification. Final report for the period 1 February 1972--October 1975* (453)

Preliminary assessments of carbide fuel pins during mild overpower transients in LMFBRs* (456)

TITLE INDEX

fuel

Study of the retention of Ba in UO_2 nuclear fuel particles by ZrO_2 as getter* (470)
 Reactor conference, Duesseldorf, 30.3.-2.4.1976. Section 2: Fuel elements, fuel cycle and isotope technique* (480)
 Phase analysis on simulated LWR fuel rods and its evaluation with the aid of constitution studies in the system uranium-zirconium-oxygen* (509)
 Phase analysis at simulated LWR fuel rods and their interpretation by constitution investigations in the uranium-zirconium-oxygen system* (510)
 Thermal properties of nuclear fuel ceramics* (613)
 Analytical methods for fissionable materials in the nuclear fuel cycle. Progress report, July 1, 1975--September 30, 1976* (690)
 Standard method of test for atom percent fission in uranium and plutonium fuel (mass spectrometric method)* (715)
 Standard method of test for atom percent fission in uranium and plutonium fuel (neodymium-148 method)* (716)
 Fuel-clad interaction. C-E thermo-structural fuel evaluation methods* (719)

fuels

Reprocessing of fast breeder fuels in France* (67)
 Review of partitioning proposals for spent nuclear fuels* (69)
 Calculation of vapor pressures of oxide fuels up to 5000 K in equilibrium and non-equilibrium evaporation* (72)
 Calculation of vapor pressures of oxide fuels up to 5,000 K for equilibrium and nonequilibrium evaporation* (73)
 Determination of the burn up and differentiation of the fission yields of uranium-thorium-fuels* (214)
 Chemical Engineering Division fuels and materials chemistry. Annual report, July 1975--June 1976* (281)
 Microprobe study of fission product behavior in high-burnup HTR fuels* (315)
 Chemical reprocessing of spent nuclear fuels* (322)
 Preparation and thermochemical stability of uranium-zirconium-carbonitrides. A contribution to nuclear fuels for advanced high-temperature reactors* (333)
 High temperature chemistry of ceramic nuclear fuels with emphasis on nonstoichiometry* (448)
 PuO_2 dissolution problem for LWR plutonium recycle and LMFBR fuels: fabrication and reprocessing problems and their resolution* (455)
 Extension of vapour pressure measurements of nuclear fuels (U, Pu) O_2 and UO_2 to 7000 K for fast reactor safety analysis* (475)
 Vapour pressure measurements to 7,000 K and equation of state of oxide fuels for fast reactor safety analysis. I. Extension of vapour pressure measurements of nuclear fuels (U, Pu) O_2 and UO_2 to 7,000 K* (477)
 Phase relations, crystal structures and physical properties of nuclear fuels* (612)
 Spectrophotometric determination of uranium in spent fuels of WWR-2 reactors* (632)

function

Calculation of the reduced partition function ratio for uranium hexafluoride* (285)

TITLE INDEX

function

Thermodynamic state, specific heat, and enthalpy function of saturated UO_2 vapor between 3000 K and 5000 K* (297)
 Theory for the derivation of a single hard-sphere potential describing the overall molecular interactions in binary liquid metal alloys as a function of temperature and composition* (525)
 Work function of gallium* (583)

functions

Thermodynamic functions and vapor pressures of uranium and plutonium oxides at high temperatures* (220)
 Spectral density functions for disordered systems* (229)

Fundamental

Fundamental studies of metal fluorination reactions. Progress report, May 1, 1976--April 31, 1977* (533)

furnace

Simple Raman cell and furnace usable at temperatures higher than 1000°C for corrosive melts* (204)

fused

Study of the Ni-Ni (II) equilibrium exchange, by radiotracer techniques in fused LiCl-KCl eutectic* (68)
 Comparison of the tendencies of complex formation of various metal ions in fused LiF-NaF-KF (eut.) and in fused LiCl-KCl (eut.)* (314)
 Determination of the Np(IV)--Np(III) normal potential in fused LiCl-CsCl (55-45%) by absorption spectrophotometry* (386)
 Raman spectra of solid ThCl_4 and fused ThCl_4 --KCl system* (488)
 Preparation of thorium metal by electrolysis of fused salt ThCl_4 --LiCl+NaCl+KCl(eut) system* (703)

Fusibility

Fusibility in $\text{K}_2\text{Nb}(\text{Ta})\text{F}_7$ --LiF-NaF systems* (298)
 Estimate of cryoscopic calculations accuracy from fusibility diagrams* (670)

fusion

Radiation effects and tritium technology for fusion reactors. Vol. IV* (529)
 Thermodynamic properties of solutions of hydrogen isotopes in metals and alloys of interest to fusion reactor technology* (663)

Ga

Thermal analysis of GaCl_3 +3In = InCl+Ga system* (4)
 Experimental determination of formation heat of gaseous Ga_2O and In_2O_3 * (105)
 Adhesion properties of melted Al, Ga, In on solid titanium* (327)
 Ga-Pi-Te system* (544)

GaCl

Thermal analysis of GaCl_3 +3In = InCl+Ga system* (4)
 Systems of NbCl_5 -- GaCl_3 and TaCl_5 -- GaCl_3 * (633)

gadolinium

Studies on some binary solvent systems for the extraction of gadolinium* (370)
 Formation enthalpy of gadolinium, terbium and dysprosium trifluorides* (603)

gallium

Thermodynamic studies of gallium--indium liquid alloys by solid state electrochemistry with oxide electrolytes* (513)

TITLE INDEX

gallium

Thermodynamic properties of liquid thorium-gallium alloys* (522)
 Work function of gallium* (583)
 Viscosities of cesium vapor to 1,620 K and of liquid gallium to 1,800 K* (622)
 Spectrophotometric determination of rare earths and yttrium in gallium base alloys* (623)
 Some structure differences of double oxides of indium and gallium with antimony(5)* (657)
 Viscosities of cesium vapor to 1620 K and of liquid gallium to 1800 K* (681)
 Eutectic point of gallium-indium system* (706)

gamma

Investigations on radioactive fission product correlations: gamma spectrometry measurements on spent fuel assemblies discharged from the Trino Vercellese reactor at the end of the 2nd irradiation cycle* (75)
 Thermal expansion of molten materials by the gamma attenuation technique* (152)
 Collection of gamma spectra data of irradiated light water moderated reactor spent fuel and study of the applicability of the methods for fuel identification. Final report for the period 1 February 1972--October 1975* (453)

gas

Prediction of the inert gas solubilities in stoichiometric molten UO_2 * (225)
 Radiolytic gas generation in plutonium contaminated waste materials* (301)
 Investigation into absorption infrared spectra of vapors under ScBr_3 , YCl_3 , LaCl_3 , GdCl_3 , ErCl_3 , LuCl_3 by means of isolation method in rare gas matrix* (497)
 Equilibrium index for fission gas bubbles* (685)

gaseous

Isotope effect on the thermodynamic quantities of gaseous uranium hexafluoride* (78)
 Experimental determination of formation heat of gaseous Ga_2O and In_2O_3 * (105)
 Thermochemistry of gaseous UF_5 and UF_4 * (238)
 Thermochemistry of gaseous metal oxides. Final report, 1 Apr 1973--31 Dec 1975* (239)
 Thermochemistry of gaseous compounds of metals. Annual summary report 5 Dec 74--4 Dec 75* (240)
 Thermochemistry of gaseous compounds of metals. Annual summary report, 5 Dec 1975--4 Dec 1976* (241)
 Study of the behavior of high activity waste produced during the regeneration of fast reactor fuel elements by the gaseous fluoride method* (311)

Gases

Gases and carbon in metals (thermodynamics, kinetics, and properties). I. Alkali metals, alkaline earth metals, light metals (Li, Na, K, Rb, Cs; Ca, Sr, Ba; Be, Mg, Al)* (192)

GaTe

GaTe-Bi₂Te₃ system* (545)

TITLE INDEX

Gd

Vibrational spectra and force constants of the elpasolites Cs_2KMF_6 ($\text{M}=\text{Sc}, \text{Y}, \text{La}, \text{Gd}, \text{Yb}$)* (41)
 ^{139}La ENDOR of $\text{LaCl}_3\cdot\text{Gd}_3$ * (130)
 Influence of small additions to the preparation and the properties of UO_2 ceramics. Part A. $\text{UO}_2/\text{Gd}_2\text{O}_3$ * (266)
 Icnic-thermocurrent study of the dipole relaxation and equilibrium in Gd-doped SrF_2 * (410)
 XPS valence bands of La, Ce and Gd and their aluminium alloys* (598)
 Gd, Tb, Dy, Ho, Er solubility in chromium* (610)
 Measurement of interfacial energies in the UO_2 - Gd_2O_3 system* (697)

GdCl

Investigation into absorption infrared spectra of vapors under ScBr_3 , YCl_3 , LaCl_3 , GdCl_3 , ErCl_3 , LuCl_3 by means of isolation method in rare gas matrix* (497)
 Thermodynamics of the lanthanide halides. II. Heat capacities and Schottky anomalies of SmCl_3 , EuCl_3 and GdCl_3 from 50 to 350 K* (594)

GeCl

BiCl_3 - GeCl_4 - TeCl_4 system* (553)

generation

Radiolytic gas generation in plutonium contaminated waste materials* (301)

GeO

Phase equilibria in the PbO - GeO_2 - V_2O_5 ternary system* (394)

getter

Study of the retention of Ba in UO_2 nuclear fuel particles by ZrO_2 as getter* (470)

Gibbs

Heat capacity of $^{242}\text{PuO}_2$ from 12 to 350°K and of $^{244}\text{PuO}_2$ from 4 to 25°K. Entropy, enthalpy, and Gibbs energy of formation of PuO_2 at 298.15°K* (184)

GIREDMET

Proceedings of GIREDMET* (136)

GIREDMETA

Nauchnye trudy GIREDMETA. Vol. 52* (115)

Glass

Glass formation in nitrate systems containing rare earth ions* (666)

glasses

Study of KBef_5 - DyF_3 cross-section of KF - BeF_2 - DyF_3 ternary system and some properties of this cross section glasses* (21)
 Anomalous quasielastic light scattering from metal oxide glasses* (183)
 New fluoride glasses* (521)
 Radiative and non-radiative transitions of rare-earth ions in glasses* (538)

TITLE INDEX

glasses

Energy transfer between Bi^{3+} yields Eu^{3+} , Bi^{3+} yields Sm^{3+} and UO_2^{2+} yields Eu^{3+} in oxide glasses* (539)

Structure of Nd^{3+} terms in nitrate glasses* (664)

gold

Magnetic susceptibility and electronic structure in metallic and non-metallic liquid cesium-gold and cesium-antimony alloys* (190)

Goldschmidt

Structural or solid state chemistry of rare earth sesquioxides from Goldschmidt to the present time* (438)

grade

Spectrographic determination of trace impurities in reactor grade aluminium* (104)

Standard methods for chemical, mass spectrometric, and spectrochemical analysis of nuclear-grade uranium dioxide powders and pellets* (711)

Standard methods for chemical, mass spectrometric, and spectrochemical analysis of nuclear-grade plutonium dioxide powders and pellets* (712)

Standard methods for chemical, mass spectrometric, and spectrochemical analysis of nuclear-grade mixed oxides (U, Pu) O_2 * (713)

Standard methods for chemical, mass spectrometric, spectrochemical, nuclear, and radiochemical analysis of nuclear-grade plutonium metal* (714)

gradient

Composition gradient during electrolysis in mixtures analogous to molten salt battery electrolytes* (234)

Oxygen redistribution in UO_2 due to a temperature gradient* (401)

gradients

The calculation of the consequences of sodium/fuel interactions with allowance for temperature gradients in the sodium* (272)

graphite

A Moessbauer spectroscopic study of intercalation compounds of graphite with ferric chloride and aluminum chloride* (627)

H

Me_2ZrCl_6 - H_2 ($\text{Me} = \text{Na}, \text{K}, \text{Rb}, \text{Cs}$) system* (45)

High-temperature microcalorimetric measurements of partial molar enthalpy of solution, $\Delta H(\text{O}_2)$, in plutonium + oxygen* (107)

Calculation and discussion of state diagrams in the system $\text{Pu}-\text{F}-\text{O}-\text{H}$ * (222)

Study of kinetics of monovalent indium ions formation in the $\text{In}-\text{InCl}_3-\text{LiCl}-\text{H}_2\text{O}$ system* (341)

Synthesis and characterization of a new uranium(V) compound: $\text{H}_3\text{UO}_6 + \text{UF}_6$ * (409)

Preparational, structural and vibrational study of $\text{M}_2\text{U}_2\text{O}_5\text{Cl}_4$, $2\text{H}_2\text{O}$ ($\text{M} = \text{Rb}, \text{Cs}$). Detection of a tetranuclear anion [$(\text{UO}_2)_4\text{O}_{12}\text{Cl}_8(\text{H}_2\text{O})_2$] $^{4-}$ * (498)

X-ray diffraction and infrared: Raman spectroscopic study of the hydrogen dinitrate ion, $(\text{O}_2\text{NO}-\text{H}-\text{ONO})$, in cesium hydrogen dinitrate* (695)

TITLE INDEX

hafnium

Properties of zirconium and hafnium tetrachlorides and their saturated solutions* (136)

Half

Half-life of ^{242}Pu from precise low-temperature heat capacity measurements* (487)

halide

Spectral properties of impurity molecules of the C_{3v} symmetry groups in solid solutions of halide salts* (44)
Phase equilibrium studies of lithium halide-containing electrolytes* (282)

Rare earth metal-metal halide systems. 19. Structural characterization of the reduced holmium chloride $\text{Ho}_{0.5}\text{Cl}_{1.5}$ * (378)

halides

Mixed valence rare-earth halides and their unusual crystal structures* (29)

Temperature effect on vibrational spectra of the impurity molecular ions in alkali halides* (46)

Conversion of protactinium monocarbide to the penta- and tetra-halides* (80)

Preparation and some properties of actinide(III) and lanthanide(III) oxide halides, and the enthalpies of formation of $\text{PuBr}_3(\text{s})$ and $\text{PuI}_3(\text{s})$ * (82)

Temperature dependence of vapor pressure of rare earth halides and their allcys* (99)

Origin of excess entropy of alkali metals halides solid solutions* (390)

Energies of valence states of atoms in halides* (499)

Thermodynamics of the lanthanide halides. II. Heat capacities and Schottky anomalies of SmCl_3 , EuCl_3 and GdCl_3 from 50 to 350 K* (594)

halogenides

Quaternary system of potassium halogenides* (90)

handbook

MATERO: Version 09. A handbook of materials properties for use in the analysis of light water reactor fuel rod behavior* (387)

hard

Theory for the derivation of a single hard-sphere potential describing the overall molecular interactions in binary liquid metal alloys as a function of temperature and composition* (525)
Application of the hard-sphere model to thermodynamic properties of liquid 3d transition metals* (615)

hazards

EUTECTIC DATA: safety, hazards, corrosion, melting points, compositions, and bibliography* (276)

heat

Thermodiffusion measurements in liquid alloys and electron transport heat calculations in the case of almost-free electrons* (33)

Experimental determination of formation heat of gaseous Ga_{20} and In_{20} * (105)

Specific heat of thorium monocarbide from 2 to 300K* (126)

Heat transfer bibliography* (158)

TITLE INDEX

Heat

- Heat capacity of $^{242}\text{PuO}_2$ from 12 to 350°K and of $^{244}\text{PuO}_2$ from 4 to 25°K. Entropy, enthalpy, and Gibbs energy of formation of PuO_2 at 298.15°K* (184)
- Specific heat of liquid metals and ceramics* (244)
- Phase transition in doped U_{4-y}O_9 . I. Heat capacity measurement* (261)
- Lambda-type heat capacity anomalies in U_3O_8 * (262)
- Partial pressures, activities and evaporation heat in the KF-ZrF_4 system* (295)
- Thermodynamic state, specific heat, and enthalpy function of saturated UO_2 vapor between 3000 K and 5000 K* (297)
- Ceramic-base heat resisting constructional material* (348)
- Heat capacities and heat content of molten cerium by levitation calorimetry* (355)
- Sodium uranium(V) trioxide, NaUO_3 : heat capacity and thermodynamic properties from 5 to 350 K* (383)
- Sodium uranium(V) trioxide, NaUO_3 : heat capacity and thermodynamic properties from 5 to 350 K* (384)
- Specific heat of Pu_{1-x}C and $\text{Pu}_{2/3}\text{C}_{1/3}$ from 10 to 300 K* (414)
- Specific heat of Pu_{1-x}C and $\text{Pu}_{2/3}\text{C}_{1/3}$ from 10 to 300 K* (415)
- Heat of formation of solid and liquid alloys of transition metals* (425)
- Periodically variable wall temperature effect on heat transfer in liquid metals* (437)
- Half-life of ^{242}Pu from precise low-temperature heat capacity measurements* (487)
- New description of liquid metal heat transfer in closed conduits* (502)
- Heat transfer measurements in a forced convection loop with two molten-fluoride salts: $\text{LiF--BeF}_2\text{--ThF}_4\text{--UF}_4$ and eutectic $\text{NaBF}_4\text{--NaF}$ * (585)
- Thermodynamics of electrolytes. 8. High-temperature properties, including enthalpy and heat capacity, with application to sodium chloride* (586)
- High-temperature heat contents and entropies of zirconium fluoride and zirconium sulfate* (591)
- Thermodynamics of the lanthanide halides. II. Heat capacities and Schottky anomalies of SmCl_3 , EuCl_3 and GdCl_3 from 50 to 350 K* (594)
- Heat of formation of europium sesquioxide and europium trichloride* (607)
- Enthalpy and heat capacity of liquid plutonium and uranium* (619)
- Enthalpy and heat capacity of liquid plutonium and uranium* (620)

Heats

- Heats of formation of solid solutions of CsCl with CsBr * (543)
- Heats of formation of holmium and terbium trichlorides* (608)
- Thermographic determination of formation heats of lower zirconium chlorides and bromides* (635)

heavier

- Comparative study of systems of heavier rare-earth metals with hydrogen [Gd , Tb , Dy , Lu]* (418)

TITLE INDEX

heavy

Revised model for 5d-electrons in the heavy rare-earth metals* (157)

hepta

Structural study and properties of the alkali metal, nitrosyl, and ammonium hepta- and octafluorouranates(VI)* (66)

hexabromotellurates

Investigation into reactions of hexabromotellurates of potassium, cesium and thallium(I) in melt* (556)

hexachloride

Some chemical reactions of uranium hexachloride and the chlorination of uranium tetrachloride* (81)

hexafluoride

Vibrational spectra of transition metal hexafluoride crystals. I. Orthorhombic MoF_6 , WF_6 , UF_6 neat crystals. Technical report* (51)

Vibrational spectra of transition metal hexafluoride crystals.

III. Exciton band structures of MoF_6 , WF_6 , and UF_6 . Technical report* (52)

Isotope effect on the thermodynamic quantities of gaseous uranium hexafluoride* (78)

Calculation of the reduced partition function ratio for uranium hexafluoride* (285)

Thermophysical and electrophysical properties of uranium hexafluoride at temperatures of $(1-11) \times 10^3$ °K and pressures of 0.1-100 atm* (302)

The spectral properties of uranium hexafluoride and its thermal decomposition products* (345)

Isotope effects in the molecular spectrum of plutonium hexafluoride* (353)

Investigation of liquid-steam equilibrium in system with diluted solutions of metal fluorides in uranium hexafluoride* (526)

hexafluorides

Thermodynamic similarity and universal equations of state of hexafluorides* (397)

hexafluorozirconates

New hexafluorozirconates(IV): BaZrF_6 , PbZrF_6 , EuZrF_6 , SrZrF_6 * (419)

HF

K_2HF_6 - K_2ZrF_6 -1.25 percent of HF system at 40°C with other solubility curves from 25°C to 70°C * (541)

Preparation and structure of transition metal fluorides $\text{U}^{(II)}\text{M}^{(IV)}\text{F}_6$ [$\text{U}^{(II)}$: Cr-Zn; $\text{M}^{(IV)}$: Hf, Zr]* (596)

HfCl

Absorption spectrum of HfCl molecule* (283)

Density and viscosity of distillable products from ZrCl_4 and HfCl_4 interaction with POCl_3 * (458)

HfF

K_2HF_6 - K_2ZrF_6 -1.25 percent of HF system at 40°C with other solubility curves from 25°C to 70°C * (541)

Hg

Cs-Hg amalgam thermodynamic properties* (339)

Thermodynamic properties of Cd-Zn-Hg liquid alloys* (340)

TITLE INDEX

HgCl

Nuclear quadrupole resonance spectra of HgCl_2 under pressure* (31)

higher

Simple Raman cell and furnace usable at temperatures higher than 1000°C for corrosive melts* (204)

Preparation and characterization of higher uranates(VI) of thallium(I)* (278)

Partly filled shells constituting anti-bonding orbitals with higher ionization energy than their bonding counterparts* (280)

Ho

Rare earth metal--metal halide systems. 19. Structural characterization of the reduced holmium chloride $\text{Ho}_{0.5}\text{Cl}_{1.1}$ * (378)

Gd, Tb, Dy, Ho, Er solubility in chromium* (610)

Phase equilibria and thermodynamic data for the ternary UO_2 - UO_3 - ReO_3 (1.5) (Ho, Tm, Lu) systems* (694)

HoF

CsF - HoF_3 system* (572)

holmium

Rare earth metal--metal halide systems. 19. Structural characterization of the reduced holmium chloride $\text{Ho}_{0.5}\text{Cl}_{1.1}$ * (378)

Investigation into reaction of rubidium and holmium fluorides* (540)

Heats of formation of holmium and terbium trichlorides* (608)

hot

Investigation of the parameters influencing the microstructure of hot-pressed boron carbide* (440)

HTGR

Formation of actinides in irradiated HTGR fuel elements* (150)

HTR

Microprobe study of fission product behavior in high-burnup HTR fuels* (315)

hydride

Pulse NMR in solids: chemical shift, lead fluoride, and thorium hydride* (363)

Hydrodynamics

Hydrodynamics of post-disassembly fuel expansions* (310)

hydrogen

Transport of hydrogen in liquid lithium* (10)

Hydrogen solubility in liquid cesium* (22)

Stability of rare earth metaniobates in hydrogen atmosphere* (164)

Some properties of hydrogen fluoride compounds with group 5 metal pentafluorides* (170)

Hydrogen solubility in rare earth intermetallic compounds* (217)

Hydrogen solubility in liquid zirconium* (287)

Hydrogen absorption of rare-earth (3d) transition intermetallic compounds* (392)

Solution behavior of hydrogen isotopes and other non-metallic elements in liquid lithium* (404)

Comparative study of systems of heavier rare-earth metals with hydrogen [Gd, Tb, Dy, Lu]* (418)

Thermodynamic study of the neptunium-hydrogen system* (428)

TITLE INDEX

hydrogen

Intensity-voltage curves of hydrogen diffusion electrodes in eutectic LiCl--KCl * (507)
 Reaction rates and electrical resistivities of the hydrogen isotopes with, and their solubilities in, liquid lithium* (529)
 Terminal solid solubility of hydrogen in the zirconium-2.5 weight % niobium alloy* (587)
 Thermodynamic properties of solutions of hydrogen isotopes in metals and alloys of interest to fusion reactor technology* (663)
 X-ray diffraction and infrared; Raman spectroscopic study of the hydrogen dinitrate ion, $(\text{O}^2\text{NO-H-ONO}^2)^+$, in cesium hydrogen dinitrate* (695)

hydrolysis

Influence of salt background on kinetics of BF_4^+ alkali hydrolysis* (504)

hyperstoichiometric

Determination of oxygen potentials of hyperstoichiometric U--Pu dioxides in the temperature range 1,500-1,850 K* (109)
 Determination of oxygen potentials of hyperstoichiometric U-Pu dioxides in the temperature range 1500-1850 K* (110)

Identification

Identification of UN in Ar matrices* (219)
 Collection of gamma spectra data of irradiated light water moderated reactor spent fuel and study of the applicability of the methods for fuel identification. Final report for the period 1 February 1972--October 1975* (453)

IKS

High-temperature unit to the IKS-21 spectrometer for the detection of molten salts absorption spectra* (208)

impedance

Complex impedance of electrochemical cells based on yttria doped thorium* (568)

impurities

Spectrographic determination of trace impurities in reactor grade aluminium* (104)
 Determination of trace impurities of titanium and zirconium in molybdenum and tungsten and their oxygen compounds by X-ray fluorescence spectrometry* (698)

impurity

Spectral properties of impurity molecules of the C_{3v} symmetry groups in solid solutions of halide salts* (44)
 Temperature effect on vibrational spectra of the impurity molecular ions in alkali halides* (46)

inception

Experimental investigation of cavitation inception and damage in a flowing sodium environment* (103)

InCl

Thermal analysis of $\text{GaCl}_3 + 3\text{In} = \text{InCl} + \text{Ga}$ system* (4)
 Tensimetric study of pressure and composition of saturated vapor in TlCl-InCl_3 system* (93)
 Study of kinetics of monovalent indium ions formation in the $\text{In}^0\text{-InCl}_3\text{-LiCl-H}_2\text{O}$ system* (341)

incongruent

Interval of congruent and incongruent sublimation of $3\text{KF} \cdot \text{ZrF}_4$ compound in the KF-ZrF_4 system* (296)

TITLE INDEX

index

Equilibrium index for fission gas bubbles* (685)

indium

Investigation into interaction of indium tribromide with dibromides of manganese, cobalt and nickel in melt* (153)

Interaction of lower indium chlorides with chloride and chloroindates of potassium* (174)

Solubility and activity of oxygen in liquid indium and copper-indium alloys* (182)

Thermodynamic study of the binary system copper-indium. I. Calorimetric study* (290)

Thermodynamic study of the binary system copper--indium. II. Potentiometric study* (291)

Study of kinetics of monovalent indium ions formation in the $\text{In}^{\text{sup } 0}\text{-InCl}^{\text{sub } 3}\text{-LiCl-H}^{\text{sub } 2}\text{O}$ system* (341)

Thermodynamic properties of diluted sodium solutions in liquid indium* (388)

Thermodynamic studies of gallium--indium liquid alloys by solid state electrochemistry with oxide electrolytes* (513)

Some structure differences of double oxides of indium and gallium with antimony(5)* (657)

Eutectic point of gallium-indium system* (706)

inert

Prediction of the inert gas solubilities in stoichiometric molten $\text{UO}^{\text{sub } 2}$ * (225)

Influence

Influence of the technology of the alloy fabrication on its properties in a liquid state* (643)

Infrared

Infrared spectroscopy of the uranium/oxygen system* (12)

Predicted infrared spectrum and the structure of the isolated $\text{UF}^{\text{sub } 5}$ molecule* (350)

Investigation of some rare earth aluminates by the method of infrared spectroscopy* (357)

Quantitative determination of $\text{UO}^{\text{sub } 2}$, and $\text{ThO}^{\text{sub } 2}$ in powdery mixtures by infrared spectrometry* (431)

Investigation into absorption infrared spectra of vapors under $\text{ScBr}^{\text{sub } 3}$, $\text{YCl}^{\text{sub } 3}$, $\text{LaCl}^{\text{sub } 3}$, $\text{GdCl}^{\text{sub } 3}$, $\text{ErCl}^{\text{sub } 3}$, $\text{LuCl}^{\text{sub } 3}$ by means of isolation method in rare gas matrix* (497)

X-ray diffraction and infrared: Raman spectroscopic study of the hydrogen dinitrate ion, $(\text{O}^{\text{sub } 2}\text{NO-H-ONO}^{\text{sub } 2})^{\text{sup } -}$, in cesium hydrogen dinitrate* (695)

inorganic

Stable inorganic matrix materials for high temperature batteries. Final report No. 1, 2 June 1967--2 February 1968* (23)

Interactions in inorganic molecular crystals - electronic spectra of $\text{ReF}^{\text{sub } 6}$ pure and mixed crystals. Technical report, 1 Jul--31 Dec 1975* (53)

Chemical Engineering Division physical inorganic chemistry. Annual report, July 1975--June 1976* (58)

input

Accurate procedure to safeguard the fissile material content of input and output solutions of reprocessing plants* (131)

TITLE INDEX

Installations

Secnd specialist meeting on sodium/fuel interaction in fast reactors, Ispra, 21-23 November 1973. Sponsored by the Committee on the Safety of Nuclear Installations, OECD Nuclear Energy Agency, Paris, France* (272)

Institute

Israel Nuclear Society symposium on "'Nuclear fuel in Israel'", Weizmann Institute of Science, Rehovot, 1.7.75* (226)

Twenty years of research in the processing of radioactive fuel and neutralization of radioactive waste at the Nuclear Research Institute (NRI)* (358)

Intensities

Intensities of lanthanide f-f transitions* (492)

Intensity

Intensity-voltage curves of hydrogen diffusion electrodes in eutectic LiCl--KCl* (507)

Use of oscillator strengths of "'supersensitive'" ion transfers of lanthanides for calculation of some parameters determining the intensity of their absorption spectra* (511)

interaction

Study of chemical interaction in $\text{Pb}_{3\text{AlF}}\text{LaF}_6$ - $\text{Ca}_{3\text{AlF}}\text{LaF}_6$ system* (26)

Interaction of uranyl chloride with zirconium tetrachloride in the melt of sodium chloride and potassium chloride* (27)

Interaction of the lithium-tin alloys with the LiCl-KCl-CsCl melted eutectic mixture* (91)

Interaction of uranium trichloride with alkaline earth metal chlorides* (139)

Interaction of uranium- and thorium tetrachlorides with sodium- and potassium chlorides* (144)

Investigation into interaction of indium tribromide with dibromides of manganese, cobalt and nickel in melt* (153)

Molybdenum and tungsten interaction with liquid aluminium* (156)

On interaction in U--S--Te system* (162)

Interaction of lower indium chlorides with chloride and chloroindates of potassium* (174)

Secnd specialist meeting on sodium/fuel interaction in fast reactors, Ispra, 21-23 November 1973. Sponsored by the Committee on the Safety of Nuclear Installations, OECD Nuclear Energy Agency, Paris, France* (272)

Interaction of atoms of rare earths-metal alloys* (320)

Interaction of lanthanide ions with fluorine-ions* (329)

On interaction of tellurium tetrachloride with copper (1), mercury (2), bismuth and iron (3) chlorides* (336)

Thermogravimetric investigation of interaction of some rare earth oxides with ammonium bifluoride* (427)

Interaction parameters in the Zn--Pb--Sn system at low zinc concentration* (436)

Interaction of rubidium and cesium fluorides with lutetium trifluorides* (444)

Density and viscosity of distillable products from ZrCl_4 and HfCl_4 interaction with POCl_3 * (458)

Lanthanum chloride interaction with magnesium oxide* (493)

Interaction of neodymium- and yttrium chlorides with magnesium oxide* (494)

TITLE INDEX

Interaction

- Interaction of thorium chloride with calcium oxide* (496)
- Interaction of potassium tetrafluoroborate with fluorides of potassium, lanthanum, neodymium in melt* (530)
- Thermographic study of iron, molybdenum, tantalum and tungsten high chlorides interaction* (555)
- Interaction in AgCl-CsCl, CsCl-KCl systems and liquidus surface of AgCl-CsCl-KCl ternary system* (659)
- Investigation of particles interaction in liquid metals and alloys* (661)
- Interaction of plutonium trichloride in melts of ternary systems* (682)
- Fuel-clad interaction. C-E thermo-structural fuel evaluation methods* (719)

Interactions

- Interactions in inorganic molecular crystals - electronic spectra of ReF_6 pure and mixed crystals. Technical report, 1 Jul--31 Dec 1975* (53)
- Knudsen cell--mass spectrometer studies of cesium--urania interactions* (120)
- Fine definition of ir spectra from high temperature interactions of $\text{U} + \text{O}_2$. Part II. Final report, 1 July 1974--30 June 1975* (218)
- The calculation of the consequences of sodium/fuel interactions with allowance for temperature gradients in the sodium* (272)
- Theory for the derivation of a single hard-sphere potential describing the overall molecular interactions in binary liquid metal alloys as a function of temperature and composition* (525)

intercalation

- A Mossbauer spectroscopic study of intercalation compounds of graphite with ferric chloride and aluminum chloride* (627)

Interdiffusion

- Interdiffusion in the CaF_2 -- YF_3 system* (669)

interfacial

- Variation of the solid--solid--nickel--thoria interfacial free energy with temperature* (5)
- Measurement of interfacial energies in the UO_2 -- Gd_2O_3 system* (697)

intergranular

- Effect of composition and structure on crevice, intergranular, and stress corrosion of some wrought Ni--Cr--Mo alloys* (604)

Interim

- Nonaqueous electrolyte systems: ionic transport in nonaqueous media. Interim technical report No. 7* (246)

interlanthanide

- Phase equilibrium in interlanthanide oxides systems* (49)

intermediate

- Statistical mechanics of intermediate-valence rare earth compounds* (279)
- Influence of intermediate products on chlorination kinetics of vanadium oxides in salt melt* (356)

intermetallic

- Hydrogen solubility in rare earth intermetallic compounds* (217)
- Hydrogen absorption of rare-earth (3d) transition intermetallic compounds* (392)

TITLE INDEX

intermetallic

Thermal decomposition of intermetallic compounds of platinum with alkaline earth and lanthanide metals at high temperatures* (399)

2. all-union conference on crystallochemistry of intermetallic compounds* (549)

2. all-union conference on crystallochemistry of intermetallic compounds* (671)

intermetallics

Fault structures in rare earth--cobalt intermetallics* (11)

Thermodynamics and kinetics of the formation of rare earth intermetallics* (137)

international

Proceedings of the international conference on liquid technology in energy production* (56)

Third international symposium on fuel cells* (77)

5th international conference on plutonium and other actinides* (84)

Proceedings of the international conference on liquid technology in energy production* (103)

5th international conference on plutonium and other actinides.

Book of abstracts* (109)

5th international conference on plutonium and other actinides* (228)

5th international conference on plutonium and other actinides* (415)

5th international conference on plutonium and other actinides* (479)

Proceedings of the international conference on liquid technology in energy production* (590)

5th international conference on plutonium and other actinides.

Book of abstracts* (620)

interpretation

Phase analysis at simulated LWR fuel rods and their interpretation by constitution investigations in the uranium-zirconium-oxygen system* (510)

Interval

Interval of congruent and incongruent sublimation of $3\text{Kf}_x\text{ZrF}_{5-x}$ compound in the $\text{Kf}-\text{ZrF}_5$ system* (296)

iodate

Determination of thorium in rare earths chloride: coprecipitation with cerium (IV) iodate and spectrophotometric determination with thoron* (447)

iodide

Solubility of sodium iodide in liquid sodium between 140 and 500°C* (154)

iodides

X-ray diffraction study into molten bromides and iodides of alkali metals* (20)

Ternary mutual system of chlorides and iodides of rubidium and cadmium* (257)

Quaternary mutual system of chlorides, bromides, iodides of sodium and rubidium* (629)

Quaternary mutual system of chlorides, bromides, iodides of potassium and cesium* (630)

Quaternary mutual system of chlorides, bromides, iodides of rubidium and cesium* (640)

TITLE INDEX

ion

Study of the compounds $M\text{sub } 3\text{UO}\text{sub } 2\text{F}\text{sub } 5$ ($M = K, Rb, Cs, NH\text{sub } 4$) by i.r. absorption and Raman diffusion spectrophotometry. III. Effect of cation on vibration frequency of the ion $\text{UO}\text{sub } 2\text{F}\text{sub } 5^{\text{sup } 3-}$ * (127)
 Thermodynamic perturbation theory for the surface tension and ion density profile of a liquid metal* (167)
 Use of oscillator strengths of "'supersensitive'" ion transfers of lanthanides for calculation of some parameters determining the intensity of their absorption spectra* (511)
 X-ray diffraction and infrared: Raman spectroscopic study of the hydrogen dinitrate ion, $(O\text{sub } 2NO-H-ONO\text{sub } 2)^{\text{sup } -}$, in cesium hydrogen dinitrate* (695)

ionic

Nonaqueous electrolyte systems: ionic transport in nonaqueous media. Interim technical report No. 7* (246)
 Ionic-thermocurrent study of the dipole relaxation and equilibrium in Gd-doped $SrF\text{sub } 2$ * (410)
 Application of molecular dynamics computations to the conformal ionic solution theory: Thermodynamics of binary molten salt mixtures* (550)
 Theoretical concepts useful in the calculation or storage of phase diagrams of ionic systems* (552)

ionization

Partly filled shells constituting anti-bonding orbitals with higher ionization energy than their bonding counterparts* (280)

ions

Optical spectra of the difluoride, dichloride, and trichloride ions in the matrix-isolated $M^{\text{sup } +}F\text{sub } 2^{\text{sup } -}$, $M^{\text{sup } +}Cl\text{sub } 2^{\text{sup } -}$, and $M^{\text{sup } +}Cl\text{sub } 3^{\text{sup } -}$ species* (16)
 Optical spectra of the dibromide and diiodide ions in the matrix-isolated $M^{\text{sup } +}Br\text{sub } 2^{\text{sup } -}$ and $M^{\text{sup } +}I\text{sub } 2^{\text{sup } -}$ species* (17)
 Temperature effect on vibrational spectra of the impurity molecular ions in alkali halides* (46)
 Comparison of the tendencies of complex formation of various metal ions in fused $LiF-NaF-KF$ (eut.) and in fused $LiCl-KCl$ (eut.)* (314)
 Interaction of lanthanide ions with fluorine-ions* (329)
 Study of kinetics of monovalent indium ions formation in the $In^{\text{sup } 0}-InCl\text{sub } 3-LiCl-H\text{sub } 2O$ system* (341)
 Progress in the spectroscopy of rare earth and transition metal ions* (412)
 Effect of fluoride ions on the kinetics of the oxidation of uranium(IV) by cerium(IV)* (424)
 Electrochemistry of niobium and tantalum, (5). Anodic dissolution of tantalum and the average valence of tantalum ions in molten KCl * (450)
 Radiative and non-radiative transitions of rare-earth ions in glasses* (538)
 Glass formation in nitrate systems containing rare earth ions* (666)
 Isorhodanide complexes of 3d-metal ions in molten salts* (672)

TITLE INDEX

ir

Fine definition of ir spectra from high temperature interactions of $U + O_2$. Part II. Final report, 1 July 1974--30 June 1975* (218)

IR spectroscopic study of dipotassium and dicesium uranyl tetrabromides* (483)

IR spectra of the $NbCl_5$, $TaCl_5$, $NbBr_5$ and $NbOCl_3$ vapors* (707)

iron

Viscosity characteristics of the iron-chromium-nickel melts* (24)

Saturation vapor pressure in iron chloride-alkaline earth metal chlorides systems* (97)

Physical models of mass transport of iron and nickel in liquid sodium systems* (128)

Kinetic peculiarities of nitrogen dissolving in iron-chromium and iron-chromium-nickel melts* (172)

Spreading of iron-nickel alloys on ZrC and nickel-cobalt alloys on ZrN* (252)

Effect of various additions on the temperature of structural transformation in liquid iron* (313)

On interaction of tellurium tetrachloride with copper (1), mercury (2), bismuth and iron (3) chlorides* (336)

Thermographic study of iron, molybdenum, tantalum and tungsten high chlorides interaction* (555)

Deoxidation thermodynamics of cerium and lanthanum in liquid iron* (654)

irradiated

Formation of actinides in irradiated HTGR fuel elements* (150)

Chemical effects of composition changes in irradiated oxide fuel materials. II. Fission product segregation and chemical equilibria* (169)

Investigation into americium extraction process from residues of irradiated nuclear fuel after its fluorination by SiF_4-NaF and SiF_4-CaF_2 melts of eutectic composition* (308)

Radial oxygen distribution in the fuel of irradiated mixed oxide fuel elements with different final stoichiometries* (316)

Collection of gamma spectra data of irradiated light water moderated reactor spent fuel and study of the applicability of the methods for fuel identification. Final report for the period 1 February 1972--October 1975* (453)

irradiation

Investigations on radioactive fission product correlations: gamma spectrometry measurements on spent fuel assemblies discharged from the Trino Vercellese reactor at the end of the 2nd irradiation cycle* (75)

isolated

Optical spectra of the difluoride, dichloride, and trichloride ions in the matrix-isolated $M^{sup} + F_2^{sup} -$, $M^{sup} + Cl_2^{sup} -$, and $M^{sup} + Cl_3^{sup} -$ species* (16)

Optical spectra of the dibromide and diiodide ions in the matrix-isolated $M^{sup} + Br_2^{sup} -$ and $M^{sup} + I_2^{sup} -$ species* (17)

Predicted infrared spectrum and the structure of the isolated UF_5 molecule* (350)

TITLE INDEX

isolation

Investigation into absorption infrared spectra of vapors under ScBr_3 , YCl_3 , LaCl_3 , GdCl_3 , ErCl_3 , LuCl_3 by means of isolation method in rare gas matrix* (497)

Isorhodanide

Isorhodanide complexes of 3d-metal ions in molten salts* (672)

Isotope

Isotope effect on the thermodynamic quantities of gaseous uranium hexafluoride* (78)

Isotope effects in the molecular spectrum of plutonium hexafluoride* (353)

Reactor conference, Duesseldorf, 30.3.-2.4.1976. Section 2: Fuel elements, fuel cycle and isotope technique* (480)

isotopes

Solution behavior of hydrogen isotopes and other non-metallic elements in liquid lithium* (404)

Reaction rates and electrical resistivities of the hydrogen isotopes with, and their solubilities in, liquid lithium* (529)

Thermodynamic properties of solutions of hydrogen isotopes in metals and alloys of interest to fusion reactor technology* (663)

Ispra

Second specialist meeting on sodium/fuel interaction in fast reactors, Ispra, 21-23 November 1973. Sponsored by the Committee on the Safety of Nuclear Installations, OECD Nuclear Energy Agency, Paris, France* (272)

Israel

Israel Nuclear Society symposium on "Nuclear fuel in Israel", Weizmann Institute of Science, Rehovot, 1.7.75* (226)

Japan

Recent development of ceramic basic science in Japan* (557)

U.S.-Japan joint seminar on defects and diffusion in solids* (611)

Johannesburg

Eleventh annual seminar on theoretical physics, Johannesburg, 5-9 July 1976* (368)

joint

U.S.-Japan joint seminar on defects and diffusion in solids* (611)

Jul

Interactions in inorganic molecular crystals - electronic spectra of ReF_6 pure and mixed crystals. Technical report, 1 Jul--31 Dec 1975* (53)

Lithium--nickel fluoride secondary battery investigation. Quarterly technical progress report No. 1, 27 Apr--27 Jul 1966* (381)

July

Chemical Engineering Division physical inorganic chemistry. Annual report, July 1975--June 1976* (58)

Fine definition of ir spectra from high temperature interactions of $\text{U} + \text{O}_2$. Part II. Final report, 1 July 1974--30 June 1975* (218)

Chemical Engineering Division fuels and materials chemistry. Annual report, July 1975--June 1976* (281)

TITLE INDEX

July

Chemical Engineering Division, reactor safety and physical property studies annual report, July 1974--June 1975* (367)
 Eleventh annual seminar on theoretical physics, Johannesburg, 5-9 July 1976* (368)
 Physical research on liquid-metal systems. Annual report, July 1975--June 1976* (405)
 Analytical methods for fissionable materials in the nuclear fuel cycle. Progress report, July 1, 1975--September 30, 1976* (690)
 Electrochemical methods for monitoring purity of molten salts. Semiannual progress report, July--December 1975* (721)

Jun

Environmental sensitivity of structural metals: liquid metal embrittlement. Annual technical progress report No. 2, 25 Jun 1970--31 May 1971* (95)
 Environmental sensitivity of structural metals: liquid metal embrittlement. Annual technical progress report No. 4, 1 Jun 1972--31 May 1973* (211)
 Environmental sensitivity of structural metals: liquid metal embrittlement. Annual technical progress report No. 2, 25 Jun 1970--31 May 1971* (212)
 Structure of lithium--boron alloys and their performance as potential anodes for high power molten salt batteries. Progress report, 1 Aug 1974--30 Jun 1975* (275)

June

Stable inorganic matrix materials for high temperature batteries. Final report No. 1, 2 June 1967--2 February 1968* (23)
 Chemical Engineering Division physical inorganic chemistry. Annual report, July 1975--June 1976* (58)
 LMFER Fuel Reprocessing Program progress report for period April 1 to June 30, 1976* (94)
 Summaries of reports of the all-union meeting on solid body chemistry. 3--5 June 1975. III. Synthesis, phase equilibrium and use of solid oxide compounds* (114)
 Fine definition of ir spectra from high temperature interactions of $O + O_{\text{sub } 2}$. Part II. Final report, 1 July 1974--30 June 1975* (218)
 Chemical Engineering Division fuels and materials chemistry. Annual report, July 1975--June 1976* (281)
 Chemical Engineering Division, reactor safety and physical property studies annual report, July 1974--June 1975* (367)
 Physical research on liquid-metal systems. Annual report, July 1975--June 1976* (405)
 Summaries of reports of the all-union meeting on solid body chemistry. 3--5 June 1975. III. Synthesis, phase equilibrium and use of solid oxide compounds* (641)

K

$Me_{\text{sub } 2}ZrCl_{\text{sub } 6}$ -- $H_{\text{sub } 2}$ (Me-Na,K,Rb,Cs) system* (45)
 Calculation of vapor pressures of oxide fuels up to 5000 K in equilibrium and non-equilibrium evaporation* (72)
 Calculation of vapor pressures of oxide fuels up to 5,000 K for equilibrium and nonequilibrium evaporation* (73)
 Calculation of the oxygen potential of $(U,Pu)O_{\text{sub } 2-x}$ with defect models and measured oxygen potential data in the temperature range from 1,000 to 1,700 K* (74)
 Phase equilibria in Li--Na--K carbonate/aluminate systems* (77)

TITLE INDEX

K

- Equilibrium between Li-K-Te alloys and LiCl-KCl molten eutectic mixture* (92)
- Determination of oxygen potentials of hyperstoichiometric U--Pu dioxides in the temperature range 1,500-1,850 K* (109)
- Determination of oxygen potentials of hyperstoichiometric U-Pu dioxides in the temperature range 1500-1850 K* (110)
- Study of the compounds $M\text{UO}_2\text{F}_5$ ($M = \text{K}, \text{Rb}, \text{Cs}, \text{NH}_4$) by i.r. absorption and Raman diffusion spectrophotometry. III. Effect of cation on vibration frequency of the ion $\text{UO}_2\text{F}_5^{3-}$ * (127)
- Phase diagrams of Li_2MoO_4 -- ZnMoO_4 , K_2MoO_4 -- ZnMoO_4 and K_2WO_4 -- ZnWO_4 systems* (160)
- Gases and carbon in metals (thermodynamics, kinetics, and properties). I. Alkali metals, alkaline earth metals, light metals (Li, Na, K, Rb, Cs; Ca, Sr, Ba; Be, Mg, Al)* (192)
- Thermodynamic state, specific heat, and enthalpy function of saturated UO_2 vapor between 3000 K and 5000 K* (297)
- Fusibility in $\text{K}_2\text{Nb}(\text{Ta})\text{F}_7$ --LiF--NaF systems* (298)
- Meltability in system of K_2TaF_7 -- NaF -- NaCl -- KCl * (299)
- Meltability in K_2NbF_7 -- NaF -- NaCl -- KCl system* (300)
- Double tungstate $\text{K}_5\text{La}(\text{WO}_4)_4$ * (338)
- Standard free energy of formation of liquid Na_2CrO_4 , 1100⁰ to 1220⁰ K* (375)
- Sodium uranium(V) trioxide, NaUO_3 : heat capacity and thermodynamic properties from 5 to 350 K* (383)
- Sodium uranium(V) trioxide, NaUO_3 : heat capacity and thermodynamic properties from 5 to 350 K* (384)
- Investigation of calcite solubility in K_2CO_3 and Na_2CO_3 solutions at temperatures of 200 to 350⁰ C* (396)
- Some thermodynamic properties of dilute solutions of actinide chlorides in (Li-K)Cl and in (Na-K)Cl eutectics* (406)
- Thermodynamic properties of dilute solutions of ThCl_4 in (Li-K)Cl and (Na-K) eutectics* (407)
- Specific heat of $\text{PuC}_{(1-x)}$ and Pu_2C_3 from 10 to 300 K* (414)
- Specific heat of $\text{PuC}_{(1-x)}$ and Pu_2C_3 from 10 to 300 K* (415)
- Thermal conductivity and diffusivity of UO_2 from 250 to 415 K* (459)
- Cesium nitrate. Standard enthalpies of solution and formation and the standard entropy at 298.15 K* (466)
- Vapor pressure studies over liquid uranium oxide and uranium plutonium oxide up to 5,000 K* (474)
- Extension of vapour pressure measurements of nuclear fuels (U, Pu) O_2 to 7000 K for fast reactor safety analysis* (475)
- Vapour pressure measurements to 7,000 K and equation of state of oxide fuels for fast reactor safety analysis. I. Extension of vapour pressure measurements of nuclear fuels (U, Pu) O_2 to 7,000 K* (477)
- Vapour pressure studies over liquid uranium plutonium oxides up to 5,000 K* (478)

TITLE INDEX

K

Practical application of conversion method in study of five-component mutual system consisting of nine salts NaF , Na_2MoO_4 , Na_2WO_4 , KF , K_2MoO_4 , K_2WO_4 , BaF_2 , BaMoO_4 , BaWO_4 * (516)

$\text{KF-KCl-K}_2\text{MoO}_4$ system* (520)

K_2HfF_6 -- K_2ZrF_6 --1.25 percent of HF system at 40°C with other solubility curves from 25°C to 70°C* (541)

New apparatus for thermophysical measurements above 2500 K* (574)

Viscosity of melts of $\text{KCl-KF-K}_2\text{ZrF}_6$ ternary system* (576)

Pressure and entropy calculations for liquid Na, K and Rb along the melting line* (584)

Thermodynamics of the lanthanide halides. II. Heat capacities and Schottky anomalies of SmCl_3 , EuCl_3 and GdCl_3 from 50 to 350 K* (594)

Some thermal properties of beryllium fluoride from 8°C to 1,200°C K* (617)

Viscosities of cesium vapor to 1,620 K and of liquid gallium to 1,800 K* (622)

Degree of thermal dissociation of MCaCl_3 (M=K, Rb, Cs) complexes during melting* (624)

Viscosities of cesium vapor to 1620 K and of liquid gallium to 1800 K* (681)

KFe

Study of KBeF_2 -- DyF_3 cross-section of KF-BeF_2 -- DyF_3 ternary system and some properties of this cross section glasses* (21)

KCl

Standard formation enthalpy of chloride alloys of ZrCl_4 -- KCl(NaCl) systems* (8)

Study of the Ni-Ni (II) equilibrium exchange, by radiotracer techniques in fused LiCl-KCl eutectic* (68)

Interaction of the lithium-tin alloys with the LiCl-KCl-CsCl melted eutectic mixture* (91)

Equilibrium between Li-K-Te alloys and LiCl-KCl molten eutectic mixture* (92)

Phase diagram of system LiCl-KCl-CaCrO_4 * (116)

Phase diagram for the ternary system CaCl_2 -- KCl-CaCrO_4 * (117)

Lattice-dynamical calculation of the thermodynamic properties of NaCl, KCl and RbCl * (231)

Structure and anodic discharge behavior of lithium-boron alloys in the LiCl-KCl eutectic melt.* (274)

Thermal investigation of NdCl_3 -- AlCl_3 -- KCl and NdCl_3 -- AlCl_3 -- CsCl system* (289)

Chronopotentiometry of U(III) and U(IV) in molten LiCl-KCl eutectic* (292)

Electrochemical behavior of U(III) and U(IV) in molten LiCl-KCl eutectic* (293)

Meltability in system K_2TaF_7 -- NaF-NaCl-KCl * (299)

Meltability in K_2NbF_7 -- NaF-NaCl-KCl system* (300)

TITLE INDEX

KCl

- Comparison of the tendencies of complex formation of various metal ions in fused LiF-NaF-KF (eut.) and in fused LiCl-KCl (eut.)* (314)
- Corrosion of liquid alloys of zinc with yttrium and with rare earth elements in LiCl--KCl melts* (335)
- Quantitative study of the chemical equilibria in solution in molten LiCl-KCl (70-30 mol.%) by absorption spectrophotometry. Application to neptunium* (385)
- Effect of temperature on the fine structure of Eu^{2+} in KCl and KI* (432)
- Electrode potentials of tantalum chlorides in a molten NaCl-KCl eutectic* (449)
- Electrochemistry of niobium and tantalum, (5). Anodic dissolution of tantalum and the average valence of tantalum ions in molten KCl* (450)
- Electrochemistry of niobium and tantalum. VI. Electrode potentials of tantalum chlorides in molten KCl* (451)
- Phase equilibrium in NdCl_3 -KCl system* (457)
- Raman spectra of solid ThCl_4 and fused ThCl_4 -KCl system* (488)
- Intensity-voltage curves of hydrogen diffusion electrodes in eutectic LiCl--KCl* (507)
- KF-KCl- K_2MoO_4 system* (520)
- MoCl_2 -NaCl and MoCl_2 -KCl binary systems* (547)
- Viscosity of melts of KCl-KF- K_2ZrF_6 ternary system* (576)
- Interaction in AgCl-CsCl, CsCl-KCl systems and liquidus surface of AgCl-CsCl-KCl ternary system* (659)
- Preparation of thorium metal by electrolysis of fused salt ThCl_4 -LiCl+NaCl+KCl(eut) system* (703)

key

CODATA recommended key values for thermodynamics, 1976. Report of the CODATA Task Group on key values for thermodynamics, 1976* (717)

KF

- Study of KBeF_2 - DyF_3 cross-section of KF- BeF_2 - DyF_3 ternary system and some properties of this cross section glasses* (21)
- Molecular vapor composition and mass spectra in KF- ZrF_4 , CsF- ZrF_4 systems* (294)
- Partial pressures, activities and evaporation heat in the KF- ZrF_4 system* (295)
- Interval of congruent and incongruent sublimation of $3\text{KF}\cdot\text{ZrF}_4$ compound in the KF- ZrF_4 system* (296)
- Comparison of the tendencies of complex formation of various metal ions in fused LiF-NaF-KF (eut.) and in fused LiCl-KCl (eut.)* (314)
- Practical application of conversion method in study of five-component mutual system consisting of nine salts NaF, Na_2MoO_4 , Na_2WO_4 , KF, K_2MoO_4 , K_2WO_4 , BaF_2 , BaMoO_4 , BaWO_4 * (516)
- KF-KCl- K_2MoO_4 system* (520)
- Viscosity of melts of KCl-KF- K_2ZrF_6 ternary system* (576)

TITLE INDEX

KI

Effect of temperature on the fine structure of Eu^{2+} in KCl and KI* (432)

Kinetic

Kinetic peculiarities of nitrogen dissolving in iron-chromium and iron-chromium-nickel melts* (172)

Kinetics

Kinetics of reaction of liquid lithium with oxygen and nitrogen* (106)

Thermodynamics and kinetics of the formation of rare earth intermetallics* (137)

Gases and carbon in metals (thermodynamics, kinetics, and properties). I. Alkali metals, alkaline earth metals, light metals (Li, Na, K, Rb, Cs; Ca, Sr, Ba; Be, Mg, Al)* (192)

Kinetics and mechanism into preparation of samarium-, europium- and ytterbium dichlorides and their some properties* (253)

Study of kinetics of monovalent indium ions formation in the $\text{In}^0\text{-InCl}_3\text{-LiCl-H}_2\text{O}$ system* (341)

Influence of intermediate products on chlorination kinetics of vanadium oxides in salt melt* (356)

Effect of fluoride ions on the kinetics of the oxidation of uranium(IV) by cerium(IV)* (424)

Influence of salt background on kinetics of BF_4^{+} alkali hydrolysis* (504)

Equilibria and kinetics in modern ceramic processing* (557)

KIC

Constitution equilibrium diagram of the $\text{LiNO}_3\text{-LiIO}_3$, $\text{NaNO}_3\text{-NaIO}_3$, $\text{KNO}_3\text{-KIO}_3$ system* (665)

KKR

KKR analysis of the Anderson-McMillan theory of electronic structure of liquid metals. Technical report* (485)

KMF

Vibrational spectra and force constants of the elpasolites Cs_2KMF_6 ($\text{M}=\text{Sc}, \text{Y}, \text{La}, \text{Gd}, \text{Yb}$)* (41)

KNO

Measurement of fluorescence of rare earths in molten salts (Nd^{3+} in $\text{NaNO}_3\text{-KNO}_3$ eutectic)* (233)

Solubility of $\text{Sr(NO}_3)_2$ in the $\text{LiNO}_3\text{-KNO}_3$ melt of variable composition* (351)

Constitution equilibrium diagram of the $\text{LiNO}_3\text{-LiIO}_3$, $\text{NaNO}_3\text{-NaIO}_3$, $\text{KNO}_3\text{-KIO}_3$ system* (665)

Knudsen

Knudsen cell--mass spectrometer studies of cesium--urania interactions* (120)

La

Thermodynamic properties of the liquid La--Mg alloys* (3)

Vibrational spectra and force constants of the elpasolites Cs_2KMF_6 ($\text{M}=\text{Sc}, \text{Y}, \text{La}, \text{Gd}, \text{Yb}$)* (41)

Density, viscosity, electric conductivity and meltability of $\text{Na}_3\text{AlF}_6\text{-La}_2\text{O}_3$ and $\text{Na}_3\text{AlF}_6\text{-Sm}_2\text{O}_3$ systems* (43)

^{139}La ENDOR of $\text{LaCl}_3\text{:Gd}^{3+}$ * (130)

Double tungstate $\text{K}_5\text{La}_2(\text{WO}_4)_4$ * (338)

XPS valence bands of La, Ce and Gd and their aluminium alloys* (598)

TITLE INDEX

laboratories

Research laboratories annual report 1975* (566)

Laboratory

Experience of the Central Control Laboratory (CCL) in accounting for and controlling nuclear material in Czechoslovakia* (349)

Argonne National Laboratory* (692)

Nuclear spectroscopy. Laboratory of chemistry and radiochemistry* (723)

LaCl

^{139}La ENDOR of $\text{LaCl}_3 \cdot \text{Gd}^{3+}$ * (130)

Investigation into absorption infrared spectra of vapors under ScBr_3 , YCl_3 , LaCl_3 , GdCl_3 , ErCl_3 , LuCl_3 by means of isolation method in rare gas matrix* (497)

LaF

Study of chemical interaction in $\text{Pb}_3\text{AlF}_6\text{-LaF}_6\text{-Ca}_3\text{AlF}_6$ system* (26)

Phases of variable composition with LaF_3 structure in $\text{MP}_2\text{-(Y,Ln)F}_3$ systems. 1. $\text{SrF}_2\text{-(Y,Ln)F}_3$ systems (thermal characteristics, formation of berthollides)* (593)

lambda

Entropy change for the lambda-type transition in nonstoichiometric $\text{U}_{4-y}\text{O}_{9-y}$ * (260)

Lambda-type heat capacity anomalies in U_3O_8 * (262)

lanthanide

Comparison of the thermodynamic properties and high temperature chemical behavior of lanthanide and actinide oxides* (2)

Preparation and some properties of actinide(III) and lanthanide(III) oxide halides, and the enthalpies of formation of $\text{PuBr}_3(\text{s})$ and $\text{PuI}_3(\text{s})$ * (82)

Interaction of lanthanide ions with fluorine-ions* (329)

Thermal decomposition of intermetallic compounds of platinum with alkaline earth and lanthanide metals at high temperatures* (399)

Intensities of lanthanide f-f transitions* (492)

Thermodynamics of the lanthanide halides. II. Heat capacities and Schottky anomalies of SmCl_3 , EuCl_3 and GdCl_3 from 50 to 350 K* (594)

lanthanides

Tetrad effect in the extraction of the lanthanides with neutral oxygen-containing compounds* (426)

Use of oscillator strengths of "supersensitive" ion transfers of lanthanides for calculation of some parameters determining the intensity of their absorption spectra* (511)

lanthanoid

Electronic structure of highly symmetric lanthanoid and actinoid compounds. VI. The electronic Raman spectrum of $\text{Cs}_2\text{NaCe(III)Cl}_6$ * (678)

lanthanoids

Electronic structure of highly symmetric complexes of lanthanoids and actinoids. Pt. 3. 'Electronic' Raman spectrum of $\text{Cs}_2\text{U(IV)Cl}_6$ * (13)

Electronic structure of highly symmetric complexes of lanthanoids and actinoids. Pt. 4. The electronic Raman spectrum of $\text{Cs}_2\text{NaEu(III)Cl}_6$ * (14)

TITLE INDEX

lanthanum

Study of lanthanum, yttrium and erbium oxychlorides in melts of alkali metal chlorides* (196)
 Lanthanum chloride interaction with magnesium oxide* (493)
 Interaction of potassium tetrafluoroborate with fluorides of potassium, lanthanum, neodymium in melt* (530)
 Deoxidation thermodynamics of cerium and lanthanum in liquid iron* (654)

LaS

Mass spectrometric determination of dissociation energies of high-temperature molecules: ScS, YS, LaS, ZrS, UO* (597)

Lattice

Lattice-dynamical calculation of the thermodynamic properties of NaCl, KCl and RbCl* (231)
 Thermal lattice dilatation of some transition metal compounds* (508)

law

Frequency spectrum and scattering law for liquid copper. A molecular dynamics investigation* (567)

lead

Density and temperature dependence of the surface tension of molten bismuth, lead, and bismuth-lead alloys* (1)
 Some observations on the compatibility of structural materials with molten lead* (25)
 Pulse NMR in solids: chemical shift, lead fluoride, and thorium hydride* (363)
 Activities of lead and bismuth in their molten alloys* (420)
 Thermophysical properties of the ternary liquid-metal system tin--lead--cadmium* (605)

levitation

Heat capacities and heat content of molten cerium by levitation calorimetry* (355)

Li

Phase equilibria in Li--Na--K carbonate/aluminate systems* (77)
 Equilibrium between Li-K-Te alloys and LiCl-KCl molten eutectic mixture* (92)
 Phase diagrams of Li_2MoO_4 -- ZnMoO_4 , K_2MoO_4 -- ZnMoO_4 and K_2WO_4 -- ZnWO_4 systems* (160)
 Gases and carbon in metals (thermodynamics, kinetics, and properties). I. Alkali metals, alkaline earth metals, light metals (Li, Na, K, Rb, Cs; Ca, Sr, Ba; Be, Mg, Al)* (192)
 Static structure factor of liquid Li* (273)
 Some thermodynamic properties of dilute solutions of actinide chlorides in (Li-K)Cl and in (Na-K)Cl eutectics* (406)
 Thermodynamic properties of dilute solutions of ThCl_4 in (Li-K)Cl and (Na-K) eutectics* (407)
 Lithium nitride (Li_3N): standard enthalpy of formation by solution calorimetry* (467)
 Phase diagrams in the pseudobinary systems of Li_2BeF_4 , alkali fluoroborates and alkali fluorides* (473)
 Solubility of nitrogen in liquid lithium and thermal decomposition of solid Li_3N * (702)

TITLE INDEX

LiBr

Phase diagrams of LiCl, CsCl, CaCl and LiBr, CsBr, CaBr ternary systems* (256)

LiBr, RbBr, CsBr ternary system* (268)

LiCl

Study of the Ni-Ni (II) equilibrium exchange, by radiotracer techniques in fused LiCl-KCl eutectic* (68)

Interaction of the lithium-tin alloys with the LiCl-KCl-CsCl melted eutectic mixture* (91)

Equilibrium between Li-K-Te alloys and LiCl-KCl molten eutectic mixture* (92)

Phase diagram of system LiCl--KCl--CaCrO₄* (116)

Phase diagram of the ThCl₄-LiCl-PbCl₂ system* (143)

Phase diagrams of LiCl, CsCl, CaCl and LiBr, CsBr, CaBr ternary systems* (256)

Structure and anodic discharge behavior of lithium-boron alloys in the LiCl-KCl eutectic melt.* (274)

Chronopotentiometry of U(III) and U(IV) in molten LiCl-KCl eutectic* (292)

Electrochemical behavior of U(III) and U(IV) in molten LiCl-KCl eutectic* (293)

Comparison of the tendencies of complex formation of various metal ions in fused LiF-NaF-KF (eut.) and in fused LiCl-KCl (eut.)* (314)

Corrosion of liquid alloys of zinc with yttrium and with rare earth elements in LiCl--KCl melts* (335)

Study of kinetics of monovalent indium ions formation in the In₂O₃-InCl₃-LiCl-H₂O system* (341)

Quantitative study of the chemical equilibria in solution in molten LiCl-KCl (70-30 mol.%) by absorption spectrophotometry.

Application to neptunium* (385)

Determination of the Np(IV)--Np(III) normal potential in fused LiCl-CsCl (55-45%) by absorption spectrophotometry* (386)

Intensity-voltage curves of hydrogen diffusion electrodes in eutectic LiCl--KCl* (507)

Preparation of thorium metal by electrolysis of fused salt ThCl₄-LiCl+NaCl+KCl(eut) system* (703)

LiF

Optical absorption spectrum of UO₂²⁺ doped LiF* (30)

Measurement of mass transfer coefficients in a mechanically agitated, nondispersing contactor operating with a molten mixture of LiF--BeF₂--ThF₄ and molten bismuth* (79)

LiF-FeF₃ system* (270)

Fusibility in K₂Nb(Ta)F₇-LiF-NaF systems* (298)

Comparison of the tendencies of complex formation of various metal ions in fused LiF-NaF-KF (eut.) and in fused LiCl-KCl (eut.)* (314)

Electric conductivity of melts of the LiF-misch metal fluoride system* (461)

Misch metal solubility in melt of LiF-BaF₂ system* (462)

Heat transfer measurements in a forced convection loop with two molten-fluoride salts: LiF--BeF₂--ThF₂--UF₄ and eutectic NaBF₄--NaF* (585)

TITLE INDEX

- life
Half-life of ^{242}Pu from precise low-temperature heat capacity measurements* (487)
- ligands
Effects of secondary ligands on the electronic structure of uranyls* (684)
- light
Electron spectrum of light alkali metals* (161)
Anomalous quasielastic light scattering from metal oxide glasses* (183)
Gases and carbon in metals (thermodynamics, kinetics, and properties). I. Alkali metals, alkaline earth metals, light metals (Li, Na, K, Rb, Cs; Ca, Sr, Ba; Be, Mg, Al)* (192)
MATERO: Version 09. A handbook of materials properties for use in the analysis of light water reactor fuel rod behavior* (387)
Collection of gamma spectra data of irradiated light water moderated reactor spent fuel and study of the applicability of the methods for fuel identification. Final report for the period 1 February 1972--October 1975* (453)
- LiH
Lines in the spectrum of ^{7}LiH (4728--5298 Å)* (374)
- LiIO
Constitution equilibrium diagram of the $\text{LiNO}_3\text{--LiIO}_3\text{--NaNO}_3\text{--NaIO}_3\text{--KNO}_3\text{--KIO}_3$ system* (665)
- limited
Lithium anode limited-cycle battery investigation. Quarterly technical progress report No. 5* (37)
Lithium anode limited-cycle battery investigation. Quarterly technical progress report No. 3* (38)
Practice of experiment planning in studying limited regions of multicomponent system* (546)
- line
Separation of alkaline earth from alkali metal fission products in the SOLAR on-line mass spectrometer* (534)
Pressure and entropy calculations for liquid Na, K and Rb along the melting line* (584)
- lines
Calculation of thermodynamic activity and liquidus lines of components in binary systems of plutonium trichloride with alkali and alkaline earth metal chlorides* (147)
Lines in the spectrum of ^{7}LiH (4728--5298 Å)* (374)
- LiNO
 $\text{LiNO}_3\text{--CsNO}_3\text{--Ca(NO}_3)_2$ and $\text{NaNO}_3\text{--CsNO}_3\text{--Ca(NO}_3)_2$ system* (255)
Solubility of $\text{Sr(NO}_3)_2$ in the $\text{LiNO}_3\text{--KNO}_3$ melt of variable composition* (351)
Constitution equilibrium diagram of the $\text{LiNO}_3\text{--LiIO}_3\text{--NaNO}_3\text{--NaIO}_3\text{--KNO}_3\text{--KIO}_3$ system* (665)
- liquid
Thermodynamic properties of the liquid La--Mg alloys* (3)
Transport of hydrogen in liquid lithium* (10)
Hydrogen solubility in liquid cesium* (22)
Thermodiffusion measurements in liquid alloys and electron transport heat calculations in the case of almost-free electrons* (33)

TITLE INDEX

liquid

- Reactions of liquid sodium with transition-metal oxides. VIII. The oxides of chromium-(III), -(IV), and -(VI) and disodium chromium(VI) tetraoxide* (35)
- Proceedings of the international conference on liquid technology in energy production* (56)
- Characterisation of the chemical activities of non-metals dissolved in liquid alkali metals* (63)
- Characterisation of the chemical activities of non-metals dissolved in liquid alkali metals* (64)
- Environmental sensitivity of structural metals: liquid metal embrittlement. Annual technical progress report No. 2, 25 Jun 1970--31 May 1971* (95)
- Nonisothermal flow of liquid metals over the surface of solid ones* (101)
- Proceedings of the international conference on liquid technology in energy production* (103)
- Kinetics of reaction of liquid lithium with oxygen and nitrogen* (106)
- Physical models of mass transport of iron and nickel in liquid sodium systems* (128)
- Determination of vibrational contribution into excessive mixing entropy of liquid amalgams. Alkali metal amalgams* (138)
- Solubility of sodium iodide in liquid sodium between 140 and 500°C* (154)
- Molybdenum and tungsten interaction with liquid aluminium* (156)
- Structure factor of liquid aluminum and calculation of transport coefficients* (159)
- Formation enthalpy of liquid tin-vanadium alloys* (166)
- Thermodynamic perturbation theory for the surface tension and ion density profile of a liquid metal* (167)
- Discrete variation of the structure and texture of liquid metals formed by components with discrete packing* (177)
- Atomic diffusion in alloys exposed to liquid sodium* (181)
- Solubility and activity of oxygen in liquid indium and copper-indium alloys* (182)
- Magnetic susceptibility and electronic structure in metallic and non-metallic liquid cesium-gold and cesium-antimony alloys* (190)
- Self-diffusion in liquid metals* (198)
- Perturbation treatment of the Percus-Yevick equation and structures of liquid magnesium and cadmium* (210)
- Environmental sensitivity of structural metals: liquid metal embrittlement. Annual technical progress report No. 4, 1 Jun 1972--31 May 1973* (211)
- Environmental sensitivity of structural metals: liquid metal embrittlement. Annual technical progress report No. 2, 25 Jun 1970--31 May 1971* (212)
- Negative temperature coefficients of electrical resistivity: the divalent liquid metals Eu, Yb and Ba* (224)
- Entropies of liquid rare earth metals* (230)
- Thermodynamic properties of solid and liquid metals and ceramics* (243)
- Specific heat of liquid metals and ceramics* (244)
- Liquid metals, 1976* (247)
- Static structure factor of liquid caesium* (248)

TITLE INDEX

liquid

- Structure factor of liquid caesium and sodium-caesium alloys* (249)
- Structure factors of liquid alkali metals* (250)
- Static structure factor of liquid Li* (273)
- Hydrogen solubility in liquid zirconium* (287)
- Report of task force on alternate structural materials for Liquid Metal Fast Breeder Reactors* (309)
- Effect of various additions on the temperature of structural transformation in liquid iron* (313)
- Study of mass transfer by liquid sodium in closed loop systems. Sodium technology development programme* (324)
- Corrosion of liquid alloys of zinc with yttrium and with rare earth elements in LiCl--KCl melts* (335)
- Thermodynamic properties of Cd-Zn-Hg liquid alloys* (340)
- Solubility of oxygen in liquid niobium* (342)
- Approximation of tabulated thermodynamic data for purposes of optimization of two-phase liquid-metal power cycles* (343)
- Electron contribution to the liquid metal viscosity* (346)
- Standard free energy of formation of liquid $\text{Na}_{20}\text{CrO}_{40}$, 1100⁰ K to 1220⁰ K* (375)
- Thermodynamic properties of diluted sodium solutions in liquid indium* (388)
- Speed of ultrasound and the thermophysical properties of the liquid metals Sn, Pb, Cd and of their binary alloys Pb:Sn and Pb:Cd* (403)
- Solution behavior of hydrogen isotopes and other non-metallic elements in liquid lithium* (404)
- Physical research on liquid-metal systems. Annual report, July 1975--June 1976* (405)
- Densities of electronic states in liquid transition metals* (422)
- Heat of formation of solid and liquid alloys of transition metals* (425)
- Current yield during separation of lithium on a liquid bismuth cathode* (430)
- Periodically variable wall temperature effect on heat transfer in liquid metals* (437)
- Calculation of thermodynamic activity of components in strontium liquid alloys* (446)
- Vapor pressure studies over liquid uranium oxide and uranium plutonium oxide up to 5,000 K* (474)
- Advances in vapour pressure studies over liquid uranium plutonium oxides up to 5000K* (476)
- Vapour pressure studies over liquid uranium plutonium oxides up to 5,000 K* (478)
- KKR analysis of the Anderson-McMillan theory of electronic structure of liquid metals. Technical report* (485)
- Liquid metals, 1976* (486)
- New description of liquid metal heat transfer in closed conduits* (502)
- Thermodynamic studies of gallium--indium liquid alloys by solid state electrochemistry with oxide electrolytes* (513)
- Thermodynamic properties of liquid thorium-gallium alloys* (522)
- Thermodynamic investigation of liquid alloys in the system Cd-In-Tl* (523)

TITLE INDEX

liquid

- Contribution to the problem of the thermodynamic properties and the structure of liquid Cd-In alloys* (524)
- Theory for the derivation of a single hard-sphere potential describing the overall molecular interactions in binary liquid metal alloys as a function of temperature and composition* (525)
- Investigation of liquid-steam equilibrium in system with diluted solutions of metal fluorides in uranium hexafluoride* (526)
- Recent developments in the chemistry of solutions of salts in liquid alkali metals* (528)
- Reaction rates and electrical resistivities of the hydrogen isotopes with, and their solubilities in, liquid lithium* (529)
- Nitrogen solubility in liquid nickel--niobium alloys* (532)
- Nuclear quadrupole relaxation and viscosity in liquid metals* (564)
- Frequency spectrum and scattering law for liquid copper. A molecular dynamics investigation* (567)
- Thermal expansion of metals over the entire liquid range* (573)
- Equation of state and transport properties of uranium and plutonium carbides in the liquid region* (577)
- Equation of state and transport properties of uranium and plutonium nitrides in the liquid region* (578)
- Densities of liquid carbides and nitrides of uranium and plutonium* (579)
- Liquid metals, 1976* (584)
- Proceedings of the international conference on liquid technology in energy production* (590)
- Density and enthalpy of liquid alkaline metals at pressures up to 1500 bar* (599)
- Thermophysical properties of the ternary liquid-metal system tin--lead--cadmium* (605)
- Thermodynamic properties of the ternary Sn-Pb-Cd liquid metal system* (606)
- Application of the hard-sphere model to thermodynamic properties of liquid 3d transition metals* (615)
- Enthalpy and heat capacity of liquid plutonium and uranium* (619)
- Enthalpy and heat capacity of liquid plutonium and uranium* (620)
- Experimental study of the temperature dependence on energy exchanges in liquid metals* (621)
- Viscosities of cesium vapor to 1,620 K and of liquid gallium to 1,800 K* (622)
- Thermodynamic properties of sulphur in liquid antimony* (642)
- Influence of the technology of the alloy fabrication on its properties in a liquid state* (643)
- Diffusion in liquid metal systems. Final report* (651)
- Deoxidation thermodynamics of cerium and lanthanum in liquid iron* (654)
- Investigation of particles interaction in liquid metals and alloys* (661)
- Viscosities of cesium vapor to 1620 K and of liquid gallium to 1800 K* (681)
- Liquid metals, 1976* (686)
- Structures of 3d-transition metals in the liquid state* (687)
- Structure of rare earth metals in the liquid state* (688)
- Structure of liquid Ce-Ni alloys* (689)
- Solubility of lithium oxide in liquid lithium* (701)

TITLE INDEX

liquid

Solubility of nitrogen in liquid lithium and thermal decomposition of solid Li_3N * (702)
Electronic states densities from structure and thermodynamic data for liquid transition metals* (705)

liquids

Structural study of liquids with strong short-range correlation in the atomic distribution* (609)
Structural study of liquids with strong short-range correlation in the atomic distribution* (653)

liquidus

Calculation of thermodynamic activity and liquidus lines of components in binary systems of plutonium trichloride with alkali and alkaline earth metal chlorides* (147)
Interaction in AgCl-CsCl , CsCl-KCl systems and liquidus surface of AgCl-CsCl-KCl ternary system* (659)
Investigation into properties of liquidus curves of binary salt system melts where chemical compounds formation. 2. Systems of monovalent and bi-valent cations with common anion* (708)

lithium

Transport of hydrogen in liquid lithium* (10)
Lithium anode limited-cycle battery investigation. Quarterly technical progress report No. 5* (37)
Lithium anode limited-cycle battery investigation. Quarterly technical progress report No. 3* (38)
Investigation into neodymium chloride state in equilibrium with metal in eutectic melt of potassium- and lithium chlorides* (40)
Studies of the electronic structure of simple metals and their alloys with the aid of Compton scattering and positron annihilation using lithium, magnesium, lithium-magnesium and aluminium-zinc as examples* (48)
Ternary systems from lithium, sodium, strontium and barium chlorides* (86)
Quaternary mutual system from fluorides and chlorides of lithium, sodium and strontium* (89)
Interaction of the lithium-tin alloys with the LiCl-KCl-CsCl melted eutectic mixture* (91)
Kinetics of reaction of liquid lithium with oxygen and nitrogen* (106)
Lithium-silver chloride secondary battery investigation. Quarterly technical progress report No. 2* (111)
Lithium-silver chloride secondary battery investigation. Final report, December 1963--November 1964* (118)
Density, surface tension and viscosity of melts of uranium trichloride-lithium chloride* (142)
Titanium dioxide solubility in melts of lithium- and alkaline earth metal chlorides* (264)
Structure and anodic discharge behavior of lithium-boron alloys in the LiCl-KCl eutectic melt.* (274)
Structure of lithium-boron alloys and their performance as potential anodes for high power molten salt batteries. Progress report, 1 Aug 1974--30 Jun 1975* (275)
Phase equilibrium studies of lithium halide-containing electrolytes* (282)
Preparation of lithium octafluorohafnates* (330)

TITLE INDEX

lithium

Phase diagram of the ternary system of lithium-, rubidium-, and cesium nitrates* (377)
 Lithium--nickel fluoride secondary battery investigation. Quarterly technical progress report No. 1, 27 Apr--27 Jul 1966* (381)
 Solution behavior of hydrogen isotopes and other non-metallic elements in liquid lithium* (404)
 Current yield during separation of lithium on a liquid bismuth cathode* (430)
 Dispersion of metallic lithium in various molten salts* (452)
 Lithium nitride (Li_3N): standard enthalpy of formation by solution calorimetry* (467)
 Electrochemical reduction of molybdenum oxychloride in molten lithium chloride - potassium chloride eutectic* (515)
 Reaction rates and electrical resistivities of the hydrogen isotopes with, and their solubilities in, liquid lithium* (529)
 Electromotive force measurements in molten lithium--magnesium alloys* (551)
 Tritium sorption in lithium--bismuth and lithium--aluminum alloys* (614)
 Anodic behavior of cadmium and cadmium-zinc alloys in eutectic melt of potassium- and lithium chlorides* (675)
 Solubility of lithium oxide in liquid lithium* (701)
 Solubility of nitrogen in liquid lithium and thermal decomposition of solid Li_3N * (702)
 Lithium alloys--chlorine secondary battery. I. Basic studies on a lithium alloys--chlorine secondary battery using a molten salt as the electrolyte* (704)

LMFBR

LMFBR Fuel Reprocessing Program progress report for period April 1 to June 30, 1976* (94)
 PuO_2 dissolution problem for LWR plutonium recycle and LMFBR fuels: fabrication and reprocessing problems and their resolution* (455)
 Analysis of LMFBR explosion model experiments by means of the SURECOM-II code* (601)
 Summary of U.S. LMFBR programs on high temperature structural design and associated materials testing* (731)

LMFBRs

Preliminary assessments of carbide fuel pins during mild overpower transients in LMFBRs* (456)

Ln

Vibration spectra of rare earth oxytungstates- Ln_2WO_6 composition* (185)
 Phases of variable composition with LaF_3 structure in MF_2 -(Y,Ln) F_3 systems. 1. SrF_2 -(Y,Ln) F_3 systems (thermal characteristics, formation of berthollides)* (593)

LnF

Structural changes in the series LnFeO_3 and LnF_3 * (197)

 LnFeO

Structural changes in the series LnFeO_3 and LnF_3 * (197)

TITLE INDEX

LnTiNbO

Physicochemical investigation of rare earth titanoniobates, LnTiNbO₆* (173)

loop

Study of mass transfer by liquid sodium in closed loop systems. Sodium technology development programme* (324)
Heat transfer measurements in a forced convection loop with two molten-fluoride salts: LiF--BeF₂--ThF₂--UF₄ and eutectic NaF--NaF* (585)

low

Preparation of uranium-233 with low uranium-232 content* (232)
Interaction parameters in the Zn--Pb--Sn system at low zinc concentration* (436)
Half-life of ²⁴²Pu from precise low-temperature heat capacity measurements* (487)
Anomalous thermal property behaviour of uranium at low temperatures* (560)

Lu

Phase equilibria and thermodynamic data for the ternary UO₂--UO₃--REO_{1.5} (Ho, Tm, Lu) systems* (694)

Lu]

Comparative study of systems of heavier rare-earth metals with hydrogen [Gd, Tb, Dy, Lu]* (418)

LuCl

Investigation into absorption infrared spectra of vapors under ScBr₃, YCl₃, LaCl₃, GdCl₃, ErCl₃, LuCl₃ by means of isolation method in rare gas matrix* (497)

luminescence

Effect of luminescence centers on optical absorption, excitation and fluorescence spectra of CsCl:Ph* (491)

lutetium

Interaction of rubidium and cesium fluorides with lutetium trifluorides* (444)

LWR

PuO₂ dissolution problem for LWR plutonium recycle and LMFBF fuels: fabrication and reprocessing problems and their resolution* (455)
Phase analysis on simulated LWR fuel rods and its evaluation with the aid of constitution studies in the system uranium-zirconium-oxygen* (509)
Phase analysis at simulated LWR fuel rods and their interpretation by constitution investigations in the uranium-zirconium-oxygen system* (510)

M

Optical spectra of the difluoride, dichloride, and trichloride ions in the matrix-isolated M⁺ + F₂²⁺-, M⁺ + Cl₂²⁺-, and M⁺ + Cl₃³⁺- species* (16)
Optical spectra of the dibromide and diiodide ions in the matrix-isolated M⁺ + Br₂²⁺- and M⁺ + I₂²⁺- species* (17)
Vibrational spectra and force constants of the elpasolites Cs₂KMF₆ (M=Sc, Y, La, Gd, Yb)* (41)

TITLE INDEX

M

Study of the compounds $M\text{UO}_2\text{F}_5$ ($M = \text{K}, \text{Rb}, \text{Cs}, \text{NH}_4$) by i.r. absorption and Raman diffusion spectrophotometry. III. Effect of cation on vibration frequency of the ion $\text{UO}_2\text{F}_5^{3-}$ (127)
 Solubility of rare earth and yttrium oxochlorides in molten salts - MCl_2 ($M = \text{Mg}, \text{Ca}, \text{Sr}, \text{Ba}$) (495)
 Preparational, structural and vibrational study of $M_2\text{UO}_5\text{Cl}_4$, $2\text{H}_2\text{O}$ ($M = \text{Rb}, \text{Cs}$). Detection of a tetranuclear anion $[(\text{UO}_2)_4\text{O}_{10}\text{Cl}_8(\text{H}_2\text{O})_2]^{4-}$ (498)
 Degree of thermal dissociation of MCaCl_3 ($M = \text{K}, \text{Rb}, \text{Cs}$) complexes during melting* (624)

machining

Electrochemical machining of the molybdenum VM-1 alloy* (129)

magnesium

Studies of the electronic structure of simple metals and their alloys with the aid of Compton scattering and positron annihilation using lithium, magnesium, lithium-magnesium and aluminium-zinc as examples* (48)
 Perturbation treatment of the Percus-Yevick equation and structures of liquid magnesium and cadmium* (210)
 Solubility of magnesium oxide in alkali metal chloride melt* (265)
 Solubility of magnesium oxide in calcium oxide--calcium chloride mixtures* (352)
 Some regularities of the corrosion behavior of binary magnesium alloys* (439)
 Lanthanum chloride interaction with magnesium oxide* (493)
 Interaction of neodymium- and yttrium chlorides with magnesium oxide* (494)
 Electromotive force measurements in molten lithium--magnesium alloys* (551)

Magnetic

Magnetic susceptibility and electronic structure in metallic and non-metallic liquid cesium-gold and cesium-antimony alloys* (190)
 Electronic and magnetic properties of rare earth- Sn_3 compounds from ^{119}Sn Moessbauer spectroscopy* (559)

manganese

Investigation into interaction of indium tribromide with dibromides of manganese, cobalt and nickel in melt* (153)

Mar

Research on a high-energy nonaqueous battery system. Quarterly progress report No. 1, 1 Dec 1965--1 Mar 1966* (720)

March

Proceedings of the symposium and workshop on advanced battery research and design, March 22--24, 1976* (269)
 Physical--chemical studies of transuranium elements. Progress report, April 1, 1976--March 31, 1977* (501)

mass

Measurement of mass transfer coefficients in a mechanically agitated, nondispersing contactor operating with a molten mixture of LiF -- BeF_2 -- ThF_4 and molten bismuth* (79)
 Knudsen cell--mass spectrometer studies of cesium--uranium interactions* (120)

TITLE INDEX

mass

- Physical models of mass transport of iron and nickel in liquid sodium systems* (128)
- Mass spectra and vapor composition of niobium pentafluoride* (216)
- Determination and correlation of mass transfer coefficients in a stirred cell* (236)
- Molecular vapor composition and mass spectra in $KF-ZrF_4$ systems* (294)
- Study of mass transfer by liquid sodium in closed loop systems. Sodium technology development programme* (324)
- Mass transport phenomena in ceramics* (401)
- Abstracts of papers and communications submitted to the 5. all-union heat and mass transfer conference* (437)
- Separation of alkaline earth from alkali metal fission products in the SOLAR on-line mass spectrometer* (534)
- Mass spectrometric determination of dissociation energies of high-temperature molecules: ScS , YS , LaS , ZrS , UO * (597)
- Standard methods for chemical, mass spectrometric, and spectrochemical analysis of nuclear-grade uranium dioxide powders and pellets* (711)
- Standard methods for chemical, mass spectrometric, and spectrochemical analysis of nuclear-grade plutonium dioxide powders and pellets* (712)
- Standard methods for chemical, mass spectrometric, and spectrochemical analysis of nuclear-grade mixed oxides (U, Pu) $O_{2.5}$ * (713)
- Standard methods for chemical, mass spectrometric, spectrochemical, nuclear, and radiochemical analysis of nuclear-grade plutonium metal* (714)
- Standard method of test for atom percent fission in uranium and plutonium fuel (mass spectrometric method)* (715)

material

- Accurate procedure to safeguard the fissile material content of input and output solutions of reprocessing plants* (131)
- Ceramic-base heat resisting constructional material* (348)
- Experience of the Central Control Laboratory (CCL) in accounting for and controlling nuclear material in Czechoslovakia* (349)

materials

- Stable inorganic matrix materials for high temperature batteries. Final report No. 1, 2 June 1967--2 February 1968* (23)
- Some observations on the compatibility of structural materials with molten lead* (25)
- Mechanical properties test data for structural materials. Quarterly progress report for period ending April 30, 1977* (57)
- Compatibility of molten rare earth metals with refractory materials* (115)
- Safeguarding nuclear materials. Vol. 2* (131)
- Thermal expansion of molten materials by the gamma attenuation technique* (152)
- Chemical effects of composition changes in irradiated oxide fuel materials. II. Fission product segregation and chemical equilibria* (169)
- Structure of the biexciton molecule and optical transition selection rules in zincblende and wurtzite materials* (186)

TITLE INDEX

materials

Chemical Engineering Division fuels and materials chemistry. Annual report, July 1975--June 1976* (281)
 Radiolytic gas generation in plutonium contaminated waste materials* (301)
 Report of task force on alternate structural materials for Liquid Metal Fast Breeder Reactors* (309)
 Safeguarding nuclear materials. Vol. 2* (349)
 Thermophysical properties of UO_2 fuel materials* (366)
 MATPRO: Version 09. A handbook of materials properties for use in the analysis of light water reactor fuel rod behavior* (387)
 Mechanical properties test data for structural materials. Quarterly progress report for period ending April 30, 1977* (471)
 Investigation on the resistance of high-refractory materials to chromium melt* (531)
 Properties of structural materials for sodium-cooled fast breeder reactor* (580)
 Refractory materials resistance to molten barium alloy* (628)
 XPS studies of actinide materials* (662)
 Analytical methods for fissionable materials in the nuclear fuel cycle. Progress report, July 1, 1975--September 30, 1976* (690)
 Mechanical properties test data for structural materials. Quarterly progress report for period ending April 30, 1977* (692)
 Summary of U.S. LMFBR programs on high temperature structural design and associated materials testing* (731)

MATPRO

MATPRO: Version 09. A handbook of materials properties for use in the analysis of light water reactor fuel rod behavior* (387)

matrices

Identification of UN in Ar matrices* (219)
 Flame emission and atomic absorption spectrometry. III. Elements and matrices* (319)

matrix

Optical spectra of the difluoride, dichloride, and trichloride ions in the matrix-isolated $M^{sup} + F^{sub 2}^{sup} -$, $M^{sup} + Cl^{sub 2}^{sup} -$, and $M^{sup} + Cl^{sub 3}^{sup} -$ species* (16)
 Optical spectra of the dibromide and diiodide ions in the matrix-isolated $M^{sup} + Br^{sub 2}^{sup} -$ and $M^{sup} + I^{sub 2}^{sup} -$ species* (17)
 Stable inorganic matrix materials for high temperature batteries. Final report No. 1, 2 June 1967--2 February 1968* (23)
 Investigation into absorption infrared spectra of vapors under $ScBr^{sub 3}$, $YCl^{sub 3}$, $LaCl^{sub 3}$, $GdCl^{sub 3}$, $ErCl^{sub 3}$, $LuCl^{sub 3}$ by means of isolation method in rare gas matrix* (497)

M₂CaCl

Degree of thermal dissociation of $M_2CaCl^{sub 3}$ (M-K, Rb, Cs) complexes during melting* (624)

MCl

Study of saturated vapor pressure above salt melts in $CoCl^{sub 2}$ -MCl systems* (98)
 Solubility of rare earth and yttrium oxochlorides in molten salts - $MCl^{sub 2}$ (M-Mg, Ca, Sr, Ba)* (495)

McMillan

KKR analysis of the Anderson-McMillan theory of electronic structure of liquid metals. Technical report* (485)

TITLE INDEX

Me

$\text{Me}_{\text{sub } 2}\text{ZrCl}_{\text{sub } 6}$ -- $\text{H}_{\text{sub } 2}$ (Me-Na, K, Rb, Cs) system* (45)

measured

Calculation of the oxygen potential of $(\text{U}, \text{Pu})\text{O}_{\text{sub } 2-x}$ with defect models and measured oxygen potential data in the temperature range from 1,000 to 1,700 K* (74)

Measurement

Measurement of mass transfer coefficients in a mechanically agitated, nondispersing contactor operating with a molten mixture of LiF -- $\text{BeF}_{\text{sub } 2}$ -- $\text{ThF}_{\text{sub } 4}$ and molten bismuth* (79)

Measurement of fluorescence of rare earths in molten salts ($\text{Nd}_{\text{sup } 3+}$ in $\text{NaNO}_{\text{sub } 3}$ -- $\text{KNO}_{\text{sub } 3}$ eutectic)* (233)

Thermodynamic study of the U-O-C system by measurement of the carbon monoxide pressures at equilibrium* (235)

Phase transition in doped $\text{U}_{\text{sub } 4}\text{O}_{\text{sub } (9-y)}$. I. Heat capacity measurement* (261)

Measurement of the cell voltage of $\text{ThO}_{\text{sub } 2}$ -- $\text{Y}_{\text{sub } 2}\text{O}_{\text{sub } 3}$ --solid electrolyte cells in the range of mixed conduction* (535)

Measurement of interfacial energies in the $\text{UO}_{\text{sub } 2}$ -- $\text{Gd}_{\text{sub } 2}\text{O}_{\text{sub } 3}$ system* (697)

measurements

Thermodiffusion measurements in liquid alloys and electron transport heat calculations in the case of almost-free electrons* (33)

Investigations on radioactive fission product correlations: gamma spectrometry measurements on spent fuel assemblies discharged from the Trino Vercellese reactor at the end of the 2nd irradiation cycle* (75)

High-temperature microcalorimetric measurements of partial molar enthalpy of solution, $\Delta H(\text{O}_{\text{sub } 2})$, in plutonium + oxygen* (107)

Some considerations in establishing thermodynamic activities from high temperature EMF measurements* (163)

Thermodynamic properties of Cr, Mn, Co and Ni dichlorides from emf measurements on cells with solid electrolytes* (200)

Extension of vapour pressure measurements of nuclear fuels $(\text{U}, \text{Pu})\text{O}_{\text{sub } 2}$ and $\text{UO}_{\text{sub } 2}$ to 7000 K for fast reactor safety analysis* (475)

Vapour pressure measurements to 7,000 K and equation of state of oxide fuels for fast reactor safety analysis. I. Extension of vapour pressure measurements of nuclear fuels $(\text{U}, \text{Pu})\text{O}_{\text{sub } 2}$ and $\text{UO}_{\text{sub } 2}$ to 7,000 K* (477)

Half-life of ^{242}Pu from precise low-temperature heat capacity measurements* (487)

Effective equation-of-state measurements on uranium dioxide* (537)

Electromotive force measurements in molten lithium--magnesium alloys* (551)

New apparatus for thermophysical measurements above 2500 K* (574)

Heat transfer measurements in a forced convection loop with two molten-fluoride salts: LiF -- $\text{BeF}_{\text{sub } 2}$ -- $\text{ThF}_{\text{sub } 2}$ -- $\text{UF}_{\text{sub } 4}$ and eutectic $\text{NaBF}_{\text{sub } 4}$ -- NaF * (585)

TITLE INDEX

measuring

Use of the SF-8 detection spectrophotometer for measuring absorption spectra of molten salts* (331)

Mechanical

Mechanical properties test data for structural materials.

Quarterly progress report for period ending April 30, 1977* (57)

Physical and mechanical properties of some uranium-containing carbides* (209)

Mechanical properties test data for structural materials.

Quarterly progress report for period ending April 30, 1977* (471)

Mechanical properties test data for structural materials.

Quarterly progress report for period ending April 30, 1977* (692)

mechanically

Measurement of mass transfer coefficients in a mechanically agitated, nondispersing contactor operating with a molten mixture of LiF -- BeF_2 -- ThF_4 and molten bismuth* (79)

mechanics

Structural mechanics in reactor technology. Vol. 1* (19)

Statistical mechanics of intermediate-valence rare earth compounds* (279)

Structural mechanics in reactor technology. Vol. 2* (601)

Mechanism

Mechanism of metals modifying* (108)

Kinetics and mechanism into preparation of samarium-, europium- and ytterbium dichlorides and their some properties* (253)

mechanisms

Basic mechanisms providing oxidation resistance in structural metals at high temperatures. II. The defect structure of nonstoichiometric rutile (TiO_2) and nonstoichiometric rutile containing 1 to 10 mole % Cb_2O_6 . Final report 21 Dec 1973-20 Dec 1974* (542)

MeCl

FeCl_3 -- MeCl systems* (435)

media

Nonaqueous electrolyte systems: ionic transport in nonaqueous media. Interim technical report No. 7* (246)

meeting

Summaries of reports of the all-union meeting on solid body chemistry. 3--5 June 1975. III. Synthesis, phase equilibrium and use of solid oxide compounds* (114)

Spring meeting, 26 and 27 April 1976, Amsterdam* (250)

Second specialist meeting on sodium/fuel interaction in fast reactors, Ispra, 21-23 November 1973. Sponsored by the Committee on the Safety of Nuclear Installations, OECD Nuclear Energy Agency, Paris, France* (272)

Specialists' meeting on the behaviour of water reactor fuel elements under accident conditions* (510)

Summaries of reports of the all-union meeting on solid body chemistry. 3--5 June 1975. III. Synthesis, phase equilibrium and use of solid oxide compounds* (641)

melt

Interaction of uranyl chloride with zirconium tetrachloride in the melt of sodium chloride and potassium chloride* (27)

TITLE INDEX

melt

- Investigation into neodymium chloride state in equilibrium with metal in eutectic melt of potassium- and lithium chlorides* (40)
- Investigation into interaction of indium tribromide with dibromides of manganese, cobalt and nickel in melt* (153)
- Solubility of magnesium oxide in alkali metal chloride melt* (265)
- Structure and anodic discharge behavior of lithium-boron alloys in the LiCl-KCl eutectic melt.* (274)
- Solubility of $\text{Sr(NO}_3)_2$ in the $\text{LiNO}_3\text{-KNO}_3$ melt of variable composition* (351)
- Influence of intermediate products on chlorination kinetics of vanadium oxides in salt melt* (356)
- Misch metal solubility in melt of LiF-BaF_2 system* (462)
- Interaction of potassium tetrafluoroborate with fluorides of potassium, lanthanum, neodymium in melt* (530)
- Investigation on the resistance of high-refractory materials to chromium melt* (531)
- Investigation into reactions of hexabromotellurates of potassium, cesium and thallium(1) in melt* (556)
- Anodic behavior of cadmium and cadmium-zinc alloys in eutectic melt of potassium- and lithium chlorides* (675)

meltability

- Density, viscosity, electric conductivity and meltability of $\text{Na}_3\text{AlF}_6\text{-La}_2\text{O}_3$ and $\text{Na}_3\text{AlF}_6\text{-Sm}_2\text{O}_3$ systems* (43)
- Meltability in system of $\text{K}_2\text{TaF}_7\text{-NaF-NaCl-KCl}$ * (299)
- Meltability in $\text{K}_2\text{NbF}_7\text{-NaF-NaCl-KCl}$ system* (300)
- Meltability diagram of RbCl-BeCl_2 system* (506)

melted

- Interaction of the lithium-tin alloys with the LiCl-KCl-CsCl melted eutectic mixture* (91)
- Adhesion properties of melted Al, Ga, In on solid titanium* (327)
- Structure of In_2Bi melted compound* (514)

melting

- Preparation of ^{147}Pm metal and the determination of the melting point and phase transformation temperatures* (18)
- Computational simulation of cladding motion after melting including resolidification processes* (19)
- Density, surface tension, and viscosity of uranium trichloride-sodium chloride melting* (141)
- Enthalpy of uranium-plutonium carbide from 298K to the melting point* (179)
- Production and properties of high-melting lower oxides* (191)
- Vacancies and melting curves of metals at high pressure* (213)
- EUTECTIC DATA: safety, hazards, corrosion, melting points, compositions, and bibliography* (276)
- Anomalous melting of f electron metals (with attention to Pu)* (318)
- Melting point of europium dichloride* (361)
- Some structural characteristics of rubidium and cesium in the vicinity of their melting points* (460)
- Thermodynamic analysis of melting point diagrams of systems consisting of alkali metal chlorides, thallium chloride and beryllium chloride* (505)

TITLE INDEX

melting

Pressure and entropy calculations for liquid Na, K and Rb along the melting line* (584)
 Investigation into outgassing from molybdenum during melting* (588)
 Degree of thermal dissociation of MCaCl_3 (M-K, Rb, Cs) complexes during melting* (624)

melts

Viscosity characteristics of the iron-chromium-nickel melts* (24)
 Investigation into thermodynamic properties of Ce-Mg melts* (39)
 Investigation into thermodynamic properties of FeCl_2 -CsCl system melts* (96)
 Study of saturated vapor pressure above salt melts in CoCl_2 - MCl systems* (98)
 Raman and i.r.-emission studies of some tungstate- and molybdate-containing melts* (102)
 Density, surface tension and viscosity of melts of uranium trichloride-lithium chloride* (142)
 Viscosity of UCl_3 - MgCl_2 binary system melts* (145)
 Atomic structure of melts of complex system with eutectic transformations* (155)
 Kinetic peculiarities of nitrogen dissolving in iron-chromium and iron-chromium-nickel melts* (172)
 Study of lanthanum, yttrium and erbium oxychlorides in melts of alkali metal chlorides* (196)
 Enthalpies of formation of melts of yttrium with aluminium* (201)
 Raman spectra of Al_2O_3 solutions in molten cryolite and other aluminum fluoride containing melts* (203)
 Simple Raman cell and furnace usable at temperatures higher than 1000°C for corrosive melts* (204)
 Electrochemical behavior of tellurium (4) in oxychloride melts* (221)
 Titanium dioxide solubility in melts of lithium- and alkaline earth metal chlorides* (264)
 Use of fluoride melts for separation of some fission products and americium from mixtures of different fluorides being the simulators of residues from fast reactor fuel fluorination* (307)
 Investigation into americium extraction process from residues of irradiated nuclear fuel after its fluorination by SiF_4 -NaF and SiF_4 - CaF_2 melts of eutectic composition* (308)
 Corrosion of liquid alloys of zinc with yttrium and with rare earth elements in LiCl - KCl melts* (335)
 Wetting of certain refractories of the system MgO - Al_2O_3 - ZrO_2 with metallic melts* (344)
 Properties of corium melts* (454)
 Electric conductivity of melts of the LiF-misch metal fluoride system* (461)
 NMR study on structural microinhomogeneity of fluoride-oxide melts* (463)
 Raman spectroscopic studies of yttrium (III) chloride-alkali metal chloride melts and of $\text{Cs}_2\text{NaYCl}_6$ and YCl_3 solid compounds* (489)
 Using molecular dynamics method for the evaluation of Fe-Ni and Al-Zn melts structure properties* (512)
 Friction in metal melts* (575)

TITLE INDEX

melts

Viscosity of melts of $KCl-KF-K_{2}ZrF_{6}$ ternary system* (576)
 Viscosity of binary of nitrate systems melts* (592)
 Interaction of plutonium trichloride in melts of ternary systems* (682)
 Investigation into properties of liquidus curves of binary salt system melts where chemical compounds formation. 2. Systems of monovalent and bi-bivalent cations with common anion* (708)

mercury

On interaction of tellurium tetrachloride with copper (1), mercury (2), bismuth and iron (3) chlorides* (336)

metal

Thermodynamic properties of diluted solutions of $MnCl_{2}$ in alkali metal chlorides* (6)
 Preparation of ^{147}Sm metal and the determination of the melting point and phase transformation temperatures* (18)
 Reactions of liquid sodium with transition-metal oxides. VIII. The oxides of chromium-(III), -(IV), and -(VI) and disodium chromium(VI) tetraoxide* (35)
 Investigation into neodymium chloride state in equilibrium with metal in eutectic melt of potassium- and lithium chlorides* (40)
 Vibrational spectra of transition metal hexafluoride crystals. I. Orthorhombic MoF_6 , WF_6 , UF_6 neat crystals. Technical report* (51)
 Vibrational spectra of transition metal hexafluoride crystals. III. Exciton band structures of MoF_6 , WF_6 , and UF_6 . Technical report* (52)
 Structural study and properties of the alkali metal, nitrosyl, and ammonium hepta- and octafluorouranates(VI)* (66)
 Environmental sensitivity of structural metals: liquid metal embrittlement. Annual technical progress report No. 2, 25 Jun 1970--31 May 1971* (95)
 Saturation vapor pressure in iron chloride-alkaline earth metal chlorides systems* (97)
 Determination of vibrational contribution into excessive mixing entropy of liquid amalgams. Alkali metal amalgams* (138)
 Interaction of uranium trichloride with alkaline earth metal chlorides* (139)
 Calculation of thermodynamic activity and liquidus lines of components in binary systems of plutonium trichloride with alkali and alkaline earth metal chlorides* (147)
 Thermodynamic perturbation theory for the surface tension and ion density profile of a liquid metal* (167)
 Some properties of hydrogen fluoride compounds with group 5 metal pentafluorides* (170)
 Anomalous quasielastic light scattering from metal oxide glasses* (183)
 Electronic conductivity and polarization effects of solid $ThO_{2}-Y_{2}O_{3}$ electrolytes and of oxygen containing Va-metal electrodes* (193)
 Structure of $NO_{3}^{\cdot -}$ radical in molten monovalent metal nitrates* (195)
 Study of lanthanum, yttrium and erbium oxychlorides in melts of alkali metal chlorides* (196)

TITLE INDEX

metal

- Environmental sensitivity of structural metals: liquid metal embrittlement. Annual technical progress report No. 4, 1 Jun 1972--31 May 1973* (211)
- Environmental sensitivity of structural metals: liquid metal embrittlement. Annual technical progress report No. 2, 25 Jun 1970--31 May 1971* (212)
- Electron structure of rare-earth metal tetraborides* (215)
- Thermochemistry of gaseous metal oxides. Final report, 1 Apr 1973--31 Dec 1975* (239)
- Titanium dioxide solubility in melts of lithium- and alkaline earth metal chlorides* (264)
- Solubility of magnesium oxide in alkali metal chloride melt* (265)
- Reaction of americium oxides with threevalent metal oxides* (303)
- Report of task force on alternate structural materials for Liquid Metal Fast Breeder Reactors* (309)
- Comparison of the tendencies of complex formation of various metal ions in fused LiF-NaF-KF (eut.) and in fused LiCl-KCl (eut.)* (314)
- Interaction of atoms of rare earths-metal alloys* (320)
- Approximation of tabulated thermodynamic data for purposes of optimization of two-phase liquid-metal power cycles* (343)
- Electron contribution to the liquid metal viscosity* (346)
- Rare earth metal--metal halide systems. 19. Structural characterization of the reduced holmium chloride $\text{Ho}_{0.5}\text{Cl}_{1.5}$ * (378)
- Thermal decomposition of solid metal oxides* (395)
- Physical research on liquid-metal systems. Annual report, July 1975--June 1976* (405)
- Progress in the spectroscopy of rare earth and transition metal ions* (412)
- Pressure and composition of vapors over alkali metal tetrachloroferrates in chlorine atmosphere* (433)
- Electric conductivity of melts of the LiF-misch metal fluoride system* (461)
- Misch metal solubility in melt of LiF-BaF₂ system* (462)
- Reactivity of transition metal fluorides. IX. Protactinium pentafluoride and tetrafluoride* (465)
- Determination of surface tension of a metal with high vapor pressure* (472)
- Raman spectroscopic studies of yttrium (III) chloride--alkali metal chloride melts and of Cs₂NaYCl₆ and YCl₃ solid compounds* (489)
- New description of liquid metal heat transfer in closed conduits* (502)
- Thermodynamic analysis of melting point diagrams of systems consisting of alkali metal chlorides, thallium chloride and beryllium chloride* (505)
- Thermal lattice dilatation of some transition metal compounds* (508)
- Theory for the derivation of a single hard-sphere potential describing the overall molecular interactions in binary liquid metal alloys as a function of temperature and composition* (525)
- Investigation of liquid-steam equilibrium in system with diluted solutions of metal fluorides in uranium hexafluoride* (526)

TITLE INDEX

metal

Fundamental studies of metal fluorination reactions. Progress report, May 1, 1976--April 31, 1977* (533)
 Separation of alkaline earth from alkali metal fission products in the SOLAR on-line mass spectrometer* (534)
 Friction in metal melts* (575)
 Preparation and structure of transition metal fluorides $U^{sup}(II)M^{sup}(IV)F_{sub 6}[U^{sup}(II) : Cr-Zn; M^{sup}(IV) : Hf, Zr]^*$ (596)
 Thermophysical properties of the ternary liquid-metal system tin--lead--cadmium* (605)
 Thermodynamic properties of the ternary Sn-Pb-Cd liquid metal system* (606)
 Diffusion in liquid metal systems. Final report* (651)
 Isorhodate complexes of 3d-metal ions in molten salts* (672)
 Accurate nuclear parameters derived from the Moessbauer spectrum of thulium metal* (696)
 Preparation of thorium metal by electrolysis of fused salt $ThCl_{sub 4}-LiCl+NaCl+KCl$ (eut) system* (703)
 High density thoria-silica-metal (III) oxide fibers* (710)
 Standard methods for chemical, mass spectrometric, spectrochemical, nuclear, and radiochemical analysis of nuclear-grade plutonium metal* (714)

metallic

Magnetic susceptibility and electronic structure in metallic and non-metallic liquid cesium-gold and cesium-antimony alloys* (190)
 Wetting of certain refractories of the system $MgO-Al_{sub 2}O_{sub 3}-ZrO_{sub 2}$ with metallic melts* (344)
 Solution behavior of hydrogen isotopes and other non-metallic elements in liquid lithium* (404)
 Dispersion of metallic lithium in various molten salts* (452)

metallocarbaborane

Actinide metallocarbaborane complex; synthesis and x-ray structure determination of the bis[$n^{sup}5-(3-1,2-dicarbollyl)dichlorouranium(IV)$ dianion* (194)

Metallurgical

Metallurgical characterization of prototypic tubesheet forging* (471)

metals

X-ray diffraction study into molten bromides and iodides of alkali metals* (20)
 Spectral emissivities of molten refractory metals* (47)
 Studies of the electronic structure of simple metals and their alloys with the aid of compton scattering and positron annihilation using lithium, magnesium, lithium-magnesium and aluminium-zinc as examples* (48)
 Characterisation of the chemical activities of non-metals dissolved in liquid alkali metals* (63)
 Characterisation of the chemical activities of non-metals dissolved in liquid alkali metals* (64)
 Environmental sensitivity of structural metals: liquid metal embrittlement. Annual technical progress report No. 2, 25 Jun 1970--31 May 1971* (95)
 Nonisothermal flow of liquid metals over the surface of solid ones* (101)
 Mechanism of metals modifying* (108)

TITLE INDEX

metals

- Compatibility of molten rare earth metals with refractory materials* (115)
- Revised model for 5d-electrons in the heavy rare-earth metals* (157)
- Electron spectrum of light alkali metals* (161)
- Discrete variation of the structure and texture of liquid metals formed by components with discrete packing* (177)
- Bonding and the electronic structure of the actinide metals* (188)
- Gases and carbon in metals (thermodynamics, kinetics, and properties). I. Alkali metals, alkaline earth metals, light metals (Li, Na, K, Rb, Cs; Ca, Sr, Ba; Be, Mg, Al)* (192)
- Self-diffusion in liquid metals* (198)
- Environmental sensitivity of structural metals: liquid metal embrittlement. Annual technical progress report No. 4, 1 Jun 1972--31 May 1973* (211)
- Environmental sensitivity of structural metals: liquid metal embrittlement. Annual technical progress report No. 2, 25 Jun 1970--31 May 1971* (212)
- Vacancies and melting curves of metals at high pressure* (213)
- Negative temperature coefficients of electrical resistivity: the divalent liquid metals Eu, Yb and Ba* (224)
- Entropies of liquid rare earth metals* (230)
- Thermochemistry of gaseous compounds of metals. Annual summary report 5 Dec 74--4 Dec 75* (240)
- Thermochemistry of gaseous compounds of metals. Annual summary report, 5 Dec 1975--4 Dec 1976* (241)
- Thermodynamic properties of solid and liquid metals and ceramics* (243)
- Specific heat of liquid metals and ceramics* (244)
- Aspects of the solution chemistry of liquid alkali metals as elucidated from electrical resistivity studies* (247)
- Liquid metals, 1976* (249)
- Structure factors of liquid alkali metals* (250)
- Anomalous melting of f electron metals (with attention to Pu)* (318)
- Diffusion coefficients in molten metals* (328)
- Origin of excess entropy of alkali metals halides solid solutions* (390)
- Thermal decomposition of intermetallic compounds of platinum with alkaline earth and lanthanide metals at high temperatures* (399)
- Speed of ultrasound and the thermophysical properties of the liquid metals Sn, Pb, Cd and of their binary alloys Pb:Sn and Pb:Cd* (403)
- Photoelectron spectroscopic studies of some transition metals and alloys* (417)
- Comparative study of systems of heavier rare-earth metals with hydrogen [Gd, Tb, Dy, Lu]* (418)
- Densities of electronic states in liquid transition metals* (422)
- Heat of formation of solid and liquid alloys of transition metals* (425)
- Periodically variable wall temperature effect on heat transfer in liquid metals* (437)
- Rare earth metals* (461)
- Rare earth metals* (462)

TITLE INDEX

metals

3. Zirconium compounds with non-metals* (464)
 KKR analysis of the Anderson-McMillan theory of electronic structure of liquid metals. Technical report* (485)
 Liquid metals, 1976* (486)
 Recent developments in the chemistry of solutions of salts in liquid alkali metals* (528)
 Basic mechanisms providing oxidation resistance in structural metals at high temperatures. II. The defect structure of nonstoichiometric rutile (TiO_{2-x}) and nonstoichiometric rutile containing 1 to 10 mole % Nb_2O_5 . Final report 21 Dec 1973-20 Dec 1974* (542)
 Rare earth metals* (546)
 Nuclear quadrupole relaxation and viscosity in liquid metals* (564)
 Thermal expansion of metals over the entire liquid range* (573)
 X-ray spectral study of the carbides and nitrides of group IV and V metals* (582)
 Liquid metals, 1976* (584)
 Density and enthalpy of liquid alkaline metals at pressures up to 1500 bar* (599)
 Application of the hard-sphere model to thermodynamic properties of liquid 3d transition metals* (615)
 Experimental study of the temperature dependence on energy exchanges in liquid metals* (621)
 Investigation of particles interaction in liquid metals and alloys* (661)
 Thermodynamic properties of solutions of hydrogen isotopes in metals and alloys of interest to fusion reactor technology* (663)
 Spectra of combination scattering and structure of 3d-metals chloride complexes in molten salts* (674)
 Electronic structure of actinides and related metals* (683)
 Structure of liquid transition metals and their alloys* (686)
 Structures of 3d-transition metals in the liquid state* (687)
 Structure of rare earth metals in the liquid state* (688)
 Electronic states densities from structure and thermodynamic data for liquid transition metals* (705)

metaniobates

- Stability of rare earth metaniobates in hydrogen atmosphere* (164)

MF

- Phases of variable composition with LaF_3 structure in $\text{MF}_2-(\text{Y}, \text{Ln})\text{F}_3$ systems. 1. $\text{SrF}_2-(\text{Y}, \text{Ln})\text{F}_3$ systems (thermal characteristics, formation of berthollides)* (593)

Mg

- Thermodynamic properties of the liquid La-Mg alloys* (3)
 Investigation into thermodynamic properties of Ce-Mg melts* (39)
 Vibrational fine structure in the uranium emission of Mg_3TeO_6 and Y_6WO_{12} * (59)
 Gases and carbon in metals (thermodynamics, kinetics, and properties). I. Alkali metals, alkaline earth metals, light metals (Li, Na, K, Rb, Cs; Ca, Sr, Ba; Be, Mg, Al)* (192)
 Solubility of rare earth and yttrium oxochlorides in molten salts - $\text{MCl}_2(\text{M-Mg, Ca, Sr, Ba})$ * (495)

TITLE INDEX

- Mg Phase diagrams of the Mg(NO₃)₃-Nd(NO₃)₃ system* (668)
- MgCl Viscosity of UCl₃--MgCl₂ binary system melts* (145)
- MgO High temperature thermal expansion of ThO₂, MgO and Y₂O₃ by X-ray diffraction* (242)
Wetting of certain refractories of the system MgO-Al₂O₃-ZrO₂ with metallic melts* (344)
- microcalorimetric High-temperature microcalorimetric measurements of partial molar enthalpy of solution, $\Delta H(O_2)$, in plutonium + oxygen* (107)
- microinhomogeneity NMR study on structural microinhomogeneity of fluoride-oxide melts* (463)
- Microprobe Microprobe study of fission product behavior in high-burnup HTR fuels* (315)
- microscopy Rare earth and actinide oxides. Thermodynamic and electron microscopy studies* (625)
- microstructure Investigation of the parameters influencing the microstructure of hot-pressed boron carbide* (440)
- mild Preliminary assessments of carbide fuel pins during mild overpower transients in LMFBRs* (456)
- misch Electric conductivity of melts of the LiF-misch metal fluoride system* (461)
Misch metal solubility in melt of LiF-BaF₂ system* (462)
- Mixed Mixed valence rare-earth halides and their unusual crystal structures* (29)
Interactions in inorganic molecular crystals - electronic spectra of ReF₆ pure and mixed crystals. Technical report, 1 Jul--31 Dec 1975* (53)
Contribution to the thermodynamic study of uranium and nickel mixed oxides* (149)
Behavior of molybdenum in mixed-oxide fuel* (202)
Radial oxygen distribution in the fuel of irradiated mixed oxide fuel elements with different final stoichiometries* (316)
Investigation of solubility and thermodynamics of dissolution of salts in mixed solvents* (347)
Fission rates, burnup, and neutron flux-fluence-spectra characterization for mixed oxide fuel experiments in the EBR-II* (416)
Measurement of the cell voltage of ThO₂--Y₂O₃-solid electrolyte cells in the range of mixed conduction* (535)
Oxygen potential and vacancy structure of the mixed oxide (UPu)O_{2-x}* (565)
Fabrication of uranium-plutonium mixed carbide* (691)

TITLE INDEX

mixed

Standard methods for chemical, mass spectrometric, and spectrochemical analysis of nuclear-grade mixed oxides (U,Pu)O₂* (713)

mixing

Determination of vibrational contribution into excessive mixing entropy of liquid amalgams. Alkali metal amalgams* (138)

mixture

Measurement of mass transfer coefficients in a mechanically agitated, nondispersing contactor operating with a molten mixture of LiF--BeF₂--ThF₄ and molten bismuth* (79)

Interaction of the lithium-tin alloys with the LiCl-KCl-CsCl melted eutectic mixture* (91)

Equilibrium between Li-K-Te alloys and LiCl-KCl molten eutectic mixture* (92)

mixtures

Viscosity of molten CaCl₂-UCl₃ mixtures* (146)

Composition gradient during electrolysis in mixtures analogous to molten salt battery electrolytes* (234)

Use of fluoride melts for separation of some fission products and americium from mixtures of different fluorides being the simulators of residues from fast reactor fuel fluorination* (307)

Solubility of magnesium oxide in calcium oxide--calcium chloride mixtures* (352)

Quantitative determination of UO₂, and ThO₂ in powdery mixtures by infrared spectrometry* (431)

Application of molecular dynamics computations to the conformal ionic solution theory: Thermodynamics of binary molten salt mixtures* (550)

Mn

Thermodynamic properties of Cr, Mn, Co and Ni dichlorides from emf measurements on cells with solid electrolytes* (200)

MnCl

Thermodynamic properties of diluted solutions of MnCl₂ in alkali metal chlorides* (6)

Mo

Effect of composition and structure on crevice, intergranular, and stress corrosion of some wrought Ni--Cr--Mo alloys* (604)

Phase behavior and thermodynamics of U--Mo--C system* (646)

Phase behavior and thermodynamics of U--Mo--C system* (647)

MoCl

MoCl₂-NaCl and MoCl₂-KCl binary systems* (547)

FeCl₃-MoCl₅-NbCl₅-WCl₆ system* (554)

model

Thermodynamic properties of (U,Pu)O_(2+x) described by a cluster model* (133)

Revised model for 5d-electrons in the heavy rare-earth metals* (157)

Thermodynamics of non-stoichiometry. A model for UC_x* (442)

Improved model for solutions of nonmetals in liquid alkali metals. Calculation of enthalpy of solution and electrical resistivity* (486)

Some trends in constitutive equation model development for high-temperature behavior of fast-reactor structural alloys* (527)

TITLE INDEX

model

Analysis of LMFBR explosion model experiments by means of the SUREFORM-II code* (601)
 Application of the hard-sphere model to thermodynamic properties of liquid 3d transition metals* (615)

models

Calculation of the oxygen potential of $(U,Pu)O_{2-x}$ with defect models and measured oxygen potential data in the temperature range from 1,000 to 1,700 K* (74)
 Physical models of mass transport of iron and nickel in liquid sodium systems* (128)

moderated

Collection of gamma spectra data of irradiated light water moderated reactor spent fuel and study of the applicability of the methods for fuel identification. Final report for the period 1 February 1972--October 1975* (453)

modern

Equilibria and kinetics in modern ceramic processing* (557)

modifying

Mechanism of metals modifying* (108)

Moessbauer

Moessbauer studies of americium compounds* (61)
 Electronic and magnetic properties of rare earth-Sn₃ compounds from ¹¹⁹Sn Moessbauer spectroscopy* (559)
 A Moessbauer spectroscopic study of intercalation compounds of graphite with ferric chloride and aluminum chloride* (627)
 Accurate nuclear parameters derived from the Moessbauer spectrum of thulium metal* (696)

MoF

Vibrational spectra of transition metal hexafluoride crystals. III. Exciton band structures of MoF₆, WF₆, and UF₆. Technical report* (52)

MoF₆

Vibrational spectra of transition metal hexafluoride crystals. I. Orthorhombic MoF₆, WF₆, UF₆ neat crystals. Technical report* (51)

mol

Quantitative study of the chemical equilibria in solution in molten LiCl-KCl (70-30 mol.%) by absorption spectrophotometry. Application to neptunium* (385)

molar

High-temperature microcalorimetric measurements of partial molar enthalpy of solution, $\Delta H(O_{2g})$, in plutonium + oxygen* (107)

mole

Basic mechanisms providing oxidation resistance in structural metals at high temperatures. II. The defect structure of nonstoichiometric rutile (TiO₂) and nonstoichiometric rutile containing 1 to 10 mole % Cb_2O_6 . Final report 21 Dec 1973-20 Dec 1974* (542)

molecular

11. european congress on molecular spectroscopy. Abstracts* (44)
 11. european congress on molecular spectroscopy. Abstracts* (46)
 Interactions in inorganic molecular crystals - electronic spectra of ReF₆ pure and mixed crystals. Technical report, 1 Jul--31 Dec 1975* (53)

TITLE INDEX

Molecular

Molecular vapor composition and mass spectra in KF-ZrF_4 systems* (294)
 Isotope effects in the molecular spectrum of plutonium hexafluoride* (353)
 Using molecular dynamics method for the evaluation of Fe-Ni and Al-Zn melts structure properties* (512)
 Theory for the derivation of a single hard-sphere potential describing the overall molecular interactions in binary liquid metal alloys as a function of temperature and composition* (525)
 Application of molecular dynamics computations to the conformal ionic solution theory: Thermodynamics of binary molten salt mixtures* (550)
 Frequency spectrum and scattering law for liquid copper. A molecular dynamics investigation* (567)

molecule

Structure of the biexciton molecule and optical transition selection rules in zincblende and wurtzite materials* (186)
 Absorption spectrum of HfCl_4 molecule* (283)
 Predicted infrared spectrum and the structure of the isolated UF_5 molecule* (350)

molecules

Spectral properties of impurity molecules of the CsCl_3 symmetry groups in solid solutions of halide salts* (44)
 Mass spectrometric determination of dissociation energies of high-temperature molecules: ScS , YS , LaS , ZrS , UO^* (597)

molten

Density and temperature dependence of the surface tension of molten bismuth, lead, and bismuth-lead alloys* (1)
 X-ray diffraction study into molten bromides and iodides of alkali metals* (20)
 Some observations on the compatibility of structural materials with molten lead* (25)
 Electrode reactions in molten sodium nitrate--barium nitrate eutectic at 350 degrees C* (42)
 Spectral emissivities of molten refractory metals* (47)
 Molten carbonate fuel cell research at ORNL* (71)
 Measurement of mass transfer coefficients in a mechanically agitated, nondispersing contactor operating with a molten mixture of LiF -- BeF_2 -- ThF_4 and molten bismuth* (79)
 Equilibrium between Li-K-Te alloys and LiCl-KCl molten eutectic mixture* (92)
 Study on conductivity, density, and viscosity of molten salt systems* (112)
 Compatibility of molten rare earth metals with refractory materials* (115)
 Viscosity of molten CaCl_2 -- UCl_3 mixtures* (146)
 Thermal expansion of molten materials by the gamma attenuation technique* (152)
 Molten-salt reactor program. Semiannual progress report for period ending February 29, 1976* (176)
 Structure of NO_3^- radical in molten monovalent metal nitrates* (195)
 Raman spectra of Al_2O_3 solutions in molten cryolite and other aluminum fluoride containing melts* (203)

TITLE INDEX

molten

- High-temperature unit to the IKS-21 spectrometer for the detection of molten salts absorption spectra* (208)
- Prediction of the inert gas solubilities in stoichiometric molten UO_2 * (225)
- Measurement of fluorescence of rare earths in molten salts (Nd^{3+} in NaNO_3 - KNO_3 eutectic)* (233)
- Composition gradient during electrolysis in mixtures analogous to molten salt battery electrolytes* (234)
- Fuel processing for molten-salt reactors* (237)
- Electrochemistry of molten salts* (263)
- Structure of lithium--boron alloys and their performance as potential anodes for high power molten salt batteries. Progress report, 1 Aug 1974--30 Jun 1975* (275)
- Molten-salt electrolytes--transport properties* (277)
- Chronopotentiometry of U(III) and U(IV) in molten LiCl-KCl eutectic* (292)
- Electrochemical behavior of U(III) and U(IV) in molten LiCl-KCl eutectic* (293)
- Dissolution of corundum in molten vanadates* (317)
- Diffusion coefficients in molten metals* (328)
- Use of the SF-8 detection spectrophotometer for measuring absorption spectra of molten salts* (331)
- Heat capacities and heat content of molten cerium by levitation calorimetry* (355)
- Structure of molten Bi--Sb alloys by means of neutron diffraction* (359)
- Redox and chemical properties of U, Np and Pu in molten alkali chlorides* (360)
- Investigations on the solvent extraction from molten salts by tracer technique* (369)
- Investigations of the solvent extraction from molten salts with the use of tracer technique* (371)
- Quantitative study of the chemical equilibria in solution in molten LiCl-KCl (70-30 mol.%) by absorption spectrophotometry. Application to neptunium* (385)
- Diffusion of uranium species in several molten chlorides* (408)
- Activities of lead and bismuth in their molten alloys* (420)
- Energy programs: molten-salt reactor program* (423)
- Electrode potentials of tantalum chlorides in a molten NaCl-KCl eutectic* (449)
- Electrochemistry of niobium and tantalum, (5). Anodic dissolution of tantalum and the average valence of tantalum ions in molten KCl* (450)
- Electrochemistry of niobium and tantalum. VI. Electrode potentials of tantalum chlorides in molten KCl* (451)
- Dispersion of metallic lithium in various molten salts* (452)
- Solubility of rare earth and yttrium oxychlorides in molten salts - MCl_2 (M-Mg, Ca, Sr, Ba)* (495)
- Electrochemical reduction of molybdenum oxychloride in molten lithium chloride - potassium chloride eutectic* (515)
- Application of molecular dynamics computations to the conformal ionic solution theory: Thermodynamics of binary molten salt mixtures* (550)
- Electromotive force measurements in molten lithium--magnesium alloys* (551)

TITLE INDEX

molten

- Solubility of oxygen in molten carbonates* (563)
- Spectrochemical method for determination of zirconium in molten salts* (566)
- Heat transfer measurements in a forced convection loop with two molten-fluoride salts: $\text{LiF--BeF}_2\text{--ThF}_2\text{--UF}_4$ and eutectic $\text{NaBF}_4\text{--NaF}$ * (585)
- Refractory materials resistance to molten barium alloy* (628)
- Molten salt chemistry : Part IV - Solubility behaviour of PuF_3 in fluoride salts of interest in molten salt reactor technology* (655)
- Isorhodate complexes of 3d-metal ions in molten salts* (672)
- Effect of second sphere cations on formation of nickel (2) chloride complexes in molten salts* (673)
- Spectra of combination scattering and structure of 3d-metals chloride complexes in molten salts* (674)
- Investigation of the solubility and diffusion of Fe atoms in Cu at high temperature using molten salt electrochemistry* (693)
- Lithium alloys--chlorine secondary battery. I. Basic studies on a lithium alloys--chlorine secondary battery using a molten salt as the electrolyte* (704)
- Electrochemical methods for monitoring purity of molten salts. Semiannual progress report, July--December 1975* (721)
- Molten-salt systems* (729)

molybdate

- Raman and i.r.-emission studies of some tungstate- and molybdate-containing melts* (102)

molybdates

- Binary molybdates of rare earths and silver (I)* (207)

molybdenum

- Electrochemical machining of the molybdenum VM-1 alloy* (129)
- Molybdenum and tungsten interaction with liquid aluminium* (156)
- Behavior of molybdenum in mixed-oxide fuel* (202)
- Some properties of molybdenum pentoxide* (305)
- Thermographic study of iron, molybdenum, tantalum and tungsten high chlorides interaction* (555)
- Investigation into outgassing from molybdenum during melting* (588)
- Thermodynamic activity of carbon in molybdenum-containing uranium carbide* (644)
- Thermodynamic activity of carbon in molybdenum-containing uranium carbide* (650)
- Determination of trace impurities of titanium and zirconium in molybdenum and tungsten and their oxygen compounds by X-ray fluorescence spectrometry* (698)

molybdenum

- Electrochemical reduction of molybdenum oxychloride in molten lithium chloride - potassium chloride eutectic* (515)

monitoring

- Electrochemical methods for monitoring purity of molten salts. Semiannual progress report, July--December 1975* (721)

monocarbide

- Thermodynamic assessment of the conversion of plutonium dioxide to plutonium monocarbide* (54)
- Thermodynamics of the conversion of plutonium dioxide to plutonium monocarbide* (55)

TITLE INDEX

monocarbide

Conversion of protactinium monocarbide to the penta- and tetra-halides* (80)
Specific heat of thorium monocarbide from 2 to 300K* (126)

monovalent

Structure of $\text{NO}_3^{\cdot-}$ radical in molten monovalent metal nitrates* (195)
Study of kinetics of monovalent indium ions formation in the $\text{In}^0\text{-InCl}_3\text{-LiCl-H}_2\text{O}$ system* (341)
Investigation into properties of liquidus curves of binary salt system melts where chemical compounds formation. 2. Systems of mono-monovalent and bi-bivalent cations with common anion* (708)

monoxide

Thermodynamic study of the U-O-C system by measurement of the carbon monoxide pressures at equilibrium* (235)

Monte

Monte Carlo analysis of diffusion and thermodynamics in UO_2 * (441)

MoO

Phase diagrams of $\text{Li}_2\text{MoO}_4\text{-ZnMoO}_4$, $\text{K}_2\text{MoO}_4\text{-ZnMoO}_4$ and $\text{K}_2\text{WO}_4\text{-ZnWO}_4$ systems* (160)
Practical application of conversion method in study of five-component mutual system consisting of nine salts NaF , Na_2MoO_4 , Na_2WO_4 , KF , K_2MoO_4 , K_2WO_4 , BaF_2 , BaMoO_4 , BaWO_4 * (516)
Thermal analysis of the system of NaF , CaF_2 , Na_2MoO_4 , CaMoO_4 * (519)
 $\text{KF-KCl-K}_2\text{MoO}_4$ system* (520)

motion

Computational simulation of cladding motion after melting including resolidification processes* (19)

Msup

Preparation and structure of transition metal fluorides $\text{U}^{\text{II}}\text{M}^{\text{IV}}\text{F}_6$ [U^{II} : Cr-Zn; M^{IV} : Hf, Zr]* (596)

multicomponent

Solubility of nitrogen in multicomponent nickel base alloys* (100)
Calculated method of eutectics of multicomponent systems on base of algorithm and digital computers* (517)
Practice of experiment planning in studying limited regions of multicomponent system* (546)

mutual

Quaternary mutual system from fluorides and chlorides of lithium, sodium and strontium* (89)
Ternary mutual system of chlorides and iodides of rubidium and cadmium* (257)
Practical application of conversion method in study of five-component mutual system consisting of nine salts NaF , Na_2MoO_4 , Na_2WO_4 , KF , K_2MoO_4 , K_2WO_4 , BaF_2 , BaMoO_4 , BaWO_4 * (516)
Projection-thermographic method for study of ternary and ternary mutual systems* (518)
Quaternary mutual system of chlorides, bromides, iodides of sodium and rubidium* (629)

TITLE INDEX

mutual

Quaternary mutual system of chlorides, bromides, iodides of potassium and cesium* (630)
 Sections of quaternary mutual system consisting of sodium, potassium and cesium chlorides and bromides* (631)
 Quaternary mutual system of chlorides, bromides, iodides of rubidium and cesium* (640)

N

Lithium nitride (Li_3N): standard enthalpy of formation by solution calorimetry* (467)
 Chemistry of elements niobium and tantalum. 87. Studies on the thermal formation and decomposition of the complexes $[\text{NbCl}_{12}]^{2+}$, $[\text{Nb}_6\text{I}_8]^{3+}$, and $[\text{Ta}_6\text{X}_{12}]^{n+}$ ($\text{X}=\text{Cl}, \text{Br}, \text{I}$)* (561)
 Solubility of nitrogen in liquid lithium and thermal decomposition of solid Li_3N * (702)

Na

Density, viscosity, electric conductivity and meltability of Na_3AlF_6 - La_2O_3 and Na_3AlF_6 - Sm_2O_3 systems* (43)
 Me_2ZrCl_6 - H_2 ($\text{Me}=\text{Na}, \text{K}, \text{Rb}, \text{Cs}$) system* (45)
 Phase equilibria in Li - Na - K carbonate/aluminate systems* (77)
 Gases and carbon in metals (thermodynamics, kinetics, and properties). I. Alkali metals, alkaline earth metals, light metals ($\text{Li}, \text{Na}, \text{K}, \text{Rb}, \text{Cs}; \text{Ca}, \text{Sr}, \text{Ba}; \text{Be}, \text{Mg}, \text{Al}$)* (192)
 Standard free energy of formation of liquid Na_2CrO_4 , 1100 to 1220 K* (375)
 Investigation of calcite solubility in K_2CO_3 and Na_2CO_3 solutions at temperatures of 200 to 350°C* (396)
 Some thermodynamic properties of dilute solutions of actinide chlorides in $(\text{Li}-\text{K})\text{Cl}$ and in $(\text{Na}-\text{K})\text{Cl}$ eutectics* (406)
 Thermodynamic properties of dilute solutions of ThCl_4 in $(\text{Li}-\text{K})\text{Cl}$ and $(\text{Na}-\text{K})$ eutectics* (407)
 Practical application of conversion method in study of five-component mutual system consisting of nine salts NaF , Na_2MoO_4 , Na_2WO_4 , KF , K_2MoO_4 , K_2WO_4 , BaF_2 , BaMoO_4 , BaWO_4 * (516)
 Thermal analysis of the system of NaF , CaF_2 , Na_2MoO_4 , CaMoO_4 * (519)
 Pressure and entropy calculations for liquid Na , K and Rb along the melting line* (584)
 System Na_2WO_4 - NaCl - CaWO_4 - CaCl_2 * (637)

NaBF

Heat transfer measurements in a forced convection loop with two molten-fluoride salts: LiF - BeF_2 - ThF_4 - UF_4 and eutectic NaBF_4 - NaF * (585)

NaCe

Electronic structure of highly symmetric lanthanoid and actinoid compounds. VI. The electronic Raman spectrum of $\text{Cs}_2\text{NaCe(III)Cl}_6$ * (678)

NaCl

Standard formation enthalpy of chloride alloys of ZrCl_4 - KCl (NaCl) systems* (8)
 Lattice-dynamical calculation of the thermodynamic properties of NaCl , KCl and RbCl * (231)

TITLE INDEX

NaCl

Meltability in system of K_2TaF_7 --NaF--NaCl--KCl* (299)
 Meltability in K_2NbF_7 --NaF--NaCl--KCl system* (300)
 $NdCl_3$ --AlCl₃--NaCl system* (434)
 Electrode potentials of tantalum chlorides in a molten NaCl-KCl eutectic* (449)
 $MoCl_2$ --NaCl and $MoCl_2$ --KCl binary systems* (547)
 $ZrCl_4$ --TiCl₃--NaCl ternary system* (548)
 Thermal analysis of NaF, NaCl, Na_2MoO_4 system* (636)
 System Na_2WO_4 --NaCl--CaWO₄--CaCl₂* (637)
 Preparation of thorium metal by electrolysis of fused salt $ThCl_4$ --LiCl+NaCl+KCl(eut) system* (703)

NaEu

Electronic structure of highly symmetric complexes of lanthanoids and actinoids. Pt. 4. The electronic Raman spectrum of $Cs_2NaEu(III)Cl_6$ * (14)

NaF

Fusibility in $K_2Nb(Ta)F_7$ --LiF--NaF systems* (298)
 Meltability in system of K_2TaF_7 --NaF--NaCl--KCl* (299)
 Meltability in K_2NbF_7 --NaF--NaCl--KCl system* (300)
 Investigation into americium extraction process from residues of irradiated nuclear fuel after its fluorination by SiF₄--NaF and SiF₄--CaF₂* melts of eutectic composition* (308)
 Comparison of the tendencies of complex formation of various metal ions in fused LiF--NaF--KF (eut.) and in fused LiCl--KCl (eut.)* (314)
 Practical application of conversion method in study of five-component mutual system consisting of nine salts NaF, Na_2MoO_4 , Na_2WO_4 , KF, K_2MoO_4 , K_2WO_4 , BaF₂, $BaMoO_4$, $BaWO_4$ * (516)
 Thermal analysis of the system of NaF, CaF₂, Na_2MoO_4 , $CaMoO_4$ * (519)
 Heat transfer measurements in a forced convection loop with two molten-fluoride salts: LiF--BeF₂*--ThF₄*--UF₄* and eutectic NaBF₄*--NaF* (585)
 Thermal analysis of NaF, NaCl, Na_2MoO_4 system* (636)
 Na_2MoO_4 --BaMoO₄--NaF--BaF₂ system* (638)

NaI

TlCl--RbCl--CsCl and NaI--TlI--CsI systems* (258)

NaIO

Constitution equilibrium diagram of the $LiNO_3$ --LiIO₃, $NaNO_3$ --NaIO₃, KNO_3 --KIO₃ system* (665)

NaMoO

Thermal analysis of NaF, NaCl, Na_2MoO_4 system* (636)
 Na_2MoO_4 --BaMoO₄--NaF--BaF₂ system* (638)

NaNO

Measurement of fluorescence of rare earths in molten salts (Nd^{3+} in $NaNO_3$ --KNO₃ eutectic)* (233)
 $LiNO_3$ --CsNO₃--Ca(NO₃)₂ and $NaNO_3$ --CsNO₃--Ca(NO₃)₂ system* (255)
 Constitution equilibrium diagram of the $LiNO_3$ --LiIO₃, $NaNO_3$ --NaIO₃, KNO_3 --KIO₃ system* (665)
 Phase diagram of $NaNO_3$ --Nd(NO₃)₃ system* (667)

TITLE INDEX

National

Argonne National Laboratory* (692)

Nauchnye

Nauchnye trudy GIREDMETA. Vol. 52* (115)

NaUO

Sodium uranium(V) trioxide, NaUO_3 : heat capacity and thermodynamic properties from 5 to 350 K* (384)

NaUO

Sodium uranium(V) trioxide, NaUO_3 : heat capacity and thermodynamic properties from 5 to 350 K* (383)

NaYCl

Raman spectroscopic studies of yttrium (III) chloride--alkali metal chloride melts and of $\text{Cs}_2\text{NaYCl}_6$ and YCl_3 solid compounds* (489)

Nb

Fusibility in $\text{K}_2\text{Nb(Ta)F}_7$ -LiF-NaF systems* (298)

NbBr

IR spectra of the NbCl_5 , TaCl_5 , NbBr_5 and NbOCl_3 vapors* (707)

NbCl

FeCl_3 - MoCl_5 - NbCl_5 - WCl_6 system* (554)
Systems of NbCl_5 - GaCl_3 and TaCl_5 - GaCl_3 * (633)

IR spectra of the NbCl_5 , TaCl_5 , NbBr_5 and NbOCl_3 vapors* (707)

NbF

Meltability in K_2NbF_7 -NaF-NaCl-KCl system* (300)

NbO

Thermodynamic properties of NbO , $\text{Fe}_{0.95}\text{O}$, CoO , NiO , Cu_2O and CuO oxides as comparative electrodes in electromotive forces method* (658)

NbOCl

IR spectra of the NbCl_5 , TaCl_5 , NbBr_5 and NbOCl_3 vapors* (707)

Nd

Measurement of fluorescence of rare earths in molten salts (Nd^{3+} in NaNO_3 - KNO_3 eutectic)* (233)

Phase diagram and properties of Nd-As alloys* (321)

Structure of Nd^{3+} terms in nitrate glasses* (664)

Phase diagram of NaNO_3 -Nd(NO_3) $_3$ system* (667)

Phase diagrams of the Mg(NO_3) $_2$ -Nd(NO_3) $_3$ system* (668)

NdCl

Thermal investigation of NdCl_3 - AlCl_3 -KCl and NdCl_3 - AlCl_3 -CsCl system* (289)

NdCl_3 - AlCl_3 -NaCl system* (434)

Phase equilibrium in NdCl_3 -KCl system* (457)

NdF

RbF - NdF_3 system* (443)

neat

Vibrational spectra of transition metal hexafluoride crystals. I. Orthorhombic MoF_6 , WF_6 , UF_6 neat crystals. Technical report* (51)

Negative

Negative temperature coefficients of electrical resistivity: the divalent liquid metals Eu, Yb and Ba* (224)

TITLE INDEX

negative

Studies on the negative alloy electrode* (269)

neodymium

Investigation into neodymium chloride state in equilibrium with metal in eutectic melt of potassium- and lithium chlorides* (40)
Interaction of neodymium- and yttrium chlorides with magnesium oxide* (494)
Interaction of potassium tetrafluoroborate with fluorides of potassium, lanthanum, neodymium in melt* (530)
Standard method of test for atom percent fission in uranium and plutonium fuel (neodymium-148 method)* (716)

neptunium

Decomposition, stoichiometry and structure of neptunium oxides* (171)
Quantitative study of the chemical equilibria in solution in molten LiCl-KCl (70-30 mol.%) by absorption spectrophotometry. Application to neptunium* (385)
Thermodynamic study of the neptunium-hydrogen system* (428)
Diffuse reflectance spectra of protactinium(IV), uranium(IV) and neptunium(IV) tetrachlorides and tetraformates* (562)
Spectrophotometric method for the determination of neptunium and plutonium in process solutions* (660)

neutral

Tetrad effect in the extraction of the lanthanides with neutral oxygen-containing compounds* (426)

neutralization

Twenty years of research in the processing of radioactive fuel and neutralization of radioactive waste at the Nuclear Research Institute (NRI)* (358)

neutron

Structure of molten Bi-Sb alloys by means of neutron diffraction* (359)
Fission rates, burnup, and neutron flux-fluence-spectra characterization for mixed oxide fuel experiments in the EBR-II* (416)

NFE

Criteria for solid solution formation in NFE alloys* (205)

NH

Study of the compounds $M\text{UO}_2\text{F}_5$ ($M = K, Rb, Cs, NH_4$) by i.r. absorption and Raman diffusion spectrophotometry. III. Effect of cation on vibration frequency of the ion $\text{UO}_2\text{F}_5^{3-}$ * (127)

Ni

Study of the Ni-Ni (II) equilibrium exchange, by radiotracer techniques in fused LiCl-KCl eutectic* (68)
Thermodynamic properties of Cr, Mn, Co and Ni dichlorides from emf measurements on cells with solid electrolytes* (200)
 R_3T_2 compounds ($R = \text{rare earth or Y; } T = Rh, Pd, Pt$) with the rhombic Er $_3$ Ni $_2$ structure type* (365)
Using molecular dynamics method for the evaluation of Fe-Ni and Al-Zn melts structure properties* (512)
Effect of composition and structure on crevice, intergranular, and stress corrosion of some wrought Ni-Cr-Mo alloys* (604)
Structure of liquid Ce-Ni alloys* (689)

TITLE INDEX

nickel

Variation of the solid--solid--nickel--thoria interfacial free energy with temperature* (5)
 Viscosity characteristics of the iron-chromium-nickel melts* (24)
 Solubility of nitrogen in multicomponent nickel base alloys* (100)
 Physical models of mass transport of iron and nickel in liquid sodium systems* (128)
 Contribution to the thermodynamic study of uranium and nickel mixed oxides* (149)
 Investigation into interaction of indium tribromide with dibromides of manganese, cobalt and nickel in melt* (153)
 Kinetic peculiarities of nitrogen dissolving in iron-chromium and iron-chromium-nickel melts* (172)
 Spreading of iron-nickel alloys on ZrC and nickel-cobalt alloys on ZrN* (252)
 Lithium--nickel fluoride secondary battery investigation. Quarterly technical progress report No. 1, 27 Apr--27 Jul 1966* (381)
 Nitrogen solubility in liquid nickel--niobium alloys* (532)
 Effect of second sphere cations on formation of nickel (2) chloride complexes in molten salts* (673)

NiC

Thermodynamic properties of NbO , $\text{Fe}_{0.95}\text{O}$, CoO , NiO , $\text{Cu}_{\text{sub}} 2\text{O}$ and CuO oxides as comparative electrodes in electromotive forces method* (658)

niobium

Thermodynamics of reactions of rare earth oxides with niobium oxide* (206)
 Mass spectra and vapor composition of niobium pentafluoride* (216)
 Structure of the thorium-(zirconium, niobium, ruthenium, rhodium)-carbon systems* (245)
 Solubility of oxygen in liquid niobium* (342)
 Electrochemistry of niobium and tantalum, (5). Anodic dissolution of tantalum and the average valence of tantalum ions in molten KCl * (450)
 Electrochemistry of niobium and tantalum. VI. Electrode potentials of tantalum chlorides in molten KCl * (451)
 Differential-spectrophotometric determination of niobium in its compounds* (490)
 Nitrogen solubility in liquid nickel--niobium alloys* (532)
 Chemistry of elements niobium and tantalum. 87. Studies on the thermal formation and decomposition of the complexes $[\text{Nb}_{\text{sub}} 6\text{Cl}_{\text{sub}} 12]^{\text{sup } 2+}$, $[\text{Nb}_{\text{sub}} 6\text{I}_{\text{sub}} 8]^{\text{sup } 3+}$, and $[\text{Ta}_{\text{sub}} 6\text{X}_{\text{sub}} 12]^{\text{sup } (n+)}$ ($\text{X}=\text{Cl}, \text{Br}, \text{I}$)* (561)
 Terminal solid solubility of hydrogen in the zirconium-2.5 weight % niobium alloy* (587)

nitrate

Electrode reactions in molten sodium nitrate--barium nitrate eutectic at 350 degrees C* (42)
 Results of research to evaluate solid plutonium nitrate as a safe shipping form* (85)
 Cesium nitrate. Standard enthalpies of solution and formation and the standard entropy at 298.15 K* (466)

TITLE INDEX

nitrate

Viscosity of binary of nitrate systems melts* (592)
 Structure of Nd^{3+} terms in nitrate glasses* (664)
 Glass formation in nitrate systems containing rare earth ions* (666)

nitrates

Investigation into reactions between trioctylammonium nitrates and rare earth nitrates by X-ray electron spectroscopy* (113)
 Zeeman effect and satellite carrier structure in absorption spectra of pure and doped praseodymium and europium double nitrates* (165)
 Structure of $\text{NO}_3^{\cdot -}$ radical in molten monovalent metal nitrates* (195)
 Ternary system of potassium, cesium and calcium nitrates* (254)
 Phase diagram of the ternary system of lithium-, rubidium-, and cesium nitrates* (377)

nitride

Theoretical prediction of the thermal conductivity of uranium nitride vapor* (354)
 Lithium nitride (Li_3N): standard enthalpy of formation by solution calorimetry* (467)

nitrides

Equation of state and transport properties of uranium and plutonium nitrides in the liquid region* (578)
 Densities of liquid carbides and nitrides of uranium and plutonium* (579)
 X-ray spectral study of the carbides and nitrides of group IV and V metals* (582)

nitrogen

Solubility of nitrogen in multicomponent nickel base alloys* (100)
 Kinetics of reaction of liquid lithium with oxygen and nitrogen* (106)
 Kinetic peculiarities of nitrogen dissolving in iron-chromium and iron-chromium-nickel melts* (172)
 Vaporization of uranium-carbon-nitrogen system* (251)
 P-T diagram of the system uranium-nitrogen* (373)
 Nitrogen solubility in liquid nickel-niobium alloys* (532)
 Solubility of nitrogen in liquid lithium and thermal decomposition of solid Li_3N * (702)

nitrosyl

Structural study and properties of the alkali metal, nitrosyl, and ammonium hepta- and octafluorocuranates(VI)* (66)

NMR

Pulse NMR in solids: chemical shift, lead fluoride, and thorium hydride* (363)
 NMR study on structural microinhomogeneity of fluoride-oxide melts* (463)

No

Stable inorganic matrix materials for high temperature batteries. Final report No. 1, 2 June 1967--2 February 1968* (23)
 Lithium anode limited-cycle battery investigation. Quarterly technical progress report No. 5* (37)
 Lithium anode limited-cycle battery investigation. Quarterly technical progress report No. 3* (38)

TITLE INDEX

No

Environmental sensitivity of structural metals: liquid metal embrittlement. Annual technical progress report No. 2, 25 Jun 1970--31 May 1971* (95)
 Lithium--silver chloride secondary battery investigation. Quarterly technical progress report No. 2* (111)
 Structure of NO_3^- radical in molten monovalent metal nitrates* (195)
 Environmental sensitivity of structural metals: liquid metal embrittlement. Annual technical progress report No. 4, 1 Jun 1972--31 May 1973* (211)
 Environmental sensitivity of structural metals: liquid metal embrittlement. Annual technical progress report No. 2, 25 Jun 1970--31 May 1971* (212)
 Nonaqueous electrolyte systems: ionic transport in nonaqueous media. Interim technical report No. 7* (246)
 $\text{LiNO}_3\text{-CsNO}_3\text{-Ca(NO}_3)_2$ and $\text{NaNO}_3\text{-CsNO}_3\text{-Ca(NO}_3)_2$ system* (255)
 Solubility of $\text{Sr(NO}_3)_2$ in the $\text{LiNO}_3\text{-KNO}_3$ melt of variable composition* (351)
 Lithium--nickel fluoride secondary battery investigation. Quarterly technical progress report No. 1, 27 Apr--27 Jul 1966* (381)
 Phase diagram of $\text{NaNO}_3\text{-Nd(NO}_3)_3$ system* (667)
 Phase diagrams of the $\text{Mg(NO}_3)_2\text{-Nd(NO}_3)_3$ system* (668)
 X-ray diffraction and infrared: Raman spectroscopic study of the hydrogen dinitrate ion, $\text{O}_2\text{N-H-ONO}_2^-$, in cesium hydrogen dinitrate* (695)
 Research on a high-energy nonaqueous battery system. Quarterly progress report No. 1, 1 Dec 1965--1 Mar 1966* (720)
 Research on electrochemical parameters for nonaqueous batteries: task I. Quarterly progress report No. 3* (727)

non

Characterisation of the chemical activities of non-metals dissolved in liquid alkali metals* (63)
 Characterisation of the chemical activities of non-metals dissolved in liquid alkali metals* (64)
 Calculation of vapor pressures of oxide fuels up to 5000 K in equilibrium and non-equilibrium evaporation* (72)
 Magnetic susceptibility and electronic structure in metallic and non-metallic liquid cesium-gold and cesium-antimony alloys* (190)
 Solution behavior of hydrogen isotopes and other non-metallic elements in liquid lithium* (404)
 Thermodynamics of non-stoichiometry. A model for $\text{UC}_{x/2}$ * (442)
 3. Zirconium compounds with non-metals* (464)
 Radiative and non-radiative transitions of rare-earth ions in glasses* (538)
 Studies of non-stoichiometric oxides by thermoanalytical methods* (626)

nonaqueous

Composition of nonaqueous solutions of potential use in high energy density batteries. Final report, 15 Nov 1969--14 Nov 1972* (119)
 Nonaqueous electrolyte systems: ionic transport in nonaqueous media. Interim technical report No. 7* (246)

TITLE INDEX

nonaqueous

Electrochemical characterization of nonaqueous systems for secondary battery application. Quarterly report, February--April 1968* (380)

Research on a high-energy nonaqueous battery system. Quarterly progress report No. 1, 1 Dec 1965--1 Mar 1966* (720)

Research on electrochemical parameters for nonaqueous batteries: task I. Quarterly progress report No. 3* (727)

noncrystalline

Basic research in crystalline and noncrystalline ceramic systems. Annual report, May 1, 1975--April 1, 1976* (728)

nondispersing

Measurement of mass transfer coefficients in a mechanically agitated, nondispersing contactor operating with a molten mixture of $\text{LiF--BeF}_2\text{--ThF}_4$ and molten bismuth* (79)

nonequilibrium

Calculation of vapor pressures of oxide fuels up to 5,000 K for equilibrium and nonequilibrium evaporation* (73)

Nonisothermal

Nonisothermal flow of liquid metals over the surface of solid ones* (101)

nonmetals

Improved model for solutions of nonmetals in liquid alkali metals. Calculation of enthalpy of solution and electrical resistivity* (486)

nonstoichiometric

Entropy change for the lambda-type transition in nonstoichiometric $\text{U}_{40}\text{O}_{(9-y)}$ * (260)

Basic mechanisms providing oxidation resistance in structural metals at high temperatures. II. The defect structure of nonstoichiometric rutile (TiO_{2-x}) and nonstoichiometric rutile containing 1 to 10 mole % $\text{Ce}_{20}\text{O}_{60}$. Final report 21 Dec 1973-20 Dec 1974* (542)

nonstoichiometry

High temperature chemistry of ceramic nuclear fuels with emphasis on nonstoichiometry* (448)

Nonstoichiometry in cubic uranium sesquinitride* (611)

normal

Determination of the Np(IV)--Np(III) normal potential in fused LiCl--CsCl (55-45%) by absorption spectrophotometry* (386)

Nov

Composition of nonaqueous solutions of potential use in high energy density batteries. Final report, 15 Nov 1969--14 Nov 1972* (119)

November

Lithium--silver chloride secondary battery investigation. Final report, December 1963--November 1964* (118)

Second specialist meeting on sodium/fuel interaction in fast reactors, Ispra, 21-23 November 1973. Sponsored by the Committee on the Safety of Nuclear Installations, OECD Nuclear Energy Agency, Paris, France* (272)

Analytical Chemistry Division annual progress report for period ending November 30, 1975* (423)

Chemistry Division annual progress report for period ending November 1, 1975* (729)

TITLE INDEX

November

Chemistry Division annual progress report for period ending November 1, 1975* (730)

Np

Redox and chemical properties of U, Np and Pu in molten alkali chlorides* (360)

Determination of the Np(IV)--Np(III) normal potential in fused LiCl-CsCl (55-45%) by absorption spectrophotometry* (386)

Optical spectroscopy of ^{237}Np in CaF_2 . Final report* (718)

NRI

Twenty years of research in the processing of radioactive fuel and neutralization of radioactive waste at the Nuclear Research Institute (NRI)* (358)

Nuclear

Nuclear quadrupole resonance spectra of HgCl_2 under pressure* (31)

Review of partitioning proposals for spent nuclear fuels* (69)

Safeguarding nuclear materials. Vol. 2* (131)

Israel Nuclear Society symposium on "Nuclear fuel in Israel", Weizmann Institute of Science, Rehovot, 1.7.75* (226)

Second specialist meeting on sodium/fuel interaction in fast reactors, Ispra, 21-23 November 1973. Sponsored by the Committee on the Safety of Nuclear Installations, OECD Nuclear Energy Agency, Paris, France* (272)

Investigation into americium extraction process from residues of irradiated nuclear fuel after its fluorination by $\text{SiF}_4\text{-NaF}$ and $\text{SiF}_4\text{-CaF}_2$ melts of eutectic composition* (308)

Chemical reprocessing of spent nuclear fuels* (322)

Preparation and thermochemical stability of uranium-zirconium-carbonitrides. A contribution to nuclear fuels for advanced high-temperature reactors* (333)

Experience of the Central Control Laboratory (CCL) in accounting for and controlling nuclear material in Czechoslovakia* (349)

Twenty years of research in the processing of radioactive fuel and neutralization of radioactive waste at the Nuclear Research Institute (NRI)* (358)

Advances in nuclear science and technology. Vol. 9* (448)

Study of the retention of Ba in UO_2 nuclear fuel particles by ZrO_2 as getter* (470)

Extension of vapour pressure measurements of nuclear fuels (U,PuO_2 and UO_2) to 7000 K for fast reactor safety analysis* (475)

Vapour pressure measurements to 7,000 K and equation of state of oxide fuels for fast reactor safety analysis. I. Extension of vapour pressure measurements of nuclear fuels (U,PuO_2 and UO_2) to 7,000 K* (477)

Nuclear quadrupole relaxation and viscosity in liquid metals* (564)

Nuclear reactor containments* (569)

Phase relations, crystal structures and physical properties of nuclear fuels* (612)

Thermal properties of nuclear fuel ceramics* (613)

Analytical methods for fissionable materials in the nuclear fuel cycle. Progress report, July 1, 1975--September 30, 1976* (690)

TITLE INDEX

nuclear

Accurate nuclear parameters derived from the Moessbauer spectrum of thulium metal* (696)
 Standard methods for chemical, mass spectrometric, and spectrochemical analysis of nuclear-grade uranium dioxide powders and pellets* (711)
 Standard methods for chemical, mass spectrometric, and spectrochemical analysis of nuclear-grade plutonium dioxide powders and pellets* (712)
 Standard methods for chemical, mass spectrometric, and spectrochemical analysis of nuclear-grade mixed oxides (U,Pu) $O_{2.5}$ * (713)
 Standard methods for chemical, mass spectrometric, spectrochemical, nuclear, and radiochemical analysis of nuclear-grade plutonium metal* (714)
 Nuclear spectroscopy. Laboratory of chemistry and radiochemistry* (723)

Nukleare

Progress report of the Teilinstitut Nukleare Festkoerperphysik. (1.6.1974--31.5.1975)* (567)

O

Calculation of the oxygen potential of (U,Pu) O_{2-x} with defect models and measured oxygen potential data in the temperature range from 1,000 to 1,700 K* (74)
 Experimental determination of formation heat of gaseous Ga_{2O} and In_{2O} * (105)
 High-temperature microcalorimetric measurements of partial molar enthalpy of solution, $\Delta H(O_{2.5})$, in plutonium + oxygen* (107)
 Chemical separation and spectrographic estimation of rare earths and yttrium in $PuO_{2.5}$ and (U,Pu) $O_{2.5}$ * (125)
 Cs--U--O phase diagram and its application to uranium--plutonium oxide fast reactor fuel pins* (175)
 Electronic conductivity and polarization effects of solid $ThO_{2.5}-Y_{2.5}O_{3.5}$ electrolytes and of oxygen containing Va-metal electrodes* (193)
 Raman spectra of $Al_{2.5}O_{3.5}$ solutions in molten cryolite and other aluminum fluoride containing melts* (203)
 Fine definition of ir spectra from high temperature interactions of U + $O_{2.5}$. Part II. Final report, 1 July 1974--30 June 1975* (218)
 Calculation and discussion of state diagrams in the system Pu-F-O-H* (222)
 Thermodynamic study of the U-O-C system by measurement of the carbon monoxide pressures at equilibrium* (235)
 High temperature thermal expansion of $ThO_{2.5}$, MgO and $Y_{2.5}O_{3.5}$ by X-ray diffraction* (242)
 Lambda-type heat capacity anomalies in $U_{3.5}O_{8.5}$ * (262)
 Influence of small additions to the preparation and the properties of $UO_{2.5}$ ceramics. Part A. $UO_{2.5}/Gd_{2.5}O_{3.5}$ * (266)
 Study of kinetics of monovalent indium ions formation in the $In^{+}O-InCl_{3.5}-LiCl-H_{2.5}O$ system* (341)
 Wetting of certain refractories of the system MgO- $Al_{2.5}O_{3.5}$ 3%-ZrC $_{2.5}$ with metallic melts* (344)

TITLE INDEX

O

- Phase equilibria in the $\text{PbO-GeO}_2\text{-V}_2\text{O}_5$ ternary system* (394)
- Synthesis and characterization of a new uranium(V) compound: $\text{Hf}_3\text{O}_8 + \text{UF}_6$ - * (409)
- Phase relations in the alkaline earth fluoride-yttria systems (in three parts). 1. The $\text{CaF}_2\text{-Y}_2\text{O}_3$ phase diagram* (468)
- Phase relations in the alkaline earth fluoride-yttria systems (in three parts). 2. The $\text{BaF}_2\text{-Y}_2\text{O}_3$ phase diagram* (469)
- Extension of vapour pressure measurements of nuclear fuels $(\text{U,Pu})\text{O}_2$ and UO_2 to 7000 K for fast reactor safety analysis* (475)
- Vapour pressure measurements to 7,000 K and equation of state of oxide fuels for fast reactor safety analysis. I. Extension of vapour pressure measurements of nuclear fuels $(\text{U,Pu})\text{O}_2$ and UO_2 to 7,000 K* (477)
- Evaporation behavior and high-temperature thermal analysis of substoichiometric plutonium oxide in the composition range from $\text{O/Pu} = 1.51$ to 2.00 * (482)
- Preparational, structural and vibrational study of $\text{M}_2\text{U}_2\text{O}_5\text{Cl}_4$, $2\text{H}_2\text{O}$ ($\text{M} = \text{Rb, Cs}$). Detection of a tetranuclear anion $[(\text{UO}_2)_2(\text{Cl})_2(\text{H}_2\text{O})_2]^{4-}$ * (498)
- Contribution to the establishment of the phase diagram for the U--O--C system by means of x-ray diffractometry at high temperatures and under controlled pressure* (503)
- Measurement of the cell voltage of $\text{ThO}_2\text{-Y}_2\text{O}_3$ - solid electrolyte cells in the range of mixed conduction* (535)
- Basic mechanisms providing oxidation resistance in structural metals at high temperatures. II. The defect structure of nonstoichiometric rutile (TiO_2) and nonstoichiometric rutile containing 1 to 10 mole % Cb_2O_6 . Final report 21 Dec 1973-20 Dec 1974* (542)
- Oxygen potential and vacancy structure of the mixed oxide $(\text{UPu})\text{O}_{2-x}$ * (565)
- Total pressures of uranium- and plutonium-bearing species above the U--Pu--C system* (618)
- Thermodynamic properties of NbO , $\text{Fe}_{0.95}\text{O}$, CoO , NiO , Cu_2O and CuO oxides as comparative electrodes in electromotive forces method* (658)
- X-ray diffraction and infrared: Raman spectroscopic study of the hydrogen dinitrate ion, $(\text{O}_2\text{NO-H-CNO})^+$, in cesium hydrogen dinitrate* (695)
- Measurement of interfacial energies in the $\text{UO}_2\text{-Gd}_2\text{O}_3$ system* (697)
- Standard methods for chemical, mass spectrometric, and spectrochemical analysis of nuclear-grade mixed oxides $(\text{U,Pu})\text{O}_2$ * (713)
- octafluorohafnates
Preparation of lithium octafluorohafnates* (330)
- octafluorouranates
Structural study and properties of the alkali metal, nitrosyl, and ammonium hepta- and octafluorouranates(VI)* (66)

TITLE INDEX

October

Collection of gamma spectra data of irradiated light water moderated reactor spent fuel and study of the applicability of the methods for fuel identification. Final report for the period 1 February 1972--October 1975* (453)

OECD

Second specialist meeting on sodium/fuel interaction in fast reactors, Ispra, 21-23 November 1973. Sponsored by the Committee on the Safety of Nuclear Installations, OECD Nuclear Energy Agency, Paris, France* (272)

ones

Nonisothermal flow of liquid metals over the surface of solid ones* (101)

ONO

X-ray diffraction and infrared: Raman spectroscopic study of the hydrogen dinitrate ion, $(O\text{sub } 2\text{NO-H-ONO}\text{sub } 2)\text{sup } -$, in cesium hydrogen dinitrate* (695)

operating

Measurement of mass transfer coefficients in a mechanically agitated, nondispersing contactor operating with a molten mixture of $LiF\text{--}BeF\text{sub } 2\text{--}ThF\text{sub } 4$ and molten bismuth* (79)

Optical

Optical spectra of the difluoride, dichloride, and trichloride ions in the matrix-isolated $M\text{sup } +F\text{sub } 2\text{sup } -$, $M\text{sup } +Cl\text{sub } 2\text{sup } -$, and $M\text{sup } +Cl\text{sub } 3\text{sup } -$ species* (16)

Optical spectra of the dibromide and diiodide ions in the matrix-isolated $M\text{sup } +Br\text{sub } 2\text{sup } -$ and $M\text{sup } +I\text{sub } 2\text{sup } -$ species* (17)

Optical absorption spectrum of $UO\text{sub } 2\text{sup } +$ doped LiF * (30)

Optical absorption Zeeman spectroscopy of rare earth orthochromites. II. Theory* (122)

Optical absorption Zeeman spectroscopy of rare earth orthochromites. I. $ErCrO\text{sub } 3$ * (123)

Structure of the biexciton molecule and optical transition selection rules in zincblende and wurtzite materials* (186)

Optical properties of rare earth trifluorides* (484)

Effect of luminescence centers on optical absorption, excitation and fluorescence spectra of $CsCl:Pb$ * (491)

Optical spectroscopy of $\text{sup } 237\text{Np}$ in $CaF\text{sub } 2$. Final report* (718)

optimization

Approximation of tabulated thermodynamic data for purposes of optimization of two-phase liquid-metal power cycles* (343)

optimizing

Parameter for optimizing the synthesis of $(U,Pu)C$ * (676)

orbitals

Partly filled shells constituting anti-bonding orbitals with higher ionization energy than their bonding counterparts* (280)

Origin

Origin of excess entropy of alkali metals halides solid solutions* (390)

ORNL

Molten carbonate fuel cell research at ORNL* (71)

TITLE INDEX

orthochromites

Optical absorption Zeeman spectroscopy of rare earth orthochromites. II. Theory* (122)
 Optical absorption Zeeman spectroscopy of rare earth orthochromites. I. ErCrO_3 * (123)

Orthorhombic

Vibrational spectra of transition metal hexafluoride crystals. I. Orthorhombic MoF_6 , WF_6 , UF_6 neat crystals. Technical report* (51)

oscillator

Use of oscillator strengths of "'supersensitive'" ion transfers of lanthanides for calculation of some parameters determining the intensity of their absorption spectra* (511)

Osub

Thermodynamic properties of $(\text{U}, \text{Pu})\text{O}_{(2+x)}$ described by a cluster model* (133)
 Entropy change for the lambda-type transition in nonstoichiometric $\text{U}_{(4-y)}\text{O}_{(9-y)}$ * (260)
 Phase transition in doped $\text{U}_{(4-y)}\text{O}_{(9-y)}$. I. Heat capacity measurement* (261)
 Thermal conductivity and diffusivity of $\text{U}_{(4-y)}\text{O}_{(9-y)}$ from 250 to 415 K* (459)

OTeF

Covalent compounds of uranium of the type $\text{F}_{(x)}\text{U}(\text{OTeF}_{(5)}\text{O})_{(6-x)}$ * (571)

outgassing

Investigation into outgassing from molybdenum during melting* (588)

output

Accurate procedure to safeguard the fissile material content of input and output solutions of reprocessing plants* (131)

overall

Theory for the derivation of a single hard-sphere potential describing the overall molecular interactions in binary liquid metal alloys as a function of temperature and composition* (525)

overpower

Preliminary assessments of carbide fuel pins during mild overpower transients in LMFBRs* (456)

oxidation

Effect of fluoride ions on the kinetics of the oxidation of uranium(IV) by cerium(IV)* (424)
 Basic mechanisms providing oxidation resistance in structural metals at high temperatures. II. The defect structure of nonstoichiometric rutile ($\text{TiO}_{(2-x)}$) and nonstoichiometric rutile containing 1 to 10 mole % $\text{Cb}_{(2)}\text{O}_{(6)}$. Final report 21 Dec 1973-20 Dec 1974* (542)

oxide

Calculation of vapor pressures of oxide fuels up to 5000 K in equilibrium and non-equilibrium evaporation* (72)
 Calculation of vapor pressures of oxide fuels up to 5,000 K for equilibrium and nonequilibrium evaporation* (73)
 Preparation and some properties of actinide(III) and lanthanide(III) oxide halides, and the enthalpies of formation of $\text{PuBr}_{(3)}(\text{s})$ and $\text{PuI}_{(3)}(\text{s})$ * (82)

TITLE INDEX

oxide

- Summaries of reports of the all-union meeting on solid body chemistry. 3--5 June 1975. III. Synthesis, phase equilibrium and use of solid oxide compounds* (114)
- Chemical effects of composition changes in irradiated oxide fuel materials. II. Fission product segregation and chemical equilibria* (169)
- Cs--U--O phase diagram and its application to uranium--plutonium oxide fast reactor fuel pins* (175)
- Anomalous quasielastic light scattering from metal oxide glasses* (183)
- Behavior of molybdenum in mixed-oxide fuel* (202)
- Thermodynamics of reactions of rare earth oxides with niobium oxide* (206)
- Solubility of magnesium oxide in alkali metal chloride melt* (265)
- Radial oxygen distribution in the fuel of irradiated mixed oxide fuel elements with different final stoichiometries* (316)
- New oxide of uranium* (325)
- Solubility of magnesium oxide in calcium oxide--calcium chloride mixtures* (352)
- Effect of basicity on ferrous oxide activity in complex oxide systems* (372)
- Fission rates, burnup, and neutron flux-fluence-spectra characterization for mixed oxide fuel experiments in the FBR-II* (416)
- NMR study on structural microinhomogeneity of fluoride-oxide melts* (463)
- Vapor pressure studies over liquid uranium oxide and uranium plutonium oxide up to 5,000 K* (474)
- Vapour pressure measurements to 7,000 K and equation of state of oxide fuels for fast reactor safety analysis. I. Extension of vapour pressure measurements of nuclear fuels (U,Pu)O₂ and UO₂ to 7,000 K* (477)
- Evaporation behavior and high-temperature thermal analysis of substoichiometric plutonium oxide in the composition range from O/Pu = 1.51 to 2.00* (482)
- Lanthanum chloride interaction with magnesium oxide* (493)
- Interaction of neodymium- and yttrium chlorides with magnesium oxide* (494)
- Interaction of thorium chloride with calcium oxide* (496)
- Thermodynamic studies of gallium--indium liquid alloys by solid state electrochemistry with oxide electrolytes* (513)
- Energy transfer between Bi³⁺ $\rightarrow$ Eu³⁺, Bi³⁺ $\rightarrow$ Sm³⁺ and UO₂²⁺ $\rightarrow$ Eu³⁺ in oxide glasses* (539)
- Oxygen potential and vacancy structure of the mixed oxide (UPu)O_{2-x}* (565)
- Electrochemical study of high-temperature stability of compounds between the rare earths and copper oxide* (634)
- Summaries of reports of the all-union meeting on solid body chemistry. 3--5 June 1975. III. Synthesis, phase equilibrium and use of solid oxide compounds* (641)
- Investigations on chemical properties of thorium oxide solid electrolyte ceramics* (652)
- Solubility of lithium oxide in liquid lithium* (701)

TITLE INDEX

oxide

High density thoria-silica-metal (III) oxide fibers* (710)

oxides

Comparison of the thermodynamic properties and high temperature chemical behavior of lanthanide and actinide oxides* (2)

Reactions of liquid sodium with transition-metal oxides. VIII. The oxides of chromium-(III), -(IV), and -(VI) and disodium chromium(VI) tetraoxide* (35)

Phase equilibrium in interlanthanide oxides systems* (49)

Contribution to the thermodynamic study of uranium and nickel mixed oxides* (149)

Decomposition, stoichiometry and structure of neptunium oxides* (171)

Production and properties of high-melting lower oxides* (191)

Thermodynamics of reactions of rare earth oxides with niobium oxide* (206)

Thermodynamic functions and vapor pressures of uranium and plutonium oxides at high temperatures* (220)

Thermochemistry of gaseous metal oxides. Final report, 1 Apr 1973--31 Dec 1975* (239)

Reaction of americium oxides with trivalent metal oxides* (303)

Influence of intermediate products on chlorination kinetics of vanadium oxides in salt melt* (356)

Thermal decomposition of solid metal oxides* (395)

Electronic properties of actinide oxides* (398)

Thermogravimetric investigation of interaction of some rare earth oxides with ammonium bifluoride* (427)

Advances in vapour pressure studies over liquid uranium plutonium oxides up to 5000K* (476)

Vapour pressure studies over liquid uranium plutonium oxides up to 5,000 K* (478)

Rare earth and actinide oxides. Thermodynamic and electron microscopy studies* (625)

Studies of non-stoichiometric oxides by thermoanalytical methods* (626)

Some structure differences of double oxides of indium and gallium with antimony(5)* (657)

Thermodynamic properties of NbO , $\text{Fe}_{0.95}\text{O}$, CoO , NiO , $\text{Cu}_{2.0}\text{O}$ and CuO oxides as comparative electrodes in electromotive forces method* (658)

Standard methods for chemical, mass spectrometric, and spectrochemical analysis of nuclear-grade mixed oxides (U , Pu) $\text{O}_{2.0}$ * (713)

oxochlorides

Solubility of rare earth and yttrium oxochlorides in molten salts - $\text{MCl}_{2.0}$ ($\text{M}=\text{Mg}$, Ca , Sr , Ba)* (495)

oxycarbides

Effect of oxygen content on the structural characteristics of uranium oxycarbides* (362)

oxychloride

Electrochemical behavior of tellurium (4) in oxychloride melts* (221)

Electrochemical reduction of molybdenum oxychloride in molten lithium chloride - potassium chloride eutectic* (515)

TITLE INDEX

oxychlorides

Study of lanthanum, yttrium and erbium oxychlorides in melts of alkali metal chlorides* (196)

oxygen

Infrared spectroscopy of the uranium/oxygen system* (12)

X-ray study of the plutonium--oxygen phase diagram at ambient temperatures up to 1100°C * (62)

Calculation of the oxygen potential of $(\text{U,Pu})\text{O}_{2-x}$ with defect models and measured oxygen potential data in the temperature range from 1,000 to 1,700 K* (74)

Kinetics of reaction of liquid lithium with oxygen and nitrogen* (106)

High-temperature microcalorimetric measurements of partial molar enthalpy of solution, $\Delta H(\text{O}_{2\text{g}})$, in plutonium + oxygen* (107)

Determination of oxygen potentials of hyperstoichiometric U--Pu dioxides in the temperature range 1,500-1,850 K* (109)

Determination of oxygen potentials of hyperstoichiometric U-Pu dioxides in the temperature range 1500-1850 K* (110)

Solubility and activity of oxygen in liquid indium and copper-indium alloys* (182)

Electronic conductivity and polarization effects of solid $\text{ThO}_{2-\text{Y}_{2\text{O}_{3\text{sub}3}}}$ electrolytes and of oxygen containing Va-metal electrodes* (193)

Radial oxygen distribution in the fuel of irradiated mixed oxide fuel elements with different final stoichiometries* (316)

Solubility of oxygen in liquid niobium* (342)

Effect of oxygen content on the structural characteristics of uranium oxycarbides* (362)

Oxygen redistribution in $\text{UO}_{2\text{g}}$ due to a temperature gradient* (401)

Tetrad effect in the extraction of the lanthanides with neutral oxygen-containing compounds* (426)

Phase analysis on simulated LWR fuel rods and its evaluation with the aid of constitution studies in the system

uranium-zirconium-oxygen* (509)

Phase analysis at simulated LWR fuel rods and their interpretation by constitution investigations in the uranium-zirconium-oxygen system* (510)

Solubility of oxygen in molten carbonates* (563)

Oxygen potential and vacancy structure of the mixed oxide $(\text{UPu})\text{O}_{2-x}$ * (565)

Determination of trace impurities of titanium and zirconium in molybdenum and tungsten and their oxygen compounds by X-ray fluorescence spectrometry* (698)

oxygenstates

Vibration spectra of rare earth oxygenstates- $\text{Ln}_{2\text{WO}_{6\text{sub}6}}$ composition* (185)

p

P--t diagram of the system uranium--nitrogen* (373)

packing

Discrete variation of the structure and texture of liquid metals formed by components with discrete packing* (177)

TITLE INDEX

papers

Abstracts of papers and communications submitted to the 5. all-union heat and mass transfer conference* (437)

Parameter

Parameter for optimizing the synthesis of (U,Pu)C* (676)

parameters

Interaction parameters in the Zn--Pb--Sn system at low zinc concentration* (436)

Investigation of the parameters influencing the microstructure of hot-pressed boron carbide* (440)

Use of oscillator strengths of "'supersensitive'" ion transfers of lanthanides for calculation of some parameters determining the intensity of their absorption spectra* (511)

Accurate nuclear parameters derived from the Moessbauer spectrum of thulium metal* (696)

Research on electrochemical parameters for nonaqueous batteries: task I. Quarterly progress report No. 3* (727)

Paris

Second specialist meeting on sodium/fuel interaction in fast reactors, Ispra, 21-23 November 1973. Sponsored by the Committee on the Safety of Nuclear Installations, OECD Nuclear Energy Agency, Paris, France* (272)

particles

Study of the retention of Ba in UO_2 nuclear fuel particles by ZrO_2 as getter* (470)

Investigation of particles interaction in liquid metals and alloys* (661)

partition

Calculation of the reduced partition function ratio for uranium hexafluoride* (285)

partitioning

Review of partitioning proposals for spent nuclear fuels* (69)

Partly

Partly filled shells constituting anti-bonding orbitals with higher ionization energy than their bonding counterparts* (280)

Pb

Study of chemical interaction in Pb_{3AlF_6} - LaF_6 - Ca_{3AlF_6} system* (26)

Speed of ultrasonic and the thermophysical properties of the liquid metals Sn, Pb, Cd and of their binary alloys Pb:Sn and Pb:Cd* (403)

Interaction parameters in the Zn--Pb--Sn system at low zinc concentration* (436)

Effect of luminescence centers on optical absorption, excitation and fluorescence spectra of $CsCl:Pb$ * (491)

Thermodynamic properties of the ternary Sn-Pb-Cd liquid metal system* (606)

PbBr

Thermogravimetric investigation of the $PbCl_2$ - $PbBr_2$ - PbI_2 and $PbCl_2$ - $PbBr_2$ - PbF_2 systems* (602)

PbCl

Phase diagram of the $ThCl_4$ - $LiCl$ - $PbCl_2$ system* (143)

Diagram of excess free energy of the $PbCl_2$ - $CsCl$ - $ZnCl_2$ system* (402)

TITLE INDEX

- PbCl**
Thermographic investigation of the $\text{PbCl}_{\text{sub } 2} - \text{PbBr}_{\text{sub } 2} - \text{PbI}_{\text{sub } 2}$ and $\text{PbCl}_{\text{sub } 2} - \text{PbBr}_{\text{sub } 2} - \text{PbF}_{\text{sub } 2}$ systems* (602)
- PbF**
Thermographic investigation of the $\text{PbCl}_{\text{sub } 2} - \text{PbBr}_{\text{sub } 2} - \text{PbI}_{\text{sub } 2}$ and $\text{PbCl}_{\text{sub } 2} - \text{PbBr}_{\text{sub } 2} - \text{PbF}_{\text{sub } 2}$ systems* (602)
- PbI**
Thermographic investigation of the $\text{PbCl}_{\text{sub } 2} - \text{PbBr}_{\text{sub } 2} - \text{PbI}_{\text{sub } 2}$ and $\text{PbCl}_{\text{sub } 2} - \text{PbBr}_{\text{sub } 2} - \text{PbF}_{\text{sub } 2}$ systems* (602)
- PbO**
Phase equilibria in the $\text{PbO} - \text{GeO}_{\text{sub } 2} - \text{V}_{\text{sub } 2}\text{O}_{\text{sub } 5}$ ternary system* (394)
Phases of high pressure in $\text{UO}_{\text{sub } 2} - \text{UO}_{\text{sub } 3} - \text{PbO}$ system* (641)
- PbWO**
Phase relations in $\text{PbWO}_{\text{sub } 4} - \text{TR}_{\text{sub } 2}(\text{WO}_{\text{sub } 4})_{\text{sub } 3}$ systems* (168)
- PbZrF**
New hexafluorozirconates(IV): $\text{BaZrF}_{\text{sub } 6}$, $\text{PbZrF}_{\text{sub } 6}$, $\text{EuZrF}_{\text{sub } 6}$, $\text{SrZrF}_{\text{sub } 6}$ * (419)
- Pd**
 $\text{R}_{\text{sub } 3}\text{T}_{\text{sub } 2}$ compounds (R = rare earth or Y; T = Rh, Pd, Pt) with the rhombohedral $\text{Er}_{\text{sub } 3}\text{Ni}_{\text{sub } 2}$ structure type* (365)
- peculiarities**
Kinetic peculiarities of nitrogen dissolving in iron-chromium and iron-chromium-nickel melts* (172)
- pellets**
Standard methods for chemical, mass spectrometric, and spectrochemical analysis of nuclear-grade uranium dioxide powders and pellets* (711)
Standard methods for chemical, mass spectrometric, and spectrochemical analysis of nuclear-grade plutonium dioxide powders and pellets* (712)
- penta**
Conversion of protactinium monocarbide to the penta- and tetra-halides* (80)
- pentafluoride**
Mass spectra and vapor composition of niobium pentafluoride* (216)
Reactivity of transition metal fluorides. IX. Protactinium pentafluoride and tetrafluoride* (465)
- pentafluorides**
Some properties of hydrogen fluoride compounds with group 5 metal pentafluorides* (170)
- pentoxide**
Some properties of molybdenum pentoxide* (305)
- Percus**
Perturbation treatment of the Percus-Yevick equation and structures of liquid magnesium and cadmium* (210)
- Periodically**
Periodically variable wall temperature effect on heat transfer in liquid metals* (437)

TITLE INDEX

perovskite

Investigation of formation thermodynamic of rare earth aluminates with perovskite structure* (326)

perovskites

Ternary UO_2 -- UO_3 -- EuO system and investigation of Eu(II) --actinide(IV)--perovskites* (50)

perturbation

Thermodynamic perturbation theory for the surface tension and ion density profile of a liquid metal* (167)

Perturbation treatment of the Percus-Yevick equation and structures of liquid magnesium and cadmium* (210)

phase

Preparation of ^{147}Pm metal and the determination of the melting point and phase transformation temperatures* (18)

Phase equilibrium in interlanthanide oxides systems* (49)

X-ray study of the plutonium--oxygen phase diagram at ambient temperatures up to 1100°C * (62)

Phase equilibria in Li -- Na -- K carbonate/aluminate systems* (77)

Phase equilibria in the system of sodium chlorides, cesium chlorides and strontium chlorides* (87)

Phase equilibria in the system of sodium chlorides, cesium chlorides and strontium chlorides* (88)

Summaries of reports of the all-union meeting on solid body chemistry. 3--5 June 1975. III. Synthesis, phase equilibrium and use of solid oxide compounds* (114)

Phase diagram of system LiCl -- KCl -- CaCrO_4 * (116)

Phase diagram for the ternary system CaCl_2 -- KCl -- CaCrO_4 * (117)

Phase diagram of the ThCl_4 -- LiCl -- PbCl_2 system* (143)

Phase diagrams of ternary systems containing sodium and potassium chlorides, thorium tetrachloride and plutonium trichloride* (148)

Phase diagrams of Li_2MoO_4 -- ZnMoO_4 , K_2MoO_4 -- ZnMoO_4 and K_2WO_4 -- ZnWO_4 systems* (160)

Phase relations in PbWO_4 -- $\text{TR}_2(\text{WO}_4)_3$ systems* (168)

Cs -- U -- O phase diagram and its application to uranium--plutonium oxide fast reactor fuel pins* (175)

Phase diagrams of LiCl , CsCl , CaCl and LiBr , CsBr , CaBr ternary systems* (256)

Phase transition in doped $\text{UO}_2(1-y)$. I. Heat capacity measurement* (261)

Phase equilibrium studies of lithium halide-containing electrolytes* (282)

Phase diagrams of BiBr_3 -- BiI_3 system* (288)

Phase diagram and properties of Nd -- As alloys* (321)

Phase diagrams of BiCl_3 -- CuCl -- TeCl_4 and CuCl -- FeCl_3 -- TeCl_4 systems* (337)

Approximation of tabulated thermodynamic data for purposes of optimization of two-phase liquid-metal power cycles* (343)

Phase diagram of the ternary system of lithium-, rubidium-, and cesium nitrates* (377)

Phase equilibria in the PbO -- GeO_2 -- V_2O_5 ternary system* (394)

TITLE INDEX

phase

- Electromodulation spectroscopy of sc and fcc phase TlCl and TlBr * (411)
- Phase equilibrium in $\text{NdCl}_3\text{-KCl}$ system* (457)
- Phase relations in the alkaline earth fluoride-yttria systems (in three parts). 1. The $\text{CaF}_2\text{-Y}_2\text{O}_3$ phase diagram* (468)
- Phase relations in the alkaline earth fluoride-yttria systems (in three parts). 2. The $\text{BaF}_2\text{-Y}_2\text{O}_3$ phase diagram* (469)
- Phase diagrams in the pseudobinary systems of Li_2BeF_4 , alkali fluoroborates and alkali fluorides* (473)
- Contribution to the establishment of the phase diagram for the U-C-C system by means of x-ray diffractometry at high temperatures and under controlled pressure* (503)
- Phase analysis on simulated LWR fuel rods and its evaluation with the aid of constitution studies in the system uranium-zirconium-oxygen* (509)
- Phase analysis at simulated LWR fuel rods and their interpretation by constitution investigations in the uranium-zirconium-oxygen system* (510)
- Theoretical concepts useful in the calculation or storage of phase diagrams of ionic systems* (552)
- Phase relations, crystal structures and physical properties of nuclear fuels* (612)
- Summaries of reports of the all-union meeting on solid body chemistry. 3--5 June 1975. III. Synthesis, phase equilibrium and use of solid oxide compounds* (641)
- Phase equilibrium study on system uranium-plutonium-tungsten-carbon* (645)
- Phase behavior and thermodynamics of U-Mo-C system* (646)
- Phase behavior and thermodynamics of U-Mo-C system* (647)
- Phase equilibria in system U-Pu-W-C * (648)
- Phase diagram of $\text{NaNO}_3\text{-Nd(NO}_3)_3$ system* (667)
- Phase diagrams of the $\text{Mg(NO}_3)_2\text{-Nd(NO}_3)_3$ system* (668)
- Phase diagrams* (679)
- Phase diagrams* (680)
- Phase equilibria and thermodynamic data for the ternary $\text{UO}_2\text{-UO}_3\text{-REO}_{1.5}$ (Ho, Tm, Lu) systems* (694)

Phases

- Phases of variable composition with LaF_3 structure in $\text{MF}_2\text{-(Y, Ln)F}_3$ systems. 1. $\text{SrF}_2\text{-(Y, Ln)F}_3$ systems (thermal characteristics, formation of berthollides)* (593)
- Phases of high pressure in $\text{UO}_2\text{-UO}_3\text{-PbO}$ system* (641)

phenomena

- Mass transport phenomena in ceramics* (401)

Photoelectron

- Photoelectron spectroscopic studies of some transition metals and alloys* (417)

photoemission

- Electronic structure of CeN studied by x-ray-photoemission spectroscopy* (28)

TITLE INDEX

physical

Chemical Engineering Division physical inorganic chemistry. Annual report, July 1975--June 1976* (58)
 Physical models of mass transport of iron and nickel in liquid sodium systems* (128)
 Physical and mechanical properties of some uranium-containing carbides* (209)
 Chemical Engineering Division, reactor safety and physical property studies annual report, July 1974--June 1975* (367)
 Physical research on liquid-metal systems. Annual report, July 1975--June 1976* (405)
 Physical--chemical studies of transuranium elements. Progress report, April 1, 1976--March 31, 1977* (501)
 Phase relations, crystal structures and physical properties of nuclear fuels* (612)
 Physical properties of thorium fluoride* (656)

Physico

Zirconium: Physico-chemical properties of its compounds and alloys* (7)
 Zirconium: Physico-chemical properties of its compounds and alloys* (464)
 Thorium: physico-chemical properties of its compounds and alloys* (679)
 Zirconium: Physico-chemical properties of its compounds and alloys* (680)

Physicochemical

Physicochemical investigation of rare earth titanoniobates, LnTiNbO_6 * (173)

Physics

Physics of semiconductors* (186)
 Physik Daten/Physics Data* (192)
 Eleventh annual seminar on theoretical physics, Johannesburg, 5-9 July 1976* (368)
 Vacuum ultraviolet radiation physics* (382)
 Vacuum ultraviolet radiation physics* (484)

Physik

Physik Daten/Physics Data* (192)

pins

Cs--U--O phase diagram and its application to uranium--plutonium oxide fast reactor fuel pins* (175)
 Preliminary assessments of carbide fuel pins during mild overpower transients in LMFBRs* (456)

planning

Practice of experiment planning in studying limited regions of multicomponent system* (546)

plants

Accurate procedure to safeguard the fissile material content of input and output solutions of reprocessing plants* (131)

platinum

Thermal decomposition of intermetallic compounds of platinum with alkaline earth and lanthanide metals at high temperatures* (399)

plutonia

Reduction of PuO_2 by CO and equilibrium partial pressures above plutonia* (304)

TITLE INDEX

Plutonium

- Plutonium 1975 and other actinides* (9)
- Thermodynamic assessment of the conversion of plutonium dioxide to plutonium monocarbide* (54)
- Thermodynamics of the conversion of plutonium dioxide to plutonium monocarbide* (55)
- X-ray study of the plutonium--oxygen phase diagram at ambient temperatures up to 1100^oC* (62)
- Plutonium 1975 and other actinides* (83)
- Segregation and vapour pressure studies in the uranium--plutonium--carbon system* (84)
- Results of research to evaluate solid plutonium nitrate as a safe shipping form* (85)
- High-temperature microcalorimetric measurements of partial molar enthalpy of solution, $\Delta H(O_{2g})$, in plutonium + oxygen* (107)
- 5th international conference on plutonium and other actinides. Book of abstracts* (109)
- Plutonium 1975 and other actinides* (110)
- Plutonium 1975 and other actinides* (133)
- Calculation of thermodynamic activity and liquidus lines of components in binary systems of plutonium trichloride with alkali and alkaline earth metal chlorides* (147)
- Phase diagrams of ternary systems containing sodium and potassium chlorides, thorium tetrachloride and plutonium trichloride* (148)
- Cs--U--O phase diagram and its application to uranium--plutonium oxide fast reactor fuel pins* (175)
- Enthalpy of uranium-plutonium carbide from 298K to the melting point* (179)
- Plutonium 1975 and other actinides* (209)
- Thermodynamic functions and vapor pressures of uranium and plutonium oxides at high temperatures* (220)
- Plutonium 1975 and other actinides* (227)
- Some constitutional studies on uranium--carbon and plutonium--carbon--rhenium and technetium systems* (228)
- Radiolytic gas generation in plutonium contaminated waste materials* (301)
- Isotope effects in the molecular spectrum of plutonium hexafluoride* (353)
- Plutonium 1975 and other actinides* (398)
- Plutonium 1975 and other actinides* (414)
- 5th international conference on plutonium and other actinides* (415)
- PuO₂ dissolution problem for LWR plutonium recycle and LMFBR fuels: fabrication and reprocessing problems and their resolution* (455)
- Vapor pressure studies over liquid uranium oxide and uranium plutonium oxide up to 5,000 K* (474)
- Plutonium 1975 and other actinides* (476)
- Vapour pressure studies over liquid uranium plutonium oxides up to 5,000 K* (478)
- Evaporation behaviour of the ternary uranium plutonium carbides* (479)
- Evaporation behaviour of the ternary uranium plutonium carbides* (480)

TITLE INDEX

plutonium

- Evaporation behaviour of the ternary uranium plutonium carbides* (481)
 Evaporation behavior and high-temperature thermal analysis of substoichiometric plutonium oxide in the composition range from $O/Pu = 1.51$ to 2.00 * (482)
 Plutonium 1975 and other actinides* (560)
 Equation of state and transport properties of uranium and plutonium carbides in the liquid region* (577)
 Equation of state and transport properties of uranium and plutonium nitrides in the liquid region* (578)
 Densities of liquid carbides and nitrides of uranium and plutonium* (579)
 Total pressures of uranium- and plutonium-bearing species above the $U-Pu-O$ system* (618)
 Plutonium 1975 and other actinides* (619)
 Enthalpy and heat capacity of liquid plutonium and uranium* (620)
 Phase equilibrium study on system uranium-plutonium-tungsten-carbon* (645)
 Spectrophotometric method for the determination of neptunium and plutonium in process solutions* (660)
 Interaction of plutonium trichloride in melts of ternary systems* (682)
 Fabrication of uranium-plutonium mixed carbide* (691)
 Standard methods for chemical, mass spectrometric, and spectrochemical analysis of nuclear-grade plutonium dioxide powders and pellets* (712)
 Standard methods for chemical, mass spectrometric, spectrochemical, nuclear, and radiochemical analysis of nuclear-grade plutonium metal* (714)
 Standard method of test for atom percent fission in uranium and plutonium fuel (mass spectrometric method)* (715)
 Standard method of test for atom percent fission in uranium and plutonium fuel (neodymium-148 method)* (716)
 Fluorination process studies of plutonium dioxide by fluid-bed* (725)

Pm

- Preparation of ^{147}Pm metal and the determination of the melting point and phase transformation temperatures* (18)

POCl

- Density and viscosity of distillable products from ZrCl_4 and HfCl_4 interaction with POCl_3 * (458)

points

- EUTECTIC DATA: safety, hazards, corrosion, melting points, compositions, and bibliography* (276)
 Some structural characteristics of rubidium and cesium in the vicinity of their melting points* (460)

polarization

- Electronic conductivity and polarization effects of solid ThO_2 - Y_2O_3 electrolytes and of oxygen containing Va-metal electrodes* (193)

polyhalides

- Thermochemistry of polyhalides. Pt.1. Caesium and rubidium dichloriodates(I)* (178)

TITLE INDEX

positron

Studies of the electronic structure of simple metals and their alloys with the aid of compton scattering and positron annihilation using lithium, magnesium, lithium-magnesium and aluminium-zinc as examples* (48)

post

Hydrodynamics of post-disassembly fuel expansions* (310)

potassium

Interaction of uranyl chloride with zirconium tetrachloride in the melt of sodium chloride and potassium chloride* (27)
 Investigation into neodymium chloride state in equilibrium with metal in eutectic melt of potassium- and lithium chlorides* (40)
 Quaternary system of potassium halogenides* (90)
 Thermographic investigation into ternary systems containing sodium and potassium chlorides, uranium trichloride and thorium tetrafluoride* (140)
 Interaction of uranium- and thorium tetrachlorides with sodium- and potassium chlorides* (144)
 Phase diagrams of ternary systems containing sodium and potassium chlorides, thorium tetrachloride and plutonium trichloride* (148)
 Interaction of lower indium chlorides with chloride and chloroindates of potassium* (174)
 Ternary system of potassium, cesium and calcium nitrates* (254)
 Electrochemical reduction of molybdenum oxychloride in molten lithium chloride - potassium chloride eutectic* (515)
 Interaction of potassium tetrafluoroborate with fluorides of potassium, lanthanum, neodymium in melt* (530)
 Investigation into reactions of hexabromotellurates of potassium, cesium and thallium(1) in melt* (556)
 Reaction of potassium superoxide in excess potassium* (590)
 Quaternary mutual system of chlorides, bromides, iodides of potassium and cesium* (630)
 Sections of quaternary mutual system consisting of sodium, potassium and cesium chlorides and bromides* (631)
 Double potassium rare earth tungstates* (639)
 Anodic behavior of cadmium and cadmium-zinc alloys in eutectic melt of potassium- and lithium chlorides* (675)

potential

Standard potential of sodium amalgam at 25°C* (32)
 Calculation of the oxygen potential of $(U,Pu)O_{2-x}$ with defect models and measured oxygen potential data in the temperature range from 1,000 to 1,700 K* (74)
 Composition of nonaqueous solutions of potential use in high energy density batteries. Final report, 15 Nov 1969--14 Nov 1972* (119)
 Structure of lithium-boron alloys and their performance as potential anodes for high power molten salt batteries. Progress report, 1 Aug 1974--30 Jun 1975* (275)
 Determination of the $Np(IV)$ -- $Np(III)$ normal potential in fused $LiCl$ - $CsCl$ (55-45%) by absorption spectrophotometry* (386)
 Theory for the derivation of a single hard-sphere potential describing the overall molecular interactions in binary liquid metal alloys as a function of temperature and composition* (525)
 Oxygen potential and vacancy structure of the mixed oxide $(UPu)O_{2-x}$ * (565)

TITLE INDEX

potentials

Determination of oxygen potentials of hyperstoichiometric U--Pu dioxides in the temperature range 1,500-1,850 K* (109)

Determination of oxygen potentials of hyperstoichiometric U-Pu dioxides in the temperature range 1500-1850 K* (110)

Electrode potentials of tantalum chlorides in a molten NaCl-KCl eutectic* (449)

Electrochemistry of niobium and tantalum. VI. Electrode potentials of tantalum chlorides in molten KCl* (451)

Potentiometric

Thermodynamic study of the binary system copper--indium. II. Potentiometric study* (291)

powders

Standard methods for chemical, mass spectrometric, and spectrochemical analysis of nuclear-grade uranium dioxide powders and pellets* (711)

Standard methods for chemical, mass spectrometric, and spectrochemical analysis of nuclear-grade plutonium dioxide powders and pellets* (712)

powdery

Quantitative determination of UO_2 , and ThO_2 in powdery mixtures by infrared spectrometry* (431)

power

Structure of lithium--boron alloys and their performance as potential anodes for high power molten salt batteries. Progress report, 1 Aug 1974--30 Jun 1975* (275)

Approximation of tabulated thermodynamic data for purposes of optimization of two-phase liquid-metal power cycles* (343)

Pr

Pressure effect on spectral properties of CeF_3 : Pr^{3+} * (677)

Practice

Practice of experiment planning in studying limited regions of multicomponent system* (546)

praseodymium

Zeeman effect and satellite carrier structure in absorption spectra of pure and doped praseodymium and europium double nitrates* (165)

precise

Half-life of ^{242}Pu from precise low-temperature heat capacity measurements* (487)

Predicted

Predicted infrared spectrum and the structure of the isolated UF_5 molecule* (350)

prediction

Theoretical prediction of the thermal conductivity of uranium carbide vapor* (124)

Prediction of the inert gas solubilities in stoichiometric molten UC_2 * (225)

Theoretical prediction of the thermal conductivity of uranium nitride vapor* (354)

Preliminary

Preliminary assessments of carbide fuel pins during mild overpower transients in LMFBRs* (456)

TITLE INDEX

Preparational

Preparational, structural and vibrational study of $M_{2U}_{5Cl_4}$, $2H_{2O}$ ($M = Rb, Cs$). Detection of a tetranuclear anion $[(UO_2)_4O_{10}Cl_8(H_2O)_2]^{4-}$ (498)

preparing

Process for preparing uranium carbonitride* (649)

pressed

Investigation of the parameters influencing the microstructure of hot-pressed boron carbide* (440)

pressure

Nuclear quadrupole resonance spectra of $HgCl_2$ under pressure* (31)

Experimental vapor pressure and critical point of sodium* (56)

Investigation of fuel vapor pressure for fast breeder safety analysis* (60)

Segregation and vapour pressure studies on the uranium-plutonium-carbon system* (83)

Segregation and vapour pressure studies in the uranium-plutonium-carbon system* (84)

Tensimetric study of pressure and composition of saturated vapor in $TlCl-InCl_3$ system* (93)

Saturation vapor pressure in iron chloride-alkaline earth metal chlorides systems* (97)

Study of saturated vapor pressure above salt melts in $CoCl_2-MCl$ systems* (98)

Temperature dependence of vapor pressure of rare earth halides and their allcys* (99)

Vacancies and melting curves of metals at high pressure* (213)

Pressure of saturated vapor of rare earth and yttrium bromides* (391)

Pressure and composition of vapors over alkali metal tetrachloroferrates in chlorine atmosphere* (433)

Determination of surface tension of a metal with high vapor pressure* (472)

Vapor pressure studies over liquid uranium oxide and uranium plutonium oxide up to 5,000 K* (474)

Extension of vapour pressure measurements of nuclear fuels $(U, Pu)O_{2.5}$ and UO_2 to 7000 K for fast reactor safety analysis* (475)

Advances in vapour pressure studies over liquid uranium plutonium oxides up to 5000K* (476)

Vapour pressure measurements to 7,000 K and equation of state of oxide fuels for fast reactor safety analysis. I. Extension of vapour pressure measurements of nuclear fuels $(U, Pu)O_{2.5}$ and UO_2 to 7,000 K* (477)

Vapour pressure studies over liquid uranium plutonium oxides up to 5,000 K* (478)

Contribution to the establishment of the phase diagram for the U-O-C system by means of x-ray diffractometry at high temperatures and under controlled pressure* (503)

Pressure and entropy calculations for liquid Na, K and Rb along the melting line* (584)

Phases of high pressure in $UO_{2.5}-UO_3-PbO$ system* (641)

TITLE INDEX

Pressure

Pressure effect on spectral properties of CeF_3Pr (677)

pressures

Calculation of vapor pressures of oxide fuels up to 5000 K in equilibrium and non-equilibrium evaporation* (72)
 Calculation of vapor pressures of oxide fuels up to 5,000 K for equilibrium and nonequilibrium evaporation* (73)
 Thermodynamic functions and vapor pressures of uranium and plutonium oxides at high temperatures* (220)
 Thermodynamic study of the U-O-C system by measurement of the carbon monoxide pressures at equilibrium* (235)
 Partial pressures, activities and evaporation heat in the KF-ZrF_4 system* (295)
 Thermophysical and electrophysical properties of uranium hexafluoride at temperatures of $(1-11) \times 10^3$ degree K and pressures of 0.1--100 atm* (302)
 Reduction of PuO_2 by CO and equilibrium partial pressures above plutonia* (304)
 Behavior of the AB-type compounds at high pressures and high temperatures* (421)
 Density and enthalpy of liquid alkaline metals at pressures up to 1500 bar* (599)
 Total pressures of uranium- and plutonium-bearing species above the U--Pu--O system* (618)

Proceedings

Proceedings of the 12th rare earth research conference. Vol. I* (11)
 Proceedings of the international conference on liquid technology in energy production* (56)
 Proceedings of the 12th rare earth research conference. Vol. II* (65)
 Proceedings of the international conference on liquid technology in energy production* (103)
 Proceedings of GIREDMET* (136)
 Proceedings of the symposium and workshop on advanced battery research and design, March 22--24, 1976* (269)
 Proceedings of the 12th rare earth research conference. Vol. II* (425)
 Proceedings of the international conference on liquid technology in energy production* (590)

products

Use of fluoride melts for separation of some fission products and americium from mixtures of different fluorides being the simulators of residues from fast reactor fuel fluorination* (307)
 The spectral properties of uranium hexafluoride and its thermal decomposition products* (345)
 Influence of intermediate products on chlorination kinetics of vanadium oxides in salt melt* (356)
 Density and viscosity of distillable products from ZrCl_4 and HfCl_4 interaction with POCl_3 * (458)
 Separation of alkaline earth from alkali metal fission products in the SOLAR on-line mass spectrometer* (534)

profile

Thermodynamic perturbation theory for the surface tension and ion density profile of a liquid metal* (167)

TITLE INDEX

programme

Study of mass transfer by liquid sodium in closed loop systems.
Sodium technology development programme* (324)

programs

Energy programs: molten-salt reactor program* (423)
Summary of U.S. LMFBR programs on high temperature structural
design and associated materials testing* (731)

progress

Lithium anode limited-cycle battery investigation. Quarterly
technical progress report No. 5* (37)
Lithium anode limited-cycle battery investigation. Quarterly
technical progress report No. 3* (38)
Mechanical properties test data for structural materials.
Quarterly progress report for period ending April 30, 1977* (57)
LMFBR Fuel Reprocessing Program progress report for period April
1 to June 30, 1976* (94)
Environmental sensitivity of structural metals: liquid metal
embrittlement. Annual technical progress report No. 2, 25 Jun
1970--31 May 1971* (95)
Lithium--silver chloride secondary battery investigation.
Quarterly technical progress report No. 2* (111)
Molten-salt reactor program. Semiannual progress report for
period ending February 29, 1976* (176)
Environmental sensitivity of structural metals: liquid metal
embrittlement. Annual technical progress report No. 4, 1 Jun
1972--31 May 1973* (211)
Environmental sensitivity of structural metals: liquid metal
embrittlement. Annual technical progress report No. 2, 25 Jun
1970--31 May 1971* (212)
Molten-salt reactor program. Semiannual progress report for
period ending February 29, 1976* (237)
Radiochemistry Division annual progress report for 1973* (271)
Structure of lithium--boron alloys and their performance as
potential anodes for high power molten salt batteries. Progress
report, 1 Aug 1974--30 Jun 1975* (275)
Lithium--nickel fluoride secondary battery investigation.
Quarterly technical progress report No. 1, 27 Apr--27 Jul
1966* (381)
Progress in the spectroscopy of rare earth and transition metal
ions* (412)
Analytical Chemistry Division annual progress report for period
ending November 30, 1975* (423)
Mechanical properties test data for structural materials.
Quarterly progress report for period ending April 30, 1977* (471)
Progress in fluorine calorimetry* (500)
Physical--chemical studies of transuranium elements. Progress
report, April 1, 1976--March 31, 1977* (501)
Fundamental studies of metal fluorination reactions. Progress
report, May 1, 1976--April 31, 1977* (533)
Progress report of the Teilinstitut Nukleare Festkoerperphysik.
(1.6.1974--31.5.1975)* (567)
Analytical methods for fissionable materials in the nuclear fuel
cycle. Progress report, July 1, 1975--September 30, 1976* (690)
Mechanical properties test data for structural materials.
Quarterly progress report for period ending April 30, 1977* (692)

TITLE INDEX

progress

Research on a high-energy nonaqueous battery system. Quarterly progress report No. 1, 1 Dec 1965--1 Mar 1966* (720)
 Electrochemical methods for monitoring purity of molten salts. Semiannual progress report, July--December 1975* (721)
 Thorium utilization program. Quarterly progress report for the period ending May 31, 1977* (722)
 Progress report 1971. Vol. 1* (723)
 Research on electrochemical parameters for nonaqueous batteries: task I. Quarterly progress report No. 3* (727)
 Chemistry Division annual progress report for period ending November 1, 1975* (729)
 Chemistry Division annual progress report for period ending November 1, 1975* (730)

project

Space Processing Applications Rocket project, SPAR 1. Final report* (536)

Projection

Projection-thermographic method for study of ternary and ternary mutual systems* (518)

property

Chemical Engineering Division, reactor safety and physical property studies annual report, July 1974--June 1975* (367)
 Anomalous thermal property behaviour of uranium at low temperatures* (560)

proposals

Review of partitioning proposals for spent nuclear fuels* (69)

protactinium

Conversion of protactinium monocarbide to the penta- and tetra-halides* (80)
 Reactivity of transition metal fluorides. IX. Protactinium pentafluoride and tetrafluoride* (465)
 Diffuse reflectance spectra of protactinium(IV), uranium(IV) and neptunium(IV) tetrachlorides and tetraformates* (562)

prototypic

Metallurgical characterization of prototypic tubesheet forging* (471)

providing

Basic mechanisms providing oxidation resistance in structural metals at high temperatures. II. The defect structure of nonstoichiometric rutile (TiO_{2-x}) and nonstoichiometric rutile containing 1 to 10 mole % CeO_{2-x} . Final report 21 Dec 1973-20 Dec 1974* (542)

pseudobinary

Phase diagrams in the pseudobinary systems of Li_2BeF_4 , alkali fluoroborates and alkali fluorides* (473)

Pt

Electronic structure of highly symmetric complexes of lanthanoids and actinoids. Pt. 3. 'Electronic' Raman spectrum of $\text{Cs}_2\text{U(IV)Cl}_6$ * (13)
 Electronic structure of highly symmetric complexes of lanthanoids and actinoids. Pt. 4. The electronic Raman spectrum of $\text{Cs}_2\text{NaEu(III)Cl}_6$ * (14)
 Thermochemistry of polyhalides. Pt. 1. Caesium and rubidium dichloriodates(I)* (178)

TITLE INDEX

Pt

R₃T₂ compounds (R = rare earth or Y; T = Rh, Pd, Pt) with the rhombohedral Er₃Ni₂ structure type* (365)

Pu

Calculation of the oxygen potential of (U,Pu)O_{2-x} with defect models and measured oxygen potential data in the temperature range from 1,000 to 1,700 K* (74)
 Determination of oxygen potentials of hyperstoichiometric U--Pu dioxides in the temperature range 1,500-1,850 K* (109)
 Determination of oxygen potentials of hyperstoichiometric U-Pu dioxides in the temperature range 1500-1850 K* (110)
 Chemical separation and spectrographic estimation of rare earths and yttrium in PuO₂ and (U,Pu)O₂* (125)
 Thermodynamic properties of (U,Pu)O_{2-x} described by a cluster model* (133)
 Distribution coefficient for U and Pu in the system (UPu)C + (UPu)₂C₃* (134)
 Calculation and discussion of state diagrams in the system Pu-F-O-H* (222)
 Anomalous melting of f electron metals (with attention to Pu)* (318)
 Redox and chemical properties of U, Np and Pu in molten alkali chlorides* (360)
 Specific heat of PuC_(1-x) and Pu₂C₃ from 10 to 300 K* (414)
 Specific heat of PuC_(1-x) and Pu₂C₃ from 10 to 300 K* (415)
 Extension of vapour pressure measurements of nuclear fuels (U,Pu)O₂ and UO₂ to 7000 K for fast reactor safety analysis* (475)
 Vapour pressure measurements to 7,000 K and equation of state of oxide fuels for fast reactor safety analysis. I. Extension of vapour pressure measurements of nuclear fuels (U,Pu)O₂ and UO₂ to 7,000 K* (477)
 Evaporation behavior and high-temperature thermal analysis of substoichiometric plutonium oxide in the composition range from O/Pu = 1.51 to 2.00* (482)
 Half-life of ²⁴²Pu from precise low-temperature heat capacity measurements* (487)
 Total pressures of uranium- and plutonium-bearing species above the U--Pu--O system* (618)
 Phase equilibria in system U--Pu--W--C* (648)
 Parameter for optimizing the synthesis of (U,Pu)C* (676)
 Standard methods for chemical, mass spectrometric, and spectrochemical analysis of nuclear-grade mixed oxides (U,Pu)O₂* (713)

PuEr

Preparation and some properties of actinide(III) and lanthanide(III) oxide halides, and the enthalpies of formation of PuBr₃(s) and PuI₃(s)* (82)

PuCsub

Specific heat of PuC_(1-x) and Pu₂C₃ from 10 to 300 K* (414)
 Specific heat of PuC_(1-x) and Pu₂C₃ from 10 to 300 K* (415)

TITLE INDEX

- PuF**
 Molten salt chemistry : Part IV - Solubility behaviour of PuF_3 in fluoride salts of interest in molten salt reactor technology* (655)
- PuI**
 Preparation and some properties of actinide(III) and lanthanide(III) oxide halides, and the enthalpies of formation of $\text{PuBr}_3(\text{s})$ and $\text{PuI}_3(\text{s})$ * (82)
- Pulse**
 Pulse NMR in solids: chemical shift, lead fluoride, and thorium hydride* (363)
- PuN**
 Fabrication and high temperature thermodynamic and transport properties of PuN* (9)
- PuO**
 Thermal expansion of $^{238}\text{PuO}_2$ * (15)
 Chemical separation and spectrographic estimation of rare earths and yttrium in PuO_2 and $(\text{U}, \text{Pu})\text{O}_2$ * (125)
 Heat capacity of $^{242}\text{PuO}_2$ from 12 to 350°K and of $^{244}\text{PuO}_2$ from 4 to 25°K. Entropy, enthalpy, and Gibbs energy of formation of PuO_2 at 298.15°K* (184)
 Reduction of PuO_2 by CO and equilibrium partial pressures above plutonia* (304)
 PuO_2 dissolution problem for LWR plutonium recycle and LMFBR fuels: fabrication and reprocessing problems and their resolution* (455)
- pure**
 Interactions in inorganic molecular crystals - electronic spectra of ReF_6 pure and mixed crystals. Technical report, 1 Jul--31 Dec 1975* (53)
 Zeeman effect and satellite carrier structure in absorption spectra of pure and doped praseodymium and europium double nitrates* (165)
- purity**
 Electrochemical methods for monitoring purity of molten salts. Semiannual progress report, July--December 1975* (721)
- purposes**
 Approximation of tabulated thermodynamic data for purposes of optimization of two-phase liquid-metal power cycles* (343)
- quadrupole**
 Nuclear quadrupole resonance spectra of HgCl_2 under pressure* (31)
 Nuclear quadrupole relaxation and viscosity in liquid metals* (564)
- Quantitative**
 Quantitative study of the chemical equilibria in solution in molten LiCl-KCl (70-30 mol.%) by absorption spectrophotometry. Application to neptunium* (385)
 Quantitative determination of UO_2 , and ThO_2 in powdery mixtures by infrared spectrometry* (431)
- Quarter**
 Quarter mutual system of chlorides, bromides, iodides of sodium and rubidium* (629)

TITLE INDEX

Quarterly

- Lithium anode limited-cycle battery investigation. Quarterly technical progress report No. 5* (37)
- Lithium anode limited-cycle battery investigation. Quarterly technical progress report No. 3* (38)
- Mechanical properties test data for structural materials. Quarterly progress report for period ending April 30, 1977* (57)
- Lithium--silver chloride secondary battery investigation. Quarterly technical progress report No. 2* (111)
- Electrochemical characterization of nonaqueous systems for secondary battery application. Quarterly report, February--April 1968* (380)
- Lithium--nickel fluoride secondary battery investigation. Quarterly technical progress report No. 1, 27 Apr--27 Jul 1966* (381)
- Mechanical properties test data for structural materials. Quarterly progress report for period ending April 30, 1977* (471)
- Mechanical properties test data for structural materials. Quarterly progress report for period ending April 30, 1977* (692)
- Research on a high-energy nonaqueous battery system. Quarterly progress report No. 1, 1 Dec 1965--1 Mar 1966* (720)
- Thorium utilization program. Quarterly progress report for the period ending May 31, 1977* (722)
- Research on electrochemical parameters for nonaqueous batteries: task I. Quarterly progress report No. 3* (727)

quasielastic

- Anomalous quasielastic light scattering from metal oxide glasses* (183)

Quaternary

- Quaternary mutual system from fluorides and chlorides of lithium, sodium and strontium* (89)
- Quaternary system of potassium halogenides* (90)
- Quaternary mutual system of chlorides, bromides, iodides of potassium and cesium* (630)
- Sections of quaternary mutual system consisting of sodium, potassium and cesium chlorides and bromides* (631)
- Quaternary mutual system of chlorides, bromides, iodides of rubidium and cesium* (640)

r

- Raman and i.r.-emission studies of some tungstate- and molybdate-containing melts* (102)
- Study of the compounds $M\text{UO}_2\text{F}_5$ ($M = \text{K, Rb, Cs, NH}_4$) by i.r. absorption and Raman diffusion spectrophotometry. III. Effect of cation on vibration frequency of the ion $\text{UO}_2\text{F}_5^{3-}$ * (127)
- R_3T_2 compounds ($\text{R} = \text{rare earth or Y; T} = \text{Rh, Pd, Pt}$) with the rhombohedral Er_3Ni_2 structure type* (365)
- Compounds with the structure of ThMn_{12} type in the R--Fe--Al systems* (671)

Radial

- Radial oxygen distribution in the fuel of irradiated mixed oxide fuel elements with different final stoichiometries* (316)

radiation

- Vacuum ultraviolet radiation physics* (382)
- Vacuum ultraviolet radiation physics* (484)

TITLE INDEX

Radiation

Radiation effects and tritium technology for fusion reactors. Vol. IV* (529)

Radiation effects and tritium technology for fusion reactors* (663)

Radiative

Radiative and non-radiative transitions of rare-earth ions in glasses* (538)

radical

Structure of $\text{NO}_3^{\cdot -}$ radical in molten monovalent metal nitrates* (195)

radioactive

Investigations on radioactive fission product correlations: gamma spectrometry measurements on spent fuel assemblies discharged from the Trino Vercellese reactor at the end of the 2nd irradiation cycle* (75)

Twenty years of research in the processing of radioactive fuel and neutralization of radioactive waste at the Nuclear Research Institute (NRI)* (358)

radiochemical

8th radiochemical conference* (347)

8th radiochemical conference* (371)

Standard methods for chemical, mass spectrometric, spectrochemical, nuclear, and radiochemical analysis of nuclear-grade plutonium metal* (714)

Radiochemistry

Radiochemistry Division annual progress report for 1973* (271)

Nuclear spectroscopy. Laboratory of chemistry and radiochemistry* (723)

Radiolytic

Radiolytic gas generation in plutonium contaminated waste materials* (301)

radiotracer

Study of the Ni-Ni (II) equilibrium exchange, by radiotracer techniques in fused LiCl-KCl eutectic* (68)

Radiotracer applications in electrometallurgical processing* (76)

Raman

Electronic structure of highly symmetric complexes of lanthanoids and actinoids. Pt. 3. 'Electronic' Raman spectrum of $\text{Cs}_2\text{U}(\text{IV})\text{Cl}_6$ * (13)

Electronic structure of highly symmetric complexes of lanthanoids and actinoids. Pt. 4. The electronic Raman spectrum of $\text{Cs}_2\text{NaEu}(\text{III})\text{Cl}_6$ * (14)

Raman and i.r.-emission studies of some tungstate- and molybdate-containing melts* (102)

Study of the compounds $\text{M}_3\text{UO}_2\text{F}_5$ ($\text{M} = \text{K}, \text{Rb}, \text{Cs}, \text{NH}_4$) by i.r. absorption and Raman diffusion spectrophotometry. III. Effect of cation on vibration frequency of the ion $\text{UO}_2\text{F}_5^{3-}$ * (127)

Raman spectra of Al_2O_3 solutions in molten cryolite and other aluminum fluoride containing melts* (203)

Simple Raman cell and furnace usable at temperatures higher than 1000°C for corrosive melts* (204)

Raman spectra of solid ThCl_4 and fused $\text{ThCl}_4\text{--KCl}$ system* (488)

TITLE INDEX

Raman

Raman spectroscopic studies of yttrium (III) chloride--alkali metal chloride melts and of $\text{Cs}_2\text{NaYCl}_6$ and YCl_3 solid compounds* (489)
 Electronic structure of highly symmetric lanthanoid and actinoid compounds. VI. The electronic Raman spectrum of $\text{Cs}_2\text{NaCe(III)Cl}_6$ * (678)
 X-ray diffraction and infrared: Raman spectroscopic study of the hydrogen dinitrate ion, $(\text{O}_2\text{NO-H-ONO}_2)^+$, in cesium hydrogen dinitrate* (695)

rare

Fault structures in rare earth--cobalt intermetallics* (11)
 Mixed valence rare-earth halides and their unusual crystal structures* (29)
 Thermodynamic properties of some rare earth carbonates and yttrium carbonates* (36)
 Proceedings of the 12th rare earth research conference. Vol. II* (65)
 Temperature dependence of vapor pressure of rare earth halides and their alloys* (99)
 Investigation into reactions between trioctylammonium nitrates and rare earth nitrates by X-ray electron spectroscopy* (113)
 Synthesis and properties of rare earth titanotungstates* (114)
 Compatibility of molten rare earth metals with refractory materials* (115)
 Optical absorption Zeeman spectroscopy of rare earth orthochromites. II. Theory* (122)
 Optical absorption Zeeman spectroscopy of rare earth orthochromites. I. ErCrO_3 * (123)
 Chemical separation and spectrographic estimation of rare earths and yttrium in PuO_2 and $(\text{U,Pu})\text{O}_2$ * (125)
 Thermodynamics and kinetics of the formation of rare earth intermetallics* (137)
 Revised model for 5d-electrons in the heavy rare-earth metals* (157)
 Stability of rare earth metaniobates in hydrogen atmosphere* (164)
 Physicochemical investigation of rare earth titanoniobates, LnTiNbO_6 * (173)
 Vibration spectra of rare earth oxytungstates- Ln_2WO_6 composition* (185)
 Thermodynamics of reactions of rare earth oxides with niobium oxide* (206)
 Binary molybdates of rare earths and silver (I)* (207)
 Electron structure of rare-earth metal tetraborides* (215)
 Hydrogen solubility in rare earth intermetallic compounds* (217)
 Entropies of liquid rare earth metals* (230)
 Measurement of fluorescence of rare earths in molten salts (Nd^{3+} in NaNO_3 - KNO_3 eutectic)* (233)
 Statistical mechanics of intermediate-valence rare earth compounds* (279)
 Rare earths* (280)
 Stability of rare earth tetrafluorides* (312)
 Rare earth elements* (319)
 Interaction of atoms of rare earths-metal alloys* (320)

TITLE INDEX

rare

- Investigation of formation thermodynamic of rare earth aluminates with perovskite structure* (326)
- Corrosion of liquid alloys of zinc with yttrium and with rare earth elements in LiCl--KCl melts* (335)
- Investigation of some rare earth aluminates by the method of infrared spectroscopy* (357)
- $R\text{sub } 3T\text{sub } 2$ compounds (R = rare earth or Y; T = Rh, Pd, Pt) with the rhombohedral $\text{Er}\text{sub } 3\text{Ni}\text{sub } 2$ structure type* (365)
- Theory of rare-earth alloys* (376)
- Rare earth metal--metal halide systems. 19. Structural characterization of the reduced holmium chloride $\text{Ho}\text{sub } 5\text{Cl}\text{sub } 11$ * (378)
- Rare earth 4d spectra in rare earth trifluorides* (382)
- Pressure of saturated vapor of rare earth and yttrium bromides* (391)
- Hydrogen absorption of rare-earth (3d) transition intermetallic compounds* (392)
- Progress in the spectroscopy of rare earth and transition metal ions* (412)
- Comparative study of systems of heavier rare-earth metals with hydrogen [Gd, Tb, Dy, Lu]* (418)
- Proceedings of the 12th rare earth research conference. Vol. II* (425)
- Thermogravimetric investigation of interaction of some rare earth oxides with ammonium bifluoride* (427)
- Electronegativities of the rare-earth elements* (429)
- Structural or solid state chemistry of rare earth sesquioxides from Goldschmidt to the present time* (438)
- Determination of thorium in rare earths chloride: coprecipitation with cerium (IV) iodate and spectrophotometric determination with thoren* (447)
- Rare earth metals* (461)
- Rare earth metals* (462)
- Optical properties of rare earth trifluorides* (484)
- Rare earths* (492)
- Solubility of rare earth and yttrium oxochlorides in molten salts - $\text{MCl}\text{sub } 2$ (M-Mg, Ca, Sr, Ba)* (495)
- Investigation into absorption infrared spectra of vapors under $\text{ScBr}\text{sub } 3$, $\text{YCl}\text{sub } 3$, $\text{LaCl}\text{sub } 3$, $\text{GdCl}\text{sub } 3$, $\text{ErCl}\text{sub } 3$, $\text{LuCl}\text{sub } 3$ by means of isolation method in rare gas matrix* (497)
- Radiative and non-radiative transitions of rare-earth ions in glasses* (538)
- Rare earth metals* (546)
- Electronic and magnetic properties of rare earth- $\text{Sn}\text{sub } 3$ compounds from $\text{sup } 119\text{Sn}$ Moessbauer spectroscopy* (559)
- Spectrophotometric determination of rare earths and yttrium in gallium base alloys* (623)
- Rare earth and actinide oxides. Thermodynamic and electron microscopy studies* (625)
- Electrochemical study of high-temperature stability of compounds between the rare earths and copper oxide* (634)
- Double potassium rare earth tungstates* (639)
- Glass formation in nitrate systems containing rare earth ions* (666)

TITLE INDEX

rare

Structure of rare earth metals in the liquid state* (688)

rates

Fission rates, burnup, and neutron flux-fluence-spectra characterization for mixed oxide fuel experiments in the EBR-II* (416)

Reaction rates and electrical resistivities of the hydrogen isotopes with, and their solubilities in, liquid lithium* (529)

ray

X-ray diffraction study into molten bromides and iodides of alkali metals* (20)

Electronic structure of CeN studied by x-ray-photoemission spectroscopy* (28)

X-ray study of the plutonium--oxygen phase diagram at ambient temperatures up to 1100°C* (62)

Investigation into reactions between trioctylammonium nitrates and rare earth nitrates by X-ray electron spectroscopy* (113)

Actinide metallocarbaborane complex; synthesis and x-ray structure determination of the bis[n^{sup}5⁺-(3)-1,2-dicarbolyl]dichlorouranium(IV) dianion* (194)High temperature thermal expansion of ThO₂, MgO and Y₂O₃ by X-ray diffraction* (242)

Contribution to the establishment of the phase diagram for the U--C--C system by means of x-ray diffractometry at high temperatures and under controlled pressure* (503)

X-ray spectral study of the carbides and nitrides of group IV and V metals* (582)

X-ray diffraction and infrared: Raman spectroscopic study of the hydrogen dinitrate ion, (O₂NO-H-ONO₂)⁺, in cesium hydrogen dinitrate* (695)

Determination of trace impurities of titanium and zirconium in molybdenum and tungsten and their oxygen compounds by X-ray fluorescence spectrometry* (698)

Rb

Me₂ZrCl₆--H₂(Me=Na, K, Rb, Cs) system* (45)Study of the compounds M₃UO₂F₅ (M = K, Rb, Cs, NH₄) by i.r. absorption and Raman diffusion spectrophotometry. III. Effect of cation on vibration frequency of the ion UO₂F₅³⁻* (127)

Gases and carbon in metals (thermodynamics, kinetics, and properties). I. Alkali metals, alkaline earth metals, light metals (Li, Na, K, Rb, Cs; Ca, Sr, Ba; Be, Mg, Al)* (192)

Preparational, structural and vibrational study of M₂U₂O₅Cl₄, 2H₂O (M = Rb, Cs). Detection of a tetranuclear anion [(UO₂)₄O₂Cl₈(H₂O)₂]⁴⁻* (498)

Pressure and entropy calculations for liquid Na, K and Rb along the melting line* (584)

Degree of thermal dissociation of M₃CaCl₃ (M=K, Rb, Cs) complexes during melting* (624)

RbBr

LiBr, RbBr, CsBr ternary system* (268)

Thermodynamics of solid solutions CsCl-RbCl and CsBr-RbBr at 25°C* (570)

TITLE INDEX

RbCl

Lattice-dynamical calculation of the thermodynamic properties of NaCl, KCl and RbCl* (231)
 TlCl-RbCl-CsCl and NaI-TlI-CsI systems* (258)
 Meltability diagram of RbCl-BeCl₂ system* (506)
 Thermodynamics of solid solutions CsCl-RbCl and CsBr-RbBr at 25⁰C* (570)

RbF

RbF-NdF₃ system* (443)

reaction

Kinetics of reaction of liquid lithium with oxygen and nitrogen* (106)
 Reaction between sodium, rubidium and strontium chlorides* (286)
 Reaction of americium oxides with trivalent metal oxides* (303)
 High-temperature calorimeter for determination of reaction enthalpy* (364)
 Reaction rates and electrical resistivities of the hydrogen isotopes with, and their solubilities in, liquid lithium* (529)
 Investigation into reaction of rubidium and holmium fluorides* (540)
 Reaction of potassium superoxide in excess potassium* (590)

Reactions

Reactions of liquid sodium with transition-metal oxides. VIII. The oxides of chromium-(III), -(IV), and -(VI) and disodium chromium(VI) tetraoxide* (35)
 Electrode reactions in molten sodium nitrate--barium nitrate eutectic at 350 degrees C* (42)
 Some chemical reactions of uranium hexachloride and the chlorination of uranium tetrachloride* (81)
 Investigation into reactions between trioctylammonium nitrates and rare earth nitrates by X-ray electron spectroscopy* (113)
 Thermodynamics of reactions of rare earth oxides with niobium oxide* (206)
 Fundamental studies of metal fluorination reactions. Progress report, May 1, 1976--April 31, 1977* (533)
 Investigation into reactions of hexabromotellurates of potassium, cesium and thallium(I) in melt* (556)

Reactivity

Reactivity of transition metal fluorides. IX. Protactinium pentafluoride and tetrafluoride* (465)

reactor

Structural mechanics in reactor technology. Vol. 1* (19)
 Investigations on radioactive fission product correlations: gamma spectrometry measurements on spent fuel assemblies discharged from the Trinc Vercellese reactor at the end of the 2nd irradiation cycle* (75)
 Spectrographic determination of trace impurities in reactor grade aluminium* (104)
 Basic problems of fast reactor fuel reprocessing* (135)
 Cs--U--O phase diagram and its application to uranium--plutonium oxide fast reactor fuel pins* (175)
 Molten-salt reactor program. Semiannual progress report for period ending February 29, 1976* (176)
 Molten-salt reactor program. Semiannual progress report for period ending February 29, 1976* (237)

TITLE INDEX

reactor

Use of fluoride melts for separation of some fission products and americium from mixtures of different fluorides being the simulators of residues from fast reactor fuel fluorination* (307)
 Study of the behavior of high activity waste produced during the regeneration of fast reactor fuel elements by the gaseous fluoride method* (311)
 Chemical Engineering Division, reactor safety and physical property studies annual report, July 1974--June 1975* (367)
 MATPRO: Version 09. A handbook of materials properties for use in the analysis of light water reactor fuel rod behavior* (387)
 Energy programs: molten-salt reactor program* (423)
 Collection of gamma spectra data of irradiated light water moderated reactor spent fuel and study of the applicability of the methods for fuel identification. Final report for the period 1 February 1972--October 1975* (453)
 Extension of vapour pressure measurements of nuclear fuels (U,Pu) $O_{2\%}$ and $UO_{2\%}$ to 7000 K for fast reactor safety analysis* (475)
 Vapour pressure measurements to 7,000 K and equation of state of oxide fuels for fast reactor safety analysis. I. Extension of vapour pressure measurements of nuclear fuels (U,Pu) $O_{2\%}$ and $UO_{2\%}$ to 7,000 K* (477)
 Reactor conference, Duesseldorf, 30.3.-2.4.1976. Section 2: Fuel elements, fuel cycle and isotope technique* (480)
 Reactor conference, Duesseldorf, 30.3.-2.4.1976. Section 1: Reactor design and experiments* (509)
 Specialists' meeting on the behaviour of water reactor fuel elements under accident conditions* (510)
 Some trends in constitutive equation model development for high-temperature behavior of fast-reactor structural alloys* (527)
 Nuclear reactor containments* (569)
 Properties of structural materials for sodium-cooled fast breeder reactor* (580)
 Structural mechanics in reactor technology. Vol. 2* (601)
 Molten salt chemistry : Part IV - Solubility behaviour of $PuF_{3\%}$ in fluoride salts of interest in molten salt reactor technology* (655)
 Thermodynamic properties of solutions of hydrogen isotopes in metals and alloys of interest to fusion reactor technology* (663)

Reactors

Westinghouse Advanced Reactors Division* (57)
 Fuel processing for molten-salt reactors* (237)
 Second specialist meeting on sodium/fuel interaction in fast reactors, Ispra, 21-23 November 1973. Sponsored by the Committee on the Safety of Nuclear Installations, OECD Nuclear Energy Agency, Paris, France* (272)
 Report of task force on alternate structural materials for Liquid Metal Fast Breeder Reactors* (309)
 Preparation and thermochemical stability of uranium-zirconium-carbonitrides. A contribution to nuclear fuels for advanced high-temperature reactors* (333)
 Radiation effects and tritium technology for fusion reactors. Vol. IV* (529)

TITLE INDEX

reactors

Spectrophotometric determination of uranium in spent fuels of WWR-2 reactors* (632)
Radiation effects and tritium technology for fusion reactors* (663)

recommended

CODATA recommended key values for thermodynamics, 1976. Report of the CODATA Task Group on key values for thermodynamics, 1976* (717)

recycle

PuO₂ dissolution problem for LWR plutonium recycle and LMFBR fuels: fabrication and reprocessing problems and their resolution* (455)

redistribution

Oxygen redistribution in UO₂ due to a temperature gradient* (401)

Redox

Redox and chemical properties of U, Np and Pu in molten alkali chlorides* (360)

reduced

Calculation of the reduced partition function ratio for uranium hexafluoride* (285)
Rare earth metal-metal halide systems. 19. Structural characterization of the reduced holmium chloride Ho₅Cl₁₁* (378)

Reduction

Reduction of PuO₂ by CO and equilibrium partial pressures above plutonia* (304)
Electrochemical reduction of molybdenum oxychloride in molten lithium chloride - potassium chloride eutectic* (515)

Ref

Interactions in inorganic molecular crystals - electronic spectra of ReF₆ pure and mixed crystals. Technical report, 1 Jul--31 Dec 1975* (53)

reflectance

Diffuse reflectance spectra of protactinium(IV), uranium(IV) and neptunium(IV) tetrachlorides and tetraformates* (562)

refractories

Wetting of certain refractories of the system MgO-Al₂O₃-ZrO₂ with metallic melts* (344)

refractory

Spectral emissivities of molten refractory metals* (47)
Compatibility of molten rare earth metals with refractory materials* (115)
Investigation on the resistance of high-refractory materials to chromium melt* (531)
Refractory materials resistance to molten barium alloy* (628)

regeneration

Study of the behavior of high activity waste produced during the regeneration of fast reactor fuel elements by the gaseous fluoride method* (311)

region

Equation of state and transport properties of uranium and plutonium carbides in the liquid region* (577)
Equation of state and transport properties of uranium and plutonium nitrides in the liquid region* (578)

TITLE INDEX

regions

Practice of experiment planning in studying limited regions of multicomponent system* (546)

regularities

Some regularities of the corrosion behavior of binary magnesium alloys* (439)

Rehovot

Israel Nuclear Society symposium on "'Nuclear fuel in Israel'", Weizmann Institute of Science, Rehovot, 1.7.75* (226)

relaxation

Ionic-thermoelectric study of the dipole relaxation and equilibrium in Gd-doped SrF_2 * (410)

Nuclear quadrupole relaxation and viscosity in liquid metals* (564)

rendus

Comptes rendus du 95* (621)

REOsub

Phase equilibria and thermodynamic data for the ternary UO_2 - UO_2 - REO (1.5) (Ho, Tm, Lu) systems* (694)

reovering

Method for recovering uranium from residual solution* (70)

representatives

New representatives of CeMn_4Al_8 structural type* (549)

Reprocessing

Reprocessing of fast breeder fuels in France* (67)

LMFBR Fuel Reprocessing Program progress report for period April 1 to June 30, 1976* (94)

Accurate procedure to safeguard the fissile material content of input and output solutions of reprocessing plants* (131)

Basic problems of fast reactor fuel reprocessing* (135)

Chemical reprocessing of spent nuclear fuels* (322)

PuO_2 dissolution problem for LWR plutonium recycle and LMFBR fuels: fabrication and reprocessing problems and their resolution* (455)

research

Proceedings of the 12th rare earth research conference. Vol. I* (11)

Proceedings of the 12th rare earth research conference. Vol. II* (65)

Molten carbonate fuel cell research at ORNL* (71)

Results of research to evaluate solid plutonium nitrate as a safe shipping form* (85)

Proceedings of the symposium and workshop on advanced battery research and design, March 22--24, 1976* (269)

Twenty years of research in the processing of radioactive fuel and neutralization of radioactive waste at the Nuclear Research Institute (NRI)* (358)

Physical research on liquid-metal systems. Annual report, July 1975--June 1976* (405)

Proceedings of the 12th rare earth research conference. Vol. II* (425)

Research laboratories annual report 1975* (566)

Research on a high-energy nonaqueous battery system. Quarterly progress report No. 1, 1 Dec 1965--1 Mar 1966* (720)

TITLE INDEX

Research

Research on electrochemical parameters for nonaqueous batteries: task I. Quarterly progress report No. 3* (727)
 Basic research in crystalline and noncrystalline ceramic systems. Annual report, May 1, 1975--April 1, 1976* (728)

residual

Method for recovering uranium from residual solution* (70)

residues

Use of fluoride melts for separation of some fission products and americium from mixtures of different fluorides being the simulators of residues from fast reactor fuel fluorination* (307)
 Investigation into americium extraction process from residues of irradiated nuclear fuel after its fluorination by SiF-NaF and SiF-CaF_2 melts of eutectic composition* (308)

resistance

Investigation on the resistance of high-refractory materials to chromium melt* (531)
 Basic mechanisms providing oxidation resistance in structural metals at high temperatures. II. The defect structure of nonstoichiometric rutile (TiO_2) and nonstoichiometric rutile containing 1 to 10 mole % Cb_2O_6 . Final report 21 Dec 1973-20 Dec 1974* (542)
 Refractory materials resistance to molten barium alloy* (628)

resisting

Ceramic-base heat resisting constructional material* (348)

resistivities

Reaction rates and electrical resistivities of the hydrogen isotopes with, and their solubilities in, liquid lithium* (529)

resistivity

Negative temperature coefficients of electrical resistivity: the divalent liquid metals Eu, Yb and Ba* (224)
 Aspects of the solution chemistry of liquid alkali metals as elucidated from electrical resistivity studies* (247)
 Improved model for solutions of nonmetals in liquid alkali metals. Calculation of enthalpy of solution and electrical resistivity* (486)

resolidification

Computational simulation of cladding motion after melting including resolidification processes* (19)

resolution

PuO_2 dissolution problem for LWR plutonium recycle and LMFBR fuels: fabrication and reprocessing problems and their resolution* (455)

resonance

Nuclear quadrupole resonance spectra of HgCl_2 under pressure* (31)

retention

Study of the retention of Ba in UO_2 nuclear fuel particles by ZrO_2 as getter* (470)

Rev

At. Energy Rev.* (7)
 At. Energy Rev.* (464)

Review

Review of partitioning proposals for spent nuclear fuels* (69)

TITLE INDEX

Rh

$R\text{sub } 3T\text{sub } 2$ compounds (R = rare earth or Y; T = Rh, Pd, Pt) with the rhombohedral $Er\text{sub } 3Ni\text{sub } 2$ structure type* (365)

rhenium

Some constitutional studies on uranium-carbon- and plutonium-carbon-rhenium and technetium systems* (227)
Some constitutional studies on uranium--carbon and plutonium--carbon--rhenium and technetium systems* (228)

rhodium

Structure of the thorium-(zirconium, niobium, ruthenium, rhodium)-carbon systems* (245)

rhombohedral

$R\text{sub } 3T\text{sub } 2$ compounds (R = rare earth or Y; T = Rh, Pd, Pt) with the rhombohedral $Er\text{sub } 3Ni\text{sub } 2$ structure type* (365)

riched

Thermodynamic properties of the U-Sn-alloys riched with tin* (284)

Rocket

Space Processing Applications Rocket project, SPAR 1. Final report* (536)

rod

MATERO: Version 09. A handbook of materials properties for use in the analysis of light water reactor fuel rod behavior* (387)

rods

Phase analysis on simulated LWR fuel rods and its evaluation with the aid of constitution studies in the system uranium-zirconium-oxygen* (509)
Phase analysis at simulated LWR fuel rods and their interpretation by constitution investigations in the uranium-zirconium-oxygen system* (510)

rubidium

Binary system of sodium, rubidium and cesium tungstates* (151)
Thermochemistry of polyhalides. Pt.1. Caesium and rubidium dichloroiodates(I)* (178)
Ternary mutual system of chlorides and iodides of rubidium and cadmium* (257)
Reaction between sodium, rubidium and strontium chlorides* (286)
Phase diagram of the ternary system of lithium-, rubidium-, and cesium nitrates* (377)
Interaction of rubidium and cesium fluorides with lutetium trifluorides* (444)
Some structural characteristics of rubidium and cesium in the vicinity of their melting points* (460)
Investigation into reaction of rubidium and holmium fluorides* (540)
Quarter mutual system of chlorides, bromides, iodides of sodium and rubidium* (629)
Quaternary mutual system of chlorides, bromides, iodides of rubidium and cesium* (640)

rules

Structure of the biexciton molecule and optical transition selection rules in zincblende and wurtzite materials* (186)

RuC

Behavior of ruthenium in fluoride-volatility process. II. Fluorination of $RuC\text{sub } 2$ by $F\text{sub } 2$ * (558)

TITLE INDEX

ruthenium

Structure of the thorium-(zirconium, niobium, ruthenium, rhodium)-carbon systems* (245)
 Behavior of ruthenium in fluoride-volatility process. II. Fluorination of RuO_2 by F_2 * (558)

rutile

Basic mechanisms providing oxidation resistance in structural metals at high temperatures. II. The defect structure of nonstoichiometric rutile (TiO_2) and nonstoichiometric rutile containing 1 to 10 mole % Nb_2O_5 . Final report 21 Dec 1973-20 Dec 1974* (542)

S

Preparation and some properties of actinide(III) and lanthanide(III) oxide halides, and the enthalpies of formation of $\text{PuBr}_3(s)$ and $\text{PuI}_3(s)$ * (82)
 On interaction in U--S--Te system* (162)
 U.S.-Japan joint seminar on defects and diffusion in solids* (611)
 Summary of U.S. LMFBR programs on high temperature structural design and associated materials testing* (731)

safe

Results of research to evaluate solid plutonium nitrate as a safe shipping form* (85)

safeguard

Accurate procedure to safeguard the fissile material content of input and output solutions of reprocessing plants* (131)

Safeguarding

Safeguarding nuclear materials. Vol. 2* (131)
 Safeguarding nuclear materials. Vol. 2* (349)

safety

Investigation of fuel vapor pressure for fast breeder safety analysis* (60)
 Second specialist meeting on sodium/fuel interaction in fast reactors, Ispra, 21-23 November 1973. Sponsored by the Committee on the Safety of Nuclear Installations, OECD Nuclear Energy Agency, Paris, France* (272)
 EUTECTIC DATA: safety, hazards, corrosion, melting points, compositions, and bibliography* (276)
 Chemical Engineering Division, reactor safety and physical property studies annual report, July 1974--June 1975* (367)
 Extension of vapour pressure measurements of nuclear fuels (U,PuO_2 and UO_2) to 7000 K for fast reactor safety analysis* (475)
 Vapour pressure measurements to 7,000 K and equation of state of oxide fuels for fast reactor safety analysis. I. Extension of vapour pressure measurements of nuclear fuels (U,PuO_2 and UO_2) to 7,000 K* (477)

salt

Study of saturated vapor pressure above salt melts in CoCl_2 - MCl systems* (98)
 Study on conductivity, density, and viscosity of molten salt systems* (112)
 Molten-salt reactor program. Semiannual progress report for period ending February 29, 1976* (176)

TITLE INDEX

salt

- Composition gradient during electrolysis in mixtures analogous to molten salt battery electrolytes* (234)
- Molten-salt reactor program. Semiannual progress report for period ending February 29, 1976* (237)
- Structure of lithium--boron alloys and their performance as potential anodes for high power molten salt batteries. Progress report, 1 Aug 1974--30 Jun 1975* (275)
- Molten-salt electrolytes--transport properties* (277)
- Influence of intermediate products on chlorination kinetics of vanadium oxides in salt melt* (356)
- Energy programs: molten-salt reactor program* (423)
- Influence of salt background on kinetics of BF_4^-sup alkali hydrolysis* (504)
- Application of molecular dynamics computations to the conformal ionic solution theory: Thermodynamics of binary molten salt mixtures* (550)
- Molten salt chemistry : Part IV - Solubility behaviour of PuF_3 in fluoride salts of interest in molten salt reactor technology* (655)
- Investigation of the solubility and diffusion of Fe atoms in Cu at high temperature using molten salt electrochemistry* (693)
- Preparation of thorium metal by electrolysis of fused salt $\text{ThCl}_4\text{-LiCl+NaCl+KCl(eut)}$ system* (703)
- Lithium alloys--chlorine secondary battery. I. Basic studies on a lithium alloys--chlorine secondary battery using a molten salt as the electrolyte* (704)
- Investigation into properties of liquidus curves of binary salt system melts where chemical compounds formation. 2. Systems of monc-moncvalent and bi-bivalent cations with common anion* (708)
- Molten-salt systems* (729)

salts

- Spectral properties of impurity molecules of the C_{3v} symmetry groups in solid solutions of halide salts* (44)
- High-temperature unit to the IKS-21 spectrometer for the detection of molten salts absorption spectra* (208)
- Measurement of fluorescence of rare earths in molten salts (Nd^{3+} in $\text{NaNO}_3\text{-KNO}_3$ eutectic)* (233)
- Electrochemistry of molten salts* (263)
- Use of the SF-8 detection spectrophotometer for measuring absorption spectra of molten salts* (331)
- Investigation of solubility and thermodynamics of dissolution of salts in mixed solvents* (347)
- Investigations on the solvent extraction from molten salts by tracer technique* (369)
- Investigations of the solvent extraction from molten salts with the use of tracer technique* (371)
- Properties of yttrium and scandium basic salts* (393)
- Dispersion of metallic lithium in various molten salts* (452)
- Solubility of rare earth and yttrium oxochlorides in molten salts - MCl_2 (M-Mg, Ca, Sr, Ba)* (495)
- Practical application of conversion method in study of five-component mutual system consisting of nine salts NaF , Na_2MoO_4 , Na_2WO_4 , KF , K_2MoO_4 , K_2WO_4 , BaF_2 , BaMoO_4 , BaWO_4 * (516)

TITLE INDEX

salts

Recent developments in the chemistry of solutions of salts in liquid alkali metals* (528)
 Spectrochemical method for determination of zirconium in molten salts* (566)
 Heat transfer measurements in a forced convection loop with two molten-fluoride salts: $\text{LiF--BeF}_2\text{--ThF}_2\text{--UF}_4$ and eutectic $\text{NaBF}_4\text{--NaF}$ * (585)
 Molten salt chemistry : Part IV - Solubility behaviour of PuF_3 in fluoride salts of interest in molten salt reactor technology* (655)
 Isorhodate complexes of 3d-metal ions in molten salts* (672)
 Effect of second sphere cations on formation of nickel (2) chloride complexes in molten salts* (673)
 Spectra of combination scattering and structure of 3d-metals chloride complexes in molten salts* (674)
 Electrochemical methods for monitoring purity of molten salts. Semiannual progress report, July--December 1975* (721)

samarium

Kinetics and mechanism into preparation of samarium-, europium- and ytterbium dichlorides and their some properties* (253)
 Formation enthalpy of di- and trifluorides of samarium* (306)

satellite

Zeeman effect and satellite carrier structure in absorption spectra of pure and doped praseodymium and europium double nitrates* (165)

saturated

Tensimetric study of pressure and composition of saturated vapor in TlCl--InCl_3 system* (93)
 Study of saturated vapor pressure above salt melts in $\text{CoCl}_2\text{--MCl}$ systems* (98)
 Properties of zirconium and hafnium tetrachlorides and their saturated solutions* (136)
 Thermodynamic state, specific heat, and enthalpy function of saturated UO_2 vapor between 3000 K and 5000 K* (297)
 Pressure of saturated vapor of rare earth and yttrium bromides* (391)

Saturation

Saturation vapor pressure in iron chloride-alkaline earth metal chlorides systems* (97)

Sb

Structure of molten Bi--Sb alloys by means of neutron diffraction* (359)

Sc

Vibrational spectra and force constants of the elpasolites Cs_2KMgF_6 ($\text{M}=\text{Sc}, \text{Y}, \text{La}, \text{Gd}, \text{Yb}$)* (41)
 Electromodulation spectroscopy of sc and fcc phase TlCl and TlBr * (411)

scale

Technical experiences in an engineering-scale facility for uranium fluorination studies* (699)
 Description of an engineering-scale facility for uranium fluorination studies* (700)

scandium

Properties of yttrium and scandium basic salts* (393)

TITLE INDEX

scattering

Studies of the electronic structure of simple metals and their alloys with the aid of Compton scattering and positron annihilation using lithium, magnesium, lithium-magnesium and aluminium-zinc as examples* (48)
 Anomalous quasielastic light scattering from metal oxide glasses* (183)
 Frequency spectrum and scattering law for liquid copper. A molecular dynamics investigation* (567)
 Spectra of combination scattering and structure of 3d-metals chloride complexes in molten salts* (674)

ScBr

Investigation into absorption infrared spectra of vapors under ScBr_3 , YCl_3 , LaCl_3 , GdCl_3 , ErCl_3 , LuCl_3 by means of isolation method in rare gas matrix* (497)

Schottky

Thermodynamics of the lanthanide halides. II. Heat capacities and Schottky anomalies of SmCl_3 , EuCl_3 and GdCl_3 from 50 to 350 K* (594)

Science

Israel Nuclear Society symposium on 'Nuclear fuel in Israel', Weizmann Institute of Science, Rehovot, 1.7.75* (226)
 Advances in nuclear science and technology. Vol. 9* (448)
 Recent development of ceramic basic science in Japan* (557)

ScS

Mass spectrometric determination of dissociation energies of high-temperature molecules: ScS , YS , LaS , ZrS , UO * (597)

secondary

Lithium--silver chloride secondary battery investigation. Quarterly technical progress report No. 2* (111)
 Lithium--silver chloride secondary battery investigation. Final report, December 1963--November 1964* (118)
 Electrochemical characterization of nonaqueous systems for secondary battery application. Quarterly report, February--April 1968* (380)
 Lithium--nickel fluoride secondary battery investigation. Quarterly technical progress report No. 1, 27 Apr--27 Jul 1966* (381)
 Effects of secondary ligands on the electronic structure of uranyls* (684)
 Lithium alloys--chlorine secondary battery. I. Basic studies on a lithium alloys--chlorine secondary battery using a molten salt as the electrolyte* (704)

Segregation

Segregation and vapour pressure studies on the uranium-plutonium-carbon system* (83)
 Segregation and vapour pressure studies in the uranium--plutonium--carbon system* (84)
 Chemical effects of composition changes in irradiated oxide fuel materials. II. Fission product segregation and chemical equilibria* (169)

selection

Structure of the biexciton molecule and optical transition selection rules in zincblende and wurtzite materials* (186)

TITLE INDEX

Self

Self-diffusion in liquid metals* (198)

Semiannual

Molten-salt reactor program. Semiannual progress report for period ending February 29, 1976* (176)

Molten-salt reactor program. Semiannual progress report for period ending February 29, 1976* (237)

Semiannual report 1975/2* (454)

Electrochemical methods for monitoring purity of molten salts.

Semiannual progress report, July--December 1975* (721)

semiconductors

Physics of semiconductors* (186)

seminar

Eleventh annual seminar on theoretical physics, Johannesburg, 5-9 July 1976* (368)

U.S.-Japan joint seminar on defects and diffusion in solids* (611)

sensitivity

Environmental sensitivity of structural metals: liquid metal embrittlement. Annual technical progress report No. 2, 25 Jun 1970--31 May 1971* (95)

Environmental sensitivity of structural metals: liquid metal embrittlement. Annual technical progress report No. 4, 1 Jun 1972--31 May 1973* (211)

Environmental sensitivity of structural metals: liquid metal embrittlement. Annual technical progress report No. 2, 25 Jun 1970--31 May 1971* (212)

separation

Chemical separation and spectrographic estimation of rare earths and yttrium in PuO_2 and $(\text{U,Pu})\text{O}_2$ * (125)

Use of fluoride melts for separation of some fission products and americium from mixtures of different fluorides being the simulacra of residues from fast reactor fuel fluorination* (307)

Current yield during separation of lithium on a liquid bismuth cathode* (430)

Separation of alkaline earth from alkali metal fission products in the SCLAR on-line mass spectrometer* (534)

September

Analytical methods for fissionable materials in the nuclear fuel cycle. Progress report, July 1, 1975--September 30, 1976* (690)

series

Structural changes in the series LnFeO_3 and LnF_3 * (197)

sesquinitride

Nonstoichiometry in cubic uranium sesquinitride* (611)

sesquioxide

Heat of formation of europium sesquioxide and europium trichloride* (607)

sesquioxides

Structural or solid state chemistry of rare earth sesquioxides from Goldschmidt to the present time* (438)

SF

Use of the SF-8 detection spectrophotometer for measuring absorption spectra of molten salts* (331)

TITLE INDEX

shells

Partly filled shells constituting anti-bonding orbitals with higher ionization energy than their bonding counterparts* (280)

shift

Pulse NMR in solids: chemical shift, lead fluoride, and thorium hydride* (363)

shipping

Results of research to evaluate solid plutonium nitrate as a safe shipping form* (85)

SiF

Investigation into americium extraction process from residues of irradiated nuclear fuel after its fluorination by SiF-NaF and SiF-CaF₂ melts of eutectic composition* (308)

silica

High density thoria-silica-metal (III) oxide fibers* (710)

silver

Lithium--silver chloride secondary battery investigation. Quarterly technical progress report No. 2* (111)

Lithium--silver chloride secondary battery investigation. Final report, December 1963--November 1964* (118)

Binary molybdates of rare earths and silver (I)* (207)

similarity

Thermodynamic similarity and universal equations of state of hexafluorides* (397)

simulated

Phase analysis on simulated LWR fuel rods and its evaluation with the aid of constitution studies in the system uranium-zirconium-oxygen* (509)

Phase analysis at simulated LWR fuel rods and their interpretation by constitution investigations in the uranium-zirconium-oxygen system* (510)

simulation

Computational simulation of cladding motion after melting including resolidification processes* (19)

simulators

Use of fluoride melts for separation of some fission products and americium from mixtures of different fluorides being the simulators of residues from fast reactor fuel fluorination* (307)

single

Theory for the derivation of a single hard-sphere potential describing the overall molecular interactions in binary liquid metal alloys as a function of temperature and composition* (525)

Sm

Density, viscosity, electric conductivity and meltability of Na₃AlF₆-La₂O₃ and Na₃AlF₆-6f-Sm₂O₃ systems* (43)

Energy transfer between Bi³⁺ yields Eu³⁺, Bi³⁺ yields Sm³⁺ and UO₂²⁺ yields Eu³⁺ in oxide glasses* (539)

SmCl

Thermodynamics of the lanthanide halides. II. Heat capacities and Schottky anomalies of SmCl₃, EuCl₃ and GdCl₃ from 50 to 350 K* (594)

TITLE INDEX

Sn

- Thermodynamic properties of the U-Sn-alloys riched with tin* (284)
 Speed of ultrascund and the thermophysical properties of the liquid metals Sn, Pb, Cd and of their binary alloys Pb:Sn and Pb:Cd* (403)
 Interaction parameters in the Zn--Pb--Sn system at low zinc concentration* (436)
 Electronic and magnetic properties of rare earth-Sn₃ compounds from ¹¹⁹Sn Moessbauer spectroscopy* (559)
 Thermodynamic properties of the ternary Sn-Pb-Cd liquid metal system* (606)

Society

- Israel Nuclear Society symposium on ''Nuclear fuel in Israel'', Weizmann Institute of Science, Rehovot, 1.7.75* (226)

sodium

- Interaction of uranyl chloride with zirconium tetrachloride in the melt of sodium chloride and potassium chloride* (27)
 Standard potential of sodium amalgam at 25⁰C* (32)
 Reactions of liquid sodium with transition-metal oxides. VIII. The oxides of chromium-(III), -(IV), and -(VI) and disodium chromium(VI) tetraoxide* (35)
 Electrode reactions in molten sodium nitrate--barium nitrate eutectic at 350 degrees C* (42)
 Experimental vapor pressure and critical point of sodium* (56)
 Ternary systems from lithium, sodium, strontium and barium chlorides* (86)
 Phase equilibria in the system of sodium chlorides, cesium chlorides and strontium chlorides* (87)
 Phase equilibria in the system of sodium chlorides, cesium chlorides and strontium chlorides* (88)
 Quaternary mutual system from fluorides and chlorides of lithium, sodium and strontium* (89)
 Experimental investigation of cavitation inception and damage in a flowing sodium environment* (103)
 Physical models of mass transport of iron and nickel in liquid sodium systems* (128)
 Thermographic investigation into ternary systems containing sodium and potassium chlorides, uranium trichloride and thorium tetrafluoride* (140)
 Density, surface tension, and viscosity of uranium trichloride-sodium chloride melting* (141)
 Interaction of uranium- and thorium tetrachlorides with sodium- and potassium chlorides* (144)
 Phase diagrams of ternary systems containing sodium and potassium chlorides, thorium tetrachloride and plutonium trichloride* (148)
 Binary system of sodium, rubidium and cesium tungstates* (151)
 Solubility of sodium iodide in liquid sodium between 140 and 500⁰C* (154)
 Atomic diffusion in alloys exposed to liquid sodium* (181)
 Electrochemical studies of the sodium--bismuth system* (187)
 Structure factor of liquid caesium and sodium-caesium alloys* (249)

TITLE INDEX

sodium

Second specialist meeting on sodium/fuel interaction in fast reactors, Ispra, 21-23 November 1973. Sponsored by the Committee on the Safety of Nuclear Installations, OECD Nuclear Energy Agency, Paris, France* (272)

Reaction between sodium, rubidium and strontium chlorides* (286)

Study of mass transfer by liquid sodium in closed loop systems. Sodium technology development programme* (324)

Sodium uranium(V) trioxide, NaUO_3 : heat capacity and thermodynamic properties from 5 to 350 K* (383)

Sodium uranium(V) trioxide, NaUO_3 : heat capacity and thermodynamic properties from 5 to 350 K* (384)

Thermodynamic properties of diluted sodium solutions in liquid indium* (388)

Properties of structural materials for sodium-cooled fast breeder reactor* (580)

Thermodynamics of electrolytes. 8. High-temperature properties, including enthalpy and heat capacity, with application to sodium chloride* (586)

Quarter mutual system of chlorides, bromides, iodides of sodium and rubidium* (629)

Sections of quaternary mutual system consisting of sodium, potassium and cesium chlorides and bromides* (631)

SOLAR

Separation of alkaline earth from alkali metal fission products in the SOLAR on-line mass spectrometer* (534)

solid

Variation of the solid--solid--nickel--thoria interfacial free energy with temperature* (5)

Spectral properties of impurity molecules of the $\text{Cs}_{3\text{v}}$ symmetry groups in solid solutions of halide salts* (44)

Results of research to evaluate solid plutonium nitrate as a safe shipping form* (85)

Nonisothermal flow of liquid metals over the surface of solid ones* (101)

Summaries of reports of the all-union meeting on solid body chemistry. 3--5 June 1975. III. Synthesis, phase equilibrium and use of solid oxide compounds* (114)

Electronic conductivity and polarization effects of solid ThO_2 - Y_2O_3 electrolytes and of oxygen containing vanadium-metal electrodes* (193)

Thermodynamic properties of Cr, Mn, Co and Ni dichlorides from emf measurements on cells with solid electrolytes* (200)

Criteria for solid solution formation in NFE alloys* (205)

Thermodynamic properties of solid and liquid metals and ceramics* (243)

Adhesion properties of melted Al, Ga, In on solid titanium* (327)

Origin of excess entropy of alkali metals halides solid solutions* (390)

Thermal decomposition of solid metal oxides* (395)

Heat of formation of solid and liquid alloys of transition metals* (425)

Structural or solid state chemistry of rare earth sesquioxides from Goldschmidt to the present time* (438)

Raman spectra of solid ThCl_4 and fused ThCl_4 -KCl system* (488)

TITLE INDEX

solid

- Raman spectroscopic studies of yttrium (III) chloride--alkali metal chloride melts and of $\text{Cs}_{2\text{NaYCl}_6}$ and YCl_3 solid compounds* (489)
- Thermodynamic studies of gallium--indium liquid alloys by solid state electrochemistry with oxide electrolytes* (513)
- Measurement of the cell voltage of ThO_2 -- Y_2O_3 solid electrolyte cells in the range of mixed conduction* (535)
- Heats of formation of solid solutions of CsCl with CsBr * (543)
- Thermodynamics of solid solutions CsCl - RbCl and CsBr - RbBr at 25°C * (570)
- Decomposition of the UN-CeN solid solution* (581)
- Terminal solid solubility of hydrogen in the zirconium-2.5 weight % niobium alloy* (587)
- Zirconia and yttria--thoria ceramics find new uses as solid electrolytes* (595)
- Summaries of reports of the all-union meeting on solid body chemistry. 3--5 June 1975. III. Synthesis, phase equilibrium and use of solid oxide compounds* (641)
- Investigations on chemical properties of thorium oxide solid electrolyte ceramics* (652)
- Solubility of nitrogen in liquid lithium and thermal decomposition of solid Li_3N * (702)

solids

- Structure and excitations of amorphous solids* (183)
- Pulse NMR in solids: chemical shift, lead fluoride, and thorium hydride* (363)
- U.S.-Japan joint seminar on defects and diffusion in solids* (611)

solubilities

- Prediction of the inert gas solubilities in stoichiometric molten UO_2 * (225)
- Reaction rates and electrical resistivities of the hydrogen isotopes with, and their solubilities in, liquid lithium* (529)

solubility

- Hydrogen solubility in liquid cesium* (22)
- Solubility of nitrogen in multicomponent nickel base alloys* (100)
- Solubility of sodium iodide in liquid sodium between 140 and 500°C * (154)
- Solubility and activity of oxygen in liquid indium and copper-indium alloys* (182)
- Hydrogen solubility in rare earth intermetallic compounds* (217)
- Titanium dioxide solubility in melts of lithium- and alkaline earth metal chlorides* (264)
- Solubility of magnesium oxide in alkali metal chloride melt* (265)
- Hydrogen solubility in liquid zirconium* (287)
- Solubility of oxygen in liquid niobium* (342)
- Investigation of solubility and thermodynamics of dissolution of salts in mixed solvents* (347)
- Solubility of $\text{Sr}(\text{NO}_3)_2$ in the LiNO_3 - KNO_3 melt of variable composition* (351)
- Solubility of magnesium oxide in calcium oxide--calcium chloride mixtures* (352)

TITLE INDEX

solubility

- Investigation of calcite solubility in K_2CO_3 and Na_2CO_3 solutions at temperatures of 200 to 350°C* (396)
- Misch metal solubility in melt of LiF-BaF₂ system* (462)
- Solubility of rare earth and yttrium oxochlorides in molten salts - MCl_2 (M-Mg, Ca, Sr, Ba)* (495)
- Nitrogen solubility in liquid nickel--niobium alloys* (532)
- K_2HF_6 -- K_2ZrF_6 --1.25 percent of HF system at 40°C with other solubility curves from 25°C to 70°C* (541)
- Solubility of oxygen in molten carbonates* (563)
- Terminal solid solubility of hydrogen in the zirconium-2.5 weight % niobium alloy* (587)
- Gd, Tb, Dy, Ho, Er solubility in chromium* (610)
- Molten salt chemistry : Part IV - Solubility behaviour of PuF_3 in fluoride salts of interest in molten salt reactor technology* (655)
- Investigation of the solubility and diffusion of Fe atoms in Cu at high temperature using molten salt electrochemistry* (693)
- Solubility of lithium oxide in liquid lithium* (701)
- Solubility of nitrogen in liquid lithium and thermal decomposition of solid Li_3N * (702)

solution

- Method for recovering uranium from residual solution* (70)
- High-temperature microcalorimetric measurements of partial molar enthalpy of solution, $\Delta H(O_2)$, in plutonium + oxygen* (107)
- Criteria for solid solution formation in NFE alloys* (205)
- Aspects of the solution chemistry of liquid alkali metals as elucidated from electrical resistivity studies* (247)
- Solution studies in the $US-UC_2$ system* (379)
- Quantitative study of the chemical equilibria in solution in molten LiCl-KCl (70-30 mol.%) by absorption spectrophotometry. Application to neptunium* (385)
- Solution behavior of hydrogen isotopes and other non-metallic elements in liquid lithium* (404)
- Cesium nitrate. Standard enthalpies of solution and formation and the standard entropy at 298.15 K* (466)
- Lithium nitride (Li_3N): standard enthalpy of formation by solution calorimetry* (467)
- Improved model for solutions of nonmetals in liquid alkali metals. Calculation of enthalpy of solution and electrical resistivity* (486)
- Application of molecular dynamics computations to the conformational ionic solution theory: Thermodynamics of binary molten salt mixtures* (550)
- Decomposition of the UN-CeN solid solution* (581)

solutions

- Thermodynamic properties of diluted solutions of $MnCl_2$ in alkali metal chlorides* (6)
- Spectral properties of impurity molecules of the C_{3v} symmetry groups in solid solutions of halide salts* (44)
- Composition of nonaqueous solutions of potential use in high energy density batteries. Final report, 15 Nov 1969--14 Nov 1972* (119)

TITLE INDEX

solutions

- Accurate procedure to safeguard the fissile material content of input and output solutions of reprocessing plants* (131)
- Properties of zirconium and hafnium tetrachlorides and their saturated solutions* (136)
- Raman spectra of Al_2O_3 solutions in molten cryolite and other aluminum fluoride containing melts* (203)
- Thermodynamic properties of diluted sodium solutions in liquid indium* (388)
- Origin of excess entropy of alkali metals halides solid solutions* (390)
- Investigation of calcite solubility in K_2CO_3 and Na_2CO_3 solutions at temperatures of 200 to 350°C* (396)
- Some thermodynamic properties of dilute solutions of actinide chlorides in (Li-K)Cl and in (Na-K)Cl eutectics* (406)
- Thermodynamic properties of dilute solutions of ThCl_4 in (Li-K)Cl and (Na-K) eutectics* (407)
- Improved model for solutions of nonmetals in liquid alkali metals. Calculation of enthalpy of solution and electrical resistivity* (486)
- Investigation of liquid-steam equilibrium in system with diluted solutions of metal fluorides in uranium hexafluoride* (526)
- Recent developments in the chemistry of solutions of salts in liquid alkali metals* (528)
- Heats of formation of solid solutions of CsCl with CsBr* (543)
- Thermodynamics of solid solutions CsCl-RbCl and CsBr-RbBr at 25°C* (570)
- Spectrophotometric method for the determination of neptunium and plutonium in process solutions* (660)
- Thermodynamic properties of solutions of hydrogen isotopes in metals and alloys of interest to fusion reactor technology* (663)

solvent

- Investigations on the solvent extraction from molten salts by tracer technique* (369)
- Studies on some binary solvent systems for the extraction of gadolinium* (370)
- Investigations of the solvent extraction from molten salts with the use of tracer technique* (371)

solvents

- Investigation of solubility and thermodynamics of dissolution of salts in mixed solvents* (347)

sorption

- Tritium sorption in lithium--bismuth and lithium--aluminum alloys* (614)

Space

- Space Processing Applications Rocket project, SPAR 1. Final report* (536)

SPAR

- Space Processing Applications Rocket project, SPAR 1. Final report* (536)

specialist

- Second specialist meeting on sodium/fuel interaction in fast reactors, Ispra, 21-23 November 1973. Sponsored by the Committee on the Safety of Nuclear Installations, OECD Nuclear Energy Agency, Paris, France* (272)

TITLE INDEX

Specialists

Specialists' meeting on the behaviour of water reactor fuel elements under accident conditions* (510)

species

Optical spectra of the difluoride, dichloride, and trichloride ions in the matrix-isolated $M^{sup} + F^{sub 2}^{sup} -$, $M^{sup} + Cl^{sub 2}^{sup} -$, and $M^{sup} + Cl^{sub 3}^{sup} -$ species* (16)
 Optical spectra of the dibromide and diiodide ions in the matrix-isolated $M^{sup} + Br^{sub 2}^{sup} -$ and $M^{sup} + I^{sub 2}^{sup} -$ species* (17)
 Diffusion of uranium species in several molten chlorides* (408)
 Total pressures of uranium- and plutonium-bearing species above the U--Pu--O system* (618)

spectra

Optical spectra of the difluoride, dichloride, and trichloride ions in the matrix-isolated $M^{sup} + F^{sub 2}^{sup} -$, $M^{sup} + Cl^{sub 2}^{sup} -$, and $M^{sup} + Cl^{sub 3}^{sup} -$ species* (16)
 Optical spectra of the dibromide and diiodide ions in the matrix-isolated $M^{sup} + Br^{sub 2}^{sup} -$ and $M^{sup} + I^{sub 2}^{sup} -$ species* (17)
 Nuclear quadrupole resonance spectra of $HgCl^{sub 2}$ under pressure* (31)
 Absorption spectra of curium (3) chloride complexes* (34)
 Vibrational spectra and force constants of the elpasolites $Cs^{sub 2}KMP^{sub 6}$ ($M=Sc, Y, La, Gd, Yb$)* (41)
 Temperature effect on vibrational spectra of the impurity molecular ions in alkali halides* (46)
 Vibrational spectra of transition metal hexafluoride crystals. I. Orthorhombic MoF_6 , WF_6 , UF_6 neat crystals. Technical report* (51)
 Vibrational spectra of transition metal hexafluoride crystals. III. Exciton band structures of $MoF^{sub 6}$, $WF^{sub 6}$, and $UF^{sub 6}$. Technical report* (52)
 Interactions in inorganic molecular crystals - electronic spectra of $ReF^{sub 6}$ pure and mixed crystals. Technical report, 1 Jul--31 Dec 1975* (53)
 Zeeman effect and satellite carrier structure in absorption spectra of pure and doped praseodymium and europium double nitrates* (165)
 Vibration spectra of rare earth oxytungstates- $Ln^{sub 2}WO^{sub 6}$ composition* (185)
 Raman spectra of $Al^{sub 2}O^{sub 3}$ solutions in molten cryolite and other aluminum fluoride containing melts* (203)
 High-temperature unit to the IKS-21 spectrometer for the detection of molten salts absorption spectra* (208)
 Mass spectra and vapor composition of niobium pentafluoride* (216)
 Fine definition of ir spectra from high temperature interactions of U + $O^{sub 2}$. Part II. Final report, 1 July 1974--30 June 1975* (218)
 Molecular vapor composition and mass spectra in $KF-ZrF^{sub 4}$, $CsF-ZrF^{sub 4}$ systems* (294)
 Use of the SP-8 detection spectrophotometer for measuring absorption spectra of molten salts* (331)
 Rare earth 4d spectra in rare earth trifluorides* (382)

TITLE INDEX

spectra

- Fission rates, burnup, and neutron flux-fluence-spectra characterization for mixed oxide fuel experiments in the EBR-II* (416)
- Collection of gamma spectra data of irradiated light water moderated reactor spent fuel and study of the applicability of the methods for fuel identification. Final report for the period 1 February 1972--October 1975* (453)
- Raman spectra of solid ThCl_4 and fused ThCl_4 --KCl system* (488)
- Effect of luminescence centers on optical absorption, excitation and fluorescence spectra of CsCl:Pb * (491)
- Investigation into absorption infrared spectra of vapors under ScBr_3 , YCl_3 , LaCl_3 , GdCl_3 , ErCl_3 , LuCl_3 by means of isolation method in rare gas matrix* (497)
- Use of oscillator strengths of 'supersensitive' ion transfers of lanthanides for calculation of some parameters determining the intensity of their absorption spectra* (511)
- Diffuse reflectance spectra of protactinium(IV), uranium(IV) and neptunium(IV) tetrachlorides and tetraformates* (562)
- Spectra of combination scattering and structure of 3d-metals chloride complexes in molten salts* (674)
- IR spectra of the NbCl_5 , TaCl_5 , NbBr_5 and NbOCl_3 vapors* (707)

Spectral

- Spectral properties of impurity molecules of the C_{3v} symmetry groups in solid solutions of halide salts* (44)
- Spectral emissivities of molten refractory metals* (47)
- Uranium carbide: thermal diffusivity, thermal conductivity and spectral emissivity at high temperatures* (121)
- Thermal diffusivity, thermal conductivity and spectral emissivity of uranium dicarbide at high temperatures* (132)
- Spectral density functions for disordered systems* (229)
- The spectral properties of uranium hexafluoride and its thermal decomposition products* (345)
- X-ray spectral study of the carbides and nitrides of group IV and V metals* (582)
- Pressure effect on spectral properties of CeF_3 : Pr^{3+} * (677)

Spectrochemical

- Spectrochemical method for determination of zirconium in molten salts* (566)
- Standard methods for chemical, mass spectrometric, and spectrochemical analysis of nuclear-grade uranium dioxide powders and pellets* (711)
- Standard methods for chemical, mass spectrometric, and spectrochemical analysis of nuclear-grade plutonium dioxide powders and pellets* (712)
- Standard methods for chemical, mass spectrometric, and spectrochemical analysis of nuclear-grade mixed oxides (U, PuO_2)* (713)
- Standard methods for chemical, mass spectrometric, spectrochemical, nuclear, and radiochemical analysis of nuclear-grade plutonium metal* (714)

TITLE INDEX

Spectrographic

Spectrographic determination of trace impurities in reactor grade aluminium* (104)

Chemical separation and spectrographic estimation of rare earths and yttrium in PuO_2 and $(\text{U}, \text{Pu})\text{O}_2$ * (125)

spectrometer

Knudsen cell--mass spectrometer studies of cesium--urania interactions* (120)

High-temperature unit to the IKS-21 spectrometer for the detection of molten salts absorption spectra* (208)

Separation of alkaline earth from alkali metal fission products in the SOLAR on-line mass spectrometer* (534)

spectrometric

Mass spectrometric determination of dissociation energies of high-temperature molecules: ScS , YS , LaS , ZrS , UO * (597)

Standard methods for chemical, mass spectrometric, and spectrochemical analysis of nuclear-grade uranium dioxide powders and pellets* (711)

Standard methods for chemical, mass spectrometric, and spectrochemical analysis of nuclear-grade plutonium dioxide powders and pellets* (712)

Standard methods for chemical, mass spectrometric, and spectrochemical analysis of nuclear-grade mixed oxides $(\text{U}, \text{Pu})\text{O}_2$ * (713)

Standard methods for chemical, mass spectrometric, spectrochemical, nuclear, and radiochemical analysis of nuclear-grade plutonium metal* (714)

Standard method of test for atom percent fission in uranium and plutonium fuel (mass spectrometric method)* (715)

spectrometry

Investigations on radioactive fission product correlations: gamma spectrometry measurements on spent fuel assemblies discharged from the Trino Vercellese reactor at the end of the 2nd irradiation cycle* (75)

Flame emission and atomic absorption spectrometry. III. Elements and matrices* (319)

Quantitative determination of UO_2 , and ThO_2 in powdery mixtures by infrared spectrometry* (431)

Determination of trace impurities of titanium and zirconium in molybdenum and tungsten and their oxygen compounds by X-ray fluorescence spectrometry* (698)

spectrophotometer

Use of the SF-8 detection spectrophotometer for measuring absorption spectra of molten salts* (331)

spectrophotometric

Determination of thorium in rare earths chloride: coprecipitation with cerium (IV) iodate and spectrophotometric determination with thoron* (447)

Differential-spectrophotometric determination of niobium in its compounds* (490)

Spectrophotometric determination of rare earths and yttrium in gallium base alloys* (623)

Spectrophotometric determination of uranium in spent fuels of WWER-2 reactors* (632)

TITLE INDEX

Spectrophotometric

Spectrophotometric method for the determination of neptunium and plutonium in process solutions* (660)

spectrophotometry

Study of the compounds $M\text{UO}_2\text{F}_5$ ($M = \text{K, Rb, Cs, NH}_4$) by i.r. absorption and Raman diffusion

spectrophotometry. III. Effect of cation on vibration frequency of the ion $\text{UO}_2\text{F}_5^{3-}$ * (127)

Quantitative study of the chemical equilibria in solution in molten LiCl-KCl (70-30 mol.%) by absorption spectrophotometry.

Application to neptunium* (385)

Determination of the $\text{Np(IV)} \rightarrow \text{Np(III)}$ normal potential in fused LiCl-CsCl (55-45%) by absorption spectrophotometry* (386)

spectroscopic

Photoelectron spectroscopic studies of some transition metals and alloys* (417)

IR spectroscopic study of dipotassium and dicesium uranyl tetrabromides* (483)

Raman spectroscopic studies of yttrium (III) chloride--alkali metal chloride melts and of $\text{Cs}_2\text{NaYCl}_6$ and YCl_3 solid compounds* (489)

A Moessbauer spectroscopic study of intercalation compounds of graphite with ferric chloride and aluminum chloride* (627)

X-ray diffraction and infrared: Raman spectroscopic study of the hydrogen dinitrate ion, $(\text{O}_2\text{NO-H-ONO})_2^{2-}$, in cesium hydrogen dinitrate* (695)

spectroscopy

Infrared spectroscopy of the uranium/oxygen system* (12)

Electronic structure of CeN studied by x-ray-photoemission spectroscopy* (28)

11. european congress on molecular spectroscopy. Abstracts* (44)

11. european congress on molecular spectroscopy. Abstracts* (46)

Investigation into reactions between trioctylammonium nitrates and rare earth nitrates by X-ray electron spectroscopy* (113)

Optical absorption Zeeman spectroscopy of rare earth orthochromites. II. Theory* (122)

Optical absorption Zeeman spectroscopy of rare earth orthochromites. I. ErCrO_3 * (123)

Investigation of some rare earth aluminates by the method of infrared spectroscopy* (357)

Electromodulation spectroscopy of sc and fcc phase TlCl and TlBr * (411)

Progress in the spectroscopy of rare earth and transition metal ions* (412)

Electronic and magnetic properties of rare earth- Sn_3 compounds from ^{119}Sn Moessbauer spectroscopy* (559)

Optical spectroscopy of ^{237}Np in CaF_2 . Final report* (718)

Nuclear spectroscopy. Laboratory of chemistry and radiochemistry* (723)

spectrum

Electronic structure of highly symmetric complexes of lanthanoids and actinoids. Pt. 3. 'Electronic' Raman spectrum of $\text{Cs}_2\text{U(IV)Cl}_6$ * (13)

TITLE INDEX

spectrum

Electronic structure of highly symmetric complexes of lanthanoids and actinoids. Pt. 4. The electronic Raman spectrum of $\text{Cs}_{2\text{NaEu(III)Cl}_6}$ * (14)
 Optical absorption spectrum of UO_2 doped LiF * (30)
 Electron spectrum of light alkali metals* (161)
 Absorption spectrum of HfCl molecule* (283)
 Predicted infrared spectrum and the structure of the isolated UF_5 molecule* (350)
 Isotope effects in the molecular spectrum of plutonium hexafluoride* (353)
 Lines in the spectrum of ^7LiH (4728--5298 Å)* (374)
 Assignments in the electronic spectrum of UF_6 * (413)
 Frequency spectrum and scattering law for liquid copper. A molecular dynamics investigation* (567)
 Electronic structure of highly symmetric lanthanoid and actinoid compounds. VI. The electronic Raman spectrum of $\text{Cs}_{2\text{NaCe(III)Cl}_6}$ * (678)
 Accurate nuclear parameters derived from the Moessbauer spectrum of thulium metal* (696)

Speed

Speed of ultrasound and the thermophysical properties of the liquid metals Sn, Pb, Cd and of their binary alloys Pb:Sn and Pb:Cd* (403)

spent

Review of partitioning proposals for spent nuclear fuels* (69)
 Investigations on radioactive fission product correlations: gamma spectrometry measurements on spent fuel assemblies discharged from the Trino Vercellese reactor at the end of the 2nd irradiation cycle* (75)
 Chemical reprocessing of spent nuclear fuels* (322)
 Collection of gamma spectra data of irradiated light water moderated reactor spent fuel and study of the applicability of the methods for fuel identification. Final report for the period 1 February 1972--October 1975* (453)
 Spectrophotometric determination of uranium in spent fuels of WWER-2 reactors* (632)

sphere

Theory for the derivation of a single hard-sphere potential describing the overall molecular interactions in binary liquid metal alloys as a function of temperature and composition* (525)
 Application of the hard-sphere model to thermodynamic properties of liquid 3d transition metals* (615)
 Effect of second sphere cations on formation of nickel (2) chloride complexes in molten salts* (673)

Spin

Spin structure and elastic properties of uranium mononitride* (368)

Sponsored

Second specialist meeting on sodium/fuel interaction in fast reactors, Ispra, 21-23 November 1973. Sponsored by the Committee on the Safety of Nuclear Installations, OECD Nuclear Energy Agency, Paris, France* (272)

Spreading

Spreading of iron-nickel alloys on ZrC and nickel-cobalt alloys on ZrN* (252)

TITLE INDEX

Spring

Spring meeting, 26 and 27 April 1976, Amsterdam* (250)

Sr

Gases and carbon in metals (thermodynamics, kinetics, and properties). I. Alkali metals, alkaline earth metals, light metals (Li, Na, K, Rb, Cs; Ca, Sr, Ba; Be, Mg, Al)* (192)
 Solubility of $\text{Sr}(\text{NO}_3)_2$ in the $\text{LiNO}_3\text{-KNO}_3$ melt of variable composition* (351)
 Solubility of rare earth and yttrium oxochlorides in molten salts - MCl_2 (M-Mg, Ca, Sr, Ba)* (495)

SrF

Ionic-thermocurrent study of the dipole relaxation and equilibrium in Gd-doped SrF_2 * (410)
 Phases of variable composition with LaF_3 structure in $\text{MF}_2\text{-(Y, Ln)F}_3$ systems. 1. $\text{SrF}_2\text{-(Y, Ln)F}_3$ systems (thermal characteristics, formation of berthollides)* (593)

SrZrF

New hexafluorozirconates(IV): BaZrF_6 , PbZrF_6 , EuZrF_6 , SrZrF_6 * (419)

Stability

Stability of rare earth metaniobates in hydrogen atmosphere* (164)
 Stability of rare earth tetrafluorides* (312)
 Preparation and thermochemical stability of uranium-zirconium-carbonitrides. A contribution to nuclear fuels for advanced high-temperature reactors* (333)
 Preparation and thermochemical stability of uranium-zirconium-carbonitrides* (334)
 Electrochemical study of high-temperature stability of compounds between the rare earths and copper oxide* (634)

stability

Preparation and thermochemical stability of uranium-zirconium-carbonitrides.* (332)

Stable

Stable inorganic matrix materials for high temperature batteries. Final report No. 1, 2 June 1967--2 February 1968* (23)

Standard

Standard formation enthalpy of chloride alloys of $\text{ZrCl}_4\text{-KCl(NaCl)}$ systems* (8)
 Standard potential of sodium amalgam at 25°C * (32)
 Standard free energy of formation of liquid Na_2CrO_4 , 1100°C to 1220°C * (375)
 Cesium nitrate. Standard enthalpies of solution and formation and the standard entropy at 298.15 K * (466)
 Lithium nitride (Li_3N): standard enthalpy of formation by solution calorimetry* (467)
 Standard methods for chemical, mass spectrometric, and spectrochemical analysis of nuclear-grade uranium dioxide powders and pellets* (711)
 Standard methods for chemical, mass spectrometric, and spectrochemical analysis of nuclear-grade plutonium dioxide powders and pellets* (712)

TITLE INDEX

Standard

Standard methods for chemical, mass spectrometric, and spectrochemical analysis of nuclear-grade mixed oxides (U,Pu) $O_{2.5}$ * (713)

Standard methods for chemical, mass spectrometric, spectrochemical, nuclear, and radiochemical analysis of nuclear-grade plutonium metal* (714)

Standard method of test for atom percent fission in uranium and plutonium fuel (mass spectrometric method)* (715)

Standard method of test for atom percent fission in uranium and plutonium fuel (neodymium-148 method)* (716)

Static

Static structure factor of liquid caesium* (248)

Static structure factor of liquid Li* (273)

Statistical

Statistical mechanics of intermediate-valence rare earth compounds* (279)

steam

Investigation of liquid-steam equilibrium in system with diluted solutions of metal fluorides in uranium hexafluoride* (526)

stirred

Determination and correlation of mass transfer coefficients in a stirred cell* (236)

stoichiometric

Prediction of the inert gas solubilities in stoichiometric molten $UO_{2.5}$ * (225)

Studies of non-stoichiometric oxides by thermoanalytical methods* (626)

stoichiometries

Radial oxygen distribution in the fuel of irradiated mixed oxide fuel elements with different final stoichiometries* (316)

stoichiometry

Decomposition, stoichiometry and structure of neptunium oxides* (171)

Thermodynamics of non-stoichiometry. A model for $UC_{x/2}$ * (442)

storage

Theoretical concepts useful in the calculation or storage of phase diagrams of ionic systems* (552)

strengths

Use of oscillator strengths of "'supersensitive'" ion transfers of lanthanides for calculation of some parameters determining the intensity of their absorption spectra* (511)

stress

Effect of composition and structure on crevice, intergranular, and stress corrosion of some wrought Ni--Cr--Mo alloys* (604)

strong

Structural study of liquids with strong short-range correlation in the atomic distribution* (609)

Structural study of liquids with strong short-range correlation in the atomic distribution* (653)

strontium

Ternary systems from lithium, sodium, strontium and barium chlorides* (86)

Phase equilibria in the system of sodium chlorides, cesium chlorides and strontium chlorides* (87)

TITLE INDEX

strontium

Phase equilibria in the system of sodium chlorides, cesium chlorides and strontium chlorides* (88)
 Quaternary mutual system from fluorides and chlorides of lithium, sodium and strontium* (89)
 Reaction between sodium, rubidium and strontium chlorides* (286)
 Calculation of thermodynamic activity of components in strontium liquid alloys* (446)

Structural

Structural mechanics in reactor technology. Vol. 1* (19)
 Some observations on the compatibility of structural materials with molten lead* (25)
 Mechanical properties test data for structural materials. Quarterly progress report for period ending April 30, 1977* (57)
 Structural study and properties of the alkali metal, nitrosyl, and ammonium hepta- and octafluorouranates(VI)* (66)
 Environmental sensitivity of structural metals: liquid metal embrittlement. Annual technical progress report No. 2, 25 Jun 1970--31 May 1971* (95)
 Structural changes in the series LnFeO_3 and LnF_3 * (197)
 Environmental sensitivity of structural metals: liquid metal embrittlement. Annual technical progress report No. 4, 1 Jun 1972--31 May 1973* (211)
 Environmental sensitivity of structural metals: liquid metal embrittlement. Annual technical progress report No. 2, 25 Jun 1970--31 May 1971* (212)
 Report of task force on alternate structural materials for Liquid Metal Fast Breeder Reactors* (309)
 Effect of various additions on the temperature of structural transformation in liquid iron* (313)
 Effect of oxygen content on the structural characteristics of uranium oxycarbides* (362)
 Rare earth metal-metal halide systems. 19. Structural characterization of the reduced holmium chloride $\text{Ho}_{0.5}\text{Cl}_{1.5}$ * (378)
 Structural or solid state chemistry of rare earth sesquioxides from Goldschmidt to the present time* (438)
 Some structural characteristics of rubidium and cesium in the vicinity of their melting points* (460)
 NMR study on structural microinhomogeneity of fluoride-oxide melts* (463)
 Mechanical properties test data for structural materials. Quarterly progress report for period ending April 30, 1977* (471)
 Preparational, structural and vibrational study of $\text{M}_2\text{U}_{2.5}\text{Cl}_4$, $2\text{H}_2\text{O}$ (M = Rb, Cs). Detection of a tetranuclear anion $[(\text{UO}_2)_4(\text{O})_2(\text{Cl})_8(\text{H}_2\text{O})_2]^{4-}$ * (498)
 Some trends in constitutive equation model development for high-temperature behavior of fast-reactor structural alloys* (527)
 Basic mechanisms providing oxidation resistance in structural metals at high temperatures. II. The defect structure of nonstoichiometric rutile (TiO_{2-x}) and nonstoichiometric rutile containing 1 to 10 mole % $\text{Ce}_{2.6}\text{O}_{6.6}$. Final report 21 Dec 1973-20 Dec 1974* (542)

TITLE INDEX

structural

- New representatives of $\text{CeMn}_{4}\text{Al}_{8}$ structural type* (549)
- Properties of structural materials for sodium-cooled fast breeder reactor* (580)
- Structural mechanics in reactor technology. Vol. 2* (601)
- Structural study of liquids with strong short-range correlation in the atomic distribution* (609)
- Structural study of liquids with strong short-range correlation in the atomic distribution* (653)
- Mechanical properties test data for structural materials.
- Quarterly progress report for period ending April 30, 1977* (692)
- Fuel-clad interaction. C-E thermo-structural fuel evaluation methods* (719)
- Summary of U.S. LMFBR programs on high temperature structural design and associated materials testing* (731)

structure

- Electronic structure of highly symmetric complexes of lanthanoids and actinoids. Pt. 3. 'Electronic' Raman spectrum of $\text{Cs}_{2}\text{U(IV)Cl}_{6}$ * (13)
- Electronic structure of highly symmetric complexes of lanthanoids and actinoids. Pt. 4. The electronic Raman spectrum of $\text{Cs}_{2}\text{NaEu(III)Cl}_{6}$ * (14)
- Electronic structure of CeN studied by x-ray-photoemission spectroscopy* (28)
- Studies of the electronic structure of simple metals and their alloys with the aid of Compton scattering and positron annihilation using lithium, magnesium, lithium-magnesium and aluminium-zinc as examples* (48)
- Vibrational fine structure in the uranium emission of $\text{Mg}_{3}\text{TeO}_{6}$ and $\text{Y}_{6}\text{WO}_{12}$ * (59)
- Atomic structure of melts of complex system with eutectic transformations* (155)
- Structure factor of liquid aluminum and calculation of transport coefficients* (159)
- Zeeman effect and satellite carrier structure in absorption spectra of pure and doped praseodymium and europium double nitrates* (165)
- Decomposition, stoichiometry and structure of neptunium oxides* (171)
- Discrete variation of the structure and texture of liquid metals formed by components with discrete packing* (177)
- Structure and excitations of amorphous solids* (183)
- Structure of the biexciton molecule and optical transition selection rules in zincblende and wurtzite materials* (186)
- Bonding and the electronic structure of the actinide metals* (188)
- Electronic structure and properties of the actinides* (189)
- Magnetic susceptibility and electronic structure in metallic and non-metallic liquid cesium-gold and cesium-antimony alloys* (190)
- Actinide metallocarborane complex; synthesis and x-ray structure determination of the $\text{bis}[\text{n}^{\text{sup}}5\text{-}(3)\text{-}1,2\text{-dicarbollyl}]\text{dichlorouranium(IV) dianion}$ * (194)
- Structure of $\text{NO}_{3}^{\text{sup}}\text{-}$ radical in molten monovalent metal nitrates* (195)
- Electron structure of rare-earth metal tetraborides* (215)

TITLE INDEX

structure

- Electron structure and bonding in ThO_2 and UO_2 * (223)
- Structure of the thorium-(zirconium, niobium, ruthenium, rhodium)-carbon systems* (245)
- Static structure factor of liquid caesium* (248)
- Structure factor of liquid caesium and sodium-caesium alloys* (249)
- Structure factors of liquid alkali metals* (250)
- Band structure and chemical bond in ThX compounds* (259)
- Static structure factor of liquid Li* (273)
- Structure and anodic discharge behavior of lithium-boron alloys in the LiCl-KCl eutectic melt.* (274)
- Structure of lithium--boron alloys and their performance as potential anodes for high power molten salt batteries. Progress report, 1 Aug 1974--30 Jun 1975* (275)
- Structure and Bonding* (280)
- Band structure studies of actinide systems* (323)
- Investigation of formation thermodynamic of rare earth aluminates with perovskite structure* (326)
- Predicted infrared spectrum and the structure of the isolated UF_5 molecule* (350)
- Structure of molten Bi--Sb alloys by means of neutron diffraction* (359)
- R_3T_2 compounds (R = rare earth or Y; T = Rh, Pd, Pt) with the rhombohedral Er_3Ni_2 structure type* (365)
- Spin structure and elastic properties of uraniummononitride* (368)
- Effect of temperature on the fine structure of Eu^{2+} in KCl and KI * (432)
- KKR analysis of the Anderson-McMillan theory of electronic structure of liquid metals. Technical report* (485)
- Structure and Bonding* (492)
- Using molecular dynamics method for the evaluation of Fe-Ni and Al-Zn melts structure properties* (512)
- Structure of In_2Bi melted compound* (514)
- Contribution to the problem of the thermodynamic properties and the structure of liquid Cd-In alloys* (524)
- Structure and Bonding* (538)
- Basic mechanisms providing oxidation resistance in structural metals at high temperatures. II. The defect structure of nonstoichiometric rutile (TiO_2) and nonstoichiometric rutile containing 1 to 10 mole % Ce_2O_3 . Final report 21 Dec 1973-20 Dec 1974* (542)
- Oxygen potential and vacancy structure of the mixed oxide $(\text{UPu})\text{O}_{2-x}$ * (565)
- Structure of $4f^{6s}$ and $4f^{8s}$ levels and Eu^{3+} and Th^{3+} configurations in short-range ultra-violet* (589)
- Phases of variable composition with LaF_3 structure in MF_2 -(Y,Ln) F_3 systems. 1. SrF_2 -(Y,Ln) F_3 systems (thermal characteristics, formation of berthollides)* (593)
- Preparation and structure of transition metal fluorides $\text{U}^{II}\text{M}^{IV}\text{F}_6$ [U^{II} : Cr-Zn; M^{IV} : Hf, Zr]* (596)
- Effect of composition and structure on crevice, intergranular, and stress corrosion of some wrought Ni--Cr--Mo alloys* (604)

TITLE INDEX

structure

- Some structure differences of double oxides of indium and gallium with antimony(5)* (657)
- Structure of Nd^{3+} terms in nitrate glasses* (664)
- Compounds with the structure of ThMn_{12} type in the R--Fe--Al systems* (671)
- Spectra of combination scattering and structure of 3d-metals chloride complexes in molten salts* (674)
- Electronic structure of highly symmetric lanthanoid and actinoid compounds. VI. The electronic Raman spectrum of $\text{Cs}_2\text{NaCe(III)Cl}_6$ * (678)
- Electronic structure of actinides and related metals* (683)
- Effects of secondary ligands on the electronic structure of uranyls* (684)
- Structure of liquid transition metals and their alloys* (686)
- Structure of rare earth metals in the liquid state* (688)
- Structure of liquid Ce-Ni alloys* (689)
- Electronic states densities from structure and thermodynamic data for liquid transition metals* (705)

structures

- Fault structures in rare earth--cobalt intermetallics* (11)
- Mixed valence rare-earth halides and their unusual crystal structures* (29)
- Vibrational spectra of transition metal hexafluoride crystals. III. Exciton band structures of MoF_6 , WF_6 , and UF_6 . Technical report* (52)
- Perturbation treatment of the Percus-Yevick equation and structures of liquid magnesium and cadmium* (210)
- Phase relations, crystal structures and physical properties of nuclear fuels* (612)
- Structures of 3d-transition metals in the liquid state* (687)

sub

- Thermal analysis of $\text{GaCl}_3 + 3\text{In} = \text{InCl} + \text{Ga}$ system* (4)
- Thermodynamic properties of diluted solutions of MnCl_2 in alkali metal chlorides* (6)
- Standard formation enthalpy of chloride alloys of ZrCl_4 and KCl(NaCl) systems* (8)
- Electronic structure of highly symmetric complexes of lanthanoids and actinoids. Pt. 3. 'Electronic' Raman spectrum of $\text{Cs}_2\text{U(IV)Cl}_6$ * (13)
- Electronic structure of highly symmetric complexes of lanthanoids and actinoids. Pt. 4. The electronic Raman spectrum of $\text{Cs}_2\text{NaEu(III)Cl}_6$ * (14)
- Thermal expansion of PuO_2 * (15)
- Optical spectra of the difluoride, dichloride, and trichloride ions in the matrix-isolated $\text{M}^{2+} + \text{F}_2^{2-}$, $\text{M}^{2+} + \text{Cl}_2^{2-}$, and $\text{M}^{2+} + \text{Br}_2^{2-}$ species* (16)
- Optical spectra of the dibromide and diiodide ions in the matrix-isolated $\text{M}^{2+} + \text{Br}_2^{2-}$ and $\text{M}^{2+} + \text{I}_2^{2-}$ species* (17)
- Study of KBeF_5 - DyF_3 cross-section of KF-BeF_2 - DyF_3 ternary system and some properties of this cross section glasses* (21)
- Study of chemical interaction in Pb_3AlF_6 - LaF_3 - CaF_2 system* (26)
- Optical absorption spectrum of NO_2 doped LiF * (30)

TITLE INDEX

sub

- Nuclear quadrupole resonance spectra of HgCl_2 under pressure* (31)
- Vibrational spectra and force constants of the elpasolites Cs_2KMgF_6 ($M = \text{Sc, Y, La, Gd, Yb}$)* (41)
- Density, viscosity, electric conductivity and meltability of Na_3AlF_6 - La_2O_3 and Na_3AlF_6 - Sm_2O_3 systems* (43)
- Me_2ZrCl_6 -- H_2 ($\text{Me} = \text{Na, K, Rb, Cs}$) system* (45)
- Ternary UO_2 -- UO_3 -- EuO system and investigation of Eu(II) --actinide(IV)--perovskites* (50)
- Vibrational spectra of transition metal hexafluoride crystals. III. Exciton band structures of MoF_6 , WF_6 , and UF_6 . Technical report* (52)
- Interactions in inorganic molecular crystals - electronic spectra of ReF_6 pure and mixed crystals. Technical report, 1 Jul--31 Dec 1975* (53)
- Vibrational fine structure in the uranium emission of MgTe_6 and Y_6WO_{12} * (59)
- Calculation of the oxygen potential of $(\text{U, Pu})\text{O}_{2-x}$ with defect models and measured oxygen potential data in the temperature range from 1,000 to 1,700 K* (74)
- Measurement of mass transfer coefficients in a mechanically agitated, nondispersing contactor operating with a molten mixture of LiF -- BeF_2 -- ThF_4 and molten bismuth* (79)
- Preparation and some properties of actinide(III) and lanthanide(III) oxide halides, and the enthalpies of formation of $\text{PuBr}_3(s)$ and $\text{PuI}_3(s)$ * (82)
- Tensimetric study of pressure and composition of saturated vapor in TlCl -- InCl_3 system* (93)
- Investigation into thermodynamic properties of FeCl_2 -- CsCl system melts* (96)
- Study of saturated vapor pressure above salt melts in CoCl_2 -- MCl systems* (98)
- Experimental determination of formation heat of gaseous Ga_2O and In_2O * (105)
- High-temperature microcalorimetric measurements of partial molar enthalpy of solution, $\Delta H(O_2)$, in plutonium + oxygen* (107)
- Phase diagram of system LiCl -- KCl -- CaCrO_4 * (116)
- Phase diagram for the ternary system CaCl_2 -- KCl -- CaCrO_4 * (117)
- Optical absorption Zeeman spectroscopy of rare earth orthochromites. I. ErCrO_3 * (123)
- Chemical separation and spectrographic estimation of rare earths and yttrium in PuO_2 and $(\text{U, Pu})\text{O}_2$ * (125)
- Study of the compounds $\text{M}_3\text{UO}_2\text{F}_5$ ($M = \text{K, Rb, Cs, NH}_4$) by i.r. absorption and Raman diffusion spectrophotometry. III. Effect of cation on vibration frequency of the ion $\text{UO}_2\text{F}_5^{3-}$ * (127)
- ^{139}La ENDOR of LaCl_3 : Gd^{3+} * (130)
- Distribution coefficient for U and Pu in the system $(\text{UPu})\text{C} + (\text{UPu})_2\text{C}_3$ * (134)
- Phase diagram of the ThCl_4 -- LiCl -- PbCl_2 system* (143)
- Viscosity of UCl_3 -- MgCl_2 binary system melts* (145)
- Viscosity of molten CaCl_2 -- UCl_3 mixtures* (146)

TITLE INDEX

sub

- Phase diagrams of Li_2MoO_4 -- ZnMoO_4 , K_2MoO_4 -- ZnMoO_4 and K_2WO_4 -- ZnWO_4 systems* (160)
- Phase relations in PbWO_4 -- $\text{TR}_2(\text{WO}_4)_3$ systems* (168)
- Physicochemical investigation of rare earth titanoniobates, LnTiNbO_6 * (173)
- Critical assessment of equation of state data for UO_2 * (180)
- Heat capacity of $^{242}\text{PuO}_2$ from 12 to 350°K and of $^{244}\text{PuO}_2$ from 4 to 25°K. Entropy, enthalpy, and Gibbs energy of formation of PuO_2 at 298.15°K* (184)
- Vibration spectra of rare earth oxytungstates- Ln_2WO_6 composition* (185)
- Electronic conductivity and polarization effects of solid ThO_2 -- Y_2O_3 electrolytes and of oxygen containing Va-metal electrodes* (193)
- Structure of NO_3 radical in molten monovalent metal nitrates* (195)
- Structural changes in the series LnFeO_3 and LnF_3 * (197)
- Cryometric studies of the system CdCl_2 -- UCl_4 * (199)
- Raman spectra of Al_2O_3 solutions in molten cryolite and other aluminum fluoride containing melts* (203)
- Fine definition of ir spectra from high temperature interactions of U + O_2 . Part II. Final report, 1 July 1974--30 June 1975* (218)
- Electron structure and bonding in ThO_2 and UO_2 * (223)
- Prediction of the inert gas solubilities in stoichiometric molten UO_2 * (225)
- Measurement of fluorescence of rare earths in molten salts (Nd^{3+} in NaNO_3 -- KNO_3 eutectic)* (233)
- Thermochemistry of gaseous UF_5 and UF_4 * (238)
- High temperature thermal expansion of ThO_2 , MgO and Y_2O_3 by X-ray diffraction* (242)
- LiNO_3 -- CsNO_3 -- $\text{Ca}(\text{NO}_3)_2$ and NaNO_3 -- CsNO_3 -- $\text{Ca}(\text{NO}_3)_2$ system* (255)
- Entropy change for the lambda-type transition in nonstoichiometric $\text{U}_{40(9-y)}\text{O}_{100}$ * (260)
- Phase transition in doped $\text{U}_{40(9-y)}\text{O}_{100}$. I. Heat capacity measurement* (261)
- Lambda-type heat capacity anomalies in $\text{U}_{30}\text{O}_{80}$ * (262)
- Influence of small additions to the preparation and the properties of UO_2 ceramics. Part A. UO_2 -- Gd_2O_3 * (266)
- Influence of small additions on the preparation and the properties of UO_2 ceramics. Part B. The UO_2 -- TiO_2 system* (267)
- LiF -- ErF_3 system* (270)
- Phase diagrams of BiBr_3 -- BiI_3 system* (288)
- Thermal investigation of NdCl_3 -- AlCl_3 -- KCl and NdCl_3 -- AlCl_3 -- CsCl system* (289)
- Molecular vapor composition and mass spectra in KF -- ZrF_4 , CsF -- ZrF_4 systems* (294)

TITLE INDEX

sub

- Partial pressures, activities and evaporation heat in the KF-ZrF_4 system* (295)
- Interval of congruent and incongruent sublimation of $3\text{KF}\cdot\text{ZrF}_4$ compound in the KF-ZrF_4 system* (296)
- Thermodynamic state, specific heat, and enthalpy function of saturated UO_2 vapor between 3000 K and 5000 K* (297)
- Fusibility in $\text{K}_2\text{Nb}(\text{Ta})\text{F}_7\text{-LiF-NaF}$ systems* (298)
- Meltability in system of $\text{K}_2\text{TaF}_7\text{--NaF--NaCl--KCl}$ * (299)
- Meltability in $\text{K}_2\text{NbF}_7\text{-NaF-NaCl-KCl}$ system* (300)
- Reduction of PuO_2 by CO and equilibrium partial pressures above plutonia* (304)
- Investigation into americium extraction process from residues of irradiated nuclear fuel after its fluorination by $\text{SiF}_4\text{-NaF}$ and $\text{SiF}_4\text{-CaF}_2$ melts of eutectic composition* (308)
- Phase diagrams of $\text{BiCl}_3\text{-CuCl-TeCl}_4$ and $\text{CuCl-FeCl}_3\text{-TeCl}_4$ systems* (337)
- Double tungstate $\text{K}_5\text{La}(\text{WO}_4)_4$ * (338)
- Study of kinetics of monovalent indium ions formation in the $\text{In}_2\text{O}_3\text{-InCl}_3\text{-LiCl-H}_2\text{O}$ system* (341)
- Wetting of certain refractories of the system $\text{MgO-Al}_2\text{O}_3\text{-ZrO}_2$ with metallic melts* (344)
- Predicted infrared spectrum and the structure of the isolated UF_5 molecule* (350)
- Solubility of $\text{Sr}(\text{NO}_3)_2$ in the $\text{LiNO}_3\text{-KNO}_3$ melt of variable composition* (351)
- R_3T_2 compounds (R = rare earth or Y; T = Rh, Pd, Pt) with the rhombohedral Er_3Ni_2 structure type* (365)
- Thermophysical properties of UO_2 fuel materials* (366)
- Standard free energy of formation of liquid Na_2CrO_4 , 1100 K to 1220 K* (375)
- Rare earth metal-metal halide systems. 19. Structural characterization of the reduced holmium chloride $\text{Ho}_5\text{Cl}_{11}$ * (378)
- Solution studies in the US-UC_2 system* (379)
- Sodium uranium(V) trioxide, NaUO_3 : heat capacity and thermodynamic properties from 5 to 350 K* (383)
- Sodium uranium(V) trioxide, NaUO_3 : heat capacity and thermodynamic properties from 5 to 350 K* (384)
- Phase equilibria in the $\text{PbO-GeO}_2\text{-V}_2\text{O}_5$ ternary system* (394)
- Investigation of calcite solubility in K_2CO_3 and Na_2CO_3 solutions at temperatures of 200 to 350 K* (396)
- Oxygen redistribution in UO_2 due to a temperature gradient* (401)
- Diagram of excess free energy of the $\text{PbCl}_2\text{-CsCl-ZnCl}_2$ system* (402)
- Thermodynamic properties of dilute solutions of ThCl_4 in (Li-K)Cl and (Na-K) eutectics* (407)
- Synthesis and characterization of a new uranium(V) compound: $\text{H}_3\text{O}^+\text{UF}_6^-$ * (409)
- Ionic-thermoelectric study of the dipole relaxation and equilibrium in Gd-doped SrF_2 * (410)
- Assignments in the electronic spectrum of UF_6 * (413)

TITLE INDEX

sub

- Specific heat of PuC_{1-x} and Pu_2C_3 from 10 to 300 K* (414)
- Specific heat of PuC_{1-x} and Pu_2C_3 from 10 to 300 K* (415)
- New hexafluorozirconates(IV): BaZrF_6 , PbZrF_6 , EuZrF_6 , SrZrF_6 * (419)
- Quantitative determination of UO_2 , and ThO_2 in powdery mixtures by infrared spectrometry* (431)
- NdCl_3 - AlCl_3 - NaCl system* (434)
- FeCl_3 - MeCl systems* (435)
- Monte Carlo analysis of diffusion and thermodynamics in UO_{2+x} * (441)
- Thermodynamics of non-stoichiometry. A model for UC_x * (442)
- RbF - NdF_3 system* (443)
- Vaporization behavior of UO_2 * (445)
- PuO_2 dissolution problem for LWR plutonium recycle and LMFBR fuels: fabrication and reprocessing problems and their resolution* (455)
- Phase equilibrium in NdCl_3 - KCl system* (457)
- Density and viscosity of distillable products from ZrCl_4 and HfCl_4 interaction with FOCl_3 * (458)
- Thermal conductivity and diffusivity of $\text{U}_{0.9}\text{O}_{2.0}$ from 250 to 415 K* (459)
- Misch metal solubility in melt of LiF - BaF_2 system* (462)
- Lithium nitride (Li_3N): standard enthalpy of formation by solution calorimetry* (467)
- Phase relations in the alkaline earth fluoride-yttria systems (in three parts). 1. The CaF_2 - Y_2O_3 phase diagram* (468)
- Phase relations in the alkaline earth fluoride-yttria systems (in three parts). 2. The BaF_2 - Y_2O_3 phase diagram* (469)
- Study of the retention of Ba in UO_2 nuclear fuel particles by ZrO_2 as getter* (470)
- Phase diagrams in the pseudobinary systems of Li_2BeF_4 , alkali fluoroborates and alkali fluorides* (473)
- Extension of vapour pressure measurements of nuclear fuels (U , Pu) $\text{O}_{0.2}$ and UO_2 to 7000 K for fast reactor safety analysis* (475)
- Vapour pressure measurements to 7,000 K and equation of state of oxide fuels for fast reactor safety analysis. I. Extension of vapour pressure measurements of nuclear fuels (U , Pu) $\text{O}_{0.2}$ and UO_2 to 7,000 K* (477)
- Raman spectra of solid ThCl_4 and fused ThCl_4 - KCl system* (488)
- Raman spectroscopic studies of yttrium (III) chloride-alkali metal chloride melts and of $\text{Cs}_2\text{NaYCl}_6$ and YCl_3 solid compounds* (489)
- Solubility of rare earth and yttrium oxochlorides in molten salts - MCl_2 ($\text{M} = \text{Mg}, \text{Ca}, \text{Sr}, \text{Ba}$)* (495)
- Investigation into absorption infrared spectra of vapors under ScBr_3 , YCl_3 , LaCl_3 , GdCl_3 , ErCl_3 , LuCl_3 by means of isolation method in rare gas matrix* (497)

TITLE INDEX

sub

- Preparational, structural and vibrational study of $M_2U_2O_5Cl_4$, $2H_2O$ ($M = Rh, Cs$). Detection of a tetranuclear anion $[(UO_2)_4O_2Cl_8(H_2O)_2]^{4-}$ (498)
- Influence of salt background on kinetics of BF_4 -alkali hydrolysis* (504)
- Meltability diagram of $RbCl$ - $BeCl_2$ system* (506)
- Structure of In_2Bi melted compound* (514)
- Practical application of conversion method in study of five-component mutual system consisting of nine salts NaF , Na_2MoO_4 , Na_2WO_4 , KF , K_2MoO_4 , K_2WO_4 , BaF_2 , $BaMoO_4$, $BaWO_4$ * (516)
- Thermal analysis of the system of NaF , CaF_2 , Na_2MoO_4 , $CaMoO_4$ * (519)
- KF - KCl - K_2MoO_4 system* (520)
- Measurement of the cell voltage of ThO_2 - Y_2O_3 -solid electrolyte cells in the range of mixed conduction* (535)
- Energy transfer between Bi^{3+} yields Eu^{3+} , Bi^{3+} yields Sm^{3+} and UO_2 yields Eu^{3+} in oxide glasses* (539)
- K_2HF_6 - K_2ZrF_6 -1.25 percent of HF system at $40^\circ C$ with other solubility curves from $25^\circ C$ to $70^\circ C$ * (541)
- Basic mechanisms providing oxidation resistance in structural metals at high temperatures. II. The defect structure of nonstoichiometric rutile (TiO_2) and nonstoichiometric rutile containing 1 to 10 mole % Cb_2O_6 . Final report 21 Dec 1973-20 Dec 1974* (542)
- $GaTe$ - Bi_2Te_3 system* (545)
- $MoCl_2$ - $NaCl$ and $MoCl_2$ - KCl binary systems* (547)
- $ZrCl_4$ - $TiCl_3$ - $NaCl$ ternary system* (548)
- New representatives of $CeMn_4Al_8$ structural type* (549)
- $BiCl_3$ - $GeCl_4$ - $TeCl_4$ system* (553)
- $FeCl_3$ - $MoCl_5$ - $NbCl_5$ - WCl_6 system* (554)
- Behavior of ruthenium in fluoride-volatility process. II. Fluorination of RuO_2 by F_2 * (558)
- Electronic and magnetic properties of rare earth- Sn compounds from ^{119}Sn Moessbauer spectroscopy* (559)
- Chemistry of elements niobium and tantalum. 87. Studies on the thermal formation and decomposition of the complexes $[Nb_6Cl_{12}]^{2+}$, $[Nb_6I_8]^{3+}$, and $[Ta_6X_{12}]^{(n+)}$ ($X=Cl, Br, I$)* (561)
- Oxygen potential and vacancy structure of the mixed oxide $(UPu)_2O_{2-x}$ * (565)
- Covalent compounds of uranium of the type $F_x/U(OTeF_5)_{6-x}$ * (571)
- CsF - HoF_3 system* (572)
- Viscosity of melts of KCl - KF - K_2ZrF_6 ternary system* (576)
- Heat transfer measurements in a forced convection loop with two molten-fluoride salts: LiF - BeF_2 - ThF_4 - UF_4 and eutectic $NaBF_4$ - NaF * (585)

TITLE INDEX

sub

Phases of variable composition with LaF_3 structure in MF_2 -(Y,Ln) F_3 systems. 1. SrF_2 -(Y,Ln) F_3 systems (thermal characteristics, formation of berthollides)* (593)

Thermodynamics of the lanthanide halides. II. Heat capacities and Schottky anomalies of SmCl_3 , EuCl_3 and GdCl_3 from 50 to 350 K* (594)

Preparation and structure of transition metal fluorides $\text{U}^{\text{II}}\text{M}^{\text{IV}}\text{F}_6$ [U^{II} : Cr-Zn; M^{IV} : Hf, Zr]* (596)

Thermogravimetric investigation of the PbCl_2 - PbBr_2 2 PbI_2 and PbCl_2 - PbBr_2 2 PbF_2 systems* (602)

Degree of thermal dissociation of MCl_3 (M-K, Rb, Cs) complexes during melting* (624)

Systems of NbCl_5 - GaCl_3 and TaCl_5 - GaCl_3 * (633)

Thermal analysis of NaF , NaCl , NaMoO_4 system* (636)

System Na_2WO_4 - NaCl - CaWO_4 - CaCl_2 * (637)

Na_2MoO_4 - BaMoO_4 - NaF - BaF_2 system* (638)

Phases of high pressure in UO_2 - UO_3 - PbO system* (641)

Molten salt chemistry : Part IV - Solubility behaviour of PuF_3 in fluoride salts of interest in molten salt reactor technology* (655)

Thermodynamic properties of NbO , $\text{Fe}_{0.95}\text{O}$, CoO , NiO , Cu_2O and CuO oxides as comparative electrodes in electromotive forces method* (658)

Constitution equilibrium diagram of the LiNO_3 - LiIO_3 , NaNO_3 - NaIO_3 , KNO_3 - KIO_3 system* (665)

Phase diagram of NaNO_3 - $\text{Nd}(\text{NO}_3)_3$ system* (667)

Phase diagrams of the $\text{Mg}(\text{NO}_3)_2$ - $\text{Nd}(\text{NO}_3)_3$ system* (668)

Interdiffusion in the CaF_2 - YF_3 system* (669)

Compounds with the structure of ThMn_{12} type in the R-Fe-Al systems* (671)

Pressure effect on spectral properties of CeF_3 : Pr^{III} * (677)

Electronic structure of highly symmetric lanthanoid and actinoid compounds. VI. The electronic Raman spectrum of $\text{Cs}_2\text{NaCe}(\text{III})\text{Cl}_6$ * (678)

Phase equilibria and thermodynamic data for the ternary UO_2 - UCl_3 - BeO (1.5) (Ho, Tm, Lu) systems* (694)

X-ray diffraction and infrared: Raman spectroscopic study of the hydrogen dinitrate ion, $(\text{O}_2\text{N}-\text{H}-\text{ONO})^+$, in cesium hydrogen dinitrate* (695)

Measurement of interfacial energies in the UO_2 - Gd_2O_3 system* (697)

Solubility of nitrogen in liquid lithium and thermal decomposition of solid Li_3N * (702)

Preparation of thorium metal by electrolysis of fused salt ThCl_4 - LiCl - NaCl - KCl (eut) system* (703)

IR spectra of the NbCl_5 , TaCl_5 , NbBr_5 and NbOCl_3 vapors* (707)

TITLE INDEX

sub

Standard methods for chemical, mass spectrometric, and spectrochemical analysis of nuclear-grade mixed oxides (U,Pu)O₂* (713)
Optical spectroscopy of ²³⁷Np in CaF₂. Final report* (718)

sublimation

Interval of congruent and incongruent sublimation of 3KFxZrF₄ compound in the KF-ZrF₄ system* (296)

substoichiometric

Evaporation behavior and high-temperature thermal analysis of substoichiometric plutonium oxide in the composition range from O/Pu = 1.51 to 2.00* (482)

sulfate

High-temperature heat contents and entropies of zirconium fluoride and zirconium sulfate* (591)

sulphur

Thermodynamic properties of sulphur in liquid antimony* (642)

Summaries

Summaries of reports of the all-union meeting on solid body chemistry. 3--5 June 1975. III. Synthesis, phase equilibrium and use of solid oxide compounds* (114)
Summaries of reports of the all-union meeting on solid body chemistry. 3--5 June 1975. III. Synthesis, phase equilibrium and use of solid oxide compounds* (641)

summary

Thermochemistry of gaseous compounds of metals. Annual summary report 5 Dec 74--4 Dec 75* (240)
Thermochemistry of gaseous compounds of metals. Annual summary report, 5 Dec 1975--4 Dec 1976* (241)
Summary of U.S. LMFR programs on high temperature structural design and associated materials testing* (731)

sup

Thermal expansion of ²³⁸PuO₂* (15)
Optical spectra of the difluoride, dichloride, and trichloride ions in the matrix-isolated M⁺F₂⁺-, M⁺Cl₂⁺-, and M⁺Cl₃⁺- species* (16)
Optical spectra of the dibromide and diiodide ions in the matrix-isolated M⁺Br₂⁺- and M⁺I₂⁺- species* (17)
Preparation of ¹⁴⁷Pm metal and the determination of the melting point and phase transformation temperatures* (18)
Optical absorption spectrum of UO₂²⁺ doped LiF* (30)
Standard potential of sodium amalgam at 25^oC* (32)
X-ray study of the plutonium--oxygen phase diagram at ambient temperatures up to 1100^oC* (62)
Study of the compounds M₃UO₂F₅ (M = K, Rb, Cs, NH₄) by i.r. absorption and Raman diffusion spectrophotometry. III. Effect of cation on vibration frequency of the ion UO₂²⁺F₅³⁻* (127)
¹³⁹La ENDOR of LaCl₃:Gd³⁺* (130)
Solubility of sodium iodide in liquid sodium between 140 and 500^oC* (154)
Heat capacity of ²⁴²PuO₂ from 12 to 350^oK and of ²⁴⁴PuO₂ from 4 to 25^oK. Entropy, enthalpy, and Gibbs energy of formation of PuO₂ at 298.15^oK* (184)

TITLE INDEX

sup

- Actinide metallocarborane complex; synthesis and x-ray structure determination of the bis[n^{sup} 5⁺-(3)-1,2-dicarbollyl]dichlorouranium(IV) dianion* (194)
- Structure of NO₃^{sup} - radical in molten monovalent metal nitrates* (195)
- Simple Raman cell and furnace usable at temperatures higher than 1000^{sup} °C for corrosive melts* (204)
- Measurement of fluorescence of rare earths in molten salts (Nd^{sup} 3⁺ in NaNO₃-KNO₃ eutectic)* (233)
- Thermophysical and electrophysical properties of uranium hexafluoride at temperatures of (1--11) x10^{sup} 3°K and pressures of 0.1--100 atm* (302)
- Study of kinetics of monovalent indium ions formation in the In^{sup} 0-InCl₃-LiCl-H₂O system* (341)
- Lines in the spectrum of ^{sup} 7LiH (4728--5298 Å)* (374)
- Standard free energy of formation of liquid Na₂CrO₄, 1100^{sup} °C to 1220^{sup} °K* (375)
- Investigation of calcite solubility in K₂CO₃ and Na₂CO₃ solutions at temperatures of 200 to 350^{sup} °C* (396)
- Synthesis and characterization of a new uranium(V) compound: H₃O^{sup} +UF₆^{sup} -* (409)
- Effect of temperature on the fine structure of Eu^{sup} 2⁺ in KCl and KI* (432)
- Half-life of ^{sup} 242Pu from precise low-temperature heat capacity measurements* (487)
- Preparational, structural and vibrational study of M₂U₂ 2₅Cl₄, 2H₂O (M = Rb, Cs). Detection of a tetranuclear anion [(UO₂)₄ 2₈Cl₈(H₂O)₂]^{sup} 4-* (498)
- Influence of salt background on kinetics of BF₄^{sup} - alkali hydrolysis* (504)
- Energy transfer between Bi^{sup} 3⁺ yields Eu^{sup} 3⁺, Bi^{sup} 3⁺ yields Sm^{sup} 3⁺ and UO₂^{sup} 2⁺ yields Eu^{sup} 3⁺ in oxide glasses* (539)
- K₂HfF₆-K₂ZrF₆ 1.25 percent of HF system at 40^{sup} °C with other solubility curves from 25^{sup} °C to 70^{sup} °C* (541)
- Electronic and magnetic properties of rare earth-Sn₃ compounds from ^{sup} 119Sn Moessbauer spectroscopy* (559)
- Chemistry of elements niobium and tantalum. 87. Studies on the thermal formation and decomposition of the complexes [Nb₆Cl₁₂]^{sup} 2⁺, [Nb₆I₈]^{sup} 3⁺, and [Ta₆X₁₂]^{sup} (n+) (X=Cl, Br, I)* (561)
- Thermodynamics of solid solutions CsCl-RbCl and CsBr-RbBr at 25^{sup} °C* (570)
- Structure of 4f^{sup} 6 and 4f^{sup} 8 levels and Eu^{sup} 3⁺ and Tb^{sup} 3⁺ configurations in short-range ultra-violet* (589)
- Some thermal properties of beryllium fluoride from 8^{sup} °C to 1,200^{sup} °K* (617)
- Structure of Nd^{sup} 3⁺ terms in nitrate glasses* (664)
- Pressure effect on spectral properties of CeF₃:Pr^{sup} 3⁺* (677)

TITLE INDEX

sup

X-ray diffraction and infrared: Raman spectroscopic study of the hydrogen dinitrate ion, $(O\text{sub } 2\text{NO-H-ONO}\text{sub } 2)\text{sup -}$, in cesium hydrogen dinitrate* (695)

Optical spectroscopy of $\text{sup } 237\text{Np}$ in $\text{CaF}\text{sub } 2$. Final report* (718)

superoxide

Reaction of potassium superoxide in excess potassium* (590)

supersensitive

Use of oscillator strengths of ''supersensitive'' ion transfers of lanthanides for calculation of some parameters determining the intensity of their absorption spectra* (511)

SURBOUM

Analysis of LMFBR explosion model experiments by means of the SUREOUM-II code* (601)

surface

Density and temperature dependence of the surface tension of molten bismuth, lead, and bismuth-lead alloys* (1)
Nonisothermal flow of liquid metals over the surface of solid ones* (101)

Density, surface tension, and viscosity of uranium trichloride-sodium chloride melting* (141)

Density, surface tension and viscosity of melts of uranium trichloride-lithium chloride* (142)

Thermodynamic perturbation theory for the surface tension and ion density profile of a liquid metal* (167)

Determination of surface tension of a metal with high vapor pressure* (472)

Interaction in AgCl-CsCl, CsCl-KCl systems and liquidus surface of AgCl-CsCl-KCl ternary system* (659)

susceptibility

Magnetic susceptibility and electronic structure in metallic and non-metallic liquid cesium-gold and cesium-antimony alloys* (190)

symmetric

Electronic structure of highly symmetric complexes of lanthanoids and actinoids. Pt. 3. 'Electronic' Raman spectrum of $\text{Cs}\text{sub } 2\text{U(IV)Cl}\text{sub } 6$ * (13)

Electronic structure of highly symmetric complexes of lanthanoids and actinoids. Pt. 4. The electronic Raman spectrum of $\text{Cs}\text{sub } 2\text{NaEu(III)Cl}\text{sub } 6$ * (14)

Electronic structure of highly symmetric lanthanoid and actinoid compounds. VI. The electronic Raman spectrum of $\text{Cs}\text{sub } 2\text{NaCe(III)Cl}\text{sub } 6$ * (678)

symmetry

Spectral properties of impurity molecules of the $\text{Csub } (3v)$ symmetry groups in solid solutions of halide salts* (44)

symposium

Third international symposium on fuel cells* (77)

Israel Nuclear Society symposium on ''Nuclear fuel in Israel'', Weizmann Institute of Science, Rehovot, 1.7.75* (226)

Proceedings of the symposium and workshop on advanced battery research and design, March 22--24, 1976* (269)

Synthesis

Summaries of reports of the all-union meeting on solid body chemistry. 3--5 June 1975. III. Synthesis, phase equilibrium and use of solid oxide compounds* (114)

TITLE INDEX

synthesis

Actinide metallocarborane complex; synthesis and x-ray structure determination of the bis[n^{sup} 5\$-(3)-1,2-dicarbollyl]dichlorouranium(IV) dianion* (194)
 Synthesis of UN coatings on uranium* (400)
 Synthesis and characterization of a new uranium(V) compound: H₃CO^{sup} + UF₆^{sup} - \$* (409)
 Summaries of reports of the all-union meeting on solid body chemistry. 3--5 June 1975. III. Synthesis, phase equilibrium and use of solid oxide compounds* (641)
 Parameter for optimizing the synthesis of (U,Pu)C* (676)

system

Thermal analysis of GaCl₃+3In = InCl+Ga system* (4)
 Infrared spectroscopy of the uranium/oxygen system* (12)
 Study of KBe₂F₅-DyF₃ cross-section of KF-BeF₂-DyF₃ ternary system and some properties of this cross section glasses* (21)
 Study of chemical interaction in Pb₃AlF₆-LaF₃-Ca₃AlF₆ system* (26)
 Me₂ZrCl₆-H₂(Me-Na,K,Rb,Cs) system* (45)
 Ternary UO₂-UO₃-EuO system and investigation of Eu(II)--actinide(IV)--perovskites* (50)
 Segregation and vapour pressure studies on the uranium-plutonium-carbon system* (83)
 Segregation and vapour pressure studies in the uranium--plutonium--carbon system* (84)
 Phase equilibria in the system of sodium chlorides, cesium chlorides and strontium chlorides* (87)
 Phase equilibria in the system of sodium chlorides, cesium chlorides and strontium chlorides* (88)
 Quaternary mutual system from fluorides and chlorides of lithium, sodium and strontium* (89)
 Quaternary system of potassium halogenides* (90)
 Tensimetric study of pressure and composition of saturated vapor in TlCl-InCl₃ system* (93)
 Investigation into thermodynamic properties of FeCl₂-CsCl system melts* (96)
 Phase diagram of system LiCl--KCl--CaCrO₄* (116)
 Phase diagram for the ternary system CaCl₂--KCl--CaCrO₄* (117)
 Distribution coefficient for U and Pu in the system (UPu)C + (UPu)₂C₃* (134)
 Phase diagram of the ThCl₄-LiCl-PbCl₂ system* (143)
 Viscosity of UCl₃-MgCl₂ binary system melts* (145)
 Binary system of sodium, rubidium and cesium tungstates* (151)
 Atomic structure of melts of complex system with eutectic transformations* (155)
 On interaction in U--S--Te system* (162)
 Electrochemical studies of the sodium--bismuth system* (187)
 Crycmetric studies of the system CdCl₂-UCl₄* (199)
 Calculation and discussion of state diagrams in the system Pu-F-O-H* (222)
 Thermodynamic study of the U-O-C system by measurement of the carbon monoxide pressures at equilibrium* (235)
 Vaporization of uranium-carbon-nitrogen system* (251)
 Ternary system of potassium, cesium and calcium nitrates* (254)

TITLE INDEX

system

- $\text{LiNO}_3\text{-CsNO}_3\text{-Ca(NO}_3)_2$ and $\text{NaNO}_3\text{-CsNO}_3\text{-Ca(NO}_3)_2$ system* (255)
 Ternary mutual system of chlorides and iodides of rubidium and cadmium* (257)
 Influence of small additions on the preparation and the properties of UO_2 ceramics. Part B. The $\text{UO}_2\text{-TiO}_2$ system* (267)
 LiBr, RbBr, CsBr ternary system* (268)
 LiF-ErF_3 system* (270)
 Phase diagrams of $\text{BiBr}_3\text{-BiI}_3$ system* (288)
 Thermal investigation of $\text{NdCl}_3\text{-AlCl}_3\text{-KCl}$ and $\text{NdCl}_3\text{-AlCl}_3\text{-CsCl}$ system* (289)
 Thermodynamic study of the binary system copper-indium. I. Calorimetric study* (290)
 Thermodynamic study of the binary system copper--indium. II. Potentiometric study* (291)
 Partial pressures, activities and evaporation heat in the KF-ZrF_4 system* (295)
 Interval of congruent and incongruent sublimation of $3\text{KF}\cdot\text{ZrF}_4$ compound in the KF-ZrF_4 system* (296)
 Meltability in system of $\text{K}_2\text{TaF}_7\text{-NaF-NaCl-KCl}$ * (299)
 Meltability in $\text{K}_2\text{NbF}_7\text{-NaF-NaCl-KCl}$ system* (300)
 Study of kinetics of monovalent indium ions formation in the $\text{In}_2\text{O}_3\text{-InCl}_3\text{-LiCl-H}_2\text{O}$ system* (341)
 Wetting of certain refractories of the system $\text{MgO-Al}_2\text{O}_3\text{-ZrO}_2$ with metallic melts* (344)
 P--t diagram of the system uranium--nitrogen* (373)
 Phase diagram of the ternary system of lithium-, rubidium-, and cesium nitrates* (377)
 Solution studies in the US-UC_2 system* (379)
 Phase equilibria in the $\text{PbO-GeO}_2\text{-V}_2\text{O}_5$ ternary system* (394)
 Diagram of excess free energy of the $\text{PbCl}_2\text{-CsCl-ZnCl}_2$ system* (402)
 Thermodynamic study of the neptunium-hydrogen system* (428)
 $\text{NdCl}_3\text{-AlCl}_3\text{-NaCl}$ system* (434)
 Interaction parameters in the Zn-Pb-Sn system at low zinc concentration* (436)
 RbF-NdF_3 system* (443)
 Phase equilibrium in $\text{NdCl}_3\text{-KCl}$ system* (457)
 Electric conductivity of melts of the $\text{LiF-misch metal fluoride}$ system* (461)
 Misch metal solubility in melt of LiF-BaF_2 system* (462)
 Raman spectra of solid ThCl_4 and fused $\text{ThCl}_4\text{-KCl}$ system* (488)
 Contribution to the establishment of the phase diagram for the U-C-C system by means of x-ray diffractometry at high temperatures and under controlled pressure* (503)
 Meltability diagram of RbCl-BeCl_2 system* (506)
 Phase analysis on simulated LWR fuel rods and its evaluation with the aid of constitution studies in the system uranium-zirconium-oxygen* (509)

TITLE INDEX

system

- Phase analysis at simulated LWR fuel rods and their interpretation by constitution investigations in the uranium-zirconium-oxygen system* (510)
- Practical application of conversion method in study of five-component mutual system consisting of nine salts NaF , Na_2MoO_4 , Na_2WO_4 , KF , K_2MoO_4 , K_2WO_4 , BaF_2 , BaMoO_4 , BaWO_4 * (516)
- Thermal analysis of the system of NaF , CaF_2 , Na_2MoO_4 , CaMoO_4 * (519)
- $\text{KF-KCl-K}_2\text{MoO}_4$ system* (520)
- Thermodynamic investigation of liquid alloys in the system Cd-In-Tl * (523)
- Investigation of liquid-steam equilibrium in system with diluted solutions of metal fluorides in uranium hexafluoride* (526)
- $\text{K}_2\text{HF}_6\text{-K}_2\text{ZrF}_6$ 1.25 percent of HF system at 40°C with other solubility curves from 25°C to 70°C* (541)
- Ga-Eu-Te system* (544)
- $\text{GaTe-Bi}_2\text{Te}_3$ system* (545)
- Practice of experiment planning in studying limited regions of multicomponent system* (546)
- $\text{ZrCl}_4\text{-TiCl}_3\text{-NaCl}$ ternary system* (548)
- $\text{BiCl}_3\text{-GeCl}_4\text{-TeCl}_4$ system* (553)
- $\text{FeCl}_3\text{-MgCl}_2\text{-NbCl}_5\text{-WCl}_6$ system* (554)
- CsF-HoF_3 system* (572)
- Viscosity of melts of $\text{KCl-KF-K}_2\text{ZrF}_6$ ternary system* (576)
- Thermophysical properties of the ternary liquid-metal system tin--lead--cadmium* (605)
- Thermodynamic properties of the ternary Sn-Pb-Cd liquid metal system* (606)
- Total pressures of uranium- and plutonium-bearing species above the U-Pu-O system* (618)
- Quaternary mutual system of chlorides, bromides, iodides of sodium and rubidium* (629)
- Quaternary mutual system of chlorides, bromides, iodides of potassium and cesium* (630)
- Sections of quaternary mutual system consisting of sodium, potassium and cesium chlorides and bromides* (631)
- Thermal analysis of NaF , NaCl , Na_2MoO_4 system* (636)
- System $\text{Na}_2\text{WO}_4\text{-NaCl-CaWO}_4\text{-CaCl}_2$ * (637)
- $\text{Na}_2\text{MoO}_4\text{-BaMoO}_4\text{-NaF-BaF}_2$ system* (638)
- Quaternary mutual system of chlorides, bromides, iodides of rubidium and cesium* (640)
- Phases of high pressure in $\text{UO}_2\text{-UO}_3\text{-PbO}$ system* (641)
- Phase equilibrium study on system uranium-plutonium-tungsten-carbon* (645)
- Phase behavior and thermodynamics of U-Mo-C system* (646)
- Phase behavior and thermodynamics of U-Mo-C system* (647)
- Phase equilibria in system U-Pu-W-C * (648)
- Interaction in AgCl-CsCl , CsCl-KCl systems and liquidus surface of AgCl-CsCl-KCl ternary system* (659)
- Constitution equilibrium diagram of the $\text{LiNO}_3\text{-LiIO}_3$, $\text{NaNO}_3\text{-NaIO}_3$, $\text{KNO}_3\text{-KIO}_3$ system* (665)
- Phase diagram of $\text{NaNO}_3\text{-Nd(NO}_3)_3$ system* (667)

TITLE INDEX

system

Phase diagrams of the $\text{Mg}(\text{NO}_3)_2\text{-Nd}(\text{NO}_3)_3$ system* (668)
 Interdiffusion in the $\text{CaF}_2\text{-YF}_3$ system* (669)
 Measurement of interfacial energies in the $\text{UO}_2\text{-Gd}_2\text{O}_3$ system* (697)
 Preparation of thorium metal by electrolysis of fused salt $\text{ThCl}_4\text{-LiCl+NaCl+KCl(eut)}$ system* (703)
 Eutectic point of gallium-indium system* (706)
 Investigation into properties of liquidus curves of binary salt system melts where chemical compounds formation. 2. Systems of mono-monovalent and bi-bivalent cations with common anion* (708)
 Cesium, calcium and barium chlorides ternary system* (709)
 Research on a high-energy nonaqueous battery system. Quarterly progress report No. 1, 1 Dec 1965--1 Mar 1966* (720)
 Adaptation of the continuous cold trap system of fluidized-bed to the fluoride volatility process* (724)

systems

Standard formation enthalpy of chloride alloys of $\text{ZrCl}_4\text{-KCl(NaCl)}$ systems* (8)
 Density, viscosity, electric conductivity and meltability of $\text{Na}_3\text{AlF}_6\text{-La}_2\text{O}_3$ and $\text{Na}_3\text{AlF}_6\text{-Sm}_2\text{O}_3$ systems* (43)
 Phase equilibrium in interlanthanide oxides systems* (49)
 Phase equilibria in Li-Na-K carbonate/aluminate systems* (77)
 Ternary systems from lithium, sodium, strontium and barium chlorides* (86)
 Saturation vapor pressure in iron chloride-alkaline earth metal chlorides systems* (97)
 Study of saturated vapor pressure above salt melts in $\text{CoCl}_2\text{-MCl}$ systems* (98)
 Study on conductivity, density, and viscosity of molten salt systems* (112)
 Physical models of mass transport of iron and nickel in liquid sodium systems* (128)
 Thermographic investigation into ternary systems containing sodium and potassium chlorides, uranium trichloride and thorium tetrafluoride* (140)
 Calculation of thermodynamic activity and liquidus lines of components in binary systems of plutonium trichloride with alkali and alkaline earth metal chlorides* (147)
 Phase diagrams of ternary systems containing sodium and potassium chlorides, thorium tetrachloride and plutonium trichloride* (148)
 Phase diagrams of $\text{Li}_2\text{MoO}_4\text{-ZnMoO}_4$, $\text{K}_2\text{MoO}_4\text{-ZnMoO}_4$ and $\text{K}_2\text{WO}_4\text{-ZnWO}_4$ systems* (160)
 Phase relations in $\text{PbWO}_4\text{-TR}_2(\text{WO}_4)_3$ systems* (168)
 Some constitutional studies on uranium-carbon- and plutonium-carbon-rhenium and technetium systems* (227)
 Some constitutional studies on uranium--carbon and plutonium--carbon--rhenium and technetium systems* (228)
 Spectral density functions for disordered systems* (229)
 Structure of the thorium-(zirconium, niobium, ruthenium, rhodium)-carbon systems* (245)

TITLE INDEX

systems

- Nonaqueous electrolyte systems: ionic transport in nonaqueous media. Interim technical report No. 7* (246)
 Phase diagrams of LiCl, CsCl, CaCl and LiBr, CsBr, CaBr ternary systems* (256)
 TlCl-RbCl-CsCl and NaI-TlI-CsI systems* (258)
 Molecular vapor composition and mass spectra in KF-ZrF_4 , CsF-ZrF_4 systems* (294)
 Fusibility in $\text{K}_2\text{Nb}(\text{Ta})\text{F}_7$ -LiF-NaF systems* (298)
 Band structure studies of actinide systems* (323)
 Study of mass transfer by liquid sodium in closed loop systems. Sodium technology development programme* (324)
 Phase diagrams of BiCl_3 - CuCl-TeCl_4 and CuCl-FeCl_3 - TeCl_4 systems* (337)
 Studies on some binary solvent systems for the extraction of gadolinium* (370)
 Effect of basicity on ferrous oxide activity in complex oxide systems* (372)
 Rare earth metal-metal halide systems. 19. Structural characterization of the reduced holmium chloride $\text{Ho}_5\text{Cl}_{11}$ * (378)
 Electrochemical characterization of nonaqueous systems for secondary battery application. Quarterly report, February--April 1968* (380)
 Physical research on liquid-metal systems. Annual report, July 1975--June 1976* (405)
 Comparative study of systems of heavier rare-earth metals with hydrogen [Gd, Tb, Dy, Lu]* (418)
 FeCl_3 - MeCl systems* (435)
 Phase relations in the alkaline earth fluoride-yttria systems (in three parts). 1. The CaF_2 - Y_2O_3 phase diagram* (468)
 Phase relations in the alkaline earth fluoride-yttria systems (in three parts). 2. The BaF_2 - Y_2O_3 phase diagram* (469)
 Phase diagrams in the pseudobinary systems of Li_2BeF_4 , alkali fluoroborates and alkali fluorides* (473)
 Thermodynamic analysis of melting point diagrams of systems consisting of alkali metal chlorides, thallium chloride and beryllium chloride* (505)
 Calculated method of eutectics of multicomponent systems on base of algorithm and digital computers* (517)
 Projection-thermographic method for study of ternary and ternary mutual systems* (518)
 MoCl_2 -NaCl and MoCl_2 -KCl binary systems* (547)
 Theoretical concepts useful in the calculation or storage of phase diagrams of ionic systems* (552)
 Viscosity of binary of nitrate systems melts* (592)
 Phases of variable composition with LaF_3 structure in MF_2 -(Y, Ln) F_3 systems. 1. SrF_2 -(Y, Ln) F_3 systems (thermal characteristics, formation of berthollides)* (593)
 Thermographic investigation of the PbCl_2 - PbBr_2 2%- PbI_2 2% and PbCl_2 - PbBr_2 2%- PbF_2 2% systems* (602)

TITLE INDEX

Systems

Systems of NbCl_5 - GaCl_3 and TaCl_5 - GaCl_3 * (633)
 Diffusion in liquid metal systems. Final report* (651)
 Interaction in AgCl - CsCl , CsCl - KCl systems and liquidus surface of AgCl - CsCl - KCl ternary system* (659)
 Glass formation in nitrate systems containing rare earth ions* (666)
 Compounds with the structure of ThMn_{12} type in the R--Fe--Al systems* (671)
 Interaction of plutonium trichloride in melts of ternary systems* (682)
 Phase equilibria and thermodynamic data for the ternary UO_2 - UO_2F_2 - $\text{REO}_{1.5}$ (Ho, Tm, Lu) systems* (694)
 Investigation into properties of liquidus curves of binary salt system melts where chemical compounds formation. 2. Systems of monc-monovalent and bi-bivalent cations with common anion* (708)
 Basic research in crystalline and noncrystalline ceramic systems. Annual report, May 1, 1975--April 1, 1976* (728)
 Molten-salt systems* (729)

T

R_3T_2 compounds (R = rare earth or Y; T = Rh, Pd, Pt) with the rhombohedral Er_3Ni_2 structure type* (365)
 P--t diagram of the system uranium--nitrogen* (373)

Ta

Fusibility in $\text{K}_2\text{Nb}(\text{Ta})\text{F}_7$ -LiF-NaF systems* (298)

tabulated

Approximation of tabulated thermodynamic data for purposes of optimization of two-phase liquid-metal power cycles* (343)

 TaCl

Systems of NbCl_5 - GaCl_3 and TaCl_5 - GaCl_3 * (633)
 IR spectra of the NbCl_5 , TaCl_5 , NbBr_5 and NbOCl_3 vapors* (707)

 TaF

Meltability in system of K_2TaF_7 --NaF--NaCl--KCl* (299)

tantalum

Electrode potentials of tantalum chlorides in a molten NaCl-KCl eutectic* (449)
 Electrochemistry of niobium and tantalum, (5). Anodic dissolution of tantalum and the average valence of tantalum ions in molten KCl* (450)
 Electrochemistry of niobium and tantalum. VI. Electrode potentials of tantalum chlorides in molten KCl* (451)
 Thermographic study of iron, molybdenum, tantalum and tungsten high chlorides interaction* (555)
 Chemistry of elements niobium and tantalum. 87. Studies on the thermal formation and decomposition of the complexes $[\text{NbCl}_{12}]^{\text{sup } 2+}$, $[\text{NbCl}_6\text{I}_8]^{\text{sup } 3+}$, and $[\text{TaCl}_6\text{X}_{12}]^{\text{sup } (n+)}$ (X=Cl, Br, I)* (561)

task

Report of task force on alternate structural materials for Liquid Metal Fast Breeder Reactors* (309)

TITLE INDEX

Task

CODATA recommended key values for thermodynamics, 1976. Report of the CODATA Task Group on key values for thermodynamics, 1976* (717)

Research on electrochemical parameters for nonaqueous batteries: task I. Quarterly progress report No. 3* (727)

Tb

Comparative study of systems of heavier rare-earth metals with hydrogen [Gd, Tb, Dy, Lu]* (418)

Structure of $4f^{10}6s$ and $4f^{10}8s$ levels and Eu^{3+} and Tb^{3+} configurations in short-range ultra-violet* (589)

Gd, Tb, Dy, Ho, Er solubility in chromium* (610)

Te

Equilibrium between Li-K-Te alloys and LiCl-KCl molten eutectic mixture* (92)

On interaction in U--S--Te system* (162)

Ga-Pi-Te system* (544)

GaTe-Bi₂Te₃ system* (545)

technetium

Some constitutional studies on uranium-carbon- and plutonium-carbon-rhenium and technetium systems* (227)

Some constitutional studies on uranium--carbon and plutonium--carbon--rhenium and technetium systems* (228)

technical

Lithium anode limited-cycle battery investigation. Quarterly technical progress report No. 5* (37)

Lithium anode limited-cycle battery investigation. Quarterly technical progress report No. 3* (38)

Vibrational spectra of transition metal hexafluoride crystals. I. Orthorhombic MoF₆, WF₆, UF₆ neat crystals. Technical report* (51)

Vibrational spectra of transition metal hexafluoride crystals.

III. Exciton band structures of MoF₆, WF₆, and UF₆. Technical report* (52)

Interactions in inorganic molecular crystals - electronic spectra of ReF₆ pure and mixed crystals. Technical report, 1

Jul--31 Dec 1975* (53)

Environmental sensitivity of structural metals: liquid metal embrittlement. Annual technical progress report No. 2, 25 Jun 1970--31 May 1971* (95)

Lithium--silver chloride secondary battery investigation.

Quarterly technical progress report No. 2* (111)

Environmental sensitivity of structural metals: liquid metal embrittlement. Annual technical progress report No. 4, 1 Jun 1972--31 May 1973* (211)

Environmental sensitivity of structural metals: liquid metal embrittlement. Annual technical progress report No. 2, 25 Jun 1970--31 May 1971* (212)

Nonaqueous electrolyte systems: ionic transport in nonaqueous media. Interim technical report No. 7* (246)

Lithium--nickel fluoride secondary battery investigation.

Quarterly technical progress report No. 1, 27 Apr--27 Jul 1966* (381)

KKR analysis of the Anderson-McMillan theory of electronic structure of liquid metals. Technical report* (485)

Technical experiences in an engineering-scale facility for uranium fluorination studies* (699)

TITLE INDEX

technology

Structural mechanics in reactor technology. Vol. 1* (19)
 Proceedings of the international conference on liquid technology in energy production* (56)
 Proceedings of the international conference on liquid technology in energy production* (103)
 Study of mass transfer by liquid sodium in closed loop systems. Sodium technology development programme* (324)
 Advances in nuclear science and technology. Vol. 9* (448)
 High temperature technology* (482)
 Radiation effects and tritium technology for fusion reactors. Vol. IV* (529)
 Proceedings of the international conference on liquid technology in energy production* (590)
 Structural mechanics in reactor technology. Vol. 2* (601)
 Influence of the technology of the alloy fabrication on its properties in a liquid state* (643)
 Molten salt chemistry : Part IV - Solubility behaviour of PuF_3 in fluoric acid salts of interest in molten salt reactor technology* (655)
 Radiation effects and tritium technology for fusion reactors* (663)

TeCl

Phase diagrams of BiCl_3 - CuCl - TeCl_4 and CuCl - FeCl_3 - TeCl_4 systems* (337)
 BiCl_3 - GeCl_4 - TeCl_4 system* (553)

Teilinstitut

Progress report of the Teilinstitut Nukleare Festkoerperphysik. (1.6.1974--31.5.1975)* (567)

tellurium

Electrochemical behavior of tellurium (4) in oxychloride melts* (221)
 On interaction of tellurium tetrachloride with copper (1), mercury (2), bismuth and iron (3) chlorides* (336)

temperature

Density and temperature dependence of the surface tension of molten bismuth, lead, and bismuth-lead alloys* (1)
 Comparison of the thermodynamic properties and high temperature chemical behavior of lanthanide and actinide oxides* (2)
 Variation of the solid--solid--nickel--thoria interfacial free energy with temperature* (5)
 Fabrication and high temperature thermodynamic and transport properties of PuN * (9)
 Stable inorganic matrix materials for high temperature batteries. Final report No. 1, 2 June 1967--2 February 1968* (23)
 Temperature effect on vibrational spectra of the impurity molecular ions in alkali halides* (46)
 Calculation of the oxygen potential of $(\text{U,Pu})\text{O}_{2-x}$ with defect models and measured oxygen potential data in the temperature range from 1,000 to 1,700 K* (74)
 Temperature dependence of vapor pressure of rare earth halides and their alloys* (99)
 High-temperature microcalorimetric measurements of partial molar enthalpy of solution, $\Delta H(\text{O}_{2\text{g}})$, in plutonium + oxygen* (107)

TITLE INDEX

temperature

- Determination of oxygen potentials of hyperstoichiometric U--Pu dioxides in the temperature range 1,500-1,850 K* (109)
- Determination of oxygen potentials of hyperstoichiometric U-Pu dioxides in the temperature range 1500-1850 K* (110)
- Some considerations in establishing thermodynamic activities from high temperature EMF measurements* (163)
- High-temperature unit to the IKS-21 spectrometer for the detection of molten salts absorption spectra* (208)
- Fine definition of ir spectra from high temperature interactions of U + O₂. Part II. Final report, 1 July 1974--30 June 1975* (218)
- Negative temperature coefficients of electrical resistivity: the divalent liquid metals Eu, Yb and Ba* (224)
- High temperature thermal expansion of ThO₂, MgO and Y₂O₃ by X-ray diffraction* (242)
- The calculation of the consequences of sodium/fuel interactions with allowance for temperature gradients in the sodium* (272)
- Effect of various additions on the temperature of structural transformation in liquid iron* (313)
- Preparation and thermochemical stability of uranium-zirconium-carbonitrides. A contribution to nuclear fuels for advanced high-temperature reactors* (333)
- High-temperature calorimeter for determination of reaction enthalpy* (364)
- Oxygen redistribution in UO₂ due to a temperature gradient* (401)
- Effect of temperature on the fine structure of Eu²⁺ in KCl and KI* (432)
- Periodically variable wall temperature effect on heat transfer in liquid metals* (437)
- High temperature chemistry of ceramic nuclear fuels with emphasis on nonstoichiometry* (448)
- High temperature technology* (482)
- Half-life of ²⁴²Pu from precise low-temperature heat capacity measurements* (487)
- Theory for the derivation of a single hard-sphere potential describing the overall molecular interactions in binary liquid metal alloys as a function of temperature and composition* (525)
- Some trends in constitutive equation model development for high-temperature behavior of fast-reactor structural alloys* (527)
- Thermodynamics of electrolytes. 8. High-temperature properties, including enthalpy and heat capacity, with application to sodium chloride* (586)
- High-temperature heat contents and entropies of zirconium fluoride and zirconium sulfate* (591)
- Mass spectrometric determination of dissociation energies of high-temperature molecules: ScS, YS, LaS, ZrS, UO* (597)
- Experimental study of the temperature dependence on energy exchanges in liquid metals* (621)
- Electrochemical study of high-temperature stability of compounds between the rare earths and copper oxide* (634)
- Investigation of the solubility and diffusion of Fe atoms in Cu at high temperature using molten salt electrochemistry* (693)

TITLE INDEX

temperature

Summary of U.S. LMFBR programs on high temperature structural design and associated materials testing* (731)

temperatures

Preparation of ^{147}Pm metal and the determination of the melting point and phase transformation temperatures* (18)

X-ray study of the plutonium--oxygen phase diagram at ambient temperatures up to 1100°C * (62)

Uranium carbide: thermal diffusivity, thermal conductivity and spectral emissivity at high temperatures* (121)

Thermal diffusivity, thermal conductivity and spectral emissivity of uranium dicarbide at high temperatures* (132)

Simple Raman cell and furnace usable at temperatures higher than 1000°C for corrosive melts* (204)

Thermodynamic functions and vapor pressures of uranium and plutonium oxides at high temperatures* (220)

Thermophysical and electrophysical properties of uranium hexafluoride at temperatures of $(1-11) \times 10^3$ degreeK and pressures of 0.1--100 atm* (302)

Investigation of calcite solubility in K_2CO_3 and Na_2CO_3 solutions at temperatures of 200 to 350°C * (396)

Thermal decomposition of intermetallic compounds of platinum with alkaline earth and lanthanide metals at high temperatures* (399)

Behavior of the AB-type compounds at high pressures and high temperatures* (421)

Contribution to the establishment of the phase diagram for the U--O--C system by means of x-ray diffractometry at high temperatures and under controlled pressure* (503)

Basic mechanisms providing oxidation resistance in structural metals at high temperatures. II. The defect structure of nonstoichiometric rutile (TiO_2) and nonstoichiometric rutile containing 1 to 10 mole % $\text{Cb}_{2/3}\text{O}_6$. Final report 21 Dec 1973-20 Dec 1974* (542)

Anomalous thermal property behaviour of uranium at low temperatures* (560)

tendencies

Comparison of the tendencies of complex formation of various metal ions in fused LiF-NaF-KF (eut.) and in fused LiCl-KCl (eut.)* (314)

Tensimetric

Tensimetric study of pressure and composition of saturated vapor in TlCl-InCl_3 system* (93)

tension

Density and temperature dependence of the surface tension of molten bismuth, lead, and bismuth-lead alloys* (1)

Density, surface tension, and viscosity of uranium trichloride-sodium chloride melting* (141)

Density, surface tension and viscosity of melts of uranium trichloride-lithium chloride* (142)

Thermodynamic perturbation theory for the surface tension and ion density profile of a liquid metal* (167)

Determination of surface tension of a metal with high vapor pressure* (472)

TITLE INDEX

TeO

Vibrational fine structure in the uranium emission of $\text{Mg}_{\text{sub } 3}\text{TeO}_{\text{sub } 6}$ and $\text{Y}_{\text{sub } 6}\text{WO}_{\text{sub } 12}$ * (59)

terbium

Formation enthalpy of gadolinium, terbium and dysprosium trifluorides* (603)

Heats of formation of holmium and terbium trichlorides* (608)

Terminal

Terminal solid solubility of hydrogen in the zirconium-2.5 weight % niobium alloy* (587)

terms

Structure of $\text{Nd}^{\text{sup } 3+}$ terms in nitrate glasses* (664)

ternary

Study of $\text{KBe}_{\text{sub } 2}\text{F}_{\text{sub } 5}-\text{DyF}_{\text{sub } 3}$ cross-section of $\text{KF}-\text{BeF}_{\text{sub } 2}-\text{DyF}_{\text{sub } 3}$ ternary system and some properties of this cross section glasses* (21)

Ternary $\text{UO}_{\text{sub } 2}-\text{UO}_{\text{sub } 3}-\text{EuO}_{\text{sub } 1}$ system and investigation of $\text{Eu(II)}-\text{actinide(IV)}-\text{perovskites}$ * (50)

Ternary systems from lithium, sodium, strontium and barium chlorides* (86)

Phase diagram for the ternary system $\text{CaCl}_{\text{sub } 2}-\text{KCl}-\text{CaCrO}_{\text{sub } 4}$ * (117)

Thermographic investigation into ternary systems containing sodium and potassium chlorides, uranium trichloride and thorium tetrafluoride* (140)

Phase diagrams of ternary systems containing sodium and potassium chlorides, thorium tetrachloride and plutonium trichloride* (148)

Ternary system of potassium, cesium and calcium nitrates* (254)

Phase diagrams of $\text{LiCl}, \text{CsCl}, \text{CaCl}$ and $\text{LiBr}, \text{CsBr}, \text{CaBr}$ ternary systems* (256)

Ternary mutual system of chlorides and iodides of rubidium and cadmium* (257)

$\text{LiBr}, \text{RbBr}, \text{CsBr}$ ternary system* (268)

Phase diagram of the ternary system of lithium-, rubidium-, and cesium nitrates* (377)

Phase equilibria in the $\text{PbO}-\text{GeO}_{\text{sub } 2}-\text{V}_{\text{sub } 2}\text{O}_{\text{sub } 5}$ ternary system* (394)

Evaporation behaviour of the ternary uranium plutonium carbides* (479)

Evaporation behaviour of the ternary uranium plutonium carbides* (480)

Evaporation behaviour of the ternary uranium plutonium carbides* (481)

Projection-thermographic method for study of ternary and ternary mutual systems* (518)

$\text{ZrCl}_{\text{sub } 4}-\text{TiCl}_{\text{sub } 3}-\text{NaCl}$ ternary system* (548)

Viscosity of melts of $\text{KCl}-\text{KF}-\text{K}_{\text{sub } 2}\text{ZrF}_{\text{sub } 6}$ ternary system* (576)

Thermophysical properties of the ternary liquid-metal system tin--lead--cadmium* (605)

Thermodynamic properties of the ternary Sn-Pb-Cd liquid metal system* (606)

Interaction in $\text{AgCl}-\text{CsCl}, \text{CsCl}-\text{KCl}$ systems and liquidus surface of $\text{AgCl}-\text{CsCl}-\text{KCl}$ ternary system* (659)

TITLE INDEX

ternary

- Interaction of plutonium trichloride in melts of ternary systems* (682)
- Phase equilibria and thermodynamic data for the ternary $\text{UO}_2\text{-UO}_3\text{-REO}_{1.5}$ (Ho, Tm, Lu) systems* (694)
- Cesium, calcium and barium chlorides ternary system* (709)

tetraborides

- Electron structure of rare-earth metal tetraborides* (215)

tetrabromides

- IR spectroscopic study of dipotassium and dicesium uranyl tetrabromides* (483)

tetrachloride

- Interaction of uranyl chloride with zirconium tetrachloride in the melt of sodium chloride and potassium chloride* (27)
- Some chemical reactions of uranium hexachloride and the chlorination of uranium tetrachloride* (81)
- Phase diagrams of ternary systems containing sodium and potassium chlorides, thorium tetrachloride and plutonium trichloride* (148)
- On interaction of tellurium tetrachloride with copper (1), mercury (2), bismuth and iron (3) chlorides* (336)

tetrachlorides

- Properties of zirconium and hafnium tetrachlorides and their saturated solutions* (136)
- Interaction of uranium- and thorium tetrachlorides with sodium- and potassium chlorides* (144)
- Diffuse reflectance spectra of protactinium(IV), uranium(IV) and neptunium(IV) tetrachlorides and tetraformates* (562)

tetrachloroferrates

- Pressure and composition of vapors over alkali metal tetrachloroferrates in chlorine atmosphere* (433)

Tetrad

- Tetrad effect in the extraction of the lanthanides with neutral oxygen-containing compounds* (426)

tetrafluoride

- Thermographic investigation into ternary systems containing sodium and potassium chlorides, uranium trichloride and thorium tetrafluoride* (140)
- Reactivity of transition metal fluorides. IX. Protactinium pentafluoride and tetrafluoride* (465)

tetrafluorides

- Stability of rare earth tetrafluorides* (312)

tetrafluoroborate

- Interaction of potassium tetrafluoroborate with fluorides of potassium, lanthanum, neodymium in melt* (530)

tetraformates

- Diffuse reflectance spectra of protactinium(IV), uranium(IV) and neptunium(IV) tetrachlorides and tetraformates* (562)

tetranuclear

- Preparational, structural and vibrational study of $\text{M}_2\text{U}_2\text{O}_5\text{Cl}_4$, $2\text{H}_2\text{O}$ (M = Rb, Cs). Detection of a tetranuclear anion $[(\text{UO}_2)_4\text{O}_{12}\text{Cl}_8(\text{H}_2\text{O})_2]^{4-}$ * (498)

tetraoxide

- Reactions of liquid sodium with transition-metal oxides. VIII. The oxides of chromium-(III), -(IV), and -(VI) and disodium chromium(VI) tetraoxide* (35)

TITLE INDEX

texture

Discrete variation of the structure and texture of liquid metals formed by components with discrete packing* (177)

tballium

Preparation and characterization of higher uranates(VI) of thallium(I)* (278)
 Thermodynamic analysis of melting point diagrams of systems consisting of alkali metal chlorides, thallium chloride and beryllium chloride* (505)
 Investigation into reactions of hexabromotellurates of potassium, cesium and thallium(I) in melt* (556)

ThCl

Phase diagram of the $\text{ThCl}_4\text{-LiCl-PbCl}_2$ system* (143)
 Thermodynamic properties of dilute solutions of ThCl_4 in (Li-K)Cl and (Na-K) eutectics* (407)
 Raman spectra of solid ThCl_4 and fused $\text{ThCl}_4\text{-KCl}$ system* (488)
 Preparation of thorium metal by electrolysis of fused salt $\text{ThCl}_4\text{-LiCl+NaCl+KCl(eut)}$ system* (703)

Theoretical

Theoretical prediction of the thermal conductivity of uranium carbide vapor* (124)
 Theoretical prediction of the thermal conductivity of uranium nitride vapor* (354)
 Eleventh annual seminar on theoretical physics, Johannesburg, 5-9 July 1976* (368)
 Theoretical concepts useful in the calculation or storage of phase diagrams of ionic systems* (552)

Theory

Optical absorption Zeeman spectroscopy of rare earth orthochromites. II. Theory* (122)
 Thermodynamic perturbation theory for the surface tension and ion density profile of a liquid metal* (167)
 Theory of rare-earth alloys* (376)
 KKR analysis of the Anderson-McMillan theory of electronic structure of liquid metals. Technical report* (485)
 Theory for the derivation of a single hard-sphere potential describing the overall molecular interactions in binary liquid metal alloys as a function of temperature and composition* (525)
 Application of molecular dynamics computations to the conformal ionic solution theory: Thermodynamics of binary molten salt mixtures* (550)

Thermal

Thermal analysis of $\text{GaCl}_3\text{-3In} = \text{InCl+Ga}$ system* (4)
 Thermal expansion of PuO_2 * (15)
 Uranium carbide: thermal diffusivity, thermal conductivity and spectral emissivity at high temperatures* (121)
 Theoretical prediction of the thermal conductivity of uranium carbide vapor* (124)
 Thermal diffusivity, thermal conductivity and spectral emissivity of uranium dicarbide at high temperatures* (132)
 Thermal expansion of molten materials by the gamma attenuation technique* (152)
 High temperature thermal expansion of ThO_2 , MgO and Y_2O_3 by X-ray diffraction* (242)

TITLE INDEX

Thermal

- Thermal investigation of $\text{NdCl}_3\text{-AlCl}_3\text{-KCl}$ and $\text{NdCl}_3\text{-AlCl}_3\text{-CsCl}$ system* (289)
- The spectral properties of uranium hexafluoride and its thermal decomposition products* (345)
- Theoretical prediction of the thermal conductivity of uranium nitride vapor* (354)
- Thermal decomposition of solid metal oxides* (395)
- Thermal decomposition of intermetallic compounds of platinum with alkaline earth and lanthanide metals at high temperatures* (399)
- Thermal conductivity and diffusivity of $\text{U}_{40}\text{Pu}_{60}$ from 250 to 415 K* (459)
- Evaporation behavior and high-temperature thermal analysis of substoichiometric plutonium oxide in the composition range from $\text{O/Pu} = 1.51$ to 2.00* (482)
- Thermal lattice dilatation of some transition metal compounds* (508)
- Thermal analysis of the system of NaF , CaF_2 , Na_2MoO_4 , CaMoO_4 * (519)
- Anomalous thermal property behaviour of uranium at low temperatures* (560)
- Chemistry of elements niobium and tantalum. 87. Studies on the thermal formation and decomposition of the complexes $[\text{NbCl}_{12}]^{2+}$, $[\text{NbI}_8]^{3+}$, and $[\text{TaX}_{12}]^{n+}$ ($\text{X}=\text{Cl}, \text{Br}, \text{I}$)* (561)
- Thermal expansion of metals over the entire liquid range* (573)
- Phases of variable composition with LaF_3 structure in $\text{MF}_2\text{-(Y, Ln)F}_3$ systems. 1. $\text{SrF}_2\text{-(Y, Ln)F}_3$ systems (thermal characteristics, formation of berthollides)* (593)
- Thermal properties of nuclear fuel ceramics* (613)
- Some thermal properties of beryllium fluoride from 800 to 1,200 K* (617)
- Degree of thermal dissociation of MCaCl_3 ($\text{M}=\text{K}, \text{Rb}, \text{Cs}$) complexes during melting* (624)
- Thermal analysis of NaF , NaCl , NaMoO_4 system* (636)
- Solubility of nitrogen in liquid lithium and thermal decomposition of solid Li_3N * (702)

thermo

- Fuel-clad interaction. C-E thermo-structural fuel evaluation methods* (719)

thermoanalytical

- Studies of non-stoichiometric oxides by thermoanalytical methods* (626)

Thermochemical

1. Thermochemical properties* (7)
- Preparation and thermochemical stability of uranium-zirconium-carbonitrides.* (332)
- Preparation and thermochemical stability of uranium-zirconium-carbonitrides. A contribution to nuclear fuels for advanced high-temperature reactors* (333)
- Preparation and thermochemical stability of uranium-zirconium-carbonitrides* (334)

thermochemistry

- Advances on thermochemistry of rare earths alloys* (65)

TITLE INDEX

Thermochemistry

- Thermochemistry of polyhalides. Pt.1. Caesium and rubidium dichloroiodates (I)* (178)
- Thermochemistry of gaseous UF_5 and UF_4 * (238)
- Thermochemistry of gaseous metal oxides. Final report, 1 Apr 1973--31 Dec 1975* (239)
- Thermochemistry of gaseous compounds of metals. Annual summary report 5 Dec 74--4 Dec 75* (240)
- Thermochemistry of gaseous compounds of metals. Annual summary report, 5 Dec 1975--4 Dec 1976* (241)

thermoelectric

- Ionic-thermoelectric study of the dipole relaxation and equilibrium in Gd-doped SrF_2 * (410)

Thermoelectric

- Thermoelectric measurements in liquid alloys and electron transport heat calculations in the case of almost-free electrons* (33)

thermodynamic

- Comparison of the thermodynamic properties and high temperature chemical behavior of lanthanide and actinide oxides* (2)
- Thermodynamic properties of the liquid La--Mg alloys* (3)
- Thermodynamic properties of diluted solutions of $MnCl_2$ in alkali metal chlorides* (6)
- Fabrication and high temperature thermodynamic and transport properties of FuN * (9)
- Thermodynamic properties of some rare earth carbonates and yttrium carbonates* (36)
- Investigation into thermodynamic properties of Ce-Mg melts* (39)
- Thermodynamic assessment of the conversion of plutonium dioxide to plutonium monocarbide* (54)
- Isotope effect on the thermodynamic quantities of gaseous uranium hexafluoride* (78)
- Investigation into thermodynamic properties of $FeCl_2$ - $CsCl$ system melts* (96)
- Thermodynamic properties of $(U,Pu)O_{2-x}$ described by a cluster model* (133)
- Calculation of thermodynamic activity and liquidus lines of components in binary systems of plutonium trichloride with alkali and alkaline earth metal chlorides* (147)
- Contribution to the thermodynamic study of uranium and nickel mixed oxides* (149)
- Some considerations in establishing thermodynamic activities from high temperature EMF measurements* (163)
- Thermodynamic perturbation theory for the surface tension and ion density profile of a liquid metal* (167)
- Thermodynamic properties of Cr, Mn, Co and Ni dichlorides from emf measurements on cells with solid electrolytes* (200)
- Thermodynamic functions and vapor pressures of uranium and plutonium oxides at high temperatures* (220)
- Lattice-dynamical calculation of the thermodynamic properties of NaCl, KCl and $RbCl$ * (231)
- Thermodynamic study of the U-O-C system by measurement of the carbon monoxide pressures at equilibrium* (235)
- Thermodynamic properties of solid and liquid metals and ceramics* (243)

TITLE INDEX

Thermodynamic

- Thermodynamic properties of the U-Sn-alloys riched with tin* (284)
- Thermodynamic study of the binary system copper-indium. I. Calcrimetric study* (290)
- Thermodynamic study of the binary system copper--indium. II. Potenticmetric study* (291)
- Thermodynamic state, specific heat, and enthalpy function of saturated UO_2 vapor between 3000 K and 5000 K* (297)
- Investigation of formation thermodynamic of rare earth aluminates with perovskite structure* (326)
- Cs-Hg amalgam thermodynamic properties* (339)
- Thermodynamic properties of Cd-Zn-Hg liquid alloys* (340)
- Approximation of tabulated thermodynamic data for purposes of optimization of two-phase liquid-metal power cycles* (343)
- Sodium uranium(V) trioxide, NaUO_3 : heat capacity and thermodynamic properties from 5 to 350 K* (383)
- Sodium uranium(V) trioxide, NaUO_3 : heat capacity and thermodynamic properties from 5 to 350 K* (384)
- Thermodynamic properties of diluted sodium solutions in liquid indium* (388)
- Thermodynamic similarity and universal equations of state of hexafluorides* (397)
- Some thermodynamic properties of dilute solutions of actinide chlorides in (Li-K)Cl and in (Na-K)Cl eutectics* (406)
- Thermodynamic properties of dilute solutions of ThCl_4 in (Li-K)Cl and (Na-K) eutectics* (407)
- Thermodynamic study of the neptunium-hydrogen system* (428)
- Calculation of thermodynamic activity of components in strontium liquid alloys* (446)
- Thermodynamic analysis of melting point diagrams of systems consisting of alkali metal chlorides, thallium chloride and beryllium chloride* (505)
- Thermodynamic studies of gallium--indium liquid alloys by solid state electrochemistry with oxide electrolytes* (513)
- Thermodynamic properties of liquid thorium-gallium alloys* (522)
- Thermodynamic investigation of liquid alloys in the system Cd-In-Tl* (523)
- Contribution to the problem of the thermodynamic properties and the structure of liquid Cd-In alloys* (524)
- Thermodynamic properties of the ternary Sn-Pb-Cd liquid metal system* (606)
- Application of the hard-sphere model to thermodynamic properties of liquid 3d transition metals* (615)
- Rare earth and actinide oxides. Thermodynamic and electron microscopy studies* (625)
- Thermodynamic properties of sulphur in liquid antimony* (642)
- Thermodynamic activity of carbon in molybdenum-containing uranium carbide* (644)
- Thermodynamic activity of carbon in molybdenum-containing uranium carbide* (650)
- Thermodynamic properties of NbO , $\text{Fe}_{0.95}\text{O}$, CoO , NiO , Cu_2O and CuO oxides as comparative electrodes in electromotive forces method* (658)

TITLE INDEX

Thermodynamic

Thermodynamic properties of solutions of hydrogen isotopes in metals and alloys of interest to fusion reactor technology* (663)
 Phase equilibria and thermodynamic data for the ternary $\text{UO}_2\text{-UO}_2\text{F}_2\text{-REO}$ (1.5) (Ho, Tm, Lu) systems* (694)
 Electronic states densities from structure and thermodynamic data for liquid transition metals* (705)

Thermodynamics

Thermodynamics of the conversion of plutonium dioxide to plutonium monocarbide* (55)
 Thermodynamics and kinetics of the formation of rare earth intermetallics* (137)
 Gases and carbon in metals (thermodynamics, kinetics, and properties). I. Alkali metals, alkaline earth metals, light metals (Li, Na, K, Rb, Cs; Ca, Sr, Ba; Be, Mg, Al)* (192)
 Thermodynamics of reactions of rare earth oxides with niobium oxide* (206)
 Investigation of solubility and thermodynamics of dissolution of salts in mixed solvents* (347)
 Monte Carlo analysis of diffusion and thermodynamics in UO_2 * (441)
 Thermodynamics of non-stoichiometry. A model for UC_x * (442)
 Application of molecular dynamics computations to the conformal ionic solution theory: Thermodynamics of binary molten salt mixtures* (550)
 Thermodynamics of solid solutions CsCl-RbCl and CsBr-RbBr at 25°C* (570)
 Thermodynamics of electrolytes. 8. High-temperature properties, including enthalpy and heat capacity, with application to sodium chloride* (586)
 Thermodynamics of the lanthanide halides. II. Heat capacities and Schottky anomalies of SmCl_3 , EuCl_3 and GdCl_3 from 50 to 350 K* (594)
 Phase behavior and thermodynamics of U-Mo-C system* (646)
 Phase behavior and thermodynamics of U-Mo-C system* (647)
 Deoxidation thermodynamics of cerium and lanthanum in liquid iron* (654)
 CODATA recommended key values for thermodynamics, 1976. Report of the CODATA Task Group on key values for thermodynamics, 1976* (717)

Thermographic

Thermographic investigation into ternary systems containing sodium and potassium chlorides, uranium trichloride and thorium tetrafluoride* (140)
 Projection-thermographic method for study of ternary and ternary mutual systems* (518)
 Thermographic study of iron, molybdenum, tantalum and tungsten high chlorides interaction* (555)
 Thermographic investigation of the $\text{PbCl}_2\text{-PbBr}_2$, $\text{PbCl}_2\text{-PbI}_2$ and $\text{PbCl}_2\text{-PbBr}_2\text{-PbF}_2$ systems* (602)
 Thermographic determination of formation heats of lower zirconium chlorides and bromides* (635)

Thermogravimetric

Thermogravimetric investigation of interaction of some rare earth oxides with ammonium bifluoride* (427)

TITLE INDEX

Thermophysical

Thermophysical and electrophysical properties of uranium hexafluoride at temperatures of $(1-11) \times 10^3$ degree K and pressures of 0.1--100 atm* (302)

Thermophysical properties of UO_2 fuel materials* (366)

Speed of ultrasound and the thermophysical properties of the liquid metals Sn, Pb, Cd and of their binary alloys Pb:Sn and Pb:Cd* (403)

New apparatus for thermophysical measurements above 2500 K* (574)

Thermophysical properties of the ternary liquid-metal system tin--lead--cadmium* (605)

ThF

Measurement of mass transfer coefficients in a mechanically agitated, nondispersing contactor operating with a molten mixture of $LiF-BiF_3$ and ThF_4 and molten bismuth* (79)

Heat transfer measurements in a forced convection loop with two molten-fluoride salts: $LiF-BiF_3$ and ThF_4-UF_4 and eutectic $NaBF_4-NaF$ * (585)

Third

Third international symposium on fuel cells* (77)

ThMn

Compounds with the structure of $ThMn_{12}$ type in the R--Fe--Al systems* (671)

ThO

Electronic conductivity and polarization effects of solid ThO_2 - Y_2O_3 electrolytes and of oxygen containing Va-metal electrodes* (193)

High temperature thermal expansion of ThO_2 , MgO and Y_2O_3 by X-ray diffraction* (242)

Quantitative determination of UO_2 and ThO_2 in powdery mixtures by infrared spectrometry* (431)

Measurement of the cell voltage of ThO_2 - Y_2O_3 -solid electrolyte cells in the range of mixed conduction* (535)

thoria

Variation of the solid--solid--nickel--thoria interfacial free energy with temperature* (5)

Electrical conduction in CaO-doped thoria electrolytes* (389)

Complex impedance of electrochemical cells based on yttria doped thoria* (568)

Zirconia and yttria--thoria ceramics find new uses as solid electrolytes* (595)

High density thoria-silica-metal (III) oxide fibers* (710)

thorium

Specific heat of thorium monocarbide from 2 to 300K* (126)

Thermographic investigation into ternary systems containing sodium and potassium chlorides, uranium trichloride and thorium tetrafluoride* (140)

Interaction of uranium- and thorium tetrachlorides with sodium- and potassium chlorides* (144)

Phase diagrams of ternary systems containing sodium and potassium chlorides, thorium tetrachloride and plutonium trichloride* (148)

Determination of the burn up and differentiation of the fission yields of uranium-thorium-fuels* (214)

TITLE INDEX

thorium

Structure of the thorium-(zirconium, niobium, ruthenium, rhodium)-carbon systems* (245)
 Pulse NMR in solids: chemical shift, lead fluoride, and thorium hydride* (363)
 Determination of thorium in rare earths chloride: coprecipitation with cerium (IV) iodate and spectrophotometric determination with thoron* (447)
 Interaction of thorium chloride with calcium oxide* (496)
 Thermodynamic properties of liquid thorium-gallium alloys* (522)
 Investigations on chemical properties of thorium oxide solid electrolyte ceramics* (652)
 Physical properties of thorium fluoride* (656)
 Thorium: physico-chemical properties of its compounds and alloys* (679)
 Preparation of thorium metal by electrolysis of fused salt $\text{ThCl}_4\text{-LiCl+NaCl+KCl}$ (eut) system* (703)
 Thorium utilization program. Quarterly progress report for the period ending May 31, 1977* (722)

thoron

Determination of thorium in rare earths chloride: coprecipitation with cerium (IV) iodate and spectrophotometric determination with thoron* (447)

threevalent

Reaction of americium oxides with threevalent metal oxides* (303)

thulium

Accurate nuclear parameters derived from the Moessbauer spectrum of thulium metal* (696)

ThX

Band structure and chemical bond in ThX compounds* (259)

ThO

Electron structure and bonding in ThO_2 and UO_2 * (223)

TI

Thermodynamic investigation of liquid alloys in the system Cd-In-Ti* (523)

TiCl

$\text{ZrCl}_4\text{-TiCl}_3\text{-NaCl}$ ternary system* (548)

time

Structural or solid state chemistry of rare earth sesquioxides from Goldschmidt to the present time* (438)

tin

Interaction of the lithium-tin alloys with the LiCl-KCl-CsCl melted eutectic mixture* (91)

Formation enthalpy of liquid tin-vanadium alloys* (166)

Thermodynamic properties of the U-Sn-alloys riched with tin* (284)

Thermophysical properties of the ternary liquid-metal system tin--lead-- cadmium* (605)

TiO

Influence of small additions on the preparation and the properties of UO_2 ceramics. Part B. The $\text{UO}_2\text{-TiO}_2$ system* (267)

Basic mechanisms providing oxidation resistance in structural metals at high temperatures. II. The defect structure of nonstoichiometric rutile (TiO_2) and nonstoichiometric rutile containing 1 to 10 mole % CeO_2 . Final report

TITLE INDEX

TiO

21 Dec 1973-20 Dec 1974* (542)

Titanium

Titanium dioxide solubility in melts of lithium- and alkaline earth metal chlorides* (264)

Adhesion properties of melted Al, Ga, In on solid titanium* (327)

Determination of trace impurities of titanium and zirconium in molybdenum and tungsten and their oxygen compounds by X-ray fluorescence spectrometry* (698)

titanoniobates

Physicochemical investigation of rare earth titanoniobates, LnTiNbO_6 * (173)

titanotungstates

Synthesis and properties of rare earth titanotungstates* (114)

TlBr

Electromodulation spectroscopy of sc and fcc phase TlCl and TlBr* (411)

TlCl

Tensimetric study of pressure and composition of saturated vapor in TlCl-InCl₃ system* (93)

TlCl-RbCl-CsCl and NaI-TlI-CsI systems* (258)

Electromodulation spectroscopy of sc and fcc phase TlCl and TlBr* (411)

TlI

TlCl-RbCl-CsCl and NaI-TlI-CsI systems* (258)

Tm

Phase equilibria and thermodynamic data for the ternary UO_2 - UO_3 - $\text{REO}_{1.5}$ (Ho, Tm, Lu) systems* (694)

TR

Phase relations in PbWO_4 - $\text{TR}_2(\text{WO}_4)_3$ systems* (168)

trace

Spectrographic determination of trace impurities in reactor grade aluminium* (104)

Determination of trace impurities of titanium and zirconium in molybdenum and tungsten and their oxygen compounds by X-ray fluorescence spectrometry* (698)

tracer

Investigations on the solvent extraction from molten salts by tracer technique* (369)

Investigations of the solvent extraction from molten salts with the use of tracer technique* (371)

transfer

Measurement of mass transfer coefficients in a mechanically agitated, nondispersing contactor operating with a molten mixture of LiF - BeF_2 - ThF_4 and molten bismuth* (79)

Heat transfer bibliography* (158)

Determination and correlation of mass transfer coefficients in a stirred cell* (236)

Study of mass transfer by liquid sodium in closed loop systems. Sodium technology development programme* (324)

Abstracts of papers and communications submitted to the 5. all-union heat and mass transfer conference* (437)

New description of liquid metal heat transfer in closed conduits* (502)

TITLE INDEX

transfer

Energy transfer between $\text{Bi}^{\text{sup } 3+}$ yields $\text{Eu}^{\text{sup } 3+}$, $\text{Bi}^{\text{sup } 3+}$ yields $\text{Sm}^{\text{sup } 3+}$ and $\text{UO}_2^{\text{sup } 2+}$ yields $\text{Eu}^{\text{sup } 3+}$ in oxide glasses* (539)
 Heat transfer measurements in a forced convection loop with two molten-fluoride salts: LiF -- BeF_2 -- ThF_2 -- UF_4 and eutectic NaBF_4 -- NaF * (585)

transfers

Use of oscillator strengths of "supersensitive" ion transfers of lanthanides for calculation of some parameters determining the intensity of their absorption spectra* (511)

transformation

Preparation of ^{147}Pm metal and the determination of the melting point and phase transformation temperatures* (18)
 Effect of various additions on the temperature of structural transformation in liquid iron* (313)

transformations

Atomic structure of melts of complex system with eutectic transformations* (155)

transients

Preliminary assessments of carbide fuel pins during mild overpower transients in LMFBRs* (456)

transition

Reactions of liquid sodium with transition-metal oxides. VIII. The oxides of chromium-(III), -(IV), and -(VI) and disodium chromium(VI) tetraoxide* (35)
 Vibrational spectra of transition metal hexafluoride crystals. I. Orthorhombic MoF_6 , WF_6 , UF_6 neat crystals. Technical report* (51)
 Vibrational spectra of transition metal hexafluoride crystals. III. Exciton band structures of MoF_6 , WF_6 , and UF_6 . Technical report* (52)
 Structure of the biexciton molecule and optical transition selection rules in zincblende and wurtzite materials* (186)
 Entropy change for the lambda-type transition in nonstoichiometric $\text{U}_{40}\text{O}_{(9-y)}$ * (260)
 Phase transition in doped $\text{U}_{40}\text{O}_{(9-y)}$. I. Heat capacity measurement* (261)
 Hydrogen absorption of rare-earth (3d) transition intermetallic compounds* (392)
 Progress in the spectroscopy of rare earth and transition metal ions* (412)
 Photoelectron spectroscopic studies of some transition metals and alloys* (417)
 Densities of electronic states in liquid transition metals* (422)
 Heat of formation of solid and liquid alloys of transition metals* (425)
 Reactivity of transition metal fluorides. IX. Protactinium pentafluoride and tetrafluoride* (465)
 Thermal lattice dilatation of some transition metal compounds* (508)
 Preparation and structure of transition metal fluorides $\text{U}^{\text{sup } (\text{II})}\text{M}^{\text{sup } (\text{IV})}\text{F}_6$ [$\text{U}^{\text{sup } (\text{II})}$: Cr-Zn; $\text{M}^{\text{sup } (\text{IV})}$: Hf, Zr]* (596)
 Application of the hard-sphere model to thermodynamic properties of liquid 3d transition metals* (615)
 Structure of liquid transition metals and their alloys* (686)

TITLE INDEX

transition

Structures of 3d-transition metals in the liquid state* (687)
Electronic states densities from structure and thermodynamic data for liquid transition metals* (705)

transitions

Intensities of lanthanide f-f transitions* (492)
Radiative and non-radiative transitions of rare-earth ions in glasses* (538)

transplutonium

All-union conference on chemistry of transplutonium elements (americium, curium, berkelium, californium)* (34)
Transplutonium 1975* (303)
All-union conference on chemistry of transplutonium elements (americium, curium, berkelium, californium)* (307)
All-union conference on chemistry of transplutonium elements (americium, curium, berkelium, californium)* (308)
Transplutonium 1975* (360)
Transplutonium 1975* (683)

transport

Fabrication and high temperature thermodynamic and transport properties of PuN * (9)
Transport of hydrogen in liquid lithium* (10)
Thermodiffusion measurements in liquid alloys and electron transport heat calculations in the case of almost-free electrons* (33)
Physical models of mass transport of iron and nickel in liquid sodium systems* (128)
Structure factor of liquid aluminum and calculation of transport coefficients* (159)
Nonaqueous electrolyte systems: ionic transport in nonaqueous media. Interim technical report No. 7* (246)
Molten-salt electrolytes--transport properties* (277)
Mass transport phenomena in ceramics* (401)
Equation of state and transport properties of uranium and plutonium carbides in the liquid region* (577)
Equation of state and transport properties of uranium and plutonium nitrides in the liquid region* (578)

transuranium

Physical--chemical studies of transuranium elements. Progress report, April 1, 1976--March 31, 1977* (501)
Chemistry of transuranium elements* (730)

trap

Adaptation of the continuous cold trap system of fluidized-bed to the fluoride volatility process* (724)

trends

Some trends in constitutive equation model development for high-temperature behavior of fast-reactor structural alloys* (527)

tribromide

Investigation into interaction of indium tribromide with dibromides of manganese, cobalt and nickel in melt* (153)

trichloride

Optical spectra of the difluoride, dichloride, and trichloride ions in the matrix-isolated $\text{M}^{\text{sup}} + \text{F}^{\text{sub 2}}^{\text{sup}} -$, $\text{M}^{\text{sup}} + \text{Cl}^{\text{sub 2}}^{\text{sup}} -$, and $\text{M}^{\text{sup}} + \text{Cl}^{\text{sub 3}}^{\text{sup}} -$ species* (16)

TITLE INDEX

trichloride

- Interaction of uranium trichloride with alkaline earth metal chlorides* (139)
- Thermographic investigation into ternary systems containing sodium and potassium chlorides, uranium trichloride and thorium tetrafluoride* (140)
- Density, surface tension, and viscosity of uranium trichloride-sodium chloride melting* (141)
- Density, surface tension and viscosity of melts of uranium trichloride-lithium chloride* (142)
- Calculation of thermodynamic activity and liquidus lines of components in binary systems of plutonium trichloride with alkali and alkaline earth metal chlorides* (147)
- Phase diagrams of ternary systems containing sodium and potassium chlorides, thorium tetrachloride and plutonium trichloride* (148)
- Heat of formation of europium sesquioxide and europium trichloride* (607)
- Interaction of plutonium trichloride in melts of ternary systems* (682)

trichlorides

- Heats of formation of holmium and terbium trichlorides* (608)

trifluorides

- Formation enthalpy of di- and trifluorides of samarium* (306)
- Rare earth 4d spectra in rare earth trifluorides* (382)
- Interaction of rubidium and cesium fluorides with lutetium trifluorides* (444)
- Optical properties of rare earth trifluorides* (484)
- Formation enthalpy of gadolinium, terbium and dysprosium trifluorides* (603)

Trino

- Investigations on radioactive fission product correlations: gamma spectrometry measurements on spent fuel assemblies discharged from the Trino Vercellese reactor at the end of the 2nd irradiation cycle* (75)

trioctylammonium

- Investigation into reactions between trioctylammonium nitrates and rare earth nitrates by X-ray electron spectroscopy* (113)

trioxide

- Sodium uranium(V) trioxide, NaUO_3 : heat capacity and thermodynamic properties from 5 to 350 K* (383)
- Sodium uranium(V) trioxide, NaUO_3 : heat capacity and thermodynamic properties from 5 to 350 K* (384)

tritium

- Radiation effects and tritium technology for fusion reactors. Vol. IV* (529)
- Tritium sorption in lithium--bismuth and lithium--aluminum alloys* (614)
- Radiation effects and tritium technology for fusion reactors* (663)

trudy

- Nauchnye trudy GIREDMETA. Vol. 52* (115)

tubesheet

- Metallurgical characterization of prototypic tubesheet forging* (471)

TITLE INDEX

tungstate

- Raman and i.r.-emission studies of some tungstate- and molybdate-containing melts* (102)
 Double tungstate $K\text{La}(\text{WO}_4)_4$ * (338)

tungstates

- Binary system of sodium, rubidium and cesium tungstates* (151)
 Double potassium rare earth tungstates* (639)

tungsten

- Molybdenum and tungsten interaction with liquid aluminium* (156)
 Thermographic study of iron, molybdenum, tantalum and tungsten high chlorides interaction* (555)
 Phase equilibrium study on system uranium-plutonium-tungsten-carbon* (645)
 Determination of trace impurities of titanium and zirconium in molybdenum and tungsten and their oxygen compounds by X-ray fluorescence spectrometry* (698)

Twenty

- Twenty years of research in the processing of radioactive fuel and neutralization of radioactive waste at the Nuclear Research Institute (NRI)* (358)

U

- Electronic structure of highly symmetric complexes of lanthanoids and actinoids. Pt. 3. 'Electronic' Raman spectrum of $\text{CsU}(\text{IV})\text{Cl}_6$ * (13)
 Calculation of the oxygen potential of $(\text{U}, \text{Pu})\text{O}_{2+x}$ with defect models and measured oxygen potential data in the temperature range from 1,000 to 1,700 K* (74)
 Determination of oxygen potentials of hyperstoichiometric U--Pu dioxides in the temperature range 1,500-1,850 K* (109)
 Determination of oxygen potentials of hyperstoichiometric U-Pu dioxides in the temperature range 1500-1850 K* (110)
 Chemical separation and spectrographic estimation of rare earths and yttrium in PuO_2 and $(\text{U}, \text{Pu})\text{O}_{2+x}$ * (125)
 Thermodynamic properties of $(\text{U}, \text{Pu})\text{O}_{2+x}$ described by a cluster model* (133)
 Distribution coefficient for U and Pu in the system $(\text{UPu})\text{C} + (\text{UPu})\text{C}_3$ * (134)
 On interaction in U--S--Te system* (162)
 Cs--U--O phase diagram and its application to uranium--plutonium oxide fast reactor fuel pins* (175)
 Fine definition of ir spectra from high temperature interactions of U + O_2 . Part II. Final report, 1 July 1974--30 June 1975* (218)
 Thermodynamic study of the U-O-C system by measurement of the carbon monoxide pressures at equilibrium* (235)
 Entropy change for the lambda-type transition in nonstoichiometric U_4O_{9-y} * (260)
 Phase transition in doped U_4O_{9-y} . I. Heat capacity measurement* (261)
 Lambda-type heat capacity anomalies in U_3O_8 * (262)
 Thermodynamic properties of the U-Sn-alloys riched with tin* (284)
 Chronopotentiometry of U(III) and U(IV) in molten LiCl-KCl eutectic* (292)

TITLE INDEX

U

- Electrochemical behavior of U(III) and U(IV) in molten LiCl-KCl eutectic* (293)
- Redox and chemical properties of U, Np and Pu in molten alkali chlorides* (360)
- Thermal conductivity and diffusivity of $U_{40}O_{60}$ from 250 to 415 K* (459)
- Extension of vapour pressure measurements of nuclear fuels (U,Pu) $O_{2.5}$ and $UO_{2.5}$ to 7000 K for fast reactor safety analysis* (475)
- Vapour pressure measurements to 7,000 K and equation of state of oxide fuels for fast reactor safety analysis. I. Extension of vapour pressure measurements of nuclear fuels (U,Pu) $O_{2.5}$ and $UO_{2.5}$ to 7,000 K* (477)
- Preparational, structural and vibrational study of $M_{2.5}U_{2.5}O_{5.5}Cl_{0.4}$, $2H_{2.5}O$ (M = Rb, Cs). Detection of a tetranuclear anion $[(UO_{2.5})_4O_{10}Cl_{0.8}(H_{2.5}O)_4]^{4-}$ * (498)
- Contribution to the establishment of the phase diagram for the U--O--C system by means of x-ray diffractometry at high temperatures and under controlled pressure* (503)
- Covalent compounds of uranium of the type $U(OTeF_5)_x$ (571)
- U.S.-Japan joint seminar on defects and diffusion in solids* (611)
- Total pressures of uranium- and plutonium-bearing species above the U--Pu--C system* (618)
- Phase behavior and thermodynamics of U--Mo--C system* (646)
- Phase behavior and thermodynamics of U--Mo--C system* (647)
- Phase equilibria in system U--Pu--W--C* (648)
- Parameter for optimizing the synthesis of (U,Pu)C* (676)
- Standard methods for chemical, mass spectrometric, and spectrochemical analysis of nuclear-grade mixed oxides (U,Pu) $O_{2.5}$ * (713)
- Summary of U.S. LMFBR programs on high temperature structural design and associated materials testing* (731)

UC

- Solution studies in the U--UC $_{2.5}$ system* (379)
- Thermodynamics of non-stoichiometry. A model for UC $_x$ * (442)

UC1

- Viscosity of UCl_3 -- $MgCl_2$ binary system melts* (145)
- Viscosity of molten $CaCl_2$ -- UCl_3 mixtures* (146)
- Cryometric studies of the system $CdCl_2$ -- UCl_4 * (199)

UF^{sup}

- Preparation and structure of transition metal fluorides $UF^{sup}(II)M^{sup}(IV)F_6$ [UF^{sup}(II) : Cr-Zn; M^{sup}(IV) : Hf, Zr]* (596)

UF

- Vibrational spectra of transition metal hexafluoride crystals. III. Exciton band structures of MoF_6 , WF_6 , and UF_6 . Technical report* (52)
- Thermochemistry of gaseous UF_5 and UF_4 * (238)
- Predicted infrared spectrum and the structure of the isolated UF_5 molecule* (350)
- Synthesis and characterization of a new uranium(V) compound: $H_3O^{+}UF_6^{sup-}$ * (409)

TITLE INDEX

UF

Assignments in the electronic spectrum of UF_{6*} (413)
Heat transfer measurements in a forced convection loop with two molten-fluoride salts: LiF -- BeF_2 -- ThF_2 -- UF_4 and eutectic NaBF_4 -- NaF * (585)

UF₆

Vibrational spectra of transition metal hexafluoride crystals. I. Orthorhombic MoF_6 , WF_6 , UF_6 neat crystals. Technical report* (51)

ultra

Structure of $4f^{6*}$ and $4f^{8*}$ levels and Eu^{3+} and Tb^{3+} configurations in short-range ultra-violet* (589)

ultrasound

Speed of ultrasound and the thermophysical properties of the liquid metals Sn, Pb, Cd and of their binary alloys Pb:Sn and Pb:Cd* (403)

ultraviolet

Vacuum ultraviolet radiation physics* (382)

Vacuum ultraviolet radiation physics* (484)

UN

Identification of UN in Ar matrices* (219)

Synthesis of UN coatings on uranium* (400)

Decomposition of the UN-CeN solid solution* (581)

und

Vacancies und melting curves of metals at high pressure* (213)

union

All-union conference on chemistry of transplutonium elements (americium, curium, berkelium, californium)* (34)

Summaries of reports of the all-union meeting on solid body chemistry. 3--5 June 1975. III. Synthesis, phase equilibrium and use of solid oxide compounds* (114)

All-union conference on chemistry of transplutonium elements (americium, curium, berkelium, californium)* (307)

All-union conference on chemistry of transplutonium elements (americium, curium, berkelium, californium)* (308)

Abstracts of papers and communications submitted to the 5.

all-union heat and mass transfer conference* (437)

2. all-union conference on crystallochemistry of intermetallic compounds* (549)

Summaries of reports of the all-union meeting on solid body chemistry. 3--5 June 1975. III. Synthesis, phase equilibrium and use of solid oxide compounds* (641)

2. all-union conference on crystallochemistry of intermetallic compounds* (671)

unit

High-temperature unit to the IKS-21 spectrometer for the detection of molten salts absorption spectra* (208)

universal

Thermodynamic similarity and universal equations of state of hexafluorides* (397)

unusual

Mixed valence rare-earth halides and their unusual crystal structures* (29)

UO

Optical absorption spectrum of UO_2 doped LiF * (30)
Ternary UO_2 -- UO_3 -- EuO system and investigation of Eu(II) --actinide(IV)--perovskites* (50)

TITLE INDEX

UC

- Study of the compounds $M\text{UO}_2\text{F}_5$ ($M = \text{K, Rb, Cs, NH}_4$) by i.r. absorption and Raman diffusion spectrophotometry. III. Effect of cation on vibration frequency of the ion $\text{UO}_2\text{F}_5^{3-}$ (127)
- Critical assessment of equation of state data for UO_2 (180)
- Prediction of the inert gas solubilities in stoichiometric molten UC_2 (225)
- Influence of small additions to the preparation and the properties of UO_2 ceramics. Part A. $\text{UO}_2/\text{Gd}_2\text{O}_3$ (266)
- Influence of small additions on the preparation and the properties of UO_2 ceramics. Part B. The $\text{UO}_2\text{-TiO}_2$ system (267)
- Thermodynamic state, specific heat, and enthalpy function of saturated UO_2 vapor between 3000 K and 5000 K (297)
- Thermophysical properties of UO_2 fuel materials (366)
- Oxygen redistribution in UO_2 due to a temperature gradient (401)
- Quantitative determination of UO_2 , and ThO_2 in powdery mixtures by infrared spectrometry (431)
- Monte Carlo analysis of diffusion and thermodynamics in UO_{2+x} (441)
- Vaporization behavior of UO_2 (445)
- Study of the retention of Ba in UO_2 nuclear fuel particles by ZrO_2 as getter (470)
- Extension of vapour pressure measurements of nuclear fuels (U, PuO_2) and UO_2 to 7000 K for fast reactor safety analysis (475)
- Vapour pressure measurements to 7,000 K and equation of state of oxide fuels for fast reactor safety analysis. I. Extension of vapour pressure measurements of nuclear fuels (U, PuO_2) and UO_2 to 7,000 K (477)
- Preparational, structural and vibrational study of $M\text{U}_2\text{O}_5\text{Cl}_4$, $2\text{H}_2\text{O}$ ($M = \text{Rb, Cs}$). Detection of a tetranuclear anion [$(\text{UO}_2)_4\text{O}_{12}\text{Cl}_8(\text{H}_2\text{O})_2$]⁴⁻ (498)
- Energy transfer between Bi^{3+} yields Eu^{3+} , Bi^{3+} yields Sm^{3+} and UO_2 yields Eu^{3+} in oxide glasses (539)
- Mass spectrometric determination of dissociation energies of high-temperature molecules: ScS , YS , LaS , ZrS , UO^* (597)
- Phases of high pressure in $\text{UO}_2\text{-UO}_3\text{-PbO}$ system (641)
- Phase equilibria and thermodynamic data for the ternary $\text{UO}_2\text{-UC}_3\text{-REO}_{1.5}$ (Ho, Tm, Lu) systems (694)
- Measurement of interfacial energies in the $\text{UO}_2\text{-Gd}_2\text{O}_3$ system (697)

UPu

- Distribution coefficient for U and Pu in the system $(\text{UPu})\text{C} + (\text{UPu})_2\text{C}_3$ (134)
- Oxygen potential and vacancy structure of the mixed oxide $(\text{UPu})\text{O}_{2-x}$ (565)

uranates

- Preparation and characterization of higher uranates(VI) of thallium(I) (278)

TITLE INDEX

urania

Knudsen cell--mass spectrometer studies of cesium--urania interactions* (120)

uranium

Infrared spectroscopy of the uranium/oxygen system* (12)
 Vibrational fine structure in the uranium emission of MgTeO_6 and Y_6WO_{12} * (59)
 Method for recovering uranium from residual solution* (70)
 Isotope effect on the thermodynamic quantities of gaseous uranium hexafluoride* (78)
 Some chemical reactions of uranium hexachloride and the chlorination of uranium tetrachloride* (81)
 Segregation and vapour pressure studies on the uranium-plutonium-carbon system* (83)
 Segregation and vapour pressure studies in the uranium--plutonium--carbon system* (84)
 Uranium carbide: thermal diffusivity, thermal conductivity and spectral emissivity at high temperatures* (121)
 Theoretical prediction of the thermal conductivity of uranium carbide vapor* (124)
 Thermal diffusivity, thermal conductivity and spectral emissivity of uranium dicarbide at high temperatures* (132)
 Interaction of uranium trichloride with alkaline earth metal chlorides* (139)
 Thermographic investigation into ternary systems containing sodium and potassium chlorides, uranium trichloride and thorium tetrafluoride* (140)
 Density, surface tension, and viscosity of uranium trichloride-sodium chloride melting* (141)
 Density, surface tension and viscosity of melts of uranium trichloride-lithium chloride* (142)
 Interaction of uranium- and thorium tetrachlorides with sodium- and potassium chlorides* (144)
 Contribution to the thermodynamic study of uranium and nickel mixed oxides* (149)
 Cs--U--O phase diagram and its application to uranium--plutonium oxide fast reactor fuel pins* (175)
 Enthalpy of uranium-plutonium carbide from 298K to the melting point* (179)
 Physical and mechanical properties of some uranium-containing carbides* (209)
 Determination of the burn up and differentiation of the fission yields of uranium-thorium-fuels* (214)
 Thermodynamic functions and vapor pressures of uranium and plutonium oxides at high temperatures* (220)
 Uranium dioxide as a ceramic nuclear fuel* (226)
 Some constitutional studies on uranium-carbon- and plutonium-carbon-rhenium and technetium systems* (227)
 Some constitutional studies on uranium--carbon and plutonium--carbon--rhenium and technetium systems* (228)
 Preparation of uranium-233 with low uranium-232 content* (232)
 Vaporization of uranium-carbon-nitrogen system* (251)
 Calculation of the reduced partition function ratio for uranium hexafluoride* (285)

TITLE INDEX

uranium

- Thermophysical and electrophysical properties of uranium hexafluoride at temperatures of $(1-11) \times 10^3$ degree K and pressures of 0.1--100 atm* (302)
- New oxide of uranium* (325)
- Preparation and thermochemical stability of uranium-zirconium-carbonitrides.* (332)
- Preparation and thermochemical stability of uranium-zirconium-carbonitrides. A contribution to nuclear fuels for advanced high-temperature reactors* (333)
- Preparation and thermochemical stability of uranium-zirconium-carbonitrides* (334)
- The spectral properties of uranium hexafluoride and its thermal decomposition products* (345)
- Theoretical prediction of the thermal conductivity of uranium nitride vapor* (354)
- Effect of oxygen content on the structural characteristics of uranium oxycarbides* (362)
- P--t diagram of the system uranium--nitrogen* (373)
- Sodium uranium(V) trioxide, NaUO_3 : heat capacity and thermodynamic properties from 5 to 350 K* (383)
- Sodium uranium(V) trioxide, NaUO_3 : heat capacity and thermodynamic properties from 5 to 350 K* (384)
- Synthesis of UN coatings on uranium* (400)
- Diffusion of uranium species in several molten chlorides* (408)
- Synthesis and characterization of a new uranium(V) compound: $\text{H}_3\text{O}^+ \text{UF}_6^-$ * (409)
- Effect of fluoride ions on the kinetics of the oxidation of uranium(IV) by cerium(IV)* (424)
- Vapor pressure studies over liquid uranium oxide and uranium plutonium oxide up to 5,000 K* (474)
- Advances in vapour pressure studies over liquid uranium plutonium oxides up to 5000K* (476)
- Vapour pressure studies over liquid uranium plutonium oxides up to 5,000 K* (478)
- Evaporation behaviour of the ternary uranium plutonium carbides* (479)
- Evaporation behaviour of the ternary uranium plutonium carbides* (480)
- Evaporation behaviour of the ternary uranium plutonium carbides* (481)
- Phase analysis on simulated LWR fuel rods and its evaluation with the aid of constitution studies in the system uranium-zirconium-oxygen* (509)
- Phase analysis at simulated LWR fuel rods and their interpretation by constitution investigations in the uranium-zirconium-oxygen system* (510)
- Investigation of liquid-steam equilibrium in system with diluted solutions of metal fluorides in uranium hexafluoride* (526)
- Effective equation-of-state measurements on uranium dioxide* (537)
- Anomalous thermal property behaviour of uranium at low temperatures* (560)
- Diffuse reflectance spectra of protactinium(IV), uranium(IV) and neptunium(IV) tetrachlorides and tetraformates* (562)

TITLE INDEX

uranium

- Covalent compounds of uranium of the type $F_{x/6-x}U(OTeF_5)_{x/6-x}$ * (571)
- Equation of state and transport properties of uranium and plutonium carbides in the liquid region* (577)
- Equation of state and transport properties of uranium and plutonium nitrides in the liquid region* (578)
- Densities of liquid carbides and nitrides of uranium and plutonium* (579)
- Methods for the accountability of uranium dioxide* (600)
- Nonstoichiometry in cubic uranium sesquinitride* (611)
- Total pressures of uranium- and plutonium-bearing species above the U--Pu--O system* (618)
- Enthalpy and heat capacity of liquid plutonium and uranium* (619)
- Enthalpy and heat capacity of liquid plutonium and uranium* (620)
- Spectrophotometric determination of uranium in spent fuels of WWER-2 reactors* (632)
- Thermodynamic activity of carbon in molybdenum-containing uranium carbide* (644)
- Phase equilibrium study on system uranium-plutonium-tungsten-carbon* (645)
- Process for preparing uranium carbonitride* (649)
- Thermodynamic activity of carbon in molybdenum-containing uranium carbide* (650)
- Fabrication of uranium-plutonium mixed carbide* (691)
- Technical experiences in an engineering-scale facility for uranium fluorination studies* (699)
- Description of an engineering-scale facility for uranium fluorination studies* (700)
- Standard methods for chemical, mass spectrometric, and spectrochemical analysis of nuclear-grade uranium dioxide powders and pellets* (711)
- Standard method of test for atom percent fission in uranium and plutonium fuel (mass spectrometric method)* (715)
- Standard method of test for atom percent fission in uranium and plutonium fuel (neodymium-148 method)* (716)
- Analysis of uranium carbides* (726)

uraniummononitride

- Spin structure and elastic properties of uraniummononitride* (368)

uranyl

- Interaction of uranyl chloride with zirconium tetrachloride in the melt of sodium chloride and potassium chloride* (27)
- IR spectroscopic study of dipotassium and dicesium uranyl tetrabromides* (483)

uranyls

- Effects of secondary ligands on the electronic structure of uranyls* (684)

US

- Solution studies in the US-UC₂ system* (379)

usable

- Simple Raman cell and furnace usable at temperatures higher than 1000°C for corrosive melts* (204)

useful

- Theoretical concepts useful in the calculation or storage of phase diagrams of ionic systems* (552)

TITLE INDEX

UO

Electron structure and bonding in ThO_2 and UO_2 * (223)

V

Sodium uranium(V) trioxide, NaUO_3 : heat capacity and thermodynamic properties from 5 to 350 K* (383)

Sodium uranium(V) trioxide, NaUO_3 : heat capacity and thermodynamic properties from 5 to 350 K* (384)

Phase equilibria in the $\text{PbO-GeO}_2\text{-V}_2\text{O}_5$ ternary system* (394)

Synthesis and characterization of a new uranium(V) compound:

$\text{H}_3\text{O}^+ + \text{UF}_6^{\text{sup}} - \text{H}^+$ * (409)

X-ray spectral study of the carbides and nitrides of group IV and V metals* (582)

Va

Electronic conductivity and polarization effects of solid $\text{ThO}_2\text{-Y}_2\text{O}_3$ electrolytes and of oxygen containing Va-metal electrodes* (193)

Vacancies

Vacancies and melting curves of metals at high pressure* (213)

vacancy

Oxygen potential and vacancy structure of the mixed oxide $(\text{UPu})\text{O}_{2-x}$ * (565)

Vacuum

Vacuum ultraviolet radiation physics* (382)

Vacuum ultraviolet radiation physics* (484)

valence

Mixed valence rare-earth halides and their unusual crystal structures* (29)

Statistical mechanics of intermediate-valence rare earth compounds* (279)

Electrochemistry of niobium and tantalum, (5). Anodic dissolution of tantalum and the average valence of tantalum ions in molten KCl * (450)

Energies of valence states of atoms in halides* (499)

XPS valence bands of La, Ce and Gd and their aluminium alloys* (598)

values

CODATA recommended key values for thermodynamics, 1976. Report of the CODATA Task Group on key values for thermodynamics, 1976* (717)

vanadates

Dissolution of corundum in molten vanadates* (317)

vanadium

Formation enthalpy of liquid tin-vanadium alloys* (166)

Influence of intermediate products on chlorination kinetics of vanadium oxides in salt melt* (356)

vapor

Experimental vapor pressure and critical point of sodium* (56)

Investigation of fuel vapor pressure for fast breeder safety analysis* (60)

Calculation of vapor pressures of oxide fuels up to 5000 K in equilibrium and non-equilibrium evaporation* (72)

Calculation of vapor pressures of oxide fuels up to 5,000 K for equilibrium and nonequilibrium evaporation* (73)

TITLE INDEX

vapor

- Tensimetric study of pressure and composition of saturated vapor in TiCl-InCl_3 system* (93)
- Saturation vapor pressure in iron chloride-alkaline earth metal chlorides systems* (97)
- Temperature dependence of vapor pressure of rare earth halides and their alloys* (99)
- Theoretical prediction of the thermal conductivity of uranium carbide vapor* (124)
- Mass spectra and vapor composition of niobium pentafluoride* (216)
- Thermodynamic functions and vapor pressures of uranium and plutonium oxides at high temperatures* (220)
- Molecular vapor composition and mass spectra in KF-ZrF_4 , CsF-ZrF_4 systems* (294)
- Thermodynamic state, specific heat, and enthalpy function of saturated UO_2 vapor between 3000 K and 5000 K* (297)
- Theoretical prediction of the thermal conductivity of uranium nitride vapor* (354)
- Pressure of saturated vapor of rare earth and yttrium bromides* (391)
- Determination of surface tension of a metal with high vapor pressure* (472)
- Vapor pressure studies over liquid uranium oxide and uranium plutonium oxide up to 5,000 K* (474)
- Viscosities of cesium vapor to 1,620 K and of liquid gallium to 1,800 K* (622)
- Viscosities of cesium vapor to 1620 K and of liquid gallium to 1800 K* (681)

Vaporization

- Vaporization of uranium-carbon-nitrogen system* (251)
- Vaporization behavior of UO_2 * (445)

vapors

- Pressure and composition of vapors over alkali metal tetrachloroferrates in chlorine atmosphere* (433)
- Investigation into absorption infrared spectra of vapors under ScBr_3 , YCl_3 , LaCl_3 , GdCl_3 , ErCl_3 , LuCl_3 by means of isolation method in rare gas matrix* (497)
- IR spectra of the NbCl_5 , TaCl_5 , NbBr_5 and NbOCl_3 vapors* (707)

vapour

- Segregation and vapour pressure studies on the uranium-plutonium-carbon system* (83)
- Segregation and vapour pressure studies in the uranium-plutonium-carbon system* (84)
- Extension of vapour pressure measurements of nuclear fuels (U,Pu) O_2 to 7000 K for fast reactor safety analysis* (475)
- Advances in vapour pressure studies over liquid uranium plutonium oxides up to 5000K* (476)
- Vapour pressure measurements to 7,000 K and equation of state of oxide fuels for fast reactor safety analysis. I. Extension of vapour pressure measurements of nuclear fuels (U,Pu) O_2 to 7,000 K* (477)

TITLE INDEX

Vapour

Vapour pressure studies over liquid uranium plutonium oxides up to 5,000 K* (478)

variable

Solubility of $\text{Sr}(\text{NO}_3)_2$ in the $\text{LiNO}_3\text{-KNO}_3$ melt of variable composition* (351)
Periodically variable wall temperature effect on heat transfer in liquid metals* (437)
Phases of variable composition with LaF_3 structure in $\text{MF}_2\text{-(Y,Ln)F}_3$ systems. 1. $\text{SrF}_2\text{-(Y,Ln)F}_3$ systems (thermal characteristics, formation of berthollides)* (593)

vapor

Study of saturated vapor pressure above salt melts in $\text{CoCl}_2\text{-MCl}$ systems* (98)

Vercellese

Investigations on radioactive fission product correlations: gamma spectrometry measurements on spent fuel assemblies discharged from the Trino Vercellese reactor at the end of the 2nd irradiation cycle* (75)

Version

MATPRO: Version 09. A handbook of materials properties for use in the analysis of light water reactor fuel rod behavior* (387)

vibration

Study of the compounds $\text{M}_3\text{UO}_2\text{F}_5$ ($\text{M} = \text{K, Rb, Cs, NH}_4$) by i.r. absorption and Raman diffusion spectrophotometry. III. Effect of cation on vibration frequency of the ion $\text{UO}_2\text{F}_5^{3-}$ * (127)
Vibration spectra of rare earth oxytungstates- Ln_2WO_6 composition* (185)

Vibrational

Vibrational spectra and force constants of the elpasolites $\text{Cs}_2\text{KM}_2\text{F}_6$ ($\text{M} = \text{Sc, Y, La, Gd, Yb}$)* (41)
Temperature effect on vibrational spectra of the impurity molecular ions in alkali halides* (46)
Vibrational spectra of transition metal hexafluoride crystals. I. Orthorhombic MoF_6 , WF_6 , UF_6 neat crystals. Technical report* (51)
Vibrational spectra of transition metal hexafluoride crystals. III. Exciton band structures of MoF_6 , WF_6 , and UF_6 . Technical report* (52)
Vibrational fine structure in the uranium emission of Mg_3TeO_6 and Y_6WO_{12} * (59)
Determination of vibrational contribution into excessive mixing entropy of liquid amalgams. Alkali metal amalgams* (138)
Preparational, structural and vibrational study of $\text{M}_2\text{U}_2\text{O}_5\text{Cl}_4$, $2\text{H}_2\text{O}$ ($\text{M} = \text{Rb, Cs}$). Detection of a tetranuclear anion $[(\text{UO}_2)_2]_4\text{O}_2\text{Cl}_8(\text{H}_2\text{O})_2^{4-}$ * (498)

violet

Structure of $4f^6$ and $4f^8$ levels and Eu^{3+} and Th^{3+} configurations in short-range ultra-violet* (589)

Viscosities

Viscosities of cesium vapor to 1,620 K and of liquid gallium to 1,800 K* (622)
Viscosities of cesium vapor to 1620 K and of liquid gallium to 1000 K* (681)

TITLE INDEX

Viscosity

- Viscosity characteristics of the iron-chromium-nickel melts* (24)
 Density, viscosity, electric conductivity and meltability of $\text{Na}_{\text{sub } 3}\text{AlF}_{\text{sub } 6}\text{-La}_{\text{sub } 2}\text{O}_{\text{sub } 3}$ and $\text{Na}_{\text{sub } 3}\text{AlF}_{\text{sub } 6}\text{-Sm}_{\text{sub } 2}\text{O}_{\text{sub } 3}$ systems* (43)
 Study on conductivity, density, and viscosity of molten salt systems* (112)
 Density, surface tension, and viscosity of uranium trichloride-sodium chloride melting* (141)
 Density, surface tension and viscosity of melts of uranium trichloride-lithium chloride* (142)
 Viscosity of $\text{UCl}_{\text{sub } 3}\text{-MgCl}_{\text{sub } 2}$ binary system melts* (145)
 Viscosity of molten $\text{CaCl}_{\text{sub } 2}\text{-UCl}_{\text{sub } 3}$ mixtures* (146)
 Electron contribution to the liquid metal viscosity* (346)
 Density and viscosity of distillable products from $\text{ZrCl}_{\text{sub } 4}$ and $\text{HfCl}_{\text{sub } 4}$ interaction with $\text{POCl}_{\text{sub } 3}$ * (458)
 Nuclear quadrupole relaxation and viscosity in liquid metals* (564)
 Viscosity of melts of $\text{KCl-KF-K}_{\text{sub } 2}\text{ZrF}_{\text{sub } 6}$ ternary system* (576)
 Viscosity of binary of nitrate systems melts* (592)

vitrecus

- Enthalpy of vitreous beryllium fluoride* (616)

VM

- Electrochemical machining of the molybdenum VM-1 alloy* (129)

volatility

- Behavior of ruthenium in fluoride-volatility process. II.
 Fluorination of $\text{RuO}_{\text{sub } 2}$ by $\text{F}_{\text{sub } 2}$ * (558)
 Adaptation of the continuous cold trap system of fluidized-bed to the fluoride volatility process* (724)

voltage

- Intensity-voltage curves of hydrogen diffusion electrodes in eutectic LiCl-KCl * (507)
 Measurement of the cell voltage of $\text{ThO}_{\text{sub } 2}\text{-Y}_{\text{sub } 2}\text{O}_{\text{sub } 3}\text{-solid electrolyte}$ cells in the range of mixed conduction* (535)

W

- Phase equilibria in system U-Pu-W-C * (648)

wall

- Periodically variable wall temperature effect on heat transfer in liquid metals* (437)

waste

- Radiolytic gas generation in plutonium contaminated waste materials* (301)
 Study of the behavior of high activity waste produced during the regeneration of fast reactor fuel elements by the gaseous fluoride method* (311)
 Twenty years of research in the processing of radioactive fuel and neutralization of radioactive waste at the Nuclear Research Institute (NRI)* (358)

water

- MATPRO: Version 09. A handbook of materials properties for use in the analysis of light water reactor fuel rod behavior* (387)

TITLE INDEX

water

Collection of gamma spectra data of irradiated light water moderated reactor spent fuel and study of the applicability of the methods for fuel identification. Final report for the period 1 February 1972--October 1975* (453)
Specialists' meeting on the behaviour of water reactor fuel elements under accident conditions* (510)

WCl

$\text{FeCl}_{3\text{sub}}-\text{MoCl}_{5\text{sub}}-\text{NbCl}_{5\text{sub}}-\text{WCl}_{6\text{sub}}$ system* (554)

weight

Terminal solid solubility of hydrogen in the zirconium-2.5 weight % niobium alloy* (587)

Weizmann

Israel Nuclear Society symposium on "Nuclear fuel in Israel", Weizmann Institute of Science, Rehovot, 1.7.75* (226)

Westinghouse

Westinghouse Advanced Reactors Division* (57)

Wetting

Wetting of certain refractories of the system $\text{MgO}-\text{Al}_{2\text{O}_{3\text{sub}}}-\text{ZrO}_{2\text{sub}}$ with metallic melts* (344)

WF

Vibrational spectra of transition metal hexafluoride crystals. III. Exciton band structures of $\text{MoF}_{6\text{sub}}$, $\text{WF}_{6\text{sub}}$, and $\text{UF}_{6\text{sub}}$. Technical report* (52)

WF6

Vibrational spectra of transition metal hexafluoride crystals. I. Orthorhombic MoF_6 , WF_6 , UF_6 neat crystals. Technical report* (51)

where

Investigation into properties of liquidus curves of binary salt system melts where chemical compounds formation. 2. Systems of monc-monovalent and bi-bivalent cations with common anion* (708)

WO

Vibrational fine structure in the uranium emission of $\text{Mg}_{3\text{sub}}\text{TeO}_{6\text{sub}}$ and $\text{Y}_{6\text{sub}}\text{WO}_{12\text{sub}}$ * (59)

Phase diagrams of $\text{Li}_{2\text{sub}}\text{MoO}_{4\text{sub}}-\text{ZnMoO}_{4\text{sub}}$, $\text{K}_{2\text{sub}}\text{MoO}_{4\text{sub}}-\text{ZnMoO}_{4\text{sub}}$ and $\text{K}_{2\text{sub}}\text{WO}_{4\text{sub}}-\text{ZnWO}_{4\text{sub}}$ systems* (160)

Phase relations in $\text{PbWO}_{4\text{sub}}-\text{TR}_{2\text{sub}}(\text{WO}_{4\text{sub}})_{3\text{sub}}$ systems* (168)

Vibration spectra of rare earth oxytungstates- $\text{Ln}_{2\text{sub}}\text{WO}_{6\text{sub}}$ composition* (185)

Double tungstate $\text{K}_{5\text{sub}}\text{La}(\text{WO}_{4\text{sub}})_{4\text{sub}}$ * (338)

Practical application of conversion method in study of five-component mutual system consisting of nine salts NaF , $\text{Na}_{2\text{sub}}\text{MoO}_{4\text{sub}}$, $\text{Na}_{2\text{sub}}\text{WO}_{4\text{sub}}$, KF , $\text{K}_{2\text{sub}}\text{MoO}_{4\text{sub}}$, $\text{K}_{2\text{sub}}\text{WO}_{4\text{sub}}$, $\text{BaF}_{2\text{sub}}$, $\text{BaMoO}_{4\text{sub}}$, $\text{BaWO}_{4\text{sub}}$ * (516)

System $\text{Na}_{2\text{sub}}\text{WO}_{4\text{sub}}-\text{NaCl}-\text{CaWO}_{4\text{sub}}-\text{CaCl}_{2\text{sub}}$ * (637)

Work

Work function of gallium* (583)

workshop

Proceedings of the symposium and workshop on advanced battery research and design, March 22--24, 1976* (269)

wrought

Effect of composition and structure on crevice, intergranular, and stress corrosion of some wrought Ni--Cr--Mo alloys* (604)

TITLE INDEX

wurtzite

Structure of the biexciton molecule and optical transition selection rules in zincblende and wurtzite materials* (186)

WWR

Spectrophotometric determination of uranium in spent fuels of WWR-2 reactors* (632)

X

X-ray diffraction study into molten bromides and iodides of alkali metals* (20)

Electronic structure of CeN studied by x-ray-photoemission spectroscopy* (28)

X-ray study of the plutonium--oxygen phase diagram at ambient temperatures up to 1100°C * (62)

Calculation of the oxygen potential of $(\text{U,Pu})\text{O}_{2-x}$ with defect models and measured oxygen potential data in the temperature range from 1,000 to 1,700 K* (74)

Investigation into reactions between trioctylammonium nitrates and rare earth nitrates by X-ray electron spectroscopy* (113)

Thermodynamic properties of $(\text{U,Pu})\text{O}_{2-x}$ described by a cluster model* (133)

Actinide metallocarborane complex; synthesis and x-ray structure determination of the bis[n^{sup}

5⁻-(3)-1,2-dicarbollyl]dichlorouranium(IV) dianion* (194)

High temperature thermal expansion of ThO_2 , MgO and Y_2O_3 by X-ray diffraction* (242)

Specific heat of PuC_{1-x} and Pu_2C_3 from 10 to 300 K* (414)

Specific heat of PuC_{1-x} and Pu_2C_3 from 10 to 300 K* (415)

Monte Carlo analysis of diffusion and thermodynamics in UO_{2+x} * (441)

Thermodynamics of non-stoichiometry. A model for UC_x * (442)

Contribution to the establishment of the phase diagram for the U--O--C system by means of x-ray diffractometry at high temperatures and under controlled pressure* (503)

Chemistry of elements niobium and tantalum. 87. Studies on the thermal formation and decomposition of the complexes $[\text{NbCl}_{12}]^{2+}$, $[\text{NbI}_8]^{3+}$, and $[\text{TaX}_{12}]^{n+}$ ($\text{X}=\text{Cl}, \text{Br}, \text{I}$)* (561)

Oxygen potential and vacancy structure of the mixed oxide $(\text{UPu})\text{O}_{2-x}$ * (565)

Covalent compounds of uranium of the type $\text{F}_x/\text{U}(\text{OTeF}_5)_{6-x}$ * (571)

X-ray spectral study of the carbides and nitrides of group IV and V metals* (582)

X-ray diffraction and infrared: Raman spectroscopic study of the hydrogen dinitrate ion, $(\text{O}_2\text{NO}-\text{H}-\text{ONO})^+$, in cesium hydrogen dinitrate* (695)

Determination of trace impurities of titanium and zirconium in molybdenum and tungsten and their oxygen compounds by X-ray fluorescence spectrometry* (698)

XPS

XPS valence bands of La, Ce and Gd and their aluminium alloys* (598)

XPS studies of actinide materials* (662)

TITLE INDEX

x10

Thermophysical and electrophysical properties of uranium hexafluoride at temperatures of $(1-11) \times 10^3$ degree K and pressures of 0.1--100 atm* (302)

Y

Vibrational spectra and force constants of the elpasolites $\text{Cs}_{2\text{KMf}}\text{F}_6$ ($\text{M}=\text{Sc}, \text{Y}, \text{La}, \text{Gd}, \text{Yb}$)* (41)

Vibrational fine structure in the uranium emission of Mg_3TeC and Y_6WO_{12} * (59)

Electronic conductivity and polarization effects of solid ThO_2 - Y_2O_3 electrolytes and of oxygen containing Va-metal electrodes* (193)

High temperature thermal expansion of ThO_2 , MgO and Y_2O_3 by X-ray diffraction* (242)

Entropy change for the lambda-type transition in nonstoichiometric $\text{U}_{4-y}\text{O}_{9-y}$ * (260)

Phase transition in doped $\text{U}_{4-y}\text{O}_{9-y}$. I. Heat capacity measurement* (261)

R_3T_2 compounds (R = rare earth or Y ; T = $\text{Rh}, \text{Pd}, \text{Pt}$) with the rhombic Er_3Ni_2 structure type* (365)

Thermal conductivity and diffusivity of $\text{U}_{4-y}\text{O}_{9-y}$ from 250 to 415 K* (459)

Phase relations in the alkaline earth fluoride-yttria systems (in three parts). 1. The CaF_2 - Y_2O_3 phase diagram* (468)

Phase relations in the alkaline earth fluoride-yttria systems (in three parts). 2. The BaF_2 - Y_2O_3 phase diagram* (469)

Measurement of the cell voltage of ThO_2 - Y_2O_3 -solid electrolyte cells in the range of mixed conduction* (535)

Phases of variable composition with LaF_3 structure in MF_2 -(Y, Ln) F_3 systems. 1. SrF_2 -(Y, Ln) F_3 systems (thermal characteristics, formation of berthollides)* (593)

Yb

Vibrational spectra and force constants of the elpasolites $\text{Cs}_{2\text{KMf}}\text{F}_6$ ($\text{M}=\text{Sc}, \text{Y}, \text{La}, \text{Gd}, \text{Yb}$)* (41)

Negative temperature coefficients of electrical resistivity: the divalent liquid metals Eu, Yb and Ba * (224)

YCl

Raman spectroscopic studies of yttrium (III) chloride--alkali metal chloride melts and of $\text{Cs}_2\text{NaYCl}_6$ and YCl_3 solid compounds* (489)

Investigation into absorption infrared spectra of vapors under ScBr_3 , YCl_3 , LaCl_3 , GdCl_3 , ErCl_3 , LuCl_3 by means of isolation method in rare gas matrix* (497)

years

Twenty years of research in the processing of radioactive fuel and neutralization of radioactive waste at the Nuclear Research Institute (NRI)* (358)

Yevick

Perturbation treatment of the Percus-Yevick equation and structures of liquid magnesium and cadmium* (210)

TITLE INDEX

YF

Interdiffusion in the CaF_2 -- YF_3 system* (669)

yield

Current yield during separation of lithium on a liquid bismuth cathode* (430)

yields

Determination of the burn up and differentiation of the fission yields of uranium-thorium-fuels* (214)

Energy transfer between Bi^{3+} -- YieldsEu^{3+} , Bi^{3+} -- YieldsSm^{3+} and UO_2^{2+} -- YieldsEu^{3+} in oxide glasses* (539)

YS

Mass spectrometric determination of dissociation energies of high-temperature molecules: ScS , YS , LaS , ZrS , UO * (597)

ytterbium

Kinetics and mechanism into preparation of samarium-, europium- and ytterbium dichlorides and their some properties* (253)

yttria

Phase relations in the alkaline earth fluoride-yttria systems (in three parts). 1. The CaF_2 -- Y_2O_3 phase diagram* (468)

Phase relations in the alkaline earth fluoride-yttria systems (in three parts). 2. The BaF_2 -- Y_2O_3 phase diagram* (469)

Complex impedance of electrochemical cells based on yttria doped thorium* (568)

Zirconia and yttria--thoria ceramics find new uses as solid electrolytes* (595)

yttrium

Thermodynamic properties of some rare earth carbonates and yttrium carbonates* (36)

Chemical separation and spectrographic estimation of rare earths and yttrium in PuO_2 and $(\text{U}, \text{Pu})\text{O}_2$ * (125)

Study of lanthanum, yttrium and erbium oxychlorides in melts of alkali metal chlorides* (196)

Enthalpies of formation of melts of yttrium with aluminium* (201)

Corrosion of liquid alloys of zinc with yttrium and with rare earth elements in LiCl -- KCl melts* (335)

Pressure of saturated vapor of rare earth and yttrium bromides* (391)

Properties of yttrium and scandium basic salts* (393)

Raman spectroscopic studies of yttrium (III) chloride--alkali metal chloride melts and of $\text{Cs}_2\text{NaYCl}_6$ and YCl_3 solid compounds* (489)

Interaction of neodymium- and yttrium chlorides with magnesium oxide* (494)

Solubility of rare earth and yttrium oxochlorides in molten salts - MCl_2 ($\text{M} = \text{Mg}, \text{Ca}, \text{Sr}, \text{Ba}$)* (495)

Spectrophotometric determination of rare earths and yttrium in gallium base alloys* (623)

Zeeman

Optical absorption Zeeman spectroscopy of rare earth orthochromites. II. Theory* (122)

Optical absorption Zeeman spectroscopy of rare earth orthochromites. I. ErCrO_3 * (123)

TITLE INDEX

Zeeman

Zeeman effect and satellite carrier structure in absorption spectra of pure and doped praseodymium and europium double nitrates* (165)

zinc

Studies of the electronic structure of simple metals and their alloys with the aid of Compton scattering and positron annihilation using lithium, magnesium, lithium-magnesium and aluminium-zinc as examples* (48)

Corrosion of liquid alloys of zinc with yttrium and with rare earth elements in LiCl--KCl melts* (335)

Interaction parameters in the Zn--Pb--Sn system at low zinc concentration* (436)

Anodic behavior of cadmium and cadmium-zinc alloys in eutectic melt of potassium- and lithium chlorides* (675)

zincblende

Structure of the biexciton molecule and optical transition selection rules in zincblende and wurtzite materials* (186)

Zirconia

Zirconia and yttria--thoria ceramics find new uses as solid electrolytes* (595)

Zirconium

Zirconium: Physico-chemical properties of its compounds and alloys* (7)

Interaction of uranyl chloride with zirconium tetrachloride in the melt of sodium chloride and potassium chloride* (27)

Properties of zirconium and hafnium tetrachlorides and their saturated solutions* (136)

Structure of the thrium-(zirconium, niobium, ruthenium, rhodium)-carbon systems* (245)

Hydrogen solubility in liquid zirconium* (287)

Preparation and thermochemical stability of uranium-zirconium-carbonitrides* (332)

Preparation and thermochemical stability of uranium-zirconium-carbonitrides. A contribution to nuclear fuels for advanced high-temperature reactors* (333)

Preparation and thermochemical stability of uranium-zirconium-carbonitrides* (334)

3. Zirconium compounds with non-metals* (464)

Phase analysis on simulated LWR fuel rods and its evaluation with the aid of constitution studies in the system uranium-zirconium-oxygen* (509)

Phase analysis at simulated LWR fuel rods and their interpretation by constitution investigations in the uranium-zirconium-oxygen system* (510)

Spectrochemical method for determination of zirconium in molten salts* (566)

Terminal solid solubility of hydrogen in the zirconium-2.5 weight % niobium alloy* (587)

High-temperature heat contents and entropies of zirconium fluoride and zirconium sulfate* (591)

Thermographic determination of formation heats of lower zirconium chlorides and bromides* (635)

Zirconium: Physico-chemical properties of its compounds and alloys* (680)

TITLE INDEX

zirconium

Determination of trace impurities of titanium and zirconium in molybdenum and tungsten and their oxygen compounds by X-ray fluorescence spectrometry* (698)

Zn

Thermodynamic properties of Cd-Zn-Hg liquid alloys* (340)

Interaction parameters in the Zn--Pb--Sn system at low zinc concentration* (436)

Using molecular dynamics method for the evaluation of Fe-Ni and Al-Zn melts structure properties* (512)

Preparation and structure of transition metal fluorides

U^{II} M^{IV} F₆ [U^{II} : Cr-Zn; M^{IV} : Hf, Zr]* (596)

ZnCl

Diagram of excess free energy of the PbCl₂-CsCl-ZnCl₂ system* (402)

ZnMoO

Phase diagrams of Li₂MoO₄--ZnMoO₄, K₂MoO₄--ZnMoO₄ and K₂WO₄--ZnWO₄ systems* (160)

ZnWO

Phase diagrams of Li₂MoO₄--ZnMoO₄, K₂MoO₄--ZnMoO₄ and K₂WO₄--ZnWO₄ systems* (160)

Zr]

Preparation and structure of transition metal fluorides

U^{II} M^{IV} F₆ [U^{II} : Cr-Zn; M^{IV} : Hf, Zr]* (596)

ZrC

Spreading of iron-nickel alloys on ZrC and nickel-cobalt alloys on ZrN* (252)

ZrCl

Standard formation enthalpy of chloride alloys of ZrCl₄--KCl (NaCl) systems* (8)

Me₂ZrCl₆--H₂(Me-Na, K, Rb, Cs) system* (45)

Density and viscosity of distillable products from ZrCl₄ and HfCl₄ interaction with POCl₃* (458)

ZrCl₄--TiCl₃--NaCl ternary system* (548)

ZrF

Molecular vapor composition and mass spectra in KF-ZrF₄, CsF-ZrF₄ systems* (294)

Partial pressures, activities and evaporation heat in the KF-ZrF₄ system* (295)

Interval of congruent and incongruent sublimation of 3KFxZrF₄ compound in the KF-ZrF₄ system* (296)

K₂HfF₆--K₂ZrF₆--1.25 percent of HF system at 40°C with other solubility curves from 25°C to 70°C* (541)

Viscosity of melts of KCl-KF-K₂ZrF₆ ternary system* (576)

ZrN

Spreading of iron-nickel alloys on ZrC and nickel-cobalt alloys on ZrN* (252)

ZrO

Wetting of certain refractories of the system MgO-Al₂O₃--ZrO₂ with metallic melts* (344)

TITLE INDEX

ZrO

Study of the retention of Ba in UO_2 nuclear fuel particles by ZrO_2 as getter* (470)

ZrS

Mass spectrometric determination of dissociation energies of high-temperature molecules: ScS, YS, LaS, ZrS, UO^* (597)

THIS PAGE
WAS INTENTIONALLY
LEFT BLANK

INTERNAL DISTRIBUTION

1-2. Central Research Library	5. Laboratory Records (RC)
3. Document Reference Section	6. Duane H. Smith
4. Patent Section	7. Park T. Owen

EXTERNAL DISTRIBUTION

- 8-9. Technical Information Center, P.O. Box 62, Oak Ridge, TN 37830
- 10. Assistant Manager for Energy Research and Development, Oak Ridge Operations, P.O. Box E, Oak Ridge, TN 37830