A. Wolf¹ and R. F. Casten²¹Negev Research Center, Beer-Sheva, Israel²Brookhaven National Laboratory, Upton, New-York, USA

ABSTRACT

Experimental $B(E2; 0_1^+ \rightarrow 2_1^+)$ values for 54 even-even nuclei in the range $A = 90 - 200$ were used to deduce effective charges of proton, neutron bosons. The results indicate that for deformed nuclei, the ratio e_n / e_p is in general agreement with the schematic Bohr and Mottelson prediction. For vibrational nuclei in the $A = 100$ and $A = 140$ regions a very significant enhancement of this ratio is found.

INTRODUCTION

It is well known that the addition of valence nucleons to the closed shells of a nucleus is accompanied by the appearance of a non-spherical field and thus the presence of an additional quadrupole moment of the protons inside the closed shell. This quadrupole core polarization effect can be renormalized as an effective charge of the nucleons added to the core. An estimate of this effect can be obtained following Bohr and Mottelson¹ and noting that the induced quadrupole moment due to the distorted field is of the order $\frac{Z}{A} Q_{sp}$, where Q_{sp} is the single particle quadrupole moment. Thus, the effective charges of the proton, neutron are expected to be :

$$e_p^{eff} = (1 + \frac{Z}{A}) e \quad (1)$$

$$e_n^{eff} = \frac{Z}{A} e, \quad (2)$$

respectively. According to this schematic model¹ :

$$e_n^{eff} / e_p^{eff} = Z / (A + Z) \quad (3)$$

Experimental information on the effective charges of protons, neutrons outside closed shells is of importance for our understanding of the core polarization effect. Such information can be extracted from data on $E2$ transition probabilities. In this work we present a systematic analysis of $B(E2; 0_1^+ \rightarrow 2_1^+)$ experimental data in even-even nuclei in the framework of the IBA-2 model.

RESULTS AND DISCUSSION

We use exact and approximate analytic formulas^{2,3} to extract the effective charges e_p , e_n of the proton, neutron bosons respectively. We used 54 B(E2) experimental values for nuclei in the range $A = 90 - 200$ from the compilation of Raman et al.⁴ These values were fitted to the analytical formulas using a least squares fit procedure. In order not to obscure local variations of e_p , e_n as a function of A , we divided the data in 11 groups of neighboring nuclei and the fits were performed separately for each group, using the appropriate analytical formula^{2,3} for each group. This procedure is different from that of Raman et al.⁵, who tried to find a single e_n / e_p value for a large number of nuclei. In Figure 1 we present as an example the fits for two groups of isotopes. The resulting e_p , e_n as a function of the average mass A in each group are presented in Figure 2. We see that around $A = 100$, the behavior of the effective charges is rather irregular, while for $A > 140$ systematic trends can be observed: e_p increases and e_n decreases with mass number. The behavior of e_n / e_p vs. A is presented in Figure 3. For most deformed nuclei, $e_n / e_p = 0.3 - 0.5$, in agreement with the expected ratio¹ $e_n / e_p = Z / (A + Z) \sim 0.3$ (eq. (3) above). Vibrational nuclei in the regions $A = 100$ and $A = 140$ have $e_n / e_p = 1.5 - 2$. The reason for the enhanced e_n / e_p ratio in these nuclei is not completely understood at present, although it is a clear feature of the data. It may be related⁶ to the fact that e_n , e_p contain a $(\text{length})^2$ factor and that in these nuclei the neutrons occupy orbitals significantly farther away from the center of the nucleus than the protons. Thus their effective charge is enhanced by the dimensions of their orbits.

The systematic trends presented in this work warrant detailed theoretical investigation in order to obtain information on the nuclear core polarization effect, which is closely related to the effective charges.

Work supported by the US DOE under Contract DE-AC02-76CH00016.

REFERENCES

1. A. Bohr and B. R. Mottelson, Nuclear Structure (Benjamin, Reading, 1975) vol I, p. 334.
2. J. N. Ginocchio and P. Van Isacker, Phys. Rev. C33, 365 (1986).
3. R. F. Casten and A. Wolf, Phys. Rev. C35, 1156 (1987).
4. S. Raman et al., At. Data Nucl. Data Tables 36, 1 (1987).
5. S. Raman et al., Phys. Rev. C37, 805 (1988).
6. W. D. Hamilton et al., Phys. Rev. Lett. 53, 2469 (1984).

Fig. 1 Fits of experimental B(E2) data to analytical formulas to deduce e_p , e_n for $^{104-112}\text{Ru}$ and $^{182-192}\text{Os}$ isotopes.

Fig. 2 Systematics of e_p , e_n vs. A . Each symbol represents the e_p or e_n extracted for one of the groups of nuclei and is plotted at the average mass of that group.

Fig. 3 Systematics of e_n / e_p vs. A . See caption to Fig. 2

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

END

DATE FILMED

12 / 10 / 90