DOE/PC/89785--8

DE92 004525

DOE/PC/89785--8

SEP 1991

THE SINGLE ELECTRON CHEMISTRY OF COALS

Report for the period
July 1 to September 30 1991

John W. Larsen
and Thomas P. Eskay

Department of Chemistry
Lehigh University
Bethlehem, PA 18105

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

PREPARED FOR THE UNITED STATES DEPARTMENT OF ENERGY

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

482

OBJECTIVES AND SCOPE OF WORKObjectives

To investigate and characterize the single electron reactions of alkyl and alkoxy aromatic compounds in order to determine the role these reactions play in the chemistry of coal.

Scope of Work

1. Attempts will be made to demonstrate that the radicals from inertinite maceral group will initiate the polymerization of 4-vinylpyridine.
2. The molecule, N,N-diphenyl-phenylenediamine, will be deposited in coals to characterize their native free radicals.
3. Tetracyanoquinodimethane (TCNQ) and tetracyanoethylene (TCNE) will be used to characterize the numbers and strengths of single electron donors in coals.

Introduction

Depolymerization of coals at low temperatures may offer advantages over thermal bond cleavage. Because bond cleavage energies of radical cations are lower than the corresponding homolytic bond cleavage energies of the same bond^{1,2}, generation of radical cations in coal may make possible depolymerization at lower temperatures.

We seek to investigate the above possibility using single molecules containing functional groups common in coals. Since the generation of a radical cation requires the removal of an electron from a neutral molecule, a primary focus of the study will be finding oxidants that will remove an electron from compounds with structural similarity to those typically found in coals. The study must also be concerned with the decomposition of radical cations and the products formed as a result of the decomposition.

Results and Discussions

We have previously reported the successful reaction of benzyl phenyl ether with tris-(1,10-phenanthroline)-iron(III) perchlorate in refluxing acetonitrile to produce the products phenol (P), benzyl acetamide (BAC), 2-benzylphenol (2BP), and 4-benzylphenol (4BP)(see reaction 1.)³. Identification of the products was achieved by gas chromatography (GC) analysis. Difficulties with the analysis made it impossible to obtain accurate mass balances.

Because accurate quantitative data is necessary to ensure that other products are not being formed, it is crucial that we perfect the analytical method and procedure before proceeding with additional reactions.

To obtain accurate quantitative data from GC analysis, the response factors of compounds relative to a chosen internal standard must be known and reproducible. Previously, using a Supelco SPB-1 (15 m. x 0.25 mm.) column for analysis with naphthalene as an internal standard, we were unable to reproduce response factors for both products and starting material of reaction 1. Large peak tailing occurred with this column. It was replaced with a J&W DB-FFAP (15 m. x 0.25 mm.) column. Standard solutions containing known amounts of the compounds in reaction 1 were made in methylene chloride the extraction solvent. Conditions were established which gave baseline separations of these compounds (see experimental section for GC conditions and solution preparation methods) and anthracene was adopted as a new internal standard. Naphthalene, the old standard, eluded in the solvent peak tail with the new column. Series of injections were made using the standard solutions and an injector split ratio of 50 to 1.

Reproducibility of response factors was not observed (see results and data section). Because split ratios below 100 to 1 have been shown to produce non linear discrimination of compounds at the injection port⁴, the split ratio was adjusted to 115 to 1 and acceptable reproducibility of response factors was achieved (see results and data section).

To test the response factors, a synthetic reaction mixture was prepared in methylene chloride and analysis of this solution produced quantitative data that agreed well with the amounts weighed into the mixture. Results of this experiment are in the data and results section in table III.

Next, a synthetic reaction mixture was prepared in acetonitrile to test the reaction workup procedure. Several portions of this mixture were carried through the workup (see table V for workup results) and one portion was used as a control. GC analysis of the control failed to produce agreement with the amounts of the compounds weighed into the mixture (see table IV.). Phenol showed the largest disagreement. The weight calculated from GC analysis was only 86 ± 1 % of the weight of phenol in the mixture. Because a change in solvent produced different results, experiments were performed to determine whether response factors using methylene chloride as the solvent and response factors using acetonitrile were different. Series of analysis were made of the reaction mixture in acetonitrile. Anthracene was not soluble in acetonitrile and response factors were based on using 2-benzyl phenol as an internal standard. These data were compared with

response factors measured in methylene chloride. Different response factors were found for the solutions in the two different solvents. In particular, phenol, had a response factor of 1.34 ± 0.01 in methylene chloride and 1.65 ± 0.05 in acetonitrile. Differences in other compounds response factors existed but were not as pronounced as phenol (see results and data section). To verify the above results a second acetonitrile solution was analyzed and gave almost identical results (see tables VI and VII.).

The effect of the two different solvents must occur before solvents and compounds are separated in the GC column. The most reasonable supposition is that injector splitter discrimination is different in acetonitrile and methylene chloride. Because our reaction analysis will contain both solvents in unpredictable ratios, we must develop an analysis system that will not be sensitive to the discrimination that the different solvents create. Work is currently underway to establish whether the current GC analysis can be modified to overcome the discrimination problems. If we are not successful, we must move to a splitless injection system.

Experimental

Materials

All compounds except benzyl acetamide (TCI), methylene chloride (Fisher A.C.S Grade), and benzyl phenyl ether (see synthesis below) were purchased from Aldrich. 4-benzylphenol was recrystallized from n-hexane and all other chemicals were used from the bottle without further purification. All compounds used showed only one peak other than solvent in GC analysis.

NMR spectra were obtained on a Jeol FX90Q FT NMR spectrometer.

Benzyl Phenyl Ether

Benzyl phenyl ether was synthesized by a modification of a procedure by Pirkla⁵. In an oven dried, nitrogen purged, three neck 50 ml. flask fitted with a condenser and equipped with a magnetic stirrbar was placed 5.29 g. (38.3 mmol.) of anhydrous K_2CO_3 , 2.7 ml. (30 mmol.) of phenol, and 35 ml. of dried acetone (dried by shaking over $CaSO_4$ and distilling under nitrogen⁶). The resulting mixture was stirred for two hours prior to the addition of 3.8 ml. (34 mmol.) of benzyl chloride. The mixture was then heated to reflux for three days. The reaction solution was then filtered repeatedly washing the solid with acetone. The filtrate was then rotary evaporated to produce a yellow oil product $rf=0.58$ on Baxter MK6F/F254 TLC plates 1:4 CH_2Cl_2 :Petroleum ether b.p. 20-40°C. Flash

chromatography of the oil with grade 60 (230-400 mesh) silica gel using the same solvent system given for TLC produced a white crystalline powder which was recrystallized from ethanol twice. ^1H NMR 6.8-7.5 (m, 10H), 5.0 (s, 1H). The spectrum agreed well with previous spectra of benzyl phenyl ether.

Chromatography Analysis and Methods

Gas chromatography was carried out using a (15 m. x 0.25 mm. ID) J&W DE-FFAP column in a Hewlett-Packard 5880A gas chromatograph equipped with a FID detector. The carrier gas was He at a flow rate of $0.5\text{ cm}^3/\text{min.}$ and the split ratio was either 115 or 50:1 (see text). The flow rate was determined by methane injections at 180°C . The analysis was carried out using the following program: 180°C for 4.5 minutes, a $10^\circ\text{C}/\text{min.}$ temperature increase to 240°C , and 15 minutes at 240°C . The above program and conditions gave the following retention times in minutes: phenol 3.80, benzyl phenyl ether 7.07, benzylacetamide 9.69, anthracene 11.58, 2-benzylphenol 14.32, 4-benzylphenol 18.31.

The following is representative of how response factor standards and synthetic reaction mixtures were prepared. 0.19967 g. of BPE 0.20455 g. of BAC, 0.09908 g. of 2BP, and 0.08374 g. of 4BP were weighed into a crimp top vial. The crimp top with a teflon lined septum was then crimped onto the vial and 0.12312 g. of phenol was then injected via syringe into the vial through the septum. This was followed by 59.39055 g. of methylene chloride also injected through the septum.

The workup procedure used for synthetic reaction mixtures carried through the reaction procedure is as follows. One ml. of a prepared synthetic reaction solution was pipetted into a 5 ml. conical vial containing 1 ml. of saturated aqueous NaHCO_3 . One ml. of CH_2Cl_2 was added to the solution along with excess sodium chloride. The CH_2Cl_2 layer was removed after shaking the vial and placed into a vial containing a known amount of anthracene. The remaining aqueous layer was then extracted with 0.6 and 0.5 ml. of CH_2Cl_2 , and the organic layers from each extraction were added to the original organic layer removed.

Response factors (RF) reported were calculated using the following equation and all data reported below are based on response factors relative to the internal standard anthracene except where otherwise noted.

$$RF = \frac{(\text{area of standard}) (\text{weight of compound})}{(\text{area of compound}) (\text{weight of standard})}$$

Data and Results

Table I. Response factors of compounds in reaction 1. split ratio 50 to 1.

Compound	RF $\pm$ standard deviation
P	1.22 $\pm$ 0.10
BPE	1.12 $\pm$ 0.04
BAC	1.58 $\pm$ 0.02
2BP	1.19 $\pm$ 0.01
4BP	1.34 $\pm$ 0.05

Table II. Response factors of compounds in reaction 1. split ratio 115 to 1.

Compound	RF $\pm$ standard deviation
P	1.610 $\pm$ 0.014
BPE	1.242 $\pm$ 0.006
BAC	1.808 $\pm$ 0.008
2BP	1.278 $\pm$ 0.006
4BP	1.385 $\pm$ 0.005

Table III. Response factor test. Weight percent of compound calculated from GC analysis divided by weight of compound weighed into solution. Methylene chloride solvent.

Compound	Weight percent
P	96 $\pm$ 2 %
BPE	100 $\pm$ 1 %
BAC	101 $\pm$ 2 %
2BP	101 $\pm$ 1 %
4BP	104 $\pm$ 1 %

Table IV. Response factor test. Weight percent of compound calculated from GC analysis divided by weight of compound weighed into solution. Acetonitrile solvent.

Compound	Weight percent
P	86 $\pm$ 1 %
BPE	96 $\pm$ 2 %
BAC	96 $\pm$ 3 %
2BP	99 $\pm$ 1 %
4BP	101 $\pm$ 1 %

Table V. Weight percent of compound recovered, after workup procedure, calculated from GC analysis divided by weight of compound weighed into solution. Results reported are from the acetonitrile solution in Table IV..

Compound	Weight percent
P	80 $\pm$ 3 %
BPE	84 $\pm$ 5 %
BAC	90 $\pm$ 2 %
2BP	93 $\pm$ 3 %
4BP	96 $\pm$ 2 %

Table VI. Response factors calculated in acetonitrile and methylene chloride using 2-benzylphenol as an internal standard.

Compound	Response factor(CH ₃ CN)	Response factor(CH ₂ Cl ₂)
P	1.65 ± 0.05	1.34 ± 0.01
BPE	1.05 ± 0.03	0.99 ± 0.02
BAC	1.39 ± 0.01	1.37 ± 0.04
4BP	1.01 ± 0.02	1.06 ± 0.03

Table VII. Response factors calculated in acetonitrile and methylene chloride using 2-benzylphenol as an internal standard. Second run with a newly prepared acetonitrile solution.

Compound	Response factor(CH ₃ CN)	Response factor(CH ₂ Cl ₂)
P	1.58 ± 0.03	1.30 ± 0.02
BPE	1.05 ± 0.01	1.032 ± 0.003
BAC	1.38 ± 0.02	1.41 ± 0.01
4BP	1.00 ± 0.01	1.020 ± 0.002

References

1. D. M. Camaioni, J. A. Franz, J. Org. Chem., 1984, 49, 1601.
2. A. Okamoto, D. R. Arnold, Can. J. Chem., 1985, 63, 2340.
3. J. W. Larsen and T. Eskay, The Single Electron Chemistry of Coals, A paper prepared for the Department of Energy, # DE-FG22-89PC89785-7, August 1991
4. W. Jennings, Gas Chromatography with Glass Capillary Columns, Academic Press; New York, 1980, 2nd. edition.
5. W. H. Pirckle and J. L. Schreiner, J. Org. Chem., 1981, 46, 4988.
6. D. D. Perrin and W. L. F. Armarego, Purification of Organic Compounds, Pergamon Press; New York, 1988, 3rd. edition.

**DATE
FILMED**

2 / 03 / 92