0047 8508120 --6

BNL 37541

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

Physics of Heavy Quark Decays and CP Violation

BNL--37541

DE86 005813

Ling-Lie Chau

Department of Physics
Brookhaven National Laboratory
Upton, New York 11973

Abstract: The phenomena of quark mixing in weak interaction are described. The current experimental status of the mixing matrix, charm and beauty particle decays, and CP violations are reported. Future interesting experiments are pointed out, especially in the event that higher generations of quarks exist.

*Invited talk at the Meson 50 Conference, Kyoto Aug. 15-17, 1985.

MASTER

The submitted manuscript has been authored under Contract No. DE-AC02-76CH00016 with the U.S. Department of Energy. Accordingly, the U.S. Government retains a nonexclusive, royalty-free license to publish or reproduce the published form on this contribution, or allow others to do so, for U.S. Government purposes.

250

Table of Contents

Introduction: Some Historical Remarks

I. The Three-Generation Quark Mixing Matrix

II. CP Violation in the Kaon and Hyperon Decays

III. Mass-Matrix Mixing and CP Violation in Heavy Quark System

III.a. Mass-Matrix Mixing and CP Violation --

Tagged-Neutral-Meson-Decay Experiments

III.b. Decay-Amplitude CP Violation --

Partial-Decay-Rate Differences

IV. Comparison of Models for CP Violation

V. Nonleptonic Decays

V.a Charm Mesons $\rightarrow$ Pseudoscalar-vector Decays

V.b Charm Mesons $\rightarrow$ Pseudoscalar-Pseudoscalar Decays

VI. Beyond the Three Generations of Quarks

VII. Concluding Remarks and Outlook

References

Acknowledgment: The advances in the subject discussed here have been made by many theorists and experimentalists, whose contributions are cited in the references. My contributions to the subject have been made in recent years in collaboration with H.-Y. Cheng (Indiana University, Bloomington), W.-Y. Keung (University of Illinois, Chicago), and also recently with F. Botella, a Fulbright Fellow from Valencia, Spain. I would like to thank Professor C.N. Yang for many enlightening discussions, and for his encouragement.

I would also like to thank Professor Z. Maki and the organizers of the Meson 50 for inviting me to give this talk and to participate in this most memorable conference.

Introduction: Some Historical Remarks

It is 51 years since Fermi⁽¹⁾ proposed the Fermi coupling for β decay, 50 years since Yukawa⁽²⁾ formulated his theory, and 35 years since Yang and Mills⁽³⁾ constructed their theory. Nature's answer to these human endeavors has been most elegant and rewarding. The electroweak unified theory of Glashow, Salam, and Weinberg⁽⁴⁾ has been confirmed by the recent discovery⁽⁵⁾ of the long-awaited intermediate bosons $W^\pm$ and Z^0 . Their masses,⁽⁶⁾ and even their production and decay characteristics, were predicted. (The observed lepton-forward-backward asymmetry, i.e., ℓ^- from the W^- moves in the proton direction, and ℓ^+ from the W^+ moves in the antiproton direction, provides a truly remarkable confirmation of the quark-proton description of the reaction.)⁽⁷⁾ This was the climax of the strikingly beautiful development of 50 plus years of history of weak interactions, as we have heard from many of the preceeding talks in this conference, and many of you in the audience have made major contributions. The list below demonstrates the beautiful interplay between theoretical ideas and experimental observations:

1930,	Nuclear β decay	----->	ν , ⁽⁸⁾
			1954 discovered, ⁽⁹⁾
1934,	Fermi coupling, ⁽¹⁾		
1935,	Nuclear force $\gg$ Fermi coupling ⁽¹⁰⁾	----->	Yukawa theory, ⁽²⁾
			π meson discovered 1947, ⁽¹¹⁾
1954	Local gauge invariance	----->	Yang-Mills theory, ⁽³⁾
1953-56	θ , τ puzzle ⁽¹²⁾	----->	parity violation, ⁽¹³⁾
			observed 1957, ⁽¹⁴⁾
1960's	Electroweak unification ⁽⁴⁾	----->	neutral current observed 1973, ⁽¹⁵⁾
			$W^\pm$, Z^0 with mass predicted, ^(4,6)
1963	Universal suppression of hyperson decays	----->	Cabibbo mixing, ⁽¹⁶⁾
1970	$K_L \rightarrow \mu^+ \mu^-$ extremely small ⁽¹⁷⁾	----->	the GIM mechanism, ⁽¹⁸⁾ and
			charm, observed 1976, ⁽¹⁹⁾

1973 CP violation from W coupling ----->

Kobayashi-Maskawa
(KM) matrix⁽²⁰⁾ and
(t), b quarks
beauty particle ob-
served 1982-1984⁽²¹⁾

In the light of this milestone of the great success of electroweak unified theory, I am going to review for you some particular areas of research⁽²²⁻²⁴⁾ in weak interactions and their outlook in our ever ongoing pursuit to understand the constituents of matter and the forces governing their interactions.

I. Status of the Quark Mixing Matrix

The mixing matrix of the three left-handed generations of quarks,

$$\begin{pmatrix} d' \\ s' \\ b' \end{pmatrix} = \begin{pmatrix} v_{ud} & v_{us} & v_{ub} \\ v_{cd} & v_{cs} & v_{cb} \\ v_{td} & v_{ts} & v_{tb} \end{pmatrix} \begin{pmatrix} d \\ s \\ b \end{pmatrix}, \quad (1.1)$$

can be determined by various experiments:

$$\begin{aligned} |v_{ud}| &= c_x [1 + O(10^{-4})] = 0.9737, & \text{from nuclear } \beta \text{ decays;}^{(25,26)} & (1.2) \\ |v_{us}| &= s_x [1 + O(10^{-4})] = 0.225, & \text{from strange particle decays;}^{(25,26)} \\ |v_{cb}| &= s_y [1 + O(10^{-4})] = 0.059, & \text{from } \tau_b = 10^{-12} \text{ sec;}^{(27,29)} \\ |v_{ub}| &= s_z & \lesssim 0.0082, \text{ from } \Gamma(b \rightarrow u)/\Gamma(b \rightarrow c) < 0.05^{(28-29)} \end{aligned}$$

Besides the few percent errors for $|v_{ud}|$, ten to twenty percent are typical errors for other v_{ij} 's. Recent news on the ancient Cabibbo fit is that the old experimental discrepancy in the lepton-momentum asymmetry in Σ decays, which gave g_1/f_1 a wrong sign, has now disappeared. A recent high-statistics experiment⁽³⁰⁾ showed $g_1/f_1 = -0.29 \pm 0.07$, which is consistent with the Cabibbo theory. In addition, a somewhat smaller value of $|v_{us}|$, $|v_{us}| = 0.225$ is obtained, which is in better agreement with older fits.⁽²⁵⁾ One interesting experimental uncertainty is still the neutron lifetime.⁽³¹⁾

Our knowledge of the quark mixing matrix has dramatically improved recently owing to the b-lifetime measurement⁽²⁷⁾ and the $\Gamma(b \rightarrow u)/\Gamma(b \rightarrow c) < 0.05$ ⁽²⁸⁾ bound from the $b \rightarrow l^+ X$ measurements, which indicated that X is mainly made out of states with charm. At present, the errors on the b lifetime are still rather large. Here we use $\tau_b = 10^{-12}$ sec as an example to obtain Eq. (1.2). Using unitarity and a phase convention⁽³²⁻³⁴⁾ such that the phase factor of the matrix elements appears with the smallest matrix elements $\lesssim 10^{-3}$, the values of the other matrix element can be obtained through unitarity

$$V = \begin{pmatrix} 1 & 0 & 0 \\ 0 & c_y & s_y \\ 0 & -s_y & c_y \end{pmatrix} \begin{pmatrix} c_z & 0 & s_z e^{-i\phi} \\ 0 & 1 & 0 \\ -s_z e^{-i\phi} & 0 & c_z \end{pmatrix} \begin{pmatrix} c_x & s_x & 0 \\ -s_x & c_x & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

$$\begin{matrix} & d & s & b \\ \begin{matrix} \hline s_x, s_y, s_z \end{matrix} & \begin{pmatrix} c_x c_z & s_x c_z & s_z e^{-i\phi} \\ -s_x c_y - c_x s_y s_z e^{i\phi} & c_x c_y - s_x s_y s_z e^{i\phi} & s_y c_z \\ s_x s_y - c_x c_y s_z e^{i\phi} & -c_x s_y - s_x c_y s_z e^{i\phi} & c_y c_z \end{pmatrix} & \begin{matrix} u \\ c \\ t \end{matrix} \end{matrix} \quad (1.3)$$

$$= \begin{pmatrix} 0.9737 & 0.225 & v_{ub} \lesssim 0.082 e^{i\phi} \\ -0.23 - 0.059 v_{ub} & 0.98 & 0.059 \\ 0.01 - v_{ub} & -0.06 + 0.228 v_{ub} & 1 \pm 0(10^{-2}) \end{pmatrix} \quad (1.4)$$

The interesting features of the mixing matrix are that the gap between the second and the third generations has been widened: $|v_{cb}|$ is about a quarter of $|v_{ud}|$. Because of such severe suppression, the 2×2 GIM matrix is almost unitary, i.e., $V_{cd}^*/V_{cs} = -V_{us}^*/V_{ud}$. It is very important to check this relation independently. Significant deviation from it can have serious implications, e.g., result in breakdown of the model or the existence of the fourth

generation. The nice feature of using the parameterization of Eq. (1.3) is that the imaginary parts appear at matrix elements with magnitudes $< 10^{-3}$. It is also clear in this parametrization that V_{ub} cannot be zero in order to explain CP violation. Actually one can show that in order to give CP violation none of the matrix elements can be zero.⁽²⁴⁾ The analysis of CP violation in the K system can give an estimate of the lower bound on V_{ub} , which is not too far below the experimental upper bound, as we shall discuss later. In addition, each angle has a direct correspondence with experimental results, as shown in Eq. (1.2). Also, its relation to ideas in quark mass matrix can also be described in a more transparent way.⁽³⁵⁾

The CP-violation effects from this 3x3 KM matrix have a quite striking feature. It was shown in Ref. (33) that if there are only three generations of quarks, the CP-violation phenomena in all different decays of three generations of quarks are characterized by one single parametrization-invariant parameter

$$X_{cp} = s_x s_y s_z s_\phi c_x c_y c_z^2 \approx 2 \cdot 10^{-4}, \quad (1.6)$$

as determined from ϵ in K decay.⁽³³⁾ Now this analysis has been carried out for more than three generations of quarks in Ref. (36), where we find that the number of invariant CP violation parameters $X_{cp,i}$ jumps to nine in the four-generation case, saturating the parameter space for generation numbers higher than three. Thus the unique X_{cp} in the three-generation case can serve as a base for searching for signals of the existence of higher generations. I shall come back to this in Section VI.

II. CP Violation in the Kaon and the Hyperon Decays

Weak interactions mediate $K^0 \leftrightarrow \bar{K}^0$, i.e., an off-diagonal term in the mass-matrix exists, so that the physical states are no more $K^0, \bar{K}^0$ but $|K_{S,L}\rangle =$

$N_{\bar{E}}[(1+\bar{E})|K^0\rangle \pm (1-\bar{E})|\bar{K}^0\rangle]$, where $N_{\bar{E}}$ is a normalization factor.

Such mixing gives unequal masses and decay width of the $K_{S,L}$ system,

$\delta m \equiv m_S - m_L$, $\delta\Gamma \equiv \Gamma_S - \Gamma_L$, $f \equiv (\Gamma_S + \Gamma_L)/2$. The mixing parameters δm , $\delta\Gamma$ can be measured in an interference experiment involving regeneration of K_S from a K_L beam. Because of $\Gamma_S \gg \Gamma_L$, the K^0 , $\bar{K}^0$ is a maximally mixed system, since no matter whether we have K^0 or $\bar{K}^0$ to begin with, soon K_S decays away and leaves only K_L which is almost an equal mixture of K^0 , $\bar{K}^0$. The parameter $\eta \equiv |1-\bar{E}|/|1+\bar{E}| \neq 1$ is a phase-convention independent indicator of the mass-matrix CP violation, and was directly measured in K_L decay

$$\begin{aligned} \delta_L &= \frac{\Gamma(K_L \rightarrow \pi^- 2^+ \bar{\nu}) - \Gamma(K_L \rightarrow \pi^+ 2^- \bar{\nu})}{\Gamma(K_L \rightarrow \pi^- 2^+ \bar{\nu}) + \Gamma(K_L \rightarrow \pi^+ 2^- \bar{\nu})} \\ &= \frac{2\text{Re}\bar{E}}{1+|\bar{E}|^2} = \frac{1-\eta^2}{1+\eta^2} = (3.3 \pm 0.12) \times 10^{-3} \quad (2.1) \end{aligned}$$

The mass-matrix CP violation can be calculated through the diagram, Fig. II.1.

After calculating the box graph in the middle, which contains X_{CP} , we still need to calculate hadronic matrix element $B_K = \langle K^0 | \{\bar{d}\gamma_\mu(1-\gamma_5)s\}^2 | \bar{K}^0 \rangle (-4/3 f_K^2 M_K)^{-1}$, which describes how the K^0 , $\bar{K}^0$ are made of quarks. This is the main uncertainty in the calculation. There is still no reliable way to calculate this B_K parameter. For a discussion on the evaluation of the values of B_K , see Ref. (38). However, we shall see later, by fitting ϵ , the range of the value B_K is rather limited in the model of three generations of quarks.

The historically first measured CP violation⁽³⁹⁾ in K decays are

$$\eta_{+-} = \frac{A(K_L \rightarrow \pi^+ \pi^-)}{A(K_S \rightarrow \pi^+ \pi^-)} = \frac{(A_{+-}/\bar{A}_{+-}) - (1-\bar{E})/(1+\bar{E})}{\text{"} + \text{"}} = \epsilon + \epsilon', \quad (2.2a)$$

$$\eta_{+-} = \frac{A(K_L \rightarrow \pi^0 \pi^0)}{A(K_S \rightarrow \pi^+ \pi^-)} = \frac{(A_{00}/\bar{A}_{00}) - (1-\bar{E})/(1+\bar{E})}{\text{"} + \text{"}} = \epsilon - 2\epsilon'. \quad (2.2b)$$

Fig. II.1 Box diagram for $K^0 \rightarrow \bar{K}^0$ transition, the B_K parameter describes the transition without the enclosed part; for details, see Ref. (38).

Fig. II.2 W-loop diagram for $K^0 \rightarrow \pi^+ \pi^-$. Its one-gluon-exchange approximation is the "Penguin" diagram. The B_K parameter describes the transition without the enclosed part; for details see Ref. (38,44).

Fig. II.3 Fixed m_t , or ϵ'/ϵ contours in the s_2 - ϕ plane, using the b lifetime to be 1 psec, and the parameters $B_K = 0.33$, $B_K' = 0.75$; for discussions on long-range effects and details see Ref. (44).

One can easily show that^(40,24) $\epsilon' = 0$ only if $A_{+-}/\bar{A}_{+-}$ as well as $A_{00}/\bar{A}_{00}$ can simultaneously be made to be one, i.e., no decay-amplitude CP violation. In that case and phase convention, $\eta_{+-} = \eta_{00} = \epsilon = \bar{\epsilon}$. Thus $|\eta_{00}/\eta_{+-}|^2 - 1 \neq 0$ is a measurement of decay-amplitude CP violation. The current experimental data give

$$|\eta_{00}/\eta_{+-}|^2 - 1 = -6 \operatorname{Re}(\epsilon'/\epsilon) = \begin{cases} -0.0046 \pm 0.0053 \pm 0.0024, & (41) \\ +0.0017 \pm 0.0082, & (42) \end{cases} \quad (2.3)$$

which are still consistent with zero. It was noted by Gilman and Wise⁽⁴³⁾ that the KM model can give a nonvanishing number of ϵ' . The W-loop diagram shown in Fig. II.2 is essential in giving such an effect. The one-gluon exchange approximation (the "Penguin") has been used to estimate the effect. Again the main uncertainty comes from the hadronic amplitude of B_K' , which describes how the K and 2π are related through quark fields, Fig. II.2. For a discussion of the values of B_K' from various calculation, see Refs. (38 and 44).

Since the KM matrix is designed to give CP violation, the CP-violation measurements ϵ , ϵ'/ϵ must give further constraint on the quark mixing matrix.

After the determination of $s_y = 0.059$ from the b lifetime, and $s_z < 0.0082$ from $|V_{ub}/V_{cb}| < 0.14$, we still have s_z , s_ϕ , m_t as independent parameters. After fitting ϵ , assuming some values for B_K , B_K' , we obtain fixed m_t , or fixed ϵ'/ϵ contours in the s_z , ϕ plane, Fig. II.3. From such analyses, we learn the following: (1) The mass of the t quark m_t cannot be too small, depending on the values of B_K' , use of m_t $\lesssim$ 40-GeV current data indicates $B_K > 0.33$. 2) other than all the other calculable factors, $|\epsilon'/\epsilon|$ depends upon B_K' , and its sign depends upon the relative sign of B_K , B_K' ; current data indicate $B_K' < 1$. 3) The lower limit on s_z from CP violation is already only a factor of 2 or 3 of the upper limit given by the experimental $|V_{ub}|$ bound.

We can see that the three-generation quark model is quite narrowed down by the current experiments.

There are now two experiments⁽⁴⁵⁾ being carried out measuring $|\eta_{00}/\eta_{+-}|$ of the K_S, K_L system with higher sensitivity. Another new type⁽⁴⁶⁾ of experiment is being planned to measure ϵ'/ϵ via the tagged $K^0, \bar{K}^0 \rightarrow 2\pi$ decays at LEAR, CERN, through the following reactions $p\bar{p} \rightarrow \pi^+ K^- K^0$, or $\pi^- K^+ \bar{K}^0$; ⁽⁴⁷⁾ we eagerly await the results from these experiments, and most importantly the observation of the top quark and its mass.

Here I would like to point out an interesting historical irony. Because of the $\Delta = 1/2$ - dominance phenomenon from $\Gamma(K^+ \rightarrow \pi^+ \pi^0)/\Gamma(K^0 \rightarrow \pi^+ \pi^-) \approx 1/670$, ϵ'/ϵ is automatically suppressed by a factor of 20. This same factor, which is now causing difficulty for experimentalists in their attempts to measure ϵ'/ϵ , actually made it possible for Professor Dalitz to have some fifteen τ events ($K^+ \rightarrow \pi^+ \pi^+ \pi^-$) in the 1950's. If there is no such suppression of $\Delta I = 3/2$ decays of $K^+ \rightarrow \pi^+ \pi^0$, the θ, τ puzzle would not have been established before 1956, and the discovery of parity violation would have been delayed for many years. Now this $\Delta I = 3/2$ suppression is making the measurement of ϵ'/ϵ difficult. So we began to ask the question whether there are decay channels of the kaon, such that this suppression rule does not apply. Indeed we found that in $K_L \rightarrow \gamma\gamma$, the decay amplitude CP-violation effect can be as big as the mass-matrix (superweak) CP-violation effect.⁽⁴⁸⁾ It is possible to observe the CP-violation effect in the K_L system only if it is not swamped by $K_S \rightarrow \gamma\gamma$. Interestingly, we found that $A(K_S \rightarrow \gamma\gamma) \approx 2.4 A(K_L \rightarrow \gamma\gamma)$, i.e., $Br(K_S \rightarrow \gamma\gamma) \approx 5 \times 10^{-6}$. The CERN ϵ'/ϵ experiment NA31 can soon measure this branching ratio. If it turns out that $K_L \rightarrow \gamma\gamma$ is not overshadowed by $K_S \rightarrow \gamma\gamma$, as predicted by our calculation, it is possible to measure the CP violation in $K_L \rightarrow \gamma\gamma$. To measure such effects in

K_L , one must do the K^0 , $\bar{K}^0$ tagging experiment as provided by the LEAR experiment.

Other decay-amplitude CP violation effects are the partial decay rate difference in $K^\pm \rightarrow (3\pi)^\pm$, and in Λ , $\bar{\Lambda}$ decays.⁽⁴⁹⁾ Convenient quantities to measure are

$$\frac{\Gamma(K^+ \rightarrow \pi^+ \pi^+ \pi^-) / \Gamma(K^+ \rightarrow \pi^+ \pi^0 \pi^0)}{\Gamma(K^- \rightarrow \pi^- \pi^- \pi^+) / \Gamma(K^- \rightarrow \pi^- \pi^0 \pi^0)} = 1 + 2\Delta_{K \rightarrow 3\pi}, \quad (2.4)$$

$$\frac{\Gamma(\Lambda \rightarrow \pi^- p) / \Gamma(\Lambda \rightarrow \pi^0 n)}{\Gamma(\bar{\Lambda} \rightarrow \pi^+ \bar{p}) / \Gamma(\bar{\Lambda} \rightarrow \pi^0 \bar{n})} = 1 + 2\Delta_\Lambda. \quad (2.5)$$

Current estimates give $\Delta_{K \rightarrow 3\pi} \sim 0.74 \text{ e}^\circ$,⁽⁵⁰⁾ and $\Delta_\Lambda \lesssim 10^{-5}$.⁽⁵¹⁾ The advantages of measuring such ratio of ratios rather than the percentage partial decay difference Δ itself are twofold: first the effects are two times as large, secondly some experimental errors tend to cancel. Another interesting CP-violation effect to look for is the difference of the momentum asymmetry parameter of the pion with respect to the polarization of Σ^+ in $\Sigma^+ \rightarrow \pi^+ n$ decay,

$$A_{\Sigma^+} = 20\alpha_{CP} \sim 4 \times 10^{-3}, \quad (2.6)$$

according to an estimate by Chau and Cheng in Ref. (49). Such experiments will be done at BNL, LEAR, or another kaon, hyperon-rich laboratory.

III. Mass-Matrix Mixing and CP Violation in Heavy Quark Decays

III.a. Mass-matrix mixing and CP violation -- Tagged-Neutral-Particle-Decay Experiments:

We are fortunate with the kaon system to have the K_S , K_L system to study mass-matrix mixing (the neutral particle-antiparticle mixing) and the superweak CP-violation parameters $\eta \equiv |1 - \bar{\epsilon}| / |1 + \bar{\epsilon}|$ in that K_L indeed lives rather long because the three-pion mass is very close to the kaon mass. However, heavier quark systems like, D^0 , $\bar{D}^0$; B^0 , $\bar{B}^0$, both mass eigenstates D_S^0 , D_L^0 ; B_S^0 , B_L^0 will be short lived. It is very difficult or

impossible, to do K_L , K_S -type of interference experiments.⁽³²⁾ So the question is whether we still can measure the mass-matrix mixing parameters δm , $\delta \Gamma$, and CP-violating parameter η separately. The answer is yes, as discussed in Refs. (24, 40). However, one has to do neutral-meson-tagged experiments. The D^0 can be tagged in $e^+e^- \rightarrow \psi(4030) \rightarrow \pi^+ D^- D^0$, $\pi^- D^+ D^0$. The three parameters δm , $\delta \Gamma$, η_D can be measured by the following three experiments: one to measure $D^0 \bar{D}^0$ mixing parameter,

$$\gamma_D = \frac{''D^0 \rightarrow \bar{D}^0''}{''D^0 \rightarrow D^0''} = \eta_D^2 \frac{(\delta m/\Gamma)^2 + (\frac{1}{2}\delta\Gamma/\Gamma)^2}{(\delta m/\Gamma)^2 + 2 - (\frac{1}{2}\delta\Gamma/\Gamma)^2}, \quad (3.1)$$

which can be measured by a tagged D^0 , $\bar{D}^0$ decays, $\gamma = ''D^0 \rightarrow \ell^- \times '' / ''\bar{D}^0 \rightarrow \ell^+ \times ''$, Or it can manifest itself in the same sign depletors productions. That $\delta m \neq 0$, $\delta \Gamma \neq 0$ are indications of mixing. Maximal mixing can happen in two ways: one way is when $\bar{\Gamma} = (\Gamma_S + \Gamma_L)/2$ is dominated by either Γ_S or Γ_L so that $(1/2) \delta\Gamma/\bar{\Gamma} \not\ll 1$, which is the case of neutral kaon decays; the other way is $\delta m/\bar{\Gamma} \gg 1$, and $\delta m/\bar{\Gamma} \gg (1/2) \delta\Gamma/\bar{\Gamma}$. We shall see that the latter situation may happen in the heavy quark systems.

Two more experiments are to measure the $\ell^\pm$ partial decay rate differences:

$$\Delta(\ell^+) = \frac{''D^0 \rightarrow \ell^+ X'' - ''\bar{D}^0 \rightarrow \ell^+ X''}{'' + ''} = - \frac{2\text{Re}\bar{\epsilon}_D}{1 + |\bar{\epsilon}_D|} + \frac{\Gamma_S \Gamma_L}{(\delta m)^2 + (\bar{\Gamma})^2}, \quad (3.2)$$

and

$$\Delta(\ell^-) = \frac{''D^0 \rightarrow \ell^- X'' - ''\bar{D}^0 \rightarrow \ell^- X''}{'' + ''} = - \frac{2\text{Re}\bar{\epsilon}_D}{1 + |\bar{\epsilon}_D|} - \frac{\Gamma_S \Gamma_L}{(\delta m)^2 + (\bar{\Gamma})^2}; \quad (3.3)$$

therefore,

$$\Delta(\ell^-) + \Delta(\ell^+) = - \frac{4\text{Re}\bar{\epsilon}_D}{1 + |\bar{\epsilon}_D|^2} \equiv 2 \frac{1 - \eta^2}{1 + \eta_D}, \quad (3.4)$$

is a pure CP violation effect, and

$$\Delta(\ell^-) - \Delta(\ell^+) = \frac{2\Gamma_S \Gamma_L}{(\delta m)^2 + (\bar{\Gamma})^2}, \quad (3.5)$$

is a pure mass-matrix mixing effect. From Eqs. (3.1)-(3.5), knowing either Γ_S , or Γ_L , δm , $\delta\Gamma$, are measurable. The mass-matrix CP-violation effects $\eta_D \neq 1$ can be measured using Eq. (3.4).

It has been quite firmly established⁽²⁹⁾ that in the case of three generations of quarks, the only appreciable neutral meson $P^0\bar{P}^0$ (mass-matrix) mixing and CP-violation effects is in the $B_S^0, \bar{B}_S^0$ mixing. Observing deviation from this prediction can provide evidence for the new physics of higher-than-three generations of quarks.

III.b. Partial-decay-rate differences

Studying the partial-decay-rate differences is a convenient way to study the decay-amplitude CP violations.^(23,49,53) As we mentioned previously, all CP-violation effects in the case of three generations of quarks are from a single CP-violating parameter X_{CP} , which with other quark-mixing matrix elements are rather well known. Given the same uncertainties in the hadronic decay amplitudes, we can search for the most likely channels where partial-decay-rate differences can be big. Consistently, we find in the B decays, charged⁽⁵⁴⁾ as well as neutral, that the partial^(53,55) decay rates can be big (few $\times 10\%$), though the branching ratios of these exclusive decays are very small (10^{-3} $\sim$ 10^{-4}). Here we have an interestingly reversed the situation, as in kaon decays, where the partial-decay-rate differences are very small ($\sim 10^{-3}$), but branching ratios large ($\sim 10\%$), since the number of events needed is inversely proportional to the square of partial-decay-rate difference, but inversely proportional to the branching ratio. Thus we have a more advantageous situation in the B decay.

Here we list some very encouraging examples, (54,55) e.g.

	Δ_{tree}	$\Delta_{\text{W-loop}}$	Br	# of events needed
$B_u^- \rightarrow D^- D^{*0}$	-1.6×10^{-2} (-0.86)	1×10^{-3} (7.3×10^{-4})	3×10^{-3} (4.1×10^{-3})	1.3×10^6 (3.3×10^2)
$B_c^- \rightarrow \pi^- D^0$	4×10^{-2} (-0.86)	3.7×10^{-3} (1×10^{-3})	2.3×10^{-4} (8.5×10^{-4})	2.7×10^6 (1.6×10^3)
$B_d^0 \rightarrow K^- \pi^+$	0	10×10^{-2}	1.6×10^{-4}	6.3×10^5
$B_S^0 \rightarrow D^- F^+$	0	4×10^{-2}	9.6×10^{-4}	5.6×10^5
$B_d^0 \rightarrow \pi^+ \pi^-$	0	25×10^{-2}	6.1×10^{-4}	2.6×10^4
$B_S^0 \rightarrow D^+ D^-$	0	-27×10^{-2}	7.0×10^{-4}	2.0×10^4

The charged B partial rate differences are mainly from the tree graphs, e.g.,

The neutral B^0 partial-rate differences can only come from interference between the tree graph and the loop group, e.g.,

For final state with no definite CP, the partial rate differences are only decay-amplitude CP violation. However, for final states with definite CP, the

partial rate differences are a combined result of mass-matrix mixing, and decay-amplitude CP violation. For details, see Refs. (54,55).

Such large partial-decay-rate differences can be searched for at CLEO II, Fermilab. LEP, SLC of current luminosity design gives barely enough B particles to do such experiments.

IV. Comparison of CP-Violating Mechanisms

Of course there is the left-right symmetric theories;⁽⁵⁶⁾ they have one more phase parameter than the KM model so that they give an easier fit to the current data.

CP violations can also come from the complex Higgs fields. The more specific model of Weinberg⁽⁵⁷⁾ with three Higgs doublet at one point was thought to be ruled out,⁽⁵⁸⁾ i.e., $|\varepsilon'/\varepsilon| \gtrsim 1/4$ too large comparing even to the old experimental bound. Recently, after reexamining the calculations,⁽⁵⁹⁾ $|\varepsilon'/\varepsilon|$ becomes small enough to be compatible with present experimental measurement. But the neutron electric dipole moment d_n ironically is found to be much larger⁽⁶⁰⁾ than previous calculations. However, its violation of the current experimental bound of $d_n = (0.3 \pm 4.8) \times 10^{-25}$ cm⁽⁶¹⁾ depends on an incalculable parameter. H.-Y. Cheng in Ref. (59) gave a very thorough analysis. For other CP-violation effects, see the talk by G. Segre in Ref. (22).

V. Nonleptonic Decays

Ever since the observation of the suppression of $K^+ \rightarrow \pi^+ \pi^0$ in 1956,⁽⁶²⁾ $\Gamma(K^+ \rightarrow \pi^+ \pi^0)/\Gamma(K^0 \rightarrow \pi^+ \pi^-) = 1/670$, i.e., the $\Delta I = 1/2$ dominance rule theorists are still struggling to understanding such phenomena in nonleptonic decays. The uncertainties in the calculation of the parameters of B_K and B_K' reflect the same difficulties. The "penguin" mechanism⁽⁶³⁾ is as elusive as ever. Many brave attempts in model calculations to estimate the nonleptonic charm

decays are challenged again and again as more data become available, and the theory is forced to be amended and modified. Disappointing as it is, this is to be expected if we are to make progress. Here I would like to point out that this process in furthering understanding of nonleptonic decays can be guided by a model-independent analysis (i.e., the quark-diagram-formulation) of the available experimental data, in the same spirit as Fermi introduced his coupling where only the then-known model-independent information was used.

One fortunate feature of charm and heavier quark decays is that there are many channels open, unlike in the K decays which have only 2π , 3π decays. As has been known for some time now,⁽⁶⁴⁾⁽²³⁾ all meson $\rightarrow$ 2 meson decays can be generally described in a model-independent way by six quark diagrams, as shown in Fig. V., multiplied by quark-mixing matrix elements, which are now quite well determined as discussed in Section I. There are twenty-plus channels of decays in D^+ , D^0 , F^+ to two-pseudoscalars, PP, decays, and double that number in pseudoscalar-vector, PV, decays. Thus here the model-independent quark-analysis can be very useful in such a quark-diagram approach. Here I shall demonstrate what we have learned from the present charm-meson-decay data, and what their implications are in this approach.

V.a Charm meson $\rightarrow$ pseudoscalar-vector decays

I shall begin with the PV decays because of the relative simplicity in presenting the discussion, though the data of PV decays are not yet as good as some of the PP decays. The simplicity in discussing the PV decays comes from an interesting feature that because of the purity of the quark content in ϕ and ω , PV many decays are given by one type of amplitude; e.g., for the quark mixing-matrix unsuppressed decays, we have^(23,65,66)

Fig. V.a The six quark diagrams for inclusive meson decay.

Fig. Vb The six quark diagrams for a meson decaying to two mesons.

$$\begin{aligned}
A(D^0 \rightarrow \phi \bar{K}^0) &= (c_1)^2 (c') ; \\
A(F^+ \rightarrow \phi \pi^+) &= (c_1)^2 (a') , \\
A(F^+ \rightarrow \rho^+ \pi^0) &= -A(F^+ \rightarrow \rho^0 \pi^+) = (c_1)^2 (d-d') , \\
A(F^+ \rightarrow \omega \pi^+) &= (c_1)^2 (d+d') .
\end{aligned} \tag{5.1}$$

for the quark mixing-matrix singly suppressed decays, we have

$$A(D^+ \rightarrow \phi \pi^+) = \sqrt{2} A(D^0 \rightarrow \phi \pi^0) = s_1 c_1 (b')$$

$$A(D^0 \rightarrow \bar{K}^{*0} K^0) = -A(D^0 \rightarrow K^{*0} \bar{K}^0) = s_1 c_1 (c'-c) . \tag{5.2}$$

(Here the primed and unprimed amplitudes differ depending on which side in the diagrams the vector boson comes out, see Ref. (66) for details.) Therefore these amplitudes can be directly obtained from experimental measurements and compared with theoretical calculations. By fitting the data

$$Br(D^0 \rightarrow \phi \bar{K}^0) = (0.7 \pm 0.2\%),^{(67)} (0.9 \pm 1.4)\%,^{(68)} (1.4 \pm 0.4)\%,^{(69)} \tag{5.3}$$

$$Br(F^+ \rightarrow \phi \pi^+) = 4.4\%,^{(68)} 11\%,^{(70)} (3.3 \pm 1.1)\%,^{(71)} \tag{5.4}$$

and the Cabibbo suppressed decay data,

$$Br(D^+ \rightarrow \phi \pi^+) = (0.93 \pm 0.26 \pm 0.16)\%,^{(67)} \tag{5.5}$$

we find the ratios $|c'|/|a'|$, $|b'|/|a'|$. From the life time measurements

$$\tau_{D^\pm} = (8.8^{+1.0}_{-0.8})10^{-13} \text{ sec}, \tau_{D^0} = (4.3^{+0.4}_{-0.3})10^{-13} \text{ sec}, \tau_F = (2.8^{+1.4}_{-0.7})10^{-13} \text{ sec},^{(72)}$$

we can obtain the absolute values

$$|a'| = 2.5 \times 10^{-6}, \quad |b'| = 3.7 \times 10^{-6}, \quad |c'| = 1.7 \times 10^{-6}$$

where the phase-space factor $\{(m_c^2 - (m_v + m_p)^2)(m_c^2 - (m_v - m_p)^2)\}^{3/2}$.

$(64\pi m_c^3 m_v^2)^{-1}$ is used ((for $|a'|$ we choose the last number in Eq. (5.4). For $|c'|$ we choose $Br(D^0 \rightarrow \phi \bar{K}^0) \sim 1\%$ in Eq. (5.3)).

From comparing the mixing-matrix non suppression data of $\bar{K}^*\pi$ decays, (67)

$$\text{Br}(D^+ \rightarrow \bar{K}^{*0}\pi^+) = (3.0 \pm 1.9 \pm 1.7)\% , \quad (5.6)$$

$$\text{Br}(D^0 \rightarrow \bar{K}^{*0}\pi^0) = (2.1 \pm 0.9 \pm 0.6)\% , \quad (5.7a)$$

$$\text{Br}(D^0 \rightarrow K^{*-}\pi^+) = (7.1 \pm 1.6 \pm 1.3)\% , \text{ from } \pi^+\pi^0 K^- \text{ mode}, \quad (5.7b)$$

$$= (7.3 \pm 1.2 \pm 0.9)\% , \text{ from } \pi^+\pi^-\bar{K}^0 \text{ mode}, \quad (5.8)$$

with the quark-diagram formula (Table IIIa,b, Ref. (66))

$$A(D^+ \rightarrow \bar{K}^{*0}\pi^+) = (C_1)^2 (a'+b') e^{i\delta_{K^*\pi}^{3/2}} , \quad (5.9)$$

assuming $e^{i\delta_{K^*\pi}^{3/2}}$ elastic (a reasonable one because $\bar{K}^{*0}\pi^+$ is an exotic channel), we obtain

$$|a'+b'| \approx 1.2 \times 10^{-6} , \quad (5.10)$$

and with,

$$\frac{\text{Br}(D^0 \rightarrow \bar{K}^{*0}\pi^0)}{\text{Br}(D^+ \rightarrow \bar{K}^{*0}\pi^+)} = \left| \left\{ 1 - \frac{a'+c'}{a'+b'} - \frac{2}{3} (1 - e^{i\Delta_{K^*\pi}}) \right\} e^{-i\delta_{K^*\pi}^{1/2}} \right|^2 \quad (5.11)$$

$$\frac{\text{Br}(D^0 \rightarrow K^{*-}\pi^+)}{\text{Br}(D^+ \rightarrow \bar{K}^{*0}\pi^+)} = \left| \left\{ 1 - \frac{a'+c'}{a'+b'} - \frac{1}{3} (1 - e^{i\Delta_{K^*\pi}}) \right\} e^{-i\delta_{K^*\pi}^{1/2}} \right|^2 \quad (5.12)$$

where $\Delta_{K^*\pi} \equiv \delta_{3/2}^{K^*\pi} - \delta_{1/2}^{K^*\pi}$ (the last terms in Eq. (5.11, 5.12) is absent if there is no SU(3) breaking and final-state-interaction), assuming elastic $\delta_{1/2}^{K^*\pi}$, we obtain (for clarity only one of the solutions is given below. For details, see Ref. (66)),

$$\Delta_{K^*\pi} \approx 51^\circ , \quad (a'+c')/(a'+b') \approx 2.1, \text{ i.e. } (b'-c')/(a'+b') \approx -1.2 . \quad (5.13)$$

(Here I would like to caution about the interpretation of phase shifts obtained here in charm decays because of the competing multiple channels available, again see discussions in Ref. (66)), we obtain

$$a' = 2.5 \times 10^{-6}, b' = -3.7 \times 10^{-6}, c' = -1.7 \times 10^{-6}. \quad (5.14)$$

Note the cancellation in $a'+b'$ and also the non-negligible amount of the amplitude c' . The most interesting and exciting results here is that we already know (up to future improved accuracies) the magnitudes and relative signs of these individual amplitudes. This implies numerous predictions (you need only to read Tables III, IV of Ref (66), e.g. from Eq. (5.2) we predict that

$$\text{Br}(D^0 \rightarrow \phi \pi^0) = 1/2 \text{Br}(D^+ \rightarrow \phi \pi^+) \simeq 0.47\%. \quad (5.15)$$

Similar analyzing the data⁽⁶⁷⁾

$$\text{Br}(D^+ \rightarrow K^0 \rho^+) = (12.2 \pm 2.8 \pm 1.9)\%; \quad (5.16)$$

$$\text{Br}(D^0 \rightarrow K^- \rho^+) = (13.7 \pm 1.3 \pm 1.5)\%, \quad (5.17)$$

$$\text{Br}(D^0 \rightarrow \bar{K}^0 \rho^0) = (1.3 \pm 0.4 \pm 0.3)\%, \quad (5.18)$$

using the quark-diagram formulation (Table II.a,b, Ref. (66))

$$A(D^+ \rightarrow K^0 \rho^+) = (c_1)^2 (a+b) e^{i\delta_{3/2}^{K\rho}} \quad (5.19a)$$

assuming elastic $e^{i\delta_{3/2}^{K\rho}}$ from exoticity, we obtain

$$|a+b| \simeq 2.6 \times 10^{-6}, \quad (5.19b)$$

and from equations similar to Eqs. (5.11, 5.12), we obtain

$$\Delta_{K\rho} = 26^\circ, (a+c)/(a+b) \simeq 1.4, \text{ i.e. } (b-c)/(a+b) \simeq -0.4, \quad (5.20)$$

as one of the two solutions. From the non suppressed decay

$$\text{Br}(D^0 \rightarrow \omega K^0) = (3.8 \pm 1.5 \pm 1.0)\%,^{(67)} \quad (5.21)$$

which is proportional to $c_1(b+c)$, using Eq. (5.20), we can obtain b , c and a individually.

$$a \approx -4.4 \times 10^{-6}, \quad b \approx 1.8 \times 10^{-6}, \quad c \approx 0.8 \times 10^{-6}, \quad (5.22)$$

as one of the four solutions. From the data of singly suppressed decay,

$$\text{Br}(D^+ \rightarrow \bar{K}^{*0} K^+) = (0.53 \pm 0.24 \pm 0.14)\%, \quad (67) \quad (5.23)$$

which is proportional to $s_1 c_1 (a'-d)$, if we ignore the SU(3) symmetry violating w-loop diagrams we can obtain amplitude d ,

$$d \approx \begin{cases} 0.2 \times 10^{-6}, & \text{for } d/a' > 0, \\ 4.8 \times 10^{-6}, & \text{for } d/a' < 0. \end{cases} \quad (5.24)$$

These are two rather extreme values, we need to have more detail understanding of the SU(3) violation before we can have a definite conclusion. From $D^0 \rightarrow \bar{K}^{*0} K^0$, $K^{*0} \bar{K}^0$ we can determine the relative sign or phase of c , c' . From $F^+ \rightarrow \rho^+ \pi^0$, $F^+ \omega \pi^+$ of Eq. (5.1) we can determine the relative sign (or phases) of d , d' . We then know all the relations among amplitudes a , b , c , d , and a' , b' , c' , d' ! The rest PV decays are predictions, up to SU(3) breaking and final-state interactions! These results must be conformed by any theoretical calculations.

V.b Charm meson $\rightarrow$ pseudoscalar-pseudoscalar decays

To complete the present story, let us go to the PP decays. Following the same procedure, from ⁽⁶⁷⁾

$$\begin{aligned} \text{Br}(D^+ \rightarrow \bar{K}^0 \pi^+) &= (3.5 \pm 0.5 \pm 0.4)\%, \\ \text{Br}(D^0 \rightarrow K^- \pi^+) &= (4.9 \pm 0.4 \pm 0.4)\%, \\ \text{Br}(D^0 \rightarrow \bar{K}^0 \pi^0) &= (2.2 \pm 0.4 \pm 0.2)\%, \end{aligned} \quad (5.25)$$

we obtain

$$|a+b|_{PP} = 1.7 \times 10^{-6} \quad (5.26a)$$

$$\Delta_{\pi\pi} = 78^\circ, \quad \{ (a+c)/(a+b) \}_{PP} = 2.8; \text{ i.e., } \{ (b-c)/(a+b) \}_{PP} = -1.8, \quad (5.26b)$$

or

$$\Delta_{K\pi} = 102^\circ, \{ (a+c)/(a+b) \}_{pp} = 2.1; \text{ i.e. } \{ (b-c)/(a+b) \}_{pp} = -1.1 \quad (5.26c)$$

Unlike the PV decays, it is much harder here to determine individual amplitudes, since none of the decays are given by one amplitude. It is interesting to point out that the nonspectator-diagram amplitudes c and d can be measured in a model-independent way by observing the following decay modes

$$\frac{-\Gamma(D^0 + \bar{K}^0 \eta_8)}{\Gamma(D^0 + \bar{K}^0 \eta_0)} = \frac{1}{2} \left| \frac{b-c}{b+2c} \right|^2, \quad \frac{\Gamma(F^+ + \eta_8 \pi^+)}{\Gamma(F^+ + \eta_0 \pi^+)} = 2 \left| \frac{a-d}{a+2d} \right|^2. \quad (5.27)$$

Next we go to the mixing-matrix singly suppressed decays. The famous result of the $\Gamma(D^0 \rightarrow K^+ k^+)/\Gamma(D^0 \rightarrow \pi^+ \pi^-) \neq 1$ first observed by the Mark II collaboration,⁽⁷³⁾ is further confirmed by Mark III measurement with greater accuracy,

$$\begin{aligned} \Gamma(D^0 \rightarrow K^- \bar{K}^+) / \Gamma(D^0 \rightarrow K^- \pi^+) &= 0.122 \pm 0.018 \pm 0.012, \\ \Gamma(D^0 \rightarrow \pi^+ \pi^-) / \Gamma(D^0 \rightarrow K^- \pi^+) &= 0.033 \pm 0.010 \pm 0.006. \end{aligned} \quad (5.28)$$

As discussed in detail in Ref. (66), the measurements of Eq. (5.28) can be explained by SU(3) and final-state interactions. It is very interesting to point out that the future measurement of $D^0 \rightarrow \pi^0 \pi^0$ can clarify in detail the importance of such mechanisms. Many other interesting reactions like $D^0 \rightarrow \bar{K}^0 \eta_8$, $\pi^0 \eta_8$, should also be measured to test these mechanisms. It is also interesting to compare future measurements of $F^+ \rightarrow \pi^+ \eta_8$ with the present measurement,

$$\text{Br}(D^+ \rightarrow \bar{K}^0 K^+) = 1.11 \pm 0.34 \pm 0.21, \quad (5.29)$$

because both are given by amplitude $(a-d)$ if there is no SU(3) breaking. Also $D^0 \rightarrow K^0 \bar{K}^0$, $\eta_0 \eta_0$ can only come from SU(3) breaking. Their observations give clear measures of the amount of SU(3) in the quark-diagram amplitudes. For many interesting results and detail analysis, you are referred to Ref. (66).

I hope that I have demonstrated to you that the general quark-diagram approach provides a framework in which experimental results can be analyzed in a model-independent way, and new experiments can be pointed out to further test certain specific model calculations.⁽⁷⁴⁾ The current experimental results have already provided much information on non-leptonic decay mechanism in term of the quark diagram amplitudes, which can be used for comparison with theoretical calculations. Many interesting predictions have resulted. The story of non-leptonic decay is very complex, but also very interesting. It will take our persistent effort, both theoretically and experimentally, to find the conclusion of the story.

As we see the experiments measuring the b lifetime, V_{ub}/V_{cb} , ϵ , ϵ'/ϵ , and the hint of m_t from UA1 experiment at CERN have really pushed the three-generation model to a corner. Future measurements of ϵ'/ϵ , $|V_{ub}/V_{cb}|$ with improved error bars, and with m_t pinpointed can really imply very specifically about B_K , B_K' , and the likelihood of the three generation model. So it's a good time to think ahead if there are four generations of quarks.

If there are fourth generation of quarks b' , t' , the number of parameters increase from the three generation 4 to 9, six angles and three phases. We generalize the parameterization of Eq. (1.3) as follows, ⁽³⁶⁾

where V_3 is the 3x3 matrix given in Eq. (1.3). The reason we put the additional phases ϕ_2, ϕ_3 where there are in Eq. (6.1) is to anticipate that the widening of the generation gap will keep its pace. Of course physical consequences are independent of ways of parameterization. What are the different physical phenomena if fourth generation do exist?

— (1) If the masses are low enough, we should search for them in e^+e^- , $p\bar{p}$, pp and cosmic rays.

(2) $|V_{cd}^*/V_{cs}| \approx |V_{us}^*/V_{ud}|$ does not have to be true. More accurate measurements of $|V_{cd}/V_{cs}|$ should be studied from semileptonic decays of $D \rightarrow \ell^\pm X$ containing s or not, and from the long predicted and $\Gamma(D^+ \rightarrow \pi^+ \pi^0)/\Gamma(D^+ \rightarrow \bar{K}^0 \pi^+) = 1/2 |V_{cd}/V_{cs}|^2$.

(3) As mentioned in Sect. I, we showed a surprising result in Ref. (36), that rather than the unique phase-convention-invariant CP violation parameter X_{CP} in the three-generation case, there are now nine of them $X_{CP,i}$, $i = 1, \dots, 9$, for generation number $n = 4$, and always saturating the whole parameter space for $n \geq 4$. So the consequences of CP violations are much more varied than the three-generation case, e.g. ϵ'/ϵ can be zero without making $B_K' = 0$; V_{ub} can be zero without annihilating the CP-violation effect ϵ . Due to the presence of more $X_{CP,i}$ variables, many more appreciable partial rate differences can happen. Those effects from the three graphs are actually only dependent on the presence of the higher generation, not on their masses. For example the partial-decay-rate difference Δ_F in mixing-matrix singly suppressed decay $F \rightarrow K^0 \pi$ can be increased to 10^{-1} , and Δ_T for t quark decays can be very large too. This makes the improvement of SPEAR experiment, and BEPC extremely important. For details, see Ref. (36).

(4) There are new contributions in the mass matrix from the new generation of b' , t' , so that the mass-matrix mixing (i.e. the neutral meson-antimeson mixing results from the three-generation model can be drastically changed. However such effects are from loop diagrams and very sensitive to the b' , t' quark masses. As shown in Ref. (75). If one picks suitable values of the quark-mixing matrix and masses of the new quarks, e.g. $D^0, \bar{D}^0$ mixing can be increased from $<10^{-4}$ to 10^{-2} if $m_b, \gtrsim 200$ Gev; $B_d^0, \bar{B}_d^0$ can also be appreciable; the mass-matrix CP violation in D , B mesons can also be large so that $(\ell^+ \ell^+ - \ell^- \ell^-) / (\ell^+ \ell^+ + \ell^- \ell^-)$, $(KK - \bar{K}\bar{K}) / (KK + \bar{K}\bar{K})$ are reasonably large. Such possibilities are extremely interesting to search for experimentally.

VII. Quark Mass Matrix

So far in the electroweak unified gauge theory, the most obscure part is the mass generating mechanism for gauge particles and for quarks. Here I would like to reiterate, agreeing with what Dr. Kobayashi had emphasized in his talk, the intimate relations between the quark mass matrix M' in the bases of quark eigen-state in weak interaction and the mixing matrix V . The mass term in the interaction Lagrangian can be written in the strong-interaction eigen quark state u , d , or equivalently the weak-interaction eigenstate u , d' ,

$$\begin{aligned} \text{mass} &= \bar{u}_L M(2/3) u_R + \bar{d}_L M(-1/3) d_R + \text{H.C.}, \\ &= \bar{u}_L M(2/3) u_R + \bar{d}_L' M'(-1/3) d_R' + \text{H.C.}, \end{aligned} \quad (7.1)$$

where $\bar{u}_L' M'(-1/3) V_R^+ = M(-1/3)$, and $V_R^+ V_R = 1 = V_L^+ V_L$.

We can show that $M'(-1/3)$ can always either chosen to be Hermitean or symmetric: ⁽⁷⁶⁾

$$\text{From} \quad M'(-1/3) = V_L^+ M(-1/3) V_R, \quad (7.2)$$

we multiply an unitary matrix W on the right hand side of Eq. (7.2),

$$M'' \equiv M' W = V_L^+ M V_R W. \quad (7.3)$$

If we require M'' to be Hermitean $M''^+ = M''$ i.e. $W^+ V_R^+ M V_L = V_L^+ M V_R W$ or $V_L W^+ V_R^+ M = M V_R W V_L^+$. A sufficient solution for W is $V_R W V_L^+ = 1$, i.e. $W = V_R^+ V_L$, and $M'' = V_L^+ M V_L$; or if we want M'' to be symmetric $M''^T = M''$, i.e. $W^T V_R^T M V_L^+ = V_L^+ M V_R W$. A sufficient solution for W is $V_R W V_L^T = 1$, i.e. $W = V_R^+ V_L^*$, and $M'' = V_L^+ M V_L^* = V_L^+ M V_L^{+T}$. All this says that we can choose a base such that $M'(-1/3)$ is either Hermitean or symmetric. Now the charged weak current is

$$J_\mu = \bar{u}_L \gamma_\mu d_L' = \bar{u}_L \gamma_\mu V d_L. \quad (7.4)$$

This V is the quark mixing matrix. So in the d_L' base the charge $-(-1/3)$ mass-matrix is

$$M_H'(-1/3) = V^+ M(-1/3) V, \quad \text{if in the Hermitian base} \quad (7.5)$$

$$M_S'(-1/3) = V^T M(-1/3) V, \quad \text{if in the symmetric base;} \quad (7.6)$$

using the V matrix given in Eq. (1.3), we obtain $M_H'(-1/3)$ and $M_S'(-1/3)$.

These are the mass matrices the weak decay phenomenology has told us. They should be conformed by model building.

We can see that the quark mixing matrix and its complex phase, the source of CP violation, is intimately related to the quark mass matrix in the base of weak-interaction quarks. Thus the mass-generating sector, i.e. the Higgs sector, is truly now the reservoir of our ignorance in the gauge theory description of the particle world. It is of great importance to make progress on it.

VIII. Concluding Remarks and Outlook

As we can see that there are many interesting experiments can be done to shed light on our understanding of the physics of heavy-quark decays and CP violation. Here I list them briefly.

VIII.a. The Measurements of the b Particle Lifetime and $\Gamma(b \rightarrow u)/\Gamma(b \rightarrow c)$ have well constrained the quark mixing matrix. It is important to further study the quark mixing matrix elements: V_{ub} via $B \rightarrow TV_\tau$, and the charmless b decays; V_{cs} , V_{cd} via $D \rightarrow l X_{s,d}$. The fitting of the CP-violation effects in K decays, ϵ and ϵ'/ϵ , has cornered the model with three generations of quarks, and has constrained the nonleptonic dynamical parameter B_K , B'_K , and the t quark mass. If V_{ub} becomes smaller than needed for CP violating effects, it's an indication of trouble with the KM scheme for CP violation, or of the exciting possibility of the existence of higher than three generations of quarks.

VIII.b. There are still many important and interesting CP violating effects to be measured in the kaon and hyperon decays: ϵ'/ϵ , ϕ_{00} ; η_{+-0} ; $\text{Br}(K_S^0 \rightarrow \gamma\gamma)$; R_+/R_- , where $R_\pm = \Gamma(K^\pm \rightarrow \pi^\pm \pi^\pm \pi^\mp)/\Gamma(K^\pm \rightarrow \pi^\pm \pi^0 \pi^0)$; R_Λ/R_Σ , where $R_\Lambda \equiv \Gamma(\Lambda \rightarrow \pi^- p)/\Gamma(\Lambda \rightarrow \pi^0 n)$, $R_\Sigma = \Gamma(\Sigma \rightarrow \pi^+ \bar{p})/\Gamma(\Sigma \rightarrow \pi^0 \bar{n})$; Pion momentum asymmetry difference in $\Sigma^\pm \rightarrow \pi^\pm n$ decays. The tagged $K^0, \bar{K}^0$ experiments for partial rate difference $\Delta(K^0, \bar{K}^0 \rightarrow 2\pi)$, $\Delta(K^0, \bar{K}^0 \rightarrow \gamma\gamma)$ are very interesting and will mark the beginning of this new type of CP experiments.

VIII.c. Some partial decay rate differences in beauty particle decays can be very large, such that only some tens of restricted beauty decays are needed:

$$\Delta(B_u^\pm \rightarrow D^\pm D^{(-)*0}), \Delta(B_d^{(-)0} \rightarrow K^\mp \pi^\pm), \Delta(B_s^{(-)0} \rightarrow D^\pm D^\mp), \Delta(B_d^{(-)0} \rightarrow \pi^\pm \pi^\mp), \\ \Delta(B_c^\pm \rightarrow \pi^\pm D^{(-)0}), \text{ etc. See Table in Section IIIb, and Refs. (54, 55).}$$

In the three-generation case, such partial decay rate differences for charm and top quarks are in general an order of magnitude smaller. This information can serve as a base for looking for hints of the existence of higher generations of quarks.

VIII.d Look for the $B_S^0, \bar{B}_S^0$ mixing effects, e.g. same-sign dileptons, which has been consistently predicted to be substantial in the case of three generations of quarks while mixing in $B_d^0, \bar{B}_d^0$ is small and sensitive to parameters, and mixing in $D^0, \bar{D}^0$ and $T^0, \bar{T}^0$ are extremely small.

VIII.e. Experiments tagging neutral mesons can provide new ways to study CP-violation effects that the conventional interference experiments can not do, e.g. measuring decay-amplitude CP violation effects in $K_L \rightarrow \gamma\gamma$; by measuring $\Delta(K^0, \bar{K}^0 \rightarrow \gamma\gamma)$; separately measuring the neutral meson $P^0, \bar{P}^0$ (mass-matrix) mixing parameters δm , δT , and the mass-matrix CP violation $\eta_p \equiv |1 - \bar{\epsilon}_p| / |1 - \epsilon_p|$ for short lived heavy quark systems.

VIII.f. Experiments should be planned to look for indications of higher than three generations of quark: the most important are the direct observations of the top particles, and the heavier ones; Since the phenomenological consequences in the case of three generations of quarks have been quite well determined, any observation of substantial deviations from them are possible indication of the existence of higher generations, e.g. large partial decay rates in charm and top quark decays, in addition to the beauty particle decays as which are the only ones calculated to be substantial in the three-generation case; large $D^0, \bar{D}^0, B_d^0, \bar{B}_d^0$ mixing in addition to $B_S^0, \bar{B}_S^0$ which is the only one calculated to be substantial in the three-generation case; large mass-matrix CP-violation in B^0, D^0 decays, e.g. $\Delta(\ell^+ \ell^+ - \ell^- \ell^-) \neq 0$; "long" lived top quark ($\tau_t \gtrsim 10^{-18}$ sec).

VIII.g. The well determination of the quark mixing matrix also helps for a systematic study on nonleptonic decays via the many channels available in charm and beauty particle decays. Recent charm decay data analyzed via the quark-diagram scheme already have shed some light on the structure of charm

decays. The quark-diagram amplitudes for charm meson $\rightarrow$ pseudoscalar-vector-meson decays are quite well determined in magnitudes and relative signs. This leads to many predictions, e.g. $\text{Br}(D^0 \rightarrow \phi \pi^0) \simeq 0.47\%$; future measurements of $D^0 \rightarrow \bar{K}^{*0} K^0$, $K^{*0} \bar{K}^0$ can shed light on amplitude $c'-c$; $F^+ \rightarrow \rho^+ \pi^0$, $\rho^0 \pi^+$ can give information on amplitude $d-d'$; $F^+ \rightarrow \omega \pi^+$ can give information on $d+d'$; and many more. For charm meson + pseudoscalar-pseudoscalar meson decays, measurements of the following few charm decays will be most helpful: $(D^+ \rightarrow \pi^+ \pi^0)$, so that the long predicted relation $\Gamma(D^+ \rightarrow \pi^+ \pi^0) / \Gamma(D^+ \rightarrow \bar{K}^0 \pi^+) = 1/2 |V_{cd}/V_{cs}|^2$ can be checked. Measurements on $D^0 \rightarrow \bar{K}^0 \eta'$ in addition to the recently measured $D^0 \rightarrow \bar{K}^0 \eta$ can test the importance of W-exchange amplitude c ; the measurement of $D^0 \rightarrow \pi^0 \pi^0$ can help to understand in details of the mechanism for the ratio $\Gamma(D^0 \rightarrow K^+ K^-) / \Gamma(D^0 \rightarrow \pi^+ \pi^-) \neq 1$; the observation of $D^0 \rightarrow K^0 \bar{K}^0$, $\eta_0 \eta_0$ can give a clear indication and measurements of SU(3) breaking.

VIII.h. Measurement of F , $D^+ \nu_\tau$, $\mu \nu_\mu$ can help to determine g_τ/g_μ and the ν_τ mass. (77)

VIII.i. Measurements of rare decays can put standard models to stringent tests, and provide windows to glimpse at possible new physics. The case of $K_L \rightarrow \mu^+ \mu^-$ ($\text{Br} = 9.1 \times 10^{-9}$) was such an example. Now there are four rare decay experiments being carried out at Brookhaven National Laboratory: (78)

Exp. 777, BNL - U. of Washington-Yale Collaboration,

$K^+ \rightarrow \pi^+ e^- \mu^+$, sensitivity $\text{Br} \sim 10^{-12}$,

Exp. 780, BNL-Yale Collaboration,

$K_L \rightarrow \mu e$, sensitivity $\text{Br} \sim 10^{-10}$,

Exp. 787, BNL-Carnegie-Mellon-Columbia-Princeton-TRILIMF Collaboration,

$K^+ \rightarrow \pi^+ \nu \bar{\nu}$, $\pi^+ " ? "$, sensitivity $\text{Br} \sim 10^{-10}$,

Exp. 791, UCLA-Los Alamos-U. of Pennsylvania-Princeton-Stanford-Temple

Collaboration,

$K_L \rightarrow \mu e$, Sensitivity 10^{-12} ,

$K_L \rightarrow \pi^0 e^+ e^-$, Sensitivity 10^{-12} ,

μ polarization $K_L \rightarrow \mu^+ \mu^-$, sensitivity $10 \sim 20\%$.

Further tightening the already very impressive bound on the neutron electric dipole moment will help to test various CP-violating models.

We can see that despite the milestone of the observation of the long-awaited intermediate bosons, there are still many fundamental questions in electroweak interactions to be answered: how the masses of gauge bosons and quarks are generated? What is the reason for the V-A nature of the weak current? Where is the origin of CP-noninvariance? Why do the quarks and leptons mix and repeat themselves? How many more are there? The apparent complexity of my discussions presented here reflects the immaturity of the field. However, this is the very nature of our pursuit: once a good question is answered, more new meaningful questions can be asked. There will be frontier research as long as there is life itself. I look forward to Meson 60 to see what the milestones will be erected and what the new frontiers will be.

REFERENCES

1. E. Fermi, Zeit. f. Phys. 88, (1934) 161.
2. H. Yukawa, Proc. Phys. Math. Soc., Japan 19, (1935) 48.
3. C.N. Yang and R.L. Mills, Phys. Rev. 96, (1954) 191.
4. S.L. Glashow, Nucl. Phys. 22, (1961) 579; S. Weinberg, Phys. Rev. Lett. 19, (1967) 1264; A. Salam in: "Elementary Particle Theory", N. Svartholm, ed., Almquist and Wiksell, Stockholm (1968); see also A. Salam and J.C. Ward, Phys. Lett. 13, (1964) 168. For renormalization of the theory see G.'t Hooft, Nucl. Phys. B33, (1963) 173.
5. UA1 collaboration, CERN, G. Arnison, et al., Phys. Lett. 122B, (1983) 103, ibid. 126B, (1983) 398; UA2 collaboration, CERN, M. Banner, et al., ibid. 122B, (1983) 476. Talk by C. Rubbia and L. Dilella at 1985 Int'l. Conf. on Lepton Photon Conf. at Kyoto, Japan, Aug. 1985.
6. See review talk by W.J. Marciano, Proceedings of the 1983 Int'l. Sym. on Lepton and Photon Interactions at High Energies, Cornell U., Aug. 4-9, 1983.
7. R.F. Peierls, T.L. Trueman and L-L. (Chau) Wang, Phys. Rev. D16, (1977) 1397. See also L-L. (Chau) Wang Theoretical Implications of ISABELLE Physics, in Proc. Int'l. School, The High Energy Limit, Erice, Italy, July 31-August 13, 1980; Plenum, Ed. A. Zickichi.
8. W. Pauli, in open letter dated Dec. 4, 1930, now reproduced in: "Collected Scientific Papers by W. Pauli", R. Kronig and V.F. Weisskopf, eds., Interscience, New York (1964).
9. Neutrino observation, C.L. Cowan, F. Reines, and F.B. Harrison, Phys. Rev. 96, (1954) 1294.
10. Ig. Tamm, Nature 133, 931 (1934); D. Iwanenko, ibid. 981 (1934).
11. The discovery of the muon, C.D. Anderson, S.H. Neddermeyer, Phys. Rev. 51, (1937) 894. The discovery of charged pions, C.M.J. Lattes, H. Muirhead, J.P.S. Occhialini, C.F. Powell, Nature 159 (1947) 694.
12. R. Dalitz, Phil. Mag. 44, (1953) 106B. E. Fabri, Nuovo Cimento II, (1954) 479.
13. T.D. Lee and C.N. Yang, Phys. Rev. 104, (1956) 254.
14. Wu, C.S., Ambler, E., Hayward, R.W., Hoppes, D.D., and Hudson, R.P., Phys. Rev. 105, (1957) 1413; Garwin, R.L., Lederman, L.M., and Weinrich, M., Phys. Rev. 105, (1957) 1415; Friedman, J.I., and Telegdi, V.L., Phys. Rev. 105, (1957) 1681.

15. Neutral current observation, F. Hasert et al., Phys. Lett. 46B, (1973) 138; Nucl. Phys. B73, (1974) 1.
16. N. Cabibbo, Phys. Rev. Lett. 10 (1963) 531; for earlier work on the subject, see M. Gell-Mann and M. Levy, Nuovo Cim. 16 (1960) 705.
17. Particle Data Group, Rev. Mod. Phys. 52 (1980) 5. Shocket et. al. Phys. Rev. Lett. 39 (1979) 59.
18. S. Glashow, J. Iliopoulos and L. Maiani, Phys. Rev. D2, (1970) 1285.
19. For implicit charm J/ψ , J.J. Aubert, et al., Phys. Rev. Lett. 33, (1974), 235; J.E. Augustin, et al., Phys. Rev. Lett. 33, (1974) 1406. For charm particle observation in V_{up} reaction in the BNL 7-foot bubble chamber, see E.G. Cazzoli, et al., Phys. Rev. Lett. 34, (1975) 1125; in e^+e^- reaction, see G. Goldhaber, et al., Phys. Rev. Lett. 37, (1976) 255, in pp reaction, see M. Basile, et al., Nuovo Cim. 63A, (1981) 230; for hint from earlier cosmic ray experiment see, K. Niu, E. Mikumo, Y. Maeda, Prog. Theor. Phys., 46 (1971) 1644.
20. M. Kobayashi and T. Maskawa, Prog. Theor. Phys. 49, (1973) 652.
21. For implicit beauty T, S.W. Herb et al., Phys. Rev. Lett. 39, (1977) 252, for explicit beauty particle observation, S. Behrends et al., Phys. Rev. Lett. 50 (1983), 881, and talk by P. Avery in Ref. 22.
22. For recent development on the subject, see talks in Proceedings of "Flavor Mixing in Weak Interactions," Europhysics Conference, March 5-10, 1984, Erice, Italy, Ed. L.-L. Chau, Plenum 1985.
23. For a general survey, see L.L. Chau, Phys. Rept. 95 (1983) 1., and
24. L.L. Chau, "Comments on Heavy Quark Decays and CP Violation," Proceedings of the G.F. Chew Jubilee (Sept. 29, 1984), to be published by the World Scientific Pub. Co.
25. For earlier fits to find V_{ud} , V_{us} , R. Shrock, and L.L. (Chau) Wang, Phys. Rev. Lett. 41 (1978) 1792; more recent fits, see the next reference.
26. J.F. Donoghue and B.R. Holstein, Phys. Rev. D25 (1982) 2015; A. Garcia and P. Kielanowski, Phys. Lett. 110B (1982) 498; and the most recent fits, WA2 experiment at CERN, M. Bourquin et al., "IV. Tests of the Cabibbo Model," CERN preprint (1983) see talk by H.W. Siebert in Ref. 22; J.F. Donoghue, B.R. Holstein, Phys. Lett. 160B (1985) 173; W.J. Marciano and A. Sirlin, BNL and NYU preprint (1985) found $V_{ud} = 0.9729 \pm 0.0012$; and A. Bohm, M. Kmicik, Phys. Rev. D31, 3005 (1985) include new Fermilab hyperon data, Ref. 30, and found $|V_{us}| = 0.225 \pm 0.002$ in better agreement with the older K_s3 fit of Ref. 25.
27. For the b lifetime, E. Fernandez, Phys. Rev. Lett. 51 (1983) 1022; N.S. Lockyer et al., Phys. Rev. Lett. 51 (1983) 1316; for more up to data information see Talks by W.T. Ford, and G.H. Trilling in Ref. 22.

28. For $\Gamma(b \rightarrow u)/\Gamma(b \rightarrow c)$, C. Klopfenstein et al., Phys. Lett. 103B, 444 (1983); A. Chen et al., Phys. Rev. Lett. 111B, 1084 (1984); and Talks by J. Lee-Franzani, and P. Avery in Ref. 22.
29. L.-L. Chau, W.-Y. Keung, M.D. Tran, Phys. Rev. D27, 2145 (1983), L.-L. Chau and W.-Y. Keung, Phys. Rev. D29, 592 (1984).
30. Fermilab Polarized Σ^- experiment, S.Y. Hsueh et al., Phys. Rev. Lett. 54, (1985), 2399.
31. See H.W. Siebert's talk in Ref. (22).
32. L. Wolfenstein, Phys. Rev. Lett. 51, (1984) 1945.
33. L.-L. Chau and W.-Y. Keung, Phys. Rev. Lett. 53, (1984) 1802.
34. L. Maiani, Phys. Lett. 62B (1976) 183; Lett. Al Nuovo Cimento, 28 (1980), 29.
35. H. Fritzsch, Max-Planck Institute preprint, MPI-PAE/PTh 42/85.
36. F. Botella and L.-L. Chau, "Anticipating the Higher Generations of Quarks from Rephasing Invariance of the Mixing Matrix," Brookhaven preprint (1985).
37. M.K. Gaillard and B.W. Lee, Phys. Rev. D10 (1974) 897.
38. See L.-L. Chau, H.-Y. Cheng and W.-Y. Keung, "A KM Model Study for ϵ , ϵ/ϵ' , and M_K ", BNL preprint BNL-35163, July '84 (unpublished); and Phys. Rev. 32, 1837 (1985); the uncertainties in the B_K parameter and the numerical findings of these papers were presented by L.-L. Chau in a talk immediately following B. Winstein at the APS Meetings at Washington, DC, 23-26, April '84. J. Bijnens, H. Sonoda and M.B. Wise, Phys. Rev. Lett. 53, (1984) 2367. A. Pich and E. de Rafael, Phys. Lett. 158B, (1985) 477.
39. J.H. Christenson, J.W. Cronin, V.L. Fitch and R. Turlay, Phys. Rev. Lett. 13, 138 (1964); and the classical theoretical discussions on CP violation. T.D. Lee, R. Oehme and C.N. Yang, Phys. Rev. 106, 340 (1957); T.T. Wu and C.N. Yang, Phys. Lett. 13, 380 (1964).
40. L.-L. Chau, "Comments on CP Violation" Brookhaven preprint, (1984).
41. J.K. Black et al., Phys. Rev. Lett. 54, (1985) 1628.
42. R.H. Bernstein et al. *ibid*, 54, (1985) 1631.
43. F.J. Gilman and M.B. Wise, Phys. Lett. 83B, 83 (1979).
44. L.-L. Chau, H.-Y. Cheng, W.-Y. Keung, "CP Violation in the Kaon Systems", Brookhaven preprint 1985.

45. The current experiments on ϵ'/ϵ : Chicago -Fermilab -Orsay-Princeton collaboration, Fermilab experiment #631; CERN-Dortmund-Edinburgh-Orsay-Pisa-Siegen collaborations, CERN NA31.
46. For discussion partial rate differences in K decays see Sect. 3.2.2 of Ref. (23); L.-L. Chau, talk at Erice Conference on "Electroweak Effects at High Energies," Feb. 1-12 '83; C. Kounnas, A.B. Lahanas and P. Pavlopoulos, Phys. Lett. 127B, 381 (1983):
47. "Physics at LEAR with Low-Energy Cooled Antiprotons," edited by U. Gastaldi and R. Klapisch, Ettore Majorana Int. Sci. Series, ed. A. Zichichi (Plenum); P. Pavlopoulos, Talk in Ref. (22); tagged K^0 experiment: Athen-Basel-ETH-Zurich-Liverpool-Saclay-Sin collaboration, CEAR, LEAR, p82.
48. L.-L. Chau, H.-Y. Cheng, Phys. Rev. Lett. 54, (1985) 1763.
49. L.-L. Chau Wang, AIP Conf. Proc. No. 72, Particle and Fields, Subseries No. 23, Virginia Polytechnic Inst. 1980, eds. G.B. Collins. L.N. Chang and J.R. Ficenec; L.-L. Chau, Proceedings of Workshop on Weak Interactions and Neutrinos, Javea, Spain, Sept. 5-11, 1983.
50. C. Avilez, Phys. Rev. D23, (1981) 1124. See also B. Grinstein, S.-J. Rey and M.B. Wise, CALT-68-1286 (1985).
51. L.-L. Chau and H.Y. Cheng, Phys. Lett. 131B, (1983) 202; T. Brown, S.F. Tuan and S. Pakvasa, Phys. Rev. Lett. 51, (1983) 1823. I would like to thank Dr. K. Kilian for illuminating discussion on the possibility of doing this experiment at LEAR.
52. For a review of CP violation in the K system see, K. Kleinknecht, Ann. Rev. Nucl. Sci. 26, 26 (1976).
53. L.M. Sehgal, and L. Wolfenstein, Phys. Rev. 162 (1976) 1362, O.E. Overseth and S. Pakvasa, Phys. Rev. 189 (1969) 1663. A. Pais and S.B. Treiman, Phys. Rev. D12, 2744 (1975); L.B. Okun, V.I. Zakharov and B.M. Pontecorvo. Lett. Nuovo Cim. 13, 218 (1975). J. Barshay and J. Geris, Phys. Lett. 84B, (1979), 319; M. Bander, D. Silverman and A. Soni, Phys. Rev. Lett. 43, (1979) 242; J. Barnabeu and C. Jarlskog, Z. Phys. C8, (1981) 233. I.I. Bigi and A.I. Sanda, Nucl. Phys. B193, (1981) 81. L. Wolfenstein, Nucl. Phys. B200 (1984), 45 and talk by K.-C. Chou in Ref. (22).
54. L.-L. Chau H.-Y. Cheng, Phys. Rev. Lett. 53, (1984) 1037.
55. L.-L. Chau, H.-Y. Cheng, "CP Violation in Decay Rates of Neutral B Mesons", Brookhaven preprint, BNL 36915 (1985), to appear in Phys. Lett.
56. For left-right symmetric theory, see talk by R.N. Mohapatra in Ref. (22).
57. See, e.g. S. Weinberg, Phys. Rev. Lett. 37, (1976) 657, Physica 96A, (1979) 327.

58. A.I. Sanda, Phys. Rev. D23 (1981) 2647; N.G. Deshpande, ibid D23 (1981) 2654.
59. Y. Dupont and T.N. Pham., Phys. Rev. D28, (1983) 2169. J.F. Donoghue and B.R. Holstein, UMHEP-213 (1984). H.-Y. Cheng, "Weinberg CP Violation Model Revisited", Brandeis preprint (1984).
60. A.A. Anselm, V.E. Bunakov, V.P. Gudkov, and N.G. Uraltsev, Leningrad preprint (1984).
61. N.F. Ramsey, Rep. Prog. Phys. 45, (1982) 95; J.M. Pendlebury et al., Phys. Lett. 136B, (1984) 327; I.S. Alterev et al., ibid 102B, (1981) 13; and B. Heckel's talk in Ref. (22).
62. Observation of $K^+ \rightarrow \pi^+ \pi^0$ suppression R.W. Birge et al., Nuovo cimento 4 (1956) 834; G. Alexander et al., Nuovo cimento 6 (1957) 478, for theoretical discussions, see M. Gell-Mann and A. Pais, Proc. Int'l Conf. of High Energy Physics, Glasgow (Pergamon Press, London 1955); Proc. Fifth Annual Rochester Conf. on High Energy Physics (NY 1955) p. 136;
63. M.A. Shifman, A.I. Vainshtein, V.I. Zakharov and Nucl. Phys. B120, 316 (1977); Sov. Phys. JETP 45 (1977) 670.
64. L.-L. Chau, talk in the Proceedings of the 1980 Quanzhou Conference (Jan. 5-10, 1980), Science Press, Beijing, Reinhold Comp., China; Van Nostr.
65. For charm $\rightarrow$ PP, see Table 2.5, Ref. (23), and footnote (2) of Ref. (24); For Charm $\rightarrow$ PV see Table 2.6, Ref. (23) and M. Gorn, Nucl. Phys. B191, (1981). X.-Y. Li, S.-F. Tuan (to be submitted to Zeits für Physik C). For treatment of SU (3) breaking and final-state-interaction, see next reference.
66. For charm $\rightarrow$ PP, PV including SU(3) breaking and final-state-interaction effects, see L.-L. Chau and H.-Y. Cheng "Exclusive Two Body Decays of Charm Meson", Brookhaven-Brandes preprint 1985; and L.-L. Chau and H.-Y. Cheng, "Quark Diagram Analysis of Charm Meson Decays", Brookhaven-Indiana preprint 1985.
67. MARK III collaboration, R. Baltrusaitis, Phys. Rev. Lett. 55, (1985) 150. R.H. Schindler, "New Results on Charmed D Meson Decay", invited talk at the 1985 SLAC Summer Institute for Particle Physics.
68. CLEO collaboration, A. Chen et al., Phys. Rev. Lett. 51, (1983) 634; see also talk by S. Stone, Int. Sym. on Lepton and Photon, Interaction at High Energies, Cornell July 83; and paper contributed to the 1985 Int'l. Lepton and photon conference at Kyoto, Japan, Aug. '85.
69. ARGUS collaboration C. Darden et al., DESY preprint 84-04-3, H. Albrecht et. al., Phys. Lett. 158B (1985) 525.
70. TASSO collaboration, M. Althoff et al., Phys. Lett. 136B, (1984) 130.

71. HRS collaboration, M. Derrick et al., Phys. Rev. Lett. 54, (1985) 2568.
72. For a review on charm lifetime see talk by K. Niu in Ref. (22). I would like to thank Professor N. Reay for very informative discussions on the subject.
73. R. Schindler et al., Phys. Rev. D24, (1981) 78. Talk by G. Goldhaber at 18th Moriond Conference, March 13-19, 1983.
74. For current comparisons with various model calculations, see discussions in Ref. (66); and for other ways of analyzing the data and comparison with calculations. See the rather exhaustive references given there in, and talk by R. Rückel in Ref. (22), and talk by B. Stech at Moriond Conference, Spring, 1985.
75. X.-G. He and S. Pakvasa Phys. Lett. 156B, (1985) 236. A.A. Anselm et al., Phys. Lett. 156B, (1985) 102. M. Gronau and J. Schechter Phys. Rev. D31, (1985) 1668. U. Türke et. al., DO-TH 84/26. I.I. Bigi PITRA 84/19.
76. P.H. Frampton and C. Jarlslog Phys. Lett. 154B (1985) 421, R.N. Mohapatra, Private communication.
77. Chao Kuang-ta, Chau Ling-Lie, Huang Tao, Du Dong-Sheng, Wu Dan-di, Chinese Phys. Lett. 2, (1985) 117.
78. See Talk by L. Littenberg in Ref. (22).