3/9/95 (2)
PREPARED FOR THE U.S. DEPARTMENT OF ENERGY,
UNDER CONTRACT DE-AC02-76-CHO-3073

PPPL-2787
UC-427

PPPL-2787

NONLOCAL HEAT TRANSPORT IN
A STOCHASTIC MAGNETIC FIELD

BY

J.M. RAX AND R.B. WHITE

December 1991

PPPL

PRINCETON
PLASMA PHYSICS
LABORATORY

PRINCETON UNIVERSITY, PRINCETON, NEW JERSEY

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

NOTICE

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial produce, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

NOTICE

This report has been reproduced directly from the best available copy.

Available to DOE and DOE contractors from the:

Office of Scientific and Technical Information
P.O. Box 62
Oak Ridge, TN 37831;
Prices available from (615) 576-8401.

Available to the public from the:

National Technical Information Service
U.S. Department of Commerce
5285 Port Royal Road
Springfield, Virginia 22161
703-487-4650

NONLOCAL HEAT TRANSPORT IN A STOCHASTIC MAGNETIC FIELD

J.M. Rax[†] and R.B. White

Princeton Plasma Physics Laboratory, Princeton University, Princeton N.J. 08543

Abstract

Heat transport in a stochastic magnetic field configuration is shown to be nonlocal. Collisional transport processes, in such a disordered media, cannot always be reduced to a standard diffusion process, and the concept of a diffusion coefficient is meaningless for a wide range of typical tokamak parameters. In the nonlocal regime the relaxation of a gradient is described by an integral equation, involving a nonlocal propagator. This propagator is calculated, and the relation to previous results is elucidated.

[†]Permanent adress: Association EURATOM-CEA sur la Fusion Contrôlée, 13108 Saint-Paul-lez-Durance, France.

MASTER

Sp

A three dimensional toroidal magnetic field is a Hamiltonian system, and the existence of magnetic surfaces in an axisymmetric configuration is a consequence of the Kolmogorov-Arnold-Moser stability theorem. The occurrence of small resonant magnetic field perturbations can lead to the onset of chaotic field line diffusion,¹ provided that the Chirikov criterion is fulfilled. This deconfinement mechanism has been recognized as a candidate to explain anomalous confinement properties of Tokamak discharges,²⁻⁴ and can be used to control the edge characteristics with ergodic divertor configurations.⁵

This letter addresses the problem of heat transport in such random fields (the results also apply to particle transport in the strong collisionality regime), and stresses the fact that models based on local diffusion cannot be used in a wide range of typical tokamak parameters. Nonlocal responses in plasma physics have already been used to study heat transport during laser plasma interaction,⁶ current transport in Reversed Field Pinch configurations,⁷ and in wave driven Tokamaks.⁸ Nonlocality in a braided magnetic field was noted by Stix³ but there was no attempt to calculate the anomalous mean squared displacement, and to relate it to a nonlocal propagator.

On the basis of a Lagrangian formulation, with Langevin equations, we show that transport cannot always be reduced to a diffusion. The relevant theoretical framework needed to describe the collisional regime in a stochastic magnetic field is an integral rather than a differential equation. The nonlocal propagator of this integral equation is calculated with the help of a Wiener functional integral over the random magnetic field. Finally a direct numerical simulation is performed to confirm the analytical results.

The most general, (linear, causal) transport theory of a passive scalar, $T(\mathbf{r},t)$ (temperature or density), in a statistically homogeneous media, is nonlocal and relates T at point $\mathbf{r}$, time t , to the source, $S(\mathbf{r}',t')$, elsewhere before t , through a propagator P :

$$T(\mathbf{r},t) = \int d\mathbf{r}' dt' P(\mathbf{r}-\mathbf{r}', t-t') S(\mathbf{r}',t'). \quad (1)$$

P is the fundamental quantity arising through first principles, the diffusion coefficient is not fundamental and may or may not exist. Let us consider the long distance behavior of

the Fourier transform, $P(\mathbf{k}, \omega)$, of $P(\mathbf{r}, t)$. If its asymptotic form for large $\mathbf{r}$ and t , i.e., near $\mathbf{k} \approx \mathbf{0}$, and $\omega \approx 0$, is

$$P^{-1}(\mathbf{k}, \omega) \sim i\omega + \mathbf{k} \cdot \mathbf{D} \cdot \mathbf{k} + \dots, \quad (2)$$

then the transport becomes local and diffusive on large space and time scales. T obeys the usual differential equation, and the mean-squared displacement $\langle \mathbf{r}(t) \mathbf{r}(t) \rangle = 2 \mathbf{D} t$ is normal. Good analytical properties of P^{-1} near $\mathbf{k} \approx \mathbf{0}$ and $\omega \approx 0$ are thus required for local diffusion to be well defined.

Recently a large number of physical problems such as diffusion in convective cells,⁹ diffusion on stochastic webs,¹⁰ diffusion on percolating clusters,¹¹ convection of a passive scalar by a turbulent flow, diffusion in disordered media, or on various kinds of fractal structures,¹² have displayed anomalous mean-squared displacement

$$\langle r^2(t) \rangle \sim t^\alpha. \quad (3)$$

The exponent, α , characterizing the fractional Brownian motion¹³ can be larger or smaller than one, with trapping or long ballistic flights often the cause of this subdiffusive or hyperdiffusive transport. When $\alpha \neq 1$, an application of the definition of the diffusion coefficient will give 0 or $+\infty$. This does not mean that the diffusion is 0 or $+\infty$, it means that *the concept of diffusion coefficient is meaningless, and that the expansion Eq. (2) does not give a simple formula*.

In the problem of fast heat diffusion along a magnetic field line combined with a Brownian wandering of this line, and a very slow collisional diffusion across the line, the exact solution of the associated Langevin equation, gives a stochastic transport across the main field with $\alpha = 1/2$, and the additional normal, $\alpha = 1$, collisional term is often negligible compared to the anomalous displacement. This solution will also display the fact that, as far as the mean-squared displacement is concerned, there are no couplings of the perpendicular purely collisional dynamics with the other processes. Thus effective diffusion coefficients D_{eff} involving parallel, perpendicular and magnetic diffusion are not directly recovered. Moreover the final result of the analysis is that, in many typical tokamak conditions, *instead of a propagator of the usual local type $P \sim \exp[-r^2/D_{\text{eff}}t]$ the*

propagator behaves as $P \sim \exp[-r^{4/3}/t^{1/3}]$, thus ruling out any attempt to define an effective diffusion coefficient.

Consider an infinite homogeneous magnetic field $\mathbf{B}$ directed along the z axis, and a small random field, $\mathbf{b}$, such that $\langle \mathbf{b} \rangle = 0$, $\nabla \cdot \mathbf{b} = 0$, and $\langle b^2 \rangle / B^2$ is a small parameter of the order of 10^{-6} - 10^{-8} ($\langle \rangle$ indicates statistical averages), the component $\mathbf{B} \cdot \mathbf{b}$, if any, is completely irrelevant to the problem. The second moments of the $\mathbf{b}$ field are characterized by the longitudinal (z) correlation length, $\lambda_{\parallel}$, and the transverse (x, y) one, $\lambda_{\perp}$ (the x and y variables describe a local slab frame near the braided magnetic surfaces). The tokamak geometry consists of nested toroidal surfaces with field lines traversing the poloidal angle, thus the statistical average is to be completed by a poloidal average. As noted by Kadomtsev and Pogutse⁴ there exist two regimes of magnetic field line diffusion, the quasilinear one, when $\lambda_{\parallel} b < \lambda_{\perp} B$, and the nonlinear one. Most studies have been restricted to the former, the ordering relevant for stochastic field line diffusion induced by resonant perturbations in a tokamak. Thus disregarding the x and y dependence of the moments compared to that in z , we consider the following Gaussian random field $\langle b_x(z)b_x(z') \rangle = \langle b_y(z)b_y(z') \rangle = \langle b^2 \rangle \exp[-(z-z')^2/2\lambda_{\parallel}^2]$ (the influence of transverse correlations is assessed at the end of the paper). This is the standard model of magnetic field line diffusion and the magnetic diffusion coefficient χ_M is:

$$\chi_M = \sqrt{\pi/2} \lambda_{\parallel} \langle b^2 \rangle / B^2. \quad (4)$$

We call $\chi_{\parallel}$ and $\chi_{\perp}$ the temperature (or density, if particle transport is diffusive rather than ballistic along one parallel correlation length) diffusion coefficients due to collisions, along and across the field. The correlation time of $\mathbf{b}$ is larger than the toroidal transit time of a thermal electron, thus $\mathbf{b}$ can be treated as static. This usual assumption is relevant to ergodic divertor configurations which are static, and to magnetic turbulence with frequency typically of the order of the electron diamagnetic frequency. The temperature diffusion equation is equivalent to Langevin equations, Eqs.(5), (6), (7):

$$\frac{dx}{dt} = \frac{dz}{dt} \frac{b_x}{B} + \eta_{\perp}, \quad (5)$$

$$\frac{dy}{dt} = \frac{dz}{dt} \frac{b_y}{B} + \eta_{\perp}, \quad (6)$$

$$\frac{dz}{dt} = \eta_{\parallel}. \quad (7)$$

The Lagrangian coordinates (x, y, z) describe pseudo-particles whose Eulerian average is the temperature distribution. The statistical characteristics of the collisional noises η are $\langle \eta_{\perp}(t) \rangle = 0$, $\langle \eta_{\perp}(t) \eta_{\perp}(t') \rangle = 2\chi_{\perp} \delta(t-t')$, $\langle \eta_{\parallel}(t) \rangle = 0$, $\langle \eta_{\parallel}(t) \eta_{\parallel}(t') \rangle = 2\chi_{\parallel} \delta(t-t')$ (δ is the Dirac distribution). Thus when $\mathbf{b} = \mathbf{0}$, we recover collisional fast diffusion along $\mathbf{B}$, and very slow diffusion across $\mathbf{B}$. Because of the very small value of the perturbing field we have neglected the difference between the curvilinear distance along the field line and the z coordinate ($ds = \sqrt{dx^2 + dy^2 + dz^2} = dz + O[\langle b^2 \rangle / B^2]$). Let us introduce $\mathbf{b}(\mathbf{k})$, the Fourier transform of $\mathbf{b}(\mathbf{z})$, and integrate Eqs. (5), (6), (7) to obtain $r^2 = x^2 + y^2$ as follows:

$$\langle r^2(t) \rangle = \int \frac{d\mathbf{k}}{k} \frac{d\mathbf{k}'}{k'} \langle \frac{\mathbf{b}(\mathbf{k}) \cdot \mathbf{b}^*(\mathbf{k}')}{B^2} \rangle \langle [e^{i\mathbf{k}z(t)} - 1][e^{i\mathbf{k}'z(t)} - 1] \rangle + \int_0^t du \int_0^u du' \langle \eta_{\perp}(u) \eta_{\perp}(u') \rangle. \quad (8)$$

Even if there are correlations between $\eta_{\parallel}(t)$ and $\eta_{\perp}(t')$ the fact that $\langle \mathbf{b} \rangle = \mathbf{0}$ leads to the cancellation of cross terms of the type $\eta_{\perp} \mathbf{b}$. Using the Gaussian properties of the collisional noise $\langle e^{i\mathbf{k}z(t)} \rangle = e^{-k^2 \chi_{\parallel} t}$, and $\langle b_x(\mathbf{k}) b_x^*(\mathbf{k}') \rangle = \langle b_y(\mathbf{k}) b_y^*(\mathbf{k}') \rangle = \langle b^2 \rangle \lambda_{\parallel} \delta(\mathbf{k} - \mathbf{k}') \exp[-k^2 \lambda_{\parallel}^2 / 2] / \sqrt{2\pi}$ we obtain the mean squared displacement :

$$\langle r^2(t) \rangle = \frac{8}{\sqrt{\pi}} \chi_M \sqrt{\chi_{\parallel} t} + 4\chi_{\perp} t. \quad (9)$$

No effective diffusion coefficient $D_{\text{eff}}(\chi_M, \chi_{\parallel}, \chi_{\perp}, \lambda_{\parallel})$ is found with the exact solution of the Langevin equation. On the basis of Eq.(9) two transport regimes are to be considered. For a given length r , if the first term on the right hand side dominates, we must accept the scaling $\langle r^2(t) \rangle = 8\chi_M \sqrt{\chi_{\parallel} t / \pi}$ and not try to define an effective local diffusivity, but explain the transport in term of a nonlocal propagator. When the anomalous $\sqrt{t}$ is smaller than the normal term an effective diffusion coefficient D_{eff} can be constructed, following the analysis proposed by Rechester and Rosenbluth,² provided that an additional mechanism like the exponential stretching of a stochastic instability is present, but such a mechanism does not exist for a pure random field.

The collisional diffusion coefficients scale differently with respect to density n , and temperature T : $\chi_{\parallel} \sim T^{5/2} n^{-1}$, $\chi_{\perp} \sim T^{-1/2} n$, so that in tokamaks we obtain a wide range of values: $\chi_{\parallel} \sim 10^9 - 10^{10} \text{ m}^2/\text{s}$, $\chi_{\perp} \sim 10^{-1} - 10^{-2} \text{ m}^2/\text{s}$. The typical width of braided magnetic domain is $r \sim 10^{-1} - 1 \text{ m}$, and $\chi_M \sim 10^{-5} - 10^{-8} \text{ m}$. Thus nonlocal transport is dominant in many situations, including the case of an ergodic divertor in a low density edge plasma, and that of localized braiding near resonant magnetic surfaces leading to a small stochastic layer. The cases of large scale stochasticity or of very small perturbations in a dense plasma fall in the second regime.

In the following we consider the first regime and neglect the ineffective slow transverse purely collisional transport due to $\chi_{\perp}$. To describe heat diffusion along the random field lines, and to calculate the propagator associated with the dominant $\sqrt{t}$ scaling we shall use a functional Wiener integration.¹⁴ The probability to go from the point $(0, 0, 0)$, to the point (x, y, z) , in a time t , averaged over all the realizations of the stochastic field lines, is given by the functional integral:

$$P(x, y, z, t) = \int_{(0,0,0)}^{(x,y,z)} D[x(u)] D[y(u)] P_1[x(u), y(u)] P_2[x(u), y(u), t]. \quad (10)$$

P_1 is the probability of a particular realization of a given line, $x(u)$, $y(u)$, linking $(0,0,0)$ and (x,y,z) , and P_2 is the probability to diffuse from $(0,0,0)$ to (x,y,z) in a time t , along this particular realization. $D[x(u)]$, $D[y(u)]$ stand for the functional integration over all the realizations of the random field. Within the quasilinear approximation, the field lines are described by a diffusion equation so that P_1 is given by the usual Gaussian formula:¹⁴

$$P(x, y, z, t) = \Theta(t) \int \frac{D[x(u)] D[y(u)]}{\sqrt{4\pi\chi_{\parallel}t}} e^{-\frac{1}{4\chi_M} \int_0^z du ([dx/du]^2 + [dy/du]^2)} e^{-\frac{1}{4\chi_{\parallel}t} \left[\int_0^z \sqrt{du^2 + dx^2 + dy^2} \right]^2}. \quad (11)$$

$\Theta(t)$ is the Heaviside step function ensuring causality. This formula exactly describes collisional diffusion along stochastic magnetic field lines, the field average being achieved

by the first exponential P_1 and the collisional diffusion being described by the second one P_2 . If we expand the exponent of this second integrand, $\sqrt{dx^2+dy^2+du^2}= du (1+ ([dx/du]^2+[dy/du]^2)/2) + O[(\langle b^2 \rangle/B^2)^2]$, the functional integration can be performed analytically up to this $O[(\langle b^2 \rangle/B^2)^2]$ order. Then $P(x,y,z,t)$ is to be summed over all the final values of z , in order to obtain the nonlocal propagator $P(x,y,t)$. Because of the very small value of $\langle b^2 \rangle/B^2$, it is in fact sufficient to work to $O[\langle b^2 \rangle/B^2]$. With very good accuracy the final z sum can be evaluated by saddle point integration, and the result is:

$$P(x,y,t) = \int P(x,y,z,t)dz = \frac{\Theta(t) e^{\left[\frac{-3(x^2+y^2)^{2/3}}{2^{2/3} 4 \chi_M^{2/3} \chi_{||}^{1/3} t^{1/3}} \right]}}{2^{2/3} \pi \sqrt{3} (x^2+y^2)^{1/3} \chi_M^{2/3} \chi_{||}^{1/3} t^{1/3}} \quad (12)$$

Equation (11) is an exact representation of P , but the normalization $\int P dx dy = 1$, has been lost during the saddle point z integration. To restore it, we have to multiply P in Eq. (12) by $\sqrt{3}/2$. Performing the poloidal average and reintroducing the proper normalization we obtain the final result :

$$P(r,t) = \frac{\Theta(t) e^{\left[-\frac{3r^{4/3}}{2^{2/3} 4 \chi_M^{2/3} \chi_{||}^{1/3} t^{1/3}} \right]}}{2\pi 2^{2/3} r^{2/3} \chi_M^{2/3} \chi_{||}^{1/3} t^{1/3}} \quad (13)$$

Equation (13) is the main result of this letter, with Eq.(1) it solves the problem of nonlocal heat transport in a stochastic magnetic field. With this propagator the evaluation of $\langle r^2(t) \rangle$ gives a value which is $\pi/2\sqrt{3}$ times the one obtained with the Langevin equation. In fact, without the saddle point approximation we obtain exactly the Langevin equation result.

In addition to the Langevin equation and to the Wiener integral solutions, we have performed a direct numerical simulation of heat diffusion in a stochastic magnetic field. We have studied the even moments of the radial temperature distribution when a Dirac distribution heat pulse is released on a given magnetic surface at time t equal to zero. The

average was performed over 10^2 stochastic field lines (because of the smoothing effect of parallel collisional diffusion the convergence, as a function of the number of lines, is very fast). The anomalous scaling $\langle r^{2p}(t) \rangle \sim t^{p/2}$, with $p=1, 2, 3\dots$ was observed in quantitative agreement with the previous theory, and is displayed on Fig. 1. The curves have been normalized to permit their display on a single plot.

Up to now we have considered random fields of the type $\mathbf{b}(z)$. Within the Hamiltonian picture of field line stochastic instability, z is the "time", and this model is relevant. However, random magnetic fields of the type $\mathbf{b}(x,y,z)$ are to be considered, in order to assess the influence of transverse finite correlations.⁴ Consider the nonlinear Langevin equations:

$$\frac{dx}{dt} = \frac{dz}{dt} \frac{b_x(x,y,z)}{B}, \quad \frac{dy}{dt} = \frac{dz}{dt} \frac{b_y(x,y,z)}{B}, \quad \frac{dz}{dt} = \eta_{\parallel}(t), \quad (14)$$

with $\langle b_y(x,y,z)b_y(x',y',z') \rangle = \langle b_x(x,y,z)b_x(x',y',z') \rangle = \langle b^2 \rangle \exp[-(z-z')^2/2\lambda_{\parallel}^2 - (x-x')^2/2\lambda_{\perp}^2 - (y-y')^2/2\lambda_{\perp}^2]$.

There are no general methods to solve this kind of nonlinear stochastic differential equations. This equation has some similarity with the problems of diffusion in turbulent flows, these problems have been widely studied and the observed intermittency¹⁵ can be studied with the help of multifractal measures.

In this letter we shall not carry out such a program, we shall simply apply a Gaussian approximation, of the resonant broadening type. Thus, what is expected is not only an anomalous scaling of the mean squared displacement with respect to the time, but also with respect to the perturbing magnetic field:⁴

$$\langle r^2(t) \rangle \sim [\langle b^2 \rangle / B^2]^{\beta} t^{\alpha}. \quad (15)$$

The usual $\beta=1$ quasilinear scaling is no longer valid because of the nonlinearity. To calculate α and β , associated with the stochastic differential equation Eq. (14), we Fourier analyze the $\mathbf{b}$ field, and perform an approximate evaluation of the resulting integral representation of $\langle dx/dt \cdot dx/dt' \rangle$ and $\langle dy/dt \cdot dy/dt' \rangle$, with a Gaussian assumption.

After some calculations we obtain the following differential equation in the long time limit:

$$\frac{d^2 \langle r^2(t) \rangle}{dt^2} \sim - \left[\frac{\langle b^2 \rangle}{B^2} \lambda_{\parallel} \lambda_{\perp}^2 \chi_{\parallel}^{1/2} \right] \langle r^2(t) \rangle^{-1} t^{-3/2}. \quad (16)$$

Bohm scaling, $\beta=1/2$, is thus recovered but with an highly anomalous diffusion exponent $\alpha = 1/4$: $\langle r^2(t) \rangle \sim \lambda_{\parallel}^{0.5} \lambda_{\perp}^{0.25} \langle b^2 \rangle^{0.5} B^{-1} t^{0.25}$. This additional slowing down is due the additional x, y disorder. In this regime, the calculation of the associated propagator P remain an open question.

Modulation experiments and heat pulse propagation studies have been used to measure the dynamical transport responses of a tokamak plasma, i.e, the propagator P. These kinds of experiments seem to be the ideal tool to distinguish local and nonlocal propagators. In addition to the dynamical response we can consider also the global one, but to address the problem of the scaling of the energy lifetime, we must know the extent of the ergodic regions and the scaling of the saturated value of the turbulent field arising through the self sustainment constraint.

We have demonstrated, discussed, and checked numerically, the fact that for a wide range of tokamak parameters heat transport in a stochastic magnetic field is described by Eq. (1), where the propagator P given by Eq. (13) must be used. An effective local diffusion can not be defined in this regime. The propagator behaves as $\exp[-r^{4/3}/t^{1/3}]$ rather than as $\exp[-r^2/t]$, and is independant of the very small cross field collisional diffusion.

Acknowledgments

This work was supported by the United States Department of energy under contract number DE-AC02-76-CHO3073. J.M. Rax would like to thank Professor N. Fisch for encouragement, advice, and stimulating discussions.

References

- 1** M.N. Rosenbluth, R.Z. Sagdeev, J.B. Taylor, and G.M. Zaslavski, Nucl. Fusion **6**, 415 (1966). N.N. Filonenko, R.Z. Sagdeev, and G.M. Zaslavski, Nucl. Fusion **7**, 253 (1967). T.H. Stix, Phys. Rev. Lett. **30**, 833 (1973). A.B. Rechester and T.H. Stix, Phys. Rev. Lett. **36**, 587 (1976).
- 2** A.B. Rechester, and M.N. Rosenbluth, Phys. Rev. Lett. **40**, 38 (1978). A.B. Rechester, M.N. Rosenbluth, and R.B. White, Phys. Rev. Lett. **42**, 1247 (1979). A. Boozer and R.B. White, Phys. Rev. Lett. **49**, 786 (1982). R.B. White "Statistical physics and chaos in fusion plasma" Horton and Reichl ed., Wiley, New York (1984). J.A. Krommes, C. Oberman and R.G. Kleva J. Plasma Phys. **30**, 11, (1985). H.E. Mynick, and J.A. Krommes Phys. Rev. Lett. **43**, 1506 (1987). P.H. Rebut, M. Brusati, M. Hugon and P. Lallia Nucl. Fusion Sup. **2** IAEA-CN-47/E-III-4 (1987). G. Laval, "From Particles to Plasmas", J.W. Van Dam ed., Addison-Wesley, New York (1989).
- 3** T.H. Stix, Nucl. Fusion **18**, 354 (1978).
- 4** E.B. Kadomtsev, and O.P. Pogutse, Nucl. Fusion Sup. **1** IAEA-CN-37/O-1 (1979).
- 5** W. Feneberg, and G.H. Wolf Nucl. Fusion **21**, 669 (1981). T.J. Martin, and J.B. Taylor, Plasma Phys. Controlled Fusion **26**, 321(1984).
- 6** J.R. Albritton, Phys. Rev. Lett. **50**, 2078 (1983). J.F. Luchiani, P. Mora, and J. Virmont, Phys. Rev. Lett. **51**, 1664 (1985).
- 7** A.R. Jacobson, and R.W. Moses, Phys. Rev. Lett. **52**, 2041 (1984).
- 8** J.M. Rax, and D. Moreau, Nucl. Fusion **29**, 1751 (1989).
- 9** W. Young, A. Pumir, and Y. Pomeau, Phys. Fluids A **1**, 462 (1989).
- 10** G.M. Zaslavski, R.Z. Sagdeev, D.K. Chaikovski, and A.A. Chernikov, Sov. Phys. JETP **68**, 995 (1989).
- 11** Y. Gefen, A. Aharony, and S. Alexander Phys. Rev. Lett. **50**, 77 (1983).
- 12** B. O'Shaughnessy, and I. Procaccia Phys. Rev. Lett. **54**, 455 (1985). S. Havlin, and D. Ben-Avraham, Advances in Phys. **36**, 695 (1987).
- 13** B.D. Hughes, E.W. Montroll, and M.F. Shlesinger J. Stat. Phys. **28**, 111 (1982). B.B. Mandelbrot "The fractal geometry of nature", Freeman, New York (1983). J. Feder, "Fractals", Plenum Press, New York (1988).
- 14** F.W. Wiegell Phys. Rep. **16**, 57 (1974). R.P. Feynman, and A.R. Hibbs "Quantum Mechanics and Path Integrals", Mac Graw Hill, New York (1965). L.S. Schulman "Techniques and applications of path integration", Wiley, New York (1981).
- 15** Y.B. Zeldovich, S.A. Molchanov, A.A. Ruzmaikin, D.D. Sokolov, Sov. J.E.T.P **62**, 1188, (1985). Y.B. Zeldovich, A.A. Ruzmaikin, and D.D. Sokoloff, "The Almighty Chance", World Scientific, Singapore (1990).

Figure caption

Fig. 1: The first even moments $\langle x^2(t) \rangle$, $\langle x^4(t) \rangle$, $\langle x^6(t) \rangle$ of the temperature distribution averaged over 100 stochastic field lines, $\chi_{||} = 10^9 \text{ m}^2/\text{s}$, $\lambda_{||} = 10 \text{ m}$, $\langle b^2 \rangle / B^2 = 10^{-6}$, in logarithmic scale.

EXTERNAL DISTRIBUTION IN ADDITION TO UC-420

Dr. F. Paoloni, Univ. of Wollongong, AUSTRALIA	Director, Japan Atomic Energy Research Inst., JAPAN
Prof. M.H. Brennan, Univ. of Sydney, AUSTRALIA	Prof. S. Itoh, Kyushu Univ., JAPAN
Plasma Research Lab., Australian Nat. Univ., AUSTRALIA	Data and Planning Center, Nagoya Univ., JAPAN
Prof. I.R. Jones, Flinders Univ, AUSTRALIA	Prof. S. Tanaka, Kyoto Univ., JAPAN
Prof. F. Cap, Inst. for Theoretical Physics, AUSTRIA	Library, Kyoto Univ., JAPAN
Prof. M. Heindler, Institut für Theoretische Physik, AUSTRIA	Prof. N. Inoue, Univ. of Tokyo, JAPAN
Prof. M. Goossens, Astronomisch Instituut, BELGIUM	S. Mori, Technical Advisor, JAERI, JAPAN
Ecole Royale Militaire, Lab. de Phy. Plasmas, BELGIUM	O. Mitarai, Kumamoto Inst. of Technology, JAPAN
Commission-European, DG. XII-Fusion Prog., BELGIUM	H. Jeong, Korea Advanced Energy Research Inst., KOREA
Prof. R. Bouciqué, Rijksuniversiteit Gent, BELGIUM	Prof. D.I. Choi, The Korea Adv. Inst. of Sci. & Tech., KOREA
Dr. P.H. Sakanaka, Instituto Fisica, BRAZIL	Prof. B.S. Liley, Univ. of Waikato, NEW ZEALAND
Instituto De Pesquisas Especiais-INPE, BRAZIL	Inst. of Plasma Physics, PEOPLE'S REPUBLIC OF CHINA
Documents Office, Atomic Energy of Canada Ltd., CANADA	Librarian, Inst. of Physics, PEOPLE'S REPUBLIC OF CHINA
Dr. M.P. Bachynski, MPB Technologies, Inc., CANADA	Library, Tsinghua Univ., PEOPLE'S REPUBLIC OF CHINA
Dr. H.M. Skarsgard, Univ. of Saskatchewan, CANADA	Z. Li, S.W. Inst. Physics, PEOPLE'S REPUBLIC OF CHINA
Prof. J. Teichmann, Univ. of Montreal, CANADA	Prof. J.A.C. Cabral, Instituto Superior Tecnico, PORTUGAL
Prof. S.R. Sreenivasan, Univ. of Calgary, CANADA	Dr. O. Petrus, AL I CUZA Univ., ROMANIA
Prof. T.W. Johnston, INRS-Energie, CANADA	Dr. J. de Villiers, Fusion Studies, AEC, S. AFRICA
Dr. R. Bolton, Centre canadien de fusion magnétique, CANADA	Prof. M.A. Hellberg, Univ. of Natal, S. AFRICA
Dr. C.R. James., Univ. of Alberta, CANADA	C.I.E.M.A.T, Fusion Division Library, SPAIN
Dr. P. Lukac, Komenského Univerzita, CZECHOSLOVAKIA	Dr. L. Stenflo, Univ. of UMEA, SWEDEN
The Librarian, Culham Laboratory, ENGLAND	Library, Royal Inst. of Technology, SWEDEN
Library, R61, Rutherford Appleton Laboratory, ENGLAND	Prof. H. Wilhelmson, Chalmers Univ. of Tech., SWEDEN
Mrs. S.A. Hutchinson, JET Library, ENGLAND	Centre Phys. Des Plasmas, Ecole Polytech, SWITZERLAND
P. Mähönen, Univ. of Helsinki, FINLAND	Bibliotheek, Inst. Voor Plasma-Fysica, THE NETHERLANDS
C. Mouttet, Lab. de Physique des Milieux Ionisés, FRANCE	M. Durgut, Vice Chairman, Middle East Tech. Univ., TURKEY
J. Radet, CEN/CADARACHE - Bat 506, FRANCE	Dr. D.D. Ryutov, Siberian Branch of Academy of Sci., USSR
Ms. C. Rinri, Univ. of Ioannina, GREECE	Dr. G.A. Eliseev, Kurchatov Inst., USSR
Dr. T. Mui, Academy Bibliographic Ser., HONG KONG	Librarian, The Ukr.SSR Academy of Sciences, USSR
Preprint Library, Hungarian Academy of Sci., HUNGARY	Dr. L.M. Kovrizhnykh, Inst. of General Physics, USSR
Dr. B. Das Gupta, Saha Inst. of Nuclear Physics, INDIA	Kernforschungsanlage GmbH, Zentralbibliothek, W. GERMANY
Dr. P. Kaw, Inst. for Plasma Research, INDIA	Bibliothek, Inst. Für Plasmaforschung, W. GERMANY
Dr. P. Rosenau, Israel Inst. of Technology, ISRAEL	Prof. K. Schindler, Ruhr-Universität Bochum, W. GERMANY
Librarian, International Center for Theo. Physics, ITALY	Dr. F. Wagner, (ASDEX), Max-Planck-Institut, W. GERMANY
Miss C. De Palo, Associazione EURATOM-ENEA, ITALY	Librarian, Max-Planck-Institut, W. GERMANY
Dr. G. Grosso, Istituto di Fisica del Plasma, ITALY	Prof. R.K. Janev, Inst. of Physics, YUGOSLAVIA
Dr. H. Yamato, Toshiba Res. & Devel. Center, JAPAN	
Prof. I. Kawakami, Atomic Energy Res.Inst., JAPAN	
Prof. K. Nishikawa, Hiroshima Univ., JAPAN	

END

**DATE
FILMED**

01/09/91