ANL-HEP-CP-90-101
For the Proceedings of the
9th International Symposium on
High Energy Spin Physics
Bonn, West Germany
10-14 September 1990

EXPERIMENTS WITH FERMILAB POLARIZED PROTON AND ANTIPROTON BEAMS^S
(For the Fermilab E-704 Collaboration*)

ANL-HEP-CP--90-101

A. Yokosawa
High Energy Physics Division, Argonne National Laboratory
Argonne, Illinois 60439 USA

DE91 006551

Abstract

We summarize activities concerning the Fermilab polarized beams. They include a brief description of the polarized-beam facility, measurements of beam polarization by polarimeters, asymmetry measurements in the π^0 production at high $p_{\perp}$ and in the Λ (Σ^0), $\pi^{\pm}$, π^0 production at large x_F , and $\Delta\sigma_L(pp, \bar{p}p)$ measurements.

Introduction

The Fermilab polarized-beam facility at 200 GeV/c became operational in 1987. At that time, we mainly concentrated on the measurements of beam polarization by polarimeters and on the studies of beam polarization monitors. Recently we carried out the polarized-beam program designated as E-704 by an international collaboration.

We summarize these activities including the discussions of some preliminary data on $p_{\perp}$ and x_F dependence.

Polarized Beam Facility

The Fermilab polarized-beam facility¹ was operated during the past TeV-II (fixed target) period which ended in February 1988.

An extracted beam from the Tevatron is delivered through the MP primary-beam line to the Meson Detector Building where a 0.73-interaction-length Be target is utilized to produce Λ and $\bar{\Lambda}$ at $\theta_{c.m.} \approx 0^\circ$. Protons and antiprotons from the Λ and $\bar{\Lambda}$ decays respectively are brought to a final target position in the MP hall through the MP secondary beam (200 GeV/c) line.

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

Polarized protons from the virtual sources as shown in Fig. 1 are focussed in the tagging section, where both the momentum and polarization are selected.¹

The typical beam flux ($\Delta p/p = \pm 5\%$) for 3×10^{12} incident protons per 20-sec spill at 200 GeV/c were: (P_{av} is average polarization)

	Tagged Beam $P_{av} = 45\%$	Total Protons (antiprotons)	Background π 's
Protons	1.0×10^7	2.0×10^7	2.0×10^6
Antiprotons	5.0×10^5	1.0×10^6	5.0×10^6

Figure 1 Virtual sources (top view).

Primakoff-Effect Measurement

The asymmetry of the nuclear coherent Coulomb π^0 production process ("Primakoff process"), was measured² for the first time with the use of the polarized-proton beam. The apparatus consisted of a lead-glass calorimeter for π^0 detection and a magnetic spectrometer for the scattered protons. A large asymmetry in the region of $|t'| < 0.001 \text{ (GeV/c)}^2$ and $1.36 < M(\pi^0 p) < 1.52 \text{ GeV/c}^2$ was observed for the reaction $p + \text{Pb} \rightarrow p + \pi^0 + \text{Pb}$, where the Coulomb process is predominant. The expected null asymmetry was observed in the larger $|t'|$ region where the diffractive-dissociation process is predominant.

The observed ϕ -angle dependence of the coherent π^0 production process may be expressed as $1 + (f T(\theta) P_B) \cos \phi$, where $T(\theta)$ is the analyzing power (target azimuthal asymmetry) for photoproduction of π^0 from a polarized proton target at c.m. polar angle θ , ϕ is the azimuthal angle, P_B is the transverse polarization, and the parameter f is a dilution factor caused by the diffractive dissociation. The raw asymmetry at ϕ is obtained as

$$A(\phi) = (N^\uparrow(\phi) - N^\downarrow(\phi)) / (N^\uparrow(\phi) + N^\downarrow(\phi)) = f T(\theta) P_B \cos \phi = \epsilon \cos \phi,$$

where $N^\uparrow(\phi)$ and $N^\downarrow(\phi)$ are the number of events at ϕ for the up and down spin direction of the incident proton, respectively.

The measured asymmetry for the Coulomb process is consistent with the analyzing power (about -70%) of the π^0 production process deduced from existing low-energy $\gamma + p \rightarrow \pi^0 + p$ data. The results demonstrate that the Primakoff process is useful for the measurement of proton and antiproton polarization at high energy.

Coulomb-Nuclear Interference Measurements

The analyzing power, A_N , of proton-proton, proton-hydrocarbon, and antiproton-hydrocarbon scattering in the Coulomb-nuclear region was measured with use of the polarized-proton and polarized-antiproton beams.³ For the elastic scattering at small $|t|$, a set of scintillation counters was utilized to detect the recoil proton which stops within a very short range in the scintillator. The results at $|t| \sim 0.003 \text{ (GeV/c)}^2$ show the value $A_N = (2.4 \pm 0.9)\%$ with the polarized-proton beam, and $A_N = (-4.6 \pm 1.9)\%$ with the polarized-antiproton beam both on a hydrocarbon target, and also $A_N = (4.5 \pm 2.8)\%$ of proton-proton scattering. These results are consistent with predictions⁴⁻⁶ based on Coulomb-nuclear interference. Recently A_N measurements were repeated with much higher statistics than those mentioned above.

Single-Spin Asymmetry in $p^\uparrow p \rightarrow \pi^0 X$ and $\bar{p}^\uparrow p \rightarrow \pi^0 X$ at High $p_\perp$

This experiment is recently completed at incident proton momentum of 200 GeV/c. Preliminary data of the $p^\uparrow p$ reaction show that the asymmetry values (A_N) at $x_F \approx 0$ are approximately zero (or small negative) up to $p_\perp = 3.5$ GeV/c and then begin to rise to $\sim +40\%$ in the region of $p_\perp = 4$ to 5 GeV/c as shown in Fig. 2. At lower energies as seen in the BNL⁷ ($p^\uparrow p \rightarrow \pi^+ X$), CERN⁸

Fig. 2 $p_\perp$ dependence of A_N at $x_F \approx 0$ (preliminary data).

($pp^\uparrow \rightarrow \pi^0 X$), Serpukhov⁹ ($\pi^- p^\uparrow \rightarrow \pi^0 X$) data, this rapid rise from zero to large positive values,* was also observed as shown in Fig. 3 although none of the data exceeded $p_\perp = 3$ GeV/c. A new finding is that all the A_N data of π^0 or π^+ production at $x_F \approx 0$ show the large positive asymmetries begin at $x_\perp = 0.4$ in the region $\sqrt{s} = 5$ to 20 GeV as shown in Fig. 4. This is strong indication that we are indeed observing asymmetries caused by hard scattering. We note that the common crossing point $x_\perp = 0.4$ was pointed out in the Ref. 9 in the

* Data taken with polarized targets (Refs. 8 and 9) are normalized to the measurements with polarized beams, that is, the sign of A_N ($hp^\uparrow \rightarrow hX$, where h represents hadron) is reversed. Note that $A_N(p^\uparrow h \rightarrow hX) = -A_N(hp^\uparrow \rightarrow hX)$ at $x_F \approx 0$.

region $\sqrt{s} = 5$ to 8.5 GeV/c. Theoretically single-spin asymmetries are discussed within the context of the QCD hard-scattering model.¹⁰ By knowing the quark content of $\pi^+ = u\bar{d}$ and $\pi^0 = (u\bar{u} - d\bar{d})/\sqrt{2}$, polarized u quark in the polarized proton beam is considered to be the carrier of the spin information.

Figure 3

Asymmetry $A_N(\%)$ vs. p_L at $P_{lab} = 1.33$ to 40 GeV/c

Figure 4 p_T and x_T positions at $A_N = 0 \rightarrow$ large positive vs. $\sqrt{s}$.

Single-spin asymmetry in $\bar{p}^\uparrow p + \pi^0 X$ shows a similar $p_\perp$ dependence as the $p^\uparrow p$ case. However, data are limited only up to $p_\perp = 3.5$ GeV/c.

x_F Dependence of Single-Spin Asymmetry in $p^\uparrow p + \pi^0 X$ and $\bar{p}^\uparrow p + \pi^0 X$

Measurements on the x_F dependence at 200 GeV/c covering $p_\perp$ up to 2 GeV/c were recently completed. Asymmetry values in the $p^\uparrow p$ reaction are consistent with zero up to $x_F = 0.3$ to 0.4, and then linearly increase to + 20% near $x_F = 1.0$ as shown in Fig. 5. The data suggest an influence of polarized u quarks at large x_F . Also they are consistent with earlier data¹¹ taken at $\langle x_F \rangle = 0.52$. This is the first x_F dependence data ever obtained in the π production. It is interesting to notice that our data resembled x_F dependence for Λ polarization in $pp + \Lambda^\uparrow X$ where polarized s quarks were considered¹² to be responsible for high polarization.

Single-spin asymmetry in $\bar{p}^\uparrow p + \pi^0 X$ shows a similar x_F dependence as $p^\uparrow p$ case.

Figure 5 x_F dependence of A_N at $p_\perp = 0.5$ to 2.0 GeV/c (preliminary data).

A Short Summary on $x_\perp$ and x_F Dependence

There are two distinct common phenomena observed in the $x_\perp$ and x_F dependence of the one-spin asymmetries. The asymmetry values are zero or small for both $x_\perp$ and $x_F < 0.3$ to 0.4. Then there is a rise from zero to

large positive values for x_L and $x_F > 0.3$ to 0.4 . It will be interesting to find out if these two phenomena are related. One strong hint is that polarized u quarks seem responsible to the rise in A_N at high x_L and high x_F . It is interesting to investigate if these phenomena may be related to the origin of proton spin. High- p_L and high- $-x_F$ scattering phenomena were interpreted as an indication for the existence of rotating color charges in polarized protons.¹³

$\Delta\sigma_L$ (pp and $\bar{p}p$), and Double-Spin Asymmetry in $p^\uparrow p^\uparrow \rightarrow \pi^0 X$ and $\bar{p}^\uparrow p^\uparrow \rightarrow \pi^0 X$

Difference in total cross sections for pure spin states, $\Delta\sigma_L$ (pp and $\bar{p}p$), was simultaneously measured at 200 GeV/c with the π^0 production up to $p_L = 3$ GeV/c. Data are currently being analyzed.

$\pi^\pm$ and Hyperon Production on Hydrogen Target With Polarized Beam

Measurements of $p^\uparrow p \rightarrow (\pi^\pm, \Lambda^0, \Sigma^0) + X$ at 200 GeV/c were completed and data are currently being analyzed.

Future Experiments

We have proposed the following simultaneous measurements using a polarized beam and a polarized target.

i) Spin dependence in direct-gamma production at high p_L .

To understand the basic question of the origin of proton spin, we may be able to determine the gluon spin distribution in the proton by measuring the spin correlation parameter A_{LL} in the direct- γ production at high p_L , with longitudinally polarized protons on longitudinally polarized target nucleons.

$$A_{LL} = (1/P_B P_T) \frac{N(\uparrow\uparrow) - N(\uparrow\downarrow)}{N(\uparrow\uparrow) + N(\uparrow\downarrow)},$$

where P_B is beam polarization, P_T is target polarization, and arrows indicate the spin direction in the laboratory system.

The QCD Compton effect, "gluon + quark $\rightarrow$ gamma + quark", is expected to be the dominant mechanism for direct- γ production at large $p_{\perp}$. The parameter A_{LL} is approximately proportional to the gluon polarization.^{14,15}

Our plan is to carry out the A_{LL} measurements up to $p_{\perp} = 5$ GeV/c with reasonable statistical accuracy. Our main detectors consist of lead-glass counters and proportional wire chambers.

ii) Spin dependence of χ_2 (3555) production.

We plan to measure the spin dependence of χ_2 (3555) production, which will be simultaneously carried out with the above mentioned direct-gamma measurements. The double-spin asymmetry, A_{LL} , in $p^{\uparrow}p^{\uparrow} \rightarrow \chi_2$ (3555) + J/ψ + γ is also expected to provide a means to study the spin dependent gluon structure function. The 15% decay branching ratio of χ_2 (3555) to J/ψ + γ allows us to analyze the helicity of the charmonium state.

There is general agreement¹⁶ theoretically that the χ_2 (3555) state is mainly produced by gluon-gluon fusion as shown below and there are promising experimental results¹⁸ suggesting that simple gluon fusion is sufficient to account for the χ_2 (3555) production in proton interactions at 200 and 250 GeV/c. The measured two-spin asymmetries (as defined below) give information on the initial gluon polarization, which can be used to reconstruct the gluon spin distribution in the polarized proton.¹⁸ By considering the fusion process, if the initial helicity state is $(+-)$, that is $(\frac{1}{2})$, the $J_z = 2$ and this state produces χ_2 .

To be exact, the observable A_{LL} is related to the distribution function of a polarized gluon in a polarized proton expressed as $G_+(x)$ and $G_-(x)$ with same- and opposite-sign helicities respectively.¹⁸

$$A_{LL}(x_F) = (1-R)/(1+R) \left[\frac{\Delta G}{G}(x_1) \cdot \frac{\Delta G}{G}(x_2) \right],$$

where x_1 , x_2 , and x_F are the longitudinal-momentum fraction of gluons, R is the ratio of matrix elements f_+ (f_-) which are the squared matrix elements for the production of χ_2 out of two gluons with same- (opposite-) sign helicities, and $\Delta G/G(x) \equiv (G_+(x) - G_-(x))/(G_+(x) + G_-(x))$.

Theoretical predictions were recently made on $A_{LL}(x_F)$ in both the J/ψ and χ_1 production based on a perturbative QCD approach.^{19,20}

REFERENCES

- S Work supported in part by the U.S. Department of Energy, Division of High Energy Physics, Contract W-31-109-ENG-38.
- * The E-581/704 Collaboration: Argonne (US), Fermilab (US), Univ. of Iowa (US), Kyoto-Kyoto Sangyo-Kyoto Education-Hiroshima (Japan), LAPP (France), Los Alamos (US), Northwestern Univ. (US), Osaka-Okayama (Japan), Rice Univ. (US), CEN Saclay (France), IHEP Serpukhov (USSR), INFN Trieste-Messina-Udine (Italy), Univ. Occup. Enviror. Health (Japan).
- 1) Grosnick et al., Nucl. Instrum. and Meth. A290 (1990) 269.
 - 2) D. C. Carey et al., Phys. Rev. Lett. 64 (1990) 357.
 - 3) N. Akchurin et al., Phys. Lett. B229 (1989) 299.
 - 4) B. Z. Kopeliovich and L. I. Lapidus, Yad. Fiz. 19 (1974) 218. (Sov. J. Nucl. Phys. 19 (1974) 114).
 - 5) N. H. Buttmore, E. Gotsman, Am. E. Leader, Phys. Rev. D18 (1978) 694.
 - 6) N. H. Buttmore, Proc. 6th Int. Symp. on High Energy Spin Phys. Marseille, J. Soffer, ed., Journal de Physique, 46, C2 (1985) p. 643.
 - 7) S. Sanoff et al., Phys. Rev. Lett. 64 (1990) 995.
 - 8) J. Antille et al., Phys. Lett. B94 (1980) 523.
 - 9) V. D. Apokin et al., Phys. Lett. B243 (1990) 461.
 - 10) D. Sivers, Phys. Rev. D41 (1990) 83; Phys. Rev., to appear.
 - 11) B. E. Bonner et al., Phys. Rev. Lett. 61 (1988) 1918.
 - 12) T. deGrand and H. I. Miettinen, Phys. Rev. D24 (1981) 2419; Phys. Rev. D31 (E) (1985) 661.
 - 13) Liang Zuo-tang and Meng Ta-chung, Phys. Rev. 42 (1990) 2380; Meng Ta-chung, Proc. 25th Rencontre de Moriond, Mar. '90, Les Arc, France.
 - 14) M. B. Einhorn and J. Soffer, Nucl. Phys. 274 (1986) 714; N. S. Craigie, K. Hidaka, M. Jacob, and F. M. Renard, Phys. Reports 99 (1983) 143.
 - 15) E. L. Berger and J. W. Qiu, Phys. Rev. D40 (1988) 788; D40 (RC) (1989) 3128.

- 16) S. D. Ellis et al., Phys. Rev. Lett. 36 (1976) 1263; B. L. Ioffe, Phys. Rev. Lett. 39 (1977) 1589; C. E. Carlson et al., Phys. Rev. D18 (1978) 760; H. Fritzsch, Phys. Lett. 67B (1977) 217; M. Gluck et al., Phys. Rev. D17 (1978) 2324; L. M. Jones et al., Phys. Rev. D17 (1978) 1782; V. Berger et al., Z. Phys. C6 (1980) 169; J. H. Kuhn, Phys. Lett. 89B (1980) 385; Y. Afek et al., Phys. Rev. D22 (1980) 86.
- 17) D. A. Bauer et al., Phys. Rev. Lett., 54 (1985) 753; Y. Lemoigne et al., Phys. Lett. 113B (1982) 509.
- 18) J. L. Cortes and B. Pire, Phys. Rev. 38 (RC) (1988) 3586.
- 19) A. P. Contogouris, S. Papadopoulos, and B. Kamal, Phys. Lett. B246 (1990) 523.
- 20) M. A. Doncheski and R. Robinett, PSU/TH/67 (1990).

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

END

DATE FILMED

02 / 05 / 91