Measurement of carbon and nitrogen biomass, and biovolume
from naturally derived marine bacterioplankton

BNL--47941
DE93 000371

SangHoon Lee
Brookhaven National Laboratory
Oceanographic and Atmospheric Sciences Division
Upton, NY 11973

SEP 28 1992

I. INTRODUCTION	2
II. MATERIALS REQUIRED	2
A. Equipment	3
B. Supplies	3
C. Solutions	3
III. PROCEDURES	3
A. Inoculation and culturing	4
B. CHN samples	4
i. Sample collection (cell harvesting)	5
ii. Controls, blanks and standards	5
C. Cell enumeration and size measurements	6
i. Cell enumeration by epifluorescence microscopy	6
ii. Cell size measurements by photomicrography	6
IV. NOTES AND COMMENTS	8
V. REFERENCES	12

This research was performed under the auspices of the United States
Department of Energy under Contract No. DE-AC02-76CH00016.

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

I. INTRODUCTION

Bacterial production or cell concentration often needs to be presented in carbon (C) or nitrogen (N) biomass. Conversions to such universal units are particularly necessary to interpret the data in terms of energy or material flow in relation to other trophic groups of organisms. For field applications, bacterial biomass is routinely estimated from cell concentrations or cell volumes (biovolume) with appropriate conversion factors.¹⁻³ There are several biomass conversion factors reported to date (Table 1) from direct determinations, using a variety of methods and organisms.³⁻¹⁰

For the direct determination of a conversion factor, in principle, three independent parameters are measured; number of cells, biovolume of the cells, and biomass. Current methods for all of those three are subject to potential errors. This is particularly true for the biovolume estimation, since errors in measuring linear dimension amplify to the power of 3 when converted to biovolume.

Described here is a method that combines the acridine orange (AO) direct count (DC) method for cell enumeration, AO epifluorescence photomicrography for the biovolume measurement, and CHN analysis for C and N biomass.⁹ Compared to other similar studies, this method has several improvements. For example, 1) bacterial cells are derived from natural seawater and grown in unsupplemented, filter-sterilized, particle-free natural seawater; 2) corrections are made for the bacterial cells passing through glass fiber GF/F filters; and 3) biovolume measurement is calibrated with fluorescent microspheres of known diameters.

II. MATERIALS REQUIRED

A. Equipment

Epifluorescence microscope and photomicrography system

CHN analyzer and related material

Slide projector

Vacuum pump

B. Supplies

Nuclepore polycarbonate filters (0.6 μ m pore size, 47 mm diameter)

Millipore filters (type GS; 0.22 μ m pore size, 47 mm diameter)

Combusted (at 450°C for > 4 hr) material:

 Glass fiber GF/F filters (24 mm diameter or smaller)

 Forceps, aluminum foil

Acid-rinsed opaque plastic bottles (1 liter)

Acid-rinsed or combusted filtration units for the 47 mm diameter type

 GS filters, and for the pre-combusted glass fiber GF/F filters

Nuclepore filters (0.2 μ m pore size, 25 mm diameter)

Microscope slides, cover slip, immersion oil

Kodak Ektachrome 400

C. Solutions

Irgalan black for staining filters (pre-stained Nuclepore filters are also available)

Acridine orange

Culture medium and inoculum, prepared from freshly collected natural sample by filtration

III. PROCEDURES

A. Inoculation and culturing

1. Soak 47 mm diameter type GS (0.22 μm pore size) Millipore filters (Millipore Corp., Bedford, Mass.) in deionized water before use. This is to screen out defective filter discs and to leach out foreign chemicals, often detergent, from filter materials. Use organic-free (acid-rinsed or combusted) labware, and rinse the containers thoroughly several times with the sample (or filtered) seawater before use.
2. Culture medium: Using a large (47 mm diameter) filtration unit (e.g., Millipore Sterifil unit), filter ca. 1 L of fresh natural seawater through the type GS filters with vacuum of <20 cm Hg; the filtrate is the culture medium. Prepare another 1 L of GS filtrate to estimate organic material that is contributed by particles shed from the GS filter (see Note and Comments).
3. Inoculum: Use the filtrate of 0.6 μm pore size Nuclepore (Costar Corp., Cambridge, Mass.) filters. These filters remove most of non-bacterial particles as well as large bacterial cells and particle-associated cells (see Notes and Comments). Using the Sterifil unit, prepare the 0.6 μm filtrate from the same seawater sample with vacuum of <20 cm Hg.
4. Start the culture by mixing about 10-20 ml of inoculum to 1000 ml of media in a dark bottle. The use of dark bottle is to suppress the growth of autotrophic cells possibly present in the filtrate. Monitor the bacterial cell concentration periodically (e.g., every 12 hrs) by the AODC method (see below). Harvest the cells at the end of the logarithmic growth, normally a few to several days after the inoculation.

B. CHN samples

i. Sample collection (cell harvesting)

1. Collect subsample for AODC from the culture before cell harvesting, and preserve with formalin (1-5% by vol). Using organic-free labware, collect the cultured cells on pre-combusted glass fiber GF/F filters (Whatman) by filtration with vacuum of <20 cm Hg. Always handle the CHN samples with combusted forceps on a sheet of combusted aluminum foil to avoid contamination. Glass fiber filters of a smaller diameter (e.g., 1.3 cm) will be better, if a CHN analyzer requires the samples to be wrapped in tiny tin boats. Save the filtrate.
2. Measure and note the volume of the filtrate, i.e., the culture filtered through the GF/F filter. Collect subsample from the filtrate for AODC samples and preserve as described above. A significant fraction of the bacterial cells pass through glass fiber filters (see Notes and Comments).
3. After the cell collection, wrap the GF/F filters with tin boats (or with combusted aluminum foil), and dry thoroughly at ca. 60°C for >6 hrs to prevent the degradation of organic materials. Store the samples in a dessicator until CHN analysis.

ii. Controls, blanks and standards

1. Filter shedding control: Filter the GS filtrate (i.e., culture media, see above) through a pre-combusted GF/F filter. It will be better if the volume of the GS filtrate that is filtered through the GF/F filter is comparable to the volume of the culture.
2. Blank: The blank may be the absolute blank, i.e., the pre-combusted GF/F, or the pre-combusted GF/F wetted with the culture media.

3. Internal standard: Apply small known amount of high-purity amino acids (e.g., alanine has a C:N ratio similar to that of bacteria) to a pre-combusted GF/F filter. Thoroughly dry and store the filters as described above. See IV for the use of the controls and blanks.

C. Cell enumeration and size measurements

i. Cell enumeration by epifluorescence microscopy

Hobbie et al.¹¹ described a standard protocol with detailed background information. Methods for direct counts are presented elsewhere in this book. Measure the cell concentrations of the unfiltered culture and the GF/F filtrate. Combining the culture volume and the cell concentrations will give the total number of cells collected on the GF/F filter.

ii. Cell size measurements by photomicrography

1. Photographic slides for the cell size measurements are prepared from the same microscope slide used for the cell enumeration. Use Ektachrome 400 (Kodak). The optimum exposure settings must be empirically determined for the particular photomicrography system being used. Normally, short exposure time gives a better contrast between the dark background and bright cells. Note that the pictures should be taken from microscope fields chosen "randomly," the same as the AODC is done.

2. Fluorescent microspheres (Covalent Technology Corp., Ann Arbor, Mich.; Polyscience, Inc., Warrington, Pa.) of known diameter may help to check your size measurement criteria. Use microspheres with similar size, color and intensity of fluorescence to those of the AO-stained bacterial cells. Note that apparent size of the fluorescent microsphere is affected by the

fluorescence intensity and the photographic setting.^{3,8,9} Therefore, an exposure setting and optical filters different from those used for the AO-stained bacterial cells may need to be tried out, in order to produce photographs similar to those of the AO-stained bacterial cells with respect to the fluorescence intensities, contrast, and overall darkness.^{3,9} The microsphere suspensions usually need to be sonicated and diluted before use. Make microscope slides and take pictures of the microspheres as described above.

3. Take pictures of a stage micrometer (10 μm spacing). It is to be used as a standard scale when projecting the photographic slides on a screen for the size measurements (see Step 4).

4. Set up the slide projector in such a way that 1 μm in the real scale is enlarged to 1 cm. Measure and record the lengths and widths of the projected cell images to the nearest 1-mm unit. Note that no cells from a slide should be excluded. However, you may measure cell images from a part of the projection, when there are too many cells in a slide. When measuring the size, it is important to use a consistent criteria for where-to-cut (see Notes and Comments). Strict cutting-off of the fuzzy "halo" of the cell images under-measures by ca. 10% in linear scale; therefore, individual operators should establish and check the cell-boundary criteria.⁹ Large sample sizes are required for differentiation between two populations (see Notes and Comments).

5. There are a few different ways to calculate cell volumes from the linear measurements. Normally, cells of the same length and width are assumed to be "spheres" (Equation 1, see below) and the other elongated forms are treated as "rods." However, in practice, it is not that simple since there are occasionally cells of some irregular shapes, e.g., crescent, curvy filament, and etc. Consideration should also be given to

whether the model rod is a cylinder with two spherical ends (Equation 2), or an oblate spheroid of a cylindrical body with tapered ends (Equation 3). Following are the formulas for:

(1) volume of sphere (formula 1) = $(4/3) \times \pi \times R^3$

(2) volume of cylindrical rod (formula 2) = $\pi \times (3L-W)/3 \times (W/2)^2$

(3) volume of oblate spheroid (formula 3) = $(4/3) \times \pi \times (L/2) \times (W/2)^2$

where R = radius, L = length, W = width, and L > W.

6. From the size measurements, cell size-frequency distribution of the cells is obtained. From the cell enumeration (cell number ml⁻¹) and the culture volume (ml), total cell numbers in the unfiltered culture and in the GF/F filtrate are obtained. Combining the total cell number and the size-frequency distributions will give the total biovolume before and after the GF/F filtration, and the difference is the biovolume that is collected on the GF/F filter.

7. The biovolume conversion factor (biovolume-to-biomass conversion) is calculated as the carbon and nitrogen biomass from the CHN analysis, divided by the biovolume of the cells collected on the GF/F filter. The biomass conversion factor for cell number (cell number-to-biomass conversion), i.e., "per cell biomass", is calculated as the biomass divided by the number of cells collected on the GF/F filter.

IV. NOTES AND COMMENTS

1. See III A.

The type GS filter is made of mixed esters of nitrocellulose, and particles shed from the filter contain significant amount of carbon and nitrogen. Controls for the particle shedding are made from the separate preparation of GS filtrate (see below for CHN controls and blanks). Alternatively, boiling for 15 min three times in distilled water was recommended to minimize the shedding.⁸

Inoculum may be prepared with larger pore-size Nuclepore filters, e. g, 0.8 μm or 1.0 μm to include larger bacterial cells. However, filtrate of the pore size $> 0.8 \mu\text{m}$ often contains substantial amount of cells other than heterotrophic natural bacteria.

2. See III B.i. Glass fiber filters by no means collect all the particles of bacteria size. The retention efficiency varies among individual filters, filter types, and brands. There is no clear-cut nominal size (diameter) for the retention. The GF/F filtrate contains bacterial cells of almost all size classes present in the unfiltered culture, but the size-frequency distribution is skewed to smaller cells.⁹ The cells collected on the GF/F filter is the fraction (from ca. 2/3 to 1/2) of the total cells in the culture. To correct for the cells passing through the GF/F, it is important to measure cell concentrations and size-frequency distributions from the GF/F filtrate as well as from the unfiltered culture (see III C ii.).

3. See III B.ii. Blanks and filter shedding controls are subtracted from samples. Internal standards are for an independent check of the CHN analysis. Since the standard contains both carbon and nitrogen, CHN analysis of the standard will produce a calibration factor (i.e., expected amount vs. measured amount) for carbon and nitrogen, separately. The two calibration factors should be reasonably close to 1, as well as close to each other. The raw CHN data are multiplied by the calibration

factor. However, there may be more than one way to incorporate those control and blank CHN samples into the biomass calculation.⁹

4. See III C.i. Because the samples have various cell concentrations, the sample volumes to be filtered for AO slides should be adjusted to yield optimum cell densities on a Nuclepore filter (normally a few 10^6 cells on a filter). The sample size for AODC is very small compared to the population size, therefore it is important to reduce the sampling error. Optimum number and the level (e.g., subsample, slide, or fields) of replicates in the direct counting method are discussed by Kirchman et al.¹² and Kirchman (this volume).

5. See III C.ii.

Since image of bacterial cells have "halo" when projected on a screen, certain criteria must be applied consistently for the cell size measurements. To aid the consistency, picture qualities (sharpness, contrast, overall darkness, etc) should be reasonably same. For this reason, photographic settings should not be changed throughout an experiment.

Size-frequency distributions typically have a wide-spread range of size classes.¹³ A certain minimum sample size is required for a resolution of statistical parameters between two populations. For a better estimation and comparison of the population statistics, a relatively large size of samples is recommended (See ref. 14 for related discussion).

6. Concluding remarks:

An accurate derivation of a biomass conversion factor is primarily

based on the accurate measurements of the cell size. Different sizing methods have different resolution power.⁵⁻⁷ Apparent cell size varies with the sample preparation and sizing method used.^{5,13,15} It is not a simple matter to isolate true variations due to those artifacts. Among the current methods for biovolume measurement, image analysis and photomicrography are generally known to yield more realistic measurements.^{3-5,9,13,15,16} However, the measuring criteria and the consistency seem equally or even more important than the method used.

Table 1 lists directly measured conversion factors published to date (conversion factors from partial measurements employing indirect estimations are not listed). There are variations between studies as well as within a study. The variation may be a true variation because of different organisms (different species compositions) and different physiological state of the cells. However, a considerable fraction of the variation may be attributable to the large potential errors in size measurements. Unfortunately, no absolute calibration is currently available for biovolume measurements. Use of per-cell biomass conversion factors, being free of the bias due to biovolume measurements, would be more practical for natural planktonic bacteria. Cellular C and N contents ratios (C/N) are relatively less variable regardless of the methods used; mostly in the range from ca. 3.5 to 6.5 except for a few cases. Those high C/N ratios may be due to interference from detritus or non-bacterial organisms.

A trend of higher conversion factors for smaller cells were reported by some, whereas no such trend was found by others (see Fig.2 of reference 8). It has been argued whether high conversion factors are physiologically possible.¹⁷ Since no inter-calibration method is currently available between studies where different methodologies and different source organisms were employed, it is not easy to compare the

conversion factors obtained in diverse studies.^{8,16} A conversion factor is most useful when applied to a set of data obtained using similar procedures, making the conversion factor internally consistent as well as convenient.

V. REFERENCES

1. Watson, S.W., Novitsky, T.J., Quinby, H.L, and Valois, F.W., Determination of bacterial number and biomass in the marine environment, *Appl. Environ. Microbiol.*, 33, 940, 1977.
2. Cho, B. C., and Azam, F., Major role of bacteria in biogeochemical fluxes in the ocean's interior, *Nature*, 332, 441, 1988.
3. Scavia, D., and Laird, G.A., Bacterioplankton in Lake Michigan: dynamics, controls, and significance to carbon flux, *Limnol. Oceanogr.*, 32, 1017, 1987.
4. Bjornsen, P.K., Automatic determination of bacterioplankton biomass by image analysis, *Appl. Environ. Microbiol.*, 51, 1199, 1986.
5. Bratbak, G., Bacterial biovolume and biomass estimation, *Appl. Environ. Microbiol.*, 49, 1488, 1985.
6. Kogure, K., and Koike, I., Particle counter determination of bacterial biomass in seawater, *Appl. Environ. Microbiol.*, 53, 274, 1987
7. Nagata, T., Carbon and nitrogen content of natural planktonic bacteria, *Appl. Environ. Microbiol.*, 52, 28, 1986.
8. Nagata, T., and Watanabe, Y., Carbon- and nitrogen-to-volume ratios of

bacterioplankton grown under different nutritional conditions, *Appl. Environ. Microbiol.*, 56, 1303, 1990.

9. Lee, S., and Fuhrman, J.A., Relationships between biovolume and biomass of naturally derived marine bacterioplankton, *Appl. Environ. Microbiol.* 53, 1298, 1987.

10. Haldal, M., Norland, S., and Tumyr, O., X-ray microanalytic method for measurement of dry matter and elemental content of individual bacteria, *Appl. Environ. Microbiol.*, 50, 1251, 1985.

11. Hobbie, J.E., Daley, R.J., and Jasper, S., Use of Nuclepore filters for counting bacteria by fluorescence microscopy, *Appl. Environ. Microbiol.*, 33, 1225, 1977

12. Kirchman, D., Sigda, J., Kapuscinski, R., and Mitchell, R., Statistical analysis of the direct count method for enumerating bacteria. *Appl. Environ. Microbiol.*, 44, 376, 1982.

13. Fuhrman, J.A., Influence of method on the apparent size distribution of bacterioplankton cell: epifluorescence microscopy compared to scanning electron microscopy, *Mar. Ecol. Prog. Ser.*, 5, 103, 1981.

14. Sokal, R.R., and Rohlf, F.J., *Biometry*, W.H. Freeman and Co., San Francisco, 1969.

15. Montesinos, E., Esteve, I., and Guerrero, R., Comparison between direct methods for determination of microbial cell volume: electron microscopy and electronic particle sizing, *Appl. Environ. Microbiol.*, 45, 1651, 1983.

16. Psenner, R., From image analysis to chemical analysis of bacteria: a long term study? *Limnol. Oceanogr.*, 35, 234, 1990.

17. Joint, I.R., and Pomroy, A.J., Activity of heterotrophic bacteria in the euphotic zone of the Celtic Sea, *Mar. Ecol. Prog. Ser.*, 41, 155, 1987.

Table 1. Conversion factors directly determined from biovolume and biomass measurements.

source ^a	culture ^b	method for biovolume ^c	biomass ^d	cell volume	conversion factor		C/N	ref
				(μm^3) mean (range)	(pg C μm^{-3}) mean (range)	(pg N μm^{-3}) mean (range)		
Pure/SW	ENR	SEM,OG,EFPM	CHN	0.376 (0.11-0.71)	567 (238-964)	104 (45-202)	5.5 (4.6- 6.7)	5
FW/EST	ENR/UN	EFM,IA	CIRA	0.195 (0.08-0.34)	351 (178-728)	ND	ND	4
FW	UN	EFM,OG	CHN	0.163 (0.09-0.26)	106 (39-188)	25 (NS- 41)	4.8 (3.3- 6.8)	7
Pure	ENR	PC	CHN	1.253 (0.52-2.02)	209 (155-292)	62 (32- 94)	3.5 (2.8- 5.9)	6
SW	UN	PC	CHN	0.290 (0.17-0.53)	186 (84-353)	30 (11- 50)	6.3 (3.4-12.2)	6
SW	UN	EFPM	CHN	0.056 (0.04-0.07)	380 (230-600)	110 (60-220)	3.7 (2.5- 4.3)	9
FW	ENR/UN	EFPM	CHN	0.545 (0.17-1.75)	136 (59-252)	27 (7- 42)	5.5 (3.4-10.5)	8

a. Pure, pure culture; SW, seawater; FW, fresh water; EST, estuary brackish water.

b. ENR, enriched medium; UN, unenriched natural (*in situ*) medium.

c. SEM, scanning electron microscopy; OG, ocular grid; EFPM, epifluorescence photomicrography; EFM, epifluorescence microscopy; IA, image analysis; PC, particle counter.

d. CHN; CHN analysis, CIRA; combustion-infra red absorption method.

**DATE
FILMED**

11 / 20 / 92