CONF-901036--1

DE90 015159

THE FATE OF RADIONUCLIDES IN SEWAGE SLUDGE APPLIED TO LAND*

H. L. Boston, H. Van Miegroet, I. L. Larsen, A. E. Walzer, J. E. Carlton.

Submitted to the 4th International Conference on Environmental Contamination

October 1 - 4, 1990, Barcelona Spain

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

*The submitted manuscript has been authored by a contractor of the U.S. Government under contract No. DE-AC05-84OR21400. Accordingly, the U.S. Government retains a nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or allow others to do so, for U.S. Government purposes.

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

THE FATE OF RADIONUCLIDES IN SEWAGE SLUDGE APPLIED TO LAND*

H.L.Boston*, H.Van Miegroet*, I.L.Larsen*, A.E.Walzer*, J.E.Carlton*

ABSTRACT

Municipal sewage sludge containing up to 12 pCi/g (dry wt) ^{137}Cs , 20 pCi/g ^{60}Co , and 300 ppm U was injected in a pasture (43 Mg/ha) and sprayed over a young pine plantation (34 Mg/ha). In the pasture, radionuclides were largely retained in the upper 15 cm of the soil, and only about 15% moved below 15 cm. Sludge rapidly infiltrated the soil on the pine plantation. One year after application, at least 85% of the ^{137}Cs , ^{60}Co , and U were found in the upper 7 cm of the pine plantation, with only about 15% moving into the 7- to 15-cm strata. On-site total added radiation dose was 2 to 6 mrem/year. Radionuclides were not detected above background in soil solutions at ~50 cm depth or in shallow down-gradient groundwater wells. Surface runoff from application areas did not have elevated radionuclide concentrations. Concentrations of these radionuclides increased slightly in vegetation on treated sites, and uranium was notably higher in earthworms.

*Research supported by the U.S. Department of Energy and the City of Oak Ridge.

*Environmental Sciences Division, Oak Ridge National Laboratory, Oak Ridge, TN 37831-6351, USA. The Oak Ridge National Laboratory is operated by Martin Marietta Energy Systems, Inc., for the U.S. Department of Energy under contract DE-AC05-84OR21400.

*Oak Ridge Associated Universities, Oak Ridge TN 37831, USA.

INTRODUCTION

Sewage treatment plants receive radionuclides from industry, hospitals, and from atmospheric fallout that may be removed and concentrated in the sewage sludge (ref 1-3). Current U.S. regulations for radionuclide releases to sanitary sewers consider neither accumulation of radionuclides in sewage sludge nor implications for sludge disposal. Therefore, radionuclides have accumulated in sludge destined for disposal by land application (ref 4). The environmental fate of sludge constituents has been the subject of numerous investigations (ref 5,6). However, the fate of radionuclides in sewage sludge applied to land has received little attention thus far.

Digested sewage sludge from the City of Oak Ridge was applied to a pasture and a 4-year-old loblolly pine plantation on the U.S. Department of Energy's Oak Ridge Reservation (Oak Ridge, TN USA) to improve soil fertility and plant growth (ref 7). During this time the sludge was contaminated with ^{137}Cs , ^{60}Co , ^{131}I , and depleted U from industrial and medical sources. We measured the activity of ^{137}Cs and ^{60}Co and the concentration of U in soils, soil water, groundwater, surface runoff, vegetation, and earthworms to determine the fate of these nuclides.

METHODS

Anaerobically digested sludge from the City of Oak Ridge was injected to 10 cm soil depth in the pasture at a rate of 43 Mg/ha from December 1986 through May 1988. The sludge contained about 12 pCi/g (dry wt) ^{137}Cs , 20 pCi/g ^{60}Co , and 110 ppm U from industrial releases. Sludge was sprayed over a 4-year-old pine plantation (34 Mg/ha) from June 1988 through June 1989. This sludge contained about 2 pCi/g ^{137}Cs , 5.7 pCi/g ^{60}Co , and 300 ppm U. In the pasture, the upper 15 cm and the 15 - 30 cm soil depth were sampled following application. Vegetation (grass) was sampled at the end of application and about 2 years later. The upper 15 cm of soil on the pine plantation was sampled following application and again in May 1990. In November 1989, the upper ~ 45 cm of soil was sampled. During May 1990, new-growth pine needles were collected.

During sludge application, soil solution at 40- to 60-cm depth was collected with suction lysimeters. Samples of surface runoff from active application areas were collected during rainstorms. Groundwater was collected from shallow wells (<10 m) up- and down-gradient of the pine plantation. Earthworms were also collected from the pine plantation. Similar samples were collected from untreated (reference) areas adjacent to the application sites.

^{137}Cs , ^{60}Co , ^{40}K , and ^7Be were determined by spectrometry (IGe detector) calibrated with standards of known activities and geometries. When possible samples were counted to reduce counting error to <10%. Total U was determined by neutron activation (sludge and soil) or ICP-mass spectrometry (water, vegetation, and earthworms) and calibrated with reference materials.

RESULTS

Aside from the radionuclide content, Oak Ridge sludge was typical of municipal sewage sludge in the U.S. (ref 7) and was applied at rates typical for land application. We found that sludge metals were rapidly immobilized and retained in the upper layers (15 cm) of the soil (ref 7), as has been observed elsewhere (ref 5). On the tree plantation, U, ^{137}Cs , and ^{60}Co were retained in the upper 15 cm of the soil (Table 1). In the pasture about 15% of the added U, ^{137}Cs , and ^{60}Co occurred below 15 cm. We could not determine if this finding reflected migration or variable injection depth. The upper 15 cm of soil on the pine plantation was sectioned (Table 1). The profile showed that radionuclides from sludge sprayed over the surface rapidly moved into the soil. Although U averaged 300 ppm in the sludge, the concentration in the surface 2 cm of the soil averaged only about 25 ppm (Table 1) when application ended. Following treatment, ^{137}Cs seemed to be retained in the upper 2 cm of soil, whereas some ^{60}Co clearly moved to the 7 to 15 cm strata. About one year after application, the majority of U, ^{137}Cs , and ^{60}Co was still in the upper 7 cm of the soil, and the extent of movement of each seemed similar. Although the average concentrations of ^{137}Cs , ^{60}Co , and U in the 7- to 15-cm depth were not significantly greater for the treated area (Table 1), there was evidence of radionuclide movement to this strata from individual

soil cores.

Mass balance calculations for sludge radionuclides on the pine plantation (based on soils collected in May 1990) accounted for all of the added radionuclides in the upper 15 cm of the soil (ref 8). Total added radiation dose (all pathways, continuous occupancy) was about 2.5 and 5.5 mrem/year for the pine plantation and pasture, respectively, and was primarily due to short-lived ^{60}Co (ref 8).

Vegetation on application sites had elevated concentrations of N, P, K, and Ca but not of heavy metals (ref 7). Grass collected from the pasture site was slightly enriched with ^{137}Cs following application and was slightly enriched in U and ^{137}Cs almost 2 years later (Table 2). New growth pine needles collected about 1 year after application had higher concentrations of ^{137}Cs , ^{60}Co , and U than plants on the reference site. The concentrations of these elements were not greatly above background levels and were small compared with naturally occurring isotopes (e.g., ^{40}K and ^7Be).

Soil water on the application sites had high NO_3^- concentrations but did not contain elevated concentrations of metals (ref 7). Soil water analyzed for ^{137}Cs and ^{60}Co on two occasions contained <5 pCi/L of each nuclide. At times, some down-gradient groundwater wells at the pine plantation contained up to 2 times the concentration of U of up-gradient wells, but concentrations were generally low (<0.1 ppm). ^{137}Cs and ^{60}Co were never detected in down-gradient wells (<2.5 pCi/L). Surface runoff from application sites was enriched in NO_3^- , P, BOD, and fecal coliform bacteria but never contained detectable ^{137}Cs or ^{60}Co (i.e., <5 pCi/L).

Earthworms and other soil organisms can accumulate metals (ref 9) and may transfer these to other organisms. Earthworms from the treated area of the pine plantation contained 0.4 pCi/g (dry wt) ^{137}Cs and 0.6 pCi/g ^{60}Co , concentrations near our limit of detection. Earthworms from the treated area contained about 3 ppm U (dry wt), compared with about 0.5 ppm U for the reference area.

Radionuclides applied to soils in sewage sludge are diluted and immobilized in the soil. On these sites the radionuclides applied with sludge increased radiation dose by only about 2 to 6 mrem/year. Concentrations of radionuclides in the vegetation were not greatly increased. Radionuclides were neither moving off-site with surface runoff nor moving into the groundwater.

Table 1 Radionuclides in soils (dry wt basis). Values are means $\pm$ 1 S.D. or ranges in [] for ^an = 6 treatment, 3 reference; ^bn = 3 treat., 3 ref.; or ^cn = 6 treat., 4 ref. samples.

	Soil depth (cm)	U (ppm)		¹³⁷ Cs (pCi/g)		⁶⁰ Co (pCi/g)	
		Treatment	Reference	Treatment	Reference	Treatment	Reference
Pasture ^a	0-15	6.2 $\pm$ 1.4	3.3 $\pm$ 0.3	1.24 $\pm$ 0.44	0.43 $\pm$ 0.13	0.26 $\pm$ 0.09	<0.07
	15-30	3.8 $\pm$ 0.4	3.3 $\pm$ 0.4	0.28 $\pm$ 0.26	<0.11	[<0.03-0.06]	<0.02
Pine Plantation ^b							
Nov. 1989	0-15	7.9 $\pm$ 0.6	3.8 $\pm$ 0.7	0.48 $\pm$ 0.06	0.43 $\pm$ 0.01	0.09 $\pm$ 0.01	<0.04
	15-30	3.6 $\pm$ 0.4	3.4 $\pm$ 0.7	0.04 $\pm$ 0.01	0.05 $\pm$ 0.02	<0.011	<0.015
	30+	3.7 $\pm$ 0.4	3.4 $\pm$ 0.5	<0.012	<0.012	<0.010	<0.010
Pine Plantation (0 - 15 cm only)							
July 1989 ^c	0-2	25.3 $\pm$ 12.2	3.9 $\pm$ 0.6	0.88 $\pm$ 0.10	0.45 $\pm$ 0.01	0.53 $\pm$ 0.16	<0.05
	2-7	11.4 $\pm$ 4.0	4.0 $\pm$ 1.4	0.49 $\pm$ 0.14	0.42 $\pm$ 0.04	0.12 $\pm$ 0.02	<0.03
	7-15	5.4 $\pm$ 1.0	4.0 $\pm$ 1.1	0.33 $\pm$ 0.19	0.43 $\pm$ 0.15	[<0.02-0.10]	<0.03
May 1990 ^c	0-2	23.0 $\pm$ 11.9	4.1 $\pm$ 1.2	0.77 $\pm$ 0.18	0.44 $\pm$ 0.04	0.31 $\pm$ 0.16	<0.03
	2-7	9.7 $\pm$ 5.3	4.0 $\pm$ 1.1	0.66 $\pm$ 0.12	0.44 $\pm$ 0.14	0.10 $\pm$ 0.07	<0.01
	7-15	5.0 $\pm$ 1.4	4.0 $\pm$ 1.1	0.52 $\pm$ 0.15	0.36 $\pm$ 0.19	[<0.01-0.05]	<0.01

Table 2 Radionuclides in vegetation (dry wt basis). Values are means $\pm$ 1 S.D. or ranges in [] for ^an = 3 treatment and 3 reference or ^bn = 6 treatment and 2 reference samples. NA = not available.

	U (ppm)	¹³⁷ Cs (pCi/Kg)	⁶⁰ Co (pCi/Kg)	⁴⁰ K (pCi/Kg)	⁷ Be (pCi/Kg)
Pasture-1988 ^a					
Treatment	NA	61 $\pm$ 52	[59-<200]	27400 $\pm$ 5800	5360 $\pm$ 2930
Reference	NA	[61-<200]	<86	18900 $\pm$ 4420	13800 $\pm$ 8200
Pasture-1990 ^a					
Treatment	[<0.01-0.03]	[<14-34]	<16	26100 $\pm$ 5200	1490 $\pm$ 662
Reference	<0.01	[<16-26]	<24	22300 $\pm$ 1660	3440 $\pm$ 383
Pines-1990 ^b					
Treatment	0.11 $\pm$ 0.08	[<15-34]	[<12-29]	3618 $\pm$ 287	1913 $\pm$ 67
Reference	0.03 $\pm$ 0.01	<15	<15	3811 $\pm$ 602	2381 $\pm$ 359

REFERENCES

1. H M Prichard, T F Gesell, E Davis, Am J Public Health **71**, 47 (1981)
2. K R Imhoff, P Koppe, F Dietz, Water Res **22**, 1059 (1988)
3. B Erlandsson, B Bjurman, S Mattsson, Water Air Soil Pollut **45**, 329 (1989)
4. U.S. Nuclear Regulatory Commission Inspection Report 70-143/86-44 (U.S. NRC, Atlanta, 1986) p 10.
5. D E Williams, J Vlamis, A H Puktie, J E Corey, Soil Sci **143**, 124 (1987)
6. D W Cole, C L Henry, W L Nutter, The Forest Alternative for Treatment and Utilization of Municipal and Industrial Wastes (University of Washington Press, Seattle, 1986) p 582.
7. H Van Miegroet, H L Boston, D W Johnson, Proceedings 12th Annual Madison Waste Conference (University of Wisconsin Press, Madison, 1989) p 571.
8. H L Boston, H Van Miegroet, I L Larsen, A E Walzer, Fate of Radionuclides in Sewage Sludge Applied to Forests and Pastures (TM-11601, Oak Ridge National Laboratory, Oak Ridge, in press).
9. M B Levine, A T Hall, G W Barrett, D H Taylor, J Environ Qual **18**, 411 (1989)

SUPERDEFORMATION IN THE MERCURY REGION

DE90 017810

R. R. Chasman

Physics Division, Argonne National Laboratory, Argonne, IL 60439

ABSTRACT

A brief history of superdeformation is presented. Calculations of superdeformed well depths, single particle energy level diagrams at $I=0$, orbital alignments as a function of rotational frequency, and dynamical moments of inertia are presented. The sensitivity of calculations to the details of the spin-orbit interaction are examined. Comparisons with experimental results are made.

Received by OSTI
SEP 27 1990

INTRODUCTION

Superdeformed nuclides (axis ratio $>1.5:1$) were first proposed¹⁾ some twenty five years ago as the explanation of the phenomenon²⁾ of fission isomerism seen in the actinides at $I=0$. The fission isomers are nuclear states associated with very elongated shapes (axis ratio of $\approx 2:1$). Although the calculations using the Strutinsky method¹⁾ give superdeformed minima in the energy surfaces of many of the actinides, these calculations do not seem to explain some of the spectroscopic information obtained in the study of the fission isomers; most notably the magnetic moment measurements³⁾ in ^{230}Pu . This failure of the calculations has given rise to much theoretical speculation about the nature of the effective potential at large deformations; however the difficulty of obtaining sufficient spectroscopic information on superdeformed actinides has made it impossible to resolve this problem. A detailed, useful treatment of superdeformation in the actinides can be found in the review⁴⁾ of Bjornholm and Lynn.

In the past few years, heavy ion reactions have been used to produce⁵⁾ several superdeformed nuclides at high spin near ^{152}Dy . Calculations using the cranked Strutinsky method suggested^{6,7,8,9)} the existence of superdeformed states in several nuclides of this region at high spin. These nuclides decay from superdeformed to less deformed shapes at high spins ($I \approx 25$) and little is known about the energy differences of rotational bandheads in this region. It is interesting to note that the same large gap in the neutron single particle spectrum ($N \approx 86$) that gives rise to these superdeformed nuclides at high spin also explains¹⁰⁾ many features of the mass yields observed in the fission of the actinides. This large shell gap gives a minimum in the energy surface at high spins because the moments of inertia are much larger for superdeformed states than they are for normally deformed states and the rotational energy is lower. In the fission case, the superdeformed shapes are favored because of the coulomb repulsion of the nascent fission fragments.

We have carried out calculations on nuclides in the $A=190$ region, using the cranked Strutinsky method, and predicted¹¹⁾ that

DISCLAIMER

MASTER
DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

The submitted manuscript has been authored by a contractor of the U. S. Government under contract No. W-31 109-ENG-38. Accordingly, the U. S. Government retains a nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or allow others to do so, for U. S. Government purposes.

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

there are many nuclides in the Hg region in which superdeformed states are experimentally accessible at moderate spins ($I \approx 40$). Based on these predictions, an experimental study¹²⁾ was made and superdeformation was first discovered in ^{191}Hg in this mass region. We also found in our calculations that this region differs substantially from the $A=150$ region in that the superdeformed minimum persists to low spin. There are other^{13,14,15)} calculations that show superdeformed minima for nuclides in this region at $I=0$; but the stability of superdeformed states at $I=0$ remains an important open theoretical and experimental question and is an area of active research. As of now, superdeformed states with $I=0$ have not been seen in this region. Our calculations suggest that the superdeformed states are at least 4 MeV higher in energy than the ground state in the $A=190$ region, at $I=0$. This is roughly twice as much excitation energy as the fission isomers seen in the Uranium isotopes and implies an order of magnitude increase in E2 decay probability from the superdeformed state if one assumes the same transition matrix element.

THE MERCURY REGION

In the Strutinsky method, the total energy is the sum of a smooth contribution to the nuclear energy, given by a liquid drop model; and a fluctuating part the "shell correction", that measures the difference between the level density for the nucleus of interest and some smoothed level density. When the level density is low at the fermi level, we get extra stabilization. In Fig. 1, we show the proton and neutron levels appropriate to the superdeformed Hg region. One can see that the shell corrections are particularly favorable for $Z=80$ and for $112 < N < 120$. In our calculations, we use the Myers-Swiatecki¹⁶⁾ parameterization of the liquid drop model and a Woods-Saxon single particle potential for the level density calculation. This parameterization⁹⁾ of the Woods-Saxon potential is based on single particle energy level spacings obtained from a many-body analysis of near yrast high spin states in the ^{148}Gd region.

The energy is calculated on a grid in deformation space. The ground state shape is that shape with lowest total energy. Excited states are predicted at all local minima in the energy surface. Angular momentum is introduced by adding a cranking term ωJ_x ¹⁷⁾ to the single-particle Hamiltonian and the calculation is extended thereby to finite angular momenta. It is worth noting that superdeformed states come down in energy relative to less deformed ones as the angular momentum increases, because they have larger moments of inertia and hence less rotational energy.

There are some caveats that apply to the Strutinsky method. Ground state masses are calculated with an accuracy of ≈ 0.8 MeV. Also, there are calculations in the $A=146$ region showing a reflection asymmetric minimum about 1.5 MeV deeper than the reflection symmetric minimum, but experimentally the nuclides do not show reflection asymmetry in their ground states. This suggests that a well depth of at least 2 MeV is needed before one has a reasonable prediction of

shape stability. On more general theoretical grounds, one expects that there will be strong mixing of configurations with different shapes when the barriers between the different shapes are small. When one goes beyond the Strutinsky method, e.g. with a generator-coordinate calculation, one finds just this sort of shape mixing. A second caveat is that one may not have looked at the region of deformation space where the true minima are found; i.e. by neglecting octupole, hexadecapole 2_5 -pole or 2_6 -pole degrees of freedom, one might get a totally erroneous picture of nuclear structure, even if the Strutinsky method were exact.

With these qualifications in mind, we turn to our¹¹⁾ calculations of superdeformation in the Hg region. These calculations include only quadrupole and hexadecapole deformations. In Fig. 2, we display the calculated superdeformed well depth at $I=40$ in the $A=190$ region. The first thing to note is that there is a large region of nuclides with a well developed superdeformed minimum at a quadrupole deformation of $\nu_2=0.5$, corresponding to an axis ratio of 5:3. We chose a 2 MeV well depth as a minimum requirement for the reasons cited above. In Fig. 3, we display the excitation energy of the superdeformed state relative to yrast at $I=40$. We have assumed that the lower the excitation energy of the superdeformed state, the easier it is to produce. Taking both of these figures into consideration, we are led to a prediction of the most experimentally accessible superdeformed states being in ^{191}Hg and ^{192}Hg . These predictions are borne out by the experimental observation^{12,18,19)} of superdeformation in these nuclides. Subsequently, superdeformed states have been found in ^{190}Hg 20), ^{193}Hg 21), ^{194}Hg 22), ^{193}Tl 23), ^{194}Tl 24), ^{194}Pb 25) and ^{196}Pb 26). All of these nuclides are expected to have superdeformed minima (cf. Fig. 2). In Fig. 4, we display the energy surface obtained with the Strutinsky method for ^{191}Hg at different values of the angular momentum. We have made a one dimensional cut of the energy surface so as to connect the minima. We get qualitatively similar results for the other known superdeformed nuclides in this region.

However, there is a serious problem! Our calculations indicate that the superdeformed minimum in ^{194}Hg and ^{196}Hg is roughly 4 MeV above yrast even at $I=40$. Conventionally, one assumes that the superdeformed band is populated in that range of angular momentum states in which it is near yrast and fission competition is not yet serious. The population of superdeformed bands in Pb is not consistent with our calculation if one makes this assumption. This suggests that either our single particle scheme is not quite right or that highly excited superdeformed states can be populated in heavy ion reactions.

Here we explore the possibility that the single particle potential is not quite right. Many years ago Rost²⁷⁾ pointed out that one could improve the agreement of Woods-Saxon calculations with experimentally determined proton energy level spacings in the heavy elements by using a spin-orbit radius parameter that is $3/4$ the value

used for the central part of the potential. As many of these actinide levels are at or even below the fermi level in the superdeformed $A=190$ region, it is worthwhile to explore the effects of this parameterization on superdeformation in the Hg region. We note that this parameterization gives results similar to our earlier work for the ground state deformations in the $A=180-190$ region. Whether or not this parameterization is valid, these calculations provide an estimate of the sensitivity of the model to reasonable changes in the parameters. We have carried out such a calculation. In Fig. 5, we compare the spherical single particle energy levels obtained with $R_{s.o.}=R_0$, and the spherical levels obtained using $R_{s.o.}=0.75R_0$. The level orderings and spacings are quite similar apart from an increased splitting of the $h_{11/2}-h_{9/2}$ pair and a decreased splitting of the $f_{7/2}-f_{5/2}$ pair with the Rost parameterization. It should be noted that there is evidence for a decreased $f_{7/2}-f_{5/2}$ splitting from the (d,p) studies of Pb and also from the observation²⁸⁾ of the $1/2-[521]$ orbital in the $A=250$ region. In Fig. 6, we display the well depths associated with the superdeformed minima at $I=40$, using the Rost spin-orbit radius. Comparing with Fig 2, one immediately sees that there are many more superdeformed nuclides with well depths greater than 2 MeV. Also, the deepest wells are on the order of 5 MeV and are ≈ 2 MeV deeper than we found with our usual parameterization, for the nuclides in the vicinity of ^{199}Po . We emphasize that the deep superdeformed well depths are retained in the Hg-Tl-Pb isotopes in which superdeformed bands are known. Unfortunately, the excitation energy of the superdeformed minima are predicted to be as high or higher in this calculation than the values shown in Fig. 3. This calculation does not provide an easy explanation of the population of the superdeformed bands in ^{194}Pb and ^{196}Pb . To put things more strongly, there is a maximum in E^* at $I=40$, for the superdeformed band for neutron numbers $110 < N < 118$, when $Z=81$ or $Z=82$. With our original parameterization, this maximum always occurs for $Z=82$. This means that the superdeformed states in Pb should be the hardest to produce according to this criterion. If the Rost parameterization is the correct one, we might argue that the decay out of the superdeformed minimum would be inhibited by the existence of a such deep wells and perhaps one might observe transitions in the superdeformed rotational bands of nuclides near ^{199}Po . Both the Rost and the conventional spin-orbit interactions predict superdeformed minima in this mass region. However, there are large differences; i.e. the results are sensitive to the details of the potential. It is then important to fix the potential as accurately as possible.

OTHER SURPRISES

There have been several other surprises in the experimental study of superdeformed bands in the Hg region. The first has been how easy it is to produce these bands in heavy ion reactions. The second is the large regular increase in the dynamic (second derivative) moment of inertia with rotational frequency. The third is the observation of almost exactly the same transition energies in superdeformed bands of different nuclides in the Hg region. This

feature was first noted in the superdeformed bands of the A=150 region.

We consider the dynamic moments of inertia. In the high spin superdeformed rotational bands of the Dy region one finds very little variation in the moment of inertia as a function of angular momentum. In the Hg region, there is in some instances a $\approx 40\%$ increase in the moment of inertia in the superdeformed bands. In the A=150 region, the superdeformed bands terminate at $I \approx 25$, while they continue to rather low spin in the Hg region. One might assume that pairing plays a rather different role in the two regions and that a pairing attenuation is related to the increase of the moment of inertia in the Hg region. The role of pairing is quite paradoxical in superdeformed nuclides. In order that there be a superdeformed minimum in a nucleus, the level density must be small near the Fermi level and pairing correlations must be small. On the other hand, a superdeformed shape has a large moment of inertia and small energy differences between members of a rotational band. This means that a small pairing correlation energy that decreases with increasing angular momentum could have a dramatic effect^{29,30)} on the moments of inertia of superdeformed states without really affecting their structure.

Because the level density is low for superdeformed shapes, a quasiparticle treatment of pairing forces is not appropriate in this case. If one uses a gap parameter to get a finite pairing correlation energy, this gives excessive pairing correlation and too low a moment of inertia at low spin. The approach that we take³¹⁾ is to calculate the pairing correlation energy accurately at $\omega_0=0$ using the method of correlated quasiparticles³²⁾ which goes beyond the quasiparticle (BCS) approximation and is particularly appropriate when pairing effects are small. Motivated by the observed smooth rise in the moment of inertia in this region, we assume that the pairing correlation energy is smoothly attenuated as a function of increasing angular momentum. The functional form that we choose is an exponential attenuation, i.e. we set

$$E_{\text{pair}}(\nu_2, \nu_4, \nu_6, \omega) = E_{\text{pair}}(\nu_2, \nu_4, \nu_6, \omega_0) e^{-(\omega/\omega_0)^2}$$

at each point in the deformation space grid. We use a single parameter ω_0 for all nuclides in the A=190 region. Comparing with the experimental data, we choose $\omega_0=0.25$ MeV ($\hbar$)-1 and get a reasonable agreement with observed moments of inertia. This means that the pair correlation energy is reduced to $\approx 40\%$ of its ground state value at $I=30$. In Fig. 7, we compare the calculated and observed moments of inertia for ^{192}Hg , where the agreement is quite good and for ^{193}Hg in Fig. 8 where the agreement is not so good. It appears that one can account for the increase in the moments of inertia quite reasonably in this way. In doing this calculation, we

included 2_6 -pole deformations in addition to quadrupole and hexadecapole deformation. This shifted the superdeformed minima somewhat; giving much smaller values of V_4 than we obtained without the inclusion of 2_6 -pole deformations. It also lowered the excitation of the superdeformed state by several hundred keV in some instances; providing a nice example of the need to use an extended deformation space in Strutinsky calculations.

The problem of almost identical transition energies in different nuclides is not easy to understand. This was first seen^{33,34)} in the $A=150$ region with almost identical transition energies in $N=86$ isotones. For obvious reasons, such bands are sometimes called twinned bands. There have been many more such bands seen in the Hg region. The differences in the transition energies are just a few keV, which is considerably less than one expects from the rigid rotor estimate of the moment of inertia. A calculation accurate to a few keV goes far beyond my assessment of the accuracy of the cranked Strutinsky method. Stephens has emphasized³⁵⁾ that the incremental alignment in the twinned bands is either integer or half integer relative to ^{192}Hg . We have examined the calculated single particle neutron alignments for the orbitals near $N=112$. These single particle alignments as a function of rotational frequency are shown in Fig. 9. We do not seem to find orbitals with constant alignments, or with half integer or integer alignment. One expects to see such alignments with a pseudo-SU(3)^{36,37)} coupling scheme, which has been suggested as an explanation of this phenomenon. However, this is known to be only an approximate symmetry in deformed nuclides.

SUMMARY

To summarize, we find that the cranked Strutinsky method, used with some care, has led us to the prediction of a new region of superdeformed nuclides that are accessible in heavy ion reactions. The experimental study of this region has raised some interesting questions: (1) why are the transition energies in superdeformed bands so similar in different nuclides; (2) why do we populate superdeformed bands, when the calculations suggest that they are highly excited; (3) why is there a large increase in the second moment of inertia with increasing spin. We believe that the last problem is at least semi-quantitatively understood; the second question may be answered within the context of the cranked Strutinsky method; and that one must go beyond the Strutinsky method and beyond such approximate symmetry concepts as pseudo-SU(3) to deal with the first. There is a reciprocal interaction between theory and experiment that is making nuclear structure the stimulating area of research that it is today. It is just this interaction that leads us to suggest experiments to clarify the role of E^* in the production of superdeformed states. This means the study of nuclides with $Z=84$ and $Z=85$, with N near 115.

This work supported by the U.S. Department of Energy, Nuclear Physics Division, under contract W-31-109-ENG-38. The calculations were carried out on the NERSC computers at Livermore.

REFERENCES

1. V. M. Strutinsky, Arkiv Fys. 36, 629(1966); Nucl. Phys. A95, 420 (1967).
- 2) S. M. Polikanov et al., JETP (USSR) 24, 1464 (1962).
- 3) H. Backe et al., Phys. Rev. Lett. 42, 490 (1979)
- 4) S. Bjornholm and J. E. Lynn, Rev. Mod. Phys. 52, 725 (1980).
- 5) P. J. Twin et al., Phys. Rev. Lett. 57, 811 (1986).
- 6) K. Neergaard and V. Paskevich, Phys. Lett B59, 218 (1975).
- 7) J. Dudek et al., Phys. Lett B112, 1 (1982).
- 8) J. Dudek and W. Nazarewicz, Phys. Rev. C 31, 298 (1985).
- 9) R. R. Chasman, Phys. Lett B187, 219 (1987).
- 10) B. D. Wilkins et al., Phys. Rev. C 14, 1832 (1976).
- 11) R. R. Chasman, Phys. Lett. B242, 317 (1990).
- 12) E. F. Moore et al., Phys. Rev. Lett. 63, 360 (1989).
- 13) C. F. Tsang and S. G. Nilsson, Nucl. Phys. A140, 275 (1970).
- 14) M. Girod, J. P. Delaroche and J. F. Berger, Phys. Rev. C 38, 1519 (1988).
- 15) P. Bonche et. al, Nucl Phys. A500, 308 (1989).
- 16) W. D. Myers and W. J. Swiatecki, Arkiv for Fysik 36. 343 (1966).
- 17) M. Brack and B. K. Jennings, Nucl. Phys. A258, 264 (1976).
- 18) D. Ye et al., Phys. Rev. C 41, 13 (1990).
- 19) J. A. Becker et al., Phys. Rev. C 41, 9 (1990).
- 20) M. W. Drigert et al. (to be published).
- 21) E. A. Henry et al., Z. Phys. A335, 361 (1990).
- 22) M. A. Riley, Nucl. Phys. A512, 178 (1990).
- 23) P. B. Fernandez et al. (to be published).
- 24) F. Azaiez et al. (to be published).
- 25) M. J. Brinkman et al., Z. Phys. A336, 115 (1990).
- 26) K. Theine et al., Z. Phys. A336, 113 (1990).
- 27) E. Rost, Phys. Lett. 26B, 184 (1968).
- 28) I. Ahmad et al., Phys. Rev. Lett. 39, 12 (1977).
- 29) W. Nazarewicz, Symposium on Nuclei at High Spin, Copenhagen, October 1989.
- 30) T. Bengtsson, Symposium on Nuclei at High Spin, Copenhagen, October 1989.
- 31) R. R. Chasman, Phys. Lett 242B, 317 (1990).
- 32) R. R. Chasman, Phys. Rev. C 5, 29 (1972).
- 33) P. Twin, Symposium on Nuclei at High Spin, Copenhagen, October 1989
- 34) T. Byrski et al., Phys. Rev. Lett. 64, 1650 (1990)
- 35) F. S. Stephens et al., Phys. Rev. Lett. 65, 301 (1990).
- 36) K. T. Hecht and A. Adler, Nucl. Phys. A137, 129 (1969).
- 37) A. Bohr, I. Hamamoto and B. R. Mottelson, Phys. Scr. 26, 267 (1982).

Figure Captions

- Fig. 1 Proton and neutron single particle levels for superdeformed Hg region.
- Fig. 2 Depth of the superdeformed minimum for nuclides near $A=190$ at $I=40$. The continuous line is the 2 MeV contour; the dashed line is the 3.5 MeV contour. The numbers inside the figure are quadrupole deformations ν_2 of the superdeformed minimum. In each row, all nuclides to the right of a written value of ν_2 have the same value.
- Fig. 3 Excitation energy (E^*) of superdeformed state at $I=40$, relative to yrast. The dashed line is the 0.5 MeV contour; the continuous line, 2 MeV; the dotted line, 4.5 MeV. The numbers inside the figure are moments of inertia calculated without pairing.
- Fig. 4 A one dimensional cut of the Strutinsky energy surface of ^{191}Hg , as a function of angular momentum.
- Fig. 5 Comparison of proton spherical single particle levels for $R_{s.o.}=R_0$ and $R_{s.o.}=0.75R_0$.
- Fig. 6 Depth of superdeformed minimum at $I=40$ obtained with Rost spin-orbit parameterization. The continuous line is a 2 MeV contour; the dashed line, the 3.5 MeV contour; the dotted line, a 4.5 MeV contour.
- Fig. 7 Dynamic moment of inertia of ^{192}Hg as a function of rotational frequency. The continuous line is the calculation.
- Fig. 8 Dynamic moment of inertia of ^{193}Hg . See caption for Fig. 6.
- Fig. 9 Single particle alignments near $N=112$ at $\nu_2=0.50$, $\nu_4=0.02$. The neutron numbers and Ω^π values are for $\Omega=0$.

SINGLE PARTICLE STATES

$$v_2 = 0.50, \quad v_4 = 0.02$$

$$\omega = 0$$

Fig. 1

Fig. 2

Fig. 4

PROTON SINGLE-PARTICLE ENERGY LEVELS

Fig. 5

Fig. 6

Fig. 7

Fig. 8

Fig. 9