FINAL REPORT TO WESTGEC

Analysis and Synthesis of Models for Effects of Climate Change on Agricultural Systems

By

S. Geng, R. Plant, R. Loomis
Department of Agronomy and Range Science
University of California, Davis

DOE/ER/61010--T5

DE93 007949

July 27, 1992

OBJECTIVE

U S T I

1. Crop Model Development

Our objectives are (1) to develop a new integrative physiological-morphological model of the wheat crop that will behave realistically in high-CO₂ environments, and (2) to update the ALFALFA model to match the wheat model's photosynthetic structures and microclimates.

2. Impact Assessment:

Assess the impacts of climate and predicted climate change on dryland spring wheat yields and rice yields in the United States.

PRODUCT

1. Crop Model Development

Documented and developed mechanistic models of wheat (a new model) and alfalfa (modification of an existing model) that incorporate known physiological effects of CO₂, air temperature and soil water content within the framework of whole-plant growth and productivity.

2. Impact Assessment

Quantified the impact of weather change on United States rice and wheat yields.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

MASTER

APPROACH

1. Crop Model Development

Existing models, ALFALFA (Denison and Loomis, 1989) and COTGRO (FACE project; Amthor, unpublished), served as the starting point for new models of alfalfa and wheat responsive to variations in $[\text{CO}_2]$. These species will serve as reasonable analogs for all small grains and most forages (and pasture). The existing models are highly structured with considerable physiological and morphological detail (e.g., individual leaves in layered canopies, autonomous tillers) allowing realistic simulation of vegetation response to environment (e.g., direct and diffuse radiation, variable soil water supply). Time is advanced hourly allowing treatment of critical diurnal sequences of temperature, water status, radiation, and other factors. The models predict in a realistic way many of the main features of crop acclimation to changing environment. Key features required for environments with altered $[\text{CO}_2]$ include simulation of canopy profiles of water vapor (as a basis for simulation of leaf temperature, conductance, and transpiration) and of $[\text{CO}_2]$ (for photosynthesis).

The plan was to correct ALFALFA in its ability to perform in hot climates and extreme drought, where its predictions were under question, and then to add advanced routines for photosynthesis. This would be based on work of Farquhar's group (Farquhar and von Caemmerer, 1982; Evans and Farquhar, 1991) as employee by Amthor in COTGRO. Having examined and rejected existing models of wheat, a new WHEAT would be based on the same general approach as ALFALFA.

2. Impact Assessment

We have developed a general method to assess climate impacts on crop yields that involves a hierarchy of three types of models: a GCM, a weather generator and a crop simulation model. Although the direction and magnitude of climate change as a result of doubling of CO_2 is uncertain, this method can be reapplied if and when there is a better prediction of the expected climate change. This method may be applied with

any GCM output and any crop model. The easily used weather generator and a reliable crop model are the keys to this assessment method. We have produced a method that shows the aspects of weather and climate change. This is much more informative for assessment and mitigation purposes than perturbing one year of observed weather data and making broad statements about expected changes based on that one year.

RESULTS

1. Crop Model Development

ALFALFA was given extended temperature-response functions, a new senescence routine, and an updated water routine, and revalidated. The new version (1.5) was employed in a study of irrigation management for conditions of the lower Colorado basin with the finding that water saving is more efficiently accomplished by reduction of area planted than by deficit irrigation (Wallinga, 1991).

WHEAT is still under development. It employs a modular structure similar to ALFALFA to simulate standard competitive plants in a uniform field. The basic model runs with an hourly time advance. Organs are initiated from discrete apices of mainstem and tillers. New leaf 'units' (leaf, axillary bud, stem internode) arise during the vegetative stage, spikelets during the reproductive phase. Tillers are generated from basal axillary buds. Developmental rate depends on nonlinear functions of day length, current temperature, and substrate level. Our present hypothesis is that an effect of water status on development is not appropriate. Rates of initiation and numbers of primordia are influenced by "genetic" rules. At this point, the 'cultivar' has the general characteristics of a modern spring wheat ('Anza').

Organ growth is empirical with limitations by temperature, substrate, and water status. Respiration related to growth is calculated dependent upon biochemical composition characteristic of each tissue (Penning de Vries et al. 1974). At present, nitrogen is considered freely available and not limiting to growth.

Water is acquired from a layered soil profile infiltrated by rain or irrigation and depleted by an expanding root system. Water uptake follows van Bavel and Ahmed (1976). Transpiration is calculated from environmental conditions (radiation and energy balances, simulated wind profile, vapor pressure, and temperature), leaf area, and stomatal conductance. (Rapid iteration in a subroutine handles the fast kinetics of these processes.) Stomatal conductance is a function a root signal (Gollan et al., 1986), light, humidity, and $[CO_2]$.

Photosynthesis follows Farquhar's approach with Amthor's (1992) data on respiration. The model intrinsically accounts for effects of $[CO_2]$, $[O_2]$, light, temperature, and leaf nitrogen status on CO_2 assimilation and respiration (Amthor, submitted). Control of $[CO_2]$ internal to leaf needs further study, but it is clear that this approach is the correct one for studies of global change.

At present, the canopy is divided into sunlit and shaded leaves only. Clumping and direct and diffuse radiation are accounted in the treatment, however (Myneni et al. 1989). Expansion into a layered model of individual leaves (as in ALFALFA) will be necessary later to account for competition among tillers as independent units.

WHEAT is composed of functional subroutines coded in FORTRAN 77 controlled by a main program. Leaf physiology/photosynthesis and soil routines are mostly finished and need only further testing and documentation. By contrast, only the skeletons of the growth modules (apex, roots, grain) have been completed. The structure of the model is presented in Appendix A.

2. Impact Assessment

The Effect of Weather and Climate Change on United States Grain Yields:

The continued increase in world population and food demand means a greater responsibility for all countries to stabilize the food supply. While advances in efficiency of food production has occurred, unfavorable weather events in recent years and possible negative climate changes have greatly reduced confidence in technology for producing a stable food supply.

To assess the of weather impact on food security three areas must be addressed: (1) the quantification of the effects of weather variation on wheat yields and yield variability, (2) quantification of the possible impact of climate change on wheat yields, (3) examination of the interaction effect due to fertilizer technology and weather on yield variability. A hierarchy of three types of simulation models is used to assess these impacts for U.S. grains crops; GCM outputs are integrated with a dynamic weather simulation model and a crop model to evaluate weather impact on crop production.

Main results are presented in the following three sections (papers can be found in Geng and Cady, 1991):

(1) The Effect of Weather and Climate Change on United States Wheat Yield:

We have connected GCM model outputs with a dynamic weather simulation model and a crop model to evaluate weather impacts on crop production, especially wheat production in the United States. Results showed that wheat yield variation is greatly intensified and exacerbated by weather variation. When CO₂ is doubled, rainfall will decrease from 8 - 37%, annual mean minimum temperature will increase from 3°C to 8°C, and yields will decrease from 8 - 61%, depending on the location. Adjustments in planting dates and alternative cultivars were also tested. In most cases, the adjustments did not significantly improve yields. It has been postulated that higher technology input increased yield variation. Our findings indicated the phenomena are much more complicated than a simple cause and effect explanation. Superficially, a higher technology input is associated with an observed increase in yield variance. The reason for this association is not necessarily due to technology input per se but that the higher technology input system allows a crop to respond to a broad range of weather conditions. As a result, the increased fluctuation of weather generates an increased fluctuation of yield.

(2) Risk Assessment Under Current and Double CO₂ Conditions for United States Wheat Yields:

There are at least two problems in quantifying weather related risks associated with crop yields. First, long-term data are

generally not available. Second, even if the data were available, they were frequently confounded with factors other than weather conditions. In this study we have used a simulation approach to quantify two types of risks associated with cropping systems under current and double CO₂ scenarios. The first order risk is defined as the probability of crop failure in a particular environment. The second order risk is the degree of uncertainty of maintaining the mean yield for a growing condition (variance of yield). Results show that, overall, with the present cropping systems the first order risk increased by 20% with doubling of CO₂. Second order risk increased at locations that presently have the highest mean wheat yields in the Great Plains. Second order risk decreased at locations that currently produce the majority of wheat in the Great Plains. However, at those locations the mean yields greatly decrease and first order risk increases substantially with doubling of CO₂. The commonly suggested mitigation methods such as adjusting cultivars and planting dates did not alleviate the weather impact and associated risk sufficiently to lead to a productive, alternative cropping system.

(3) Weather Impact Assessment on U.S. Rice Yields and California Wheat Yields:

The U.S. produces 7.2 million metric tons of rice, 28% of which is exported to developing countries. In the U.S. rice is produced in California and the Gulf Coast states. It is anticipated that global climate change will have a major influence in agricultural practices and crop selection in these states. This study utilizes simulation techniques to quantify the potential magnitude of this influence. In addition, the impact of climate change on dryland spring wheat in California is evaluated. Results indicate that rice yields decrease by between 14% and 24% in the Gulf Coast states and between 11% and 21% in California. In both regions the decrease in rice yields were due primarily to the large increase in summer temperatures. On the other hand, dryland fall planted spring wheat yields in California increase by 62% and 125%. This is because of the increased rainfall and temperatures during the winter months in California.

LITERATURE CITED

- Amthor, J.S. 1992. CO₂ inhibits respiration in leaves of Rumex crispus L. Plant Physiol. 98:757-760.
- Amthor, J.S. 199X. A model of interactions among respiration, photosynthesis, and CO₂ in the mesophyll of C₃ leaves. Aust. J. Plant Physiol. (submitted)
- Denison, R.F. and R.S. Loomis. 1989. An integrative physiological model of alfalfa growth and development. Publ. No. 1926. Div. Agric. Nat. Res., Univ. Calif., Oakland, California. 73 p.
- Evans, J.R. and G.D. Farquhar. 1991. Modeling canopy photosynthesis from the biochemistry of the C₃ chloroplast. pp. 1-15. In K.J. Boote and R.S. Loomis (eds.). Modeling crop photosynthesis -- from biochemistry to canopy. CSSA Spec. Publ. 19, Am. Soc. Crop Sci., Madison, Wisconsin. 140 p.
- Farquhar, G.D. and S. von Caemmerer. 1982. Modeling of photosynthetic response to environmental conditions. p. 549-587. In A. Pirson and M.H. Zimmerman (eds.). Physiological plant ecology. II. Encyclopedia of Plant Physiology, New Series. Vol. 12B. Springer-Verlag, Berlin.
- Geng, S. and C.W. Cady. 1991. Climatic Variations and Change: Implications for Agriculture in the Pacific Rim. The Public Service Research and Dissemination Program, University of California, Davis, California, USA.
- Gollan, T., J.B. Passioura and R. Munns. 1986. Soil water status affects stomatal conductance of fully turgid wheat and sunflower leaves. Aust. J. Plant Physiol. 13:459-464.
- Penning de Vries, F.W.T., A.H.M. Brunsting, and H.H. Van Laar. 1974. Products, requirements and efficiency of biosynthesis: a quantitative approach. J. theor. Biol. 45:339-377.
- van Bavel, C.H.M. and J. Ahmed. 1976. Dynamic simulation of water depletion in the root zone. Ecol. Model. 2:189-212.
- Wallinga, J. 1991. Improving simulation of alfalfa growth and development with the ALFALFA model in water limiting and hot climate situations. Res. Rep., Theoretical Production Ecology, Agricultural University Wageningen, The Netherlands. 74 p.

APPENDIX A

Structure of WHEAT, a multilevel explanatory model of wheat crops responsive to variations in $[\text{CO}_2]$.

Initialize: plant and soil characteristics

Enter daily loop

Enter hourly loop

Read/calculate environmental conditions (temperature, dewpoint, wind, longwave radiation, direct and diffuse shortwave and PAR)

Enter dynamic, short time-step soil module (iterate several times h^{-1})

Calculate time step (s to min) from present conditions

Add rain or irrigation water

Redistribute water between layers (pseudo-Darcian flow)

Redistribute heat between layers

Move water between soil and roots (if present)

End soil loop

Calculate light interception by leaves

Enter leaf physiology module

Do until CO_2 flux is stable:

Get starting value of conductance

Calculate radiation, transpiration, and sensible heat exchange and get canopy temperature

Calculate canopy photosynthesis and maintenance respiration

Calculate new conductance accounting for $[\text{CO}_2]$ and temperature

End Do

Update plant carbohydrate (CH_2O) pools for photosynthesis and maintenance

End leaf physiology module

Calculate root maintenance respiration and update root

CH_2O pool

Calculate root growth in each layer (limit T, H_2O , CH_2O)

Do over tillers (mainstem to youngest):

Add/subtract CH_2O to/from stem pool

Grow apex (limit daylength, T, H_2O , CH_2O)

If apex is vegetative and big enough, initiate leaf unit (new leaf, axillary bud, stem internode)


```

    If apex is reproductive and big enough, initiate
    spikelet
      If conditions are right, initiate tiller
      If leaves are still growing:
        Calculate leaf growth
        Determine canopy structure (LAI, angle)
      End If
      If reproductive organs have been initiated at the apex:
        Calculate reproductive structure growth (limit T,
        H2O, CH2O)
        Update CH2O pools partitioned to reproduction
      End If
    End Do
    Calculate death of individual organs, mobilize remains
  End hourly loop
  Carbon balance check (inputs, outputs, growth, CH2O)
End daily loop
Output present status of plant (mass, developmental stage, etc.)
Output soil conditions by layer
Output process rates and total (photosynthesis, evapotranspiration,
etc.)

```

END

DATE
FILMED
3/19/93