NOTICE

**CERTAIN DATA
CONTAINED IN THIS
DOCUMENT MAY BE
DIFFICULT TO READ
IN MICROFICHE
PRODUCTS.**

Laboratory for Atomic, Molecular, and Radiation Physics (M/C 273)
Department of Physics
College of Liberal Arts and Sciences
Box 4348, Chicago, Illinois 60680-4348
(312) 996 4668 Fax (312) 996-8824

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

10 December 1992

PROGRESS REPORT:

Grant#: DE-FG02-91ER12108

Period of Performance: 15 December-30 September 1992

MOLECULAR DESIGN CONCEPT FOR X-RAY LASER RESEARCH

Principal Investigator:

Charles K. Rhodes

Co-Investigators:

Ting Shan Luk
Armon McPherson
Keith Boyer

Prepared for:

U.S. Department of Energy
ER-16, GTN
19901 Germantown Road
Germantown, MD 20874

Received by OSTI

DEC 17 1992

MASTER

EP

DO NOT WRITE IN THESE SPACES

TABLE OF CONTENTS

ABSTRACT	1
I. DISCUSSION OF RESEARCH	2
Appendix A: Comparison of Kinetic Energy Distributions and Ionic Fragment Yields of CO ₂ and N ₂ O Arising from Coulomb Explosions Induced by Multiph- oton Ionization and Fast Ion Impact	3

ABSTRACT

The goal of this program is the construction of an x-ray laser in the kilovolt regime. Recent experimental results indicate that a new technique for the generation of strong amplification of x-ray wavelengths is feasible. It involves the combination of (1) a new ultrahigh brightness subpicosecond laser technology and (2) a recently discovered unique mode of strong-field interaction, particularly applicable to molecules. A concept of molecular x-ray design emerges from these considerations which matches the mode of excitation to the structure of the molecular system. The molecular approach enables the combination of a very highly electronically excited conditions with an environment characteristic of dense cold matter, a general situation exceptionally conducive to x-ray amplification. Both high efficiency and wavelength tunability are intrinsic features of this method. Recent results discussed in this report are revealing important characteristics of the molecular strong-field coupling pertinent to our goal. A continued program of research is proposed to evaluate this method for the production of x-ray amplification in the kilovolt region.

I. DISCUSSION OF RESEARCH

The availability of high peak power ultraviolet (UV) lasers has revolutionized experimental research on the behavior of matter very strong electric fields ($E \geq e/a_0^2$). Moreover, the situation is such that presently available theories are not effective in describing many aspects of the new regime of strong-field coupling being explored. This is particularly true for many-electron systems, a class of materials that includes all molecules. Despite our imperfect understanding of the mechanisms involved in these high-field interactions, recent experimental results suggest that these multiphoton processes can result in a considerable transfer of energy to the irradiated system. For an intensity of $\sim 10^{17}$ W/cm² a total energy transfer exceeding 1 keV has been observed. Consequently, it is expected that total energy transfers considerably above 10 keV will be characteristic of the higher intensities ($> 10^{19}$ W/cm²) that will become available in the near future.

Molecules are emerging as a very interesting class of systems for study for at least two reasons. They are the multitude of species available and the experimental finding that the dynamics of the multiphoton coupling is sensitive to the molecular structure of the system. Accordingly, data concerning several molecules, including N₂, O₂, CO, N₂O, CO₂, and OCS, are under active study.

Recent results are giving detailed information on strong-field interactions relevant to our research goal. They involve the comparison of ion collisions and multiphoton processes that is discussed in Appendix A.

APPENDIX A

Comparison of Kinetic Energy Distributions and Ionic Fragment Yields of CO_2 and N_2O Arising from Coulomb Explosions Induced by Multiphoton Ionization and Fast Ion Impact

D. A. Tate*, T. S. Luk, K. Boyer, and C. K. Rhodes

Laboratory for Atomic, Molecular, and Radiation Physics
Department of Physics, University of Illinois at Chicago
P. O. Box 4348, Chicago, Illinois 60680

ABSTRACT

The comparison of kinetic energy distributions and ionic fragment yields produced from CO_2 and N_2O in fast ion collisions and multiphoton ionization reveals three comparative and distinguishing features of the interactions. They are (1) similarities in the O^{q+} (specifically, O^{3+}) energy distributions from CO_2 arising from correspondingly similar magnitudes of the peak electric fields occurring in the interactions, (2) a strongly dissimilar pattern of site-dependent ionization as observed in N_2O , and (3) a marked similarity in the production of anomalously energetic C^{2+} ions arising from high-lying CO^{2+} precursors having a substantial charge-transfer character.

* Colby College, Department of Physics, Mayflower Hill Drive, Waterville, ME 04901

I. INTRODUCTION

Corresponding aspects of strong-field multiquantum processes and collisional events, particularly ionic collisions with atoms [1-4] and molecules [5,6], have been previously examined. Several conclusions emerged from these discussions. Among them were (1) that certain gross features at the strong-field multiquantum interaction, for intensities in $10^{16} - 10^{21}$ W/cm² range, could be understood by analogy with collisional processes [4] and (2) that the multiquantum coupling can uniquely produce entirely new forms of excited matter [6]. Experimental evidence supporting both of these conclusions is presented and discussed below.

II. MULTIPHOTON EXPERIMENTAL STUDIES

The subpicosecond (~ 600 fs) KrF⁺ (248 nm) laser source [7] and time-of-flight ion apparatus [8-10] used in these studies have been fully described elsewhere. Generally, the material under study (CO₂ and N₂O) was introduced into the chamber at a density corresponding to $1-2 \times 10^{-8}$ ion, a value sufficiently low that distortion of the measured ionic kinetic energy distributions by the influence of space-charge could be neglected [8]. The region of peak intensity studied spanned the $\sim 3 \times 10^{16} - 2 \times 10^{17}$ W/cm² range and was achieved, on the low and high ends of this interval, respectively, by the use of a CaF₂ lens ($f/10$) and a parabolic mirror ($f/6$).

A. O³⁺ Kinetic Energy Distributions

We consider initially the measured kinetic energy spectrum of the O³⁺ ions produced from CO₂ by multiphoton ionization. This distribution, shown in Fig. 1(a), corresponds to an intensity of $\sim 3 \times 10^{16}$ W/cm², namely, for a peak field strength of approximately one atomic

unit. A distribution of fragment energies, having most of the yield below ~ 65 eV, but extending to a maximum value of ~ 100 eV, is observed.

The data illustrated in Fig. 1(a) can be compared, as shown in Fig. 1(b), to a corresponding spectrum produced [11] by Si^{8+} ion impact on CO_2 at ~ 3 MeV/u. In Fig. 1(a) and Fig. 1(b), the overall envelopes of the distributions carry the significant information and no importance is attributed to the narrow structures which are influenced by counting statistics.

An overlay of these two distributions reveals, in aggregate, that their shapes coincide remarkably well. Furthermore, analysis [11] of the observed collisional data for the O^{3+} distribution indicates the production of the $(\text{CO}_2)^{6+}$ molecular ion in the collisional interaction. This finding is generally consistent with other results on the production [12] of energetic oxygen ions from CO_2 by ion impact, specifically, by Ar^{12+} collisions at ~ 1.05 MeV/u. This latter study, which comparatively considers the collisional behavior of CO and CO_2 , has also been examined in the light of corresponding multiphoton-induced coulomb explosions of these two molecules. [13,14]

The multiphoton produced O^{3+} distribution exhibits an appreciable intensity dependence. [14] Figure 1(c) illustrates the kinetic energy spectrum of the O^{3+} fragment produced at a peak intensity of $\sim 2 \times 10^{17}$ W/cm². In comparison to the distribution shown in Fig. 1(a), a clear shift to higher energy is seen and a signal is observable to energies considerably above 100 eV. For this distribution, using the equilibrium interatomic distances characteristic of the CO_2 molecule, a simple analysis of the coulomb energies implies the multiphoton production of the $(\text{CO}_2)^{q+}$ ion with a corresponding charge given by $q \approx 8 \pm 1$, a range of values slightly higher than that indicated in the collisional production [12,14].

If we assume that the level of molecular ionization produced is governed mainly by the peak value [15,16] of the electric field generated in the interaction, the similarity of the observed ionic kinetic energy distributions illustrated above leads to the conclusion that the electric forces, giving rise to the data shown in Fig. 1, are comparable for both the collisional and multiphoton interactions. In this picture, similar electric force strengths give rise to similar rapidly produced levels of molecular ionization, and hence, to closely corresponding fragment energy distributions. From the data pictured in Fig. 1, the effective field involved in both the multiphoton and collisional interactions is estimated to be between one and two atomic units. A field of this value for the collisional case implies a characteristic impact parameter $b \cong (2.5 \pm 0.5) a_0$ and corresponding geometric cross section $\sigma \cong (5.2 \pm 1.7) \times 10^{-16} \text{ cm}^2$, magnitudes which are consistent with a reasonable range for such collisional interactions.

The production of a nascent system of the form CO_2^{q+} with a charge q representing several electrons, perhaps with a value of $q \sim 8$, points to an important basic consideration. Significantly, this multiply charged system is produced in the multiphoton case by an interaction causing a negligible transfer of momentum to the molecule. Core state (K-shell) excitation with x-rays followed by Auger decay can produce multiple ionization, but generally that mechanism generates for this material mainly doubly ionized material [17,18] with, perhaps, a small fraction of triply ionized ions. The maximum level of ionization that can be attained by the x-ray induced Auger mechanism for molecules composed of atoms having relatively low atomic numbers is modest. Energetic heavy ion collisions [19-21] can generate considerably higher levels of ionization, but this interaction generally imparts a very substantial recoil momentum to the constituents of the molecular targets [5]. However, the combination of a recoil-free

interaction and a high level of ionization can be quite simply achieved by the strong-field multiquantum mechanism. This method, therefore, enables the unique production of forms of kinetically cold highly excited matter [22] that are essentially impossible to produce by other available methods [5]. Basically, these results give evidence that entirely new forms of excited material can be generated by strong-field coupling.

B. Observed Pattern of Molecular Site-Dependent Ionization

The relative energy-integrated magnitudes of the observed ionic signals can give further information on the interactions. Ion collisional studies [12,23] (Ar^{12+} , 1.05 MeV/u) on CO_2 and N_2O have shown that the central atom is more easily ionized than the terminal atoms in these linear systems. Specifically, for CO_2 , this tendency reveals itself in the collisional interaction through observation of the fact that the C^{q+} ions ($q = 2 - 4$) are considerably more intense than the corresponding O^{q+} ions. The dynamics of these collisions is such that the ions arising from the ends of the CO_2 molecule appear to recruit electrons from the central carbon ion. A similar conclusion follows from collisional data [23] on N_2O in which the signal strengths associated with multiply ionized N^{q+} ions (e.g. $q = 2,3,4$) are more than one order of magnitude greater than those of the corresponding O^{q+} ions.

The ionic fragments produced in molecular coulomb explosions arising from multiphoton ionization give evidence for the opposite tendency. Studies [10] of isotopically substituted nitrous oxide ($^{14}\text{N}^{15}\text{N}^{16}\text{O}$) have shown, if anything, that the central atom (^{15}N) is less easily ionized than the terminal atoms. Examples of this behavior can be seen in the production of both singly (N^+) and triply ionized (N^{3+}) atoms as shown in Fig. 2. In this sense, in contrast

to the similarities in behavior illustrated in Fig. 1, the collisional and multiphoton processes clearly exhibit a substantial difference.

The differing time durations of the collisional and multiphoton events, the known anisotropy of the molecular multiphoton coupling [24-26], and the angular geometry of the force pattern characteristic of a collisional encounter may be factors influencing this difference in behavior. Specifically, with respect to consideration of the angular properties of the interactions, the electric field of linearly polarized laser radiation represents a force acting along a fixed line in space. Previous work [24-26] has shown that the multiphoton coupling strength varies appreciably with respect to the angle between this space-fixed axis and the molecular axis with maximum coupling occurring with the coincidence of these axes. Hence, in the multiphoton case, the interaction occurs mainly with molecules oriented along the polarization vector leading to a concentration of its action along the molecular axis. In this situation, the electric field can be effective in driving the electrons into states correlating with charge-asymmetric dissociation limits [9,26]. Charged-particle collisions, on account of the rapidly varying direction of the force with respect to the molecular axis that occurs during the interaction, are clearly less effective in concentrating the action of the force along the molecular axis. Indeed, most of the phase space associated with such collisions is associated with a large component of the field perpendicular to the molecular axis. Therefore, although the peak magnitudes of the forces may be comparable in both the multiphoton and collisional interactions, the angular variation of these forces in the fixed molecular frame differ greatly. This difference in the angular properties of the externally applied force field may also contribute to the reason why certain unusual molecular excited states [27] are generated with substantial probability in strong-field

experiments, but not in charged-particle collisions. This point is considered further below in the discussion concerning the details of the C^{2+} kinetic energy distribution.

C. C^{2+} Kinetic Energy Distribution

The kinetic energy distributions observed for the central carbon atom reveal additional features of the collisional and multiphoton interactions. Consider the case of C^{2+} , which exhibited an anomalously broad distribution arising from the $Si^{8+} + CO_2$ collisions. [11] Fragmentation of an ionized CO_2^{q+} system can produce carbon ions through two paths. These are

and

with $\alpha + \beta + \gamma = q$ and $x = \beta + \gamma$. As pointed out in the analysis [11] of Mathur et al., both processes, on account of the cancellation of recoil momenta affecting the $C^{\beta+}$ ions, tend to produce slow carbon ions. Naturally, this assumes that no appreciable rotation of the $(CO)^{x+}$ system occurs prior to dissociation in process (2b).

In consideration of the C^{2+} ion, the simple coulomb picture was unable to account for the magnitude of the maximum recoil energy observed [11] on the $Si^{8+} + CO_2$ collisions. The distribution of these collisional recoils, which is illustrated below in Fig. 3(a), shows ion production up to a maximum energy of 14 ± 2 eV. A consistent explanation [11] of the high energy component does emerge, however, if the collisional process produces inner-electron excitation in the carbon monoxide ionic fragment so that a subsequent Auger decay of the

collisionally excited system produces high-lying states of CO^{2+} . With this mechanism, the energetic C^{2+} ions can then be produced by the reactions

in which the $(\text{CO}^+)^*$ is an inner-shell excited species and the $(\text{CO}^{2+})^*$ precursor designates strongly repulsive high-lying states [28] of CO^{2+} produced by the Auger decay. Importantly, independent Auger studies [29] of CO have demonstrated the production of C^{2+} ions up to a maximum kinetic energy of 16 ± 8 eV. The positions of these Auger channels [29] are indicated on the data illustrated in Fig. 3(a) and they are seen to coincide well with the high energy component of the collisionally generated C^{2+} ions.

Significantly, the kinetic energy distribution of C^{2+} ions produced by multiphoton ionization also exhibits a high energy component, as shown in Fig. 3(b), the position of which corresponds well to the same CO Auger channels. The similarity of the distributions of the energetic C^{2+} ions, produced by both the collisional and multiphoton interactions, and their correspondence with the known Auger channels, supports the conclusion that the C^{2+} fragments originate in both cases from a common CO^{2+} precursor. This precursor would correspond to high-lying repulsive states [28] of CO^{2+} produced by Auger decay in the collisional case and direct multiphoton excitation in the high-intensity radiative interaction. In this picture, the C^{2+} ions would arise from channels of the form

namely, from CO^{2+} states involving an appreciable charge transfer character. Due to the very similar electronic structures of N_2 and CO, these states appear to be the direct analogues of the

corresponding N_2^{2+} charge-transfer states associated with new radiative transitions discovered in the N_2^{2+} ion under comparable conditions of irradiation [27]. The dynamics of the multiphoton process appears to favor the generation of such charge-asymmetric species, a point noted in earlier studies of ion production [9,26] for both N_2 and I_2 .

III. CONCLUSIONS

The comparison of kinetic energy distributions and ionic fragment yields produced from CO_2 and N_2O in fast ion collisions and multiphoton ionization reveals three comparative and distinguishing features of the interactions. They are (1) similarities in the O^{q+} (specifically, O^{3+}) energy distributions from CO_2 arising from correspondingly similar magnitudes of the peak electric fields occurring in the interactions, (2) a strongly dissimilar pattern of site-dependent ionization observed in N_2O that may be influenced to the different angular characteristics and time durations of the electric force fields generated in the two classes of interactions considered, and (3) a marked similarity in the production of energetic C^{2+} ions arising from high-lying CO^{2+} precursors having a substantial charge-transfer character. The latter feature points to a similarity with recent studies of N_2 involving both ion production and the generation of molecular ionic radiating states. Overall, these observations support the conclusion that new forms of excited molecular species can be generally produced by intense subpicosecond interactions.

REFERENCES

- [1] T. Åberg, A. Blomberg, J. Tulkki, and O. Goscinski, "Maximum Entropy Theory of Recoil Charge Distributions in Electron-Capture Collisions," *Phys. Rev. Lett.* 52, 1207 (1984).
- [2] X.-D. Mu, T. Åberg, A. Blomberg, and B. Crasemann, "Production of Multiply Charged Ions by Strong UV Laser Pulses: Theoretical Evidence for Stepwise Ionization," *Phys. Rev. Lett.* 56, 1909 (1986).
- [3] K. Boyer and C. K. Rhodes, "Atomic Inner-Shell Excitation Induced by Coherent Motion of Outer-Shell Electrons," *Phys. Rev. Lett.* 54, 1490 (1985).
- [4] K. Boyer, G. Gibson, H. Jara, T. S. Luk, T. A. McIntyre, A. McPherson, R. Rosman, J. C. Solem, and C. K. Rhodes, "Corresponding Aspects of Strong-Field Multiquantum Processes and Ion-Atom Collisions," *IEEE Trans. Plasma Sci.* 16, 541 (1988).
- [5] T. S. Luk, A. McPherson, G. N. Gibson, K. Boyer, and C. K. Rhodes, "Molecular X-Ray Laser Research," *Izvestia Academia Nauk, Seria Fizicheskaya*, T. 55, 768 (1991) (in Russian), English translation in Short Wavelengths Lasers and Their Applications, edited by V. V. Korobkin and M. Yu. Romanovsky (Nova Science Publishers, INC., Commack, NY, 1992) p. 1.
- [6] T. S. Luk, A. McPherson, K. Boyer, and C. K. Rhodes, "Mechanisms of Short Wavelength Generation," in Atoms in Intense Radiation Fields, edited by M. Gavrilu (Academic Press, Inc., New York, in press).
- [7] T. S. Luk, A. McPherson, G. Gibson, K. Boyer, and C. K. Rhodes, "Ultra-High Intensity KrF⁺ Laser System," *Opt. Lett.* 14, 1113 (1989).

- [8] T. S. Luk and C. K. Rhodes, "Multiphoton Dissociative Ionization of Molecular Deuterium," *Phys. Rev. A* 38, 6180 (1988).
- [9] K. Boyer, T. S. Luk, J. C. Solem, and C. K. Rhodes, "Kinetic Energy Distributions of Ionic Fragments Produced by Subpicosecond Multiphoton Ionization of N_2 ," *Phys. Rev. A* 39, 1186 (1989).
- [10] T. S. Luk, A. M. Moriarty, A. Awashti, K. Boyer, and C. K. Rhodes, "Isotopic Studies of Atomic Site Selectivity in Molecular Multiphoton Ionization of N_2O ," *Phys. Rev. A* 45, 6744 (1992).
- [11] D. Mathur, E. Krishnakumar, F. A. Rajgara, U. T. Raheja, and Vidhya Krishnamurthi, "Energy Distributions of Recoil Ions Produced in 100 MeV Collisions of Si^{8+} with CO_2 and CS_2 Molecules," *J. Phys. B* 25, 2997 (1992).
- [12] T. Matsuo, T. Tomura, M. Kase, T. Kambara, H. Kumagai, and H. Tawara, "Production of Multiply Charged Ions from CO and CO_2 Molecules in Energetic Heavy-Ion Impact," *Chem. Phys.* 121, 93 (1988).
- [13] T. S. Luk, A. McPherson, B. E. Bouma, K. Boyer, and C. K. Rhodes, "Studies of the Interaction of Molecules and Solids with Intense Subpicosecond 248 nm Radiation," in Multiphoton Processes, edited by G. Mainfray and P. Agostini (CEA, Saclay, 1991) p. 217.
- [14] T. S. Luk, D. A. Tate, A. McPherson, K. Boyer, and C. K. Rhodes, "Excitation of Molecules and Solids with Intense Subpicosecond Ultraviolet Radiation," in Coherence Phenomena in Atoms and Molecules in Laser Fields, edited by André D. Bandrauk and

Stephen C. Wallace, NATO ASI, Vol. B287 (Plenum Press, New York/London, 1992)
p. 175.

- [15] S. Augst, D. Strickland, D. D. Meyerhofer, S. L. Chin, and J. A. Eberly, "Tunneling Ionization of Noble Gases in a High-Intensity Laser Field," *Phys. Rev. Lett.* 63, 2212 (1989).
- [16] G. Gibson, T. S. Luk, and C. K. Rhodes, "Tunneling Ionization in the Multiphoton Regime," *Phys. Rev. A* 41, 5049 (1990).
- [17] J. Murakami, M. C. Nelson, S. L. Anderson, and D. M. Hanson, "Fragmentation of Nitrous Oxide by Monochromatic Soft X-Rays," *J. Chem. Phys.* 85, 5755 (1986).
- [18] R. Murphy and W. Eberhardt, "Site Specific Fragmentation in Molecules: Auger-Electron Ion Coincidence Studies on N_2O ," *J. Chem. Phys.* 89, 4054 (1988).
- [19] K.-H. Scharmer, "Highly Charged Recoil Ions," in Fundamental Processes in Energetic Atomic Collisions, edited by H. O. Lutz, J. S. Briggs, and H. Kleinpoppen, Series B, Vol. 103 (Plenum Press, New York, 1983) p. 637.
- [20] C. L. Cocks and R. E. Olsen, "Recoil Ions," *Phys. Reports* 205, 153 (1991).
- [21] A. S. Schlachter, W. Groh, A. Müller, H. F. Beyer, R. Mann, and R. E. Olson, "Production of Highly Charged Rare-Gas Recoil Ions by 1.4-MeV/amu U^{44+} ," *Phys. Rev. A* 26, 1373 (1982).
- [22] A. McPherson, T. S. Luk, G. Gibson, K. Boyer, and C. K. Rhodes, "Ion Production and Molecular Excitation Occurring in Multiphoton Processes," *Nucl. Instr. and Methods* B43, 468 (1989).

- [23] H. Tawana, T. Tonuma, K. Baba, T. Matsuo, M. Kase, T. Kambara, H. Kumagai, and I. Kohno, "Multiply Charged Ion Production Through Coulomb Explosion of Molecular Ions in Heavy Ion-Molecule Collisions," Nucl. Instr. Methods B 23, 203 (1987).
- [24] K. Codling, L. J. Frasinski, and P. A. Hatherly, "On the Field Ionization of Diatomic Molecules by Intense Laser Fields," J. Phys. B 22, L321 (1989).
- [25] J. J. Frasinski, K. Codling, P. Hatherly, J. Barr, I. N. Ross, and W. T. Toner, "Femtosecond Dynamics of Multielectron Dissociative Ionization by Use of a Picosecond Laser," Phys. Rev. Lett. 58, 2424 (1987).
- [26] D. T. Strickland, Y. Beaudoin, P. Dietrich, and P. B. Corkum, "Optical Studies of Inertially Confined Molecular Iodine Ions," Phys. Rev. Lett. 68, 2755 (1992).
- [27] G. Gibson, T. S. Luk, A. McPherson, K. Boyer, and C. K. Rhodes, "Observation of a New Inner-Orbital Molecular Transition at 55.8 nm in N_2^{2+} Produced by Multiphoton Coupling," Phys. Rev. A 40, 2378 (1989).
- [28] R. W. Weimore, R. S. LeRoy, and R. K. Boyd, "Theoretical Investigation of the Diatomic Dication CO^{2+} ," J. Phys. Chem. 88, 6318 (1987).
- [29] W. Eberhardt, "Core Electron Excitation and Decay in Molecules," Phys. Scr. T17, 28 (1987).

Figure Captions

- Figure 1: Comparison of the kinetic energy spectra for O^{3+} from CO_2 produced by multiphoton ionization and ion impact. (a) Distribution observed with subpicosecond (~ 600 fs) irradiation at 248 nm at a peak intensity of $\sim 3 \times 10^{16}$ W/cm². Most ions have an energy below ~ 65 eV. (b) Distribution produced by ion impact (Si^{8+}/CO_2) at a collision energy of ~ 3 MeV/u. Data from Ref. 11, presented with permission. (c) Distribution observed with subpicosecond irradiation, as in Fig. 1(a) above, with a peak intensity of 2×10^{17} W/cm². The peak in the kinetic energy distribution occurs at ~ 65 eV and an observable ion yield extends to energies > 100 eV.
- Figure 2: (a) Charge-state fragments observed by time-of-flight measurement from N_2O . Spectrum of N^+ from $^{14}N^{15}N^{16}O$ occurring in the range $13 \leq M/q \leq 17$. The abundance ratio of $^{14}N:^{15}N:^{16}O$ for these data is 2:08:1:1.25, respectively. A small Ar^{3+} signal is present for calibration. (b) Time-of-flight ion spectrum observed from $^{14}N^{15}N^{16}O$ in the region illustrating N^{3+} production. The abundance ratio of $^{14}N^{15}N^{16}O$ for these data is 1.7:1:1.05, respectively. See text for discussion. Data from Ref. 10.
- Figure 3: C^{2+} kinetic energies characteristic of collisional, multiphoton, and Auger processes. (a) C^{2+} ion kinetic energy distribution observed from $Si^{8+} + CO_2$ collisions at ~ 3 MeV/u. Data from Ref. 11 is shown with permission. The positions of Auger channels identified in Ref. 29 are also indicated. (b) C^{2+} kinetic energy distribution generated by multiphoton ionization of CO_2 showing the presence of a high energy component up to an energy of ~ 24 eV. The corresponding locations of Auger channels from Ref. 29 are also indicated.

END

**DATE
FILMED**

2 / 9 / 93