SANTA CRUZ INSTITUTE FOR PARTICLE PHYSICS (SCIPP)

TASK A

GRANT RENEWAL

1 November 1992 – 31 October 1993

	Page
1. INTRODUCTION	1
2. EXPERIMENTAL EFFORTS	3
2.1 ZEUS Project	3
2.2 SDC Effort	7
2.3 ALEPH Experiment	16
2.4 SLD Experiment	35
2.5 Experiment E791 at Fermilab	37
2.6 MARK III Analysis	40
3. BIBLIOGRAPHY	46
4. STATUS OF FACILITIES FOR COMPUTING	50
5. PERSONNEL AND BUDGET DETAILS	51
5.1 University Contributions	51
5.2 Budget and Personnel	63
5.3 Computing Equipment	56

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

MASTER

TASK A

Principal Investigators: P. Burchat, D. Dorfan, A. Litke, C. Heusch,
H. Sadrozinski, T. Schalk, A. Seiden

1. INTRODUCTION

This is a proposal for the second year of funding of a four-year grant. The grant supports the Task A experimental physics activities of the Santa Cruz Institute for Particle Physics (SCIPP) as well as work in theoretical physics. The Task A effort is divided into four subgroups: the ZEUS/SDC group, the ALEPH group, the SLD group and the E-791 group. The E-791 group is mostly funded by sources external to this grant (through a PYI grant and a special NSF grant for women in science.) The four subgroups are led by 4, 2, 1, and 1 faculty members, respectively. The groups share computer, laboratory, and secretarial resources as well as maintaining a close relation to foster the exchange of ideas, optimize planning, and make best use of individual unique talents.

Work for the coming year is a logical continuation of the efforts of the past year. Some special highlights of this past year have been:

- (1) The move onto beamline and start of ZEUS data taking.
- (2) The completion of the SDC technical proposal including a detailed long-term plan for construction.
- (3) Continuing publication of very detailed physics results from ALEPH concerning τ and b physics, and a precision measurement of electroweak and QCD parameters.
- (4) Completion of very successful data taking for E-791 at Fermilab, with nearly twice as many events recorded as initially proposed.
- (5) First measurement of beam polarization at the SLC.

These efforts have led to about 15 physics publications this past year centered mainly on topics related to QCD, couplings of flavors to the Z^0 , and heavy flavor decays. Taken as a whole, the results in jets from LEP, the ratio of hadronic to leptonic decays of the τ , the leptonic branching fraction of the J/ψ , and the charmonium mass spectrum provide a very consistent set of values of α_s at a variety of scales. In particular, they show the running of α_s by a factor of about three from m_τ to m_z . Results from LEP also provide evidence of the triple gluon vertex.

Similarly, the measurement of the $b\bar{b}$ fraction of Z^0 decays, from the MARK II as well as LEP, provide increasingly better measurements of the Z^0 coupling to

b quarks. Combined with earlier precision measurements of the Z^0 mass, width, and leptonic branching fractions, the Z^0 decays continue to provide a very precise verification of the Standard Model.

In addition to physics results, the efforts of the past year have led to about 15 contributions in a variety of areas in instrumentation. These include electronics design and development, radiation damage measurements for silicon detectors, development of simulation programs for silicon detector and electronics performance, and detailed measurements on drift chamber gases aimed at very low multiple scattering.

As part of the effort for physics at the SSC we have completed detailed simulations of tracking in the SDC aimed at Higgs reconstruction and top quark studies. These indicate excellent performance in the Higgs search even at six times the machine design luminosity. Such high luminosities may be needed in case of a very heavy Higgs or in the case of supersymmetry. Finally, we have participated in a theoretical study delineating rates and potential for Higgs discovery in the case of supersymmetry.

The work discussed above is presented in more detail in the next sections, along with plans for next year. This is followed by the bibliography of work completed this past year. The last sections deal with manpower and funding issues for the next year.

2. EXPERIMENTAL EFFORTS

2.1. ZEUS PROJECT

2.1.1. SCHEDULE AND STATUS OF HERA AND ZEUS

The ZEUS project at DESY in Hamburg made a big step forward on December 3, 1991 with the observation of e-p collisions inside HERA using the ZEUS luminosity monitors. This happened shortly before a scheduled four-month long shut-down for machine improvement and for the move on-line of the experiments. The experiments are now in their interaction regions and, starting April, are waiting for collisions. The initial plan for HERA operations looks as follows:

- April 1992 - Circulating beams.
- May 1992 - Start of spring data run (number
 - of bunches limited to about 10).
- July 20–August 17, 1992 - Machine development.
- August 18–November 30, 1992 - Full data run.
- November 30–December 31, 1992 - Fiscal shut-down.
- 1993 - Full data run.

The ZEUS detector has all the hardware in place to do the initial physics at low luminosity. There is a problem with procuring the drift chamber electronics (FADC's) which are slow coming out of RAL due to funding delays. This problem will be alleviated while the run is progressing.

2.1.2. PHYSICS GROUPS

In order to be prepared for data analysis this year, ZEUS has formed groups which concentrate on making sure that for physics topics in reach of low-luminosity running, both hardware and software are ready. The most important groups cover deep-inelastic scattering (DIS) and low- x physics. In the future, gluon structure functions and heavy quark physics will play an important role. We have participated in the low- x and DIS groups on the occasion of our DESY presence. The participation will intensify soon when we locate more personnel permanently at DESY.

2.1.3. SANTA CRUZ PERSONNEL AT DESY

Since February 1992, J. Ng, who joined UC Santa Cruz after getting his Ph.D. from Harvard with CDF, has been stationed permanently at DESY. He has taken over one of the "Collaboration Tasks": managing the online histogramming. He is

taking expert shifts and participates in physics groups as much as this duty allows him.

2.1.4. HARDWARE STATUS OF THE LEADING PROTON SPECTROMETER (LPS)

Coinciding with J. Ng's move to Hamburg, we delivered the first assembly of detector-chips-hybrid board, assembled and tested at UC Santa Cruz. All the components constitute prototypes of the final parts. We were able to read the data into a controller developed at CERN and via VME into the computer. Both the testing at UC Santa Cruz and the trouble-shooting at DESY were important steps before assembly and testing of the production line parts. We were able to assess the need for facilities at DESY for support of the LPS installation.

In the following we give the status of the different components:

- (1) *Silicon Strip Detectors.* The 36 detectors for the momentum analysing stations (S4,S5,S6) have been tested in Turin and are ready for mounting and bonding. The detectors for other stations (S1,S2,S3) await funding through INFN later. It is interesting to note that detectors were procured from MICRON, ENERTECHNIQUE and CANBERRA allowing comparison between different companies. The most difficult part of the detector manufacturing is the unusual shape with elliptical cut-outs for the outgoing proton beam which had to be cut either with a diamond saw or with a laser.
- (2) *Bipolar Amplifier-Comparator Chip.* The design of the amplifier-comparator chip was submitted to Tektronix for fabrication in their bipolar SHPi line in September 1991. After some delay in the layout phase the production went smoothly and we received the first die in April 1992. In the future, we will do our own layout in-house, using the Tektronix software, to control the schedule better and get better characterization of the whole circuit. We have now tested the first chips and anticipate good yield. We have analyzed the frequency behavior of the first amplifier stage and found good agreement between simulation and measurement (Fig. 1). The noise was determined injecting a known charge at the amplifier input and varying the threshold: the width of the counting rate curve is a measure of the noise of the whole circuit (see Fig. 2). We extract an equivalent noise charge of $ENC=860\text{ e}$ for 3.3pF input capacitance, an excellent result for a shaping time of 25ns . The signal is $25,000\text{ e}$. The radiation hardness of the SHPi process has been studied extensively by us (Ref. 1). Note that this amplifier-comparator chip fulfills the SSC requirements of speed, low-noise and radiation hardness. Only its power is a factor two too large. Testing of the whole lot will be done at UC Santa Cruz and we will get a good look

Fig. 1. Measured and simulated gain of the first stage of the Tektronix Amplifier-Comparator Chip.

Fig. 2. Threshold curve of the Tektronix Amplifier-Comparator Chip.

at the yield for this process which will ultimately determine the price for a large SSC-production run.

- (3) *Digital Pipeline Chip.* The digital time slice chip (DTSC) is a two-buffer SRAM being manufactured in radiation-hardened CMOS by UTMC. We have characterized the radiation hardness of the process extensively and have build and tested several prototypes. The DTSC is scheduled to be shipped to UC Santa Cruz on May 25, 1992 for initial testing. Full production testing will be done at the foundry.
- (4) *Hybrid Board.* The mixture of analog and digital signals on the hybrid board, combining detectors, analog chips and digital chips requires special care of noise suppression involving by-passing and layout. Another constraint is the need for good heat conduction to prevent heating up of the detectors. We have done the design and layout of a 12-layer board for fabrication as a Cu-Invar sandwich at CERN. The production of the hybrid board will be finished at the end of June 1992.
- (5) *Assembly and Testing.* The assembly and bonding of the chips to the hybrid boards have to be done in the U.S. because of the clause in our export license for the rad-hard CMOS chips. We will have this done in one of the Silicon Valley companies specializing in board assembly which we used before for similar work in the MARK II Silicon Vertex Detector Project. We have found that comparable work would cost much more if performed in Europe. We have agreed to pay for the work done in the U.S. to be able to supervise the work properly; in exchange our European collaborators will pay our ZEUS Common Fund charges at DESY.

The schedule of assembly is paced by the arrival of hybrids and the chip testing. Testing the electronics assembly after bonding will be done at UC Santa Cruz. We anticipate to deliver the first detector planes to DESY in the fall. H. Sadrozinski and B. Hubbard will spend part of the summer at DESY to set up test stations there and take ZEUS shifts. In the late fall, students will move to DESY permanently.

2.1.5. PREPARATION FOR ZEUS ANALYSIS AT UC SANTA CRUZ

Since last year, we have built up at UC Santa Cruz a computer system which allows us to support the ZEUS analysis locally. All ZEUS programs are running on our UNIX-based RS6000 IBM RISC stations. DESY is using RISC stations from DEC, Silicon Graphics, Apollo and IBM. We anticipate no real problem with staying interfaced with DESY. Interestingly enough, we have received requests to do Monte Carlo production on our computer system and send the output tapes

back to DESY. After initial tests we would like to help out as much as possible. We hope that T. Schalk will be able to spend some time at DESY in the fall to work on the efficient use of computer resources.

Our students have done, up to now, physics studies using stand alone Monte Carlo generators to explore the physics potential of HERA and get familiar with physics topics. Working first on the level of four vectors is efficient since the complete generation and reconstruction of a DIS event takes 5 minutes in the ZEUS software. J. Leslie has explored the determination of gluon structure functions with the production of $J/\psi \rightarrow \mu^+\mu^-$ events via the photon-gluon-fusion process. In order to check her programs, she has reproduced the existing data from the EMC collaboration. N. Cartiglia has worked on charm production and different charm tagging techniques. Two more students are expected to get their thesis with ZEUS physics, most likely on topics requiring more data than expected from the initial run in 1992.

2.2. SDC EFFORT

2.2.1. STATUS OF EXPERIMENT

The design of the SDC has moved ahead this year at a rapid pace. A concrete design has been completed as well as detailed subsystem cost estimates, construction schedules, and performance simulations. The SDC Technical Design Report has now been submitted to the SSC lab and will be defended by the collaboration over the next few months.

Santa Cruz has contributed to the evolution of the overall design through participation on the SDC technical board, executive board, and resource committee which is responsible for negotiating foreign financial contributions. Specific responsibilities have been the design and R&D for the silicon tracking system and also editing of the whole tracking section of the Technical Design Report. In addition, we have completed an extremely detailed Conceptual Design Report for the SDC silicon tracking system.

Santa Cruz has completed a very broad program of R&D for a large silicon tracker. Most of the work has been presented publicly in various forums and this body of work has strongly influenced other efforts such as the D0 silicon upgrade proposal and silicon tracking projects in Europe. A brief list of some of the projects of the past year is as follows:

- (1) Development of a program to simulate the behavior of silicon detectors including electrostatics, magnetic field effects, Landau fluctuations, electronic pulse shaping, and discriminator behavior.
- (2) Development of the first VLSI low-power, low-noise, bipolar readout chip for silicon detectors.
- (3) Development of the first data storage pipeline chip in radiation-hard CMOS.
- (4) Detailed radiation characterizations of the above chips.
- (5) Measurement of the behavior of various DC and AC coupled silicon detectors after irradiation with various types of particles (neutrons, proton and photons) and at varying temperatures.
- (6) Detector design aimed at minimizing capacitance and maximizing radiation hardness.
- (7) Development of a detailed track reconstruction program in the SDC GEANT framework.

In the coming year we are planning to continue with the program outlined above. Primary goals are:

- (1) Completion of the detector design.
- (2) Completion of the readout chip design. Fabrication and testing of the bipolar and CMOS chips.
- (3) Test of full detector modules, i.e., detectors with chips, cables, control signals, and shielding.
- (4) Begin work on a data acquisition system using fiber optics for the readout. This will require the definition of readout protocol and interfaces to the digital data storage chip as well as choices for the fiber drivers.
- (5) Definition and plan for all the cabling and fiber readout for the SDC silicon detector.
- (6) Development of an x-ray based alignment system which allows alignment and calibration of all detectors after complete assembly.
- (7) Improvement in our tracking program, emphasizing tracking in jets.
- (8) Responsibility for cost and schedule for the silicon tracker. This includes responsibility for meeting milestones, scheduling meetings, and dissemination of information to about 100 collaborators.

The hardware development mentioned above requires several types of beam tests which we are planning. Radiation testing will proceed at Los Alamos as in the past two years. We are also initiating a beam test program at KEK to begin next October and to continue periodically for several years. These should allow the final crucial measurements to verify our detailed design. To meet the SSC installation schedule we must begin final fabrication of major components (detectors, electronics, and mechanical parts) in 1994. This leaves only one year to complete the R&D design work in a large number of areas.

2.2.2. PATTERN RECOGNITION AND PHYSICS STUDIES

This past year a large effort has gone into performance simulation for SDC tracking. Figure 3 depicts the design of the silicon tracker. The simulation allows for a flexible description of the detector, in order to investigate a wide variety of designs. In the GEANT framework, detectors and inert material are described by a hierarchical volume structure. For the silicon system, the lowest level volumes are the detector wafers, which are positioned within layer volumes that simulate the support material. Each wafer has axial (or radial) strips on one side and stereo strips on the other, and an inactive region (default 500 μm) at the edges. The detector configurations involve wafers which are bonded together into larger "readout units" and which are "shingled" to avoid excess dead space.

The GEANT routines transport particles through the apparatus and generate decays and interactions with material, including conversions, bremsstrahlung, and delta rays. When a charged particle crosses a silicon wafer, the amount of dE/dx energy deposited in each strip is recorded and compared with a threshold. Clusters of strips with energies above the threshold are recorded in the raw data banks, with no analog information kept.

Fig. 3. Silicon detector array.

The raw data are reconstructed using an algorithm briefly summarized here. The coordinate reconstruction matches an axial strip with a stereo strip (if possible) within a single double-sided wafer and the results are expressed in the cylindrical coordinates r , ϕ and z . Segments are formed between two adjacent layers, connecting coordinates that are close in ϕ , and requiring the extrapolated z_0 to be within the range $|z_0| < 25$ cm. At this point the segment parameters ϕ_0 , ρ (curvature), z_0 and $s = \tan \lambda$ ($\lambda = \text{dip angle}$) are determined with respect to the distance of

closest approach to the origin. For the forward disks, the segment parameters are combinations of the above parameters which are more appropriate for the geometry.

The track reconstruction program used has several steps. The first is to take the segments from the silicon system and make clusters from segments that have close values of ϕ_0, ρ and z_0, s . The association is a χ^2 match based on the estimated error in the parameters and an estimate of multiple scattering. The barrel and forward segments are clustered separately as they have different parameter sets. Candidate tracks are clusters with at least two segments.

The second step is to take the list of hits in the candidate tracks and perform an iterative five parameter fit (the parameters ϕ_0, ρ, z_0, s and the impact parameter b_0). The fit includes multiple scattering in the silicon layers. Next, the algorithm tries to delete points that contribute the most to the χ^2 if the result from the first fit has a large χ^2 . A road technique is then used to pick up hits that were missed in the silicon as well as add information from other systems and also to join tracks which cross both the silicon barrel and forward regions. Hits near the track are picked up individually and then the track is refit, to guarantee the best possible estimate when extrapolating.

In the following sections, the silicon system and an outer system of straw tubes was used to model performance of the full SDC tracking system. The straw tube system consisted of three axial superlayers and two superlayers with 3° stereo angle.

Higgs Boson

The decay of a neutral Higgs boson (mass = 300 GeV) $H^0 \rightarrow Z^0 Z^0 \rightarrow e^+ e^- \mu^+ \mu^-$ has been studied with the combined silicon and straw system. The ISAJET generator was used to produce the Higgs event and PYTHIA minimum bias events were added as background. The design luminosity was simulated by a Poisson-distributed mean of 1.6 background events for each bunch crossing, with a total of seven crossings (-4 to $+2$ from the crossing with the Higgs) used to account for loopers and long drift-times in the straw system.

Figure 4 shows the reconstructed efficiency vs η for all tracks with transverse momenta above 1 GeV for 200 Higgs events at design luminosity. Tracks were required to have at least 4 axial silicon hits or 3 axial silicon hits and 1 straw segment. For tracks with $P_\perp > 20$ GeV, the efficiency (averaged over all η) is 99%. The number of fake tracks (tracks which have more than 2 hits from another Monte Carlo track) with $P_\perp > 5$ GeV is found to be 0.03 ± 0.01 per event.

To identify the gauge boson decay products, tracks are assumed to be identified as either an electron or a muon, using the Monte Carlo information in place of the calorimeter or muon system. Leptons are required to have $P_\perp > 10$ GeV and electrons to have $P_{\text{gen}}/P_{\text{fit}}$ within the range 0.7 to 1.4 to simulate the calorimeter

Fig. 4. Efficiency of single tracks with $P_{\perp} > 1$ GeV as a function of pseudorapidity.

energy and track momentum matching cut. In the case of electrons, the tracking momentum for these cuts is based on a refitted value using only the silicon information in order to minimize the effects of bremsstrahlung. Note that all leptons passing the above cuts were used in the analysis; therefore such leptons from b decays and conversions were considered.

To pair the leptons in order to make Z^0 candidates, the following algorithm was used. All pairs of same type, opposite sign leptons were made and the invariant mass was calculated. The invariant mass of the electrons was formed using the direction information from tracking but smearing the generated energies by the calorimeter resolution, since this typically provides the most accurate value of the electron energies. The muon invariant mass was calculated using the reconstructed momentum from the central tracking system only. The list of Z^0 candidates was made by ordering the pairs by their closeness to the Z^0 mass and then accepting pairs so that each lepton was used only once. The invariant mass of the electron pairs is shown in Fig. 5a, and the muon pairs in Fig. 5b. A cut requiring the invariant mass to be within 10 GeV of the Z^0 mass was then used to further restrict the list before the Higgs was reconstructed.

The four lepton invariant mass is plotted in Fig. 5c for events with two Z^0 candidates. Figures 5a–c are for the design luminosity of 10^{33} for which the efficiency for reconstructing the Higgs is 84%. This includes a small loss for events where the leptons did not all pass the 10 GeV $P_{\perp}$ cut. Table 1 summarizes the detector performance for this decay mode. In addition, the analysis was performed at luminosities of 3×10^{33} and 6×10^{33} . All results are shown in Table 1 and the invariant mass plots for the 6×10^{33} luminosity case are in Figs. 5d–f.

Fig. 5. a) Dilepton invariant mass distribution for electrons that have passed the P_{gen}/P_{fit} cut. b) Dilepton invariant mass distribution for muons. c) Reconstructed (solid) and generated (dashed) four-lepton invariant mass distribution for a 300 GeV Higgs. (d)-(f) similarly for $6 \times 10^{33} \text{ cm}^{-2} \text{ s}^{-1}$.

It is seen that the tracking detector should allow excellent performance for Higgs reconstruction well beyond the design luminosity. This is the single most important design goal for the detector. The track reconstruction was also performed without using information from silicon barrel layers 1 and 2 (at $r=9$ and 12 cm respectively) to check that the design is robust against some inefficiency. Note that occupancy in the silicon system is not a problem even at six times design luminosity and thus does not impair high luminosity performance.

Table 1. Summary of efficiencies and number of fake tracks for $H^0 \rightarrow e^+e^-\mu^+\mu^-$ events for various configurations.

Luminosity/ Configuration	Fakes per event with $P_{\perp} > 5\text{GeV}$	Track Efficiency $P_{\perp} > 20\text{GeV}$	Electron E/p Efficiency $0.7 < E/p < 1.4$	M_Z cut Efficiency		Higgs Recon- struction Efficiency
				e	μ	
1×10^{33} 8 layer STS + straws	0.03 ± 0.01	0.991	0.96 ± 0.01	0.99 ± 0.01	0.99 ± 0.01	0.84 ± 0.04
1×10^{33} 6 layer STS + straws	0.02 ± 0.01	0.963	0.93 ± 0.01	0.99 ± 0.01	0.97 ± 0.01	0.71 ± 0.04
3×10^{33} 8 layer STS + straws	0.04 ± 0.02	0.983	0.96 ± 0.01	1.00 ± 0.01	0.97 ± 0.02	0.83 ± 0.04
6×10^{33} 8 layer STS + straws	0.18 ± 0.03	0.972	0.93 ± 0.01	1.00 ± 0.01	0.93 ± 0.02	0.75 ± 0.04
6×10^{33} 6 layer STS + straws	0.06 ± 0.02	0.956	0.93 ± 0.01	1.00 ± 0.01	0.90 ± 0.02	0.65 ± 0.04

Capability for b -jet Tagging With Secondary Vertices

The capability to identify b -quark jets is expected to be an important tool for studying the $t\bar{t}$ system, by reducing the non- $t\bar{t}$ background substantially, and also reducing combinatoric background for measuring the mass of the t quark. The efficiency for tagging b jets using secondary vertices has been investigated with the GEANT simulation and full event reconstruction. A simple algorithm is used, which is to count for each jet the number of reconstructed tracks with significant impact parameter. Figure 6 shows a simulated $t\bar{t}$ event where both W bosons from the t decays subsequently decay leptonically.

ISAJET is used to generate b and “generic” jets in separate jobs, and tracks were reconstructed with the combined silicon and straw system. Tracks with measured $P_{\perp} > 0.8$ GeV within the cone $\Delta R < 0.5$ to the jet axis are associated to each jet. With the algorithm used, the reconstruction efficiency for these tracks ranges from 95% (jet $P_{\perp} = 50$ GeV) to 80% (jet $P_{\perp} = 500$ GeV). Tracks are further required to have $\chi^2/DOF < 5$ and $b_0 < 1$ cm (in order to discriminate against reconstruction errors and secondary interactions), and the impact parameter is required to have the correct sign to be consistent with originating from the decay of a particle moving in the jet direction.

Fig. 6. GEANT simulation of an Isajet $t\bar{t} \rightarrow e\mu\nu b\bar{b}$ event with $M_{\text{top}} = 150 \text{ GeV}$, showing the beam pipe, inner tracking gas enclosure, and first barrel silicon layer at 80% of actual size. Tracks reconstructed using data from the silicon and straw systems with $P_{\perp} > 2 \text{ GeV}$ and $|\eta| < 2.5$ are plotted. Charged leptons and b jets from the $t\bar{t}$ decay are indicated with their generated momenta; the muon is actually inside one of the b jets. Arrows indicate the true decay vertices of the b and c mesons within the b jets. The secondary vertices are visible by eye from the reconstructed tracks.

Figure 7 shows the tagging efficiency for b jets compared to that for a mix of jets from gluons and u , d , s and c quarks by requiring 3 or more tracks with impact parameter greater than 3σ . For b jets with $40 \text{ GeV} < p_{\perp} < 200 \text{ GeV}$, the efficiency is approximately 30%, and the rejection of the non- b jets is of order 100:1. Using these b tagging results from the full GEANT simulation, the efficiency for tagging one or both b jets from $t\bar{t}$ decay is estimated to be about 50%, assuming only secondary vertices are used.

The SDC design will allow b tagging over a very broad range in jet momentum. This is relevant not only to the $t\bar{t}$ studies, but could be of interest if new physics is discovered which populates preferentially the heavy quark sector.

Fig. 7. b tagging efficiency vs. jet $p_{\perp}$ for the combined system using the method described in the text.

2.3 ALEPH Experiment

This year marks a transition for our group, led by Alan Litke. Since 1984, our efforts had been devoted to pioneering the development and implementation of silicon detectors with VLSI readout, particularly in connection with the MARK II Silicon Strip Vertex Detector. This work has been completed, as described in Section 2.3.1. Now our emphasis has shifted to exploiting the marvelous physics opportunities opened up by the silicon vertex detector in the ALEPH experiment at LEP-I and, eventually, LEP-II. For our work on the ALEPH experiment, we seek continued funding during the period 1 November 1992 through 31 October 1993.

During the past year the ALEPH effort within SCIPP has been greatly strengthened by the addition of a new assistant professor, Robert Johnson, and a second postdoc, Gary Taylor. Both Johnson and Taylor have brought to this group considerable experience with the ALEPH experiment. Johnson worked on ALEPH for more than 5 years as a postdoc in the University of Wisconsin group, and made major contributions to the ALEPH hardware (Time Projection Chamber), software, and analysis efforts. Taylor did his thesis work on ALEPH, during which time he also contributed to the construction and operation of the inner drift chamber, and afterwards worked for a year as a postdoc with the Imperial College group. In addition, our first postdoc, James Wear, did his thesis work on ALEPH in the Wisconsin group. Finally, two graduate students have been recruited at UCSC, who are beginning to contribute both to our hardware and analysis activities. Thus we believe that we now have a strong group which can be expected to make significant contributions to the ALEPH experiment by capitalizing both on its many years of experience within the ALEPH collaboration and the extensive experience and expertise of Litke in the field of silicon vertex detectors at e^+e^- colliders.

As explained in Section 2.3.2, the Santa Cruz group has during the past year made important contributions to the commissioning and operation of the ALEPH silicon vertex detector (VDET), which performed well during the 1991 LEP data run. ALEPH now is beginning its 1992 run, after an overhaul of the VDET during the past months. Analysis of the 1991 VDET data has begun, as reviewed in Section 2.3.5, and we are becoming heavily involved, both at Santa Cruz and at CERN, in developing methods to extract physics results from the vertexing capability of our new detector. We are looking forward to having at our disposal several million events at the Z peak from LEP-I and plan also to participate in ALEPH data taking and analysis at LEP-II, scheduled to begin at an energy above the W -pair threshold in 1994.

We begin with a summary of the status of our work on the MARK II experiment in the next section. The remainder of this report is devoted to our work on ALEPH, for which we describe the ALEPH vertex detector and our contributions to it, briefly summarize the recent ALEPH physics results, and describe the first steps toward physics with the VDET.

2.3.1 Status of the MARK II Experiment at the SLC

During the past year, the MARK II experiment at the SLC was essentially completed. Analysis of the data from the last run relied on the precision tracking information from the Silicon Strip Vertex Detector (SSVD)—the main contribution of our group to the MARK II detector. This analysis gave a measurement of $R_{b\bar{b}} \equiv \Gamma(Z \rightarrow b\bar{b})/\Gamma(Z \rightarrow \text{hadrons})$ using, for the first time, precision vertex information. We find $R_{b\bar{b}} = 0.251 \pm 0.049 \pm 0.030$, in good agreement with other measurements and the Standard-Model prediction. This result has been published in Physical Review Letters.¹

The successful operation of the SSVD marked the first time a precision silicon vertex or tracking system had been implemented in a colliding beam environment. Our group played a major role in the SSVD, including the project leadership. We were responsible for the development of the silicon detectors, as well as the design, assembly, and testing of the silicon detector modules. We developed an x -ray alignment system and were responsible for the data acquisition electronics. A long paper on the design, assembly, operation, and final performance of the SSVD has recently been published in Nuclear Instruments and Methods.² We hope it will prove useful to the many groups that are now beginning work with silicon detectors.

The SSVD was a great technical success and an important pioneering effort that demonstrated the viability and power of this new technology. Unfortunately, due to the very limited data sample provided by the SLC (only 294 Z^0 events recorded), it produced only limited physics results. However, it gave us a taste for the physics potential of a precision vertex detector looking at Z^0 decays. This led to our decision to join the silicon vertex detector effort in the ALEPH experiment at LEP.

¹R. Jacobsen *et al.*, *Measurement of the $b\bar{b}$ Fraction in Hadronic Z^0 Decays with Precision Vertex Detectors*, Phys. Rev. Lett. **67** (1991) 3347.

²C. Adolphsen *et al.*, *The MARK-II Silicon Strip Vertex Detector*, Nucl. Instrum. Methods A **313** (1992) 63.

Figure 8: The geometry of the ALEPH silicon vertex detector.

2.3.2 The ALEPH Silicon Vertex Detector

Description: The ALEPH silicon vertex detector consists of two cylindrical layers of double-sided detectors. The geometry is shown in Fig. 8. The first layer is at a radius of 6.3 cm and has 9 faces; the second layer has a radius of 11.0 cm with 15 faces. The angular coverage extends over the range $|\cos \theta| < 0.85$ for the inner layer and $|\cos \theta| < 0.67$ for the outer layer.

All faces are identical and are made from two detector modules which are placed end-to-end. The layout of a module is shown in Fig. 9. Each module consists of two $5 \times 5 \text{ cm}^2$ silicon detectors. These double-sided detectors have ≈ 2000 $r\phi$ strips with a pitch of $25 \mu\text{m}$ on the p -side and ≈ 1000 z strips with a pitch of $50 \mu\text{m}$ on the n -side. The two detectors are wire-bonded together on the $r\phi$ side.

The readout is done with CAMEX-64 chips (64 channels/chip) at a pitch of $100 \mu\text{m}$. Each read-out strip is AC-coupled to its corresponding amplifier channel via a 210 pF capacitor. These capacitors are specially fabricated on a quartz substrate as an “AC coupling chip,” with 64 channels per chip. A total of 72,000 channels are read out from the VDET.

Figure 9: The layout of the ALEPH double-sided silicon detector module.

Performance: After alignment, the spatial resolution for tracks at normal incidence is measured to be $12\mu\text{m}$ in $r\phi$ and $12\mu\text{m}$ in z . The impact parameter resolutions for $45\text{ GeV}/c$ tracks from $Z^0 \rightarrow \mu^+\mu^-$ events are $\sigma_{b_{r\phi}} = 25\mu\text{m}$ and $\sigma_{b_z} = 29\mu\text{m}$. These are to be compared to $\sigma_{b_{r\phi}} = 123\mu\text{m}$ and $\sigma_{b_z} = 960\mu\text{m}$ when VDET track information is not used. Also, the VDET improves the momentum resolution at $45\text{ GeV}/c$ by 33%. The effect of the improved tracking and vertexing on some physics analyses are described in Section 2.3.5.

Santa Cruz Role: Since joining the ALEPH collaboration, the Santa Cruz group has played an important part in the activities concerning the vertex detector. During the winter, 1990-91 LEP shutdown, we helped set up the system for performing optical alignment measurements of the VDET and assisted in the detector's assembly, installation, and testing. During last year's data run, we served part time as VDET run coordinator. During and after the run, we were part of a small group of troubleshooters that analyzed and investigated the non-obvious problems that arose during the commissioning of this new device, which is based on the largely untested technology of double-sided silicon detectors. These problems and our studies are described in the next section.

Due to concerns about possible radiation damage to the VDET, and actual damage from sudden beam losses (see below), the Santa Cruz group took responsibility for the design, fabrication, installation, and operation of a silicon radiation monitoring system. This system is the specific hardware responsibility of the Santa Cruz group in ALEPH. It is described in Section 2.3.3.

Problems and Solutions: During the 1991 data run, two quite unexpected problems arose in the operation of the VDET which had important implications for the future running of the detector. More generally, these problems are of concern to experiments (such as the SDC detector at the SSC) which employ, or plan to employ, silicon trackers.

1. **Sudden beam loss damage:** In four separate incidents during data taking, sudden loss of the LEP beams resulted in damage to a total of about 7% of the VDET channels. This was of great concern to us, since these beam losses were sure to continue in the future and could lead to an on-going deterioration of the detector.

Upon investigation, we arrived at the following scenario for the damage:

- A burst of ionizing radiation generates current in the silicon detector.
- This current produces a voltage spike across the AC coupling capacitor which breaks down the capacitor and produces a short.
- This short leads to saturation of the affected channel, and, through coupling via interstrip capacitance, to saturation of ≈ 10 neighboring channels.

To test this scenario, our group studied the breakdown voltage distribution of the coupling capacitors when subjected to a short (1 msec) pulse. Our results were consistent with the above hypothesis.

Our surprising conclusion is that the damage resulted from a total radiation dose which could be as low as 0.6 rad! However, this dose must occur on a time scale which is short compared to the charging time of the detector bias circuit (≈ 1 msec in our case).

To reduce the effects of the damage, we added an extra bias voltage which took the neighboring strips (but not the strip with the shorted capacitor) out of saturation. This reduced the number of saturated channels by an order of magnitude, from 7% to 0.7%.

To attempt to reduce or eliminate this problem in the future, the Santa Cruz group has installed a radiation monitor (see Section 2.3.3) which will be able to detect beam losses near the VDET on a short time scale ($< 100 \mu\text{sec}$). If radiation approaching a dangerous level is detected, it will be able to supply a fast signal to LEP which can be used to dump the beams in a safe manner.

2. **Dying line driver chips:** During the data run we noticed that the line driver chips (used to send the module signals from 256 channels to the data acquisition system) were dying at the rate of 1.2 chips per month. Again, we were concerned by the possibility of an on-going deterioration of the detector. To investigate this mystery, we decided to remove VDET from ALEPH during the winter, 1991-92 shutdown. We found a problem which can explain the line-driver deaths:

- On 9 faces, a special two-component epoxy had been improperly applied to protect the wire bonds connecting the module hybrid circuit to a thin cable. This led to incomplete mixing/curing of the epoxy and to microscopic craters still filled with fluid more than 9 months after application!

- The flow of cooling air sprayed this fluid as a fine mist onto the line driver chips.
- The epoxy-component residue on the chips resulted in their slow death (this part is not understood in detail).

During the shutdown we repaired or replaced all of the damaged modules.

Present Status: The vertex detector has been reinstalled in ALEPH, with all modules working properly. We have now begun to accumulate data again with LEP colliding beams.

2.3.3 The ALEPH Silicon Radiation Monitoring System

In order to measure the radiation levels near the VDET, the Santa Cruz group has designed, fabricated, and installed a radiation monitoring system in ALEPH. It is based on a set of silicon PIN diodes, each 1 cm^2 in area. These are located on a plastic ring, centered on the beamline, 8 cm in radius, and displaced 36 cm from the interaction point.

Two complementary systems have been implemented:

- The *SLOW* system is used to monitor the radiation level on a time scale of seconds. It provides a measurement of the integrated dose to the VDET and gives ALEPH the ability to evaluate background conditions close to the interaction point. It provides a warning signal when excessive radiation levels are detected, so that ALEPH can notify the LEP operators to take corrective action before significant damage is done to VDET. This system is operational all the time; even under beam conditions that make it necessary to turn off all the other sub-detectors. It is extremely reliable and easily calibrated.
- The *FAST* system will be used to study sudden beam losses of the type that we believe caused damage to the AC-coupling capacitors of the VDET. Its integration time of $100\text{ }\mu\text{sec}$ is comparable to the LEP beam orbit time. We have implemented a self-contained data acquisition system (based on a Macintosh) that is able to record a “snapshot” of the time development of signals from the PIN diode channels in each radiation incident. This system can also be used to supply LEP with a signal to implement a “safe” beam dump when the radiation approaches a level potentially dangerous to VDET.

2.3.4 ALEPH Physics Results

During the past year, ALEPH has continued to update those analyses which can benefit from the increase in statistical significance afforded by the factor of 3 increase in events with respect to the data sample accumulated during 1989 and 1990. In addition, work has begun on analyses which rely on the new silicon vertex detector. Most of the results which have been achieved to date were reported at the March 1992 Moriond conferences.

In the area of searches, most analyses were even before this year limited by the beam energy from further improvement. The notable exceptions are the Higgs searches and the searches for rare decays, such as $Z \rightarrow \gamma\gamma\gamma$, for which we now limit the branching ratio to less than $1 \cdot 10^{-5}$. With the addition of the 1991 data, our limit on the mass of the Standard-Model Higgs has improved from $48 \text{ GeV}/c^2$ to $54 \text{ GeV}/c^2$. It is clear that dramatic improvements in the searches await the start of LEP-II.

Electroweak physics has continued to benefit from the increase in data, especially in the area of asymmetries. The UCSC involvement has been in the measurement of the forward backward asymmetry in $Z \rightarrow b\bar{b}$, where Johnson was one of the early leaders in the continuing work to extract this quantity from a fit to the inclusive lepton spectrum. (Also, both Johnson and Taylor have made major contributions to the calibration and understanding of the lepton identification, for both electrons and muons.) With the addition of the 1991 data, the uncertainty in the measured asymmetry has decreased from 0.031 to 0.019. Similar improvements have been made in the leptonic forward-backward asymmetries and the tau polarization asymmetry. Measurements of the Z lineshape have also been updated. A major improvement in this area has come from improved knowledge of the beam energy, by use of measurements based on resonant depolarization. The uncertainty in the ALEPH Z mass is now $\pm 11 \text{ MeV}/c^2$.

Activity in the area of heavy flavor physics has increased in the past year, due both to the larger data sample and the advent of the new silicon vertex detector. ALEPH has recently published a signal for Λ_b production, using correlations between Λ baryons and leptons in the same jet.³ From the impact parameters of the leptons in this signal, a first measurement of the Λ_b lifetime has been made: $\tau_{\Lambda_b} = 1.12^{+0.34}_{-0.32} \text{ ps}$, as reported at Moriond. Another more direct signal, using correlations between leptons and Λ_c^+ baryons (detected in the $pK^-\pi^+$ mode), is nearing publication, along with a measurement of B_s^0 production, from correlations between leptons and D_s^- mesons. As an

³D. Decamp *et al.*, *Evidence for B Baryons in Z Decays*, CERN-PPE-91-229, Submitted to Phys. Lett. B.

Figure 10: The two signals for D_s^- used in the ALEPH search for B_s production: (a) D_s^- signal in the $\phi\pi^-$ decay mode. (b) D_s^- signal in the $K^{*0}K^-$ decay mode. At this point of the analysis, no requirement has yet been made that there be a lepton in the same jet.

example of the performance of our spectrometer, Fig. 10 shows the two signals for D_s^+ used in this analysis. Taylor and Johnson have contributed to the latter work, especially in the area of identification of the kaons from the ϕ decay, where special techniques are necessary to improve the efficiency of the dE/dx measurement, due to the small opening angle of the two kaons.

2.3.5 Physics with VDET—The First Steps

Physics results which rely on the new vertex detector still are limited, due to the fact that correctly calibrated and reconstructed data have only been available for about three months. During the summer and autumn of 1991, a large effort was made to understand the calibration and alignment of the

Figure 11: Measured decay lengths for 3-prong decays of the tau lepton, in centimeters. At least two of the three tracks in each event are required to have hits in the silicon vertex detector.

three tracking detectors such that the best possible use could be made of the vertex detector precision. Systematic problems in the time projection chamber (TPC) were of particular importance, since the direction information from the outer tracking chambers is responsible for reducing the error on a track impact parameter by about a factor of two with respect to what the vertex detector could do alone. Johnson and Taylor contributed to the effort in solving these problems. In particular, Taylor made a study of the t_0 and drift-velocity calibration of the TPC which resulted in big improvements in the systematic error in the measurement of the polar-angle of tracks (he will continue this year to contribute to this aspect of the tracking calibration). As a result, in the 1991 data set, which was completely reprocessed offline during December 1991, the impact parameter resolution of high-momentum tracks is $25\,\mu\text{m}$ in $r\phi$ and $29\,\mu\text{m}$ in z .

The vertex detector data have already been used to measure the tau lifetime both by impact-parameter methods in the 1-prong sample and by decay-length measurements in the 3-prong sample. Figure 11 shows a histogram of the measured decay lengths in the 3-prong sample. Since the resolution on the decay length ($\approx 800\,\mu\text{m}$) is considerably less than $\beta\gamma c\tau \approx 2300\,\mu\text{m}$, the

Figure 12: Measured impact parameters of high- $p_{\perp}$ leptons in hadronic Z decays. Each lepton candidate is required to have at least one hit in the silicon vertex detector.

exponential decay distribution is very obvious. Santa Cruz has contributed to the 3-prong analysis through work on developing new pattern recognition techniques for the closely spaced tracks that one finds in such decays.

The ALEPH measurement of the average b -hadron lifetime from impact parameters of high- $p_{\perp}$ leptons has also been updated using vertex-detector data. Figure 12 shows a histogram of the measured impact parameters. Not only the resolution, but also the background rejection, has been improved with respect to the earlier ALEPH publication.⁴ Again, the exponential decay is evident in the measured distribution. With the vertex detector, we are clearly in a situation where the statistical error on the result is completely

⁴D. Decamp *et al.*, *Measurement of the B Hadron Lifetime*, Phys. Lett. B **257** (1991) 492.

dominated by the number of events measured, as opposed to the experimental resolution. That has the added advantage in such analyses of reducing systematic errors due to uncertainty in the resolution function, since we are no longer in a situation of trying to measure a slightly displaced mean in an almost gaussian distribution.

2.3.6 Future Plans in ALEPH

The ALEPH Experiment: Some increase in LEP luminosity is expected this year as the machine goes from four-bunch to eight-bunch operation. Thus one hopes for at least 700K Z^0 events per experiment on tape from the 1992 running and at least that number in the 1993 run. Combined with the present data sample of almost 500K events, ALEPH may have a total of almost two million Z^0 hadronic decays to analyze by the end of the contract period. As one example, this would correspond to more than 800K B 's produced, including more than 100K B_s mesons.

There will be two improvements to ALEPH in the next 1-1/2 years. First, a new small-angle luminosity monitor (SICAL) will be installed this year. Second, there will be a general improvement program for the data acquisition system (to improve reliability and maintainability) in 1993-94. The VDET DAQ is part of this program, as described in the next section.

The ALEPH physics program on precision electroweak measurements, heavy flavor physics, QCD studies, and searches for new particles, will continue. The anticipated future hardware and physics analysis activities of the Santa Cruz group are described in the next three sections.

The Silicon Vertex Detector: We have decided *not* to upgrade the existing detector, but to concentrate on maintaining it in good working order, and to learn how best to use it to extract interesting physics. The VDET is a new instrument, providing precision spatial measurements in both coordinates, and it will take some time to learn how to fully exploit its potential for physics.

One activity connected with the maintenance and reliability of the VDET is the improvement of the data acquisition system. The present readout system depends on Fastbus-based event builders for which spares and expertise are becoming increasingly hard to find. As a general improvement program for the ALEPH data acquisition system, all readout will be standardized with commercially available, VME-based 68040 processor modules. Santa Cruz will help implement the conversion to the VME-based readout system for the VDET.

The Silicon Radiation Monitoring System: The fast silicon radiation monitoring system, as described in section 2.2.2, will be used during the coming run to monitor beam losses in LEP and to evaluate the potential of each beam loss to damage the VDET. In collaboration with the LEP machine development group, we expect to conduct studies of how different types of beam losses can affect the VDET, and to evaluate the effectiveness of controlled “safe” beam dumps.

Of particular concern is the implementation this year of electrostatic plate separators which are necessary for the conversion from four-bunch to eight-bunch operation. These electrostatic plates are prone to sparking, which can lead to sudden beam losses and therefore to damage to the AC-coupling capacitors of the VDET.

We hope we can detect beam losses early with the radiation monitoring system and then implement a safe beam dump (all of this needs to be thoroughly tested!). Alternatively, we can try to turn off safely and rapidly the VDET detector voltage. In either case, we need to design and build an extremely reliable system for producing a “dump” signal, based on the information supplied by the radiation monitor.

The Santa Cruz Data Analysis Effort: In the coming year we will concentrate the majority of our effort on data analyses which will lead to published physics results. Our hardware obligations at CERN should soon begin to decrease, thus allowing Litke and Wear to devote more time to offline analysis. Taylor will remain stationed at CERN and will devote about 75% of his time to analysis. At Santa Cruz, Johnson has completed installation of the ALEPH and CERN software and the ALEPH data on the SCIPP cluster of IBM RS6000 workstations and has already begun some analysis work there along with our two graduate students, McNeil and Rennels.

One of our main efforts is directed toward the eventual extraction of B_s and Λ_b mass peaks from our data. It is clear that to attain sufficient background rejection for such an analysis, we will have to rely heavily on the vertex detector, in contrast with our current results on B_s and Λ_b , which rely on a high- $p_\perp$ lepton to select semileptonic b decays. Toward this end, we have been working both to increase our efficiency in reconstruction of charm mesons and to study tagging of vertices by the vertex detector. For reconstruction of the reaction $D^{*+} \rightarrow D^0 \pi^+$, we have added to our existing set of D^0 decay modes, $D^0 \rightarrow K^- \pi^+$, $D^0 \rightarrow K^- \rho^+$, and $D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-$ (shown in Fig. 13), the mode $D^0 \rightarrow K^0 \pi^+ \pi^-$ (Fig. 14), which had not yet been observed within ALEPH. It is not likely that there are any other useful modes for our purpose; however, these four modes alone comprise almost 25%

Figure 13: (a) The D^* , D^0 mass difference for $D^0 \rightarrow K^- \pi^+$ candidates from $D^{*+} \rightarrow D^0 \pi^+$, where the D^0 mass has been required to be between $1.835 \text{ GeV}/c^2$ and $1.895 \text{ GeV}/c^2$. (b) The D^* , D^0 mass difference for the mode $D^0 \rightarrow K^- \rho^+$, where the K^- , ρ^+ mass is required to be between $1.815 \text{ GeV}/c^2$ and $1.915 \text{ GeV}/c^2$. (c) The D^* , D^0 mass difference for the mode $D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-$, where the D^0 candidate mass is required to be between $1.825 \text{ GeV}/c^2$ and $1.905 \text{ GeV}/c^2$. The data are taken from about 170 000 hadronic Z decays. In (a) and (b), the D^* candidate is required to have at least 25% of the beam energy, while for (c) the cut is at 35%. The overplotted smooth curves represent a fit of a single gaussian plus a second-order polynomial. The fitted gaussian parameters are printed on the plots.

Figure 14: The D^* , D^0 mass difference for the mode $D^0 \rightarrow \bar{K}^0 \pi^+ \pi^-$. The solid curve represents a fit to the data using the D^0 sidebands to define the background shape and a gaussian curve to describe the signal.

of all D^0 decays. For the D_s^+ sector life is more difficult, due to the small splitting between the D_s^* and the D_s ($\Delta m < m_\pi$). We are trying to find, however, other B_s decay channels to exploit besides those with a D_s^+ meson. For example, we are currently studying the reaction $D_{s1}(2536)^+ \rightarrow D^{*+} K^0$, which is something that nobody yet in ALEPH has looked for. We will also search for decays with a baryon-anti-baryon pair in the final state.

Tagging of vertices has been studied using a number of charm decays. For example, Figure 15.a shows the signals for $D^0 \rightarrow K^- \pi^+$ and $D^+ \rightarrow K^- \pi^+ \pi^+$, using *no* D^* or lepton tag, but requiring that the two or three tracks in the decay form a displaced vertex. For comparison, we also show the results for the case where no cuts on decay length or vertex quality are made—use of the vertex detector clearly provides a large improvement in the signal-to-noise ratio. This indicates the true power of a vertex detector which is read out in both dimensions: even for 2-track decays (which always form a perfect vertex if one looks only in the x - y plane) the combinatorial background can be greatly reduced by requiring that the two candidate tracks form a good 3-dimensional vertex which is displaced from the origin in the direction of flight.

Both Johnson and Taylor have had a deep involvement during the past years in the ALEPH analyses of inclusive lepton production in hadronic Z decays. This work has led to publications of branching ratios,⁵ B - $\bar{B}$ mixing,⁶ and forward-backward asymmetry in $Z \rightarrow b\bar{b}$.⁷ With the larger data sample now being accumulated, continuation of the measurement of forward-backward asymmetry remains a priority within ALEPH due to its importance as a precision measurement of the weak mixing angle. This measurement should continue to improve with increased statistics. However, greater attention must be paid to systematic effects. We plan to begin using the vertex detector in this analysis to make crosschecks of our understanding of the various sources of leptons in the data, and in particular to differentiate between leptons from heavy-flavor decays versus those from light-quark events. The vertex detector may also provide some information on the production of higher resonances than the $D^*(2010)^+$ in semileptonic B decays, which is important to understand due to the effect such decays would have on the transverse-momentum spectrum of the leptons.

⁵D. Decamp, *et al.*, *Heavy Flavour Production in Z Decays*, Phys. Lett. B **244** (1990) 551.

⁶D. Decamp, *et al.*, *Measurement of B - $\bar{B}$ Mixing at the Z*, Phys. Lett. B **258** (1991) 236.

⁷D. Decamp, *et al.*, *Measurement of the Forward-Backward Asymmetry in $Z \rightarrow b\bar{b}$ and $Z \rightarrow c\bar{c}$* , Phys. Lett. B **263** (1991) 325.

Figure 15: Examples of using the vertex detector to enhance signals for the charm decays $D^0 \rightarrow K^-\pi^+$ (left) and $D^+ \rightarrow K^-\pi^+\pi^+$ (right). The D^0 candidates were required to have at least 20% of the beam energy and to have a vertex at least 2σ from the beam spot and a vertex $\chi^2/\text{dof} < 2$. The D^+ candidates were required to have at least 30% of the beam energy and to have a vertex at least 3σ from the beam spot and a vertex $\chi^2/\text{dof} < 2$. The dashed histograms show the corresponding signals in the case where the cuts on decay length and vertex quality are not used.

Figure 16: Signal for the decay $\Xi^- \rightarrow \lambda\pi^-$, obtained from 190 000 hadronic Z decays. Both the Ξ and λ vertices are required to be displaced from the origin by at least 3 cm, and the λ vertex must be at least 2 cm further out than the Ξ vertex.

A topic which we are contemplating investigating is a measurement of $\bar{B}^0 \rightarrow \tau^- \bar{\nu}_\tau D^+$, which has never before been observed, due to the difficulty in isolating a signal. The vertex detector may allow a signal to be found in the 3-prong tau decay mode by vertexing both the tau and the D^{*+} or D^+ candidate, followed by vertexing the tau with the D . Such a measurement, aside from being the first direct observation of this process, could rule out some unusual models in which a virtual charged Higgs plays a large role in mediating the decay.

One of our students is working at Santa Cruz on a measurement of Λ , Ξ , and Ω production in hadronic Z decays. This is an analysis which can be completed on a short timescale using the data already available. It makes extensive use of vertexing (although the silicon detector is not essential, due to the long decay lengths) and is an important measurement for our understanding of s -quark production and baryon production in the fragmentation process. Figure 16 shows the Ξ^- signal that we have obtained from about 13% of our data. Work is under way to analyze the remaining data to try to obtain a signal for $\Omega^- \rightarrow \lambda K^-$.

In the future, when data from LEP-II are available, we plan to concentrate on the search for the Higgs boson, which will rely heavily on the vertex detector to isolate the dominant $b\bar{b}$ decays of the Higgs. The goal is either to detect or rule out the Higgs up to the Z mass. Such work will draw heavily on the experience we are gaining at this time with the vertex detector data.

2.4. SLD EXPERIMENT

2.4.1. EXPERIMENT STATUS

The SLD is very close to having all subsystems complete and fully functional. The one remaining system to bring on-line is the endcap CRID (Cerenkov Ring Imaging Detector). The hardware is all in place, however the electronics is yet to be installed. This year the complete CCD vertex detector has been installed and its readout is a routine part of the data acquisition system. In the enclosed SLD one-event picture (Fig. 17) the first generation vertex off-line software was able to link 19 tracks from the central drift chamber into the vertex detector. The drift chamber systems now have a measured intrinsic resolution of less than 70 microns and work is progressing on a second generation tracking and calibration code. The Barrel CRID has seen Cerenkov rings from both the liquid trays and the gas and the CRID group is working hard on calibration software. When the SLC is not delivering beam to the interaction point (e.g., for installation of the new polarized gun) we run cosmic data to provide calibration data for these two systems. The LAC and WIC, which make up the major calorimeter and muon detector, were the first functional systems and are very stable. The luminosity subsystem has also proved to be quite stable. For the 1992 data run, which started in February, we were running a second-generation acquisition system which was significantly more robust and functional than that of the previous year. This system incorporates close to 500 separate and independent CPU's and now works quite smoothly. We now routinely have multiple triggers which can be tuned to match the SLC background conditions. One last system needs to be mentioned here, that of the energy spectrometer. Technically, it is a part of the SLC rather than the SLD, however, G. Blaylock from SCIPP has made this system work to the point that both SLC and the experiment rely on it for accurate energy measurements in a very routine fashion.

2.4.2. SLC/SLD DATA

The SLD has logged over 1000 Z's in the month prior to a shutdown on April 6, 1992 to install the new polarized gun. The associated muon backgrounds in these events is roughly 0.7 which is half of what it was during the 1991 engineering run. We have had several days of smooth running with average luminosity of 15 Z's per hour. Figure 18 shows the integrated Z rate for this last month of running (both the SLC delivered Z's and the SLD Z's to tape). During the first couple of days of running with the new polarized gun it has delivered 8×10^{10} electrons to the end of the linac with a measured polarization of $> 25\%$ in the switch yard.

After a week or two of spin transport studies we will start data taking again with this new gun with the intent of producing > 1000 polarized Z's onto SLD tapes during May. June through September running will concentrate on both running SLD and sufficient machine luminosity to achieve 10,000 Z's on tape. We expect a significant polarized Z run for the SLD in 1993.

Fig. 17. SLD 1-event display with URTX-linked tracks.

Fig. 18. Delivered and logged SLC luminosity.

2.5. EXPERIMENT E791 AT FERMILAB

2.5.1. INTRODUCTION

Experiment E791 is a fixed-target, hadroproduction experiment based on a large-acceptance, two-magnet spectrometer in the Tagged Photon Laboratory at Fermilab. SCIPP joined E791 in the fall of 1990 under the leadership of P. Burchat. The collaboration completed the data-taking phase of the experiment in January of this year, surpassing its goal of writing 10 billion events to tape by a factor of two.

The principal goal of E791 is to extend the studies of the decay properties of charm mesons and baryons begun by the photoproduction experiment E691 with a twenty-fold increase in data-sample size, a larger silicon microstrip detector system (23 planes instead of 9), and larger angular acceptance of the silicon and spectrometer (partially achieved by moving the target closer to the spectrometer). To handle the high data rate associated with an intense 500 GeV π^- beam and a fairly open trigger,* the data acquisition was completely rebuilt for E791. The new system consisted of a 640-MB event buffer interfaced to 42 Exabyte 8-mm tape drives through event builders in six VME crates. In the five-month period of data-taking, twenty billion events were recorded by this system on 24,000 8-mm tapes. This data set is now being analysed at parallel computing facilities at The Ohio State University, University of Mississippi and Fermilab. Development of offline reconstruction and analysis code is being carried out on computers at most collaborating institutions, including Santa Cruz.

2.5.2. SCIPP'S CONTRIBUTION TO E791

The SCIPP effort on E791 currently consists of P. Burchat, part-time efforts of K. Sugano, and G. Blaylock and P. Gagnon. Sugano and Blaylock are both senior postgraduate research physicists and Gagnon is a fourth-year graduate student. Gagnon worked at Fermilab full-time for the duration of the E791 data-taking. Sugano was at Fermilab for most of the run. Blaylock and Burchat commuted to the experiment for shifts and collaboration meetings.

The contributions of the SCIPP group to E791 have been in two areas: the calorimeters and tracking for the silicon detectors. SCIPP is sharing (with the Illinois Institute of Technology) responsibility for the lead-liquid scintillator electromagnetic calorimeter and the lead-plastic scintillator hadronic calorimeter. Gagnon was extensively involved in maintaining the hardware during the 1991 run

* The main trigger requirement is that the event contain a certain minimum amount of transverse energy in the calorimeters.

and calibrating the calorimeters using special muon and electron runs. Burchat and Gagnon have studied the energy resolution of the calorimeters and are now implementing recursive partitioning for nonparametric classification^[2] for electron identification which should have higher efficiency and lower pion misidentification compared to the method used by E691.

Blaylock has studied the performance of E791's silicon strip vertex detector with measurements of the signal efficiency and noise rates (and patterns) intrinsic to the detector. This has led to a simple but realistic model for the Si detector in the E791 Monte Carlo program. Furthermore, he has written a dictionary-based pattern recognition program for reconstructing tracks from the silicon strip information. He has studied the performance of this track-finder and others, measuring efficiencies for track reconstruction and rates for spurious track production using an assortment of reconstruction algorithms. The results of these studies point to small but significant improvements to be made in E791 silicon reconstruction.

Both Blaylock and Sugano are playing important roles in software certification committees for the offline reconstruction code. Sugano is chairman of the committee to certify the code which formats the output for data summary tapes. Blaylock is on the committee which will certify the drift chamber reconstruction code.

2.5.3. STATUS OF OFFLINE RECONSTRUCTION

Although the offline reconstruction code is not in its final form, we have a version in which data from all parts of the detector is analysed and summary tapes are made. This code is updated nightly on a wide variety of computers at the collaborating institutions, including an IBM RS/6000 at Santa Cruz. Code development is done locally on all of these computers. Changes are then made to the official version which resides on a VAX at Fermilab and is maintained through a code management system. Data is distributed on 8-mm tapes.

Analysis of large numbers of data tapes is done on 'farms' of parallel processors. The offline reconstruction code has been successfully run on 10^8 out of 2×10^{10} triggers (approximately 100 tapes). From this data, we have extracted a charm signal from which we can estimate the number of charm decays we will be able to reconstruct in our entire data set. We estimate that this number will be at least 200,000 – a factor of 20 more than E691.

We are now working on finalizing the offline reconstruction code and improving the efficiency of running code on the farms. We will then start the production run of offline reconstruction of the entire data sample.

2.5.4. PHYSICS TOPICS

We will host an E791 physics workshop at Santa Cruz for four days in July, 1992. At this meeting, we will review the theoretical and experimental aspects of the many physics topics we expect to study with the E791 data sample.

Two studies in which we are particularly interested and which will benefit greatly from the increased sample size of E791 are $D^0\bar{D}^0$ mixing and the form factors in semileptonic D decays.

The twenty-fold increase in data sample size of E791 will allow a more sensitive study of D^0 mixing than that carried out by E691.^[3] With the larger data sample, possible backgrounds can be studied in more detail and perhaps eliminated to enhance the sensitivity of the study. In addition, the larger acceptance of the silicon system and spectrometer will allow increased reconstruction efficiency. It should be possible to extend the sensitivity to mixing down to the 10^{-3} level. In addition, the E791 sample should be large enough to see the first positive signal for the double-Cabibbo-suppressed decays of the neutral D meson.

Form factors in semileptonic D decays have recently received a great deal of attention. Experiment E691 measured the three form factors governing the semileptonic decay $D^+ \rightarrow \bar{K}^{*0}e^+\nu_e$.^[4] The results are significantly different from the predictions of various theoretical models. The E691 study is based on a four-dimensional fit to the decay distributions of 204 candidate events. With such a small data sample and so many free parameters, it is very difficult to estimate the uncertainty on each measured parameter. This sort of analysis will benefit greatly from a twentyfold increase in the size of the data sample.

As well as the two particular analyses discussed above, there is a rich program of other topics including a precise measurement of the D_s lifetime, D_s decay rates (in particular, $D_s \rightarrow \phi e \nu$ from which one can extract an absolute branching fraction when compared to measured D^0 and $D^\pm$ semileptonic decay rates), rare decays of charmed mesons to multi-body final states, excited D mesons and charmed baryons.

2.6. MARK III ANALYSIS

2.6.1. OVERVIEW

The current MARK III author list consists of 47 physicists. Approximately 18 physicists are still actively involved in data analysis or preparing publications. In radiative J/ψ decays we have completed our investigation of the $\eta(1430)$ (a former glueball candidate) and are in the process of publishing our amplitude analyses of $\gamma\pi^+\pi^-$ and γK^+K^- decays. These latter two channels provide access to states with $J^{PC} = \text{even}^{++}$. Scalar mesons, heretofore unseen in radiative J/ψ decays, are now observed. The $f_0(1440)$ is produced in $\gamma\pi^+\pi^-$ and the $f_2(1730)$, produced in γK^+K^- , is measured to be predominantly spin-zero.

We have published a measurement of the 1^{++} axial vector mixing angle using our measured values for the branching ratios $B(J/\psi \rightarrow \gamma f_1(1285))$ and $B(J/\psi \rightarrow \gamma f_1(1420))$. The mixing angle derived from these rates agrees with that obtained from the quadratic mass formula. These results provide the first measurements of axial vector production in radiative J/ψ decays. The rates for $f_1(1285)$ and $f_1(1420)$ production are comparable to those for the $f_2(1270)$ and $f_2(1525)$. This is in contradiction with the naive expectation that spin-one states should be suppressed in the annihilation of two massless gluons.

We have published a precision measurement of the leptonic branching ratio of the J/ψ by measuring $\psi(2S) \rightarrow \pi^+\pi^- J/\psi$. This measurement allows a determination of the strong coupling constant, α_s .

A detailed paper on the resonant substructure in $D \rightarrow \bar{K}\pi\pi\pi$ decays has shown that these decays proceed through quasi-two-body vector-vector and axial vector-pseudoscalar intermediate states. For two modes, the branching fractions for all three isospin combinations have been measured, allowing a determination of the isospin phase shifts. This in turn, allows an improved comparison of the data with theoretical models.

In the following section we provide some detail about the publications in which Santa Cruz physicists as a whole played a major role.

2.6.2. THE $\eta(1430)$

As outlined in last year's report, the $\eta(1430)$, observed as a broad $K\bar{K}\pi$ enhancement in the decay

$$J/\psi \rightarrow \gamma K\bar{K}\pi, \tag{1}$$

was once touted as a glueball candidate. However, an isobar model partial wave analysis of (1) by us has shown the $\eta(1430)$ to be a composite of three states, two

of which are most probably quark-based. These are the $0^{-+}a_0(980)\pi$ near 1.41 GeV/ c^2 (most likely a radial excitation of the $\eta(958)$, henceforth referred to as the $\eta(1400)$), and the $1^{++}K^*K$ near 1.44 GeV/ c^2 (the $f_1(1420)$). In a companion partial-wave analysis of the decay

$$J/\psi \rightarrow \gamma\eta\pi^+\pi^- \quad (2)$$

the $\eta(1400)$ has been observed proceeding through the same partial wave ($a_0(980)\pi$) as in decay (1). This analysis, performed by Mark Burchell (now at CERN), has been accepted for publication in Physical Review Letters.

A prediction for the ratio of branching ratios

$$r = \frac{B(X \rightarrow a_0(980)\pi) \cdot B(a_0(980) \rightarrow K\bar{K})}{B(X \rightarrow a_0^\pm(980)\pi^\mp) \cdot B(a_0^\pm(980) \rightarrow \eta\pi^\pm)} \quad (3)$$

can be made for the decay of X ($X = \eta(1400)$ or $f_1(1285)$) as a function of the coupling strength of the $a_0(980)$ to $K\bar{K}$ relative to $\eta\pi$, (g_K^2/g_η^2). This can be compared with r obtained from the two partial wave analyses. The details of this analysis, performed by William Lockman, were described in last year's report. We are currently writing a paper on this topic, to be submitted to Physical Review D Brief Reports.

2.6.3. LEPTONIC WIDTH AND TESTS OF QCD

The missing mass spectrum for events in which the $\pi^+\pi^-$ recoil spectrum contains a lepton pair,

$$\psi(2S) \rightarrow \pi^+\pi^- J/\psi, J/\psi \rightarrow l^+l^-, \quad (4)$$

is shown in Figures 19(a-b). The inclusive missing mass spectrum opposite the $\pi^+\pi^-$ in the decay

$$\psi(2S) \rightarrow \pi^+\pi^- J/\psi, J/\psi \rightarrow \text{anything} \quad (5)$$

is shown in Figures 19(c-d). Determining the leptonic branching ratio of the J/ψ from the ratio of the number of events in the J/ψ in Figures 19(a) and 19(b) to the number of events in the J/ψ in Figures 19(c) and 19(d) and correcting for the

relative detection efficiency yields

$$B(J/\psi \rightarrow e^+e^-) = (5.92 \pm 0.15 \pm 0.20)\%, \quad (6)$$

and

$$B(J/\psi \rightarrow \mu^+\mu^-) = (5.90 \pm 0.15 \pm 0.19)\%. \quad (7)$$

Assuming lepton universality we obtain

$$B(J/\psi \rightarrow l^+l^-) = (5.91 \pm 0.11 \pm 0.20)\%. \quad (8)$$

where the first error is statistical and the second is systematic. The systematic error from this measurement is almost a factor of four smaller than the previous world average. The derivation of the systematic error is discussed in detail in our publication.

The strong coupling constant α_s is determined by fitting the measured ratios $[\Gamma(J/\psi \rightarrow ggg)/\Gamma(J/\psi \rightarrow \mu^+\mu^-)]$ and $[\Gamma(\Upsilon(1S) \rightarrow ggg)/\Gamma(\Upsilon(1S) \rightarrow \mu^+\mu^-)]$ with the leading-order QCD prediction multiplied by a factor $(1 + \mathcal{A}v^2/c^2)$ to account for relativistic corrections. The free parameters in the fit are $\mathcal{A}$ and the QCD scale factor, $\Lambda_{\overline{MS}}^{(4)}$. We use $v^2/c^2 = 0.24$ and 0.073 for the charmonium and upsilon systems, respectively. The results are

$$\Lambda_{\overline{MS}}^{(4)} = (240 \pm 10)\text{MeV} \quad (9)$$

$$\alpha_s(1.5\text{GeV}/c^2) = 0.30 \pm 0.01 \quad (10)$$

$$\alpha_s(4.9\text{GeV}/c^2) = 0.195 \pm 0.002 \quad (11)$$

where the errors are experimental only. The theoretical errors are unknown and may be large.

Fig. 19 The missing mass distribution opposite the $\pi^+\pi^-$ system for process (4). (a) Data set A. (b) Data set (b). The missing mass distribution opposite the $\pi^+\pi^-$ system for process (5). (c) Data set A. (d) Data set B.

Using the average value of $[\Gamma(\Upsilon \rightarrow \gamma gg)/\Gamma(\Upsilon \rightarrow ggg)] = 2.79 \pm 0.15$ determined from ARGUS, CLEO and CUSB data on inclusive photon decays of $\Upsilon(1S)$ yields $\alpha_s(4.9\text{GeV}/c^2) = 0.179 \pm 0.009$, in qualitative agreement with the results presented above. Running α_s from (10) out to the Z^0 yields $\alpha_s(92\text{GeV}/c^2) = 0.114 \pm 0.001$,

2.6.4. FUTURE PUBLICATIONS

In this section, we list the MARK III papers which are currently under internal review for publication. As can be seen from the list, the results continue to be highly topical and interesting, provided that they can be published in a timely manner. On the other hand, many collaborators have taken up new positions elsewhere. The need for timely publication while meeting the high quality standards set by previous MARK III papers in the light of diminishing manpower makes the role of coordinating the publication effort more crucial than ever. William Lockman has accepted the responsibility for this coordination role because of his expertise, physics judgment and excellent working relations with those at SLAC and throughout the collaboration. We continue to count on his leadership role in this important part of our program.

The following topics are currently under internal review:

1. Amplitude Analysis of $J/\psi \rightarrow \gamma K \bar{K}$ and $J/\psi \rightarrow \gamma \pi^+ \pi^-$.
2. Doubly-Cabibbo-suppressed decays of the charged D meson and mixing in neutral D decays.
3. Study of the $a_0(980)$ Meson in Radiative J/ψ Decays.
4. Evidence for $\rho_1(1600)$ from the decay $J/\psi \rightarrow \pi^+ \pi^- \pi^0$.
5. Decays of J/ψ into a nonstrange vector and Tensor Meson Pair.
6. Hadronic and radiative decays of the $\psi(3770)$ to non- $D\bar{D}$ final states.
7. Production of the $f_0(975)$ meson in J/ψ decays.
8. Measurement of the time-like ρ form factor at $\sqrt{s} = 3.095 \text{ GeV}/c^2$.
9. Measurement of the inclusive photon spectrum from the J/ψ .

REFERENCES

1. SCIPP 92-05, N. Cartiglia *et al.*. Submitted to IEEE Nucl. Symp. 1992.
2. J.H.Friedman, *A Recursive Partitioning Decision Rule for Nonparametric Classification*, IEEE Trans. Comp., 404 (1977).
3. J.C.Anjos, *et al.*, *Phys. Rev. Lett.* **13**, 1239 (1988).
4. J.C.Anjos, *et al.*, *Phys. Rev. D* **62**, 722 (1989)); J.C.Anjos, *et al.*, *Phys. Rev. Lett.* **65**, 2630 (1990)).

3. BIBLIOGRAPHY

MARK III Physics

1. Observation of $f_1(1285) \rightarrow \pi^+\pi^-\pi^+\pi^-$ in radiative J/ψ Decays, T. Bolton, *Phys. Lett.* **B278**, 495 (1992).
2. Resonant substructure in $\bar{K}\pi\pi\pi$ Decays of D Mesons. D. Coffman *et al.*, *Phys. Rev.* **D45**, 2196 (1992).
3. A Direct Measurement of the J/ψ Leptonic Branching Fraction, D. Coffman *et al.*, *Phys. Rev. Lett.* **68**, 282 (1992).
4. Measurement of the Inclusive Decay Properties of Charmed Mesons, D. Coffman *et al.*, *Phys. Lett.* **B263**, 135 (1991).
5. Two Variations on a ρ' Theme: Data from LASS and MARK III, W.M. Dunwoodie *et al.*, presented at Joint Institute Lepton-Photon Symposium and Euromphysics Conference on High Energy Physics, Geneva, Switzerland, July 25–August 1, 1991, SLAC-PUB-5675 (November 1991).
6. An Amplitude Analysis of the $K\bar{K}$ and $\pi^+\pi^-$ Systems ($m < 2 \text{ GeV}/c^2$) Produced in J/ψ Radiative Decay. L. Chen *et al.*, presented at International Conference on Hadron Spectroscopy, College Park, MD, August 12-16, 1991, SLAC-PUB-5669 (October 1991).
7. Evidence for $\rho_1(1600)$ from the decay $J/\psi \rightarrow \pi^+\pi^-\pi^0$ L. Chen *et al.*, presented at International Conference on Hadron Spectroscopy, College Park, MD, August 12-16, 1991, SLAC-PUB-5674 (October 1991).
8. Partial Wave Analysis of $J/\psi \rightarrow \gamma\eta\pi^+\pi^-$, T. Bolton *et al.*, SLAC-PUB-5632 (January 1992). Accepted by *Phys. Rev. Lett.*

MARK II Physics

9. Measurement of the $b\bar{b}$ Fraction in Hadronic Z^0 Decays with Precision Vertex Detectors, R. Jacobsen *et al.*, *Phys. Rev. Lett.* **67**, 3347 (1991).
10. The MARK II Silicon Strip Vertex Detector, C. Adolphsen *et al.*, *Nucl. Instrum. & Methods* **A313**, 63 (1992).
11. Measurement of the Charged Multiplicity of Events Containing Bottom Hadrons at $E(\text{cm}) = 91 \text{ GeV}$, B.A. Schumm *et al.*, SLAC-PUB-5727 (January 1992). Submitted to *Phys. Rev. Lett.*

ALEPH Experiment

12. Recent Results and Running Experience of the New ALEPH Vertex Detector, G. Batignani *et al.*, presented at the IEEE Nuclear Science Symposium, Santa Fe, November, 1991, ALEPH Report 91-158 (December 1991).
13. The ALEPH Double Sided Silicon Strip Vertex Detector, E. Focardi *et al.*, presented at the International Lepton-Photon Symposium, Geneva, July, 1991, ALEPH Report 91-146 (October 1991).
14. Measurement of α_s in Hadronic Z Decays Using All Orders Resummed Predictions, D. Decamp *et al.*, CERN-PPE-92-033 (February 1992). Submitted to Phys. Lett. B.
15. Measurement of $B-\bar{B}$ at the Z Using a Jet Charge Method, D. Decamp *et al.*, CERN-PPE-92-48 (March 1992). Accepted by Phys. Lett. B.
16. Evidence for the Triple Gluon Vertex from Measurements of the QCD Color Factors in Z Decay into Four Jets, D. Decamp *et al.*, CERN-PPE-92-31 (February 1992). Accepted by Phys. Lett. B.
17. Measurement of the tau Lepton Lifetime, D. Decamp *et al.*, CERN-PPE-92-003 (January 1992). Submitted to Phys. Lett. B.
18. Evidence for B Baryons in Z Decays, D. Decamp *et al.*, CERN-PPE-91-229 (December 1991). Submitted to Phys. Lett. B.

Superconducting Super Collider

19. Type Inversion in Silicon Detectors, D. Pitzl *et al.*, *Nucl. Instrum. Methods* **A311**, 98 (1992).
20. Silicon Strip Front End Using a Low-noise Analog CMOS Process, W. Dabrowski and J. DeWitt, presented at the Sixth European Symposium on Semiconductor Detectors, Milan, Italy, February 24-26, 1992, SCIPP 92/02 (February 1992).
21. Pulse Shapes in Silicon Strip Detectors as a Diagnostic Tool, S. Gadomski, M. Turala, N. Cartiglia, J. Leslie and H.F.-W. Sadrozinski, presented at the Sixth European Symposium on Semiconductor Detectors, Milan, Italy, February 24-26, 1992, SCIPP 92/03 (February 1992).
22. Temperature Effects on Radiation Damage to Silicon Detectors, E. Barberis, J.A. Ellison *et al.*, presented at the Sixth European Symposium on Semiconductor Detectors, Milan, Italy, February 24-26, 1992, UCR/SSC/92-01 (February 1992).

23. Noise Measurements on Radiation-Hardened CMOS Transistors, W. Dabrowski *et al.*, presented at the IEEE Nuclear Science Symposium, Santa Fe, NM, November 5-9, 1991, SCIPP 91/24 (January 1992).
24. High Precision Tracking: Silicon Detectors, Gas Microstrips, A. Seiden, presented at 1991 Symposium on the Superconducting Super Collider, Corpus Christi, TX, October 13-17, 1991, SCIPP 91/36 (November 1991).
25. New Experimental Techniques and New Detectors, H.F.-W. Sadrozinski, presented at the 3rd Topical Seminar on Heavy Flavors, San Miniato, Italy, June 17-21, 1991, SCIPP 91/35 (November, 1991).
26. Temperature Effects in the Radiation Damage of Silicon Detectors under Proton Irradiation, H. Ziock *et al.*, presented at the IEEE Nuclear Science Symposium, Santa Fe, NM, November 5-9, 1991.
27. Physics Goals for the SDC Tracking System, B. Hubbard *et al.* (SDC Collaboration), presented at Division of Particles and Fields Meeting of the APS, Vancouver, Canada, August 18-22, 1991. SCIPP 91/19 (August 1991).
28. Tracking Strategies for Magnetic Detectors at High Luminosity and High Energy, A. Seiden, presented at 5th Pisa Meeting on Advanced Detectors, La Biodola, Italy, May 26-31, 1991, SCIPP 91/17 (July 1991).
29. The Silicon Tracking Device for the SSC, H.F.-W. Sadrozinski, presented at 15th International Lepton Photon Symposium at High Energies, Geneva, Switzerland, July-August 1, 1991, SCIPP 91/45 (June 1991).
30. Silicon Tracking Conceptual Design Report (1992), A.J. Weinstein *et al.*, SCIPP 92/04 (March 1992).
31. Radiation Damage to Tektronics SHPi Transistors, N. Cartiglia *et al.*, SCIPP 92/05 (March 1992).
32. Inefficiencies in the Frontend Electronics of the Silicon Tracking System, H.F.-W. Sadrozinski, SCIPP 92/01 (January 1992).
33. *b* Jet Tagging with the SDC Tracking System, B. Hubbard, SDC Note 92-212 (1992).
34. Design and Testing of Fast, Low-Power, Low-Noise Amplifier-Comparator VLSI Circuits, W. Dabrowski, H.F.-W. Sadrozinski and J. DeWitt, SCIPP 91/31 (October 1991). Submitted to Nucl. Instrum. Methods A.
35. Subsystem R&D Progress Report for a Silicon Tracking System: FY 1990-1991, A. Weinstein *et al.*, SCIPP 91/28 (September 1991).

*Budget info removed.
ppp 21-55*

Other Physics

36. Measurement of the Direct Photon Spectrum from $\Upsilon(1s)$ Decays, A. Bizzeti *et al.*, *Phys. Lett.* **B267**, 286 (1991).
37. Status of Gluonium Searches, C.A. Heusch, Proceedings of the 5èmes Rencontres de Physique de La Vallée D'Aoste, La Thuile, Italy, Mar 3-9, 1991, Results and Perspectives in Particle Physics (Editions Frontières, Gif-Sur-Yvette, 1992), CERN-PPE-91-225 (December 1991).
38. Studies for a Tau-Charm Factory, C.A. Heusch, presented at 2nd Symposium on Particles, Strings and Cosmology, Boston, MA, March 25-30, 1991, SCIPP 91/42 (October 1991).
39. Alignment of the SLC Final Focus System Using Beam Orbits, Yu-chiu Chao *et al.*, presented at IEEE Particle Accelerator Conference, San Francisco, CA, May 6-9, 1991, SLAC-PUB-5617 (August 1991).
40. Physics at an Asymmetric Energy B Meson Factory, P. Burchat, invited lectures, Lake Louise Winter Institute: Particle Physics – the Factory Era, Lake Louise, Canada, February 17-23, 1991, SCIPP 91/11 (May 1991).
41. The Search for Scalar Gluonium, C.A. Heusch, SCIPP 92/46 (December 1991). To be published in Nucl. Phys. B (Proc. Suppl).
42. Studies of Helium Gas Mixtures in Drift Chambers, P. Burchat, J. Hiser, A. Boyarski, D. Briggs, SLAC-PUB-5626 (September 1991). Submitted to Nucl. Instrum. Methods A.
43. Analysis of the $\eta\pi^0\pi^0$ Final State in Photon-Photon Collisions, K. Karch *et al.*, DESY-91-126 (October 1991).
44. Search for D^0 and B^0 Decays into $\pi^0\pi^0$. D. Antreasyan *et al.*, DESY-91-076 (January 1991). Submitted to Z. Phys. C.

Silicon Technology Applied to Other Scientific Fields

45. Retinal Readout Array, A. Litke and M. Meister, *Nucl. Instrum. Methods* **A310**, 389 (1991).

4. STATUS OF FACILITIES FOR COMPUTING

Over the past year we have put together a modest UNIX cluster based on the IBM RS/6000 workstations. To date, DoE has provided ~\$100K and the University ~\$50K for the system. As it currently stands, we have four workstations (three IBM and one Sun Sparc2) with seven local X-terminals and ten remote X-terminals (either at SLAC or home). The attached peripherals include: four 8mm tape drives, two Exabyte stacker/robots, six Gbytes of disk space, two uninterruptable power supplies to smooth out the brown UCSC power, a postscript printer, a CD-rom reader, a 1/4" tape drive, modems for the remote X-terminals and extended warranties to cover all the equipment for 3-4 years. We are running an almost exclusive X11R5-based windowing system with TeX, TOPDRAW, XFIG, EMACS, FLIST, MAPLE V, GHOSTSCRIPT (a postscript previewer), and most of the GNU software.

This new cluster is already supported over 20 people doing code development and analysis for ALEPH, SDC, ZEUS, E791, and the theory group. We have operational versions of CERNLIB, the ZEUS libraries, all the E791 code and the ALEPH analysis programs. We are also supporting rad-hard chip layout, the SPICE simulation code and a locally developed pulse simulation package used to support the silicon strip detector development.

The smooth integration and operations of this UNIX cluster is in no small way due to the efforts of F. Staats, our system administrator. A recent graduate from the CIS and Physics Departments at Santa Cruz, he has taken the flow of equipment and made a smoothly running functional system with it. He is currently in the process of porting the NQS batch system to help with teaching the SCIPP group the ins and outs of UNIX which we had very little experience with. With Staats's help we are becoming quite functional with the system. We could not have progressed nearly so far without his expertise, and he is necessary for supporting the system.

This coming year we will add one more workstation (with requisite disk) with an emphasis on 3-D visualization ability. With the complexity of the SDC events and the ZEUS events it is very clear that we now need a workstation with high end 3-D capability to handle the information density present. We will also add a network resident color printer during this year.

We are still in need of X-terminals for the office support staff as well as the labs. With the anticipated heavy usage this year we will learn first-hand what other holes we have in this system.

SANTA CRUZ INSTITUTE FOR PARTICLE PHYSICS (SCIPP)

TASK C

GRANT RENEWAL

1 November 1992 – 31 October 1993

	Page
1. SUMMARY OF RECENT AND ONGOING RESEARCH	1
1.1 NMC/NA-37	1
1.2 SMC/NA-47	2
1.3 Tasks for the Santa Cruz Contingent	10
2. PLANS FOR CONTRACT YEAR 1992/3	18
3. BUDGET AND STAFFING	19
ATTACHMENT	20

TASK C

1. SUMMARY OF RECENT AND ONGOING RESEARCH

DEEPLY INELASTIC LEPTON SCATTERING

This task presently comprises work on two CERN-based muon scattering experiments, NA-37 (the New Muon Collaboration, NMC) and NA-47 (the Spin Muon Collaboration, SMC).

1.1. NMC/NA-37

NA-37 data taking was completed, and the data are in the process of physics analysis. Almost all running periods of the three years of production have by now undergone the basic reconstruction and quality check work prior to the extraction of physics results, and a number of topics have been brought to the publication stage (cf. Appendix).

The principal ambition of NA-37 was the study of structure functions with sufficient accuracy, both statistical and systematic, that important and controversial QCD parameters could be extracted with good confidence. Our prime effort in this context was a precision determination of $F_2^{(p)}$ and $F_2^{(n)}$ over a large range of the scaling variable x . Previous experiments had shown systematic differences: Fig. 1 shows our final results for this ratio,^[1] extending down to $x = 0.003$.

The data are precise enough to reveal a sizable departure of the Q^2 dependence at $0.1 \leq x \leq 0.4$ from perturbative QCD predictions, and permit higher-twist parameterizations that see differences between proton and neutron targets (Fig. 2).

An unexpected result of potentially great importance is again due to the accuracy with which $F_2^{(p)}$, $F_2^{(n)}$ were extracted from the data: the Gottfried sum,^[2] which weights the quark momentum distributions inside the nucleon by their charges, is predicted by the quark-parton model to integrate out to one-third. Our experiment^[3] produces a sizable and significant difference from that prediction:

$$\begin{aligned}
 S_q &\equiv \int_0^1 \sum_i e_i^2 [q_i^p(x) + q_i^n(x) - \bar{q}_i^p(x) - \bar{q}_i^n(x)] dx \\
 &= 0.33 \quad \text{(QPM) ,} \\
 &= 0.240 \pm 0.016 \text{ (experiment) .}
 \end{aligned}
 \tag{1.1}$$

The data are shown in Fig. 3; they are most easily interpreted in terms of a breaking of isospin symmetry ($\bar{q}_i^n \neq \bar{q}_i^p$), or of a flavor-asymmetric sea. There is plenty of theoretical speculation on the origin of our data.

Other NMC results are based on an investigation of the gluon distributions by means of J/ψ production.^[4] We published data on free nucleons (from hydrogen and deuterium), and are presently in the process of doing the same for nucleons bound in heavier nuclei. Clean $\mu^+\mu^-$ mass peaks show the J/ψ signal in Fig. 4, and the resulting gluon distribution is shown together with our earlier EMC results in Fig. 5. The extraction of the momentum fraction of the nucleus carried by the gluons is usually fitted in terms of a function

$$xG(x) = \frac{\eta + 1}{2} (1 - x)^\eta, \quad (1.2)$$

where we determine $\eta = 5.1 \pm 0.9$.

Further significant results have been published both on structure functions in nuclei^[6] – which permit a detailed study of the sensitivity of parton distributions to the size and the mean density of the nuclei in question – and on exclusive vector meson production, both in hydrogen and in heavier nuclei.^[7] This has permitted us to determine $p_\perp$ and t slopes for clean ρ^0 samples (Fig. 6). An interesting conclusion is that the $p_\perp$ slopes at high Q^2 are compatible with a Pomeron exchange mechanism for this exclusive process.

The special trigger we built into the experiment for the investigation of lepton flavor conservation limits ran for the last two periods of data-taking, asking for no outgoing muons but a one-cluster energy deposition in the calorimeter that might indicate an outgoing e^+ or τ^+ ($\rightarrow e^+\nu\bar{\nu}$) in the final state.

A number of tapes with events of this trigger is presently being processed. To avoid radiative backgrounds, they are all on H_2 and D_2 targets, and contain the highest-energy incident beams ($E_\mu = 200, 280$ GeV) for maximal sensitivity to Q^2 -dependent effects beyond the Standard Model. Analysis will continue.

1.2. SMC/NA-47

The presently active experiment, NA-47, is in its second year of execution. A shake-down run at low muon energy (90 GeV) will take until July 1, 1992. At that time, we will take over as prime users of the beam and hope to start serious data taking.

In 1992, the polarized target is still the old EMC target, upgraded to be used with deuterated butanol; it should give us $\sim 28\%$ polarization for neutrons – but the necessary subtraction procedures are liable to lead to an effectively lower polarization.

Fig. 1. The structure function ratio F_2^n/F_2^p averaged over Q^2 as a function of x for the 90 and 280 GeV data. The statistical errors are represented by the error bars. The bands at the bottom indicate the systematic uncertainties.

Fig. 2. The difference between twist-four coefficients for the proton and neutron obtained from the NMC and renormalized SLAC and BCDMS data. The statistical uncertainties are indicated with error bars, the systematic uncertainty by the error band.

Fig. 3. The difference $F_2^p(x) - F_2^n(x)$ (solid symbols and the scale to the right) and $\int_{x_{\min}}^1 (F_2^p - F_2^n) dx/x$ (open symbols and the scale to the left) at $Q^2 = 4 \text{ GeV}^2$. The circles are from the linear fit in $\log Q^2$ and triangles from a fit by L. Whitlow (unpublished). The extrapolated result S_G is indicated by the bar. The simple quark-parton-model (QPM) prediction is also shown.

Fig. 4. Invariant mass distribution of selected $\mu^+\mu^-$ pairs for deuterium and hydrogen. The smooth curves are the sum of a gaussian and an exponentially falling background fitted to the observed distributions.

Fig. 5. The nucleon distribution $xG(x)$ from the present experiment and from the EMC. The solid curve is a parameterization of $xG(x)$ discussed in the text.

Fig. 6. Acceptance corrected distribution of the two-pion invariant mass. The data are for muon scattering on deuterium at 200 GeV and satisfy the cuts listed in Table 1 and the inelasticity cut (1). The full line is a result of a fit assuming a superposition of a ρ^0 p-wave Breit-Wigner and a non-resonant contribution parameterized as $B(m_{\pi\pi}) = a_0(m_{\pi\pi} - 2M_\pi)^{a_1} e^{-a_2 m_{\pi\pi}}$, where a_0, a_1, a_2 are fitted parameters and M_π is the pion mass.

The physics program that we expect to attack this year is mainly a first shot at the spin-dependent neutron structure function $g_1^n(x)$. We recall here that the EMC (with one of us as a collaborator) produced a puzzling set of evidence on the spin-dependent structure function $g_1^p(x)$ of the proton. In the familiar quark-parton, model this function is related to the quark distributions inside the proton by

$$g_1(x) = \frac{1}{2} \sum_i e_i^2 [q_i^\uparrow(x) - q_i^\downarrow(x)] . \quad (1.3)$$

The $q_i^\downarrow, q_i^\uparrow$ symbols stand for the probabilities that the i^{th} quark, of electric charge e_i , have spin down or spin up, respectively, at fractional momentum x . The EMC^[5] had taken data on the asymmetry in the scattering of polarized muons off polarized protons. The measured asymmetry is related to the virtual photon asymmetry

$$A_1 = \frac{\sigma_{1/2} - \sigma_{3/2}}{\sigma_{1/2} + \sigma_{3/2}} , \quad (1.4)$$

where the subscripts refer to anti-alignment and alignment, respectively, of virtual photon and parton spins.

The data, which in the above framework can be seen to scan the spin “content” of the proton in terms of its quarks, led to a major surprise: Fig. 7 shows them together with a straightforward parton model prediction.^[8,9] At x values below 0.15, there is serious disagreement between data and prediction.

This discrepancy – if proven experimentally correct (and this proof is one of the aims of the SMC) – is of major importance. Extrapolating to $x = 0$, as shown in Fig. 8, the integral amounts to

$$\int_0^1 g_1^p(x) dx = 0.113 \pm .012 \pm .025 \quad (1.5)$$

$$\text{for } \langle Q^2 \rangle = 10.7 \text{ GeV}/c^2 .$$

But the well-known Ellis-Jaffe sum rule^[9] (which makes the assumption that the strange quark sea is unpolarized) predicts this integral to be ~ 0.186 , well outside any explainable error.

When we now call the Bjorken Sum Rule^[10] for help—which is not quark-model-based but derives from more rigorous ideas of light-cone current algebra,

$$\int_0^1 (g_1^p - g_1^n) dx = \frac{1}{6} \frac{|g_A|}{|g_V|} \left(1 - \frac{\alpha_s}{\pi}\right) , \quad (1.6)$$

Fig. 7. The asymmetry A_1 in deeply inelastic scattering of polarized muons on polarised protons. The dotted line is a parton model expectation from Ref. 8.

Fig. 8. $xg_1(x)$ (right-hand axis) as a function of x for the proton. The left hand axis shows $\int_{x_m}^a g_1(x) dx$ where x_m is the lower edge of each bin. The smooth curves represent parameterizations of the data.

we can use the EMC data and this rule (where all the quantities on the RHS have their customary significance, and α_s its usual Q^2 dependency) to predict the corresponding integral for the neutron structure function in terms of the quark contents:

$$\int_0^1 g_1^n(x) dx = -0.08 \pm .028 . \quad (1.7)$$

This prediction of a small but significantly negative integral will be tested in the next two SMC runs. One can thus use the two relations (1.5) and (1.7) to solve for individual quark contributions (assuming the s, c seas to be unpolarized), and construct the proton spin that is due to quarks, S_q^p , out of the data:

$$\begin{aligned} S_q^p &= \frac{1}{2} \int_0^1 (\Delta u(x) + \Delta d(x)) dx \\ &= \frac{1}{2} (0.12 \pm 0.20) , \\ &\quad \text{with } \Delta u(x) \equiv u^\uparrow(x) - u^\downarrow(x), \text{ etc.} \end{aligned} \quad (1.8)$$

Only about one-eighth of the proton spin is due to the quarks, and maybe none at all – a pretty startling result that the SMC wants to check. How?

The task is three-fold: (1) check the correctness of the EMC proton result; this will be done not only by an independent new measurement, but also by reaching much lower x values than otherwise available: the planned experiments at SLAC (E-142, -143) can reach $x = 0.015$, with a later 50 GeV electron beam, and 0.04 for the earlier experiments, whereas we are sensitive to $x = 2 \times 10^{-3}$. It has been pointed out^[11] that the gluon “sea” at infinite energy ($x \rightarrow 0$) is particularly dangerous for extrapolation to $x \sim 0$, and the inherent error in the extraction of the Ellis-Jaffe sum may have been strongly underestimated. Note that the EMC results are compatible with lower-energy, higher- x SLAC data^[12], as seen in Fig. 7, but deviate from the prediction only at lower x values. Extending the x range is, therefore, of considerable importance. Note that this extension is kinematically excluded at upcoming SLAC experiments. (2) Measure the neutron value for $\int_0^1 g_1^n(x) dx$, which is predicted to be small but significantly negative. This task, again with the need to extend the integral to the smallest available (x) values, is being pursued by SMC measurements in 1992/3. (3) Search for specific origins of the discrepancies between quark-model-based expectations and data (aside from confirming them). This can be done by several measurements that check on specific models. If the origin of the surprise lies in the assumption of flavor symmetry in

the “sea” (*i.e.*, the assumption that $\bar{d}(x) = \bar{s}(x)$), it is possible to do semi-inclusive meson measurements singling out K or η mesons that have valence $s, \bar{s}$ content. Recall that our recent NMC data on the Gottfried Sum Rule,^[3] probably establish a lack of flavor symmetry^[13] between $\bar{u}(x)$ and $\bar{d}(x)$.^[14] If the fact that the value of

$$\int_0^1 g_1^p(x) dx \sim \langle p | \bar{q} \gamma^3 \gamma_5 q | p \rangle, \quad (1.9)$$

(which is the expectation value of the axial singlet current) does not meet expectations, is related to the axial anomaly,^[14] an inclusive measurement of η' and η meson production may shed light on details. If the most obvious explanation – that the nucleon spin is due to a polarized gluon component of the wave function – is borne out, a measurement of J/ψ production via the well-established photon-gluon fusion process will yield the decisive answer on the gluon distributions. It becomes evident then that the third task will be to look into semi-inclusive production of various mesons. This task is one of the specific aims of our own involvement, as we will detail below.

There is a need for a properly functioning shower detection system that can address the η^0 and J/ψ production questions. This detector can, however, also serve to realize another, far more speculative aim of our involvement: use the deeply inelastic scattering process for a novel test of lepton flavor conservation. This is an issue of very fundamental importance, to which we have devoted considerable effort in our MARK II and MARK III programs. This proponent pointed out in 1986,^[15] that there are no tests of this unexplained tenet of our Standard Model that involve high- Q^2 processes, whereas minimal extensions of the Standard Model would increase strongly with Q^2 . Bigi *et al.*^[16] made quantitative determinations of the sensitivities that could be reached in μN and eN deeply inelastic scattering by looking for processes of the type

$$\mu N \rightarrow e(\tau)X; \quad (1.10)$$

here, the $\mu \rightarrow \tau$ transition is more promising than $\mu \rightarrow e$, but experimentally not as straightforward (in practice, the $\tau \rightarrow e\nu\nu$ decay has to be folded into the detection process). It turns out that sensitivities of order 1.5 TeV (for the mass scale of a new lepton-flavor-violating interaction beyond the Standard Model) can be reached at the highest available muon energies. Indeed, a special trigger using the existing electromagnetic calorimeter was built into NMC experimentation (although no funds were available to optimize that search by means of a more highly resolving pre-shower detector); the data taken with this trigger are expected to yield a first indication of the sensitivity of this method.

In the meantime, this proponent had also pointed out^[17] the complementarity of this (space-like) search for lepton flavor violating boson exchange with other searches involving time-like photons: a subsequent analysis of MARK II data at SLAC^[18] yielded the most sensitive search to date for lepton-flavor-violating processes involving τ leptons at high Q^2 , pushing new-interaction mass scales to about 1.2 TeV. Three of the four LEP Collaborations have adopted the approach of Ref. 18 to launch similar searches, but data taken on the Z^0 pole are less sensitive in terms of new mass scales. For the present purposes, we intend to use the running of the world's highest $-(Q^2, \nu)$ muon beam for a continuation of these studies, but will concentrate some of our effort on the complicated analysis of our data taken during NMC running in the past few years.

There is a third issue involving electromagnetic shower detection in μN scattering that deserves attention: the EMC found^[19] evidence for an anomalous yield of direct photons in deeply inelastic μp scattering, at a level of 0.13 per interaction, at $E_\mu = 200$ GeV. These photons appear mostly at low transverse momentum, $p_\perp$, and low fractional energies, z , unlike the spectra predicted from hadron decay (see Fig. 9). They can be parameterized by

$$\frac{1}{N_\mu} \frac{dN}{dp_\perp} = (2.2 \pm 0.2) \exp((-16 \pm 2)p_\perp^2). \quad (1.11)$$

Fig. 9 shows that the shape of the $p_\perp$ distribution differs markedly from the hadron decay photon distribution; its strong correlation with the current axis is indicative of a quark-level origin, although the magnitude is well above that of expected quark radiation (also indicated).

This effect is of interest not only *per se*, but also due to the fact that the overall signal is on the order of the (unresolved) $\eta \rightarrow 2\gamma$ yield, although of a different shape. Should a connection exist to η production, this would be another process expected to yield asymmetries for different target polarization data (cf. above). If there is no asymmetry, a quark radiation effect becomes more credible.

1.3. TASKS FOR THE SANTA CRUZ CONTINGENT

Continued responsibility of the Santa Cruz group includes two key parts of the spectrometer, the calorimeter H-2 and the beam profile monitor H-6. For present SMC running, the importance of the calorimeter's electromagnetic and hadronic sections has increased; an upgrading of the front section (for better spatial resolution and e/hadron rejection) is discussed below. We are the only "experts" on this massive device and on the triggers incorporating its information; we are presently working on a revamping of its software tasks and its documentation for the Collaboration.

Fig. 9 Direct photon yield from EMC data,^[19] the full line is the expected yield from hadron decay, the dashed lines would be expected for deeply inelastic γ^*q scattering, with 3 MeV, 300 MeV the assumed quark masses.

The beam profile monitor, H-6, is a mosaic of ultra-fast scintillator segments with special readout; its action can be appreciated by a look at Fig. 10. It provides not only shape information to the experiment operators at all times, but also an independent normalization of the beam intensity.

The calorimeter H-2 is a massive device consisting of two interleaved electromagnetic sections and two sequential hadronic sections, of similar construction. The basic method used is a scintillator/absorber sandwich configuration as shown in Figs. 11 a,b. The advantages are a sturdy, large-solid-angle device (with transverse linear dimensions $2.80 \times 5.60\text{m}^2$), arranged in a total of three longitudinal sections: the electromagnetic unit has a total of 40 interleaving horizontal and vertical $280 \times 28\text{ cm}^2$ "fingers" between lead absorbers, to yield redundant information on location and pulse height; the two hadronic sectors consist of horizontal

Fig. 10. Beam profile as measured with the finely segmented scintillator mosaic H-6. Intensity per site is indicated by the size of the symbol.

and vertical Fe/scintillator sandwiches. The electromagnetic thickness is $19X^0$, the hadronic about 4.5λ . The readout has one amplified and one direct channel. From the first, we show a minimum-ionizing (muon) peak in Fig. 11c.

This device has the disadvantages of being crudely segmented ($28 \times 28 \text{ cm}^2$ effective bin size), and of limited energy resolution; in addition, there is a signal attenuation along the 2.80 m length that has to be carefully entered into the readout and analysis algorithms. Nevertheless, H-2 provides valuable information on all of the topics of our particular physics emphasis. To optimize these applications, this year's running will refine our plans to implement a $5X^0$ preshower wall that will permit good spatial resolution of shower trajectories as well as two-shower separation for $\pi^0 \rightarrow 2\gamma$, $\eta^0 \rightarrow 2\gamma$, etc...recognition as well as electron/hadron rejection. The hadronic parts are vital for the recognition of typical deeply inelastic topologies, where the hadronic flux for the target fragmentation and beam fragmentation should be identifiable.

The value of the calorimeter for the experiment has risen from EMC days through two developments. First, we installed a special readout logic that permits triggering on deposition of one or more shower energy clusters in the electromag-

Fig. 11(a). Overall layout of electromagnetic calorimeter H2. Behind each layer of lead, there are a horizontal and a vertical scintillator strip.

Fig. 11(b). Details of joint light collection onto one phototube for the 19 X° thick H2 calorimeter.

Fig. 11(c). Muon spectrum (amplified) in electromagnetic section of H2.

netic section; second, the removal of the Čerenkov counter has left only H-2 for the identification of some hadrons (by shower build-up) and for μ/e hadron separation.

In particular, operation of H-2 permits us to perform the physics tasks mentioned above:

- (a.) Semi-inclusive meson production as a check on isospin (u-d) or flavor (u-d, d-s) symmetry in the spin-dependent quark distributions $\Delta u, \Delta d, \Delta s$. This topic has been partially addressed by Frankfurt *et al.*,^[20] who calculated charged pion asymmetries. Depending on the inaccuracies introduced by the non-identification of charged hadrons, and on the “dilution factors” due to the non-polarized part of the target structure, useful discrepancies can be established to tell the difference between various models. The π^0 content can, for all intents and purposes, be assumed to be $\sim \frac{1}{2} (\pi^+ + \pi^-)$; it will be useful to check inclusive photon production (cf. Fig. 9) against this prediction. A subsequent analysis of any discrepancy can then be related to a conjectured $\eta(\rightarrow 2\gamma)$ contribution. Only the data can decide whether it makes sense to interpret such effects in terms of the η meson distributions, which would be expected to show sizable asymmetries for different polarizations if the strange quark sea were polarized. Again, this topic would be much easier to address if we were to build a pre-shower wall—but an initial attempt is certainly justified.
- (b) Inclusive J/ψ production as a probe of the gluon spin content: a number of theoretical interpretations of the EMC spin anomaly assume a sizable gluon spin alignment. A measurable quantity

$$\Delta\Gamma = \frac{\alpha_S}{\pi} \Delta g, \quad (1.12)$$

where $\Delta g = g_1^p(x) - g_1^n(x)$, would be a natural source for an asymmetry in J/ψ production.^[22] Model-dependent calculations have been made by Glück and Reya^[23] and by Guillet.^[21] Note that while photon-gluon fusion calculations have been able to fit the energy-dependent cross-sections well, the normalizations have been problematic. For our asymmetry measurements, however, these are without interest. What is needed is a background-free sample and good statistics. Our own recent NMC measurements^[24] demonstrate (Fig. 12) that we can easily isolate J/ψ signals in the $\mu^+\mu^-$ mode. The inclusion of the e^+e^- mode with the help of H-2 will not only double the critical statistical sample, but the resolution in the electron mode is naturally superior. A model calculation of the asymmetry to be expected for a reasonable choice of parameters is given in Fig. 13.

Fig. 12. J/ψ signals in dimuon mass spectrum as observed by our NMC experiment,^[24] in both hydrogen and deuterium.

Fig. 13. The expected J/ψ production asymmetry in the appropriate range of kinematics, from Guillet,^[21] calculation. This curve is for $E_\mu = 120$ GeV, $z = 0.7$ (z is the fractional energy).

- (c) Connected with the above shower measurements that demand the absence of charged tracks in the direction of the presumed photons, we will be able to watch out for signals of the prompt-photon yield discussed in the previous section: if they are due to hadron decay, they will display a potential asymmetry as under (b), above; if they are a sign of hard photon-quark scattering, they will depend on the charge but not on the helicity of these quarks, and display no asymmetry.
- (d) Lastly, we will retain the “Trigger 6” or Lepton Flavor Trigger from the NMC logic. This trigger is defined by

$$T_6 \equiv \mu_{\text{in}} \star \overline{\mu_{\text{out}}} \star e_{\text{cand}} , \quad (1.13)$$

where “ e_{cand} ” stands for an electron candidate from possible $\mu \rightarrow e, \mu \rightarrow \tau$ conversion with final-state $\tau \rightarrow e\nu\bar{\nu}$ decay. It is implemented by the demand that there be one and only one cluster of shower energy above a given energy threshold. Data analysis on candidate events taken on H_2 and D_2 targets is presently in progress on the NMC data; we note that the higher- Z materials in the target add potential radiative backgrounds, and that the highest possible Q^2 and E_μ values would be beneficial for these studies.

REFERENCES

1. D. Allesia *et al.*, *Phys. Rev. Lett.* **B249**, 366 (1990).
2. P. Amaudruz *et al.*, *Nucl. Phys. B* in publication (1992).
3. P. Amaudruz *et al.*, *Phys. Rev. Lett.* **66**, 2712 (1991).
4. D. Allesia *et al.*, *Phys. Rev.* **B258**, 493 (1991).
5. J. Ashman *et al.*, *Phys. Rev. Lett.* **B206**, 367 (1988).
6. P. Amaudruz *et al.*, *Z. Physik* **C53** 73 (1992).
7. P. Amaudruz *et al.*, *Z. Physik*. To be published.
8. R. Carlitz, J. Kaus, *Phys. Rev. Lett.* **38**, 673 (1976); J. Kaus, *Nucl. Phys.* **B128**, 219 (1977).
9. J. Ellis, R. Jaffe, *Phys. Rev.* **D9**, 1444 (1974); and E. **D10** 1669 (1974).
10. J. D. Bjorken, *Phys. Rev.* **148**, 1467 (1966), *Phys. Rev.* **D1**, 1376 (1970).
11. F. E. Close, R. G. Roberts, RAL-88-007.
12. G. Baum *et al.*, *Phys. Rev. Lett.* **51**, 1135 (1988).
13. J. Soffer, CPT-91-8-2588 (1991).
14. U. Ellwanger, B. Stech, *Z. Physik* **C49**, 683 (1991).
15. C. A. Heusch, Testing Lepton Flavor Conservation at High Q^2 . *Progress in Electroweak Interactions*, Ed. Frontieres (1986).
16. I.I. Bigi, G. Koepp, P.M. Zerwas, *Phys. Rev. Lett.* **166B**, 238 (1986).
17. C. A. Heusch, SLAC Report 306 (1987); C. A. Heusch and J. J. Gomez-Cadenas, in: *High Energy Physics in the 1990's*, World Scientific (1989).
18. J. J. Gomez-Cadenas, C. A. Heusch *et al.*, *Phys. Rev. Lett.* **66**, 1007 (1991).
19. J.J. Aubert *et al.*, *Phys. Rev. Lett.* **B228**, 248 (1989).
20. L. Frankfurt *et al.*, *Phys. Rev. Lett.* **B230**, 141 (1989).
21. J. Ph. Guillet, *Z. Physik* **C39** 75 (1988).
22. G. Altarelli, G.G. Ross, *Phys. Rev. Lett.* **B212**, 391 (1988).
23. M. Glück, E. Reya, DO/TH 88/4 (1988).
24. D. Allasia *et al.*, *Phys. Rev. Lett.* **B249**, 366 (1991).

2. PLANS FOR CONTRACT YEAR 1992/3

For the new contract year, we envision the following tasks:

- (a) NA-37 – Finalize physics analysis on the lepton-flavor sensitive “Trigger-6” subset of 200, 280 GeV μp , μd data. As has previously been argued, these data will be the first ever to set limits on new-interaction phenomena involving lepton flavor change in high- Q^2 space-like exchanges.

This is a very sensitive analysis task, and has to build on reconstruction, alignment, and efficiency results of other triggers. The presence of NA-37 experts on site at CERN is vital for this work, and one Santa Cruz person will have to be in residence at CERN to participate in all analysis and publication discussions while finalizing these results.

- (b) NA-47 – This calendar year’s NA-47 running extends until December 20, and we expect to have one postdoc, plus one thesis student, participating in data-taking, and responsible for calorimeter maintenance.

To improve the resolution of the H2 calorimeter segmentation, we will attempt to cover the inner 2m x 2m around the beamhole with existing limited streamer chamber devices that are presently still available at CERN. Assembling and testing a prototype module in the muon beam, and getting an appropriate mechanical mount prepared, will cost on the order of \$26,800 (SFr 40,000) – including the procurement, testing, and assembly of the necessary readout electronics. This contribution to SMC apparatus will help in improving both the identification of individual mesons decaying into radiative final states ($J/\psi \rightarrow e^+e^-$, $\eta \rightarrow 2\gamma$, $\omega \rightarrow \pi^0\gamma$), and the electron/hadron separation that is of vital importance for lepton flavor studies.

We hope to complete this improvement work by the time of beam turn-on expected for April, 1993.

The 1993 run is expected to have the new polarized proton/deuteron target available, and acquires major importance for the experiment. We request funding for the presence of two people (one post-doc, one thesis student) at CERN full-time, one graduate student all summer and frequent participation by the P.I. (who will simultaneously be responsible for all senior undergraduates theses in U.C. Santa Cruz, making a full-time presence at CERN impossible (see below)).

Attachment

Recent Santa Cruz/Task C Papers Based on CERN Muon Work.

1. Ratio of J/ψ Production Cross-Sections in Deep Inelastic Muon Scattering from Tin and Carbon. By New Muon Collaboration (P. Amaudruz, *et al.*), Cern-PPE-91-198, Nov. 1991. 20pp. Submitted to *Nucl. Phys. B*.
2. The Ratio $F_2(n)/F_2(p)$ in Deep Inelastic Muon Scattering. By New Muon Collaboration (P. Amaudruz, *et al.*), Cern-PPE-91-167, Oct. 1991. 42pp. Submitted to *Nucl. Phys. B*.
3. Precision Measurement of Structure Function Ratios for Li-6, C-12 and C-40. By New Muon Collaboration (P. Amaudruz, *et al.*), Cern-PPE-91-147, Sept. 1991. 11pp. Submitted to *Z. Phys. C*.
4. Precision Measurement of the Structure Function Ratios $F_2(He)/F_2(D)$, $F_2(C)/F_2(D)$ and $F_2(CA)/F_2(D)$. By New Muon Collaboration (P. Amaudruz, *et al.*), Cern-PPE-91-52, Mar. 1991. 19pp. Published in *Z. Phys. C* **51** 387-394 (1991).
5. The Gottfried Sum from the Ratio $F_2(n)/F_2(p)$. By New Muon Collaboration (P. Amaudruz, *et al.*), Cern-PPE-91-05, Jan. 1991. 10pp. Published in *Phys. Rev. Lett.* **66**, 2712-2715 (1991).
6. Inelastic J/ψ Production in Deep Inelastic Scattering from Hydrogen and Deuterium and the Gluon Distribution of Free Nucleons. By New Muon Collaboration (NMC) (D. Allasia, *et al.*), Cern-PPE-90-178, Nov. 1990. 14pp. Published in *Phys. Rev. Lett.* **B258**, 493-498 (1991).
7. Measurement of the Neutron and the Proton $F(2)$ Structure Function Ratio. By New Muon Collaboration (NMC) (D. Allasia, *et al.*), Cern-PPE-90-103, Jul. 1990. 13pp. Published in *Phys. Rev. Lett.* **B249**, 366-372 (1990).
8. Production of Transverse ρ^0 Mesons in Deeply Inelastic Muon Scattering, D. Allasia *et al.* Submitted for publication (1992).

SANTA CRUZ INSTITUTE FOR PARTICLE PHYSICS (SCIPP)

TASK F

GRANT RENEWAL

1 November 1992 – 31 October 1993

	Page
I. Introduction	1
II. Description of Research–Howard Haber	2
A. Introduction	2
B. Summary of Recent Research Activities	2
C. Ongoing and Future Research Activities	8
III. Description of Research–Michael Dine	12
A. Introduction	12
B. Baryon Number Violation at High Temperature & Energy	13
C. Baryon Asymmetry at the Electroweak Transition	15
D. Electroweak Phase Transition	17
E. Low Energy Supersymmetry	18
F. String Theory	19
G. Ongoing Research and Future Plans	21
IV. Description of Research–Robert Leigh	23
A. Summary of Recent and Ongoing Research Activities	23
B. Future Research Plans	24
V. Description of Research–Alex Pomarol	25
A. Summary of Recent and Ongoing Research Activities	25
B. Future Research Plans	26
VI. BIBLIOGRAPHY	33
VII. STAFFING AND BUDGET	36
A. Staffing	36
B. Budget Justification	37

TASK F

I. Introduction

This is a proposal for continued funding of the Task F effort of the Santa Cruz Institute for Particle Physics (SCIPP) for the period November, 1992—October, 1993. The funding request for the coming year, covering regular operations during the period November 1, 1992 through October 31, 1993, is \$179,768.

In this proposal, we describe a program of theoretical particle physics research. The Principal Investigators are Michael Dine and Howard Haber. In addition, the Task F grant directly supports one post-doctoral position, which is currently occupied by Robert Leigh. We propose to conduct a search in early 1993 for a post-doctoral research associate who will replace Leigh starting in September, 1993. Finally, our group attracts a number of outside theory visitors who are supported by other sources of funding. At present, Alex Pomarol occupies a post-doctoral research position here at Santa Cruz, supported by a Spanish government fellowship. We anticipate that he will remain at Santa Cruz until September, 1993. We are presently considering a post-doc application from Dietrich Bödeker who would be supported by a German government fellowship starting in the fall of 1992. In addition we plan to have two short-term visitors during September–December, 1992: Michael Peskin (supported in part by sabbatical leave from SLAC) and Antonio Dobado (supported by a special grant from the Spanish government).

Howard Haber has worked on phenomenological and theoretical problems associated with the search for the origin of electroweak symmetry breaking. In the Standard Model, electroweak symmetry breaking is generated by the dynamics of an elementary scalar (Higgs) sector which is coupled to the gauge bosons and fermions. Much theoretical attention has also been given to approaches which go beyond the Standard Model, such as supersymmetry and technicolor. Haber has worked extensively on evaluating and developing strategies for detecting phenomena associated with the electroweak symmetry breaking sector at present and future colliders. He has co-authored two major reviews: a *Physics Reports* article surveying the theory and phenomenology of low-energy supersymmetric models, and *The Higgs Hunter's Guide* which is a comprehensive guide to the physics of Higgs bosons.

Michael Dine has worked on non-perturbative behavior of the Standard Model at high energies and temperatures, on “low-energy” supersymmetry and on superstring theory. In the Standard Model, Dine and collaborators have presented arguments that baryon number is badly violated at high temperature, but that such effects are completely negligible in high energy scattering processes. He has

electroweak phase transition using such effects; various problems with this idea have also been elucidated. With Leigh and others, Dine has studied a number of aspects of the electroweak phase transition, which bear on the baryogenesis problem. He is currently constructing models with dynamical supersymmetry breaking and investigating their properties. Dine has been investigating a number of problems in superstring theories, including most recently questions associated with discrete symmetries and strong CP violation.

II. Description of Research—Howard Haber

A. Introduction

My major research activities can be summarized as the examination of methods by which departures from the Standard Model (which encompasses quantum chromodynamics and the Glashow-Weinberg-Salam model of electroweak interactions) may be discovered at colliders which are either presently in operation or at the next generation of accelerators which are now being designed or built. Discovering the physics involved in the spontaneous breaking of the electroweak symmetry is crucial for completing our understanding of the Standard Model and pointing the direction toward the theory that will supersede it. The main focus of this work involves the physics of Higgs bosons, which encompasses the study of the minimal Higgs boson of the Standard Model and the extension to non-minimal models. In particular, much of my attention has been devoted to the study of theoretical techniques which can be used to probe non-minimal Higgs sectors at present and future colliders. In considering the possibility of physics beyond the Standard Model, a very useful prototypical model employs “low-energy” supersymmetry. The search for evidence of such new physics will be an important part of the program envisioned for the next generation of high energy physics colliders which are currently being developed or planned.

B. Summary of Recent Research Activities

During the past year, I have continued my investigations of the physics of Higgs bosons and their implications for Higgs searches at present and future colliders.⁽¹⁾ Despite the enormous success of the Standard Model of strong and electroweak interactions, the exact mechanism by which masses for the gauge bosons and fermions are generated remains uncertain. In the Standard Model, the Higgs boson provides the only direct experimental access to this crucial aspect of the theory. More generally, the Higgs sector (or equivalent) can be expected to provide the most detailed tests of the correctness of any unified gauge theory. It is generally accepted that the mass scale of the phenomena responsible for electroweak symmetry breaking must not exceed the masses of the W and Z bosons by a large factor. Thus, the

ability to either discover or establish the non-existence of Higgs bosons with mass below the TeV scale is a crucial goal for the next generation of hadron and e^+e^- colliders. The appropriate techniques and experiments for fully exploring the Higgs boson spectrum are, however, model dependent.

Despite the simplicity of the Higgs mechanism, the existence of fundamental scalars in field theory is problematical. If the electroweak model is embedded in a more fundamental structure characterized by a much larger energy scale (*e.g.*, the Planck scale, which must appear in any theory including gravity), the Higgs boson would tend to acquire mass of order the large scale due to radiative corrections. Only by adjusting (*i.e.*, “fine-tuning”) the parameters of the Higgs potential “unnaturally” can one arrange a large hierarchy between the Planck scale and the scale of electroweak symmetry breaking. This implies that it is not “natural” to have a Higgs boson that is relatively light unless the size of radiative corrections to scalar masses can be controlled by the structure of the theory. Supersymmetry is one of the most promising theoretical ideas proposed to explain the origin of the scale of electroweak interactions. The minimal supersymmetric extension of the Standard Model (MSSM) is the most economical among models of this type,^[2] and deserves close examination as a candidate for a model of physics beyond the Standard Model.

The Higgs sector of the MSSM is particularly well-constrained and may provide the crucial test which can verify or exclude the simplest model.^[3] The MSSM contains precisely two Higgs doublets and no additional Higgs multiplets. The model possesses five physical Higgs particles: two CP-even scalars (H^0 and h^0 , with $m_{h^0} \leq m_{H^0}$), one CP-odd scalar (A^0) and a charged Higgs pair ($H^\pm$). The parameters of the Higgs sector are fixed at tree level once the ratio of vacuum expectation values (VEVs), $\tan \beta$, and one physical Higgs mass is specified. Here, $\tan \beta = v_2/v_1$, where v_1 (v_2) is the VEV of the Higgs field that couples to down-type (up-type) quarks and leptons. As an example, the tree-level relations among Higgs masses and $\tan \beta$ impose an upper bound on the mass of the lightest CP-even Higgs scalar: $m_{h^0} \leq m_Z |\cos 2\beta| \leq m_Z$. Such a bound, if reliable, would have significant implications for future experiments at LEP-II. In principle, experiments running at LEP-II operating at $\sqrt{s} = 200$ GeV and design luminosity would either discover the Higgs boson (via $e^+e^- \rightarrow h^0 Z$) or rule out the MSSM. (Whether this is possible to do in practice depends on whether Higgs bosons with $m_{h^0} \approx m_Z$ can be detected.^[4]) However, $m_{h^0} \leq m_Z$ need not be respected when radiative corrections are incorporated. In the radiative corrections to the neutral CP-even Higgs squared-mass matrix, the 22-element is shifted by a term proportional to $(g^2 m_t^4 / m_W^2 \sin^2 \beta) \ln(M_t^2 / m_t^2)$.^[5-7] Such a term arises from an incomplete cancellation between top-quark and top-squark loop contributions to the neutral

Higgs boson self-energy. If m_t is large, this term significantly alters the tree-level predictions.

Ralf Hempfling and I computed the exact one-loop expression for the light Higgs mass bound, as a function of all the relevant supersymmetric parameters.^[5] For values of m_t above 120 GeV, the shift in the upper bound of the light Higgs mass can be substantial. For example, if $M_{\tilde{t}} = 1$ TeV and $m_t = 150$ GeV, we find $m_{h^0} - m_Z = 22$ GeV. For $m_t = 200$ GeV, this mass difference exceeds 50 GeV. We concluded that the upper bound of the mass of the lightest Higgs boson of the MSSM will remain in doubt until the top quark is discovered and its mass is determined. Unless the top quark mass is somewhat close in mass to the present CDF limit of 91 GeV,^[8] we would conclude that LEP-200 will not be sensitive to a substantial region of the Higgs sector parameter space of the MSSM. Higher energy colliders will be required to discover or rule out the minimal supersymmetric model.

One can also compute radiative corrections to the full CP-even Higgs mass-squared matrix. Various calculations employing various approximations have appeared in the literature.^[7,9-14] In ref. 10, Hempfling and I present a calculation of the mass-squared matrix in which all leading logarithmic terms are included. The method goes as follows. We take the supersymmetry breaking scale (M_{SUSY}) to be somewhat larger than the electroweak scale. For simplicity, we assume that the masses of all supersymmetric particles (squarks, sleptons, neutralinos and charginos) are roughly degenerate and of order M_{SUSY} . This means that various soft-supersymmetry breaking parameters such as the diagonal squark mass parameter ($M_{\tilde{Q}}$) and the gaugino Majorana mass terms, as well as the supersymmetric Higgs mass parameter are all roughly equal to M_{SUSY} .

One starts with the tree-level expressions for the Higgs masses in terms of the Higgs self-coupling parameters, λ_i of a general non-supersymmetric two-Higgs doublet model. The leading logarithmic expressions for Higgs masses in the MSSM are obtained from the tree-level expressions of the non-supersymmetric model by treating the λ_i as running parameters evaluated at the electroweak scale, m_{weak} . Supersymmetry is then invoked as a boundary condition at the scale M_{SUSY} . That is, at $\mu = M_{\text{SUSY}}$, the λ_i are not independent; instead they are related to gauge couplings as dictated by supersymmetry. Below M_{SUSY} , supersymmetry is broken. Thus, we use the renormalization group equations (RGEs) of the non-supersymmetric two-Higgs-doublet model to evolve the λ_i (and the gauge couplings which appear in the boundary conditions) down to m_{weak} . The resulting $\lambda_i(m_{\text{weak}})$ are then used to compute the Higgs masses and CP-even mixing angle in terms of m_{A^0} and $\tan \beta$.

By making use of the one-loop RGEs (which are solved numerically), the Higgs masses and mixing angle obtained in this way correctly include the leading log-

arithmetic corrections summed to all orders of perturbation theory. One can also derive a simple set of analytic expressions by solving the RGEs iteratively at one-loop. The resulting expressions for the Higgs masses and mixing angle then include the one-loop leading log terms. These expressions serve as a useful check of the full calculation of the one-loop radiatively corrected Higgs masses. Since the complete one-loop calculation is quite complex, it is interesting to investigate the accuracy of the leading logarithmic results. First consider the case of the charged Higgs mass. Marco Diaz and I computed the exact one-loop contribution to $m_{H^\pm}$ due to a loop of fermions and their supersymmetric partners.^[15] (Contributions from Higgs and gauge bosons and their supersymmetric partners are known to be small.^[16]) The largest term in the radiative corrections (in the absence of squark mixing) grows as $m_t^2 \ln(M_t^2/m_t^2)$. However, non-leading logarithmic corrections are not negligible and can lead to significant modifications of the leading logarithmic results. Moreover, if squark mixing is important, large negative corrections to the charged Higgs mass can result in certain regions of parameter space which are proportional to the squark mixing parameters. However, it should be emphasized that the radiative corrections to the charged Higgs mass are significant only for $\tan \beta < 1$, a region of MSSM parameter space not favored in supersymmetric models.

The radiative corrections to the neutral CP-even Higgs mass matrix are more dramatic due to the leading $(g^2 m_t^4 / m_W^2 \sin^2 \beta) \ln(M_t^2/m_t^2)$ correction to the 22-element of the squared mass matrix as previously mentioned. The leading-log formulae obtained by the methods outlined above are expected to be accurate as long as: (i) the scale characterizing supersymmetric masses, M_{SUSY} , is large and sufficiently separated from m_Z (say, $M_{\text{SUSY}} \gtrsim 500$ GeV), (ii) m_t is somewhat above m_Z (say, $m_t \gtrsim 125$ GeV) while still being small compared to M_{SUSY} , and (iii) the squark mixing parameters are not unduly large. In particular, (ii) is an important condition—it is the dominance of the leading $m_t^4 \ln(M_{\text{SUSY}}^2/m_t^2)$ term that guarantees that the non-leading logarithmic terms are unimportant. (In contrast, there is no leading logarithmic contribution to $m_{H^\pm}^2$ that grows with m_t^4 . As a result, the non-leading logarithmic terms tend to be more important as discussed earlier.)

I have checked the reliability of the leading-log formulae for the neutral CP-even Higgs masses in two ways. First, I compared the leading-log results with those of ref. 9 which includes non-leading logarithmic contributions to the scalar mass-squared matrix. Second, Marco Diaz and I have performed an exact one-loop calculation of the neutral Higgs masses for values of $\tan \beta$ near 1. This calculation leads to some major simplifications in the analysis, since at $\tan \beta = 1$, the tree-level light Higgs mass is zero. We confirmed that for $\tan \beta$ near 1, large radiative corrections to the light Higgs mass can easily push m_{h^0} to values beyond the current LEP experimental lower bound of (about) 60 GeV.^[19] Thus, the possibility that

the light Higgs mass is due entirely to radiative corrections is not ruled out!

One may wonder whether the large radiative corrections alluded to above are reliable. Perhaps the perturbative calculations are not trustworthy when the one-loop Higgs mass shift is as large or larger than the tree-level value. In fact, one can show that these large radiative shifts are reliable for values of $m_t \lesssim 250$ GeV. This can be done by estimating the largest corrections that would emerge from a two-loop calculation.^[20] Such an analysis demonstrates that the effective expansion parameter is $g^2 m_t^2 / 4\pi m_W^2$. The surprisingly large mass shifts in the neutral Higgs sector enters due to an extra overall factor of m_t^2 which arises for the first time at the one-loop level. Additional leading logarithmic enhancements can be summed to all orders using RGE techniques described above.

Apart from the large mass shifts to CP-even Higgs masses, a number of other features of the radiatively corrected Higgs masses are noteworthy. There exists a range of parameters in which the tree-level bound, $m_{h^0} \leq m_{A^0}$ is violated. In particular, in the region of small $\tan\beta$ and small m_{A^0} , it is possible to have $m_{h^0} > 2m_{A^0}$, which would permit the decay $h^0 \rightarrow A^0 A^0$. This has been investigated in detail by Ralf Hempfling, Yosef Nir and me.^[21] Once allowed, $h^0 \rightarrow A^0 A^0$ is almost certainly the dominant decay mode, with branching ratios as large as 80% or more when m_{A^0} is small. As m_{A^0} increases beyond 30 GeV, the region of parameter space quickly shrinks where this decay is permitted. The possibility of $h^0 \rightarrow A^0 A^0$ has only recently been incorporated into the LEP Higgs searches. At present, the ALEPH collaboration can only rule out $m_{A^0} > 20$ GeV (at 95% CL) independent of the value of $\tan\beta$.^[22]

Given the significance of the radiative corrections to the Higgs parameters of the MSSM, it is very important to assess the impact of these results on MSSM Higgs phenomenology. In particular, all such analyses performed before 1991 were based on tree-level predictions. The impact of the radiative corrections on Higgs searches at LEP-II is very important as mentioned above. In particular, one can no longer rule out the MSSM if a CP-even Higgs scalar is not discovered at LEP-II. The obvious question to ask is whether the MSSM can be ruled out by virtue of a Higgs search at future colliders. At the next e^+e^- linear collider (NLC), with center-of-mass energy of 300 GeV or higher, the LEP-II Higgs searches can be extended and the complete MSSM Higgs parameter space can be probed.^[23] However, there is as yet no official proposal to build the NLC. Thus, it is more appropriate to first consider the potential for Higgs phenomenology at hadron supercolliders such as the SSC (and LHC). In a series of two papers with Jack Gunion and other collaborators, we explored a variety of techniques for discovering the Higgs bosons of the MSSM at the SSC and LHC.^[24,25] We focused on the production and decays of the CP-even and CP-odd neutral Higgs bosons. Production cross-sections and decay branching

ratios were computed including the one-loop leading logarithmic corrections to Higgs masses and couplings as described above. We then explored the range of parameter space for which the gold-plated signature $Higgs \rightarrow ZZ \rightarrow \ell^+ \ell^- \ell^+ \ell^-$ could be used to detect the Higgs boson. In this process, one of the two Z bosons is off-shell if the Higgs mass lies between m_Z and $2m_Z$. The MSSM Higgs bosons provide a challenge for Higgs hunters for a number of reasons. The light CP-even scalar is lighter than $2m_Z$, and can only be detected in the gold-plated mode if it is heavier than about 130 GeV. Here, the radiative corrections help immensely, since for large m_t , the light Higgs mass is shifted significantly above m_Z . For the heavier CP-even Higgs state, the gold-plated mode is viable only if m_{H^0} is not too heavy, since for $m_{H^0} \gg m_Z$, the $H^0 ZZ$ coupling approaches zero. For the CP-odd Higgs boson, there is no tree-level $A^0 ZZ$ coupling. However, this coupling is induced at one-loop and there exists a region of parameter space (albeit small) in which the A^0 can be detected in its gold-plated mode. In addition, with a complete set of Higgs branching ratios in hand, other techniques for Higgs detection at the SSC can be explored. The techniques with the greatest chance of success are similar to those developed for the Standard Model Higgs boson in the intermediate Higgs mass regime (roughly between m_W and $2m_Z$). Based on our analyses^[24,25] and those of refs. 26-29, it follows that nearly all of the MSSM Higgs parameter space can be explored at the SSC. How well one can ultimately do depends critically on the value of the top quark mass (which enters the analysis in part through its effects on the radiative corrections). Still, there appears to be a region of parameter space which continues to defy all attempts for MSSM Higgs detection; work is progressing to see if it is possible to close this gap.

After two years of running at LEP and nearly two million Z^0 events accumulated, the Standard Model of particle physics continues to provide a remarkably detailed and complete description of all observed high energy physics phenomena. Based on the precision electroweak measurements performed at LEP, one can now say that the Standard Model has been verified to within an accuracy of better than 1%. This places significant constraints on possible new physics beyond the Standard Model. On the other hand, the arguments of naturalness and hierarchy suggest that new physics must enter at around the 1 TeV scale or less. Thus, the precision electroweak measurements may already provide interesting constraints on the nature of this new physics. By computing one-loop radiative corrections to various physical observables, one can indirectly obtain information on "heavy" physics which can enter as virtual particles in the one-loop graphs.

As described above, the one-loop radiative corrections have a dramatic effect on the Higgs sector of the MSSM. Ironically, this effect is in a direction which makes it harder to find evidence for the MSSM at LEP. In addition, we can consider the one-loop effects of supersymmetric particles on various electroweak observables

(ρ , Δr , m_W and A_{LR}). These effects can also be parameterized (in the simplest version of the model) by relatively few parameters. Barbieri, Frigeni, Giuliani and I^[30] examined the effect of radiative corrections on various electroweak quantities as a function of the MSSM parameters. Although some of this work had appeared previously in the literature, our analysis was the first systematic survey of the full parameter space of the model. We demonstrated that the magnitude of the corrections in supersymmetry were too small to be observed over a rather large range of supersymmetric parameter space.

The effects of new heavy particles on electroweak radiative corrections can be neatly summarized by two different combinations of vector boson self-energies. T is proportional to the shift in the ρ -parameter and S counts the number of very massive degenerate chiral weak multiplets. It has been shown that S can be reliably estimated in a class of technicolor models.^[31] New heavy technifermion doublets do not decouple from S , so precision electroweak measurements can potentially rule out such models. In order to better understand the results I obtained in collaboration with Barbieri *et. al.*, I computed S and T in the MSSM as a function of the supersymmetric parameters.^[32] In contrast to technicolor, the MSSM is rather benign: its contributions to S and T are generally quite small and smoothly decouple as the supersymmetric mass scale grows large. I obtained analytic asymptotic forms for the contributions of the various supersymmetric (and the Higgs) sectors to S and T . By examining these results, one can understand why the contributions of supersymmetry to electroweak radiative corrections are small.

C. Ongoing and Future Research Activities

In this section, I shall describe a number of projects which are in progress or under consideration. First, in the consideration of the phenomenology of the MSSM, the parameter $\tan \beta$ is a very important quantity. For example, as mentioned a number of times above, for $\tan \beta < 1$ the Higgs top-quark coupling is enhanced which increases the effects of the top-quark loops on the radiative corrections. It is often said that models of low-energy supersymmetry based on supergravity strongly favor $\tan \beta > 1$.^[33] One can obtain an interesting constraint on $\tan \beta$ by imposing requirement that Higgs-fermion couplings remain finite when running from the electroweak scale to some large energy scale Λ . (Beyond Λ , one assumes that new physics enters.) The limits on $\tan \beta$ depend on m_t and the choice of the high energy scale Λ . Using the renormalization group equations for the gauge and Higgs-fermion Yukawa couplings, one integrates from the electroweak scale to Λ (allowing for the possible existence of a supersymmetry-breaking scale, $m_Z \leq M_{\text{SUSY}} \leq \Lambda$), and determines the region of $\tan \beta$ - m_t parameter space in which the Higgs-fermion Yukawa couplings remain finite.^[34,35] (The t , b and τ are

all included in the analysis.) Based on a one-loop analysis, Fabio Zwirner and I^[34] have obtained a lower and upper bound on $\tan\beta$ as a function of m_t , M_{SUSY} and Λ . For example, if there is no new physics (other than perhaps minimal supersymmetry) below the grand unification scale of 10^{16} GeV, then based on the CDF limit^[8] of $m_t > 91$ GeV, we concluded that $0.5 \lesssim \tan\beta \lesssim 55$. The lower limit on $\tan\beta$ becomes even sharper if the top-quark mass is heavier. Remarkably, the limits on $\tan\beta$ do not get substantially weaker for Λ as low as 100 TeV, where the $\tan\beta$ limits relax to $0.3 \lesssim \tan\beta \lesssim 90$. It would be interesting to repeat this analysis using two loop renormalization group equations. In this case, one would also need the two-loop RGEs for the Higgs self-couplings as well. Although there is no expectation of any major qualitative change, there are examples in the literature in which the quantitative results based on two-loop RGEs does lead to noticeable shifts from results based on the one-loop RGEs. The two-loop RGE analysis will be done in collaboration with D.R.T. Jones and F. Zwirner. Limits on $\tan\beta$ can also be obtained by demanding that all tree-level amplitudes satisfy perturbative unitarity. C.P. Yuan, Tom Weiler and I are reviewing the relevant calculations in the literature and intend to derive an appropriate set of limits for the two-Higgs doublet model (with or without supersymmetry). Limits obtained in this manner simply indicate at what point perturbative calculations are no longer trustworthy.

Radiative corrections to the Higgs sector of the MSSM has had a profound impact on Higgs phenomenology, as discussed above. In particular, very large corrections that grow as m_t^4 have been isolated in the CP-even Higgs mass matrix. A complete one-loop calculation of the radiatively corrected CP-even Higgs mass matrix has recently appeared from two groups.^[9,13] Marco Diaz and I have also been involved in this computation, which generalizes the calculation we performed for $\tan\beta$ near 1.^[14] In the calculation with $\tan\beta$ arbitrary, there are a number of very interesting technical questions that could use some clarification. The observable $\tan\beta$ must be carefully defined in order to fix its counterterm. For example, in ref. 18, a minimally subtracted $\tan\beta$ is used. This is fine for the calculation at hand. Eventually, when a larger set of Higgs phenomena (beyond the CP-even Higgs masses) is measured, one will need to relate this definition of $\tan\beta$ to an experimentally observable quantity. Second, in the analysis of the CP-even mass matrix, one must perform a diagonalization to obtain the two Higgs mass eigenvalues. At tree-level, the diagonalizing angle α is uniquely defined. At one-loop, the procedure is not unique. Again, one must proceed carefully in order to define quantities that are directly accessible to experiment. Diaz and I are attempting to clarify and elucidate these issues.

The leading-logarithmic approximation to the Higgs parameters of the MSSM has proved to be very useful. It provides a rather accurate summary of the largest

corrections to the CP-even Higgs masses (and to the charged Higgs mass to a lesser extent). However, in some cases, certain non-leading logarithmic corrections can also be important. Two examples are: effects due to squark mixing and terms that grow as m_t^2 (with no additional logarithmic enhancement). The possible dominance of squark mixing effects in the corrections to the charged Higgs mass was explicitly shown in the calculation by Marco Diaz and me.^[15] Ralf Hempfling and I are modifying our leading logarithmic analysis in order to include the leading effects of squark mixing. This can be done by altering the supersymmetric boundary conditions to the RGEs at the supersymmetry breaking scale to explicitly include the largest squark mixing terms. Other improvements to this approach can also be implemented. The end result will be a manageable set of formulae which will accurately reproduce the results of the complete (RGE-improved) one-loop computation.

Assuming that Higgs bosons are discovered and experimentally measured with precision at either LEP-II or an NLC, a complete set of radiative corrections to production and decay processes will be needed. For the Standard Model Higgs boson, Bernd Kniehl has been one of the leaders in this area.^[36] For the two-doublet Higgs model and the MSSM, such a program is also important. In particular, assuming one CP-even Higgs state is found, it will be crucial to determine whether such a state is unique or whether it is part of a nonminimal Higgs sector. To answer this question may require precision measurements and the attendant radiatively corrected theory. In the two-doublet model, there are many “natural” relations among couplings. For example, assuming the parameters $\tan\beta$ and the CP-even Higgs mixing angle α have been measured, then $g_{h^0 ZZ}^2/g_{H^0 ZZ}^2 = \tan^2(\beta - \alpha)$. All such natural relations are modified by *finite* amounts when radiative corrections are taken into account. In the MSSM, there are even more such cases, since many tree-level two-Higgs doublet parameters are related by supersymmetry. Bernd Kniehl and I are intending to examine some of the radiative corrections to the Higgs couplings to ZZ in the MSSM and more general two-doublet model. Some preliminary work along these lines is also under way in collaboration with Ralf Hempfling and Thomas Weiler.

In all the projects listed above, the Higgs sector that is being investigated is assumed to conserve CP. Although this is probably a good first approximation, it is worthwhile to return to this assumption and consider the consequences of relaxing it. How would evidence of Higgs-generated CP-violation be detected? One possible approach is to take the most general two-Higgs doublet model with CP-violation and thoroughly explore its consequences. We are particularly interested in whether there are techniques which utilize precision measurements at future e^+e^- colliders that may be sensitive to such CP-violating effects. This work is in progress; collaborators include N. Deshpande, Jack Gunion, Tony Mendez and

Alex Pomarol. As a related problem, I will try to extend an idea previously put forth by Nir and me^[37] in which the magnitude of the CP-violation in the Higgs sector is directly tied to new physics which occurs at an energy scale somewhat (but not too far) above the scale of electroweak symmetry breaking. If a new source of CP-violation were to be found in nature (*e.g.*, the electric dipole moment of the neutron, or perhaps some particular signal of CP-violation in the B -sector which did not fit into the Standard Model), it may be possible to obtain an *upper* bound for the scale of this new physics.

Looking beyond the SSC, information about the origin of electroweak symmetry breaking and physics beyond the Standard Model will eventually come from a very high energy e^+e^- linear collider (NLC). There is now a worldwide effort to develop the necessary technology to make such a machine feasible. In addition, there has also been a concerted effort by various groups to consider the physics potential of such a collider. I am involved in various efforts (centered at SLAC) to investigate in detail the physics potential of the NLC. In September, 1991, I gave a plenary talk at the International Workshop on Physics and Experiments with Linear Colliders in Finland on the theory and phenomenology of Higgs bosons at future e^+e^- linear colliders. The follow-up international meeting is planned for April, 1993 in Hawaii. In order to prepare for this next conference, three two-day workshops on future colliders will be organized in the fall of 1992. In December, 1992, I will be involved in organizing the third of these workshops at Santa Cruz. The topic of this workshop will be Higgs bosons and electroweak symmetry breaking. We hope to attract physicists interested in hadron supercolliders (SSC and LHC) in addition to those working on e^+e^- future colliders. During the next few months, Jack Gunion and I intend to complete our work on the potential for Higgs physics at a $\gamma\gamma$ collider. It may be possible to run the NLC in a $\gamma\gamma$ collision mode via Compton backscattering of laser photons off the linear collider electrons (and positrons).^[38,39] The Higgs boson produced in such collisions would provide an accurate measurement of the $\phi^0\gamma\gamma$ coupling. This coupling is induced at the one-loop level and receives contributions from all virtual charged particles that get mass via the Higgs mechanism. Thus, the measurement of the $\phi^0\gamma\gamma$ coupling would provide significant information on the mechanism of mass generation. Gunion and I have computed the expected event rate for the detection for Higgs bosons of the Standard Model and the MSSM.^[40] Our more recent analysis includes the effects of the MSSM Higgs radiative corrections. We are presently investigating the effectiveness of polarization for enhancing the Higgs signal.

III. Description of Research—Michael Dine

A. Introduction

There is virtually no reason to doubt that the standard model completely and correctly describes physics at currently accessible energies. With the advent of LEP, physicists find themselves performing high-precision tests of the electroweak theory; the planning for the SSC is based on the conviction that we understand thoroughly the high energy interactions of quarks and gluons. However, there are many questions which the standard model leaves unanswered: What is the origin of the large hierarchy between the W mass and the Planck mass? Why do the quark and lepton masses take the values they do? Why does the cosmological constant vanish? Reflecting these puzzles, my interests in particle physics have ranged not only over aspects of the standard model but over possible extensions which might resolve some of these questions. In recent years, my work on standard model physics has focused on two areas. The first of these involves questions related to non-perturbative phenomena, and the roles these might play at high temperatures and energies. This has lead, most strikingly, to an understanding of how the electroweak theory itself might be the origin of the asymmetry between matter and antimatter.^[41] The second area of activity has involved trying to improve our understanding of the phase transition in the electroweak theory. In collaboration with A. Linde (Stanford) and P. Huet (SLAC), R. Leigh and I have demonstrated that within the minimal standard model the phase transition is first order, provided perturbation theory is valid.^[42] We have also made a first effort to understand the propagation of the bubbles which drive the phase transition and whose structure is crucial to understanding baryogenesis in the electroweak theory.

My recent investigations of possible physics beyond the standard model have principally involved supersymmetry and superstrings. I have a longstanding interest in supersymmetric models of particle physics. In the last few years I have concentrated on the study of certain naturalness questions, particularly flavor changing neutral currents and CP violation,^[43] and model building in the framework of dynamical supersymmetry breaking.^[44] Apart from specific models, these efforts have also lead us to reconsider certain aspects of the so-called μ -problem, and the models of continuous R symmetries considered by Hall and Randall.^[45] We have also considered in this context certain questions concerning axions and strong CP which have been raised recently.^[46]

I also have an abiding interest in superstring theory. While from my early work on the subject I have serious doubts that these theories, at least as currently understood, can completely describe nature, they remain extremely interesting as paradigms for unification. It is possible to address in these theories numerous questions which cannot be studied in conventional field theories. During the past year,

with T. Banks (Rutgers) I have used string theories as a framework in which to investigate certain questions about discrete symmetries.^[48] In particular, a number of authors have argued that discrete symmetries, if and when they arise, should be gauged,^[50] and have attempted to write consistency conditions (analogs of conventional anomaly cancellation conditions) which such symmetries should obey.^[51] String theory sheds light on these issues. We have discovered that in consistent string theories, gauged discrete symmetries obey weaker conditions than those which have been proposed. At the same time, we have also discovered that global discrete symmetries do naturally arise, which are only broken by exponentially small non-perturbative effects. Related investigations have also lead Banks, N. Seiberg and I to propose a novel solution to the strong CP problem (and perhaps the problem of the cosmological constant).^[49]

Below I review some of my recent efforts in these areas. The first section reviews earlier work on baryon number violation at high energy and temperature, the second discusses the baryon asymmetry. These are followed by sections on the electroweak phase transition, supersymmetric models, and superstrings.

B. Baryon Number Violation at High Temperature and Energy

Several years ago, Kuzmin, Rubakov and Shaposnikov,^[52] following on earlier suggestions of Dimopoulos and Susskind^[53] and Klinkhammer and Manton^[54] argued that at high temperatures, baryon and lepton number violation would be unsuppressed in the standard model. The basic idea is extremely simple, and can be understood by analogy with a particle in a periodic potential. At zero temperature or low energy, if the barrier is large, the associated barrier penetration factor is exponentially small. However, at high temperatures, one populates states of higher and higher energy, and for sufficiently high energy states the barrier penetration factor is of order one.

If this view is correct, it implies, first, that any baryon asymmetry produced prior to the electroweak phase transition can be severely altered. For example, in theories where $B - L$ is conserved, both the baryon and lepton numbers would be wiped out; in the presence of modest majorana neutrino masses, there is no way to preserve *any* asymmetry.^[55] Perhaps more strikingly, it is possible that this violation of baryon number could be used to understand the origin of the matter-antimatter asymmetry; this will be described in section C below.

While the one dimensional model is suggestive, there has been some debate whether this is really an appropriate analogy for field theory. Many papers, arguing this question pro or con, have appeared since the original work of Rubakov et al.^[56-60] With Lechtenfeld and Sakita at City College, and Fischler and Polchinski at the University of Texas, I investigated this question.^[61] We found that indeed

baryon and lepton numbers are badly violated at high temperatures. We showed, for example, that the field configurations associated with the top of the barrier, called “sphalerons,”^[54] do indeed come to thermal equilibrium on a short time scale. We also showed, contrary to various assertions in the literature, that baryon number is violated in these transitions. We gave general arguments for the behavior of the system above the phase transition temperature, and described a simple model where it is possible, in principle, to perform calculations for all temperatures. Many of our observations were anticipated by earlier work.^[52,57,58] We believe our analysis, however, to be in some respects simpler and more complete. In particular, it makes clear why the problem of baryon number violation at high temperature is different than in scattering of two particles at high energies. For example, Kuzmin et. al.^[52] have suggested that baryon number violation might be enhanced in decays of extremely heavy particles. We showed that, generically, this is not the case. On the other hand, Rubakov^[62] has provided a simple example in which a straightforward calculation shows no suppression of B violation. This example, however, is in some sense the “exception which proves the rule:” in his models, the heavy particle is a very large solitonic state; its form factor, for momenta of order its mass, is exponentially small. Thus it is very hard to *produce* this state. These insights were important in understanding certain surprising results obtained from instanton calculations described below.

One interesting outcome of this analysis was anticipated in numerical studies by Rubakov and collaborators.^[63] These authors had argued that at high temperatures, classical configurations such as monopole-antimonopole pairs would be in thermal equilibrium. We advanced a simple argument that this view is correct. This observation also has cosmological implications, which I plan to explore in the future.

In their original paper, Kuzmin, Rubakov and Shaposhnikov suggested that not only might baryon number violation be significant at high temperature, but it might also be enhanced at high energies.^[52] The underlying idea is again, quite simple: if one has a lot of energy available, it may become easier for the system to pass over the barrier. In refs. 64-66, it has been shown that instanton calculations give B-violating cross sections which, while exponentially small at low energies, grow exponentially rapidly with energy, and it has been suggested that these rates may be of order one at energies of order 10's of TeV. Such a result, if true, would clearly have dramatic implications for machines such as the SSC.

In collaboration with D. Karabali and B. Sakita at City College, and T. Banks and G. Farrar at Rutgers, I investigated this question.^[69] We showed that this problem can be understood by truncating the full field theory to a quantum mechanics model with a single degree of freedom. Semiclassical calculations in the reduced model reproduce exactly the leading instanton results for the Green's functions of

the full field theory. Thus *nothing* is lost in going to this simple system. Indeed it is easy to show, directly in Minkowski space, that this system reproduces the exponential growth of the cross section with energy. The amplitude is proportional to a barrier penetration factor, which grows with energy, times a piece which describes the coupling of the initial state to the reduced system. Because the classical instanton configurations are smooth functions, this coupling vanishes exponentially with energy. These two factors combine to give, for a range of energy, precisely the instanton result. However, this calculation also makes clear that the instanton analysis is very misleading for energies approaching the barrier height. The point is simply that the barrier penetration factor is bounded by one, while the initial and final state overlaps continue to decrease exponentially. Of course, the truncation is not a sensible approximation to the theory for these energies. However, as this simple picture explains completely the instanton result in the energy regime where the instanton calculation is valid, and gives a small result for all energies, we believe no compelling argument has been made that the cross section ever becomes large. Indeed, considering the truncated model one can see that any effort to improve the instanton calculation at these energies faces serious difficulties. While many papers have appeared criticizing the calculations of ref. 64 and 66, our work provides a simple picture of why this analysis is correct at low energies, why it breaks down at high energies, and of what the high energy behavior should in fact be.

The work which has been done on this problem, in my view, raises many interesting questions in field theory. It is certainly not clear that we understand how to compute S-matrix elements in weakly coupled field theories, when there are large numbers of particles in the final state. I hope to pursue some of these more general issues in the near future.

C. Baryon Asymmetry at the Electroweak Transition

Given that the electroweak theory itself violates baryon number at high temperature, it is natural to ask whether or not one can produce the baryon asymmetry purely through electroweak processes, at temperatures of order the weak scale. This would be interesting, not only because the asymmetry might be purely a consequence of low energy physics, but also because it would help resolve various problems in cosmology. For example, in conventional inflationary universe scenarios, it is necessary to reheat to extremely high temperatures in order to produce the asymmetry; in practice, such reheating is difficult to achieve. Moreover, one often finds that various processes (such as heavy particle decays) which occur relatively late can produce entropy and wash out any preexisting asymmetry. Indeed, through the years a number of mechanisms have been proposed for producing the asymmetry at low temperatures, including one by myself and I. Affleck.^[70]

If one is to produce baryons from electroweak effects, it is necessary to have departures from equilibrium. This will be the case, as stressed in the original work of Kuzmin et al, if the weak phase transition is first order. One also needs CP violation. Shaposhnikov,^[71] McLerran^[72] and others have explored the possibility of obtaining the asymmetry in a first order transition, using only sources of CP violation which might be present in the minimal standard model, or the minimal standard model with axions. Shaposhnikov's scenario requires that some rather exotic physics operate; McLerran's scenario fails because CP violation associated with axions is exponentially small at the relevant temperatures. With L. Susskind (Stanford) and two graduate students, P. Huet (presently at SLAC) and R. Singleton (now at MIT), I considered the possibility that the asymmetry might be generated in extensions of the standard model with more CP violation. Our work was stimulated in part by a paper of Turok and Zdrozny.^[73] (Related work was performed at the same time by McLerran, Shaposhnikov, Turok and Voloshin.^[74]) We realized that in a certain limit, the problem could be analyzed in great generality, using elementary thermodynamic notions. We found that generically, in this limit, the asymmetry is of order α_W^6 times CP violating phases and numerical factors depending on the details of the phase transition. Here four factors of α_W come from the baryon number violating rate itself. The additional factors come from an effect which had not been stressed in other work on this subject: the baryon number violating process turns off when the scalar field is of order $\alpha_W T$. In fact, in the two Higgs model analyzed by other authors, the asymmetry is potentially even further suppressed. In our work, we noted that models with singlets can lead to an asymmetry which is substantially larger than α_W^6 .

Subsequent to our initial work on the subject, Huet, Singleton and I continued to refine our estimates and to understand various issues more deeply.^[75] We attempted to obtain improved estimates of transition rates taking account of various finite-temperature effects and the top quark. We also examined two dimensional models in order to study certain troubling theoretical issues.

Many puzzles remain in this subject, and we are some distance from the goal of being able to compute the asymmetry in any given model as a function of the model parameters. I recently reviewed the situation in a talk at the Texas Symposium on these topics (Yale, 1992).^[77] I stressed the need for improved simulations of baryon number violating rates (numerical simulations are necessary for establishing not only the value of the high temperature rate, but also for determining how serious is the suppression discussed in the work with Huet, Singleton and Susskind), for a better understanding of the phase transition, and for further study of the various mechanisms which can generate the asymmetry. My recent and planned efforts devoted to understanding the phase transition will be described in the next section.

Last summer, during a visit to the Hebrew University in Jerusalem, I devoted a great deal of time to studying existing simulations and ways to improve them in discussions with S. Solomon. While this subject has been on the “back burner” recently, I hope to return to it in the near future. As far as mechanisms for producing the asymmetry go, I am trying to understand better a number of ideas which have been proposed by Cohen, Kaplan and Nelson^[52] which appear capable of producing a large asymmetry. I am also trying to understand better some results recently reported by Turok et al, which suggest that the suppression we discussed may not be important.^[78] Some thoughts on all of these questions are described in my Yale talk.

D. Electroweak Phase Transition

The electroweak phase transition was the prototype for studies of cosmological phase transitions in particle physics models.^[79] However, at the time this transition seemed of little observational relevance, so the topic was not thoroughly investigated. With the possibility that this transition might be responsible for the baryon asymmetry this subject has taken on renewed importance. A number of issues connected with the transition were touched on in refs. 41 and 74 (the nucleation of bubbles and some aspects of their propagation, for example). However, the subject remained in a rather unsatisfactory stage. More recently, papers have appeared arguing variously that the phase transition is substantially more first order than originally thought,^[81] that it is substantially less first order,^[80] and that the transition does not proceed through formation of critical bubbles and is unlikely to be relevant to baryogenesis.^[82] Widely disparate estimates have appeared for the bubble wall velocity. While in our first paper on the asymmetry,^[41] for example, we gave an estimate (which I now believe to be incorrect) suggesting that the wall was quite non-relativistic, Turok gave an estimate (which he has since retracted) that the wall becomes ultrarelativistic!^[83]

P. Huet and A. Linde, R. Leigh and I devoted a major effort to trying to resolve these many confusions. We were able to conclusively resolve the question of the order of the phase transition in the minimal model, at least so far as perturbation theory is valid. We showed that the transition is indeed first order, though somewhat less so than the usual one-loop analysis suggests. We also showed that subcritical bubbles are not relevant in the range of parameters for which the conventional analysis would suggest that baryogenesis can occur. Finally, we made a serious, if still somewhat crude, attempt to determine the properties of the bubble. We found, in fact, that the bubble is likely to be on the border between relativistic and non-relativistic behavior, and that it is likely to be rather thick. Similar results have been obtained by different methods by Liu, McLerran and Turok.^[84] All of

these efforts remain crude and incomplete. We are continuing to study this problem, and are likely to collaborate with McLerran et al on improved calculations in the immediate future.

With another student, J. Bognasco, I am also following up some aspects of the perturbative analysis of the effective potential. We are evaluating a number of higher order corrections, in order to better assess the reliability of the perturbation expansion.

E. Low Energy Supersymmetry

In low energy supersymmetric models, in order to adequately suppress various flavor changing processes, it is necessary that the squark and slepton masses exhibit a high degree of degeneracy (at least one part in $10^2 - 10^3$). There exist models for which such degeneracy is natural, but these theories are rather special.^[85] It is not at all clear, for example, how string theory might yield such a degeneracy. With A. Kagan and S. Samuel at CCNY, I investigated the question “under what circumstances might this degeneracy be natural.”^[43] Within the framework of the “minimal supersymmetric standard model” (a model in which one simply adds superpartners for the known fields, and suitable soft breaking terms) we considered theories in which, at some large scale, such as the Planck scale, there is no degeneracy among scalar quark and lepton masses. Then we noted that if, at this scale, the gaugino masses were somewhat larger than squark and slepton masses, renormalization group evolution of these masses to low energies automatically leads to a high degree of degeneracy. It turns out that, imposing certain naturalness requirements at high energy, it is possible to obtain adequate suppression of all flavor changing processes not involving CP violation (e.g. the real part of the $K^0 - \bar{K}^0$ mass matrix). CP violation, however, imposes more serious constraints. These, it appears, can be avoided only by accepting a certain amount of fine tuning or by considering various complications of the model. Some extensions of the minimal model were considered which yield adequate suppression. An alternative, and perhaps more attractive possibility, is that CP violation is somehow suppressed. For example, CP might be conserved in the microscopic theory, but violated spontaneously at some energy a few orders of magnitude below the Planck scale; some further possibilities are mentioned below.

More recently, with a student at UCSC, Doug MacIntyre, I have been considering the problem of constructing realistic models with dynamically broken supersymmetry. Some time ago, Ian Affleck, Nathan Seiberg and I showed that non-renormalization theorems are frequently violated by non-perturbative effects, and that in some theories supersymmetry is dynamically broken and the ground state is stable.^[86] However, at the time we found it difficult to construct realistic models

with dynamical supersymmetry breaking. We considered a variety of approaches. One possibility is to break supersymmetry at scales of order the weak scale. This seems generally to run afoul with dangerous axions and goldstone bosons. Alternatively, one can try to use one of these theories with dynamical supersymmetry breaking as the hidden sector of a supergravity model. Here at first sight one runs into a problem with obtaining adequate mass for gauginos. The difficulty arises when one considers the effective lagrangian at energies somewhat above M_W . If one examines the general form of the supergravity lagrangian given in ref. 87, one finds that gaugino masses can only arise through very high dimension operators. This is because, in general, all of the fields in the hidden sector are charged under some of the gauge symmetries of the theory. (This is not the case for conventional models, and a simple operator of dimension five is allowed which has precisely the correct effect.) Assuming that the extra dimensions are made up by powers of M_P , one finds that gluino masses are of order small fractions of an eV .

Inspired by some recent work of Hall and Randall,^[45] we have reconsidered this question.^[44] These authors argued that one should consider models with continuous R symmetries. Such symmetries, in general, forbid gaugino mass terms; to obtain an acceptable phenomenology, it is necessary to add additional fields, such as color octets. The addition of such fields can also solve the problems described above. However, both in our context and in that of Hall and Randall, we noted that there is a serious problem. We showed that a gluino mass can only arise in this framework if the hidden sector possesses a $U(1)$ gauge field whose auxiliary (" D ") component has an expectation value. However, under these circumstances there is the danger of developing a large Fayet-Iliopoulos term for hypercharge.^[68] We argued that such a term can be avoided if hypercharge is unified at some scale into a large group, and discussed some examples. In the end, the models we have developed are not beautiful, but they tend to make interesting predictions for additional light particles.

We have also considered a variety of issues in models of the type considered by Hall and Randall. We have argued that naturalness dictates a somewhat different field content than they considered, explored the resulting phenomenology. Necessarily one predicts a variety of comparatively light (mass of order $\frac{M_Z}{2}$) particles in this framework. We have also extended their analysis to theories with discrete R symmetries, and have noted some curious aspects of the cosmological constant problem in these models.

F. String Theory

I am interested in string theory, not only as a candidate unified theory of interactions, but also as a model in which one can address questions which cannot

be settled in conventional field theory. In the past I have worked on a range of problems: the problem of the dilaton, non-renormalization theorems for string perturbation theory, the role of non-perturbative corrections in the σ -model analysis, questions of naturalness and the existence of $(0,2)$ string models, aspects of the moduli space of $(2,2)$ conformal field theories, the meaning of $R \rightarrow \frac{1}{R}$ duality, and others. Here I describe some more recent activities.

Over the past year, I have published two papers on string-related questions. The first, with Tom Banks, addressed certain questions related to discrete symmetries in string theory. It has been argued (most compellingly by Krauss and Wilczek^[50]) that global discrete symmetries are unnatural, particularly in any framework which includes gravity. For example, wormholes would be expected to violate such symmetries. In this view, only gauged discrete symmetries appear reasonable. Ibanez and Ross^[51] attempted to formulate a set of conditions which any gauged discrete symmetry must satisfy. Banks and I noted that the full set of conditions discussed by these authors are not satisfied in many consistent string compactifications, and understood the hidden assumption in the Ibanez-Ross work. In the end, it turns out that the only valid constraints are those that can be understood in terms of instantons of the low energy field theory.

Perhaps more interestingly, in the course of this work, we were lead to ask the following question: is it possible to find anomalous discrete symmetries in string theory? In other words, could one find examples of gauged discrete symmetries (e.g. general coordinate transformations) which were violated by low energy instantons. If so, one would have found a non-perturbative consistency condition (analogous to Witten's $SU(2)$ anomaly) for string theory. A quick survey revealed many examples of such symmetries. However, in all cases it was possible to cancel the anomaly by assigning a non-trivial transformation law to the (so-called "model-independent") axion under the symmetry. This is disappointing, since such a consistency condition might have significantly narrowed the space of allowed compactifications. But the result is interesting from another point of view. While the non-anomalous symmetry is spontaneously broken in these cases, the original symmetry of perturbation theory is broken only by non-perturbative effects. Thus, from the point of view of string theory, weakly broken, global discrete symmetries seem completely reasonable.

With T. Banks and N. Seiberg, I considered certain aspects of the strong CP and cosmological constant problems.^[49] This work was not specifically string theoretic in character, but I present it here since it is perhaps even more speculative. The goal of the exercise was to exhibit a solution of the strong CP (and perhaps cosmological constant) problems which did not imply the existence of a light particle such as an axion. Indeed, the model we considered possessed an axion, but

this axion coupled to two strong gauge groups, one with a scale much larger than the QCD scale. Ordinarily, such an axion would have no effect on the θ problem of QCD. However, here we assumed that the periodicities of the two strong groups with respect to this axion were not relatively rational. As a result, it is simple to show that the theory possesses an infinity of vacua with arbitrarily small values of θ . We showed that anthropic considerations would tend to favor states with extremely small θ (though not necessarily quite small enough values of the cosmological constant). Extensions of these ideas can in principle also solve the cosmological constant problem. We were not, however, able to find a convincing example, either in conventional field theory or string theory, of such an axion, though we listed some possibilities in string theory.

Recently, motivated by questions in non-critical string theories, Lance Dixon and I have been considering some issues involving tachyons in superstring theories. We begin with the observation that in many superstring compactifications, tachyons appear at some special value of the moduli. At these points, one can compute the potential for the tachyon systematically; close to these points, one can investigate whether a stable solution (with non-vanishing cosmological constant) exists. We have computed the potential in many cases, and are presently exploring various aspects of the resulting solutions. We are also comparing our results with related work of Antoniadis and Kounnas.^[89]

G. Ongoing Research and Future Plans

A number of ongoing research projects have been mentioned above. I intend to continue my efforts to better understand the phase transition, and to further understand baryogenesis in various extensions of the standard model. I am particularly interested in constructed theories in which the transition is more strongly first order, and the adiabatic types of analysis are invalid. I hope also to return to the question of improved numerical simulations of baryon-number violating rates.

The work described above on dynamical supersymmetry breaking is largely completed. In the course of this work a number of interesting questions, related to supersymmetry and the computation of certain masses and couplings have arisen which I plan to investigate further with my student Doug MacIntire.

There are a number of questions I wish to pursue in string theory. I have already mentioned the problem of tachyons in superstring theories. We hope to apply the results of this analysis to various problems both in critical and non-critical string theories. My student, Doug MacIntire, has continued the search for possible anomalous discrete symmetries in string theory, and is now examining large classes of models by computer. In addition, I plan to return to a set of problems raised earlier in work with Banks, Dykstra and Fischler.^[90] In that

paper, we constructed monopole solutions of string theory and considered some of their properties. The existence of these solutions poses a number of intriguing questions, which are not easily answered in the standard conformal field theory framework. For example, how does one describe, from the point of view of the two-dimensional, world sheet theory which describes these configurations, the soliton mass and charge? How does one introduce collective coordinates for translations and gauge transformations? What role do enlarged supersymmetries play in the case of solutions to the heterotic string theory and superstring theories? In the intervening period, a large literature on classical solutions has developed, and E. Witten (private communication, unpublished) has made some interesting suggestions about these questions. The resolution of these questions should take one a bit beyond the standard first-quantized framework for string theory, without requiring a full second quantized treatment. I am also following other recent developments concerning classical solutions (concerning black hole solutions, in particular).

I plan, as well, to continue to examine various phenomenological questions in string theory. For example, I intend to study recent suggestions concerning supersymmetry breaking due to Dixon, Kaplunovsky, Louis and Peskin.^[91] Some time ago, N. Seiberg and I had argued that it would be difficult to find stable ground states of string theory with broken supersymmetry at weak coupling.^[92] These authors, exploiting the various loopholes we had listed to our arguments, appear to have done just that. Without describing this ingenious idea in detail, let me simply note that there are a number of interesting questions which it raises. First, there is the question of whether corrections to this result are small. Second, one would like to understand the “phenomenology” of these ground states (for example what is the value of the cosmological constant). Third, one would like to explore other variations of this idea, bringing to bear previous experience with supersymmetry breaking.

During a recent talk at the Cincinnati Symposium on Quantum Gravity (in honor of the retirement of Louis Witten) I spoke on certain questions of naturalness and corresponding lessons from string theory. In the course of preparing this talk, I considered a number of issues related to the strong CP problem, which I hope to pursue further. First, after noting that string theory automatically gives Planck scale axions, I showed that discrete symmetries could rather easily give rise to accidental Peccei-Quinn symmetries, with much smaller decay constants, suppressing symmetry-violating operators up to very high dimension. I intend to pursue further the question of whether string models exhibit such symmetries, and whether one obtains any other generic predictions from such scenarios. I also considered in this talk the question of whether string theory might be an example of a theory with spontaneous CP violation, in which θ is completely calculable. With my student, Doug MacIntire, and with Rob Leigh, we have already been

obtained by Choi, Kaplan and Nelson.^[94] It turns out that in string theory, CP is a gauge symmetry and can only be spontaneously broken. Of course, while θ may be calculable, it is likely to be large in general. We are investigating particular models in the hope of understanding under what circumstances the effective θ may be small enough. Again, the hope is that some generic predictions might emerge from such scenarios.

In conclusion, I should add (as should any theorist) that it is quite likely that topics of interest will arise which are not described in this proposal. In the past, I have worked on many topics not mentioned above (axions, standard model perturbation theory, strong CP violation, other aspects of string theory, etc.). Most importantly, I expect to become involved in projects more directly supportive of SCIPP and SLAC experimental efforts.

IV. Description of Research—Robert Leigh

A. Summary of Recent and Ongoing Research Activities

Electric Dipole Moments and CP Violation: Recent work by Weinberg suggests that a dimension-six operator could induce large contributions to the electric dipole moment of the neutron. With four collaborators, I calculated this effect in extensions of the Standard Model with softly broken supersymmetry.^[95,96] The effect is suppressed by renormalization group effects, but is similar in size to other contributions.

A second work^[97] was inspired by that of Barr and Zee who suggested that the electron electric dipole moment in multi-Higgs models receives a contribution from a two-loop graph that is larger than the standard one-loop result. I, with two collaborators, discovered and computed the full set of such graphs. These results can have important consequences for model-building as experimental accuracy increases.

Constraints on the Baryogenesis Scale:^[98,99,96] In recent years it has been realized that at temperatures above the electroweak symmetry breaking scale (such as occurred in the early universe), there are unsuppressed interactions that violate baryon and lepton number through electroweak anomalies. It is thus important to ensure that, whatever the origin of baryogenesis, the baryon asymmetry should survive this epoch. Since the anomalous electroweak processes preserve $B - L$, we can derive bounds on the scale at which $B - L$ asymmetry must have been produced in the early universe. I, with three collaborators, examined in particular the out-of-equilibrium decays of heavy right-handed neutrinos as appear in many GUTs, as well as the effects of $B - L$ violation in other extensions of the Standard Model.

Majoron Physics:^[100] Recently, corroborating evidence for the existence of a neutrino of mass 17 keV has been obtained. The existence of such a particle creates theoretical problems, some of which may be alleviated by introducing right-handed singlet neutrinos along with a spontaneously broken lepton number symmetry. The Goldstone boson associated with the symmetry is called the Majoron. I, with several collaborators, have been investigating a wide range of astrophysical, cosmological and laboratory bounds that might place limits on the properties of these new particles.

Electroweak Phase Transition:^[42,101] It is also possible, and theoretically appealing, that anomalous electroweak processes may have produced the observed baryon asymmetry at 'low' energies of order the electroweak scale in the early universe. A greater understanding of how the phase transition associated with spontaneous breaking of $SU(2) \times U(1)$ proceeds is of fundamental importance in a determination of the resulting baryon asymmetry. Recently I, with several collaborators, have made important improvements to the calculation of the finite temperature effective potential for the order parameter of the transition. The transition is found reliably to be of the first order, and proceeds by nucleation and subsequent expansion of critical bubbles of true vacuum. The baryon asymmetry in these scenarios is typically produced in, or just in front of, the walls of these bubbles and the total number of baryons produced is a function of various properties of the wall, notably the speed at which it propagates through the medium, and its thickness. We have studied these properties using simple approximate methods. We have found that the wall tends to be rather thick, being set by a momentum scale which is small at the transition. Furthermore, we find that a net damping force is present when equilibrium of the surrounding plasma can not be maintained at the wall, and that this typically leads to wall velocities of order several tenths of the speed of light. We are now pursuing a more detailed understanding, particularly the effects of the above on the produced baryon asymmetry, including possible new effects arising from instabilities leading to viscous fingering of the bubble walls.

B. Future Research Plans

I plan to pursue many aspects of the research detailed above. In particular, I am now working on a more detailed understanding of the properties of bubble walls at the electroweak phase transition. It is now roughly understood what effects are important for determining the speed of the wall and refinements in this very difficult calculation are being carried out. There are also new phenomena that may be important and that have been neglected up to now. For example, in all treatments the bubbles are assumed to be spherical with smooth walls. There is some evidence from the quark-hadron transition that this may not be so in the electroweak case as well, namely that there are important instabilities that could

treatments the bubbles are assumed to be spherical with smooth walls. There is some evidence from the quark-hadron transition that this may not be so in the electroweak case as well, namely that there are important instabilities that could lead to a dendritic or foamy wall profile. We are now investigating whether or not this effect is important for baryogenesis.

It is also important to more fully understand the limits of calculability of the effective potential at finite temperature. The perturbation expansion breaks down in a certain range of parameter space due to infrared divergences. It has recently been understood how to improve the convergence of the expansion, and in our previous work, we have found important results in this regard. More detailed work is necessary to understand exactly the limits of applicability of the expansion for calculations of the baryon asymmetry, *etc.*

In recent years it has been suggested that certain string theories in non-critical dimension are exactly solvable up to some non-perturbative input parameters. The presence of such parameters is of fundamental importance, in that, if they are not present, then one expects, at least in principle, that all effects are completely calculable in string theory. Recently, we have been studying a particular class of such parameters in heterotic superstring theory, namely θ -parameters associated with the strong CP problem.

V. Description of Research—Alex Pomarol

A. Summary of Recent and Ongoing Research Activities

Radiative Corrections to Higgs bosons decays in the two-Higgs-doublet model: Experimental results indicate that the t -quark mass is large^[8] and can therefore induce large virtual effects in processes involving Higgs bosons. In an initial paper,^[102] A. Mendez and I computed the corrections to the charged Higgs hadronic width arising from loops involving a heavy quark. In a second paper^[103] we extended the analysis to the neutral Higgs bosons and computed the $\mathcal{O}(\alpha m_t^2/m_W^2)$ corrections to their couplings to fermions and gauge bosons.

CP Violation in the Higgs sector: Higgs induced CP-violating effects can appear in theories with a non-minimal Higgs sector. A general two-Higgs doublet extension of the Standard Model provides the simplest setting for Higgs-induced CP-violation. In such a model, either explicit or spontaneous CP-violation is possible. Mendez and I have examined possible ways of unraveling a CP non-invariant Higgs sector through the interactions of Higgs bosons with gauge bosons.^[104] For example, by studying Higgs pair production and Higgs- Z associated production, one could determine that no set of CP-quantum numbers was consistent with the

observed data. We also have derived some general sum rules for the Higgs-gauge bosons couplings which can be used to obtain information about the scalar content of the Higgs sector.

Spontaneous CP violation in supersymmetric models: The supersymmetric extension of the Standard Model contains new sources of CP violation. It has been known for a long time that the contribution of these new sources to the neutron electric dipole moment is two or three orders of magnitude larger than the present experimental limit if the phases that parametrize the CP violation, φ , are of $\mathcal{O}(1)$. Thus a fine-tuning is required, $\varphi \lesssim 10^{-2} - 10^{-3}$. Since these CP violating phases come from different sectors of the supersymmetric model, these multiple fine-tunings seem to be unnatural. One possible and very elegant solution to this problem is to require that CP is spontaneously broken.

We study this possibility in the Minimal Supersymmetric Model (MSSM). Spontaneous breaking cannot occur at tree-level but may be possible if radiative corrections to the Higgs potential are taken into account. We analyze the physical spectrum of this model and show that a light scalar is required in order to be consistent with spontaneous CP invariance. Considering recent LEP Higgs searches, we show that this can only take place in a very small and disfavored region of the MSSM parameter space. In a second work we consider an extension of the MSSM in which one adds an $SU(2) \times U(1)$ singlet Higgs superfield. In this case, spontaneous CP violation is possible over a large region of the parameter space. Papers on both these models are currently in preparation.

B. Future Research Plans

Future plans consist of completing the projects outlined above. It is of interest to determine whether it is possible to construct a non-minimal supersymmetric model in which all CP-violating phenomena can be explained in terms of one relative phase between Higgs vacuum expectation values. In a related project, I intend to consider the idea of spontaneous CP violation in supersymmetric models with radiative fermion masses.

A second project currently in progress concerns the analysis of the radiative corrections in the Higgs sector using the global symmetries of the Higgs potential. The basic idea is to first identify a (partially respected) symmetry responsible for some (natural) tree-level relation of the theory. By investigating the nature of the symmetry breaking one can determine that the contribution of certain sectors to the radiative corrections to the tree-level relation must vanish in some appropriate limit of the model parameters. This allows one to estimate the size of the radiative corrections in a simple way. A well-known example in the Standard Model is the so-called *custodial* isospin symmetry which plays a crucial role in the analysis of

the radiative corrections of the ρ -parameter. In the MSSM, Haber and I have shown that an approximate extended *custodial* symmetry does not allow radiative corrections to the charged Higgs mass of $\mathcal{O}(g^2 m_t^4 / m_W^2)$ in the absence of squark mixing. Other interesting examples of similar type are now being examined.

REFERENCES

1. For a review and a guide to the literature, see: J.F. Gunion, H.E. Haber, G.L. Kane, and S. Dawson, *The Higgs Hunter's Guide*, (Addison-Wesley, Redwood City, CA, 1990).
2. H.P. Nilles, *Phys. Rep.* **110** (1984) 1; H.E. Haber and G.L. Kane, *Phys. Rep.* **117** (1985) 75.
3. J.F. Gunion and H.E. Haber, *Nucl. Phys.* **B272** (1986) 1; **B278** (1986) 449; **B307** (1988) 445.
4. Z. Kunszt and W.J. Stirling, *Phys. Lett.* **B242** (1990) 507; N. Brown, *Z. Phys.* **C49** (1991) 657; V. Barger, R.J.N. Phillips and K. Whisnant, *Phys. Rev.* **D43** (1991) 1110; D.J. Summers, *Phys. Lett.* **B274** (1992) 209.
5. H.E. Haber and R. Hempfling, *Phys. Rev. Lett.* **66** (1991) 1815.
6. Y. Okada, M. Yamaguchi and T. Yanagida, *Prog. Theor. Phys.* **85** (1991) 1; *Phys. Lett.* **B262** (1991) 54; J. Ellis, G. Ridolfi and F. Zwirner, *Phys. Lett.* **B257** (1991) 83; **B262** (1991) 477; R. Barbieri, M. Frigeni, and F. Caravaglios, *Phys. Lett.* **B258** (1991) 167.
7. R. Barbieri and M. Frigeni, *Phys. Lett.* **B258** (1991) 395; A. Yamada, *Phys. Lett.* **B263** (1991) 233.
8. F. Abe *et. al.* [CDF Collaboration], *Phys. Rev. Lett.* **68** (1992) 447.
9. P.H. Chankowski, S. Pokorski and J. Rosiek, *Phys. Lett.* **B274** (1992) 191.
10. H.E. Haber and R. Hempfling, SCIPP-91/33 (1992); H.E. Haber, SCIPP-92/11 (1992) to appear in the Proceeding of the International Workshop on Electroweak Symmetry Breaking, Hiroshima, Japan, 12-15 November 1991.
11. D.M. Pierce, A. Papadopoulos, and S. Johnson, LBL-31416 (1991).
12. K. Sasaki, M. Carena and C.E.M. Wagner, MPI-Ph/91-109 (1991).
13. A. Brignole, CERN-TH-6366/92 (1992).
14. M.A. Diaz and H.E. Haber, SCIPP-91/13 (1992).
15. M.A. Diaz and H.E. Haber, SCIPP-91/14 (1991), to be published in *Phys. Rev. D*.
16. J.F. Gunion and A. Turski, *Phys. Rev.* **D39** (1989) 2701; **D40** (1989) 2333.
17. A. Brignole, J. Ellis, G. Ridolfi and F. Zwirner, *Phys. Lett.* **B271** (1991) 123; [E: **B273** (1991) 550].
18. A. Brignole, preprint DFPD/91/TH/28 (1991).

19. S.L. Wu, invited talk given at The SSC Physics Symposium, Madison, WI, 13-15 April 1992.
20. J.R. Espinosa and M. Quiros, *Phys. Lett.* **B267** (1991) 27.
21. H.E. Haber, R. Hempfling and Y. Nir, SCIPP-91/40 and WIS-92/5/Feb-PH (1992).
22. D. Decamp *et. al.* [ALEPH Collaboration], *Phys. Lett.* **B265** (1991) 475.
23. H.E. Haber, SCIPP-92/10 (1992), to be published in the Proceedings of the Workshop on Physics and Experiments with Linear Colliders, Saariselkä, Finland, 9-14 September 1991.
24. J.F. Gunion, R. Bork, H.E. Haber and A. Seiden, UCD-91-29 and SCIPP-91/34 (1991), *Phys. Rev. D*, in press.
25. J.F. Gunion, H.E. Haber, and C. Kao, UCD-91-32 and SCIPP-91/44 (1991), *Phys. Rev. D*, in press.
26. J.F. Gunion and L.H. Orr, UCD-91-15 (1991).
27. V. Barger, M.S. Berger, A.L. Stange, and R.J.N. Phillips, MAD/PH/680 (1991).
28. Z. Kunszt and F. Zwirner, in *Proceedings of the Large Hadron Collider Workshop*, Vol. II, edited by G. Jarlskog and D. Rein, CERN 90-10 (1990), p. 578; CERN-TH.6150/91 and ETH-TH/91-7 (1991).
29. H. Baer, M. Bisset, C. Kao and X. Tata, FSU-HEP-911104 (1991).
30. R. Barbieri, M. Frigeni, F. Giuliani, and H.E. Haber, *Nucl. Phys.* **B341** (1990) 309.
31. M.E. Peskin and T. Takeuchi, *Phys. Rev. Lett.* **65** (1990) 964; B. Holdom and J. Terning, *Phys. Lett.* **B247** (1990) 88; M. Golden and L. Randall, *Nucl. Phys.* **B361** (1991) 3.
32. H.E. Haber, preprint in preparation.
33. See *e.g.*, G.F. Giudice and G. Ridolfi, *Z. Phys.* **C41** (1988) 447; M. Olechowski and S. Pokorski, *Phys. Lett.* **B214** (1988) 393; M. Drees and M.M. Nojiri, *Nucl. Phys.* **B369** (1992) 54.
34. H.E. Haber and F. Zwirner, unpublished.
35. G.M. Asatryan, A.N. Ioannisyan and S.G. Matinyan, *Sov. J. Nucl. Phys.* **53** (1991) 371 [*Yad. Fiz.* **53** (1991) 592].
36. B. Kniehl, *Nucl. Phys.* **B352** (1991) 1; **B357** (1991) 439; DESY-91-128 (1991); DESY-92-021 (1992).
37. H.E. Haber and Y. Nir, *Nucl. Phys.* **B335** (1990) 363.

38. T.L. Barklow, SLAC-PUB-3564 (1990), to appear in *The Proceedings of the 1990 DPF Summer Study on High Energy Physics*, Snowmass, CO, June 25-July 13, 1990.
39. D.L. Borden, D.A. Bauer, and D.O. Caldwell, SLAC-PUB-5715 (1992).
40. J.F. Gunion and H.E. Haber, SCIPP-90/22 (1990), to appear in *The Proceedings of the 1990 DPF Summer Study on High Energy Physics*, Snowmass, CO, June 25-July 13, 1990.
41. M. Dine, P. Huet, R. Singleton and L. Susskind, *Phys. Lett.* **B257** (1991) 351.
42. M. Dine, P. Huet, R. Leigh and A. Linde, SLAC-PUB-5741 (1992); SLAC-PUB-5740 (1992).
43. M. Dine, A. Kagan and S. Samuel, *Phys. Lett.* **B243** (1990) 250.
44. M. Dine and D. MacIntire, SCIPP-91/16(1992).
45. L. Hall and L. Randall, *Nucl. Phys.* **B352** (1991) 289.
46. M. Kamiankowski and J. March-Russel, Institute for Advanced Study preprint IASSNS-HEP-92-9 (1992); R. Holman, S. Hsu, T. Kephart, E. Kolb, R. Watkins and L. Widrow NSF-ITP-92-06 (1992); S.M. Barr and D. Seckel, Bartol preprint BA-91-11 (1992).
47. M. Dine, Invited talk presented at the Cincinnati Symposium in Honor of Louis Witten, SCIPP preprint in preparation.
48. T. Banks and M. Dine, *Phys. Rev.* **D45** (1992) 1424.
49. T. Banks, M. Dine and N. Seiberg, *Phys. Lett.* **B273** (1991) 105.
50. L. Krauss and F. Wilczek, *Phys. Rev. Lett.* **62** (1989) 1221.
51. L. Ibanez and G. Ross, CERN preprint CERN-TH-6000 (1991).
52. V. Kuzmin, V. Rubakov and M. Shaposhnikov, *Phys. Lett.* **155B** (1985) 36.
53. S. Dimopoulos and L. Susskind, *Phys. Rev.* **D18** (1978) 4500.
54. N. Manton, *Phys. Rev.* **D28** (1983) 2019; F. Klinkhammer and N. Manton, *Phys. Rev.* **D30** (1984) 2212.
55. J. Harvey and M. Turner, *Phys. Rev.* **D42** (1990) 3344; S. Barr and A. Nelson, *Phys. Lett.* **B246** (1990) 141; W. Fischler, R. Giudice, R. Leigh and S. Paban, *Phys. Lett.* **B258** (1991) 45.
56. J. Ellis, R.A. Flores, S. Rudaz and D. Seckel, *Phys. Lett.* **194B** (1987) 241.
57. P. Arnold and L. McLerran, *Phys. Rev.* **D36** (1987) 581.
58. S. Yu Khlebnikov and M.E. Shaposhnikov, *Nucl. Phys.* **B308** (1988) 885.

59. E. Mottola and A. Wipf, *Phys. Rev.* **D39** (198) 588.
60. A.G. Cohen, M.J. Dugan, A.V. Manohar, *Phys. Lett.* **B222** (1989) 91.
61. M. Dine, W. Fischler, O. Lechtenfeld, J. Polchinski, and B. Sakita, *Nucl. Phys.* **B342** (1990) 381.
62. V.A. Rubakov, *JETP Lett.* **41** (1988) 266.
63. D. Yu Grigoriev and V.A. Rubakov, *Nucl. Phys.* **B299** (1988) 67.
64. A. Ringwald, *Nucl. Phys.* **B330** (1990) 1.
65. O. Espinosa, *Nucl. Phys.* **B343** (1990) 310.
66. L. McLerran, A. Vainshtein and M. Voloshin, *Phys. Rev.* **D42** (1990) 171.
67. T. Banks, C.M. Bender and T.T. Wu, *Phys. Rev.* **D8** (1973) 3346; T. Banks and C.M. Bender, *Phys. Rev.* **D8** (1973) 3366.
68. J.-L. Gervais and B. Sakita, *Phys. Rev.* **D16** (1977) 3507; H.J. de Vega, J. L. Gervais and B. Sakita, *Phys. Rev.* **D19** (1979) 604.
69. T. Banks, M. Dine, G. Farrar, D. Karabali and B. Sakita, *Nucl. Phys.* **B347** (1990) 581.
70. I. Affleck and M. Dine, *Nucl. Phys.* **B249** (1985) 361.
71. M.E. Shapohnikov, *Nucl. Phys.* **B299** (1988) 797.
72. L. McLerran, *Phys. Rev. Lett.* **62** (1075) 1989.
73. L. Turok and J. Zadrozny, *Phys. Rev. Lett.* **65** (1990).
74. L. McLerran, M. Shaposhnikov, N. Turok and M. Voloshin, *Phys. Lett.* **B256** (1991) 451.
75. M. Dine, P. Huet and R. Singleton, SCIPP preprint SCIPP-91/08 (1991).
76. A. Cohen, D. Kaplan, and A. Nelson, *Phys. Lett.* **B245** (1990) 561; *Nucl. Phys.* **B349** (1991) 727; *Phys. Lett.* **bf B263** (1991) 86; UCSD preprint UCSD-PTH-91-20 (1991).
77. M. Dine, SCIPP preprint in preparation, to appear in the Proceedings of the TEXAS Workshop on Electroweak Baryon Number Violation, Yale University, 1992.
78. D. Grigoriev, M. Shaposhnikov and N. Turok, Princeton preprint PUPT-91-1275 (1991).
79. D.A. Kirshnitz and A.D. Linde, *Ann. Phys.* **101** (1976) 195.
80. D. Brahm and S. Hsu, Harvard preprint HUTP-91-A064 (1991); Caltech preprint CALT-68-1705 (1991).
81. M. Shaposhnikov, CERN preprint CERN-TH-6319-91 (1991).

82. M. Gleiser and R. Kolb, Fermilab preprint FERMILAB-PUP-91-305-A (1991); N. Tetradis, DESY preprint DESY-91-151 (1991).
83. N. Turok, Princeton University preprint (1991).
84. B. Liu, L. McLerran, N. Turok, Minnesota preprint to appear.
85. M. Dine and W. Fischler, *Nucl. Phys.* **B204** (1982) 346.
86. I. Affleck, M. Dine and N. Seiberg, *Nucl. Phys.* **B256** (1985) 557.
87. E. Cremmer, B. Julia, J. Scherk, S. Ferrara, L. Girardello and P. van Nieuwenhuizen, *Nucl. Phys.* **B147** (1979) 105.
88. J. Polchinski and L. Susskind, *Phys. Rev.* **D26** (1982) 3661.
89. I. Antoniadis and C. Kounnas, *Phys. Lett.* **B261** (1991) 369.
90. T. Banks, M. Dine, M. Dykstra and W. Fischler, *Phys. Lett.* **212B** (1988) 45.
91. L. Dixon, V. Kaplunovsky, J. Louis and M. Peskin, SLAC preprint to appear.
92. M. Dine and N. Seiberg, *Phys. Lett.* **B162** (1985) 299.
93. M. Dine, R. Leigh and D. MacIntire, SCIPP preprint 92/16 (1992).
94. K. Choi, D. Kaplan and A. Nelson, UCSD preprint PTH 92-11 (1992).
95. J. Dai, H. Dykstra, R.G. Leigh, S. Paban and D. Dicus, *Phys. Lett.* **B237** (1990) 216; **B242** (1990) 547.
96. R.G. Leigh, Ph.D. thesis, University of Texas at Austin, Aug. 1991.
97. R.G. Leigh, S. Paban, and R.-M. Xu, *Nucl. Phys.* **B352** (1991) 45.
98. W. Fischler, G.F. Giudice, R.G. Leigh and S. Paban, *Phys. Lett.* **258B** (1991) 45.
99. W. Fischler, G. Giudice, R.G. Leigh and S. Paban, in Proceedings of "Quantum Mechanics of Fundamental Systems III", Santiago, Chile, January, 1990.
100. W. Fischler, G. Giudice, R.G. Leigh, S. Paban and S. Thomas *Phys. Lett.* **266B** (1991) 408.
101. R.G. Leigh, to appear in the Proceedings of the TEXAS Workshop on Electroweak Baryon Number Violation, Yale University, 1992.
102. A. Méndez and A. Pomarol, *Phys. Lett.* **B265** (1991) 177.
103. A. Méndez and A. Pomarol, *Phys. Lett.* **B279** (1992) 98.
104. A. Méndez and A. Pomarol, *Phys. Lett.* **B272** (1991) 313.
105. A. Pomarol, Ph.D. thesis, Universitat Autònoma de Barcelona, June 1991.

VI. List of Publications: July, 1991–June, 1992

A. Howard Haber

Howard E. Haber

1. Heavy Fermion Effects in $e^+e^- \rightarrow ZH$ and $Z \rightarrow H\nu\bar{\nu}$, S. Dawson and H.E. Haber, *Phys. Rev. D* **D44** (1991) 53.
2. $H^\pm \rightarrow W^\pm\gamma$ and $H^\pm \rightarrow W^\pm Z$ in Two Higgs Doublet Models: Large Fermion Mass Limit, M. Capdequi Peyranere, H.E. Haber and P. Irulegui, *Phys. Rev. D* **D44** (1991) 191.
3. Lectures on Electroweak Symmetry Breaking, H.E. Haber, in *Testing the Standard Model*, Proceedings of the 1990 Theoretical Advanced Study Institute in Elementary Particle Physics, edited by M. Cvetič and Paul Langacker (World Scientific, Singapore, 1991) pp. 340–475.
4. Higgs Physics: Theory and Phenomenology, H.E. Haber, in *The Standard Model and Beyond*, Proceedings of the 1990 Mt. Sorak Symposium on Theoretical Physics, edited by J.E. Kim, (World Scientific, Singapore, 1991) pp. 28–104.
5. Higgs Boson Low-Energy Theorems and Their Applications, S. Dawson and H.E. Haber, *Int. J. Mod. Phys. A* **7** (1992) 107.
6. One Loop Radiative Corrections to the Charged Higgs Mass of the Minimal Supersymmetric Model, by M.A. Díaz and H.E. Haber, SCIPP-91/14 (1991), *Phys. Rev. D* **D45** (1992), in press.
7. Searching for CP-Even Higgs Bosons of the Minimal Supersymmetric Model at Hadron Supercolliders, by J.F. Gunion, R. Bork, H.E. Haber, and A. Seiden, UCD-91-29 and SCIPP-91/34 (1991), *Phys. Rev. D*, in press.
8. Searching for the CP-Odd Higgs Boson of the Minimal Supersymmetric Model at Hadron Supercolliders, by J.F. Gunion, H.E. Haber, and C. Kao, UCD-91-32 and SCIPP-91/44 (1991), *Phys. Rev. D*, in press.
9. The Decay $h^0 \rightarrow A^0 A^0$ in the Minimal Supersymmetric Model, by H.E. Haber, R. Hempfling, and Y. Nir, SCIPP-91/40 and WIS-92/5/Feb-PH (1992), submitted to *Phys. Rev. D*.
10. Can the Higgs Mass be Entirely due to Radiative Corrections?, by M.A. Díaz and H.E. Haber, SCIPP-91/13 (1992), submitted to *Phys. Rev. D*.
11. The Renormalization-Group Improved Higgs Sector of the Minimal Supersymmetric Model, by H.E. Haber and R. Hempfling, SCIPP-91/33 (1992), submitted to *Phys. Rev. D*.

12. Higgs Theory and Phenomenology at Future e^+e^- Linear Colliders, by H.E. Haber, SCIPP-92/10 (1992), to appear in *Proceedings of the Workshop on Physics and Experiments with Linear Colliders*, Saariselkä, Finland, 9–14 September, 1991.
13. The Higgs Sector in the Minimal Supersymmetric Model: Radiative Corrections and Their Implications, by H.E. Haber, SCIPP-92/11 (1992), to appear in *Proceedings of the International Workshop on Electroweak Symmetry Breaking*, Hiroshima, Japan, 12–15 November, 1991.

B. Michael Dine

1. Creating the Baryon Asymmetry at the Electroweak Transition (with P. Huet, R. Singleton and L. Susskind) *Phys. Lett.* **B257** (1991) 351.
2. Baryogenesis at the Electroweak Scale (with P. Huet and R. Singleton, Santa Cruz preprint SCIPP-91-08 (1991) to appear in *Nucl. Phys. B*.
3. Electroweak Baryogenesis, in *Proceedings of the Vancouver Meeting, Particles and Fields '91*, edited by D. Axen, D. Bryman and M. Comyn (World Scientific, Singapore, 1992), p. 831.
4. Irrational Axions as a Solution of the Strong CP Problem in an Eternal Universe (with T. Banks and N. Seiberg), *Phys. Lett.* **B273** (1991) 105.
5. Note on Discrete Gauge Anomalies (with T. Banks), *Phys. Rev.* **D45** (1992) 1424.
6. Comments on the Electroweak Phase Transition (with R. Leigh, P. Huet, A. Linde and D. Linde), SLAC-PUB-5470, to appear in *Phys. Lett. B*.
7. Towards the Theory of the Electroweak Phase Transition (with R. Leigh, P. Huet, A. Linde and D. Linde), SLAC-PUB-5741 to appear in *Phys. Rev. D*.
8. Supersymmetry, Naturalness and Dynamical Supersymmetry Breaking (with D. MacIntire), SCIPP preprint SCIPP-91/16 (1992).
9. Electroweak Baryogenesis: Where are we Now?, Invited Talk at the Texas Workshop on Electroweak Baryon Number Violation (Yale, 1992), SCIPP preprint in preparation.
10. Some Problems of Naturalness: Lessons from String Theory, Invited Talk at the Cincinnati Symposium in Honor of Louis Witten, SCIPP preprint in preparation.
11. Of CP and other Gauge Symmetries in String Theory (with R. Leigh and Doug MacIntire), SCIPP preprint SCIPP-92/16 (1992).

C. Robert Leigh

1. Electric Dipole Moment of the Electron (with S. Paban and R.-M. Xu), *Nucl. Phys.* **B352** (1991) 45.
2. Constraints on the Baryogenesis Scale from Neutrino Masses (with W. Fischler, G.F. Giudice, R.G. Leigh and S. Paban), *Phys. Lett.* **258B** (1991) 45.
3. The Invisible Majoron (with W. Fischler, G. Giudice, S. Paban and S. Thomas) *Phys. Lett.* **266B** (1991) 408.
4. Comments on the Electroweak Phase Transition (with M. Dine, P. Huet, A. Linde and D. Linde), SLAC-PUB-5470, to appear in *Phys. Lett. B*.
5. Towards the Theory of the Electroweak Phase Transition (with M. Dine, P. Huet, A. Linde and D. Linde), SLAC-PUB-5741 to appear in *Phys. Rev. D*.
6. Bubble Propagation at the Electroweak Phase Transition, in Proceedings of the TEXAS Workshop on Electroweak Baryon Number Violation, Yale University, 1992.
7. Of CP and other Gauge Symmetries in String Theory, SCIPP preprint SCIPP-92/16 (1992).

D. Alex Pomarol

1. t-quark loop corrections to the charged Higgs boson hadronic decay, by A. Méndez and A. Pomarol, *Phys. Lett.* **B265** (1991) 177.
2. t-quark loop corrections to the neutral Higgs couplings in the two-Higgs-doublet model, by A. Méndez and A. Pomarol, *Phys. Lett.* **B279** (1992) 98.
3. Signals of CP violation in the Higgs sector, by A. Méndez and A. Pomarol, *Phys. Lett.* **B272** (1991) 313.

*Budget info.
removed
pp 36-38*

END

**DATE
FILMED**

12 / 23 / 92