Wetland Vegetation Establishment in L-Lake

by

Steven R. Kroeger
Savannah River Ecology Laboratory

July, 1990

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

for

EXECUTIVE SUMMARY

Wetland vegetation was transplanted from PAR Pond to L-Lake between January and August, 1987. Approximately 100,000 individual plants representing over 40 species were transplanted along the southern shoreline. Three zones of vegetation were created: 1) submersed/floating-leaved, 2) emergent, 3) upper emergent/shrub. During the summers of 1987, 1988, 1989, the Savannah River Ecology Laboratory sampled the vegetation in 54 permanent transects located in planted (N=32) and unplanted areas (N=22). The 1989 vegetation data from L-Lake were compared to 1985 data from PAR Pond.

Submersed and Floating-leaved Vegetation

Nine species of submersed and floating-leaved plants were transplanted into this zone. American lotus (*Nelumbo lutea*), a floating-leaved species, and water celery (*Vallisneria americana*), a submersed plant, were the surviving species in 1989. Wave action and low initial planting apparently made establishment of some species difficult.

Colonization has occurred rapidly in this zone. In 1987, 95% of the plots sampled had no vegetation present. By 1989, the percentage of empty plots had declined to 62%, due to expansion of water celery and American lotus. In 1987, mean cover per plot was 1%, however this increased to 22% by 1989. No submersed or floating-leaved vegetation was found in the unplanted areas.

Comparing the vegetation of L-Lake with PAR Pond shows that there are few similarities. American lotus is well established in both reservoirs but PAR Pond has a greater species diversity. Water celery is well established in L-Lake but was not sampled in PAR Pond. Many emergent species occur in the deeper water in PAR Pond and these

same species are expected to occur in the deeper waters of L-Lake given time. Cattail (*Typha latifolia*) already is established in L-Lake in water greater than 30 cm (1 ft).

Emergent Vegetation

Most species were successfully transplanted in this zone. In 1987, 32% of the plots sampled in planted areas were without vegetation, but by 1989 this had declined to 16%. Mean cover per plot increased from 22% in 1987 to 40% in 1988 and remained the same for 1989. Emergent vegetation is becoming established slowly in unplanted shoreline areas. Alligator weed (*Alternanthera philoxeroides*), water pennywort (*Hydrocotyle umbellata*) and cattail were the only emergent species in unplanted areas with absolute frequencies greater than 5% in 1989. The frequency of empty plots remained relatively constant since 1987, approximately 85% per year.

The emergent vegetation of L-Lake is similar in composition to the emergent vegetation of PAR Pond. The *Panicum/Sacciolepis* (*P. hemitomon* and *S. striata*) grasses were the most important component of the vegetation in both sites. Water celery was important in L-Lake but not sampled in PAR Pond.

Upper Emergent/Shrub Vegetation

All species planted in 1987 were present in 1989. Willow (*Salix nigra*) and the *Panicum/Sacciolepis* grasses were the most important species in this vegetation zone. This zone was sufficiently vegetated with terrestrial species so that the proportion of barren plots was always very low. Mean cover per plot in planted areas increased from 59% in 1987 to 69% in 1988 and then decreased to 55% in 1989. Damage to vegetation

by feral pigs was noted in 1989, but additional unknown factors may have contributed to the decrease in cover. Willow was the dominate shrub.

Conclusions

1.) American lotus and water celery are well established along some shoreline areas where it was planted.

2.) Additional time is necessary for the vegetation in water greater than 30 cm (1 ft) deep to develop and become similar in composition to the vegetation in PAR Pond.

3.) No vegetation in water greater than 30 cm deep is present along areas of shoreline which were not planted.

4.) Emergent vegetation is well established only along areas of shoreline where it was planted.

5.) Emergent vegetation is becoming established slowly along areas of shoreline which were not planted.

6.) Emergent vegetation of L-Lake in planted areas is similar to emergent vegetation in PAR Pond.

7.) Willow is the dominate shrub.

TABLE of CONTENTS

Executive Summary	i
Introduction	1
Methods	4
Planting.....	4
Vegetation Sampling.....	7
Data Analysis.....	10
Vegetation Sampling at PAR Pond.....	11
Results and Discussion	11
Submersed and Floating-leaved Vegetation.....	11
Transplant Survival.....	12
Changes from 1987 to 1989 in Planted Areas.....	12
Comparison Between Planted and Unplanted Areas.....	16
Comparison with Submersed/Floating-leaved Vegetation at PAR Pond.....	16
Emergent Vegetation.....	19
Transplant Survival.....	19
Changes from 1987 to 1989 in Planted Areas.....	19
Comparison Between Planted and Unplanted Areas.....	22
Comparison with Emergent Vegetation at PAR Pond.....	24
Upper Emergent/Shrub Vegetation.....	24
Transplant Survival.....	26
Changes from 1987 to 1989 in Planted Areas.....	26
Comparison Between Planted and Unplanted Areas.....	28
Summary	28
References	31
 APPENDIX A. Maps of Shoreline Vegetation	32
Species codes used on the topographic maps.....	33
APPENDIX B. Vegetation sampled during 1987, 1988, and 1989	45

INTRODUCTION

The Department of Energy's (DOE) decision to restart L-Reactor resulted in the construction of L-Lake, a 400 ha (1,000 acre) once-through cooling reservoir to cool thermal effluents before entry into Steel Creek. To obtain necessary federal and state permits for thermal effluent discharge into the lake, DOE was required to meet requirements mandated by Section 316b of the Water Pollution Control Act Amendments of 1972. One requirement was the establishment of a balanced biological community (BBC). A primary component of a BBC is vegetation because it forms the base for aquatic food chains, provides habitat for animals and functions in nutrient cycling.

To augment natural establishment of a wetland plant community, DOE requested that the University of Georgia's Savannah River Ecology Laboratory (SREL) implement planting of wetland vegetation along the shores of L-Lake. A panel of experts in the areas of wetland ecology and restoration was invited to the Savannah River Site and developed a management plan for establishing an appropriate wetland plant community (Whigham et al. 1985). In addition, the panel recommended that plants for L-Lake be transplanted from PAR Pond, a recirculating cooling reservoir.

The panel recommended that plants for L-Lake be transplanted from PAR Pond for the following reasons. First, its location near L-Lake facilitated logistics. Second, its naturally established vegetation, portions of which have been exposed to thermal effluents for 30 years, may be more thermally tolerant than plants from natural sources. Finally, species found in PAR Pond are representative of natural wetland species in the region.

L-Lake was completed in October, 1985. The Savannah River Ecology Laboratory requested competitive bids from private contractors to implement the transplanting project in 1985. A topographic survey of the shoreline of the southern half of the lake was made in 1986 to aid the contractor in transplanting.

Southern Tier Consulting, a firm specializing in the establishment of wetland plant communities, was selected for the project. Between January and August, 1987, approximately 4200 meters (14,000 ft) of L-Lake shoreline were planted with 100,000 plants representing over 40 species.

The panel also proposed that zones of vegetation be established to represent species patterns found in PAR Pond and natural lakes (Whigham et al. 1985). These zones are differentiated by species composition and may be controlled by water depth. Three zones of vegetation were established (Fig 1.). Trees, shrubs and some emergents formed the upper emergent/shrub zone. This zone lies above the waterline, but can flood periodically. The emergent zone consists of erect species (e.g. cattail) which are present mostly in shallow water. The third zone consists of floating-leaved (e.g. water lilies) and submersed species, and occurs in deeper water.

In August 1987, SREL sampled the shoreline vegetation of L-Lake to assess the survival of transplants and the rate of colonization of new areas. Vegetation was also sampled during the summers of 1988 and 1989 along permanent transects located in both planted and unplanted areas of shoreline. This report focuses on 1) plant survival since 1987, 2) changes in community composition between 1987 and 1989 in planted areas, 3) differences in vegetation structure between planted and unplanted areas in 1989, and 4)

FIGURE 1. Plant zonation at L-Lake.

a comparison of the vegetation in the submersed/floating-leaved and emergent zones of L-Lake and PAR Pond.

METHODS

Planting

The management plan for establishment of littoral/wetland vegetation on the L-Lake shoreline (Whigham et al. 1985) was used as a guide for planting. The (spatially) continuous planting option was adopted; this option recommended that all available portions of the southern half of L-Lake be planted. The northern half of the lake was not planted because it is thermally impacted and survival of the transplants was expected to be low. About 10% of the southern shoreline was not planted because it had prior designated use for regulatory compliance monitoring (i.e. fish electroshocking and vegetation monitoring).

PAR Pond and commercial nurseries were used as sources of plant material. Perennial herbaceous plants were excavated by hand from PAR Pond, but trees, one emergent herb (*Sagittaria latifolia*) and seed of some grasses were obtained from commercial sources. Species planted are listed in Table 1. Maps of planting locations are in Appendix A. Each individual was planted with a ten gram tablet of timed-release Agriform fertilizer (20-10-5). Planting occurred from January through August, 1987. Emergent plants were spaced at 60 cm (2 ft) centers and floating-leaved species at 1.2 m (4 ft) centers. Most emergents were planted in blocks containing one species. A block was a rectangular area 15 m (50 ft) long and the width varied according to the slope of the shoreline. Seed densities for grasses are not known since they were spread by hand.

TABLE 1. Species and quantities planted at L-Lake by Southern Tier Consulting between January and August, 1987. (*Planted as seed.)

<u>Scientific name</u>	<u>Common name</u>	<u>Quantity planted</u>
Submersed/floating-leaved zone		
<i>Brasenia schreberi</i>	Water shield	<1000
<i>Eleocharis acicularis</i>	Spike rush	<2000
<i>Najas gracillima</i>	Bushy pondweed	<100
<i>Nelumbo lutea</i>	American lotus	<1000
<i>Nymphaea odorata</i>	White waterlily	>2000
<i>Nymphoides aquatica</i>	Floating heart	<100
<i>Potamogeton pulcher</i>	Pondweed	<1000
<i>Potamogeton vaseyi</i>	Pondweed	<100
<i>Vallisneria americana</i>	Water celery	>2000
Emergent zone		
<i>Axonopus</i> sp.	Carpet grass	10 lbs *
<i>Bacopa caroliniana</i>	Bacopa	>2000
<i>Carex comosa</i>	Sedge	<2000
<i>Carex glaucescens</i>	Sedge	<100
<i>Dulichium arundinaceum</i>	Three-way sedge	<100
<i>Echinochloa crusgalli</i>	Wild millet	25 lbs *
<i>Echinodorus cordifolius</i>	Burhead	<100
<i>Eleocharis equisetoides</i>	Spike rush	<2000
<i>Eleocharis quadrangulata</i>	Spike rush	<2000
<i>Erianthus giganteus</i>	Beard grass	<100
<i>Glyceria striata</i>	Manna grass	<100
<i>Hydrochloa caroliniensis</i>	Grass	<1000

TABLE 1 continued.

<u>Scientific name</u>	<u>Common name</u>	<u>Quantity planted</u>
<i>Hydrocotyle umbellata</i>	Water pennywort	< 100
<i>Juncus acuminatus</i>	Rush	< 100
<i>Juncus brachycarpus</i>	Rush	< 100
<i>Juncus effusus</i>	Soft rush	> 2000
<i>Juncus diffusissimus</i>	Rush	< 100
<i>Leerisa oryzoides</i>	Rice cutgrass	> 2000
<i>Lycopus rubellus</i>	Water horehound	< 1000
<i>Panicum hemitomon</i>	Panic grass	> 2000
<i>Panicum virgatum</i>	Switchgrass	10 lbs*
<i>Paspalum distichum</i>	Knotgrass	< 1000
<i>Polygonum sp.</i>	Smartweed	< 2000
<i>Pontederia cordata</i>	Pickereelweed	< 1000
<i>Sagittaria latifolia</i>	Arrowhead	2000
<i>Scirpus cyperinus</i>	Bulrush	< 2000
<i>Sparganium americanum</i>	Bur reed	< 100
<i>Typha domingensis</i>	Cattail	< 1000
<i>Typha latifolia</i>	Cattail	> 2000
Upper emergent/shrub zone		
<i>Acer rubrum</i>	Red maple	> 2000
<i>Cephalanthus occidentalis</i>	Buttonbush	> 2000
<i>Mikania scandens</i>	Climbing hempweed	< 100
<i>Nyssa sylvatica</i>	Blackgum	> 2000
<i>Salix nigra</i>	Black willow	> 2000
<i>Taxodium distichum</i>	Cypress	> 2000

Along some areas of shoreline individuals of *Nelumbo lutea* were spaced nine meters (30 ft) apart because of limited quantities. *Vallisneria americana*, a submersed species, was planted by weighting a plant and a fertilizer tablet in a cheesecloth bag and tossing the bags from a boat.

Once plants were introduced into L-Lake it was expected that they would reproduce and fill unoccupied areas. Plant reproduction could occur sexually through seed production and dispersal, or asexually by the separation of a plant into a fragment capable of reforming a whole plant or by spreading from the roots. Asexual reproduction is also called vegetative reproduction and occurs commonly.

The presence of the human pathogenic amoeba, *Naegleria fowleri* in L-Lake made working in water hazardous. This amoeba may cause the fatal disease Primary Amoebic Meningoencephalitis which results if the amoeba migrates up the nasal passages, through the olfactory nerve, and into the brain. Therefore, floating-leaved and submersed species could not be transplanted safely to water more than knee deep. Only the shallow areas of the submersed/floating-leaved zone were planted.

Vegetation Sampling

Permanent belt transects, 2 m wide, were established in a stratified random pattern, perpendicular to the shore of L-Lake. Fifty-four transects were sampled, thirty-two in planted areas and twenty-two in unplanted ones (Fig. 2). Species presence was recorded and percent cover was estimated in paired one square meter sample plots beginning at an elevation approximately 30 cm (1 ft) above the waterline and continuing at one meter intervals to a water depth of one meter. Trees greater than 2.5 cm (1 in) diameter at the

FIGURE 2. Location of permanent sample transects. Transects marked with a dot (•) are located in unplanted areas.

base were not sampled because the sampling design employed here was appropriate only for herbaceous plants and woody seedlings. Since the wetland vegetation in L-Lake was in an early stage of development, there were many sample plots with no plants. This absence of vegetation was recorded and was especially common in the submersed/floating-leaved zone.

Cover was estimated as the percentage of ground surface covered by vegetation when looking down from above the vegetation. Cover represents the vertical projection of plant parts (leaves and stems) onto the ground (Bonham 1989) and approximates the area over which a species exerts influence. Cover for a species may range from 1 to 100% in each plot. Because of overlap, total cover for all species in any plot may be greater than 100%. Small individuals and seedlings were assigned a cover value of one percent.

Taxonomonic nomenclature follows that of Radford, Ahles and Bell (1968). In some cases, the lack of flowers and fruits prevented identification of a plant to the species level. Since *Panicum hemitomom* and *Sacciolepis striata* share the same habitat and are difficult to differentiate vegetatively, these grasses were combined into one group called *Panicum/Sacciolepis*.

The elevation of each transect was surveyed at one meter intervals from approximately 40 cm above the waterline to one meter water depth. These elevational data, based upon the normal high water mark, were used to demarcate the three vegetation zones. The normal high water mark was determined by measuring the elevation of water marks on rocks and on a pier.

Data Analysis

Measurements of how often a species occurred (absolute frequency) and how much area it influenced (absolute cover) were obtained during sampling. Absolute frequency and cover were converted to relative frequency and relative cover and expressed as a percentage. The relative frequency of a species is the absolute frequency of that species divided by the sum of the absolute frequencies of all species. Likewise, the relative cover of a species is the ratio of its coverage to the total coverage for all species.

Both absolute and relative values are helpful in interpreting plant community data. For example, if four species were found in each of ten sample plots, then the absolute frequency for each species is 100%, whereas the relative frequency for each species is 25%. The first value expresses the probability of occurrence in a sample plot, and the second expresses the probability of occurrence among the species sampled. Cover and frequency for a species may occur in various combinations. For example, a species may occur in many plots but have low cover, or in few plots and have high cover. Integrating frequency and cover is achieved by summing the values for relative frequency and relative cover. This sum is commonly called an importance value and, in the results presented here, has a maximum value of 200. A species with a high importance value is not necessarily an ecologically important one, but it is notable given the sample data.

Importance values were used both to compare vegetation in the planted areas among the three years of study and to compare vegetation structure and development between planted and unplanted areas. This latter comparison was made between categories of species such as submersed, floating-leaved, emergent, terrestrial or shrub. Wetland species

which became established naturally were assigned to the appropriate category using Reed (1986). A terrestrial species was defined as an herbaceous plant not listed in Reed (1986). Appendix B provides the number sampled and coverage for all species by zone in both planted and unplanted areas of L-Lake.

Vegetation Sampling at PAR Pond

Wetland vegetation of PAR Pond has become established naturally since the lake was constructed in 1958. There are well developed zones of vegetation that served as the model for the vegetation structure of L-Lake. Data on the structure and composition of the emergent vegetation were collected in 1985 using belt transects one meter wide (Harrison, unpublished); thirty-two transects were sampled, sixteen in the north arm and 16 in the cold arm. Species presence was noted and species cover was estimated in one square meter sample plots at one meter intervals along these transects from the waterline to a point in the water where the emergent vegetation ended. The focus of the PAR Pond study was on emergent vegetation, therefore much of the submersed/floating-leaved zone was neglected. Since water depth was measured the vegetation data was analyzed in the same way as the L-Lake data. This allowed comparisons of the vegetation between L-Lake and PAR Pond to be made.

RESULTS and DISCUSSION

Submersed and Floating-leaved Vegetation

The shoreline between 30 cm (1 ft) and 100 cm (3.3 ft) water depth defines this zone of submersed and floating-leaved vegetation. Nine species of submersed and floating-leaved plants were planted (Table 1). A total of 940 plots in planted areas and 786 in

unplanted areas were sampled each year (Table 2).

Transplant Survival

Of the nine species planted, American lotus (*Nelumbo lutea*), a floating-leaved species, and water celery (*Vallisneria americana*), a submersed plant, survived in 1989. Deep water and the presence of the pathogenic amoeba, *N. fowleri*, made this an unsafe zone to plant and sample. Wave action, combined with the initial planting of fewer than 100 individuals of some species (Table 1), apparently made establishment of some species difficult.

Few studies have addressed the establishment of transplanted submersed and floating-leaved species. Kollar (1985) had no success transplanting the submersed species *Potamogeton pectinatus* and *P. perfoliatus* in the Chesapeake Bay. He suggested that high turbidity and algal epiphytes may have prevented these species from becoming established. L-Lake has high turbidity and algal blooms during the summer that may have prevented successful transplantation of some species. Kollar (1985) successfully transplanted *V. americana* in the Chesapeake Bay, and it is also becoming well established in L-Lake. This species and *N. lutea* are colonizing unoccupied areas in the submersed zone of L-Lake quickly (Fig. 3).

Changes from 1987 to 1989 in Planted Areas

Colonization is occurring rapidly in this zone. In 1987, 95% of the plots sampled had no vegetation present. By 1989, the percentage of empty plots had declined to 62% (Table 2). Much of this reduction in the number of empty plots was due to expansion of *V. americana* and *N. lutea*. Concomitant with the colonization of empty plots has been a

TABLE 2. Comparison of frequency and cover estimates during 1987-1989.

Site	N ¹	Species occurrences ²		Empty plots (%)		Mean cover/plot				
		1987	1988	1989	1987	1988	1989	1987	1988	1989
Submersed/floating-leaved zone										
Planted	940	75	177	401	94.8	82.9	62.3	1.0	7.8	21.8
Unplanted	786	0	1	6	100.0	99.9	99.4	0.0	<0.1	<0.1
Emergent zone										
Planted	340	379	576	825	32.1	22.3	15.6	21.9	39.7	39.5
Unplanted	216	54	53	85	85.2	88.9	83.8	3.5	4.3	2.4
Upper emergent/shrub zone										
Planted	196	1001	1878	1784	0.0	0.0	1.0	58.8	68.7	55.1
Unplanted	118	364	769	910	18.6	8.5	5.1	39.3	60.5	49.3

¹Total number of plots sampled.

²Sum of the number of species (not individuals) in each plot.

FIGURE 3. Importance values for species planted in the submersed/floating-leaved zone.

substantial increase in plant cover. In 1987 mean cover per plot was 1%, however, this increased to 22% by 1989 (Table 2). Cattails (*Typha latifolia*) are expanding into this zone from the emergent zone (Fig 3).

Vallisneria americana is present on both the east and west shores of L-Lake and is found on clay and sand substrates. When transplanted, it soon drops its leaves and exists as a tuber in the sediment. This transplant response probably explains its apparent absence in 1987 (Fig. 3). In 1989, *V. americana* was present in 8.8% of the plots sampled and its frequency may have been underestimated due to lake turbidity and water depths of nearly one meter. Only vegetative reproduction has been observed.

The establishment of *Vallisneria americana* (a major winter food for waterfowl; Jones and Drobney 1986) suggests that L-Lake may become a feeding area for wintering waterfowl. Although it is a common plant nationally, it is uncommon in South Carolina. On the Savannah River Site, it is found only in PAR Pond and L-Lake (Knox and Sharitz 1990).

Nelumbo lutea is also a quick colonizer. In 1987, it occurred in only 3.9% of sampled plots, but by 1989 its absolute frequency had increased to 26.7%. Radial growth of 14 meters (45 ft) in one growing season has been reported for patches in Tennessee Valley Authority reservoirs (Hall and Penfound 1944). Although radial growth of *N. lutea* has not been measured in L-Lake it has occurred very rapidly. It is reproducing vegetatively and is present only along portions of the eastern shoreline. It did not become established along the western shoreline. Although it does produce seeds, there has been no evidence of seedling establishment.

During 1987, the white water lily, *Nymphaea odorata*, shared dominance with *N. lutea* (Fig. 3). In 1989, however, only one *N. odorata* individual was found in an unsampled and protected area in the northwestern portion of the lake. In PAR Pond, *N. odorata* and water shield (*Brasenia schreberi*) occur in protected areas such as coves. Water flow from reactor discharge may prevent these species from occurring near discharge points (Grace 1984). Thus the extirpation of *N. odorata* from planted areas in L-Lake may have been caused by wave action which uprooted or damaged the plants.

Comparison Between Planted and Unplanted Areas

No submersed or floating-leaved vegetation was present in unplanted sites in 1989, thus planting appears essential in promoting the establishment of this zone (Fig 4). Four years after lake construction was completed, nearly all plots in unplanted areas remained unvegetated (Table 2). Five plots contained cattail (*T. latifolia*) or water pennywort (*Hydrocotyle umbellata*). These two emergents accounted for 100% of the relative cover in the unplanted areas (Fig. 4).

Comparison with Submersed/Floating-leaved Vegetation at PAR Pond

There are few similarities between the vegetation of this zone in L-Lake with the vegetation in PAR Pond. *Nelumbo lutea* is common in both reservoirs (Fig. 5). PAR Pond has a greater species diversity, however water pennywort (*Bacopa caroliniana*), spike rush (*Eleocharis quadrangulata*), pickerelweed (*Pontederia cordata*) and cattail (*Typha latifolia*) are emergent species (Fig. 5). It will take some time before these species, present in the emergent zone of L-Lake, expand into the submersed/floating-leaved zone. Although present in PAR Pond, *V. americana* was not sampled.

FIGURE 4. A comparison of vegetation between planted and unplanted areas of L-Lake in the submersed/floating-leaved zone.

FIGURE 5. A comparison of vegetation between L-Lake and PAR Pond in the submersed/floating-leaved zone.

Emergent Vegetation

This zone lies between the waterline and 30 cm (1 ft) of water. Approximately 30 species were planted (Table 1). A total of 340 plots in planted areas and 216 plots in unplanted areas were sampled (Table 2).

Transplant Survival

Most species planted in this zone were still present in 1989. Spikerushes (*Eleocharis equisetoides* and *E. quadrangulata*), water pennywort (*H. umbellata*), soft rush (*Juncus effusus*), grasses, smartweeds (*Polygonum persicaria* and *P. hydropiperoides*) and cattail (*T. latifolia*) were the most common emergents present in 1989 (Fig. 6).

Emergent species that were initially planted but not recorded in any subsequent sampling year include *Dulichium arundinaceum*, *Glyceria striata*, *Juncus acuminatus*, *J. brachycarpus*, *J. diffusisimus* and *Typha domingensis*. Two of these species, *D. arundinaceum* and *T. domingensis*, have been observed along areas of shoreline outside the sample transects. The *Juncus* spp. may have failed to survive or may have been missed in the samples because of problems with identification.

Changes from 1987 to 1989 in Planted Areas

The number of species occurrences increased from 379 in 1987 to 825 in 1989. Accompanying this increase has been a decrease in the number of empty plots. In 1987, 32% of the 340 plots sampled were empty, but by 1989 the percentage declined to 16%. Mean cover per plot increased from 22% from 1987 to 40% in 1988 and remained similar for 1989 (Table 2). Noteworthy changes for emergent species included a gradual increase in relative frequency of *Eleocharis* spp. and *T. latifolia*, a large increase of *H. umbellata*

FIGURE 6. Importance values for species planted in the emergent zone.

and *V. americana* and a major decrease in relative frequency and cover of the shoreline grasses.

Water depth was the major factor separating two distinct groups of grasses. Forming one group were *Panicum hemitomon*, a planted species, and *Sacciolepis striata*, an unplanted species. Both species grew outward from the shore into shallow water. Because of similar growth patterns and the difficulty in distinguishing them, these species were considered as one group. All other planted grass species comprise the second group called shoreline grasses. They were growing at or near the shoreline and not out into the water. The lack of flowers prevented identification of many shoreline grasses to species.

Changes in grass composition between 1987 and 1989 include a large decrease in relative frequency and relative cover of the shoreline grasses and a gradual increase in these parameters for the *Panicum/Sacciolepis* group (Fig. 6). In 1987, relative cover of shoreline grasses was 27%, but by 1989 had decreased to 1.4%. Most of this decrease in relative cover resulted from a decline in the absolute cover from 20.2 m² in 1987 to 1.9 m² in 1989. A 46 cm (1.5 ft) drop in the water level of L-Lake during summer 1987 may have aided the establishment of shoreline grasses. The subsequent decrease in frequency and coverage for shoreline grasses was probably due to lake levels returning to normal.

Most emergent species are reproducing vegetatively, however reproduction by seed was observed of *H. umbellata*, *T. latifolia*, *J. effusus* and *Pontederia cordata*. During 1987, blocks of *P. cordata* occurred only where the species was planted. However, by 1989, all of these blocks had been destroyed by beavers or pigs. In 1989, individual plants, not

blocks, of *P. cordata* had become established along both east and west shores; one plant was found in an unplanted area.

Cattail (*T. latifolia*) was present in the lake before transplanting began, however most of the individuals sampled in planted areas were transplants. Cattails transplanted in winter did not survive as well as those transplanted during spring after the emergence of new growth. Ristich et al. (1976) found significant differences in survival of *T. angustifolia* planted at different times, with the highest survival rates occurring for individuals transplanted during late spring (May-June). Observations of *T. latifolia* at L-Lake are consistent with his results.

In 1987, cattail was present in 9% of the 340 plots sampled. By 1989, its absolute frequency had increased to 29%. Although cattail seedlings have been observed, most of its expansion was due to vegetative reproduction. It has spread both laterally along the shore and into the deeper water of the submersed and floating-leaved zone.

Nelumbo lutea and *V. americana* were also important components of this zone. These two species have grown into the emergent zone from the submersed and floating-leaved zone.

Comparison Between Planted and Unplanted Areas

Emergent vegetation is becoming established slowly in unplanted shoreline areas (Fig 7). Alligator weed (*Alternanthera philoxeroides*), *H. umbellata* and *T. latifolia* were the only emergent species in unplanted areas with absolute frequencies greater than 5% in 1989. The number of empty plots in unplanted areas has remained relatively constant since 1987 (Table 2). In 1989, 84% of the 216 plots sampled had no vegetation present.

FIGURE 7. A comparison of vegetation between planted and unplanted areas of L-Lake in the emergent zone.

No submersed or floating-leaved vegetation was present in unplanted areas.

Planting of emergent species greatly promoted the establishment of this vegetation type. Most plots in the unplanted areas remained empty or had low species diversity. In planted areas, 15.6% of the plots remained empty since planting of vegetation in 1987.

Comparison with Emergent Vegetation at PAR Pond

The *Panicum/Sacciolepis* grasses were the most important component of the vegetation in both sites (Fig. 8). *Vallesneria americana* was important in L-Lake but was not sampled in PAR Pond. Its absence from PAR Pond may be because it is a submersed species and the PAR Pond sampling emphasized emergent species. Species not sampled at L-Lake but present in PAR Pond included *Erianthus giganteus* and *Hypericum* spp. Both species were present outside the sample transects at L-Lake but in very low numbers. *Scirpus cyperinus* and *Lycopus* spp. occur more commonly above the water line at L-Lake.

Upper Emergent/Shrub Vegetation

The portion of shore from the waterline to approximately 30 cm (1 ft) above the water line delineates this zone. This area represents a portion of the shrub zone defined by Whigham et al (1985) as occurring between the water line and 0.9 m (3 ft) above the water line. The results presented here focus on herbs and shrubs, since trees greater than 2.5 cm diameter at the base were not sampled. A total of 196 plots was sampled in planted areas and 118 plots in unplanted areas (Table 2). The distinction between planted and unplanted areas is not useful for shrubs in this zone because shrubs were planted along most of the southern two-thirds of the lake. The distinction is accurate, however, for the herbaceous transplants in the other two zones.

FIGURE 8. A comparison of vegetation between L-Lake and PAR Pond in the emergent zone.

Transplant Survival

All species planted in 1987 were present in 1989. The emergents *J. effusus*, *Polygonum* spp. *S. latifolia*, *T. latifolia*, the shoreline grasses and *Panicum/Sacciolepis* were also important components of this zone (Fig. 9).

Changes from 1987 to 1989 in Planted Areas

This zone was sufficiently vegetated with terrestrial species that the proportion of barren plots was always very low (Table 2). Mean cover per plot increased from 59% in 1987 to 69% in 1988 and then decreased to 55% in 1989 (Table 2). Damage to vegetation by feral pigs was noted in 1989, but additional unknown factors may have contributed to the decrease in cover.

The most striking change was a large decrease in relative frequency and relative cover of the shoreline grasses. Probable explanations for these decreases are the same as those for the decline in shoreline grasses in the emergent zone. The *Panicum/Sacciolepis* group had a gradual increase in cover and frequency. Most individuals of cattail (*T. latifolia*) planted in this zone experienced shading by willow (*S. nigra*) and died. An overall decrease in relative frequency and relative cover of cattail has occurred (Fig. 9).

Willow was the dominant shrub. In many instances the 30-40 cm (1+ ft) shoots planted in 1987 were 5-8 meters (16-26 ft) tall in 1989. Willow reproduction occurred vegetatively. Beavers have dispersed fragments of willow that became established as new individuals elsewhere.

FIGURE 9. Importance values for species planted in the upper emergent/shrub zone.

Buttonbush (*Cephalanthus occidentalis*) does not reproduce vegetatively as well as willow. Although its relative frequency and relative cover have increased since 1987, it was still one of the least important shrub species (Fig. 9).

Comparison Between Planted and Unplanted Areas

Facultative emergent and terrestrial species were the most important components of the unplanted areas (Fig. 10). The importance values of these groups are about the same in both planted and unplanted areas, however no *J. effusus*, *Polygonum* spp. nor *Panicum/Sacciolepis* were found in unplanted areas.

Willow was a dominant shrub in the unplanted areas. It occurred in 47% of the 118 plots sampled. In the planted areas it occurred in 36% of the 196 plots sampled.

SUMMARY

The submersed plant *V. americana* is found along both shores of L-Lake and is expanding rapidly. The floating-leaved plant *N. lutea* is only found along the southern half of the eastern shore, but it also is colonizing new areas quickly. No other species are components of the submersed/floating-leaved zone. This zone was difficult to establish resulting in a species diversity below expectations. The species present are, however, well established only where they were planted, and are reproducing vegetatively. In unplanted areas there is no submersed and floating-leaved vegetation. Attempts to establish this type of vegetation in other wetlands and lakes are not known.

The establishment of emergent vegetation in L-Lake exceeded expectations. Nearly all species planted in 1987 were present in 1989. Species diversity, in planted areas, is comparable to PAR Pond. Emergent vegetation is very slowly becoming established in

FIGURE 10. A comparison of vegetation between planted and unplanted areas of L-Lake in the upper emergent/shrub zone.

comparable to PAR Pond. Emergent vegetation is very slowly becoming established in unplanted areas, and species diversity in these areas is low.

Willow is the dominate shrub in the upper emergent/shrub zone. Vegetation in this zone is a blend of emergent and terrestrial species. Woody vegetation is in an early growth stage.

Acknowledgements

Sally Landaal, Kevin Moorhead, Maria Vella, Gene Schupp, Rebbeca Sharitz and Kay Kirkman provided editorial advice. Aline Clement prepared the illustration (Figure 2) depicting plant zonation. I appreciate the help provided by these individuals. Most of all I am thankful for Priscilla Stanford's help in vegetation sampling and data entry.

REFERENCES

- Bonham, Charles D. 1989. Measurements for Terrestrial Vegetation. John Wiley and Sons. 338 pp.
- Grace, James. 1984. A survey of the wetland vegetation of PAR Pond. Unpublished. 39 pp.
- Hall, T.F. and W.T. Penfound. 1944. The biology of American Lotus, *Nelumbo lutea* (Willd.) Pers. Am. Midl. Naturalist 31:744-758.
- Jones, J.J. and R.D. Drobney. 1986. Winter feeding ecology of Scaup and Common Goldeneye in Michigan. J. Wildl. Manage. 50(3):446-452.
- Knox, J. and R. Sharitz. 1990. Endangered, Threatened, and Rare Vascular Flora of the Savannah River Site. National Environmental Research Park Program, U.S. Department of Energy. SRO-NERP-20 147 pp.
- Kollar, S.A. Jr. 1985. SAV reestablishment results-upper Chesapeake Bay. In: Coastal Zone '85: Proceedings of the Fourth Symposium on Coastal and Ocean Management. O.T. Magoon et al. editors. American Society of Civil Engineers, New York, N.Y. pp. 759-777.
- Radford, A.E., H.E. Ahles and C.R. Bell. 1968. Manual of the Vascular Flora of the Carolinas. The University of North Carolina Press, Chapel Hill. 1183 pp.
- Reed, Porter B. Jr. 1986. Wetland Plants of the State of South Carolina 1986. U.S. Fish and Wildlife Service WELUT-86/W12.40. 37 pp.
- Ristich, S.S., S.W. Fredrick and E.H. Buckley. 1976. Transplantation of *Typha* and the distribution of vegetation and algae in a reclaimed estuarine marsh. Bull. Torrey Botanical Club. 103(4):157-164.
- Whigham, D., H. Allen, G. Davis, E. Garbisch, W. Kitchens, R. Sharitz, and L. Lee. 1985. A Management Plan for Establishment of Littoral/Wetland Vegetation on the Shoreline of L-Lake on the Savannah River Plant, South Carolina. 42 pp.

Appendix A: Maps of Shoreline Vegetation

Topographic maps of portions of the shoreline of L-Lake were made by Besson and Pope in 1986. Six maps represent the southern one-half of the eastern shore and four depict the southern one-third of the western shore. Surveying stakes were placed at 200 foot intervals along the shore at about the 190 foot contour. Stakes were numbered consecutively beginning at the dam and are symbolized by a filled triangle on the maps. A stake numbered E-12 marks the twelfth 200 foot interval from the dam along the eastern shore.

The topographic maps and the surveyors' stakes were used by Southern Tier Consulting (STC) in planting. The maps denote the locations of species planted and vegetation sample transects. Sample transects are denoted by numbers and lines perpendicular to the shore.

Codes are used to designate the locations of emergent species on the maps. The first two characters represent the genus, the last two represent the specific epithet. Submersed, floating-leaved, shrub and tree species are not coded. Codes are listed on the next page.

Species codes used on the topographic maps:

<u>Code</u>	<u>Species</u>
BACA	<i>Bacopa caroliniana</i>
CACO	<i>Carex comosa</i>
DUAR	<i>Dulichium arundinaceum</i>
ELEQ	<i>Eleocharis equisetoides</i>
ELQU	<i>Eleocharis quadranulata</i>
GLSP	<i>Glyceria</i> sp.
HYCA	<i>Hydrochloa caroliniensis</i>
JUAC	<i>Juncus acuminatus</i>
JUEF	<i>Juncus effusus</i>
LYRU	<i>Lycopus rubellus</i>
PAHE	<i>Panicum hemitomon</i>
PADI	<i>Paspalum distichum</i>
POCO	<i>Pontederia cordata</i>
POSP	<i>Polygonum</i> spp.
SALA	<i>Sagittaria latifolia</i>
SCCY	<i>Scirpus cyperinus</i>
SPAM	<i>Sparganium americanum</i>
TYLA	<i>Typha latifolia</i>
VAAM	<i>Vallisneria americana</i>

SHORELINE TOPOGRAPHICAL SURVEY

OF

L-REACTOR COOLING LAKE, SAVANNAH RIVER PLANT

AIKEN COUNTY SOUTH CAROLINA

FOR

SAVANNAH RIVER ECOLOGY LABORATORY

No planting was done in this section, E-1.

Appendix B: Vegetation sampled during 1987, 1988 and 1989

This appendix summarizes all species sampled in each of the three zones between planted and unplanted areas. Species are listed alphabetically by zone. Frequency, the number of plots in which a species occurred, is denoted by N. Cover refers to the sum of the species cover. For example, in the submersed/floating-leaved zone in planted areas, *Alternanthera philoxeroides* occurred in six plots in 1988; its total cover in 1988 was 125%.

Submersed/Floating-leaved Zone, Planted

<u>Species</u>	<u>Year</u>	<u>N</u>	<u>Cover</u>
<i>Alternanthera philoxeroides</i>	1987	0	0
	1988	6	125
	1989	10	84
<i>Aneilema keisak</i>	1987	0	0
	1988	0	0
	1989	1	1
<i>Eleocharis equisetoides</i>	1987	2	2
	1988	1	10
	1989	0	0
<i>Eleocharis quadrangulata</i>	1987	1	5
	1988	0	0
	1989	2	2
Empty plots	1987	891	.
	1988	779	.
	1989	589	.
<i>Hydrocotyle umbellata</i>	1987	0	0
	1988	4	125
	1989	19	305
<i>Nelumbo lutea</i>	1987	37	322
	1988	134	6505
	1989	251	14087
<i>Nymphaea odorata</i>	1987	34	565
	1988	4	85
	1989	0	0
<i>Paspalum</i> sp.	1987	0	0
	1988	0	0
	1989	3	9
<i>Polygonum</i> sp.	1987	1	5
	1988	0	0
	1989	0	0
<i>Sacciolepis striata</i>	1987	0	0
	1988	1	5
	1989	2	2
<i>Sagittaria latifolia</i>	1987	0	0
	1988	0	0
	1989	1	2

Submersered/Floating-leaved Zone, Planted

<u>Species</u>	<u>Year</u>	<u>N</u>	<u>Cover</u>
<i>Typha latifolia</i>	1987	0	0
	1988	6	82
	1989	29	288
<i>Vallisneria americana</i>	1987	0	0
	1988	21	399
	1989	83	5674

Submersered/Floating-leaved Zone, Unplanted

<u>Species</u>	<u>Year</u>	<u>N</u>	<u>Cover</u>
Empty plots	1987	786	.
	1988	785	.
	1989	781	.
<i>Hydrocotyle umbellata</i>	1987	0	0
	1988	1	5
	1989	1	1
<i>Typha latifolia</i>	1987	0	0
	1988	0	0
	1989	5	37

Emergent Zone, Planted

<u>Species</u>	<u>Year</u>	<u>N</u>	<u>Cover</u>
<i>Alternanthera philoxeroides</i>	1987	22	158
	1988	41	768
	1989	72	610
<i>Aneilema keisak</i>	1987	3	30
	1988	5	221
	1989	8	111
<i>Bacopa caroliniana</i>	1987	4	115
	1988	7	290
	1989	9	299
<i>Bidens</i> sp.	1987	3	11
	1988	5	8
	1989	0	0
<i>Boehmeria cylindrica</i>	1987	2	40
	1988	0	0
	1989	1	1
<i>Carex comosa</i>	1987	0	0
	1988	2	95
	1989	2	40
<i>Carex</i> sp.	1987	0	0
	1988	0	0
	1989	2	2
<i>Cephalanthus occidentalis</i>	1987	1	1
	1988	1	5
	1989	1	3
<i>Cyperus esculentus</i>	1987	3	9
	1988	1	8
	1989	0	0
<i>Cyperus</i> sp.	1987	1	25
	1988	2	8
	1989	1	1
<i>Digitaria</i> sp.	1987	2	6
	1988	3	13
	1989	0	0
<i>Echinochloa crusgalli</i>	1987	1	3
	1988	14	245
	1989	0	0

Emergent Zone, Planted

<u>Species</u>	<u>Year</u>	<u>N</u>	<u>Cover</u>
<i>Eclipta alba</i>	1987	3	22
	1988	7	12
	1989	2	2
<i>Eleocharis equisetoides</i>	1987	11	42
	1988	17	536
	1989	10	64
<i>Eleocharis quadrangulata</i>	1987	35	877
	1988	53	2362
	1989	103	1893
Empty plots	1987	109	.
	1988	77	.
	1989	53	.
<i>Erechtites hieracifolia</i>	1987	0	0
	1988	1	2
	1989	2	2
<i>Eupatorium</i> sp.	1987	0	0
	1988	1	1
	1989	1	1
<i>Galium</i> sp.	1987	0	0
	1988	0	0
	1989	10	19
<i>Gerardia</i> sp.	1987	1	5
	1988	0	0
	1989	0	0
<i>Gnaphalium</i> sp.	1987	0	0
	1988	1	1
	1989	0	0
<i>Hydrochloa caroliniensis</i>	1987	4	75
	1988	6	95
	1989	10	83
<i>Hydrocotyle umbellata</i>	1987	4	19
	1988	40	540
	1989	94	390
<i>Juncus effusus</i>	1987	27	199
	1988	32	675
	1989	39	1036

Emergent Zone, Planted

<u>Species</u>	<u>Year</u>	<u>N</u>	<u>Cover</u>
<i>Juncus megacephalus</i>	1987	0	0
	1988	0	0
	1989	1	1
<i>Juncus</i> sp.	1987	2	10
	1988	2	11
	1989	3	9
<i>Leersia oryzoides</i>	1987	1	10
	1988	2	18
	1989	23	99
<i>Ludwigia alternifolia</i>	1987	0	0
	1988	0	0
	1989	3	5
<i>Ludwigia decurrens</i>	1987	0	0
	1988	6	39
	1989	0	0
<i>Ludwigia leptocarpa</i>	1987	8	54
	1988	8	22
	1989	5	16
<i>Ludwigia</i> sp.	1987	1	35
	1988	0	0
	1989	1	1
<i>Lycopus rubellus</i>	1987	0	0
	1988	0	0
	1989	3	13
<i>Lycopus virginicus</i>	1987	0	0
	1988	0	0
	1989	2	2
<i>Mikania scandens</i>	1987	2	105
	1988	5	10
	1989	7	33
<i>Nelumbo lutea</i>	1987	31	294
	1988	58	1284
	1989	59	1032
<i>Nymphaea odorata</i>	1987	10	151
	1988	1	5
	1989	0	0

Emergent Zone, Planted

<u>Species</u>	<u>Year</u>	<u>N</u>	<u>Cover</u>
<i>Panicum dichotomiflorum</i>	1987	0	0
	1988	2	6
	1989	0	0
<i>Panicum hemitomon</i>	1987	44	543
	1988	60	1759
	1989	64	1759
<i>Paspalum distichum</i>	1987	16	377
	1988	0	0
	1989	0	0
<i>Paspalum sp.</i>	1987	0	0
	1988	7	178
	1989	1	2
<i>Polygonum hydropiperoides</i>	1987	0	0
	1988	24	889
	1989	35	502
<i>Polygonum persicaria</i>	1987	0	0
	1988	15	601
	1989	4	6
<i>Polygonum sagittatum</i>	1987	0	0
	1988	0	0
	1989	1	5
<i>Polygonum sp.</i>	1987	34	879
	1988	0	0
	1989	0	0
<i>Pontederia cordata</i>	1987	8	303
	1988	17	158
	1989	20	119
<i>Rubus sp.</i>	1987	0	0
	1988	0	0
	1989	2	5
<i>Sacciolepis striata</i>	1987	7	271
	1988	8	85
	1989	29	921
<i>Sagittaria latifolia</i>	1987	12	127
	1988	23	123
	1989	20	126

Emergent Zone, Planted

<u>Species</u>	<u>Year</u>	<u>N</u>	<u>Cover</u>
<i>Salix nigra</i>	1987	11	240
	1988	12	317
	1989	14	293
<i>Scirpus cyperinus</i>	1987	2	38
	1988	1	5
	1989	1	1
<i>Setaria</i> sp.	1987	0	0
	1988	2	3
	1989	3	3
<i>Sparganium americanum</i>	1987	1	2
	1988	0	0
	1989	0	0
<i>Typha latifolia</i>	1987	31	782
	1988	64	1804
	1989	100	1115
Unknown Apiaceae	1987	0	0
	1988	1	5
	1989	0	0
Unknown Cyperaceae	1987	0	0
	1988	1	1
	1989	0	0
Unknown Poaceae	1987	20	1551
	1988	1	2
	1989	1	1
Unknown Rosaceae	1987	0	0
	1988	0	0
	1989	1	1
Unknowns	1987	11	49
	1988	0	0
	1989	0	0
<i>Vallisneria americana</i>	1987	0	0
	1988	17	289
	1989	55	2804

Emergent Zone, Unplanted

<u>Species</u>	<u>Year</u>	<u>N</u>	<u>Cover</u>
<i>Alternanthera philoxeroides</i>	1987	6	276
	1988	14	355
	1989	15	241
<i>Ambrosia artemisiifolia</i>	1987	0	0
	1988	0	0
	1989	1	1
<i>Ampelopsis arborea</i>	1987	0	0
	1988	1	3
	1989	0	0
<i>Andropogon</i> sp.	1987	0	0
	1988	0	0
	1989	2	2
<i>Aneilema keisak</i>	1987	0	0
	1988	1	3
	1989	1	1
<i>Bacopa caroliniana</i>	1987	0	0
	1988	0	0
	1989	1	2
<i>Berchemia scandens</i>	1987	0	0
	1988	0	0
	1989	1	1
<i>Boehmeria cylindrica</i>	1987	0	0
	1988	0	0
	1989	1	1
<i>Cassia fasciculata</i>	1987	0	0
	1988	0	0
	1989	1	1
<i>Cyperus esculentus</i>	1987	9	55
	1988	1	3
	1989	0	0
<i>Cyperus</i> sp.	1987	1	1
	1988	1	1
	1989	0	0
<i>Digitaria</i> sp.	1987	2	2
	1988	2	110
	1989	0	0

Emergent Zone, Unplanted

<u>Species</u>	<u>Year</u>	<u>N</u>	<u>Cover</u>
<i>Eclipta alba</i>	1987	0	0
	1988	0	0
	1989	1	1
Empty plots	1987	184	.
	1988	192	.
	1989	181	.
<i>Eupatorium</i> sp.	1987	1	3
	1988	0	0
	1989	1	1
<i>Galium</i> sp.	1987	0	0
	1988	0	0
	1989	1	1
<i>Gelsemium sempervirens</i>	1987	2	26
	1988	1	30
	1989	1	1
<i>Hydrocotyle umbellata</i>	1987	0	0
	1988	7	241
	1989	14	28
<i>Juncus</i> sp.	1987	0	0
	1988	1	2
	1989	3	23
<i>Leersia oryzoides</i>	1987	1	10
	1988	1	25
	1989	5	70
<i>Lespedeza</i> sp.	1987	0	0
	1988	0	0
	1989	1	1
<i>Lonicera japonica</i>	1987	0	0
	1988	0	0
	1989	1	10
<i>Ludwigia leptocarpa</i>	1987	11	67
	1988	3	7
	1989	1	1
<i>Lycopus</i> sp.	1987	0	0
	1988	0	0
	1989	1	3

Emergent Zone, Unplanted

<u>Species</u>	<u>Year</u>	<u>N</u>	<u>Cover</u>
<i>Panicum dichotomiflorum</i>	1987	0	0
	1988	2	2
	1989	2	2
<i>Panicum hemitomon</i>	1987	0	0
	1988	0	0
	1989	1	1
<i>Paspalum</i> sp.	1987	0	0
	1988	1	1
	1989	0	0
<i>Polygonum hydropiperoides</i>	1987	0	0
	1988	1	1
	1989	0	0
<i>Polygonum sagittatum</i>	1987	0	0
	1988	0	0
	1989	1	1
<i>Pontederia cordata</i>	1987	0	0
	1988	0	0
	1989	1	1
<i>Poplar</i> sp.	1987	0	0
	1988	0	0
	1989	1	1
<i>Rubus</i> sp.	1987	3	3
	1988	1	5
	1989	1	2
<i>Sacciolepis striata</i>	1987	0	0
	1988	1	1
	1989	0	0
<i>Sagittaria latifolia</i>	1987	6	90
	1988	4	24
	1989	1	2
<i>Salix nigra</i>	1987	0	0
	1988	2	4
	1989	5	7
<i>Solidago</i> sp.	1987	0	0
	1988	0	0
	1989	1	1

Emergent Zone, Unplanted

<u>Species</u>	<u>Year</u>	<u>N</u>	<u>Cover</u>
<i>Typha latifolia</i>	1987	0	0
	1988	6	97
	1989	15	109
Unknown Asteraceae	1987	0	0
	1988	1	1
	1989	0	0
Unknown Fabaceae	1987	0	0
	1988	0	0
	1989	1	1
Unknown Poaceae	1987	8	221
	1988	1	10
	1989	1	1
Unknowns	1987	4	7
	1988	0	0
	1989	1	1

Upper Emergent/Shrub Zone, Planted

<u>Species</u>	<u>Year</u>	<u>N</u>	<u>Cover</u>
<i>Acalypha gracilens</i>	1987	0	0
	1988	0	0
	1989	2	2
<i>Acer negundo</i>	1987	0	0
	1988	8	13
	1989	9	17
<i>Acer rubrum</i>	1987	13	26
	1988	30	94
	1989	20	127
<i>Alternanthera philoxeroides</i>	1987	13	98
	1988	92	482
	1989	105	365
<i>Ambrosia artemisiifolia</i>	1987	5	9
	1988	5	9
	1989	10	10
<i>Ambrosia</i> sp.	1987	1	1
	1988	3	3
	1989	0	0
<i>Ammannia coccinea</i>	1987	0	0
	1988	4	4
	1989	0	0
<i>Ampelopsis arborea</i>	1987	0	0
	1988	3	13
	1989	5	10
<i>Andropogon</i> sp.	1987	18	199
	1988	58	456
	1989	99	865
<i>Aneilema keisak</i>	1987	0	0
	1988	28	59
	1989	33	209
<i>Baccharis halimifolia</i>	1987	0	0
	1988	0	0
	1989	1	1
<i>Bacopa caroliniana</i>	1987	12	82
	1988	22	141
	1989	14	147

Upper Emergent/Shrub Zone, Planted

<u>Species</u>	<u>Year</u>	<u>N</u>	<u>Cover</u>
<i>Berchemia scandens</i>	1987	4	35
	1988	5	27
	1989	4	14
<i>Bidens</i> sp.	1987	20	126
	1988	39	100
	1989	9	13
<i>Boehmeria cylindrica</i>	1987	65	944
	1988	123	1264
	1989	142	1425
<i>Carex comosa</i>	1987	0	0
	1988	1	2
	1989	2	8
<i>Carex</i> sp.	1987	4	13
	1988	7	33
	1989	11	25
<i>Carya tomentosa</i>	1987	2	4
	1988	3	11
	1989	2	3
<i>Cassia fasciculata</i>	1987	4	17
	1988	8	15
	1989	6	6
<i>Celtis laevigata</i>	1987	1	3
	1988	0	0
	1989	1	2
<i>Celtis</i> sp.	1987	0	0
	1988	1	3
	1989	0	0
<i>Centrosema virginianum</i>	1987	0	0
	1988	0	0
	1989	14	28
<i>Cephalanthus occidentalis</i>	1987	13	51
	1988	27	133
	1989	30	114
<i>Clitoria mariana</i>	1987	2	25
	1988	9	42
	1989	0	0

Upper Emergent/Shrub Zone, Planted

<u>Species</u>	<u>Year</u>	<u>N</u>	<u>Cover</u>
<i>Cornus florida</i>	1987	0	0
	1988	1	3
	1989	1	5
<i>Crataegus</i> sp.	1987	0	0
	1988	0	0
	1989	1	1
<i>Cyperus esculentus</i>	1987	37	114
	1988	5	23
	1989	0	0
<i>Cyperus</i> sp.	1987	6	18
	1988	24	35
	1989	11	13
<i>Digitaria</i> sp.	1987	88	1719
	1988	55	1425
	1989	12	50
<i>Diodia</i> sp.	1987	0	0
	1988	4	4
	1989	10	16
<i>Diospyros virginiana</i>	1987	1	5
	1988	2	23
	1989	4	39
<i>Echinochloa crusgalli</i>	1987	11	253
	1988	0	0
	1989	0	0
<i>Echinodorus cordifolius</i>	1987	3	11
	1988	2	11
	1989	5	6
<i>Eclipta alba</i>	1987	71	1159
	1988	45	80
	1989	16	17
<i>Eleocharis acicularis</i>	1987	0	0
	1988	0	0
	1989	1	3
<i>Eleocharis quadrangulata</i>	1987	2	2
	1988	9	60
	1989	6	16

Upper Emergent/Shrub Zone, Planted

<u>Species</u>	<u>Year</u>	<u>N</u>	<u>Cover</u>
Empty plots	1987	1	.
	1988	0	.
	1989	2	.
<i>Erechtites hieracifolia</i>	1987	0	0
	1988	57	242
	1989	51	60
<i>Erianthus giganteus</i>	1987	0	0
	1988	0	0
	1989	2	3
<i>Erigeron</i> sp.	1987	0	0
	1988	59	186
	1989	0	0
<i>Eupatorium</i> sp.	1987	6	49
	1988	95	519
	1989	88	452
<i>Euphorbia</i> sp.	1987	2	25
	1988	0	0
	1989	4	8
<i>Galium</i> sp.	1987	7	39
	1988	10	50
	1989	11	15
<i>Gelsemium sempervirens</i>	1987	2	10
	1988	23	110
	1989	33	164
<i>Gerardia</i> sp.	1987	3	9
	1988	20	125
	1989	4	13
<i>Gnaphalium</i> sp.	1987	1	1
	1988	14	15
	1989	10	32
<i>Habenaria repens</i>	1987	0	0
	1988	0	0
	1989	9	15
<i>Helenium amarum</i>	1987	0	0
	1988	0	0
	1989	11	16

Upper Emergent/Shrub Zone, Planted

<u>Species</u>	<u>Year</u>	<u>N</u>	<u>Cover</u>
<i>Heterotheca</i> sp.	1987	0	0
	1988	0	0
	1989	1	1
<i>Hydrocotyle umbellata</i>	1987	14	68
	1988	47	478
	1989	47	83
<i>Hypericum gentianoides</i>	1987	1	5
	1988	3	3
	1989	3	5
<i>Hypericum</i> sp.	1987	0	0
	1988	5	15
	1989	9	10
<i>Hypericum walteri</i>	1987	0	0
	1988	0	0
	1989	3	3
<i>Impatiens capensis</i>	1987	8	40
	1988	7	19
	1989	6	8
<i>Juncus biflorus</i>	1987	0	0
	1988	0	0
	1989	26	108
<i>Juncus effusus</i>	1987	9	53
	1988	13	210
	1989	23	317
<i>Juncus megacephalus</i>	1987	0	0
	1988	0	0
	1989	1	1
<i>Juncus scirpoides</i>	1987	0	0
	1988	0	0
	1989	1	1
<i>Juncus</i> sp.	1987	12	77
	1988	33	174
	1989	11	23
<i>Lactuca</i> sp.	1987	0	0
	1988	0	0
	1989	6	7

Upper Emergent/Shrub Zone, Planted

<u>Species</u>	<u>Year</u>	<u>N</u>	<u>Cover</u>
<i>Leersia oryzoides</i>	1987	11	64
	1988	27	148
	1989	31	161
<i>Lespedeza</i> sp.	1987	0	0
	1988	1	8
	1989	5	5
<i>Liquidambar styraciflua</i>	1987	1	5
	1988	0	0
	1989	2	4
<i>Lonicera japonica</i>	1987	0	0
	1988	0	0
	1989	12	15
<i>Lonicera</i> sp.	1987	2	6
	1988	0	0
	1989	0	0
<i>Ludwigia alternifolia</i>	1987	0	0
	1988	3	8
	1989	11	16
<i>Ludwigia decurrens</i>	1987	0	0
	1988	3	4
	1989	1	2
<i>Ludwigia leptocarpa</i>	1987	102	1199
	1988	61	175
	1989	18	22
<i>Ludwigia</i> sp.	1987	0	0
	1988	1	1
	1989	0	0
<i>Lycopus rubellus</i>	1987	14	40
	1988	14	66
	1989	18	90
<i>Lycopus</i> sp.	1987	5	48
	1988	1	5
	1989	6	10
<i>Lycopus virginicus</i>	1987	0	0
	1988	10	21
	1989	24	48

Upper Emergent/Shrub Zone, Planted

<u>Species</u>	<u>Year</u>	<u>N</u>	<u>Cover</u>
<i>Mikania scandens</i>	1987	8	286
	1988	44	177
	1989	63	317
<i>Modiola caroliniana</i>	1987	0	0
	1988	0	0
	1989	1	1
<i>Myrica cerifera</i>	1987	11	93
	1988	21	194
	1989	23	337
<i>Nelumbo lutea</i>	1987	0	0
	1988	1	10
	1989	0	0
<i>Nymphaea odorata</i>	1987	1	1
	1988	0	0
	1989	0	0
<i>Nyssa sylvatica</i>	1987	0	0
	1988	0	0
	1989	1	2
<i>Panicum dichotomiflorum</i>	1987	0	0
	1988	9	29
	1989	1	1
<i>Panicum hemitomom</i>	1987	30	248
	1988	64	929
	1989	98	1507
<i>Panicum sp.</i>	1987	14	83
	1988	26	77
	1989	25	68
<i>Parthenocissus quinquefolia</i>	1987	0	0
	1988	1	2
	1989	4	17
<i>Paspalum distichum</i>	1987	15	345
	1988	0	0
	1989	0	0
<i>Paspalum sp.</i>	1987	0	0
	1988	22	337
	1989	13	25

Upper Emergent/Shrub Zone, Planted

<u>Species</u>	<u>Year</u>	<u>N</u>	<u>Cover</u>
<i>Passiflora incarnata</i>	1987	2	6
	1988	2	17
	1989	2	2
<i>Petalostemum pinnatum</i>	1987	2	2
	1988	0	0
	1989	0	0
<i>Physalis</i> sp.	1987	2	10
	1988	0	0
	1989	0	0
<i>Pinus taeda</i>	1987	2	2
	1988	20	26
	1989	25	50
<i>Platanus occidentalis</i>	1987	5	15
	1988	5	39
	1989	8	81
<i>Pluchea purpurascens</i>	1987	1	8
	1988	0	0
	1989	0	0
<i>Polygonum densiflorum</i>	1987	0	0
	1988	0	0
	1989	1	1
<i>Polygonum hydropiperoides</i>	1987	0	0
	1988	39	209
	1989	33	58
<i>Polygonum persicaria</i>	1987	0	0
	1988	59	194
	1989	32	34
<i>Polygonum sagittatum</i>	1987	0	0
	1988	1	1
	1989	4	4
<i>Polygonum</i> sp.	1987	17	118
	1988	4	40
	1989	1	1
<i>Polypremum procumbens</i>	1987	7	36
	1988	20	60
	1989	8	14

Upper Emergent/Shrub Zone, Planted

<u>Species</u>	<u>Year</u>	<u>N</u>	<u>Cover</u>
<i>Pontederia cordata</i>	1987	2	15
	1988	7	22
	1989	8	13
<i>Poplar</i> sp.	1987	0	0
	1988	0	0
	1989	1	1
<i>Potentilla</i> sp.	1987	0	0
	1988	0	0
	1989	1	1
<i>Prunus</i> sp.	1987	0	0
	1988	1	1
	1989	3	5
<i>Quercus nigra</i>	1987	1	1
	1988	0	0
	1989	0	0
<i>Quercus</i> sp.	1987	0	0
	1988	1	5
	1989	3	8
<i>Rhus copallina</i>	1987	4	30
	1988	10	57
	1989	11	31
<i>Rhus radicans</i>	1987	0	0
	1988	1	3
	1989	1	1
<i>Rhus</i> sp.	1987	4	20
	1988	0	0
	1989	0	0
<i>Rubus</i> sp.	1987	9	62
	1988	27	279
	1989	60	596
<i>Rumex</i> sp.	1987	0	0
	1988	6	13
	1989	1	1
<i>Sacciolepis striata</i>	1987	6	26
	1988	6	63
	1989	16	74

Upper Emergent/Shrub Zone, Planted

<u>Species</u>	<u>Year</u>	<u>N</u>	<u>Cover</u>
<i>Sagittaria latifolia</i>	1987	29	380
	1988	27	245
	1989	16	45
<i>Salix nigra</i>	1987	46	948
	1988	65	1532
	1989	70	1616
<i>Scirpus cyperinus</i>	1987	10	69
	1988	13	127
	1989	16	50
<i>Scutellaria</i> sp.	1987	0	0
	1988	1	2
	1989	6	6
<i>Setaria</i> sp.	1987	0	0
	1988	1	2
	1989	0	0
<i>Sida rhombifolia</i>	1987	9	13
	1988	10	24
	1989	5	7
<i>Smilax</i> sp.	1987	0	0
	1988	1	2
	1989	2	11
<i>Solanum</i> sp.	1987	0	0
	1988	0	0
	1989	1	1
<i>Solidago</i> sp.	1987	5	8
	1988	9	17
	1989	6	11
<i>Specularia perfoliata</i>	1987	1	1
	1988	0	0
	1989	0	0
<i>Spigelia marilandica</i>	1987	0	0
	1988	0	0
	1989	1	1
<i>Taxodium distichum</i>	1987	2	6
	1988	2	10
	1989	2	17

Upper Emergent/Shrub Zone, Planted

<u>Species</u>	<u>Year</u>	<u>N</u>	<u>Cover</u>
<i>Typha latifolia</i>	1987	46	911
	1988	48	775
	1989	18	78
<i>Ulmus</i> sp.	1987	0	0
	1988	3	3
	1989	3	3
Unknown Apiaceae	1987	0	0
	1988	7	29
	1989	0	0
Unknown Asteraceae	1987	16	107
	1988	16	25
	1989	27	41
Unknown Cyperaceae	1987	0	0
	1988	2	2
	1989	0	0
Unknown Fabaceae	1987	10	85
	1988	19	126
	1989	31	222
Unknown Juncaceae	1987	0	0
	1988	1	2
	1989	0	0
Unknown Lamiaceae	1987	5	13
	1988	23	56
	1989	1	1
Unknown Poaceae	1987	27	300
	1988	53	386
	1989	27	77
Unknown Rubiaceae	1987	8	17
	1988	2	2
	1989	0	0
Unknown Solanaceae	1987	3	8
	1988	0	0
	1989	0	0
Unknowns	1987	45	608
	1988	30	51
	1989	14	15

Upper Emergent/Shrub Zone, Planted

<u>Species</u>	<u>Year</u>	<u>N</u>	<u>Cover</u>
<i>Vaccinium</i> sp.	1987	0	0
	1988	2	20
	1989	2	18
<i>Vitis</i> sp.	1987	0	0
	1988	14	70
	1989	13	59
<i>Wahlenbergia marginata</i>	1987	0	0
	1988	2	4
	1989	1	1

Upper Emergent/Shrub Zone, Unplanted

<u>Species</u>	<u>Year</u>	<u>N</u>	<u>Cover</u>
<i>Acalypha gracilens</i>	1987	0	0
	1988	0	0
	1989	2	2
<i>Acer negundo</i>	1987	0	0
	1988	0	0
	1989	1	1
<i>Acer rubrum</i>	1987	4	11
	1988	4	33
	1989	6	39
<i>Alternanthera philoxeroides</i>	1987	27	197
	1988	39	346
	1989	72	326
<i>Ambrosia artemisiifolia</i>	1987	0	0
	1988	5	5
	1989	13	16
<i>Ambrosia</i> sp.	1987	2	20
	1988	1	1
	1989	1	1
<i>Ammannia coccinea</i>	1987	0	0
	1988	1	2
	1989	0	0
<i>Ampelopsis arborea</i>	1987	0	0
	1988	3	9
	1989	1	1
<i>Andropogon</i> sp.	1987	6	31
	1988	17	131
	1989	54	572
<i>Aneilema keisak</i>	1987	0	0
	1988	10	280
	1989	20	57
<i>Bacopa caroliniana</i>	1987	1	2
	1988	5	8
	1989	4	9
<i>Berchemia scandens</i>	1987	0	0
	1988	1	1
	1989	3	7

Upper Emergent/Shrub Zone, Unplanted

<u>Species</u>	<u>Year</u>	<u>N</u>	<u>Cover</u>
<i>Bidens</i> sp.	1987	13	198
	1988	23	115
	1989	7	7
<i>Boehmeria cylindrica</i>	1987	21	89
	1988	43	406
	1989	59	501
<i>Carex</i> sp.	1987	0	0
	1988	0	0
	1989	3	7
<i>Cassia fasciculata</i>	1987	0	0
	1988	4	29
	1989	9	22
<i>Cassia obtusifolia</i>	1987	0	0
	1988	0	0
	1989	1	1
<i>Cephalanthus occidentalis</i>	1987	0	0
	1988	5	13
	1989	6	11
<i>Crataegus</i> sp.	1987	0	0
	1988	1	20
	1989	1	5
<i>Cyperus esculentus</i>	1987	5	21
	1988	4	12
	1989	0	0
<i>Cyperus</i> sp.	1987	13	46
	1988	36	160
	1989	21	38
<i>Digitaria</i> sp.	1987	28	771
	1988	24	904
	1989	10	203
<i>Diodia</i> sp.	1987	0	0
	1988	2	2
	1989	3	3
<i>Echinochloa crusgalli</i>	1987	3	8
	1988	0	0
	1989	0	0

Upper Emergent/Shrub Zone, Unplanted

<u>Species</u>	<u>Year</u>	<u>N</u>	<u>Cover</u>
<i>Eclipta alba</i>	1987	24	319
	1988	15	25
	1989	10	11
Empty plots	1987	22	.
	1988	10	.
	1989	6	.
<i>Erechtites hieracifolia</i>	1987	0	0
	1988	18	35
	1989	13	27
<i>Erigeron</i> sp.	1987	0	0
	1988	38	265
	1989	1	3
<i>Eupatorium</i> sp.	1987	6	84
	1988	46	275
	1989	59	299
<i>Fuirena squarrosa</i>	1987	0	0
	1988	0	0
	1989	2	11
<i>Galium</i> sp.	1987	0	0
	1988	2	2
	1989	0	0
<i>Gelsemium sempervirens</i>	1987	5	145
	1988	13	314
	1989	8	24
<i>Gerardia</i> sp.	1987	0	0
	1988	6	21
	1989	1	1
<i>Gnaphalium</i> sp.	1987	1	1
	1988	7	7
	1989	2	3
<i>Habenaria repens</i>	1987	0	0
	1988	0	0
	1989	1	1
<i>Helenium amarum</i>	1987	0	0
	1988	0	0
	1989	8	8

Upper Emergent/Shrub Zone, Unplanted

<u>Species</u>	<u>Year</u>	<u>N</u>	<u>Cover</u>
<i>Hydrocotyle umbellata</i>	1987	2	3
	1988	9	49
	1989	21	61
<i>Hypericum gentianoides</i>	1987	0	0
	1988	8	14
	1989	1	1
<i>Hypericum sp.</i>	1987	1	1
	1988	2	2
	1989	5	5
<i>Impatiens capensis</i>	1987	0	0
	1988	3	3
	1989	0	0
<i>Juncus biflorus</i>	1987	0	0
	1988	0	0
	1989	8	20
<i>Juncus effusus</i>	1987	0	0
	1988	1	1
	1989	9	40
<i>Juncus megacephalus</i>	1987	0	0
	1988	0	0
	1989	4	8
<i>Juncus sp.</i>	1987	0	0
	1988	16	91
	1989	15	81
<i>Leersia oryzoides</i>	1987	3	31
	1988	11	90
	1989	18	73
<i>Lespedeza sp.</i>	1987	0	0
	1988	0	0
	1989	2	6
<i>Lonicera japonica</i>	1987	0	0
	1988	0	0
	1989	9	166
<i>Lonicera sempervirens</i>	1987	0	0
	1988	0	0
	1989	1	2

Upper Emergent/Shrub Zone, Unplanted

<u>Species</u>	<u>Year</u>	<u>N</u>	<u>Cover</u>
<i>Ludwigia alternifolia</i>	1987	0	0
	1988	0	0
	1989	5	19
<i>Ludwigia decurrens</i>	1987	0	0
	1988	6	32
	1989	4	4
<i>Ludwigia leptocarpa</i>	1987	40	384
	1988	49	562
	1989	22	37
<i>Ludwigia</i> sp.	1987	1	5
	1988	0	0
	1989	1	1
<i>Lycopus rubellus</i>	1987	1	1
	1988	3	7
	1989	9	22
<i>Lycopus</i> sp.	1987	0	0
	1988	0	0
	1989	11	36
<i>Lycopus virginicus</i>	1987	0	0
	1988	16	63
	1989	20	75
<i>Mikania scandens</i>	1987	0	0
	1988	6	11
	1989	9	10
<i>Myrica cerifera</i>	1987	0	0
	1988	2	15
	1989	8	27
<i>Panicum dichotomiflorum</i>	1987	0	0
	1988	13	47
	1989	6	86
<i>Panicum hemitomom</i>	1987	0	0
	1988	3	7
	1989	30	81
<i>Panicum</i> sp.	1987	1	3
	1988	9	34
	1989	9	35

Upper Emergent/Shrub Zone, Unplanted

<u>Species</u>	<u>Year</u>	<u>N</u>	<u>Cover</u>
<i>Paspalum distichum</i>	1987	0	0
	1988	1	1
	1989	0	0
<i>Paspalum sp.</i>	1987	0	0
	1988	9	90
	1989	10	20
<i>Pinus taeda</i>	1987	0	0
	1988	8	9
	1989	8	14
<i>Platanus occidentalis</i>	1987	2	3
	1988	2	13
	1989	2	6
<i>Polygonum hydropiperoides</i>	1987	0	0
	1988	8	31
	1989	22	61
<i>Polygonum persicaria</i>	1987	0	0
	1988	34	166
	1989	30	46
<i>Polygonum sagittatum</i>	1987	2	15
	1988	5	26
	1989	9	114
<i>Polygonum sp.</i>	1987	8	44
	1988	2	4
	1989	0	0
<i>Polypremum procumbens</i>	1987	4	5
	1988	4	13
	1989	2	2
<i>Pontederia cordata</i>	1987	0	0
	1988	1	5
	1989	1	5
<i>Poplar sp.</i>	1987	0	0
	1988	0	0
	1989	1	1
<i>Quercus sp.</i>	1987	0	0
	1988	2	54
	1989	3	58

Upper Emergent/Shrub Zone, Unplanted

<u>Species</u>	<u>Year</u>	<u>N</u>	<u>Cover</u>
<i>Rhus copallina</i>	1987	1	3
	1988	3	32
	1989	9	32
<i>Rubus</i> sp.	1987	7	69
	1988	12	191
	1989	35	503
<i>Rumex</i> sp.	1987	0	0
	1988	2	6
	1989	0	0
<i>Sacciolepis striata</i>	1987	0	0
	1988	3	21
	1989	0	0
<i>Sagittaria latifolia</i>	1987	17	639
	1988	18	180
	1989	22	85
<i>Salix nigra</i>	1987	20	602
	1988	39	1118
	1989	55	1368
<i>Scirpus cyperinus</i>	1987	0	0
	1988	3	31
	1989	5	18
<i>Scutellaria</i> sp.	1987	0	0
	1988	1	1
	1989	1	1
<i>Sida rhombifolia</i>	1987	6	14
	1988	1	1
	1989	0	0
<i>Smilax</i> sp.	1987	0	0
	1988	0	0
	1989	1	1
<i>Solidago</i> sp.	1987	3	20
	1988	0	0
	1989	5	9
<i>Specularia perfoliata</i>	1987	4	11
	1988	0	0
	1989	0	0

Upper Emergent/Shrub Zone, Unplanted

<u>Species</u>	<u>Year</u>	<u>N</u>	<u>Cover</u>
<i>Typha latifolia</i>	1987	7	64
	1988	15	51
	1989	19	99
Unknown Apiaceae	1987	0	0
	1988	4	75
	1989	0	0
Unknown Asteraceae	1987	0	0
	1988	2	6
	1989	5	5
Unknown Cyperaceae	1987	1	1
	1988	3	10
	1989	2	2
Unknown Fabaceae	1987	10	87
	1988	11	54
	1989	11	156
Unknown Juncaceae	1987	0	0
	1988	1	3
	1989	0	0
Unknown Lamiaceae	1987	9	37
	1988	3	3
	1989	2	2
Unknown Poaceae	1987	39	569
	1988	23	339
	1989	10	34
Unknown Solanaceae	1987	1	10
	1988	0	0
	1989	0	0
Unknowns	1987	15	70
	1988	14	19
	1989	7	19
<i>Vitis</i> sp.	1987	0	0
	1988	4	140
	1989	4	37
<i>Wahlenbergia marginata</i>	1987	0	0
	1988	1	2
	1989	0	0

END

DATE
FILMED
7/9/93