DOE/PC/89785--T3

DOE/PC/89785--T3

DE93 008537

THE SINGLE ELECTRON CHEMISTRY OF COALS

Report for the period
January 1, 1991 to March 31, 1991

John W. Larsen and
Robert A. Flowers II

Department of Chemistry
Lehigh University
Bethlehem, PA 18015

April 22, 1991

RECEIVED
DOE/PC/89785--T3
APR 22 1991
PM 1:10

PREPARED FOR THE UNITED STATES DEPARTMENT OF ENERGY

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

MASTER

ds
DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

OBJECTIVES AND SCOPE OF WORK

Objectives

To investigate and characterize the single electron reactions of coals in order to determine the role these reactions play in the chemistry of coal.

Scope of Work

1. Attempts will be made to demonstrate that the radicals from the inertinite maceral group will initiate the polymerization of 4-vinylpyridine.
2. The molecule N,N-diphenyl-phenylenediamine will be deposited in coals to characterize their native free radicals.
3. Tetracyanoquinodimethane (TCNQ) and tetracyanoethylene (TCNE) will be used to characterize the numbers and strengths of single electron donors in coals.

DETAILED DESCRIPTION OF TECHNICAL PROGRAM

Introduction

A small aromatic molecule such as benzene can act as a weak donor or a very weak acceptor in a charge transfer complex.¹ Donor ability increases with decreasing ionization potential I ; and acceptor ability increases with increasing electron affinity E . I decreases and E increases as an aromatic molecule becomes larger. $I=E$ in graphite and therefore it is as good an electron donor as it is an electron acceptor. Aside from increasing or decreasing aromatic ring size, a molecule can become a good single electron acceptor or donor by adding electron withdrawing or releasing groups to it.

Coals are known to have aromatic molecules as a part of their macromolecular structure. Low and medium rank coals are highly substituted with electron withdrawing and releasing groups.² Higher rank coals contain large aromatic ring systems.³ Coals of varying rank have the potential to donate an electron to a good single electron acceptor.

Tetracyanoquinodimethane (TCNQ) and tetracyanoethylene (TCNE) are good single electron acceptors with comparable electron affinities.⁴ The nitrile shift in the infrared spectrum of TCNQ and TCNE is a sensitive indicator of charge transfer.^{5,6} TCNQ and TCNE will be used to study the single electron donating ability of a series of Argonne Premium coal samples.

It should be noted that Retcofsky, employing Curie Law behavior has demonstrated that the native coal radicals are not

due to charge transfer.⁷ The objective of this work is to demonstrate whether new interactions within the coal produce radicals through charge transfer interactions.

Experimental

100% TCNQ or TCNE loadings (one electron acceptor per one polynuclear aromatic)³ were prepared by adding a stock solution of the electron acceptor dissolved in pyridine to a coal sample. A 0.03 M solution of TCNQ in pyridine was prepared by weighing 6.316 grams of TCNQ into a tared one liter volumetric flask and dissolving it in pyridine. The solution was purged with dry nitrogen and sealed with a septum. 100 ml of this solution would correspond to a 100% loading of TCNQ on one gram of Illinois #6 coal. The volume of solution placed into coal to correspond to a 100% loading based on PNA's could be varied for the different Argonne premium coal samples depending on average aromatic cluster size and percent aromaticity.

A typical sample preparation involved weighing one gram of coal into a roundbottom flask containing a magnetic stirrer and sealing it with a septum. The flask was removed from the drybox and approximately 100 ml of the 0.03 M stock solution was added via a syringe. The mixture was stirred for one week. The solution was rotovapped for one week and then dried in a vacuum oven at room temperature until it reached constant weight. It could not be determined whether the TCNQ was contained in the bulk of the coal or on the surface.

Infrared Spectra were recorded on a Mattson Series 100 Fourier transform infrared spectrometer equipped with Harrick optics and a sampling stage for diffuse reflectance. Spectra were recorded from the co-addition of 1000 scans of 9000 data points at 4 cm^{-1} resolution against a background of KCl. A typical sample preparation involved separately grinding 200 mg of coal and KCl in a Wig-L-Bug for 3 minutes, followed by co-grinding a 5% mixture of coal and KCl for an additional 3 minutes.⁸ All operations were carried out in a Vacuum Atmosphere drybox in dry N_2 free of O_2 .

Results and Discussion

Figure 1 contains the diffuse reflectance Fourier transform infrared (DRIFT) spectra of TCNQ and TCNE. The nitrile band of TCNQ is located at 2227 cm^{-1} . The nitrile band of TCNE is located at 2236 cm^{-1} . It should be noted that these bands are narrow and sharp. Upon transfer of electron density from a donor to these acceptors, these bands become very broad.

Figure 2 shows the DRIFT spectra of Illinois #6 coal with a 100% loading of TCNQ and TCNE based on the PNA's in the coal. In the case of TCNQ, the nitrile shift corresponds to complete electron transfer from the coal to TCNQ. The TCNE Illinois #6 spectrum shows a 36 cm^{-1} shift. Figures 3 through 5 show the 1-1 complexes of TCNQ and TCNE (based on PNA's) for Beulah-Zap, Pittsburgh #8 and Lewiston-Stockton coals. The nitrile stretch in these spectra also show the same behavior discussed for Illinois #6 coal.

One major feature of all of the spectra containing a 100% loading of TCNQ is the presence of a structureless absorption between 3700 to 2300 cm^{-1} . This band may be due to an electronic transition that lies in the infrared region. If some type of PNA-TCNQ stack is formed, the energy difference between the HOMO's and LUMO's in such a stack may be sufficiently small for electronic transitions to lie in the infrared region.

Figures 6 and 7 display the spectra for Upper Freeport and Pocahontas #3 with a 100% loading of TCNQ and TCNE based on PNA's. It is apparent that in the case of TCNQ, only a small amount of the acceptor is interacting with the coal. The parent $\text{C}\equiv\text{N}$ stretch at 2227 cm^{-1} dominates the spectrum and only a small amount appears to be completely shifted. The TCNE however, displays a large shift in these coals consistent with the other coals discussed above. Coals above 87% C do not swell appreciably and it is possible that the TCNQ molecule may be too large to penetrate these coals. The TCNE is smaller and even a small amount of swelling may afford diffusion of the TCNE into the coal network.

Conclusions

The simplest explanation for these shifts is there exists in coal single electron donors which are capable of transferring an electron to TCNQ in the ground state. All of the TCNQ placed in the coal appears to be converted to the radical anion as displayed in the IR spectrum for all of the coals except for the 100% loading

in the two high rank coals. The low loadings of TCNQ (20 mg acceptor/g of coal) contain enough TCNQ to form a complex with 3% of the PNA's in coal. It is very possible that 3% of the PNA's in coal have a sufficiently low ionization potential to transfer an electron to TCNQ. However, in a 100% loading, it is highly unlikely that all or most of the PNA's in coal have a sufficiently low ionization potential to donate an electron to TCNQ.

Another explanation for the shift is there are cooperative interactions between the TCNQ or TCNE with the aromatic species in the coal. The TCNQ or TCNE forms extensive stacks with the aromatic groups in coals. This interaction has led to the formation of an extended valence band structure with the TCNQ LUMO being half filled or in other words occupied with one electron. In the case of TCNE, its LUMO is approximately 30% occupied.

References

1. Mulliken, R.S.; Person, W.B. 1969, "Molecular Complexes", Wiley-Interscience, New York, p 4.
2. Shinn, J.H. Fuel, 1987, 63, 1187.
3. Solum, M.S.; Pugmire, R.J.; Grant, D.M. Energy and Fuels, 1989, 3, 187.
4. Hester, R.E. in "Advances in Infrared and Raman Spectroscopy, vol 4", Clark, R.J.H., Hester, R.E. Eds., Heyden & Son Ltd., 1980, London, UK.
5. Chappel, J.S.; Bloch, A.N.; Bryden, W.A.; Maxfield, M.; Poehler, T.O.; Cowan, D.O. J. Am. Chem. Soc., 1981, 103, 2442.
6. Moore, J.C.; Smith, D.; Youhne, Y.; Devlin, J.P. J. Phys. Chem., 1971, 75, 325.
7. Retcofsky, H.L.; Hough, M.R.; Macguire, M.M.; Clarkson, R.B. in "Coal Structure", Gorbaty, M.L. and Ouchi, K. Eds. ACS Adv. Chem. Ser. 1981, 192, 37.
8. Fuller, M.P.; Griffiths, P.R. Anal. Chem., 1978, 50, 1906.

List of Figures

- Figure 1. a) DRIFT Spectrum of TCNQ; b) DRIFT Spectrum of TCNE.
- Figure 2. a) Illinois #6 coal With a 100% Loading of TCNQ (based on PNA's); Illinois #6 Coal With a 100% Loading of TCNE (based on PNA's).
- Figure 3. a) Beulah-Zap Lignite With a 100% Loading of TCNQ (based on PNA's); Beulah-Zap Lignite With a 100% Loading of TCNE (based on PNA's).
- Figure 4. a) Pittsburgh #8 Coal With a 100% Loading of TCNQ (based on PNA's); b) Pittsburgh #8 Coal With a 100% Loading of TCNE (based on PNA's).
- Figure 5. a) Lewiston-Stockton Coal With a 100% Loading of TCNQ (based on PNA's); Lewiston-Stockton Coal With a 100% Loading of TCNE (based on PNA's).
- Figure 6. a) Upper Freeport Coal With a 100% Loading of TCNQ (based on PNA's); b) Upper Freeport Coal With a 100% Loading of TCNE (based on PNA's).
- Figure 7. a) Pocahontas #3 Coal With a 100% Loading of TCNQ (based on PNA's); b) Pocahontas #3 Coal With a 100% Loading of TCNE (based on PNA's).

Figure 1 (a) DRIFT spectrum of TCNQ. (b) DRIFT spectrum of TCNE.

Figure 2 (a) DRIFT Spectrum of Illinois #6 Coal With a 100% Loading of TCNQ. (based on PNA's) (b) DRIFT Spectrum of Illinois #6 Coal With a 100% Loading of TCNE. (based on PNA's)

Figure 2 (a) DRIFT Spectrum of Beulah-Zap Lignite With a 100% Loading of TCNQ. (based on PNA's) (b) DRIFT Spectrum of Beulah-Zap Lignite With a 100% Loading of TCNE. (based on PNA's)

Figure 4 (a) DRIFT Spectrum of Pittsburgh #8 Coal With a 100% Loading of TCNQ. (based on PNA's) (b) DRIFT Spectrum of Pittsburgh #8 Coal With a 100% Loading of TCNE. (based on PNA's)

Figure 5 (a) DRIFT Spectrum of Lewiston-Stockton Coal With a 100% Loading of TCNQ. (based on PNA's) (b) DRIFT Spectrum of Lewiston-Stockton Coal With a 100% Loading of TCNE. (based on PNA's)

Figure 6 (a) DRIFT Spectrum of Upper Freeport Coal With a 100% Loading of TCNQ. (based on PNA's) (b) DRIFT Spectrum of Upper Freeport Coal With a 100% Loading of TCNE. (based on PNA's)

Figure 7 (a) DRIFT Spectrum of Pocahontas #3 Coal With a 100% Loading of TCNQ. (baed on PNA's) (b) DRIFT Spectrum of Pocahontas #3 With a 100% Loading of TCNE. (based on PNA's)

END

**DATE
FILMED**

4 / 1 / 93

.