NOTICE

**CERTAIN DATA
CONTAINED IN THIS
DOCUMENT MAY BE
DIFFICULT TO READ
IN MICROFICHE
PRODUCTS.**

DE-FG03-92ER25129/A001

U.C. Berkeley

Project Title

Stability, Bifurcation, and Control of Hamiltonian Systems

DOE/ER/25129--T1

DE93 012103

1 Principal Investigators

Jerrold E. Marsden
Department of Mathematics
University of California
Berkeley, CA 94720
510-642-5229

DISCLAIMER

and

Tudor S. Ratiu
Department of Mathematics
University of California
Santa Cruz, CA 95034.
408-459-4180

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

2 Principal Project Personnel

Jerrold E. Marsden

A. Role in the project. Principal Investigator/Researcher

B. Principal Areas of Research and Expertise. Mechanics, Dynamics, Bifurcation theory.

C. Percentage of time. Approximately 20 %

D. Education. PhD Princeton, 1968

E. Employment History. University of California, Berkeley, 1968-present.

F. Relevant Professional Societies and Honors. Editor for the Journal of Nonlinear Science and Springer's Applied Mathematical Sciences Series, Member, AMS, SIAM, APS; Gravity Research Foundation prize, 1973, Carnegie Fellowship, 1977, Killam Scholar, 1979, Wiener Prize, 1990, Humboldt Prize, 1991, Fairchild Scholar, 1992.

G. Relevant prior publications not emanating from this award:

Holmes, P.J., J.E. Marsden and J. Scheurle [1988] Exponentially small splittings of separatrices with applications to KAM theory and degenerate bifurcations, *Cont. Math.* **81**, 213-244.

Oh, Y.G., N. Sreenath, P.S. Krishnaprasad and J.E. Marsden [1989] The Dynamics of Coupled Planar Rigid Bodies Part 2: Bifurcations, Periodic Solutions, and Chaos, *Dynamics and Diff. Eq'ns.* **1**, 269-298.

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

- Bloch, A.M. and J.E. Marsden [1989] Controlling homoclinic orbits, *Theoretical and Computational Fluid Mechanics* **1**, 179-190.
- Bloch, A.M. and J.E. Marsden [1990] Stabilization of rigid body dynamics by the energy-Casimir method, *Syst. and Cont. Lett.* **14**, 341-346.
- Simo, J.C., T.A. Posbergh and J.E. Marsden [1990] Stability of coupled rigid body and geometrically exact rods: block diagonalization and the energy-momentum method, *Physics Reports* **193**, 280-360.
- Simo, J.C., D.R. Lewis and J.E. Marsden [1991] Stability of relative equilibria I: The reduced energy momentum method, *Arch. Rat. Mech. Anal.* **115**, 15-59.
- Chern, S.J. and J.E. Marsden [1990] A note on symmetry and stability for fluid flows, *Geo. Astro. Fluid. Dyn.* **51**, 1-4.
- Marsden, J.E., T.S. Ratiu and G. Raugel [1991] Symplectic connections and the linearization of Hamiltonian systems, *Proc. Roy. Soc. Ed. A* **117**, 329-380.
- Holm, D.D. and J.E. Marsden [1991] The rotor and the pendulum, *Symplectic Geometry and Mathematical Physics*, ed. by P. Donato, C. Duval, J. Elhadad, and G.M. Tuynman, Birkhäuser, Boston, 189-203.
- J.E. Marsden [1992], *Lectures on Mechanics* London Mathematical Society Lecture note series, **174**, Cambridge University Press.

Tudor S. Ratiu

A. Role in the project. Principal Investigator/Researcher

B. Principal Areas of Research and Expertise. Mechanics, Dynamics, Integrable Systems.

C. Percentage of time. One summer month.

D. Education. PhD Berkeley, 1980

E. Employment History. University of Michigan, 1980-83, University of Arizona, Tucson, 1983-87, University of California, Santa Cruz, 1987-present.

F. Relevant Professional Societies and Honors. AMS; Sloan Fellowship, NSF Postdoctoral Fellowship, and Miller Fellowship Award, U.C. Berkeley, Spring, 1994.

G. Relevant prior publications not emanating from this award:

Holm, D.D., J.E. Marsden, T.S. Ratiu and A. Weinstein [1985] Nonlinear stability of fluid and plasma equilibria, *Phys. Rep.* **123**, 1-116.

Marsden, J.E. and T.S. Ratiu [1986] Reduction of Poisson manifolds, *Lett. in Math. Phys.* **11**, 161-170.

Abarbanel, H.D.I., D.D. Holm, J.E. Marsden and T.S. Ratiu [1986] Nonlinear stability analysis of stratified fluid equilibria, *Phil. Trans. R. Soc. Lond. A* **318**, 349-409; also *Phys. Rev. Lett.* **52** [1984] 2352-2355.

Lewis, D.R., J.E. Marsden, R. Montgomery and T.S. Ratiu [1986] The Hamiltonian structure for dynamic free boundary problems, *Physica D* **18**, 391-404.

- Lewis, D.R., J.E. Marsden and T.S. Ratiu [1987] Stability and bifurcation of a rotating liquid drop, *J. Math. Phys.* **28**, 2508-2515.
- Mazer, A. and T. Ratiu [1989] Hamiltonian formulation of adiabatic free boundary Euler flows. *J. Geom. and Phys.* **6**, 271-291.
- Mazer, A. and T. Ratiu [1989] Formal stability of two-dimensional self-gravitating rotating disks. *Cont. Math.* **99**, 233-258.
- Bloch, A.M., R.W. Brockett and T.S. Ratiu [1990] A new formulation of the generalized Toda Lattice equations and their fixed point analysis via the momentum map, *Bull. Amer. Math. Soc.* **23**, 477-485.
- Marsden, J.E., R. Montgomery and T.S. Ratiu [1990] *Reduction, symmetry, and phases in mechanics*. Memoirs AMS **430**.
- Lewis, D.R., J.E. Marsden, T.S. Ratiu and J.C. Simo [1990] Normalizing connections and the energy-momentum method, Proceedings of the CRM conference on *Hamiltonian systems, Transformation Groups, and Spectral Transform Methods*, CRM Press, Harnad and Marsden (eds.), 207-227.

3 Additional Project Personnel

Graduate students supported in the past on DOE funds included Debra Lewis and Richard Montgomery (now at Santa Cruz), Gloria Sanchez (normally in Venezuela, but currently visiting Berkeley) Mark Alber (for travel), now at Notre Dame, and current students receiving support include C.Y. Xu at Berkeley. (Funds were not sufficient to support more students or students at Santa Cruz).

Collaborators at other Universities were not supported by these funds, but include Juan Simo (Applied Mechanics, Stanford), Anthony Bloch (Mathematics, Ohio State), Darryl Holm (Center for Nonlinear Studies, Los Alamos), Genevieve Raugel (Paris), Jürgen Scheurle (Hamburg) and P.S. Krishnaprasad (Systems Research Center, Maryland)

4 Project Overview

A. Project Objectives.

1. **Past.** In 1985 Holm, Marsden, Ratiu and Weinstein, in 1986 Abarbanel, Holm, Marsden, and Ratiu, and in 1987 Krishnaprasad and Marsden, the Arnold or energy-Casimir method underwent major development and was applied to a wide variety of fluid, plasma, and elastic systems to obtain specific stability results for relative equilibria. This work builds on the results of several people on Hamiltonian structures for mechanical systems with symmetry. To overcome limitations of this method for elastic systems, the energy-momentum method was developed by Marsden, Simo, Lewis, and Posbergh. In the course of this development a remarkable block diagonal structure was discovered which not only overcame the intended difficulty, but also made the method more implementable in complex examples, by giving a normal form to the second variation of the Hamiltonian and the symplectic structure and effectively providing a separation of rotational and internal modes in rotating systems. It is especially useful for applications to large scale systems since it avoids the inversion of the large matrices that can occur in the Lagrangian to Hamiltonian conversion, for instance. Although the full implications are still a subject of current research, this theory will have a major impact. One of the major themes of our research is to connect these developments in reduction theory with developments in control strategies (work of Krishnaprasad, Bloch, Marsden, Sanchez) and with work in Geometric phases.

In 1989, Bloch and Marsden developed a basic method for using control theory for dynamical systems with homoclinic orbits. (This is not unrelated to current work in control of chaos!). It was developed for boundary layer bursting in fluid dynamics specifically, but the methods apply rather generally. Since homoclinic orbits and cycles play such a large role in a variety of mechanical system, such as rigid body systems, we plan to continue this study in that context. In addition, the effect of geometric phases will be brought into the picture.

In 1992, Bloch, Krishnaprasad, Marsden, and Sanchez showed how to design control strategies to stabilize otherwise unstable equilibria (such as rotation of a rigid body with internal rotors about a middle axis) using reduction theory to design the control law. Now that the basic mechanism is understood, and is implemented in at least one nontrivial example, we propose to link these ideas with those described in the previous section. Since the methods are general, and ultimately are based on symmetry ideas (including the block diagonal results) we have every expectation that our results will extend from the case of a rigid body with internal rotors to more complex systems such as rigid bodies with flexible components.

For the single rotor case, the preceding reference proves stabilization of the system about the intermediate axis for sufficiently large torque by using the Energy-Casimir method. Related to its Hamiltonian structure is the existence of conserved quantities in other forced systems, such as the kinematic chains of Baillieul and Baillieul and Levi. The question of stabilization of Hamiltonian control systems has been considered, although in a rather different fashion by van der Schaft. It also shows, using the geometric phases ideas of Marsden, Montgomery and Ratiu, and Krishnaprasad, how an attitude drift can occur if the body is perturbed away from a stable equilibrium. Generalizing a formula of Montgomery, we have shown how to calculate this drift precisely and indicate how to compensate for it. Again, one of the objectives is to put the experience gained from examples like this into a more general context provided by Lagrangian reduction theory.

2. Current. One of the main current pieces of work being completed is the work on dissipation induced instabilities (Bloch, Krishnaprasad, Marsden, Ratiu). Here we show that the predictions of the energy-momentum method are optimal if one is willing to add on small amounts of dissipation. In other words, if one has a saddle point of the second variation of the augmented energy (as determined by the block diagonalization method) then with the addition of physical dissipation, the system spectrally destabilizes. This phenomenon has been observed in a number of specific situations, such as rotating shafts, but we prove it for general systems. It also shows how the energy-momentum methods predictions are sharp. Consider a Hamiltonian system whose linearization at an equilibrium (or relative equilibrium) takes the form

$$\dot{x} = v, \quad \dot{v} = -\Lambda x + Sv$$

where $S = -S^T$ gives the gyroscopic forces and $\Lambda = \Lambda^T$. This form can be obtained from the block-diagonalization method. Assume that this Hamiltonian system is unstable for $S = 0$ and neutrally stable for $S \neq 0$, i.e., that some of the eigenvalues of Λ are in the left half-plane but that the eigenvalues of the entire system return to the imaginary axis for certain non-zero S ; this phenomenon goes under the name of *gyroscopic stabilization*. The question one raises is if the stability of this system is destroyed by the addition of a small damping term to the equations of motions. Precisely, we assume that $dH/dt = -2\mathcal{R}$, where the Rayleigh dissipation function is given by $2\mathcal{R} = v^T R v$ for $R = R^T \geq 0$.

It is shown that the answer to the above question is affirmative and therefore that gyroscopic stabilization is destroyed by the addition of small damping. This is true for both finite dimensional and certain classes of infinite dimensional systems, such as rotating rods and strings.

This result is proved by a combination of the techniques of Chetaev (to show Liapunov instability) and Hahn (to show from this that there is some eigenvalue in the right half plane). The above system appears as a linearization of a Hamiltonian system with symmetry if the symmetry group is abelian. In the non-abelian case the linearized system involves the interaction between the "rigid" and "internal" variables. We have also obtained the result in this case, and have applied it to the problem of a rigid body with internal momentum wheels, for example. What we are now in a position to do is to combine these techniques with the others in this proposal to enable one to

better understand the control of these mechanical systems. For example, one of the first things is to use this method along with the stabilizing feedback controls of (Bloch, Krishnaprasad, Marsden, and Sanchez, 1992) to show that one gets asymptotically stable solutions, by simulating damping (positive or negative) in the control law. (This idea was independently arrived at by Kammer.)

The proof of these results for the nonabelian case requires a substantial extension of the abelian case. The strategy here is to create a nontrivial modification of the Chetaev functional; the key point is that this modification must take into account the interaction between the internal modes and the overall rotational modes.

Examples like the classical ozone molecule, or the rigid body with internal rotors has a nonabelian group $SO(3)$ and has the interesting feature of a discrete symmetry group as well. This example is interesting for a study of the Euler-Lagrange-Poincaré equations, for bifurcations of relative equilibria, and for testing the effect of dissipation and controls in the nonabelian case.

Infinite dimensional examples for dissipation induced instabilities also need further development. Two relevant examples to which it formally applies are the rotating liquid drop following the 1987 set up of Lewis, Marsden, and Ratiu, and the second is the Richardson number criterion for stratified flow, following Abarbanel, Holm, Marsden, and Ratiu. In each case we have interesting regimes in which there are eigenvalues on the imaginary axis, yet the second variation of the (augmented) Hamiltonian is indefinite. Other examples of Mahalov will also be considered in the context of Hamiltonian bifurcations considered next.

We have also completed, with Jürgen Scheurle, a work on Lagrangian reduction, that has its own interest and was used to give a bifurcation analysis of the double spherical pendulum. This work has sparked interest in the theory of nonholonomic constraints. We are also currently working with Genevieve Raugel on Euler flow in thin domains, such as on S^2 . This is motivated by work of Chern and Marsden on spherical ABC flows, and by work of Sell, Hale and Raugel on thin domain methods. We are also discussing these ideas with Holm in his work on the shallow water equations and ocean dynamics.

3. Planned Future Work. We plan to develop the Euler-Lagrange-Poincaré equations, recently discovered by Marsden and Scheurle (an unconstrained version of the work mentioned in the last paragraph) and relate it to the work described above. For example these equations are closely related to the Lagrangian side of the energy momentum method (as developed by Lewis) and to work of Koiller, Krishnaprasad, and Yang on nonholonomic constrained systems (of the sort arising in robotics). Also, we will be developing methods useful for extending the dissipation induced instabilities results to some infinite dimensional systems and to semidirect product systems (such as stratified fluids), and to include resonance bifurcations. In particular (with Mary Silber and Vivien Kirk) we have developed a geometric phase formula (building on work of Marsden, Ratiu and Montgomery) for pinched spheres. Some of the ideas are related to work of Holm on polarization laser dynamics. These sorts of ideas will help us to extend the dissipation induced instabilities results to include the effects of the nonlinear terms and normal forms. We have also just completed a work on the generic movement of eigenvalues (with Dellnitz and Melbourne) and on bifurcations in the heavy top (with Simo, and Lewis) and we plan to use these results in the general development of Hamiltonian bifurcations. In this connection, the work of Lewis on stability and bifurcation in singular cases is quite important, and we plan to link our approach with her results.

Here are a few more details: In 1991-22, Marsden and Scheurle studied Lagrangian reduction in the context of systems with a momentum map constraint. This is a Lagrangian analogue of Hamiltonian symplectic reduction. The Lagrangian description focuses on the variational principle for reduced systems. We will be developing this theory for unconstrained systems, which is the analogue of Poisson reduction for Hamiltonian systems. Our NSF work develops the differential geometry and pure mathematical side of this project, while the DOE work will focus on the applications side.

For systems with the momentum map $\mathbf{J}$ constrained to a specific value μ , the key to the construction is the modification of the Lagrangian L to the Routhian R^μ , which is obtained from the Lagrangian by subtracting off the mechanical connection paired with the constraining value μ of the momentum map. A basic ingredient needed for the Euler-Lagrange-Poincaré equations is a

velocity shift in the Lagrangian (the shift is given by a choice of connection, often the mechanical connection—recalled below), so this velocity shifted Lagrangian plays the role that the Routhian does in the constrained theory.

For the Euler-Lagrange-Poincaré equations, the context is a principle (left) G -bundle $Q \rightarrow S$ as the configuration space, and one starts with a G invariant Lagrangian L on TQ . One divides the variations up into horizontal and vertical parts using the given connection, which divides up the original Euler-Lagrange equations. Remarkably, although this is, in principle, a simple conceptual idea, it has largely gone unnoticed (except perhaps for Cendra, Ibrort, and Marsden in 1987), but will play a role for many of our investigations. The resulting coupled system of equations are as follows:

$$\frac{d}{dt} \frac{\partial}{\partial \dot{x}^i} l(x^i, \dot{x}^i, \xi) - \frac{\partial}{\partial x^i} l(x^i, \dot{x}^i, \xi) = \frac{\partial l}{\partial \xi^a} \Omega_{ij}^a \dot{x}^j \quad (4.1)$$

$$\frac{d}{dt} \frac{\partial}{\partial \xi^a} l(x^i, \dot{x}^i, \xi) = c_{ba}^c \frac{\partial l}{\partial \xi^c} \xi^b \quad (4.2)$$

Here, l is the Lagrangian $l: TQ/G \rightarrow \mathbb{R}$ induced by the given Lagrangian L on TQ , where we regard TQ/G as a $\mathfrak{g}$ (the Lie algebra of G) bundle over TS . The coordinates x are in the base space S and the variables ξ are on $\mathfrak{g}$. Also, Ω is the curvature form and c_{ba}^c are the structure constants of $\mathfrak{g}$. The second set of equations, the Poincaré equations, were discovered by Poincaré in 1901. However, how the Poincaré equations couple to the Euler-Lagrange equations through the connection is new. If one writes the Euler-Lagrange equations on TQ directly in a local trivialization, one gets the above system without the curvature term on the Euler-Lagrange side. However, this representation of the equations is not intrinsic, while with the curvature, it is, representing the original variational principle relative to horizontal and vertical variations. One gets from one form to the other by means of the velocity shift given by the connection.

The variables in the equations are related to the rigid (ξ) and internal (x) variables. As has been shown by Simo, Lewis, and Marsden, the division of variables into internal and rigid parts has deep implications for both stability theory and for bifurcation theory, again, continuing along lines developed originally by Riemann, Poincaré and others. The main way this new insight is achieved is through a careful split of the variables, using the mechanical connection as a main ingredient. This split puts the second variation of the augmented Hamiltonian at a relative equilibrium as well as the symplectic form into normal form. It is remarkable that they are put into this form simultaneously. The method enables one to obtain stability results that would otherwise be intractable. One of our longer term goals is to link the Euler-Lagrange-Poincaré equations with this theory in the hopes of obtaining a division of the variables into rigid and internal for the nonlinear theory as well as for the linearized theory, and to examine the singular case from this point of view as well.

Abelian Lagrangian reduction is due to Routh (1860). Marsden and Scheurle give a geometrization and a generalization of the Routh procedure to the nonabelian case. One incorporates the conservative gyroscopic forces into the variational principle in the sense of Lagrange and d'Alembert and employ a Dirac constraint construction to include the cases in which the reduced space is not a tangent bundle. The nonabelian case is illustrated by the rigid body.

Given $\mu \in \mathfrak{g}^*$, define the Routhian $R^\mu: TQ \rightarrow \mathbb{R}$ as follows:

$$R^\mu(q, v) = L(q, v) - \langle \alpha(q, v), \mu \rangle \quad (4.3)$$

where α is the mechanical connection. The Routhian has the form of a *Lagrangian with a gyroscopic term*.

Solutions of the Euler-Lagrange equations for L can be regarded as solutions of the Euler-Lagrange equations for the Routhian, with the addition of "magnetic forces". One proves that the Euler-Lagrange equations for L are equivalent to the Euler-Lagrange-d'Alembert equations for the Routhian. This Routhian is closely related to the classical Routhian as Marsden and Scheurle show: $R_{\text{classical}}^\mu = R^\mu + \mu_c g_{ac} \dot{q}^a \mathbb{I}^{bc}$. Here, $\mathbb{I}$ is the locked inertia tensor.

One can drop the variational principle to the quotient space Q/G_μ , with $\mathcal{L} = R^\mu$. In this principle, the variation of the integral of R^μ is taken over curves satisfying the fixed endpoint

condition; this variational principle therefore holds in particular if the curves are also constrained to satisfy the condition $J(q, v) = \mu$. Then we find that the variation of the function R^μ restricted to the level set of J satisfies the variational condition. The restriction of R^μ to the level set equals

$$R^\mu = \frac{1}{2} \|\text{hor}(q, v)\|^2 - V_\mu \quad (4.4)$$

In the resulting variational principle, the endpoint conditions can be relaxed to the condition that the ends lie on orbits rather than be fixed. This is because the kinetic part now just involves the horizontal part of the velocity, and so the endpoint conditions in the variational principle, which involve the contraction of the momentum p with the variation of the configuration variable δq vanish if $\delta q = \zeta_Q(q)$ for some $\zeta \in \mathfrak{g}$, i.e., if the variation is tangent to the orbit. The condition that (q, v) be in the μ level set of J means that the momentum p vanishes when contracted with an infinitesimal generator on Q .

The function R^μ restricted to the level set defines a function on the quotient space $T(Q/G_\mu)$ - that is, it factors through the tangent of the projection map $\tau_\mu : Q \rightarrow Q/G_\mu$. The variational principle also drops, therefore, since the curves that join orbits correspond to those that have fixed endpoints on the base. Note, also, that the magnetic term defines a well-defined two form on the quotient as well, as is known from the Hamiltonian case, even though α_μ does not drop to the quotient in general. In terms of the coordinate representation for the special case of a torus action, and cyclic variables, this can be seen from the fact that α_μ depends on the θ -variables, whereas β does not, because the 1-form $\mu_a d\theta^a$ is closed. However, on the quotient we have the well defined magnetic two form $\beta = d(g_{t\alpha} \mu_a \tilde{\tau}^{ab} dq^\alpha)$.

To summarize, suppose that $q(t)$ satisfies the Euler-Lagrange equations for L with $J(q, \dot{q}) = \mu$, then the induced curve on Q/G_μ satisfies the **reduced Lagrangian variational principle**, i.e., the variational principle of Lagrange-d'Alembert on Q/G_μ with magnetic term β and the Routhian dropped to $T(Q/G_\mu)$. For a torus action, i.e., with cyclic variables, this reduced variational principle is equivalent to the Euler-Lagrange equations for the classical Routhian, which agrees with the classical procedure of Routh.

There is a well defined reconstruction procedure for these systems. One can horizontally lift a curve in Q/G to a curve $q(t)$ in Q (which therefore has zero angular momentum) and then one rotates it by the group action by a time dependent group element solving the equation $\dot{g}(t) = g(t)\xi(t)$ where $\xi(t) = \alpha(q(t))$, as is used in the development of geometric phases.

There are some interesting systems for which the Legendre transformation is degenerate. For example, this occurs for the Vlasov-Poisson equations of plasma physics (Marsden and Weinstein, 1982). The Vlasov-Poisson equations are purely Lie-Poisson (using the group of canonical transformations), but the inverse Legendre transformation to the Poincaré equations is degenerate. For this system, Marsden and Morrison (in progress) have a scheme for understanding this phenomenon, using a Tulczyjew's generalized Legendre transformation and the Dirac theory of constraints to arrive at the Low variational principle. It would be of interest to extend this to the Maxwell-Vlasov equations using the ideas proposed here of the Euler-Lagrange-Poincaré equations.

We shall also be investigating the link between the geometry of the Euler-Lagrange-Poincaré equations and those of nonholonomic constrained mechanical systems. These systems, characterized by rolling and grasping constraints are basic in robotics and other areas of control theory. This part of the proposal is a result of joint discussions and work in progress with Richard Murray, Tony Bloch, and P.S. Krishnaprasad. It was already pointed out by Krishnaprasad and Yang and Bloch and Crouch, how the Routh reduction work of Marsden and Scheurle described above is related to nonholonomic systems. Angular momentum constraints can be called *dynamic non-holonomic constraints* since they are preserved by the basic Euler-Lagrange or Hamilton equations and reduction theory (either Lagrangian or Hamiltonian) is a well developed method for dealing with them.

On the other hand, *kinematic non-holonomic constraints* are those imposed by the kinematic set up, such as rolling constraints. Often these constraints are given by linear constraints on the velocity. An example is a wheelbarrow or a car. Some systems involve dynamic and kinematic nonholonomic constraints; an example is a penny rolling on a turntable (angular momentum of the whole system is conserved). One of the goals of the proposal is to investigate problems in control and

optimal control from a unified viewpoint. For example, problems of both car parking and satellite reorientation involve geometric phases. Thus, one would like to utilize the theory developed for dynamic nonholonomic constraints for the kinematic case.

B. C. How this work relates to the DOE mission and other projects. We have worked extensively with Darryl Holm at Los Alamos over the last decade, and will continue this fruitful collaboration in the future. Our work has been useful in a number of basic technological applications, such as the development of improved codes for solid and fluid dynamics, mainly by Juan Simo and his group at Stanford. We have a close ongoing collaboration with him. Also, our control ideas (with Bloch, Krishnaprasad, and Sanchez) have played a useful and influential role in the attitude control of rigid bodies with internal structure (such as internal rotors for attitude control). Another example is the work on polarization laser dynamics and work on ocean dynamics by Holm and collaborators at Los Alamos. His work is directly influenced to a great extent by our ideas, and he has implemented them in a very effective way; this includes ideas on stability (coming out of our joint works), geometric phases, and bifurcations.

D. Project History. This project has been supported in the past largely by DOE, and to smaller extents, by NSF and AFOSR. In particular, Marsden and Weinstein held, for several years, a joint DOE contract, that ended in 1991.

Current and Pending Support

Current Support:

- DOE contract FDDE-FG93-92ER25129 for the period 7/1/92-6/30/93 for \$ 50,001 entitled *Stability, bifurcation, and control of Hamiltonian systems*. This grant is shared with Santa Cruz (35K to Berkeley and 15K to Santa Cruz; supports one summer month and one student at Berkeley).
- NSF grant DMS-89-22704 for the period 5/15/90-4/30/93 awarded for \$ 153,300 (about 51K per year, including overhead) on *Reduction theory in Mechanics and Classical Relativistic Fields* to J. Marsden. Provides summer salary and support for one student.
- NSF grant DMS-91-22708 for the period 5/31/92-5/31/93 awarded for \$ 138,598 (about 46K per year, including overhead) to Ratiu, Lewis and Montgomery.
- Subcontract to USN contract with the University of California-San Diego (Henry Abarbanel) UCSC-FDN00014-92-J-1545 for 4/1/92-12/31/94 awarded for 24K (12K per year) entitled *Symmetric mode truncations in inviscid geophysical fluid dynamics*

Pending Support:

- Proposal in preparation to NSF for renewal of the current proposal above (Marsden)
- Proposal to the AFOSR/DOE as a supplement to the current DOE proposal to support the postdoctoral visit of Hans Peter Kruse (Hamburg) for 6/1/93-5/31/95. (Marsden).
- Proposal to NSF-intelligent control initiative for 10/1/92-9/30/95 entitled *Control, Dynamics, Algorithms, and Mechanical Systems*, joint with Sastry, Bloch, and Brockett.
- Faculty associate on Knobloch's proposal to AFOSR/ARO for 1/1/93-12/31/95 on *Parametric instabilities in mechanical systems* (Marsden).
- The Fields Institute (Marsden). Institute funded by NSERC (the Canadian NSF) and the Province of Ontario for 1.8M. This Institute funding does not support Marsden's own personal research. His term as director will probably end 7/1/93.

5 Scientific and Technical Content

A. Schedule of Major Research Activities. The work on dissipation induced instabilities will be completed in the finite dimensional case by the end of 1992. Work on the infinite dimensional case should see progress by the end of 1993. Work on Hamiltonian bifurcations should see some preliminary papers by the end of 1993. Work on the Euler-Lagrange-Poincaré equations (supported by NSF) will be written up by early in 1993 and work will begin on using it in this DOE project.

B. Scientific and technical issues currently being addressed. One of the main issues to be overcome is, as mentioned above, the extension of the dissipation induced instabilities results to the infinite dimensional case. Here the problem is the infinite spectrum and PDE difficulties (unbounded operators, for example). However, with our past experience with such problems (eg, our earlier work with Kato and Ball), we expect to be able to overcome these. Once the Euler-Lagrange-Poincaré equations are finished, we will face a significant job in properly relating it to the block diagonalization and stability work, and to the separation of internal and rotational modes, but we think this is possible, and will lead to new insight. We also expect to have spin-offs applicable to our NSF work on classical relativistic fields, such as relativistic fluids, plasmas, and elasticity.

C. Experimental and Theoretical Approach. We are in good contact with people doing numerical implementation and simulation of these ideas, such as Simo (Stanford) and Krishnaprasad. Also, we consult with people with robotic experimental facilities that feed back to the theory, such as Krishnaprasad (Maryland), Sastry (Berkeley), and Murray (Caltech). Our theoretical approach is based on geometric analysis, dynamical systems, and nonlinear functional analysis.

D. Importance of Solving the Problem. Our research is fundamental in the sense that it has, and should continue to interact with concrete engineering applications that help these applications in a basic way. For example, our theory has played a key role in the fundamental improvement of numerical codes for rods, shells, and fluids by Simo. It has also been an important ingredient in the robotics work of Krishnaprasad and the work of many others. It is crucial to keep supporting this fundamental mathematical work, or else the important spin-offs to these applications would cease. The time scale for this direct influence is varied, but over the years, has shortened, as geometric mechanics matures.

6 Project Output

A. Major recent accomplishments. Perhaps the nicest recent accomplishment is the work on dissipation induced instabilities proving that the block diagonalization method is sharp. Other major works include the beginnings of a bifurcation theory of Hamiltonian systems with symmetry, with the paper of Marsden, Dellnitz and Melbourne, and that of Lewis, Marsden, Ratiu and Simo on bifurcations in the heavy top.

B. Bibliography of at most 5 relevant publications.

1991 Stability of relative equilibria (Simo, Lewis, Marsden, and Posbergh), *Arch. for Rat. Mech. and An.* **115**, 15-100.

1992 The heavy top: a geometric treatment (Lewis, Marsden, Ratiu, and Simo), *Nonlinearity*, **5**, 1992, 1-48.

1992 Stabilization of rigid body dynamics by internal and external torques, (Bloch, Krishnaprasad, Marsden and Sanchez de Alvarez), *Automatica*, **28**, 745-756.

1993 Lagrangian reduction and bifurcations in the double spherical pendulum (Marsden and Scheurle), *to appear in ZAMP*.

1993 Dissipation induced instabilities, (Bloch, Krishnaprasad, Marsden, and Ratiu), *preprint*.

END

DATE
FILMED

5 / 14 / 93