177
1015-178
LA-7429-MS

Informal Report

16 572

**On Plasma Transport and Flow
in the Fluid Picture**

University of California

LOS ALAMOS SCIENTIFIC LABORATORY

Post Office Box 1663 Los Alamos, New Mexico 87545

LA-7429-MS
Informal Report
UC-20f
Issued: August 1978

On Plasma Transport and Flow in the Fluid Picture

Guthrie Miller

ON PLASMA TRANSPORT AND FLOW IN THE FLUID PICTURE

by

Guthrie Miller

ABSTRACT

The fluid equations are applied to cylindrical pinches, and the role of resistivity, heat conductivity, and viscosity is examined. Analytical solutions for special cases are reviewed. One-dimensional partial differential equations suitable for numerical solution are derived that describe: 1) the radial evolution of a pinch and 2) theta-pinch end loss (for $r=0$).

I. INTRODUCTION

The fluid model offers the simplest means for a quantitative description of plasma behavior. In this report the fluid model is applied to cylindrical pinches (mostly to the theta pinch). The complete set of equations is too complicated for a general analytical solution, but in a few special cases this is possible, and the analytical solutions known to the author are collected here. Analytical solutions are heuristically useful and are also useful as check cases for numerical solutions.

The main object in this report has been to derive tractable one-dimensional partial differential equations. Nowadays, with the use of computers, if a problem can be so formulated, it can be regarded as solved. The other object was to review the implications of the fluid picture, including the role of transport coefficients. Viscosity, in particular, is rather

complicated and this has led to a tendency to ignore it. In this report two cases where classical viscosity is of importance but that do not seem widely known are pointed out. These are: 1) for flow in the z direction with long mean free path (end loss or $m=0$ oscillations) and 2) for long wavelength $m > 2$ flow. For the second case it is shown that classical viscosity gives "finite-Larmor-radius" stabilization identical to that obtained from Vlasov-fluid theory.

II. FLUID MODEL

Starting with the fluid equations for a simple $Z = 1$ plasma as given, for example, by Braginskii,¹ the quasi-neutral approximation $n_e = n_i = n$ is used, and the electron inertia and viscosity terms are neglected. Because $\mathbf{v}_e = \mathbf{v}_i - \mathbf{j}/(en)$, where $\mathbf{j} = \nabla \times \mathbf{B}$, there are then nine variables: n , $\mathbf{v} = \mathbf{v}_i$, s_e , s_i (entropies per particle for electrons and ions), and $\mathbf{B}$ (magnetic field). These are governed by the following nine equations.

$$\left. \frac{\partial n}{\partial t} \right|_{r_0} = -n \nabla \cdot \mathbf{v} \quad , \quad (1)$$

$$m n \left. \frac{\partial \mathbf{v}}{\partial t} \right|_{r_0} = -\nabla p - \text{div } \Pi + \mathbf{j} \times \mathbf{B} \quad , \quad (2-4)$$

$$n T_i \left. \frac{\partial s_i}{\partial t} \right|_{r_0} = \text{div} (\kappa_i \nabla T_i) + \frac{3}{2} n \frac{T_e - T_i}{\tau_{eq}} - \Pi_{\alpha\beta} \frac{\partial v_\alpha}{\partial r_\beta} \quad , \quad (5)$$

$$\begin{aligned} n T_e \left. \frac{\partial s_e}{\partial t} \right|_{r_0} &= \text{div} (\kappa_e \nabla T_e) + \frac{3}{2} n \frac{T_i - T_e}{\tau_{eq}} \\ &+ \frac{T_e}{e} \mathbf{j} \cdot \nabla s_e + \mathbf{j} \cdot \sigma^{-1} \mathbf{j} \quad , \end{aligned} \quad (6)$$

$$n \frac{\partial}{\partial t} \left(\frac{B}{n} \right)_{r_0} = B \cdot \nabla v - \nabla \times \left\{ \sigma^{-1} j + \frac{j \times B - \nabla p_e}{en} \right\} \quad (7-9)$$

Equations (7-9) are obtained from the equation of motion of the electron fluid by solving for the electric field E , taking the curl, and using $\nabla \times E = -\partial B / \partial t$.

The time derivatives are taken with the variable r_0 fixed, where r_0 is the initial coordinate of a fluid element. The relation between r_0 and r is as follows:

$$r = r_0 + \int v dt,$$

$$\frac{\partial}{\partial t} \Big|_{r_0} = \frac{\partial}{\partial t} \Big|_r + v \cdot \nabla \quad (10)$$

The reason for using r_0 as a variable is that Eq. (1) is automatically satisfied for $n = n_0 (dV_0/dV)$, where n_0 is the initial density and (dV_0/dV) the Jacobian of the transformation between r_0 and r . This can be seen mathematically as follows:

$$\begin{aligned} J &\equiv \frac{dV}{dV_0} \equiv \frac{\partial(x, y, z)}{\partial(x_0, y_0, z_0)} \quad , \\ \frac{\partial J}{\partial t} \Big|_{r_0} &= \frac{\partial(v_x, y, z)}{\partial(x_0, y_0, z_0)} + \frac{\partial(x, v_y, z)}{\partial(x_0, y_0, z_0)} + \frac{\partial(x, y, v_z)}{\partial(x_0, y_0, z_0)} \\ &= \left(\frac{\partial v_x}{\partial x} + \frac{\partial v_y}{\partial y} + \frac{\partial v_z}{\partial z} \right) \frac{\partial(x, y, z)}{\partial(x_0, y_0, z_0)} = \nabla \cdot v J \quad . \end{aligned}$$

Therefore,

$$\frac{\partial n}{\partial t} \Big|_{r_0} = \frac{\partial}{\partial t} \left(\frac{n_0}{J} \right)_{r_0} = - \frac{n_0}{J^2} \frac{\partial J}{\partial t} \Big|_{r_0} = - n \nabla \cdot v \quad .$$

The electron and ion pressure and temperature are obtained from the entropy per particle using the perfect gas relation $T = [n \exp(s)]^{2/3}$. Braginskii's notation for the various transport coefficients is followed with the exception that $\tau_{eq} \equiv M\tau_e/(2\pi)$. Also note that κ_i , κ_e , Π , and σ^{-1} are tensor quantities depending on the magnetic field direction. For simplicity, the thermal force and additional terms in the electron and ion heat fluxes are ignored.

The strict conditions for validity of these equations are $t > \tau_e, \tau_i$, where t is a characteristic time, and $L > \lambda_e, \lambda_i$, where L is a characteristic length. However, the fluid equations are used in practice (perhaps for good reason) when these conditions are violated.

III. RADIAL EVOLUTION OF A PINCH

In this section configurations having a dependence only on r and t are considered. These configurations are described by $n(r)$, $v_r(r)$, $s_i(r)$, $s_e(r)$, $B_z(r)$, and $B_\theta(r)$. The case with $v_\theta(r) \neq 0$ is taken up briefly at the end of this section. The configuration is assumed basically to be in equilibrium (except for slow changes of the applied fields and transport processes) and therefore the acceleration terms in Eqs. (2-4) are dropped. The equations giving the time evolution of the system are as follows:

$$nT_i \left. \frac{\partial s_i}{\partial t} \right|_{r_0} = \frac{1}{r} \frac{\partial}{\partial r} \left(r\kappa_i \frac{\partial T_i}{\partial r} \right) + \frac{3}{2} n \frac{T_e - T_i}{\tau_{eq}}, \quad (11)$$

$$\begin{aligned} nT_e \left. \frac{\partial s_e}{\partial t} \right|_{r_0} = & \frac{1}{r} \frac{\partial}{\partial r} \left(r\kappa_e \frac{\partial T_e}{\partial r} \right) + \frac{3}{2} n \frac{T_i - T_e}{\tau_{eq}} \\ & + \left[\frac{1}{r} \frac{\partial}{\partial r} (rB_\theta) \right]^2 \left(\frac{B_z^2}{B^2 \sigma_{\parallel}} + \frac{B_\theta^2}{B^2 \sigma_{\perp}} \right) \end{aligned}$$

$$\begin{aligned}
& + \left(\frac{\partial B_z}{\partial r} \right)^2 \left(\frac{B_z^2}{B^2 \sigma_{\perp}} + \frac{B_{\theta}^2}{B^2 \sigma_{\parallel}} \right) \\
& - \frac{2}{r} \frac{\partial}{\partial r} (r B_{\theta}) \frac{\partial B_z}{\partial r} \frac{B_{\theta} B_z}{B^2} \left(\frac{1}{\sigma_{\parallel}} - \frac{1}{\sigma_{\perp}} \right) \quad , \quad (12)
\end{aligned}$$

$$\begin{aligned}
\frac{\partial}{\partial t} \left(\frac{B_z}{n} \right)_{r_0} &= \frac{1}{nr} \frac{\partial}{\partial r} \left[r \frac{\partial B_z}{\partial r} \left(\frac{B_z^2}{B^2 \sigma_{\perp}} + \frac{B_{\theta}^2}{B^2 \sigma_{\parallel}} \right) \right. \\
& \left. - \frac{\partial}{\partial r} (r B_{\theta}) \frac{B_{\theta} B_z}{B^2} \left(\frac{1}{\sigma_{\parallel}} - \frac{1}{\sigma_{\perp}} \right) \right] \quad , \quad (13)
\end{aligned}$$

$$\begin{aligned}
\frac{\partial}{\partial t} \left(\frac{B_{\theta}}{nr} \right)_{r_0} &= \frac{1}{nr} \frac{\partial}{\partial r} \left[\frac{1}{r} \frac{\partial}{\partial r} (r B_{\theta}) \left(\frac{B_z^2}{B^2 \sigma_{\parallel}} + \frac{B_{\theta}^2}{B^2 \sigma_{\perp}} \right) \right. \\
& \left. - \frac{\partial B_z}{\partial r} \frac{B_{\theta} B_z}{B^2} \left(\frac{1}{\sigma_{\parallel}} - \frac{1}{\sigma_{\perp}} \right) \right] \quad . \quad (14)
\end{aligned}$$

The equilibrium relation from Eqs. (2-4) is

$$\frac{\partial}{\partial r} \left(p + \frac{B^2}{2} \right) + \frac{B_{\theta}^2}{r} = 0 \quad .$$

At any instant of time this equation and the continuity relation, $\partial r^2 / \partial r_0^2 = n_0 / n$, are two equations that allow $r(r_0)$ and $n(r)$ to be determined from the known values of s_i , s_e , B_z / n , and $B_{\theta} / (nr)$.

For the low-pressure case, $p \ll B^2$, with $\sigma_{\perp} = \sigma_{\parallel} = \text{const}$, there is an interesting analytic solution of Eqs. (13) and (14) given by

$$\begin{aligned} B_z &= B_0 J_0(kr) \exp(-k^2 t / \sigma) , \\ B_\theta &= B_0 J_1(kr) \exp(-k^2 t / \sigma) , \end{aligned} \quad (15)$$

with $n(r)$ constant in time.

Only the case $B_\theta = 0$ (theta pinch) will be considered further. In this case the equilibrium relation is

$$p + \frac{B^2}{2} = \frac{B_e^2}{2} , \quad (16)$$

where B_e is the external magnetic field far from the plasma. Equation (16) by itself determines $n(r_0)$, with $r(r_0)$ then obtained using $\partial r^2 / \partial r_0^2 = n_0 / n$.

The simplest special case of Eqs.(11-14) is when all terms on the right-hand sides are small (e.g., for high temperatures). In this case s_i , s_e , and B_z/n are constants on-axis, since $v_r=0$ for $r=0$. The conditions on-axis during adiabatic compression or expansion are determined from Eq. (16),

$$\beta_0 \left(\frac{n}{n_0} \right)^{5/3} + (1-\beta_0) \left(\frac{n}{n_0} \right)^2 = \left(\frac{B_e}{B_0} \right)^2 ,$$

where β_0 is the initial beta on-axis, n_0 the initial density on-axis, and B_0 the initial applied magnetic field.

It is useful to make order-of-magnitude estimates for the various transport processes described on the right-hand sides of Eqs.(11-14). The electrons and ions cool by radial heat conduction in times $t \approx n a^2 / \kappa \sim B_a^2 T^{1/2} / (n m^{1/2})$. * These times become infinite if the plasma is surrounded by a vacuum region ($n = 0$). Also, ion heat conduction is a factor $(M/m)^{1/2}$ more important than electron heat conduction. The electron and ion temperatures approach a common value in a time $t \approx \tau_{eq} \sim T_e^{3/2}$. Finite

* $\omega \tau \gg 1$ is assumed throughout this report.

conductivity apparently causes field diffusion with a characteristic time $t \approx a^2 \sigma_{\perp} \sim a^2 T_e^{3/2}$.

To more carefully examine diffusion, a useful first approximation is to assume that the conductivity is constant. This corresponds to a flat initial T_e profile and short times so that resistive heating is small. The latter requirement is not restrictive since the time for resistive heating is the same as the time for the plasma to diffuse away. The time t for appreciable heating is determined by $j^2 t / \sigma \approx p$, where $jB \approx p/a$ so that $t \approx \sigma a^2 B^2 / p$, which is the diffusion time (B is the external field), as will be seen later.

For constant temperature the diffusion process is described by the simple equation,

$$\frac{\partial}{\partial t} \left(\frac{B_z}{n} \right)_{r_0} = \frac{1}{\sigma n r} \frac{\partial}{\partial r} \left(r \frac{\partial B_z}{\partial r} \right).$$

A further fairly good approximation, due to the fact that $v_r = 0$ at $r = 0$, is to take $\partial/\partial t|_{r_0} \approx \partial/\partial t|r$. There are two simple limiting cases: 1) high β , where $n \approx \text{const}$, which gives

$$\frac{\partial B_z}{\partial t} = \frac{1}{\sigma r} \frac{\partial}{\partial r} \left(r \frac{\partial B_z}{\partial r} \right)$$

and a characteristic diffusion time σa^2 , where a is the plasma characteristic radius, and 2) low β , where $B \approx \text{const}$, which gives (using $\partial B/\partial r = -1/B \partial p/\partial r$)

$$\frac{\partial n}{\partial t} = \frac{p}{B^2 \sigma r} \frac{\partial}{\partial r} \left(r \frac{\partial n}{\partial r} \right),$$

and a characteristic diffusion time $\sigma a^2 B^2 / p$. This time with B the external field applies for both high and low beta since in the high-beta case $B^2/p \approx 1$.

Order-of-magnitude estimates can be given for the importance on the diffusion time scale of the terms neglected in Eqs. (7-9). These terms are 1) electron inertia $\approx (m/e^2 n)/a^2 (p/B^2)$, 2) electron viscosity $\approx (m/e^2 n)/a^2 (n T_e/B^2)$,

and 3) thermal force $\approx nT_e/p$. Although not clear from this rough estimate, the thermal force is not of importance in theta pinches with $T_e < T_i$ and moderate temperature gradients.

Finally, if $v_\theta \neq 0$ the r component of Eqs.(2-4) becomes

$$\frac{\partial}{\partial r} \left(p + \frac{B^2}{2} \right) - \frac{Mn v_\theta^2}{r} = 0 \quad .$$

The centrifugal force term is only significant if $Mn v_\theta^2 \approx p$ or $v_\theta \approx$ thermal speed, which means that the ion Larmor orbits are significantly displaced during one period. Rotation-driven instabilities become important for smaller rotation speeds² so this term is usually dropped.

The θ -component of Eqs. (2-4), which describes the viscous damping of the ion rotation, is

$$-Mn \frac{\partial v_\theta}{\partial t} = \frac{1}{r^2} \frac{\partial}{\partial r} (r^2 \Pi_{r\theta}) \quad .$$

From Braginskii,¹ the r, θ component of the viscous stress tensor is given by

$$\Pi_{r\theta} = -\eta_1 \left(\frac{\partial v_\theta}{\partial r} - \frac{v_\theta}{r} \right) + \eta_3 \left(\frac{\partial v_r}{\partial r} - \frac{v_r}{r} \right) \quad .$$

The effect of the η_3 viscosity component (gyro-viscosity) will be discussed later on, and if this term is simply ignored there results an equation for v_θ ,

$$Mn \frac{\partial}{\partial t} \left(\frac{v_\theta}{r} \right) = \frac{1}{r^3} \frac{\partial}{\partial r} \left[\eta_1 r^3 \frac{\partial}{\partial r} \left(\frac{v_\theta}{r} \right) \right] \quad .$$

An order-of-magnitude estimate for the time for the ion velocity distribution to decay to a rigid rotation is

$$t = \frac{10}{3} \frac{M n a^2}{\eta_1} = \frac{\omega_1^2 a^2 M \tau_1}{T_1} \sim B^2 a^2 T^{1/2} / (n M^{1/2})$$

IV. LONGITUDINAL FLOW AND TRANSPORT

The main confinement limitation for the θ pinch is flow out the ends due to lack of equilibrium. Longitudinal heat conduction may also be of importance (e.g., with end plugs). The situation is actually two-dimensional (z and r) but useful one-dimensional equations can be obtained by neglecting radial transport and considering only the axis of the system ($r = 0$). These equations are as follows:

$$M n \left. \frac{\partial^2 z}{\partial t^2} \right|_{z_0} = - \frac{\partial p}{\partial z} + \frac{4}{3} \frac{\partial}{\partial z} \left(\eta_0 \frac{\partial v_z}{\partial z} \right) \quad , \quad (17)$$

$$n T_1 \left. \frac{\partial s_1}{\partial t} \right|_{z_0} = \frac{\partial}{\partial z} \left(\kappa_1 \frac{\partial T_1}{\partial z} \right) + \frac{3}{2} n \frac{T_e - T_1}{\tau_{eq}} + \frac{4}{3} \eta_0 \left(\frac{\partial v_z}{\partial z} \right)^2 \quad , \quad (18)$$

$$n T_e \left. \frac{\partial s_e}{\partial t} \right|_{z_0} = \frac{\partial}{\partial z} \left(\kappa_e \frac{\partial T_e}{\partial z} \right) + \frac{3}{2} n \frac{T_1 - T_e}{\tau_{eq}} \quad . \quad (19)$$

From the continuity equation

$$n = \frac{n_0}{\frac{\partial(z, r^2)}{\partial(z_0, r_0^2)}} \quad , \quad (20)$$

with

$$\frac{\partial(z, r^2)}{\partial(z_0, r_0^2)} = \left(\frac{\partial z}{\partial z_0}\right) \left(\frac{\partial r^2}{\partial r_0^2}\right)$$

for $r=0$. The radial equilibrium relation is needed to determine $\partial r^2/\partial r_0^2$ and close the system,

$$\frac{n_0^{5/3} [\exp(\frac{2}{3}\epsilon_e) + \exp(\frac{2}{3}\epsilon_1)]}{\left(\frac{\partial z}{\partial z_0}\right)^{5/3} \left(\frac{\partial r^2}{\partial r_0^2}\right)^{5/3}} + \frac{B_0^2 (1-\beta_0)}{2 \left(\frac{\partial r^2}{\partial r_0^2}\right)^2} = \frac{B_e^2}{2}, \quad (21)$$

where B_0 is the initial value of the external field B_e , and β_0 the initial beta value on-axis.

Neglecting heat conduction, equilibration of electrons and ions, and ion viscosity, and assuming small amplitude disturbances i.e., $\partial z/\partial z_0 = 1 + \epsilon$, and constant external field, Eq. (17) becomes

$$\frac{\partial^2 \epsilon}{\partial t^2} = - \frac{1}{Mn_0} \frac{\partial^2 p}{\partial z^2} = - \frac{c^2 u_A^2}{c^2 + u_A^2} \frac{\partial^2 \epsilon}{\partial z^2}. \quad (22)$$

The small signal propagation velocity on-axis is thus $v^2 = c^2 u_A^2 / (c^2 + u_A^2)$, where $c^2 = 5p_0 / (3Mn_0)$ and $u_A^2 = (1-\beta_0) B_0^2 / (Mn_0)$. Another analytic result can be obtained for a simple isentropic rarefaction in the low- β limit. In this case $\partial r^2/\partial r_0^2 = 1$ and $n = n_0 \partial z_0/\partial z$. The equation of motion, Eq. (17), is then

$$\left. \frac{\partial^2 z}{\partial t^2} \right|_{z_0} = - \frac{1}{Mn_0} \frac{\partial p}{\partial z_0}.$$

Taking derivatives with respect to z_0 , this becomes

$$\frac{\partial^2}{\partial t^2} \left(\frac{n_0}{n} \right)_{z_0} = \frac{\partial^2}{\partial t^2} \left(\frac{p_0}{p} \right)_{z_0}^{3/5} = - \frac{1}{Mn_0} \frac{\partial^2 p}{\partial z_0^2} .$$

The solution* of this equation for $p|_{t=0}=F(z)$ is

$$p = F(z_0 \pm c_0 \left(\frac{p}{p_0} \right)^{4/5} t) ,$$

where $c_0^2 = 5p_0/(3Mn_0)$. For a situation with the pressure initially constant from $z=0$ to $z=L$ the pressure takes the value p at point z_0 given by

$$z_0 = L - c_0 \left(\frac{p}{p_0} \right)^{4/5} t ;$$

therefore,

$$p = p_0 \left(\frac{L - z_0}{c_0 t} \right)^{5/4} .$$

The complete solution is obtained using

$$z = L - c_0 t + \int_{L - c_0 t}^{z_0} \left(\frac{n_0}{n} \right) dz_0 = z_1 + \int_{z_1}^{z_0} \left(\frac{p_0}{p} \right)^{3/5} dz_0 ,$$

*A solution of the equation $\partial^2 f / \partial x^2 = \alpha \partial^2 f^\lambda / \partial y^2$ is $f = F(y \pm (\alpha \lambda)^{1/2} f^{(\lambda-1)/2} x)$, where F is an arbitrary function. As can be seen $F(y)$ gives the value of $f(x, y)$ for $x = 0$.

where $z_1 \equiv L - c_0 t$, which gives

$$\left(\frac{L-z_0}{c_0 t}\right)^{1/4} = 1 - \left(\frac{z-z_1}{4c_0 t}\right),$$

and thus

$$p = \begin{cases} p_0, & 0 < z < z_1, \\ p_0 \left[1 - \frac{z-z_1}{4c_0 t}\right]^5, & z > z_1. \end{cases} \quad (23)$$

The propagation velocity of the rarefaction (c_0) agrees with the small signal velocity ($v=c$ for $u_A \rightarrow \infty$).

It is important to note that a quantitative treatment of the flow requires a two-dimensional treatment. Figure 1 gives the radially integrated plasma density histories at various points in a theta pinch of half-length L as calculated by Brackbill for ideal two-dimensional flow. The rarefaction propagation velocity is $v=1.7 v_i$ from Fig. 1 ($T_e=T_i$, $\beta=0.7$, $v_i \equiv (2T_i/M)^{1/2}$) which is quite different from that given by Eq.(22), $v=0.75 v_i$.

Order-of-magnitude estimates of the transport terms in Eqs. (17-19) can be made in a straightforward way. One observation is that the times associated with viscous damping and with ion heat conduction are of the same order and are given by $t \approx L^2/(v_i \lambda_i)$, where λ_i is the ion mean free path and v_i the ion thermal velocity. Thus these processes are of dominating importance in simple end loss ($t < L/v_i$) only for $\lambda_i > L$, which violates the original assumption on which the fluid description is based. Also, with $\lambda_i > L$, the heat and momentum fluxes exceed their kinetic limits nTv_i and nMv_i^2 . The fluid equations are commonly used to describe situations with long mean free path, and in these cases the ion heat flux and momentum flux (perhaps with flux limits applied) are of importance.

There are several analytical solutions for longitudinal heat conduction. The first (DeSilva, Chu, and Johnson) covers the interesting situation $\partial p / \partial z = 0$ and $T_e = T_i$. Neglecting ion heat conductivity, Eqs.(18) and (19) become in the case

$$2nT \frac{\partial s}{\partial t} = \frac{\partial}{\partial z} \left(\kappa_e \frac{\partial T}{\partial z} \right) , \quad (24)$$

where $s = \log(T^{3/2}/n)$, and $\kappa_e = \alpha T^{5/2}$ with $\alpha = \text{const.}$ A solution is

$$T = T_0 [1 - (z/L)^2]^{2/7} f(t) \quad (25)$$

with

$$f(t) = \left[1 + 1.24 \frac{10 \alpha T_0^{5/2}}{21 n_0 L^2} t \right]^{-2/5} .$$

The density profile is constant in time,

$$n(z) = \frac{n_0}{1.24} [1 - (z/L)^2]^{-2/7} ,$$

and satisfies

$$\int_0^L n \, dz = n_0 L .$$

*If the electron-ion equilibration time is long, these formulas with $t \rightarrow 2t$ describe the decay of the electron temperature.

Another solution (Dreicer, Chu, and Johnson) covers the case $n = \text{const}$ and is interesting as an illustration of the small effect of the density profile. For this case Eq.(24) becomes

$$3n \frac{\partial T}{\partial t} = \alpha \frac{\partial}{\partial z} \left(T^{5/2} \frac{\partial T}{\partial z} \right) , \quad (26)$$

which is satisfied by

$$T = T_0 f(z)^{2/7} \left(1 + 2.13 \frac{5 \alpha T_0^{5/2}}{21 n L^2} t \right)^{-2/5} ,$$

where

$$\frac{d^2 f}{dz^2} = - \frac{2.13}{L^2} f^{2/7}$$

and $f(0)=1$, $f(L)=0$. Since $f^{2/7} \approx 1$, $f \approx 1 - (z/L)^2$.

There is a non-separable solution of the electron heat conduction problem which describes the establishment of a temperature profile when a column with initial temperature T_0 is connected to a cold heat sink at one end. This is given by

$$T = T_0 f \left[\left(\frac{z^2 n}{\alpha T_0^{5/2} t} \right)^{1/2} \right] ,$$

where $f(x)$ is determined from the equation

$$\frac{2}{7} \frac{d^2 f^{7/2}}{dx^2} + \frac{3}{4} x \frac{df}{dx} = 0^*$$

with the boundary conditions $f(0) = 0$, $f(\infty) = 1$. The function f is plotted in Fig. 2. The temperature profile for x in the range 0 to 2 closely matches the shape $[1-(L-z)^2/L^2]^{2/7}$ for $z = xL/2$; therefore, the time to set up a steady profile in a column of half-length L is approximately $t \approx L^2 n / (4 \alpha T_0^{5/2})$. Chu and Johnson³ obtain a similar estimate with the 4 in the denominator replaced by 3.7. When thermal losses are important (e.g., T_e large) this time tends to be shorter than the time for mechanical motion or electron-ion equilibration to take place, thus the assumptions of constant density and no equilibration are valid.

It is possible to reduce Eqs.(18-19) to zero dimensions by assuming the temperature profiles are given by Eq. (25). By symmetry, $v_z = 0$ at $z=0$, $\partial/\partial t|_{z=0} = \partial/\partial t|_z$ and therefore, Eqs.(18-19) become ordinary differential equations describing the point $r = 0$, $z = 0$.

V. GYRO-VISCOSITY

In this section the effect of gyro-viscosity ($\eta_3 \neq 0$) on long wavelength small oscillations of a theta pinch is examined. This type of viscosity does not dissipate energy, yet it results in "finite-Larmor-radius" stabilization of $m \geq 2$ modes in a theta pinch as will be seen. Since the energy of motion does not die away with this type of viscosity, the stabilized modes must be purely oscillatory.

First, it will be shown that gyro-viscosity drops out of the equation of motion for small oscillations of a constant-pressure plasma (sharp-boundary model). Therefore, the viscosity enters in only through the boundary conditions at the plasma surface. For long wavelength motion the z -dependence is small and the x and y components of the equations of motion, Eqs. (2-4), contain additional terms owing to viscosity, e.g.,

$$(\text{div } \Pi)_x = \frac{\partial \Pi_{xx}}{\partial x} + \frac{\partial \Pi_{xy}}{\partial y} \quad .$$

*The electron-ion equilibration time is here assumed to be long. If it is short, $f(x) \rightarrow f(2^{1/2}x)$.

Fig. 1.

Line density histories at various points in a theta pinch of half-length L , from a two-dimensional computation by Brackbill assuming adiabatic flow.

Fig. 2.

A plot of the function $f(x)$, where $x = (z^2/(Dt))^{1/2}$, with D the thermal diffusion coefficient, which describes the propagation of a "cooling wave".

Because long wavelength motion is approximately divergenceless potential flow in the sharp-boundary model, i.e., $v = \nabla\phi$, $\nabla^2\phi = 0$, the additional viscosity terms are zero,

$$(\text{div } \Pi)_x = \eta_3 \left(\frac{\partial^2 v_y}{\partial x^2} + \frac{\partial^2 v_y}{\partial y^2} \right) = 0 \quad .$$

The boundary condition with viscosity is

$$\left[p + \frac{B^2}{2} + \Pi_{rr} \right]_{\text{in}} = \left[\frac{B^2}{2} \right]_{\text{out}} \quad , \quad (27)$$

where

$$\Pi_{rr} = - \eta_3 \left(\frac{1}{r} \frac{\partial v_r}{\partial \theta} - \frac{v_\theta}{r} + \frac{\partial v_\theta}{\partial r} \right) \quad . \quad (28)$$

To apply the boundary condition it is convenient to use a complex formalism, where

$$\begin{aligned} \xi_r &= \text{Re} \left\{ \left(\frac{r}{a} \right)^{m-1} \xi_0 \exp[-i(m\theta + \omega t)] \right\} \quad , \\ \xi_\theta &= \text{Re} \left\{ -i \left(\frac{r}{a} \right)^{m-1} \xi_0 \exp[-i(m\theta + \omega t)] \right\} \quad , \end{aligned} \quad (29)$$

with ξ_0 a complex number. This can be seen to represent divergenceless potential flow. The components of the boundary condition, Eq. (27), are as follows.⁴ ($\text{Re}\{\xi_0 \exp[-i(m\theta + \omega t)]\}$ is factored out in each case):

$$\left(p + \frac{B^2}{2}\right)_{in} = \frac{anM\omega^2}{m} ,$$

$$\left(\frac{B^2}{2}\right)_{out} = 0 , \quad (30)$$

while using Eqs.(28) and (29) and $\eta_3 = nT_1/(2\omega_1)$,

$$\pi_{rr} = \frac{(m-1)nT_1}{\omega_1 a} \omega , \quad (31)$$

where $\omega_1 = eB_0(1-\beta)^{1/2}/M$.

The boundary condition, Eq.(27), leads to the dispersion relation

$$\omega^2 + \frac{m(m-1)T_1}{\omega_1 a^2 M} \omega = 0 , \quad (32)$$

which is identical to that obtained by Turner⁵. The viscosity term is roughly equal to $\omega\Omega^*$, where Ω^* is the diamagnetic rotation frequency.

To appreciate the importance of viscosity one must imagine a situation that is otherwise unstable (because of additional magnetic fields or rotation). The dispersion relation is then of the form

$$\omega^2 + \eta\omega + \gamma^2 = 0 ,$$

which has stable (purely oscillatory) roots for $\eta > 2\gamma$.

REFERENCES

1. S. I. Braginskii, "Transport Processes in a Plasma", in Reviews of Plasma Physics, M. A. Leontovich, Ed. (Consultants Bureau, New York, 1965), Vol. I, p. 205.

2. J. P. Freidberg and L. D. Pearlstein, "Rotational Instabilities in a Theta Pinch", Phys. Fluids 20, 1207 (1978).
3. T. K. Chu and L. C. Johnson, "Conduction Heat Loss Scaling in Open Field Line Geometries", Phys. Fluids 20, 1684 (1977).
4. G. Miller, "Motion of a Plasma Column in a Perturbing Magnetic Field", Phys. Fluids 18, 1704 (1975).
5. L. Turner, "Finite-Larmor-Radius Stabilization in a Sharp-Boundary Vlasov-Fluid Screw Pinch", Phys. Fluids 20, 654 (1977).