MARTIN MARIETTA

**ENVIRONMENTAL
RESTORATION
PROGRAM**

**Preliminary Report on the Ecological
Assessment of Waste Area Grouping 5
at Oak Ridge National Laboratory,
Oak Ridge, Tennessee**

**T. L. Ashwood
G. W. Suter II
A. J. Stewart**

**MANAGED BY
MARTIN MARIETTA ENERGY SYSTEMS, INC.
FOR THE UNITED STATES
DEPARTMENT OF ENERGY**

UCN-17580 (6 7-91)

ENERGY SYSTEMS

This report has been reproduced directly from the best available copy.

Available to DOE and DOE contractors from the Office of Scientific and Technical Information, P.O. Box 62, Oak Ridge, TN 37831; prices available from 615-576-8401, FTS 626-8401.

Available to the public from the National Technical Information Service, U.S. Department of Commerce, 5285 Port Royal Rd., Springfield, VA 22161.

Environmental Restoration Division
ORNL Environmental Restoration Program

**Preliminary Report on the Ecological Assessment of Waste Area
Grouping 5 at Oak Ridge National Laboratory,
Oak Ridge, Tennessee**

T. L. Ashwood
G. W. Suter II
A. J. Stewart

Date Issued—September 1992

Prepared by
Environmental Sciences Division
Oak Ridge National Laboratory
ESD Publication 3950

Prepared for
U.S. Department of Energy
Office of Environmental Restoration and Waste Management
under budget and reporting code EW 20

OAK RIDGE NATIONAL LABORATORY
Oak Ridge, Tennessee 37831-6285
managed by
MARTIN MARIETTA ENERGY SYSTEMS, INC.
for the
U.S. DEPARTMENT OF ENERGY
under contract DE-AC05-84OR21400

MASTER

JP

Author Affiliations

T. L. Ashwood, G. W. Suter II, and A. J. Stewart are members of the Environmental Sciences Division, Oak Ridge National Laboratory, Martin Marietta Energy Systems, Inc.

CONTENTS

FIGURE	v
TABLES	v
ACRONYMS	vii
EXECUTIVE SUMMARY	ix
1. INTRODUCTION	1
1.1 BACKGROUND	1
1.2 SUMMARY OF ECOLOGICAL ASSESSMENT PLAN	1
2. SCREENING-LEVEL ECOLOGICAL RISK ASSESSMENT	4
2.1 ECOLOGICAL HAZARD DEFINITION	4
2.1.1 Sources and Routes of Exposure	4
2.1.2 Exposed Environment	5
2.2 ECOLOGICAL ENDPOINTS	7
2.3 EXPOSURE ASSESSMENT	7
2.3.1 Water	7
2.3.2 Soil	7
2.3.3 Sediments	8
2.4 EFFECTS ASSESSMENT	8
2.5 RISK CHARACTERIZATION	8
2.5.1 Aquatic Risks	8
2.5.2 Terrestrial Risks	9
2.6 DATA AND ASSESSMENT NEEDS	10
2.6.1 Biological Sampling, Analysis, and Testing	10
2.6.2 Baseline Assessment Strategy	10
3. RARE AND ENDANGERED SPECIES AND WETLANDS	12
4. AMBIENT TOXICITY TESTING	13
4.1 MATERIALS AND METHODS	13
4.2 RESULTS AND DISCUSSION	14
5. WILD TURKEY CONTAMINANT SCREENING	18
5.1 METHODS	18
5.1.1 Trapping	18
5.1.2 Radiological Screening	19
5.2 RESULTS AND DISCUSSION	19
6. FUTURE ACTIVITIES	21
7. ACKNOWLEDGEMENTS	22

8. REFERENCES	23
APPENDIX—ENDANGERED SPECIES AND WETLANDS SURVEY REPORTS	25

FIGURE

1	Waste Area Grouping 5 and environs, including ambient toxicity testing locations	2
---	--	---

TABLES

1	Federal and state listed threatened and/or endangered species and species designated "in need of management" by the State of Tennessee known or expected to occur on the Oak Ridge Reservation	6
2	Results of <i>Ceriodaphnia</i> toxicity tests of WAG 5 water samples during test period 1 (December 12-18, 1991) and test period 2 (February 27-March 5, 1992)	14
3	Results of <i>Ceriodaphnia</i> toxicity test of water from WAG 5 site 6 (April 30-May 7, 1992)	15
4	Summary of results of water quality analyses, WAG 5 sites, for toxicity tests conducted during December 12-18, 1991	16
5	Summary of results of water quality analyses, WAG 5 sites, for toxicity tests conducted during February 27-March 5, 1992	16
6	Summary of results of water quality analyses, WAG 5 site 6, for the toxicity tests conducted during April 30-May 7, 1992	17
7	Radiological screening results for wild turkeys	20

ACRONYMS

BMAP	Biological Monitoring and Abatement Program
BNI	Bechtel National, Inc.
EPA	U.S. Environmental Protection Agency
ESD	Environmental Sciences Division
LLW	low-level waste
NBS	National Bureau of Standards
NPDES	National Pollutant Discharge Elimination System
OHF	Old Hydrofracture Facility
ORNL	Oak Ridge National Laboratory
ORR	Oak Ridge Reservation
PCB	polychlorinated biphenyl
RI	remedial investigation
SOP	standard operating procedure
SWSA	solid waste storage area
SWMU	solid waste management unit
TWRA	Tennessee Wildlife Resources Agency
WAG	waste area grouping
WOC	White Oak Creek

EXECUTIVE SUMMARY

In support of the remedial investigation for Waste Area Grouping (WAG) 5, staff of the Environmental Sciences Division of Oak Ridge National Laboratory have conducted preliminary ecological assessment activities. A screening-level ecological risk assessment has been completed, ambient toxicity tests have been conducted on streams and seeps within WAG 5, WAG 5 has been surveyed for rare and endangered species and wetlands, and wild turkeys that may feed on contaminated vegetation and insects in WAG 5 have been screened for beta-emitting isotopes and ^{137}Cs .

The screening-level ecological risk assessment identified some data gaps that were addressed in the ecological assessment plan. These include gaps in data on the toxicity of surface water and soil within WAG 5 and on the status of rare and endangered species. In addition, the screening-level risk assessment identified the need for data on the level of contaminants in wild turkeys that may be consumed by predatory wildlife and humans.

Three rounds of ambient toxicity tests on six streams and seeps, using the microcrustacean *Ceriodaphnia*, have identified potential toxicity in three of the sample sites. Further tests are required to identify the toxicant.

No rare or endangered animal species have been identified in the WAG 5 area.

Limited success in trapping wild turkeys during the winter of 1992 has resulted in an inability to draw strong conclusions about contaminant levels in these animals. Three hens thought to be part of a flock that occasionally feeds in WAG 5 were collected and screened for beta-emitting isotopes and ^{137}Cs . No birds contained measurable levels of ^{137}Cs , and only one bird contained beta radiation significantly above background.

Data collection activities will continue in FY 1993 to fill data gaps identified in the screening-level risk assessment. Specifically, soil toxicity tests will be conducted, and additional ambient toxicity tests will be conducted on streams and seeps. If feasible, additional turkey samples will be collected.

1. INTRODUCTION

This report presents a summary of ecological risk assessment activities for Waste Area Grouping (WAG) 5 conducted by staff of the Environmental Sciences Division (ESD) of Oak Ridge National Laboratory (ORNL) during FY 1992. These activities were conducted in support of the baseline ecological risk assessment to be conducted by the Bechtel National, Inc. (BNI) team that is performing the remedial investigation (RI) for WAG 5. ESD's role in support of the BNI team is defined in the ecological assessment plan (Ashwood 1992).

1.1 BACKGROUND

WAG 5 is one of the groups of solid waste management units (SWMUs) defined by ORNL for purposes of remedial action planning. Wag 5 consists of 13 SWMUs covering an area of ~20 ha, >75% of which is included in Solid Waste Storage Area (SWSA) 5 (Fig. 1). Most of the area is covered by grass, but deciduous forests fringe SWSA 5. Although White Oak Creek (WOC) and Melton Branch border the SWMUs within WAG 5, these streams and most of their floodplains lie outside the defined boundaries of WAG 5. Surface water resources within the WAG include two surface impoundments [the Old Hydrofracture Facility (OHF) pond and the process waste sludge basin] and intermittent drainages D1 and D2 (Fig. 1).

From 1959 to 1973, SWSA 5 was used for disposal of solid low-level waste (LLW) and some transuranic wastes. The portion of SWSA 5 known as SWSA 5 North has been used since 1970 for storage of transuranic wastes. Liquid LLW was disposed of in both hydrofracture facilities (Fig. 1) and is currently stored in the LLW concentrate tanks (Fig. 1). Thus, the primary radionuclides present in WAG 5 are transuranic and LLW constituents (e.g., ^3H , ^{60}Co , ^{90}Sr , ^{137}Cs , $^{233,235}\text{U}$, $^{238,239}\text{Pu}$, ^{241}Am , and ^{244}Cm). In addition, metals (e.g., lead, mercury, and selenium) and some organics [e.g., chlorinated solvents and polychlorinated biphenyls (PCBs)] are probably present, although data on these contaminants is limited.

1.2 SUMMARY OF ECOLOGICAL ASSESSMENT PLAN

The ecological assessment plan (Ashwood 1992) defines six tasks that are required to provide sufficient data to conduct a baseline ecological risk assessment. The first task is a screening-level ecological risk assessment to verify the need for the remaining five tasks and to identify any additional data needs. Results of the screening-level risk assessment, conducted in December 1991, are presented in Sect. 2. of this report.

The second task consists of rare and endangered species surveys and wetlands surveys. Results of these surveys are reported in Sect. 3.

Fig. 1. Waste Area Grouping 5 and environs, including ambient toxicity testing locations.

Task 3 is a series of ambient toxicity tests on water from the two impoundments, the surface streams, and those seeps with sufficient flow to provide a sample (Fig. 1). Three rounds of toxicity tests have been conducted, and the results are included in Sect. 4 of this report.

Task 4 is a set of toxicity tests on contaminated soil from WAG 5. The soil is to be collected at the same time and locations and times as soil samples collected by BNI for measurement of soil contamination. At the time of this report (July 1992), the BNI team have not yet collected their soil samples, so the soil toxicity tests have not been conducted.

Task 5 consists of collecting samples of wild turkeys that forage on WAG 5 to determine whether the turkeys are significantly contaminated. A limited number of turkeys have been sampled at the time of this report, and screening results are reported in Sect. 5.

2. SCREENING-LEVEL ECOLOGICAL RISK ASSESSMENT

The purpose of this screening-level ecological risk assessment is to serve as a preliminary hazard identification for the baseline risk assessment for WAG 5. Toward that end it will, as far as possible, identify ecological hazards associated with wastes in WAG 5, prioritize them with respect to their potential risks, and identify data needs on the basis of that analysis. The ecological assessment is concerned with risks to populations and communities of nonhuman organisms that occur on the site or may occur there in the future. A separate screening assessment addresses risks to human health posed by WAG 5. Ecological risks were not a component of the original RI plan for WAG 5 (Bechtel 1988), but are being incorporated into the remedial investigation through a separate ecological assessment plan (Ashwood 1992).

Although radionuclides are the principal contaminants of concern at WAG 5 with respect to human health, they are not emphasized in this screening assessment for two reasons. First, unlike chemical contaminants, they do not have modes of action in nonhuman organisms that differ from those that are treated in human health risk assessments. Second, they are the primary focus of the RI because of human health concerns and could not conceivably be screened out of further consideration by this ecological assessment. Third, extensive analysis of media for radionuclides was planned at WAG 5 before the preliminary assessment and other planning activities for the ecological portion of the sampling and analysis plan were begun. Therefore, they are left to the baseline ecological assessment for which much better radionuclide exposure data will be available.

This assessment is organized in the format of a standard risk assessment with a hazard definition phase, followed by parallel exposure and effects assessments, and ending with risk characterization. The results are used to draw conclusions about data needs and assessment needs for the WAG 5 baseline risk assessment.

2.1 ECOLOGICAL HAZARD DEFINITION

The purpose of the hazard definition phase of a risk assessment is to define the scope of the assessment, the sources being assessed, and the endpoints of the assessment. The combination of a source of potentially toxic exposure with a valued and potentially susceptible receptor (the endpoint organisms) in a particular environment constitutes a hazard.

2.1.1 Sources and Routes of Exposure

The ultimate sources of contamination on WAG 5 are radioactive and mixed wastes buried or stored on the site, the wastes generated by the OHF, and any spills or releases of incidental materials such as oils or pesticides. Except for radionuclides in some specific facilities such as the transuranic storage facilities and the OHF pond, these sources are poorly characterized (Shoun 1987, Bechtel 1988).

The proximate sources of contamination on WAG 5 include the buried wastes, contaminated soil and water, water and sediments in waste ponds, and contaminated plants or animals that may be consumed by other animals. Potential routes of exposure include

direct ingestion of soil or water in ponds and seeps, external exposure to wastes or contaminated soil, uptake by plant roots from the waste or contaminated soil, and consumption of contaminated plants or animals.

Organisms and ecosystems near WAG 5 may also be exposed to contaminants from WAG 5. Contaminated water and silt may enter WAG 2 from WAG 5 runoff and leachate. In addition, organisms may become contaminated on WAG 5 and subsequently move off the WAG.

2.1.2 Exposed Environment

WAG 5 is a waste disposal area located on the Oak Ridge Reservation (ORR) south of ORNL. The ecology of this area is described in Loar (1987-1991), Loar et al. (1981), Boyle et al. (1982), Mann and Kitchings (1982), Oakes et al. (1982), and Sherwood and Loar (1987). WAG 5 is largely a mowed lawn with other areas consisting of roads and structures related to waste management. It includes small wooded areas along the periphery, between facilities, and along drainages that could support some small birds and mammals. It is also used incidentally by the wildlife of the surrounding forest. Although a formal wildlife survey of the site has not been conducted, informal observations and surveys of similar nearby waste sites suggest that there are few resident amphibians, reptiles, mammals, or nesting birds on the site.

The scope of this assessment is limited to the waste site. The stream ecosystems receiving runoff and leachate from WAG 5 (WOC and Melton Branch) and the floodplains of those streams are within WAG 2, which is being assessed separately. However, the WAG 5 assessment will consider the contribution of WAG 5 to contamination of WAG 2 and environmental data from WAG 2 will be used to evaluate the potential significance of those contributions. Similarly, the contribution of WAG 5 to the exposure experienced by the terrestrial ecosystem that extends across the reservation will be estimated, but risks to that ecosystem will be assessed in the reservation-wide risk assessment.

There are no aquatic communities on WAG 5. The drainages on the site are ephemeral, and the only permanent surface water is in waste ponds. (Although there are undoubtedly some hearty organisms—particularly microbes—in the waste ponds, the ponds are not considered to be part of the natural environment that is subject to assessment and protection.)

Although several federally-listed threatened or endangered species occur in the region (Kroodsma 1987), none are known to occur on WAG 5. However, the endangered Indiana bat (*Myotis sodalis*) is suspected to occur on the lower East Fork Poplar Creek, and therefore may occur in the area of WAG 5. Several state-listed species occur on the ORR and may occur on WAG 5 (Table 1).

Wetlands and floodplains are protected by Executive Orders 11990 and 11988, respectively. No wetlands survey has been conducted for WAG 5. WAG 5 lies above the 100-year floodplains of WOC and Melton Branch.

Table 1. Federal and state listed threatened and/or endangered species and species designated "in need of management" by the State of Tennessee known or expected to occur on the Oak Ridge Reservation

Common Name	Scientific Name	Administrative Status ^a
Fish		
Tennessee dace	<i>Phonixus oreas</i>	INM
Birds		
Black-crowned night heron	<i>Nycticorax nycticorax</i>	INM
Black vulture	<i>Coragyps atratus</i>	INM
Common barn owl	<i>Tyto alba</i>	INM
Cooper's hawk	<i>Accipiter cooperii</i>	ST
Grasshopper sparrow	<i>Ammodramus savannarum</i>	ST
Osprey	<i>Pandion haliaetus</i>	SE
Red-shouldered hawk	<i>Buteo lineatus</i>	ST
Sharp-shinned hawk	<i>Accipiter striatus</i>	ST
Mammals		
Indiana bat	<i>Myotis sodalis</i>	FE, SE
Plants		
Spreading false foxglove	<i>Aureolaria patula</i>	ST
Appalachian bugbane	<i>Cimicifuga rubifolia</i>	ST
Pink lady's slipper	<i>Cypripedium acaule</i>	SE
Tall larkspur	<i>Delphinium exaltatum</i>	SE
Northern bush honeysuckle	<i>Diervilla lonicera</i>	ST
Nuttall waterweed	<i>Elodea nuttallii</i>	SS
Mountain witch-alder	<i>Fothergilla major</i>	ST
Golden seal	<i>Hydrastis canadensis</i>	ST
Butternut	<i>Juglans cinerea</i>	ST
Canada lily	<i>Lilium canadense</i>	ST
Michigan lily	<i>Lilium michiganense</i>	ST
Fen orchid	<i>Liparis loeselii</i>	SE
Ginseng	<i>Panax quinquefolium</i>	ST
Tuberclad rein orchid	<i>Platanthera flava var herbiola</i>	ST
Purple fringeless orchid	<i>Platanthera peranoena</i>	ST
Carey saxifrage	<i>Saxifraga careyana</i>	SS
Lesser ladies' tresses	<i>Spiranthes ovalis</i>	SS

^aFE = Federally Endangered; FT = Federally Threatened; SE = State Endangered; ST = State Threatened; SS = State Special Concern; INM = In Need of Management according to the State of Tennessee

2.2 ECOLOGICAL ENDPOINTS

The endpoints for this assessment are those described and justified in the *Approach and Strategy for Performing Ecological Risk Assessments for the Department of Energy Oak Ridge Field Office Environmental Restoration Program* (Suter et al. 1992). In brief they are (1) no reduction greater than 20% in the abundance or productivity of populations of fishes, birds, mammals, or vascular plants, (2) no injury of individuals of threatened or endangered species sufficient to impair survival or reproduction, and (3) no loss of wetland or floodplain communities. The 20% figure in the first endpoint is derived from an analysis of Environmental Protection Agency (EPA) regulatory criteria and is intended to approximate the agency's *de facto* but unstated level of protection (Suter et al. 1992). No particular species are chosen to represent the endpoints because most-sensitive species cannot be identified *a priori*, and in most cases the toxicity and exposure data do not permit discrimination among individual members of taxa or trophic groups.

2.3 EXPOSURE ASSESSMENT

2.3.1 Water

Maximum aqueous exposures to WAG 5 contaminants would occur in seeps or at the mouths of ephemeral drainages flowing from WAG 5 into Melton Branch or WOC (WAG 2). These sources have not been characterized with respect to nonradionuclide contamination. However, data are available for the receiving streams (Blaylock et al. 1992).

Fish and invertebrates inhabiting the water column in WAG 2 are assumed to be exposed to chemicals in the water primarily through respiratory uptake. This assumption is accurate for nearly all chemicals except for the few that bioaccumulate through food chains to a significant extent (i.e., PCBs, methyl mercury, and selenium). For those species, measured body burdens are more reliable measures of exposure. In addition, the sample preparation and analysis techniques used to measure concentrations in water are intended to measure total contaminant concentrations rather than bioavailable concentrations. Therefore, the reported concentrations may be very conservative estimates of aqueous exposure.

2.3.2 Soil

The principal source of contaminant exposure on WAG 5 is likely to be the soil. Several radionuclides have been identified in WAG 5 soil but nonradiological contamination has not been determined (Shoun 1987). The same is true of the soils of the receiving system, WAG 2 (Blaylock et al. 1992). Chemicals in soil may be directly ingested by animals, taken up by plants, indirectly ingested with prey or plant matter. In addition, the radionuclides may cause external radiation exposure. Plant uptake is primarily a function of concentration in the aqueous phase, but other exposure routes are expected to be a function of total concentration.

2.3.3 Sediments

The only sediments on WAG 5 are those in the waste ponds. These are considered to be waste materials and therefore are treated as an ultimate source rather than as a contaminated ecosystem component.

2.4 EFFECTS ASSESSMENT

Three types of effects data (laboratory toxicity data for individual chemicals, biological survey data, and media toxicity test data) are potentially useful in ecological risk assessments, but none can be used for assessment of effects on WAG 5. First, because neither source terms nor nonradiological contaminant concentrations in ambient media are available for WAG 5, laboratory toxicity data cannot be used to estimate effects on WAG 5. Second, because no biological surveys have been conducted on WAG 5, effects based on the biological properties of the site cannot be estimated. Finally, neither soils nor waters from WAG 5 have been tested for toxicity. As a result of the ORNL Biological Monitoring and Abatement Program (BMAP) program, all three of these types of effects data are available for WAG 2. The results of the WAG 2 ecological screening risk assessment, as they relate to WAG 5, are discussed below.

2.5 RISK CHARACTERIZATION

2.5.1 Aquatic Risks

The screening assessment for WAG 2 provides indirect information about contaminants of concern because WAG 2 borders WAG 5 on the west and south and because WAG 2 receives leachate and runoff from WAG 5 and from similar waste burial sites in the WOC watershed. Mercury and PCBs were the primary contaminants of concern in WAG 2. These contaminants exceeded water quality criteria and toxicological benchmarks in water and were estimated to exceed those values in sediment pore water. In addition PCB and mercury concentrations in fish flesh exceeded the lower screening criteria for piscivorous wildlife. These hazardous levels were found in both WOC and Melton Branch. In addition, Al, Cd, Cr, and Pb in water, Ag, Ba, Co, Cr, and Zn in sediments, and Se in fish flesh were high enough in some samples to suggest hazards to aquatic life. Many compounds were not detected but had detection limits higher than toxicological benchmarks. In particular, detection limits for Ag, As, and Se in water exceeded national ambient water quality criteria. PCBs, Hg, and the other elements found to constitute potential hazards in WAG 2 waters have all been found in ground water samples from WAG 5 (Shoun 1987 Bechtel 1988).

Radionuclide concentrations in WAG 2 are sufficiently low that effects on populations of aquatic animals should not occur (Blaylock et al. 1992).

The toxicity of surface waters in WOC and Melton Branch have been tested on a regular basis since 1986. Because of the major changes that have occurred in effluent input and toxicity in that interval, this discussion will emphasize the most recent results (Loar 1991). The tests employed are the standard 7-d static renewal tests measuring the growth and survival of fathead minnow (*Pimephales promelas*) larvae and the survival and reproduction

of the daphnid cladoceran *Ceriodaphnia dubia* (Weber et al. 1985). During 1990, one or both of these tests were performed monthly with water taken from WOC just above the confluence of Melton Branch and from Melton Branch just above its mouth. In all cases, survival and reproduction of *C. dubia* were high. Fathead minnow survival was less than 70% for the Melton Branch samples in four months and for the WOC samples in one month. However, in each case of low survival, minnow growth was good and variance in survival among replicates was high. Hence, the observations of low minnow survival were attributed to disease rather than toxicity by the authors (A. J. Stewart and L. A. Kszos in Loar 1991). Results in 1989 were similar but with a higher frequency of tests with low minnow survival (Loar 1990).

Biological survey data available from BMAP (Loar 1987-1991, Loar et al. 1981) indicate that WAG 2 supports a diverse fish community and that, although the condition index of redbreast sunfish is lower than in reference streams, fish are growing as well as in reference streams and appear to be reproducing well. Liver detoxification enzymes are elevated in redbreast sunfish (*Lepomis auritus*) from WAG 2, suggesting that the fish are exposed to organic contaminants, and creatinine levels are elevated, suggesting possible kidney damage. A fish and amphibian kill occurred in Melton Branch in 1990 but was attributed to high water temperatures. The macroinvertebrate community in WAG 2 appears to be degraded and to contain high frequencies of pollution tolerant species, but insufficient reference sampling has been performed and the cause of this condition is unclear.

In sum, measurements of contaminant concentrations suggest that toxic effects could be occurring in WAG 2 waters and sediments. However, the screening criteria are conservative and the contaminant concentrations are total concentrations rather than bioavailable concentrations, which compounds the conservatism. Toxicity tests with WAG 2 water do not support the occurrence of aquatic toxic effects in WAG 2, but they include only two species and only one life stage of the fish, and they are performed with grab samples, so they may miss episodic toxicity. The biological survey data do not indicate severe toxic effects in the aquatic environment, but the benthic invertebrate community may be modified by contamination, and fish may be experiencing subtle toxic effects.

The consequence of these results for WAG 5 is that, because toxic effects may be occurring in Melton Branch and WOC, contributions from WAG 5 cannot be dismissed. However, because WAG 5 contributes relatively little of the flow of these streams, the contaminant concentrations and toxicity of seepage and runoff from WAG 5 would need to be relatively high to significantly contribute to the problem; therefore, they should be readily detectable by chemical analysis and toxicity testing of runoff and seep waters.

2.5.2 Terrestrial Risks

No information is available concerning exposure of plants and animals of WAG 5 to nonradionuclide chemicals or effects of such exposures on plants and animals on WAG 5. Effects of radiation exposure on terrestrial organisms cannot be precluded but cannot be reasonably assessed with existing data. The mowed lawns that dominate WAG 5 and that cover the buried wastes constitute an unnatural vegetation that provides poor habitat for the small amphibians, reptiles and mammals that might form distinct populations on an area the size of WAG 5; therefore, biological surveys of WAG 5 are likely to be uninformative. However, larger species such as deer and wild turkey have been observed to use WAG 5.

Individuals of these species may become contaminated on WAG 5, and could spread contaminants across the reservation, serve as vectors to humans, and conceivably experience significant radiation effects.

2.6 DATA AND ASSESSMENT NEEDS

2.6.1 Biological Sampling, Analysis, and Testing

Surveys should be conducted for threatened, endangered, or state-listed species and for wetlands. Floodplains should not occur on WAG 5.

Because the contaminant composition of WAG 5 is unknown, a reasonable approach to ecological risk assessment is to begin by performing toxicity tests. The contribution of WAG 5 to aquatic toxicity occurs through runoff and groundwater seepage. Therefore, water from the six identified seeps and the two ephemeral tributaries should be sampled and tested for toxicity to *C. dubia* in the 7-d survival and reproduction test. Soil samples should be collected for toxicity testing from areas with relatively high surface-radiation levels (on the assumption that nonradionuclide chemicals are associated with the radioactive wastes) and from any areas in which vegetation reestablishment seems to be inhibited. These samples should be used in earthworm and seed germination tests.

Wildlife contamination surveys should be conducted to determine if wildlife are exposed to toxic contaminants on WAG 5 in a manner that results in detectable bioaccumulation. The species should be chosen for their societal relevance and potential for exposure. Game species are appropriate because they are valued by the public and because they are potential vectors for human exposure. Deer are already monitored for radionuclide contamination by the Tennessee Wildlife Resources Agency (TWRA) check-in of deer killed by hunters. Waterfowl, including the Canada geese that may graze on WAG 5, are monitored by BMAP. The remaining game species that is known to use WAG 5 is the wild turkey. Turkeys from flocks that use WAG 5 should be trapped and analyzed for radionuclides, PCBs, mercury, and trace metals. In addition, if burrowing mammals are found within the burial grounds, burrows should be surveyed for radioactivity. If the burrow entrance is radioactive, soil samples from the excavated material should be analyzed.

2.6.2 Baseline Assessment Strategy

Unless wetlands or threatened, endangered, or state-listed species are found on WAG 5, the site should be treated primarily as a current source of contaminants to the aquatic ecosystem of WAG 2 and to the terrestrial ecosystem of the ORR. WAG 2 and the reservation will each be assessed separately using the information from WAG 5 and the other source WAGs as input. For this purpose, the WAG 5 baseline risk assessment should identify the input of WAG 5 to WAG 2 in terms of contaminant fluxes, in terms of toxic units, and in terms of contributions to the contaminant levels in reservation wildlife.

Secondarily, WAG 5 should be assessed as a terrestrial site that would, if abandoned, revert to a natural biotic community to the extent that site toxicity allows. The risks to future vegetative communities can be assessed on the basis of the soil phytotoxicity and earthworm

test results, soil composition, and the current state of vegetation on the site. The risks to wildlife can be assessed on the basis of the results of wildlife contamination studies and simple bioaccumulation models like those used in the human health risk assessment.

3. RARE AND ENDANGERED SPECIES AND WETLANDS

Surveys of rare and endangered animal species and of wetlands were conducted for WAG 5. No rare or endangered animals have been identified in WAG 5. The borders of wetlands identified within WAG 5 are included in the map in the appendix.

4. AMBIENT TOXICITY TESTING

Bioassays are tests that use responses of living organisms to quantify environmental conditions. In this study, tests with a freshwater microcrustacean, *Ceriodaphnia dubia*, were used to assess the quality of water samples collected from various locations in WAG 5. The procedures for conducting *Ceriodaphnia* tests were developed initially by the EPA; the method was developed to estimate the toxicity of wastewaters and pure chemicals, largely for purposes of environmental compliance under the National Pollutant Discharge and Elimination System (NPDES) permit program (Mount and Norberg 1984). As toxicity testing with *Ceriodaphnia* became incorporated more extensively into the NPDES program, additional studies showed that this test method could be used reliably to (1) quantify toxicity in ambient waters (cf. Burton et al. 1987, Stewart et al. 1990), (2) identify point- or area-source contributors to ambient toxicity (Nimmo et al. 1990), and (3) identify constituents in ambient and effluent waters that contributed to toxicity (cf. Norberg-King et al. 1991; Kszos et al. 1992). *Ceriodaphnia* tests provide an estimate of toxicity by measuring the survival and reproduction of the animals. Because survival and reproduction are processes fundamental to all organisms, *Ceriodaphnia* test results can be used in ecological risk assessments.

4.1 MATERIALS AND METHODS

Ten sites to be evaluated for toxicity initially were identified in the Ecological Assessment Plan for WAG 5 (Fig. 1). Preliminary investigation showed that two of the sites (the OHF pond, site 5; and the process waste sludge basin, site 10) were too radiologically contaminated to be tested in the aquatic toxicology laboratory. A grab sample of water from a seepage site on the border of SWSA 5 South (site 6) also proved to be too contaminated for toxicity testing, even after the water had been filtered to remove suspended sediment. Water from site 1, a seepage site on the bank of WOC, was not accessible to toxicology laboratory personnel during the first two test periods. Thus, during the first two toxicity test periods (December 17-19, 1991, and February 27-March 5, 1992), daily grab water samples from each of six locations (sites 2, 3, 4, 7, 8, and 9) were evaluated. A 3-L grab sample of water from site 1 was collected on April 29, and tested for toxicity during April 30-May 7, 1992, during the third test period.

Toxicity assessments and chemical water quality analyses of the water samples were made using U.S. Environmental Protection Agency (EPA) procedures (EPA 1979). The toxicity assessments were made using 7-d static-renewal tests based on the survival and reproduction of *Ceriodaphnia dubia*; these procedures are described by Weber et al. (1989). The chemical water quality measurements included assessments of pH, conductivity (YSI Model 32 conductance meter), alkalinity (EPA Method 130.1), hardness (EPA Method 130.2), and dissolved oxygen (VWR dissolved oxygen meter).

Each *C. dubia* test was initiated by putting one neonate (<24 h old) into each of 10 20-mL polystyrene beakers containing 16 to 19 mL of test solution. Within a test, neonates did not differ in age by more than 8 h. A control treatment, consisting of a set of 10 beakers, each containing dilute mineral water [a 2:8 ratio (volume-to-volume) of Perrier® water and deionized distilled water augmented with trace metals] and one neonate, was used with each

test. When the water was renewed each day, the animals were fed 100 μL of a mixture of trout chow, yeast, Cerophyl and green algae. The temperature was maintained at $25 \pm 1^\circ\text{C}$ in each test; the temperature of the samples was adjusted to $25 \pm 2^\circ\text{C}$ before use. On each day of the test, the animals were inspected visually for survival prior to being transferred to fresh test solution. When the animals achieved reproductive maturity, the number of neonates produced by each animal was counted immediately after the adult had been transferred. The results of the tests were summarized statistically using SAS (SAS Institute 1985) on a personal computer.

4.2 RESULTS AND DISCUSSION

The results of the *Ceriodaphnia* tests conducted during the first two test periods are summarized in Table 2. In each of these tests, the control was acceptable and there was no evidence for acute toxicity (i.e., survival of the daphnids was consistently greater than 60%). However, significant reductions in *Ceriodaphnia* reproduction were detectable (based on Tukey's test) in at least one of the WAG 5 sites in each test period, suggesting the presence of chronic toxicity. In each test period, mean reproduction values for the *Ceriodaphnia* were ranked (1 for highest, 7 for lowest; ties were assigned a mean rank value). The mean ranks for each of the sites, based on the two test periods, were as follows: controls, 1.75; site 2, 2.25; site 3, 3; site 7, 3.5; site 8, 4.5; site 4, 6; and site 9, 7. The high mean rank values for sites 4 and 9 suggested strongly that water from these two sites adversely affected *Ceriodaphnia* during both test periods. Water from these two sites shows good evidence for chronic toxicity.

Table 2. Results of *Ceriodaphnia* toxicity tests of WAG 5 water samples during test period 1 (December 12-18, 1991) and test period 2 (February 27-March 5, 1992)

Site	Test Period 1		Test Period 2	
	Survival (%)	Reproduction (mean $\pm$ SE)	Survival (%)	Reproduction (mean $\pm$ SE)
Control	100	19.4 $\pm$ 3.5	100	19.2 $\pm$ 1.0
2	80	19.4 $\pm$ 3.4	90	16.8 $\pm$ 1.8
3	80	21.1 $\pm$ 2.5	90	13.1 $\pm$ 1.9 ^a
4	90	10.7 $\pm$ 2.8	70	9.3 $\pm$ 0.7 ^a
7	100	14.5 $\pm$ 2.3	80	15.9 $\pm$ 1.1
8	80	12.9 $\pm$ 1.0	100	14.8 $\pm$ 1.3
9	70	0.9 $\pm$ 0.6 ^a	100	8.2 $\pm$ 0.9 ^a

^aSignificantly lower than the corresponding test-period control based on Tukey's test ($P < 0.05$).

The results of the third toxicity test, which was used to assess water from WAG 5 site 6, are summarized in Table 3. In this test, the animals had relatively low survival (50%), but the daphnids that survived had a level of reproduction typical for controls (21.2 ± 1.5 offspring per female). In ambient testing situations, such as that used for assessing the surface water and seepage water samples from WAG 5, the probability of getting a *Ceriodaphnia* survival value less than or equal to 60% is only about 1.5% in tests where survival in the control is 80% to 100% (Loar 1991). Thus, the results of the site 6 test could be considered to be equivocal. However, ambient tests conducted concurrently with the site 6 test (including tests of filtered and nonfiltered water from the outfall of Lake Reality, a settling basin at the Oak Ridge Y-12 Plant) had high survival and very high reproduction (cf. Table 3). Thus, in comparison to the test results for the two Lake Reality samples, the water from site 6 (WAG 5) might be considered to be toxicologically problematic. The results of confirmatory tests would reduce uncertainty about this possibility.

Table 3. Results of *Ceriodaphnia* toxicity test of water from WAG 5 site 6 (April 30–May 7, 1992)^a

Site	Survival (90%)	Reproduction (mean $\pm$ SE)
Control ^a	20	25.0 ± 3.0
LR-o (nonfiltered)	90	47.7 ± 3.3
LR-o (filtered) ^b	100	47.5 ± 1.5
WAG 5 Site 6	50	21.2 ± 1.5

^aThe diluted mineral water control for this test was unacceptable. Thus, the results of an ambient toxicity test of filtered and nonfiltered water from the outfall of Lake Reality (LR-o; the Oak Ridge Y-12 Plant settling basin) conducted during the same period are provided for comparative purposes.

^bWater sample was filtered through a 0.5- μ m pore size glass fiber filter to remove particulate matter prior to testing.

Data for the chemical water quality factors measured during the three toxicity test periods are summarized in Tables 4, 5, and 6. Rain events occurred between the second and third sampling days during the first test period. A temporary dilution of conservative water quality factors (such as conductivity, alkalinity, and hardness) was caused by the rainfall; this situation resulted in greater variation of these factors during the February test than during the December test (Table 4 vs Table 5). With each site-date sample considered individually, none of the sites had pH, conductivity, alkalinity, or hardness levels that would be expected to adversely affect *Ceriodaphnia* survival or reproduction, based on much experience in ambient and effluent testing with this organism. Conductivity, hardness, and pH values for the samples were similar to those measured for other streams near ORNL (e.g., WOC and Melton Branch). Additionally, the concentrations of dissolved oxygen in the water samples was, in every instance, over 7 mg/L. A concentration of oxygen of 7 mg/L is high enough to be nonlimiting to *Ceriodaphnia*. Thus, none of the measured water quality factors, singly or in combination, appeared to be adverse enough to lower either the survival or reproduction of *Ceriodaphnia*. This consideration suggests that reductions in either survival or reproduction in these tests could provide reliable evidence for acute or chronic toxicity, respectively.

Table 4. Summary of results of water quality analyses, WAG 5 sites, for toxicity tests conducted during December 12-18, 1991^a

Water Quality Factor	Site 2	Site 3	Site 4	Site 7	Site 8	Site 9
pH (standard units)	7.78 ± 0.22	7.75 ± 0.12	7.72 ± 0.16	7.82 ± 0.16	7.74 ± 0.20	7.68 ± 0.2
Conductivity (μ S)	375 ± 81	341 ± 45	498 ± 26	480 ± 28	473 ± 75	596 ± 6
Alkalinity (mg/L CaCO ₃)	124 ± 28	127 ± 29	234 ± 9	225 ± 13	213 ± 35	299 ± 5
Hardness (mg/L CaCO ₃)	197 ± 39	170 ± 15	245 ± 19	224 ± 9	216 ± 19	295 ± 11
Dissolved oxygen (mg/L)	8.2 ± 0.1	8.3 ± 0.1	8.1 ± 0.1	8.2 ± 0.1	8.1 ± 0.1	7.8 ± 0.2

^aValues are mean ± SE.

Table 5. Summary of results of water quality analyses, WAG 5 sites, for toxicity tests conducted during February 27-March 5, 1992^a

Water Quality Factor	Site 2	Site 3	Site 4	Site 7	Site 8	Site 9
pH (standard units)	7.91 ± 0.09	8.11 ± 0.03	8.14 ± 0.03	8.19 ± 0.03	7.87 ± 0.16	8.12 ± 0.06
Conductivity (μ S)	278 ± 18	315 ± 11	522 ± 11	541 ± 4	650 ± 8	605 ± 5
Alkalinity (mg/L CaCO ₃)	102 ± 6	114 ± 5	260 ± 7	269 ± 4	354 ± 24	301 ± 2
Hardness (mg/L CaCO ₃)	135 ± 10	159 ± 7	278 ± 5	269 ± 3	294 ± 4	285 ± 8
Dissolved oxygen (mg/L)	8.1 ± 0.2	8.3 ± 0.1	8.0 ± 0.2	7.9 ± 0.2	7.9 ± 0.2	8.0 ± 0.2

^aValues are mean ± SE.

Table 6. Summary of results of water quality analyses, WAG 5 site 6, for the toxicity tests conducted during April 30-May 7, 1992^a

Water Quality Factor	Site 6
pH (standard units)	7.42
Conductivity (μS)	170
Alkalinity (mg/L $CaCO_3$)	72
Hardness (mg/L $CaCO_3$)	96
Dissolved oxygen (mg/L)	8.4

^aValues are for single measurements made from the grab sample on the first day of the test.

5. WILD TURKEY CONTAMINANT SCREENING

Wild turkeys were reintroduced to the ORR by TWRA in 1984. Subsequently, the population of these animals has increased to the point that TWRA routinely removes turkeys from the ORR and transports them to other areas of the state for restocking programs. In addition, some turkeys probably leave the ORR on their own.

Wild turkeys have been observed feeding in WAG 5 and in waste burial areas in other WAGs. Although the present potential for transfer of site contaminants through turkeys to humans is low because no nearby areas are open for turkey hunting, it is possible that turkey hunting on the ORR will be permitted in the future (J. W. Evans, TWRA, personal communication to T. L. Ashwood, ORNL, July 1992). Moreover, under a no action scenario, wild turkeys may serve as a pathway to humans who gain access to the site after the termination of institutional control.

5.1 METHODS

Anecdotal reports and sightings by project personnel suggested that a large flock, or perhaps two flocks, were feeding in the WAG 5 area. During the period of trapping, however, one part of the flock(s) was found to be frequenting an area of the White Oak Lake floodplain very near to the lake, and a second group from the same flock(s) was found in the vicinity of the water towers north of Melton Valley Drive and west of the Experimental Gas-Cooled Reactor Access Road. Trapping was attempted in both locations. In order to provide a reference site, turkeys were trapped from an uncontaminated site on Lou Cagle Road west of state highway 95. Trapping was attempted from January through February 1992. Turkeys cannot be effectively live-trapped during warm months because trap mortality increases with temperature.

5.1.1 Trapping

Turkeys were lured into specific trap sites by cracked corn placed at the sites for several days prior to the trapping event in order to accustom the turkeys to finding bait at the site. The time of day when the turkeys took the bait was noted during this accustomization period so that the trap could be set well in advance of the arrival of the animals on the day of trapping.

A rocket net was used to trap turkeys on Lou Cagle Road. The rocket net utilizes three explosive charges fired from tubes to launch the net up and out over the clustered turkeys. This technique requires a large clear area where the turkeys can cluster and where the rocket net can be fired without obstruction. Because no such areas were available in the contaminated area, the rocket net was not used for trapping potentially contaminated animals.

In areas where sufficient clearance was not available to fire the rocket net, turkeys were anesthetized with bait that had been saturated with tribromoethanol, a chemical that interferes with the transmission of nerve impulses. Anesthetized birds were placed in burlap bags for transfer to the TWRA checking station.

5.1.2 Radiological Screening

Turkeys were screened for radiological contamination at the TWRA checking station established to support the deer hunts on the ORR. Screening of wild turkeys was similar to screening of deer killed during the hunts.

Leg bones were screened with a Ludlum Model 2200 single-channel scaler system with a Ludlum Model 44-1 beta scintillation probe. The beta screening system is calibrated at the beginning of each analysis session using a ^{90}Sr source traceable to the National Bureau of Standards (NBS). Live turkey legs were held under the detector for 1 min. Leg bones from dead turkeys were also held immediately under the detector head for 1 min. In one case, the skin was stripped from the dead turkey leg to determine whether the skin reduced the number of counts.

Live turkeys received a 5-min whole body gamma count on an NaI detector coupled through an analog-to-digital converter to a 386SX computer that is programmed to act as a multichannel analyzer. The gamma counting system is calibrated at the beginning of each analysis session with an NBS-traceable ^{137}Cs standard solution.

The first attempts to count live turkeys by simply resting the bagged bird on the detector resulted in the birds falling off the detector head. Subsequent counts were made with the bird in a Lucite® box that prevented the bird from moving off the detector. Dead turkeys were not gamma counted because they were too big and too rigid to fit into the detector space.

5.2 RESULTS AND DISCUSSION

Trapping success was quite low. Three female turkeys were collected from the potentially contaminated flock in the vicinity of White Oak Lake. Two male turkeys were collected from the uncontaminated site on Lou Cagle Road. The two male turkeys were killed by the rocket net and were kept frozen.

Table 7 contains results of the screening analyses performed on the five turkeys. No turkey contained significant gamma-emitting isotopes. Only one turkey (WOL2, Table 7) contained beta-emitting isotopes substantially above background. This turkey, a hen, was sacrificed. The remaining females were taken by TWRA personnel for release as part of wild turkey reintroduction programs in other areas of east Tennessee.

These limited data do not provide sufficient sample size to assess the distribution of contaminants within the wild turkey population that feeds on WAG 5. In addition, data are needed on other nonradiological contaminants.

Additional attempts to live-trap wild turkeys cannot be made until cold weather. It is not clear at the time of this report if sufficient time is available after the start of cold weather to collect the samples and obtain the results before the data are needed for the baseline risk assessment. Nevertheless, further attempts to live-trap wild turkeys will be made because the data are important for risk assessments in all WAGs where the potential exists for turkeys to become contaminated.

Table 7. Radiological screening results for wild turkeys

Turkey ID ^a	Tissue Analyzed	Analyte ^b	Result ^c
LC01	Bone	Beta	15 cpm
	Bone	Beta	13 cpm
	Bone	Beta	18 cpm ^d
LC02	Bone	Beta	24 cpm
	Bone	Beta	19 cpm
WOL1	Whole body	Gamma	0.033 ± 0.013 pCi/g ^e
	Whole body	Gamma	0.027 ± 0.013 pCi/g ^e
	Whole body	Gamma	0.047 ± 0.013 pCi/g ^f
	Bone	Beta	24 cpm
	Bone	Beta	19 cpm
WOL2	Whole body	Gamma	0.054 ± 0.011 pCi/g ^f
	Bone	Beta	28 cpm
	Bone	Beta	35 cpm
	Bone	Beta	40 cpm
WOL3	Whole Body	Gamma	0.063 ± 0.013 pCi/g ^f
	Bone	Beta	19 cpm
	Bone	Beta	22 cpm

^aWOL = turkeys captured in floodplain near White Oak Lake; LC = turkeys captured on Lou Cagle Road.

^bBeta = beta radiation detected by counter in 1-min count; Gamma = gamma radiation detected by Ge (Li) detector in 5-min count and interpreted as ¹³⁷Cs.

^ccpm = counts per minute.

^dCount was made with skin stripped from leg bone.

^eTurkey (in burlap bag) was placed directly on gamma detector, but turkey had fallen off detector by end of counting time.

^fTurkey was placed in Lucite® box and box was placed on gamma detector.

6. FUTURE ACTIVITIES

The primary planned activity not yet completed is toxicity testing of the soil from WAG 5 (Ashwood 1992). Also, additional ambient toxicity tests of the seeps and streams are required to confirm earlier indications of toxicity and to identify a potential toxicant. If feasible, additional turkey samples will be collected, and existing samples will be analyzed for trace metals and PCBs as indicated in the original plan (Ashwood 1992). The screening-level ecological risk assessment has identified the need for radiological screening of woodchuck burrows.

7. ACKNOWLEDGEMENTS

The authors thank Peggy Braden, Linda Wicker, Gail Morris, and Diana Hicks for assistance in collecting and analyzing the ambient toxicity samples. Barbara Rosenstiel and Roger Kroodsma conducted the rare and endangered species surveys. Jim Evans and Dick Hurd, both of TWRA, collected the wild turkeys. Norm Teasley and Carolyn Grainger, both of the Analytical Chemistry Division, conducted the radiological screening of the wild turkeys. Lynn Kszos and George Southworth reviewed a draft of this report.

8. REFERENCES

- Ashwood, T. L. 1992. Ecological assessment plan for Waste Area Grouping 5. ORNL/M-1871. Oak Ridge National Laboratory.
- Bechtel National, Inc. 1988. Remedial investigation plan for ORNL Waste Area Grouping 5. ORNL/RAP/Sub-87/99053/8&V1. Oak Ridge National Laboratory, Oak Ridge, Tennessee.
- Blaylock, B. G., M. L. Frank, F. O. Hoffman, L. A. Hook, G. W. Suter, and J. A. Watts. 1992. Screening of contaminants in Waste Area Grouping 2 at Oak Ridge National Laboratory, Oak Ridge, Tennessee. ORNL/ER-62. Oak Ridge National Laboratory, Oak Ridge, Tennessee.
- Boyle, J. W., R. Blumberg, S. J. Cotter, G. S. Hill, C. R. Kerley, R. H. Ketelle, R. L. Kroodsmas, D. W. Lee, R. C. Martin, R. D. Roop, D. N. Secora, W. P. Staub, and R. E. Thoma. 1982. Environmental analysis of the operation of the Oak Ridge National Laboratory (X-10 Site). ORNL-5870. Oak Ridge National Laboratory, Oak Ridge, Tennessee.
- Burton, G. A., D. Nimmo, D. Murphey, and F. Payne. 1987. Stream profile determinations using microbial activity assays and *Ceriodaphnia*. Environ. Toxicol. Chem. 6:505-513.
- EPA (U.S. Environmental Protection Agency). 1979. Methods for chemical analysis of water and wastes. EPA-600/4-79-020. Environmental Monitoring and Support Laboratory, U.S. Environmental Protection Agency, Cincinnati, Ohio.
- Kroodsmas, R. L. 1987. Resource management plan for the Oak Ridge Reservation, Volume 24: Threatened or endangered animal species. ORNL/ESH-1/24. Oak Ridge National Laboratory, Oak Ridge, Tennessee.
- Kszos, L. A., A. J. Stewart, and P. A. Taylor. 1992. An evaluation of nickel toxicity to *Ceriodaphnia dubia* and *Daphnia magna* in a contaminated stream and in laboratory tests. Environ. Toxicol. Chem. vol. 11 (in press).
- Loar, J. M., J. A. Solomon, and G. F. Cada. 1981. Technical background information for the ORNL Environmental and Safety Report, Vol. 2: A description of the aquatic ecology of the WOC watershed and Clinch River below Melton Hill Dam. ORNL/TM-7509/V2. Oak Ridge National Laboratory, Oak Ridge, Tennessee.
- Loar, J. M., S. M. Adams, H. Amano, J. B. Berry, B. G. Blaylock, H. L. Boston, M. L. Frank, C. T. Garten, M. A. Huston, B. D. Jimenez, B. L. Kimmel, J. T. Kitchings, L. Meyers-Schone, D. A. Mohrbacher, C. R. Olsen, M. G. Ryon, J. G. Smith, G. R. Southworth, A. J. Stewart, S. S. Talmage, and B. T. Walton. 1987. First annual report on the Biological Monitoring and Abatement Program at Oak Ridge National Laboratory. ORNL/TM-10399. Oak Ridge National Laboratory, Oak Ridge, Tennessee.
- Loar, J. M. (ed.). 1988. Second annual report of the ORNL Biological Monitoring and Abatement Program. ORNL/TM-10804. Oak Ridge National Laboratory, Oak Ridge, Tennessee.
- Loar, J. M. (ed.). 1989. Third annual report of the ORNL Biological Monitoring and Abatement Program. ORNL/TM-11358. Oak Ridge National Laboratory, Oak Ridge, Tennessee.
- Loar, J. M. (ed.). 1990. Fourth annual report of the ORNL Biological Monitoring and Abatement Program. ORNL/TM-11544. Oak Ridge National Laboratory, Oak Ridge, Tennessee.

- Loar, J. M. (ed.). 1991. Fifth annual report of the ORNL Biological Monitoring and Abatement Program. ORNL/TM-12143. Oak Ridge National Laboratory, Oak Ridge, Tennessee.
- Mann, L. K. and J. T. Kitchings. 1982. Resource data inventory for the Oak Ridge area. ORNL/TM-7941. Oak Ridge National Laboratory, Oak Ridge, Tennessee.
- Mount, D. I., and T. J. Norberg. 1984. A seven-day life cycle cladoceran toxicity test. *Environ. Toxicol. Chem.* 3:425-434.
- Nimmo, D. W., M. H. Dodson, P. H. Davies, J. C. Greene, and M. A. Kerr. 1990. Three studies using *Ceriodaphnia* to detect nonpoint sources of metals from mine drainage. *Res. J. Water Pollut. Control Fed.* 62:7-15.
- Norberg-King, T. J., E. J. Durhan, and G. T. Ankley. 1991. Application of toxicity identification evaluation procedures to the ambient waters of the Colusa basin drain, California. *Environ. Toxicol. Chem.* 10:891-900.
- Oakes, T. W., B. A. Kelly, W. O. Ohnesorge, J. S. Eldridge, J. C. Bird, K. E. Shanks, and F. S. Tsakeres. Technical background information for the ORNL Environmental and Safety Report, Vol. 4: White Oak Lake and Dam. ORNL/TM-7509/V4. Oak Ridge National Laboratory, Oak Ridge, Tennessee.
- SAS Institute. 1985. SAS User's Guide: Basics, Version 5 Edition. SAS Institute, Inc., Cary, NC.
- Sherwood, C. B. and J. M. Loar. 1987. Environmental data for the White Oak Creek/White Oak Lake watershed. ORNL/TM-4662. Oak Ridge National Laboratory, Oak Ridge, Tennessee.
- Shoun, R. R. 1987. Environmental data package for ORNL Waste Area Grouping—Five (WAG-5), Solid Waste Storage Area—Five (SWSA-5). ORNL/RAP-19. Oak Ridge National Laboratory, Oak Ridge, Tennessee.
- Stewart, A. J., L. A. Kszos, B. C. Harvey, L. F. Wicker, G. J. Haynes, and R. D. Bailey. 1990. Ambient toxicity dynamics: assessments using *Ceriodaphnia dubia* and fathead minnow (*Pimephales promelas*) larvae in short-term tests. *Environ. Toxicol. Chem.* 9:367-379.
- Suter, G. W., A. Redfearn, R. K. White, and R. A. Shaw. 1992. Approach and strategy for performing ecological risk assessments for the Department of Energy Oak Ridge Field Office environmental restoration program. ES/ER/TM-33. Oak Ridge National Laboratory.
- Weber, C. I., W. H. Peltier, T. J. Norberg-King, W. B. Horning II, F. A. Kessler, J. R. Menkedick, T. W. Neiheisel, P. A. Lewis, D. J. Klemm, Q. H. Pickering, E. L. Robinson, J. M. Lazorchak, L. J. Wymer, and R. W. Freyberg. 1989. Short-term methods for estimating the chronic toxicity of effluents and receiving waters to freshwater organisms, 2nd ed. EPA 600/4-89-001. U.S. Environmental Protection Agency, Cincinnati, Ohio.

APPENDIX

ENDANGERED SPECIES AND WETLANDS SURVEY REPORTS

Internal Correspondence

MARTIN MARIETTA ENERGY SYSTEMS, INC.

July 16, 1992

T. L. Ashwood

Threatened and Endangered Animal Species at WAG 5

No threatened or endangered animal species (aquatic and terrestrial invertebrates and vertebrates) or critical habitat listed, or proposed to be listed, by the U.S. Fish and Wildlife Service is known to be present on the Oak Ridge Reservation.^{1,2,3} However, the endangered Indiana bat is a possible summer resident along East Fork Poplar Creek and must be included in environmental considerations for proposed construction projects in this area.

Several animal species listed by the State of Tennessee^{1,4,5} as threatened (T), endangered (E), or in need of management (INM) are known to occur on the Oak Ridge Reservation. Environmental considerations for any proposed project that would disturb habitats where a T, E, or INM species occurs must include the potentially affected species.

The following list indicates the locations where T, E, or INM species may occur on the Oak Ridge Reservation. Any proposed project that would not affect habitats in these locations would not be expected to affect the species.

<u>Species</u>	<u>Status</u>	<u>Location</u>
<u>U.S. Fish and Wildlife Service</u>		
Indiana bat	E	Vicinity of East Fork Poplar Creek
<u>State of Tennessee</u>		
Tennessee dace	INM	Bear Creek drainage, Ish Creek, and tributaries of East Fork Poplar Creek (siltation due to erosion in the vicinity of the creeks must be avoided)
Osprey	E	Vicinity of Melton Hill Lake and Clinch River
Sharp-shinned hawk	E	Area between Jones Island, Grubb Island, and 500-kV power line
Cooper's hawk	T	Entire Oak Ridge Reservation ⁶
Grasshopper sparrow	T	Large grassy fields
Black-crowned night-heron	INM	Clinch River
Black vulture	INM	Entire Oak Ridge Reservation ⁶ ; an active nest site is located on Chestnut Ridge between Highway 95 and New Zion Patrol Road
Red-shouldered hawk	INM	Forests and fields along the Clinch River and relatively large streams ⁶
Common barn owl	INM	Freel's bend and vicinity

No threatened or endangered animal species is known to occur on WAG 5.

REFERENCES AND FOOTNOTES

¹Kroodsma, R. L. 1987. Resource Management Plan for the Oak Ridge Reservation, Volume 24: Threatened and Endangered Animal Species. ORNL/ESH-1/V24, Oak Ridge National Laboratory, Oak Ridge, Tennessee. 24 pgs.

²Barclay, L. A. (U.S. Fish and Wildlife Service, Cookeville, TN). June 13, 1990. Letter to R. L. Kroodsma (ORNL).

³Bay, R. T. (U.S. Fish and Wildlife Service, Cookeville, TN). March 7, 1991. Letter to R. L. Kroodsma (ORNL).

⁴Hatcher, R. M. 1990. Help for Tennessee's rare and endangered wildlife. Tennessee Wildlife Resources Agency, Nashville, Tennessee

⁵TWRC. 1990. Wildlife in need of management. Tennessee Wildlife Resources Commission Proclamation No. 86-29 as amended by Proc. No. 90-2, dated 2-1-90, Nashville, Tennessee

⁶During survey or construction activities at a project site, project personnel should report any observations of hawks or vultures indicating possible nesting on the site or vicinity.

R. Kroodsma, Bldg. 1505, MS-6038, Phone: 4-7310

OAK RIDGE NATIONAL LABORATORY
OPERATED BY MARTIN MARIETTA ENERGY SYSTEMS, INC.
FOR THE U.S. DEPARTMENT OF ENERGY

POST OFFICE BOX 2008
OAK RIDGE, TENNESSEE 37831

March 9, 1992

T.L. Ashwood, Bldg 1505, MS 6036, ORNL

PRELIMINARY WETLANDS INVESTIGATION - WAG 5 REMEDIAL
INVESTIGATION

I. INTRODUCTION

The preliminary wetlands investigation of the above referenced site has been completed. A field investigation of the site was conducted on February 6, 1992.

Field methodology followed procedures established in the Federal Manual for Identifying and Delineating Jurisdictional Wetlands (Federal Interagency Committee for Wetland Delineation, 1989). Use of this methodology is required on Department of Energy property by the US Army Corps of Engineers. In order to be identified as a wetland, an area must have hydrophytic vegetation, hydric soils, and be saturated by groundwater or inundated by surface water for at least seven days during the growing season. In certain situations an area can be identified as a wetland without possessing all three wetland parameters.

Wetlands were identified in six drainage areas. The locations and approximate boundaries of these wetland areas are shown on the accompanying map and are designated as D-1, D-1a, D-1b, D-2, D-2a, and D-3. The wetlands are in stream floodplain areas and are controlled hydrologically by stream overflow and groundwater discharge.

WAG 5
page two

II. WETLAND AREA DESCRIPTIONS

A. Area D-1

This area was surveyed from the head of the drainage to the gravel road on the western boundary of WAG 5. Wetlands were observed along the entire length of the tributary. A short section at the head of the drainage is dominated by scrub/shrub and emergent vegetation including black willow (Salix nigra; OBL), sycamore, and red maple saplings, and a herbaceous layer dominated by cattail (Typha sp.; OBL), sedges (Carex sp., Scirpus sp., and Cyperus sp.; these genera are primarily OBL and FACW), and seedbox (Ludwigia sp.; OBL). The dominant canopy species in the drainage are sweetgum, red maple, and sycamore, with smaller numbers of tulip poplar present. The subcanopy is similar in composition except for the absence of tulip poplar and presence of ironwood (Carpinus caroliniana; FAC). The ground cover includes nepal microstegium (Microstegium vimineum; FAC), sedges, and cane (Arundinaria gigantea; FACW). A portion of this area is similar in general appearance to sites in which orchids (i.e., Plantanthera flava v. herbiok) have been found on the Reservation. The presence or absence of rare plants in this area will be investigated during the rare plant survey to be conducted during the growing season.

Most of the area to the north of the drainage has been formerly cleared and is now dominated by maintained grasses. The slope to the south of the drainage is forested. Common species on this slope include American beech and shortleaf pine on the lower and upper slopes, respectively.

B. Area D-1a

A scrub/shrub wetland was identified in drainage D-1a. This drainage runs parallel to Melton Valley Drive and is just outside of the northern boundary of WAG 5. The wetland area is located near the toe of slope formed by the WAG 5 entrance road and is just north of building 7831A. The canopy species include sycamore (Platanus occidentalis;

WAG 5
page 3

FACW), sweetgum (Liquidambar styraciflua; FAC+), red maple (Acer rubrum; FAC), and tulip poplar (Liriodendron tulipifera; FAC). The subcanopy included all of the listed canopy species except for tulip poplar. The dominant understory species is common pawpaw (Asimina triloba; FAC). Vines in and adjacent to the wetland include Japanese honeysuckle (Lonicera japonica, FAC) and greenbriar (Smilax sp.). Due to the season, the herbaceous community could not be fully identified, however, a sedge (Cyperus spp.; most are FACW) is one of the species included in this layer. Additional wetlands downstream of this small wetland are confined to narrow, discontinuous strips of level alluvial soils immediately adjacent to the stream.

The remainder of this drainage area is dominated by non-hydrophytic vegetation, has non-hydric soils, and lacks wetland hydrology. The dominant vegetation in the upland areas includes sweetgum, tulip poplar, sycamore, red maple, shortleaf pine (Pinus echinata; FACU), white oak (Quercus alba; FACU), American beech (Fagus grandifolia; FACU), and red elm (Ulmus rubra; FAC). Understory species include sweetgum, sycamore, red maple, beech, buckeye (Aesculus flava; FACU), and black walnut (Juglans nigra; FACU). Shrub species include flowering dogwood (Cornus florida; FACU) and witchhazel.

C. Area D-1b

This is a small drainage area just south of the process waste sludge basin. It is connected to the D-1 drainage through a culvert. A scrub/shrub - emergent wetland was identified where the stream enters the culvert. In the wetland area, the dominant species are sycamore, sweetgum, and black willow saplings, which form an open subcanopy layer, and sedges, rushes (Juncus spp.) and cattails in the herbaceous layer. Another survey, which includes soil sampling, of the area immediately upstream of the culvert wetland should be conducted because the vegetation, by itself, is not sufficient evidence for a wetland determination. The vegetation in this upstream area includes sweetgum, red maple, sycamore, and Japanese honeysuckle, all of which are able to grow in hydric, as well as nonhydric, soils. The ground cover is sparse in

WAG 5
page 4

this area. A greater number of herbaceous species may be found during the growing season.

D. Area D-2

This area was surveyed from the head of the drainage to the WAG 5 boundary. Soil samples were not taken in this area.

The upper end of the drainage is not wetland. The dominant species in this upland area are tulip poplar, sycamore, sweetgum and red cedar (Juniperus virginiana; FACU). Nepal microstegium and Japanese honeysuckle are abundant in the herbaceous layer. A short distance downstream, the canopy shifts to tulip poplar, sweetgum and red maple. A seep is located approximately 1/4 of the distance down the drainage. Continuing downstream from the seep area, the wetland area generally follows topographic contours. The area is dominated by hydrophytic vegetation. Saturated soils and ponded water were observed. Black willow appears in the canopy, and ironwood, dogwood (Cornus amomum or C. stolonifera; FACW), and brookside alder (Alnus serrulata; FACW+) are abundant in the understory. Nepal microstegium dominates the herbaceous layer throughout the drainage, however, due to the season it was not possible to fully identify species in this strata.

The upland slopes adjacent to the drainage are forested with tulip poplar, sweetgum, and red cedar dominating. The areas immediately beyond the slopes are open maintained fields dominated by grasses.

E. Area D-2a

This area consists of a series of seeps located along the southern boundary of WAG 5. These areas were not closely surveyed due to high contamination levels. Most of the areas designated as seeps on the maps are dominated by hydrophytic vegetation. Species include black willow, sweetgum, sedges, and cattail. Standing water was present in most of the seep areas.

WAG 5
page 5

F. Area D-3

This area is outside of the WAG 5 boundary. Soils were not sampled due to equipment breakage. With the exception of one small area, hydrophytic vegetation was not present. The streambed in this drainage area is broad and flat, and the stream channel has steep banks. This suggests that the water table is moderately deep and that surface water may infiltrate and drain out of the area at a fairly rapid rate. The area adjacent to the stream is dominated by sweetgum, tulip poplar, and white oak. Some sycamore and ironwood are also present. Near the head of the drainage, there is an increase in the number of sycamore, and black willows appear in the canopy. If a wetland determination in this area is required, further field investigation will be necessary. This may be conducted at the same time as the rare plant survey.

III. SOIL AND HYDROLOGY

Several soil samples were taken to determine the presence or absence of hydric soils in most of the areas investigated. (Equipment failure and possible soil contamination precluded soil sampling in all areas). Hydric soils characterized by low chroma colors and mottling were found in the areas dominated by hydrophytic vegetation. Hydric soil was not found outside of these wetland areas. The wetland areas displayed saturated soils, a high water table, and ponding. Morphologic wetland indicators include drift lines, evidence of scouring, matted and silted leaves, and surface roots.

WAG 5

page 6

The results of this investigation are preliminary. A detailed wetlands delineation would need to be performed in order to accurately establish the limits of wetlands in the project area. Please feel free to contact me if you have any questions or comments.

Barbara A. Rosensteel

Barbara A. Rosensteel, Coordinator, Rare Plant and Wetland Surveys,
Environmental Sciences Division, Bldg 1506, MS-6034, ORNL (6-7544)
8123

cc: H.L. Boston w/o map
A.J. Kuharda

DISTRIBUTION

1-5. T. L. Ashwood	29-30. P. T. Owen
6. L. W. Barnthouse	31. P. D. Parr
7. L. D. Bates	32. D. E. Reichle
8. H. L. Boston	33. G. E. Rymer
9. D. T. Bell	34. F. E. Sharples
10. J. B. Cannon	35. D. S. Shriner
11. R. B. Clapp	36. G. R. Southworth
12. R. B. Cook	37-39. A. J. Stewart
13. J. H. Cushman	40. S. H. Stow
14. M. F. P. DeLozier	41-43. G. W. Suter
15. D. E. Fowler	44. K. B. Swain
16. C. W. Gehrs	45. R. I. Van Hook
17. C. D. Goins	46. L. D. Voorhees
18. P. J. Halsey	47. D. R. Watkins
19. D. S. Hicks	48. R. K. White
20. S. G. Hildebrand	49. P. A. Schrandt
21. P. Kanciruk	50. Central Research Library
22. B. L. Kimmel	51-55. ESD Library
23. L. A. Kszos	56-59. ER Document Management Center
24. J. M. Loar	50-51. Laboratory Records Dept.
25. V. Legg	52. ORNL Patent Section
26-28. D. M. Matteo	53. ORNL Y-12 Technical Library
-
54. Office of Assistant Manager for Energy Research and Development, DOE Oak Ridge Field Office, P.O. Box 2001, Oak Ridge, TN 37831-8600
 55. J. W. Evans, Tennessee Wildlife Resources Agency, P. O. Box 2008, Oak Ridge, TN 37831-6034
 56. J. F. Franklin, Bloedel Professor of Ecosystem Analysis, College of Forest Resources, University of Washington, Anderson Hall (AR-10), Seattle, WA 98195
 57. R. C. Harriss, Institute for the Study of Earth, Oceans, and Space, Science and Engineering Research Building, University of New Hampshire, Durham, NH 03824
 58. G. Y. Jordy, Director, Office of Program Analysis, Office of Energy Research ER-30, G-226, U. S. Department of Energy, Washington, DC 20545
 - 59-60. R. L. Nace, Branch Chief, Nonenrichment Facilities, Oak Ridge Program Division, Office of Eastern Area Programs, Office of Environmental Restoration, EM-423, Trevion 2, U.S. Department of Energy, Washington, DC 20585
 61. R. H. Olsen, Professor, Microbiology and Immunology Department, University of Michigan, Medical Sciences II, #5605, 1301 East Catherine Street, Ann Arbor, MI 48109-0620
 62. A. Patrinos, Director, Environmental Sciences Division, Office of Health and Environmental Research, ER-74, U.S. Department of Energy, Washington, DC 20585

- 63-64. R. C. Sleeman, DOE Oak Ridge Field Office, P.O. Box 2001, Oak Ridge, TN 37831-8540
- 65-66. J. T. Sweeney, DOE Oak Ridge Field Office, P.O. Box 2001, Oak Ridge, TN 37831-8541 number of copies needed may vary]
- 67. D. W. Swindle, Radian Corporation, 120 South Jefferson Circle, Oak Ridge, TN 37830
- 68-69. H. M. Thron, Chief, Enrichment Facilities, Oak Ridge Program Division, Office of Eastern Area Programs, Office of Environmental Restoration, EM-423, Trevion 2, U.S. Department of Energy, Washington, DC 20585.
- 70. F. J. Wobber, Environmental Sciences Division, Office of Health and Environmental Research, Office of Energy Research, ER-74, U.S. Department of Energy, Washington, DC 20585
- 71-72. Office of Scientific and Technical Information, P.O. Box 62, Oak Ridge, TN 37831

**DATE
FILMED
12/18/92**