DOE REPORT DOE/ER/60941-1

FINAL PROGRESS REPORT
 DOE CONTRACT NO. DE-FG02-90ER60941
 TITLE: PATHOGENIC REGULATION OF
 RUNNING WATER INVERTEBRATE POPULATIONS
 PI: KENNETH W. CUMMINS
 DEPARTMENT OF BIOLOGICAL SCIENCES
 UNIVERSITY OF PITTSBURGH
 PITTSBURGH, PA 15260 (412-624-5491)

DOE/ER/60941--1

DE92 015675

PREFACE

This final report reflects the fact that what was anticipated as a 3 year funding period was cut to 2 years and the amount of the award was cut in half for year two, which became the final year. thus, full completion of the project as originally proposed was not possible.

RESEARCH RESULTS

Our previous observations on stream invertebrate mortality related to growth (Cummins and Wilzbach 1988) led to experiments in which we evaluated patterns of mortality and growth in the trichopteran shredder *Pycnopsyche guttifer* (Limnephilidae) under varying conditions of animal density, exposure to potential predation, and food quality. The experiments involved a combination of laboratory culture and field enclosure techniques, and free-ranging animal releases into natural stream reaches. All the larvae used in the experiments were derived from field collected egg masses, which are laid terrestrially and contain an average each of about 250 eggs.

In laboratory cultures, larvae were reared individually (with no potential for competition or predation), with excess food of different qualities (fast, medium, and slowly decomposing leaf litter discs, in which fast = high quality). In the field, entire egg masses were either held in enclosures (excess food, no predation, potential for intra-specific competition), or were introduced free into the environment. The field studies were conducted in two 100 m reaches, separated by a 500 m. buffer, of a second-order woodland stream which did not contain the species. In the upper reach, 20 egg masses were placed along the shore, two masses every 10 m one on each side of the stream, and replicate enclosures were started with one egg mass each. With an expected mortality (from previous lab studies) of 90% from egg to adult, this stocking density would potentially yield a typical terminal instar density of about one larva per meter square. In the downstream reach, both the free-ranging and enclosed larvae were introduced at 5 times the density of the upper reach. That is, 100 egg masses were introduced, with one mass each meter on alternating sides of the stream; enclosures were started with five egg masses each. In lab and in field experiments, larvae were periodically censused to measure survival and growth. Censuses of free-released larvae involved establishing 10 randomly located 0.5 m- wide transects across the channel width at each census period, in which all CPOM (coarse particulate organic matter - primarily leaf litter) was collected with 500 µm mesh nets. The total CPOM collected averaged several hundred grams dry weight per transect). The litter was sorted by hand to fast, medium, or slow processing categories (i. e. leaf species that have known in-stream processing rates) and *Pycnopsyche* and other shredders were removed, weighed, and instar determined. Each field sample required approximately 20 person hours to process.

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

yes

Among the free-released larvae, initial survival was greater in the upper (1x density) than in the lower (5x density) reach (Fig. 1). Differential downstream displacement between the two reaches has been ruled out as a possible explanation for the higher upstream survival because no significant trends in downstream displacement were detected in either reach. By 500 degree-days, however, differences were statistically indistinguishable; in both cases survival was less than 10%.

The field enclosures showed the same survivorship pattern as the free-ranging population (Fig. 2). Mortality rapidly erased initial differences in stocking densities; after 400 degree-days, % survival was the same in upstream and downstream reaches. A comparison of the total (upstream and downstream reaches combined) free-ranging population and the enclosed larvae revealed that the pattern of survivorship was similar and statistically indistinguishable after 400 degree-days.

Fig. 1. Survivorship of free-ranging larvae in upper (1x density) and lower (5x density) 100 m stream sections.

Fig. 2. Survivorship of larvae confined in field enclosures.

There was greater survival initially in the lab cultures fed the lower quality food, but by 700 degree-days, differences in survival between *Pycnopsyche* larvae fed low and high quality food disappeared (Fig. 3). This same inverse relationship between growth/molting and survivorship observed initially in the lab cultures of *P. guttifer* had been reported earlier in studies with mayfly species (Cummins and Wilzbach 1988). Food quality did, however, significantly effect temperature-specific mean individual weight at the end of the experiment.

When all treatments are compared, the general pattern of mortality was similar, as shown by the exponential fit in Fig. 4. By 500-600 degree-days, survivorship was less than 10%, and about 1% in all treatments after 1000 degree-days (essentially the entire growth period). The majority of the mortality occurs in the first two or three larval instars. Given that the same general survivorship pattern occurs irrespective of larval density, food availability, and the presence of potential predators, our conclusion is that mortality in these invertebrates cannot be driven solely by factors like predation or competition. This is not to say that predation or competition are not important in nature; our data suggest, however, that when the potential for predation, etc. is prevented - the same mortality curve occurs as when these factors may potentially operate. Therefore another regulatory agent must be involved, perhaps acting in concert with other potential mortality agents; our hypothesis is that this involves pathogenesis.

Fig. 3. Survivorship of lab-cultured larvae related to food quality.

Fig. 4. Survivorship curves for lab cultures, field enclosures, and free-ranging populations of *P. guttifer*.

We have begun some preliminary studies to identify the microbial inhabitants of *Pycnopsyche* digestive tracts. Microbes were found in the fore- mid- and hindguts in dissections of live, 2nd-3rd instar *Pycnopsyche*. Our qualitative observations are that the microbial community within the midgut differs in composition among the fore, mid, and anterior portions.

LITERATURE CITED

Cummins, K.W. and M.A. Wilzbach. 1988. Do pathogens regulate stream invertebrate populations? *Verh. Internat. Verein. Limnol.* 23: 1232-1243.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

END

**DATE
FILMED**

8 / 25 / 92