DOE/ER/13880-5

DOE/ER/13880--5

DE93 004650

Late Transition Metal Oxo and Imido Complexes

Progress Report

for

Budget Period

May 15, 1992 to May 14, 1993

**Paul R. Sharp
University of Missouri-Columbia**

December 1992

Prepared for

**THE U. S. DEPARTMENT OF ENERGY
GRANT NO. DE-FG02-88ER13880**

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

for

Abstract

A range of new phosphine Au(I) oxo complexes, $[(\text{LAu})_3(\mu\text{-O})]^+$, were prepared, testing the limits of the phosphine steric and electronic properties. Their reaction with reducing agents (hydrazines, CO and RNC) give a variety of new and known gold clusters in unprecedented high yields and establish the steric and electronic relationships between the phosphine and the nuclearity of the product Au cluster. The $[\text{L}_2\text{Pt}(\mu\text{-O})]_2$ class of dioxo complexes has been expanded to include the first anionic oxo complex, $\{[(\text{dppm-H})\text{Pt}(\mu\text{-O})]_2\}\text{Li}_2$ and the first Pt imido complexes, $[(\text{L}_2\text{Pt})_2(\mu\text{-O})(\mu\text{-NR})]$ and $[\text{L}_2\text{Pt}(\mu\text{-NR})]_2$. The previously observed Rh imido/amido A-frame chemistry has been extended to Ir. The p-MePh-imido complex has been characterized by X-ray diffraction and clearly shows the resonance delocalization of the nitrogen lone pair into the aryl ring which is responsible for the previously observed electrophilic ring addition and oxidative ring coupling reactions.

Project Description

This project is a study of the synthesis and reactivity of oxo, imido and related complexes of the second and third series metals of groups 9-11. Our interest in these complexes arises from 1) our desire to model surface species present in catalytic reactions (ethylene epoxidation on Ag,¹ NH_3 oxidation on Pt,² NO reduction by CO on Rh,³ O_2 electroreduction on Pt⁴) and 2) the unique reactivity associated with complexes of "hard" ligands with "soft" metals.⁵

Results and Discussion

Au Complexes. Last year we reported the remarkable ability of the gold oxo complex, $[(\text{LAu})_3(\mu\text{-O})]^+$ ($\text{L} = \text{PPh}_3$), to produce hydrazido complexes which decompose in high yields to the Au cluster, $[(\text{LAu})_6]^{2+}$. We have continued to investigate this chemistry and have synthesized a very large range of oxo complexes by variations of the phosphine ligand L. This series covers almost the whole available range of steric and electronic properties for monodentate phosphines. These new oxo complexes also react with hydrazines, CO and RNC to produce new clusters. We have been able to study the phosphine dependence of the cluster produced and have found that steric factors predominate, large phosphines give Au_6 clusters and small phosphines give Au_{11} clusters (Scheme I). Electronic factors do not seem to be significant. Although, with the more

Scheme I

unusual phosphines the nature of the clusters remains to be determined. This chemistry was presented this past summer at an international conference, attended by renowned experts on gold cluster chemistry (H. Schmidbaur and M. Mingos), and was very well received.

Pt Complexes. The $[L_2Pt(\mu-O)]_2$ class of oxo complexes⁶ has been expanded to include the first anionic oxo complex, $\{[(dppm-H)Pt(\mu-O)]_2\}Li_2$ (eq 1 and Figure 1a) and the first Pt

Figure 1

imido complexes, $[(L_2Pt)_2(\mu-O)(\mu-NR)]$ and $[L_2Pt(\mu-NR)]_2$ (Scheme II). Although the

Scheme II

imido complexes have not yet been structurally characterized, the precursor amido complexes, which are novel complexes themselves, have been (Figure 1b). The anionic complex forms good crystals allowing us to determine the solid-state structure of the second oxo complex of this type. In addition, we suggested previously the possibility of forming anionic oxo complexes in other systems. The discovery of this Pt complex encourages us to try to produce anionic oxo complexes in other metal systems.

The synthesis of the imido complexes, $[(L_2Pt)_2(\mu-O)(\mu-NR)]$ and $[L_2Pt(\mu-NR)]_2$, is very exciting. We know from our A-frame Rh and Ir work that the imido complexes have a much richer chemistry than the oxo analogues. This is a result of a number of factors, one of which is the apparent weaker M-N bond. Thus, we anticipate much greater reactivity for these imido complexes.

Ir Imido/Amido A-frame Complexes. As reported last year, tautomeric mixtures of $\text{Rh}_2(\mu\text{-NR})(\text{CO})_2(\mu\text{-dppm})_2$ and $\text{Rh}_2(\mu\text{-NHR})(\text{CO})_2(\mu\text{-dppm-H})(\mu\text{-dppm})$ (**1/2**, $\text{R} = \text{Ph}$, $p\text{-MePh}$) add carbon based electrophiles to the para position of the imido aryl ring to give the ring addition products, $[\text{Rh}_2(\mu\text{-N}(4,4\text{-R,R'}\text{-C}_6\text{H}_4)(\text{CO})_2(\mu\text{-dppm})_2]^+\text{X}^-$ **3** (eq 2). We have extended this

chemistry to Ir and have determined the solid state structure of the imido tautomer which clearly shows the electron density transfer from the nitrogen lone pair into the ring which results in the activation of the ring to electrophilic attack. We have also begun exploring the reaction chemistry of the Ir imido/amido complexes with CO and find that the isocyanate complexes are much less stable than they were with Rh. This could result in much higher yields of free isocyanate. This has important implications for the synthesis of isocyanates from the carbonylation of nitro compounds.

References

1. Grant, R. B.; Lambert, R. M. *J. Chem. Soc., Chem. Commun.* **1983**, 662-663.
2. Asscher, M.; Guthrie, W. L.; Lin, T.-H.; Somorjai, G. A. *J. Phys. Chem.* **1984**, 88, 3233-3238.
3. Lorimer, P.; Bell, A. T. *J. Catal.* **1979**, 59, 223. Hecker, W. L.; Bell, A. T. *J. Catal.* **1983**, 84, 200.
4. Hoare, J. P. *The Electrochemistry of Oxygen*; Wiley-Interscience: New York, 1968; pp 143-151.
5. Pearson, R. G. *J. Am. Chem. Soc.* **1963**, 85, 3533 and *J. Chem. Ed.* **1968**, 54, 643.
6. Li, W.; Barnes, C. L.; Sharp, P.R. *J. Chem. Soc., Chem. Commun.* **1990**, 1634-1646.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

Publications for Period May 15, 1992 to May 14, 1993 (copies attached)

"Late Transition Metal μ -Oxo and μ -Imido Complexes. 8. Rhodium Imido/Amido Coupling Reactions with Carbon Monoxide" Ge, Y.-W.; Sharp, P.R. *Inorg. Chem.* **1992**, *31*, 379-384.

"Preparation and Decomposition of Gold(I) Hydrazido Complexes: Gold Cluster Formation" Ramamoorthy, V.; Wu, Z.; Yi, Y.; Sharp, P. R. *J. Am. Chem. Soc.*, **1992**, *114*, 1526-1527.

"Rhodium Arylimido/Arylamido Complexes: Electrophilic Additions" Ge, Y.-W.; Sharp, P.R. *Inorg. Chem.* in press.

"Chemistry of $[\text{Cp}^*\text{Rh}(\mu\text{-Cl})_2]$ and Its Dioxygen and Nitrosobenzene Insertion Products" Hoard, D. W.; Sharp, P. R. *Inorg. Chem.* in press.

"Late Transition Metal Oxo and Imido Complexes. 11. Gold (I) Oxo Complexes" Yang, Y.; Sharp, P.R. *Inorg. Chem.* submitted.

Preprints & reprints
removed & cycled
separately

END

**DATE
FILMED**

1 / 11 / 93