006123

1 OF 1

DOE/ER/54071-

DOE/ER/54071--T3

DE93 006123

FINAL REPORT

DOE PROJECT TITLE: SALIENT ISSUES OF EDGE PHYSICS
PERTAINING TO LOSS OF CONFINEMENT:
A RESISTIVE MHD ANALYSIS

DOE GRANT #: DE-FG03-90ER54071

PI: DR. DAVID R. THAYER

ORGANIZATION: SAIC, SCIENCE APPLICATIONS
INTERNATIONAL CORP.

DATE: APRIL 23, 1992

CONTENTS:

FY91 PROGRESS REPORT

FY90 PROGRESS REPORT

THERMALLY DRIVEN TURBULENCE AND
TRANSPORT IN A SHEARED SYSTEM REPORT

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

for

TECHNICAL PROGRESS REPORT - FY90

DOE PROJECT TITLE: ALIEN ISSUES OF EDGE PHYSICS PERTAINING TO LOSS OF CONFINEMENT: A RESISTIVE MHD ANALYSIS

DOE GRANT #: DE-FG03-90ER54071

PI: DR. DAVID R. THAYER

**ORGANIZATION: APPLIED PLASMA PHYSICS DIVISION
SCIENCE APPLICATIONS INTERNATIONAL CORP.**

TECHNICAL PROGRESS:

In general we have made significant contributions towards the ultimate goal of creating a complete theory of edge turbulence and transport in magnetic fusion devices. Our main focus has been to utilize a resistive MHD model. This analysis includes: 1) "rippling" type models in which the current fluctuations are decoupled and the resistivity fluctuations are fundamental, and 2) "drift" type models in which the ambient current can be small (no resistivity fluctuations needed) and the current fluctuations are fundamental. In these turbulence frameworks, the important atomic physics based edge sources have been included (impurity radiation, ionization, and charge exchange effects).

In collaboration with our analytic theoretical analysis are: experiments which have been conducted on the TEXT tokamak (Fusion Research Center - University of Texas at Austin -- Dr. Alan Wootton's group) and the ATF stellarator (Oak Ridge National Laboratory), numerical simulations using 3D MHD codes which run on the MFE Cray computer system (Oak Ridge National Laboratory -- Dr. Ben Carreras' group), and theory conducted at the University of California at San Diego (Prof. Pat Diamond's group). It should be noted that this three-way collaboration on edge physics (theory-simulation- experiment) has been very successful in the transport task force (TTF) program [refer to the synopses of the TTF meetings in Austin (January, 1989), San Diego (August, 1989), and Hilton Head (February, 1990)]; in fact, the edge group has been noted as a good paradigm for other groups that should form to address different regions of the confinement problem.

The research topic which we have primarily focused on is the impurity radiation

effects on the edge fluctuations and transport. The important conclusion which we have ascertained is that the condensation part of the impurity radiation driven edge turbulence is not present in the edge of fusion devices (note: the results of this analysis are detailed in the enclosed report -- which is to be published shortly). This is a striking result since the condensation piece of the impurity radiation enhancement has been previously thought of as the "robust" part of the instability (note the publications: Thayer, et al., J. Nucl. Mater., 1987; Drake, Phys. Fluids, 1987; Thayer and Diamond, Phys. Fluids, 1987; Drake, et al., Phys. Fluids, 1988; and Thayer, et al., Plasma Phys. and Contr. Nucl. Fus. Res., 1989). It is important to realize that the literature in both the solar physics and the fusion physics areas (for the past twenty years) has indicated that there are two major parts to the radiative instabilities (i.e., the thermal instability and the condensation instability). Since the thermal instability depends on the gradient of the radiation profile (with respect to temperature), and thus depends on the details of the radiation profile, this drive has been thought of as the "fickle" instability. In particular, if one considers the realistic multiple impurity model (where various impurities are dominant at different radii), then the thermal instability drive can be positive or negative, depending on the particular circumstance. However, the condensation instability has been previously thought of as quite "robust", since the drive is only dependent on the presence of the radiation rate (and thus is not dependent on the details of the impurity radiation cooling profile). The upshot of our analysis is that we realized that the assumption (parallel pressure balance) used in deriving the condensation instability for the edge fluctuation region was not correct. In this case the parallel pressure balance was violated since the phase velocity of the waves are greater than the sound speed over most of the eigenmode structure.

In order to rectify the problem with the condensation part of the impurity radiation drive (stated above), we have enhanced our fundamental model of edge turbulence and transport by formulating the radiative driven drift-rippling mode model. In essence, this fluctuation model contains equations which govern the coupling between the potential, the temperature, the density, and the parallel velocity fluctuations. The key ingredient in this more complete analysis is that the parallel pressure balance is not assumed a priori. In this case (for sheared magnetic fields in fusion devices) the parallel velocity equation evolves the parallel pressure gradient (through the dynamics of the temperature and the density) so that this important term is not trivially set equal to zero. This fundamental model of the edge includes the temperature gradient, the density gradient, the impurity radiation drive, and the parallel conduction dissipations. Given this model, we calculate (using a renormalized theory) the nonlinear saturated state of fluctuations and transport. The interesting result is that the condensation effect is shown to be lower than the thermal instability by an order of magnitude for typical TEXT parameters. The analytic expressions for the

fluctuation amplitudes and the transport coefficients have been derived for the saturated turbulent state, locally, about any mode rational surface position. In addition, we have calculated the results globally by using the TEXT profile information (current, temperature, density, and impurity radiation) for the entire edge region. Note that the detailed results are presented in the accompanying research report. Our results compare favorably with the fluctuation levels and transport coefficients determined from probe measurements on the TEXT tokamak. The recent simulation results from ORNL indicate consistency with our numerical values. In addition, from the simulation results there is evidence that the edge spectra peaks at the thermal instability drive position (maximum radiation gradient) and not the condensation drive position (maximum radiation). Consequently, the simulation has produced evidence that the condensation instability is in fact small, which is in agreement with our analytic theory. However, it should be noted that more simulations need to be performed in order to confirm this effect. Also, since the radiation profiles from the experiments are still in question as to the detailed structure, we cannot, at this point, state that there is concrete experimental evidence that the condensation instability is not present in the edge of fusion devices. Nevertheless, it is clear from our calculations that for the situation in which the condensation effect is small and the "fickle" thermal instability is the remaining dominant impurity radiation enhancement, that it is important to explore other edge sources and the appropriate edge models which lead to other significant edge transport enhancements.

In addition to our detailed calculations performed on the radiative drift-rippling mode system, we have performed preliminary calculations for the radiative drift wave model and the ionization drift wave model. The radiative drift wave model includes the thermal instability in an edge model which does not assume the existence of an ambient current. Consequently, this model is relevant for stellarators (such as ATF). The fluctuation and transport results from this theory are very similar to the radiative dominant limit of the radiative rippling mode model; however, the mode frequency corresponds to the drift wave frequency and the mode is heavily damped in the absence of the radiative drive. The ionization drift wave model includes the density source from the ionization of neutrals and the momentum sink from the charge exchange with the neutrals. In addition to the presence of the drift wave frequency and the growth rate corresponding to the ionization rate, the most interesting result is an ionization induced inward particle flux. This result would indicate the existence of an inward particle flux resulting after the occurrence of a gas puff (due to the enhanced neutral density and the resultant enhanced ionization rate).

In conclusion, the status of our comprehensive edge fluctuations and transport model is quite good. Our comparison with the experiments and the simulation results indicate reasonable agreement. However, it is very important to continue this strong collaborative effort in order that ultimately a complete edge theory can be obtained

which will demonstrate a major success for the understanding of magnetic fusion transport. Consequently, the success in edge transport could then be used as a major stepping stone to approach the theoretical understanding of confinement region transport. In order to achieve this goal of a complete theory of edge fluctuations and transport, we suggest that the following topics should be vigorously pursued: 1) Continue the successful work on the radiative drift-rippling mode model with multiple impurities and make comparisons with the simulations and the experiments, 2) Perform an in-depth atomic physics analysis to determine the detailed radiation profiles which should be used for the theoretical edge models and which correspond with the experimental measurements, 3) Continue work on additional atomic physics effects on the edge models (e.g., ionization particle sources and charge exchange momentum sinks), and include the impurity gradient drive in our drift-rippling model, 4) Construct a two-point theory of the edge model in order a) to calculate the predicted spectrum under the driven environment and b) to determine the cross correlations (between potential and density, between potential and temperature, and between density and temperature). It should be noted that item 4) is particularly important since the experimental measurements of all these quantities are now becoming readily available so that this evidence can be used as a critical comparison which will enable us to pinpoint the correct edge turbulence and transport theory.

The work performed under this DOE grant has been presented (up to this date) at the following meetings and seminars:

- Tokamak Edge Turbulent Transport, D. R. Thayer, P. H. Diamond, B. A. Carreras, J. A. Holmes, J. N. Leboeuf, Ch. P. Ritz, and A. J. Wootton, APS meeting, Anaheim, CA (1989).
- Thermally Driven Convective Cell Turbulence at the Edge of TEXT, J. A. Holmes, B. A. Carreras, J. N. Leboeuf, D. K. Lee, Ch. P. Ritz, A. J. Wootton, D. R. Thayer, and P. H. Diamond, APS meeting, Anaheim, CA (1989).
- Resistivity Gradient Driven Turbulence at the Edge of TEXT, J. N. Leboeuf, B. A. Carreras, J. A. Holmes, D. K. Lee, C. P. Ritz, A. J. Wootton, D. R. Thayer, and P. H. Diamond, APS meeting, Anaheim, CA (1989).
- Radiative Enhanced Fluctuation Induced Transport in Sheared Systems (Rippling and Drift Wave Turbulence), Transport Task Force Meeting seminar, Hilton Head, SC (1990).
- Enhanced Edge Transport in Sheared Systems, D. R. Thayer, P. H. Diamond, A. S. Ware, B. A. Carreras, C. Hidalgo, J. A. Holmes, J. N. Leboeuf, Ch. P. Ritz, and A. J. Wootton, Sherwood Theory Meeting, Williamsburg, VI (1990).
- Role of Fueling-Induced Instabilities in Tokamak Edge Turbulence, A. S. Ware, P. H. Diamond, D. R. Thayer, B. A. Carreras, and J. N. Leboeuf, Sherwood Theory Meeting, Williamsburg, VI (1990).
- Modeling of Fluctuations at the Edge of the TEXT Tokamak, J. N. Leboeuf, B. A. Carreras, J. A. Holmes, D. K. Lee, V. E. Lynch, Ch. P. Ritz, A. J. Wootton, P. H. Diamond, A. S. Ware, and D. R. Thayer, Sherwood Theory Meeting, Williamsburg, VI (1990).
- Radiative Enhanced Turbulence and Transport in a Sheared System, UCLA, Institute of Plasma and Fusion Research seminar, Los Angeles, CA (1990).

**DATE
FILMED**

2/10/93