PROGRESS REPORT

MASTER

UNIVERSITY OF MINNESOTA
SCHOOL OF PHYSICS AND ASTRONOMY

EXPERIMENTAL AND THEORETICAL HIGH ENERGY PHYSICS
U. S. DEPARTMENT OF ENERGY CONTRACT NO. EY-76-C-02-1764

OCTOBER 1, 1979 - SEPTEMBER 30, 1980

DISCLAIMER

This book was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency Thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

1. Introduction

During 1980, the high energy physics program at the University of Minnesota experienced a qualitative change in its character. The experimental program expanded to include major efforts on frontier experiments in three different laboratories. The Fermilab charged-neutral spectrometer published its first definitive results on the electromagnetic couplings of vector mesons, while securing approval for a new effort, aimed at use of the Tevatron, to test QCD by measuring direct photon production. The first Minnesota baryon decay detector was installed in the unique Soudan Underground Research Site in northeastern Minnesota in order to test unified field theories by studying the long-term stability of matter. Construction of equipment was also begun for Brookhaven AGS experiment 755, which also seeks to test QCD through the examination of large transverse momentum exclusive processes. Simultaneously, the extremely fruitful, decade-long program at the Argonne ZGS was concluded by the publication of new results obtained with the polarized proton beam.

This large increase in experimental activity has been complemented by additional theoretical efforts to understand data measured by the Minnesota and other experimental groups. This higher level of activity in both theoretical and experimental high energy physics has been recognized by the University's authorization of new faculty positions in each area. The theorist, now being sought, would mark the first faculty addition to this group in a decade.

It is useful at this turning point to recount briefly the historical context for this rapid increase in activity. High energy physics at

Minnesota started in 1960 as an appendage to the theoretical nuclear physics effort. The first experimental efforts began in 1963 and were limited to analysis of bubble chamber film. Between 1968 and 1975, three counter/chamber experiments were performed at the Argonne ZGS. In these experiments, the group gained expertise in the use of modern, online computer-controlled proportional chamber techniques. Simultaneously, it became clear that a small bubble chamber group could not make a meaningful contribution, so this activity was discontinued. During the latter 1970's the group performed a series of experiments at the Argonne ZGS.

Particularly notable were a detailed measurement of pion-proton elastic scattering both to search for cross-section fluctuations and to measure the s dependence of the large-angle cross-section, an experiment to measure direct electron production and a series of experiments with the polarized proton beam to investigate the effects of spin on the strong interaction. One of these polarized experiments revealed previously unsuspected asymmetries in the inclusive production of pions in polarized proton-proton collisions. Another series of spin-dependent experiments revealed considerable detailed information about nucleon-nucleon scattering amplitudes and the multiple scattering of protons from deuterium and helium at intermediate energies. In 1978, the Minnesota group began a Fermilab experiment on the electromagnetic couplings of vector mesons, which complemented work done at Argonne in 1968. At the same time, Minnesota began to play a significant role in the long-standing Fermilab hyperon collaboration.

One of the reasons for the success of the Minnesota experimental group has been its ability to resist fragmentation of its efforts. Minnesota is quite unique in that all of its high energy physicists play a role in each

major experimental project. This approach leads to a flexibility of response, which is a major asset in research whose schedule is often controlled by external forces. At the current time, the group's focus is on the Soudan baryon decay experiment, with other activities targeted at Fermilab and Brookhaven. Over a period of months, in response to new needs, this focus could change in a continuous and efficient way.

A description of both the theoretical and experimental efforts under this contract for the year 1980 is included in this report.

Nonabelian Stokes Theorem (S. Gasiorowicz with P. Fishbane, U of Virginia and P. Kaus, U. C. Riverside).

As part of our study of the Wilson loop operator

$$W(C) = \text{Tr} \left(P \exp \oint_C dx^\mu A_\mu^a(x) \lambda^a/2 \right)$$

we have found a nonabelian generalization of the Stokes theorem. Two approaches were used: In the first, we used a differential equation technique to deform the contour C to include an infinitesimal element of area $d\sigma_{\mu\nu}(x)$ on an arbitrary surface spanning the contour. Using

$$P \exp \oint_{(y)} dx^\mu \underline{A}_\mu \approx 1 + d\sigma_{\mu\nu}(y) \underline{G}^{\mu\nu}(y)$$

where

$$\underline{G}^{\mu\nu} = \partial^\mu \underline{A}^\nu - \partial^\nu \underline{A}^\mu - [\underline{A}^\mu, \underline{A}^\nu] ; \quad \underline{A} = A^a \lambda^a/2$$

for the infinitesimal loop, we find

$$W(C) = \text{Tr} \int d\sigma_{\mu\nu}(y) U_{Ay} \underline{G}^{\mu\nu}(y) U_{yA}$$

where

$$U_{Ay} = P \exp \int_y^A dx^\mu \underline{A}_\mu = U_{yA}^{-1}$$

is a line integral along an arbitrary path from y to an arbitrary fixed point A . This reduces to the usual Stokes' theorem for an abelian theory. The surface integral is independent of the choice of surface for fields satisfying the field equations

$$D^\mu \underline{G}_{\mu\nu} = \partial^\mu \underline{G}_{\mu\nu} - [\underline{A}^\mu, \underline{G}_{\mu\nu}] = 0$$

In the second approach we used a series of Baker-Hausdorff type identities to obtain a particularly simple expansion that sums to the expression

obtained above. For a rectangular area in the (1,2) plane, for example, with one corner at (x_1, x_2) and the opposite corner at $(x_1 + a, x_2 + b)$ the expansion has the simple form

$$W(C) = \text{Tr} \exp \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} \frac{a^m}{m!} \frac{b^n}{n!} D_1^{m-1} D_2^{n-1} G_{12}$$

A paper describing these results is in preparation.

Connection between Asymptotic Behavior and Bound States in QCD (S. Gasiorowicz with P. Fishbane, U of Virginia and P. Kaus, U. C. Riverside).

We have been continuing our work on extracting information on the hadronic spectrum from the two-point function. The basic method is one of constructing a function that is cut-less and that differs from the true function asymptotically less than any power of q^2 . The result is an infinite set of moment conditions; when CDD-type ambiguities are removed, these can be solved and in lowest order place the masses at locations of the zeros of a Bessel function.¹

We have improved the basic method by extracting branch-point singularities that are related to the threshold behavior. In particular this necessitates a treatment of spin 1/2 constituents that is somewhat more complicated than the treatment for spin 0 constituents, basically because threshold behavior at $E = \pm m$ needs to be treated properly. The new moment conditions lead to solutions that mimic the naive bag model. We have also calculated corrections due to 4th order perturbative modifications of the two-point function. A paper describing this work is in the final stages of preparation.

¹D. Beavis, S. Y. Chu and P. Kaus, Ann. Phys. N. Y. (in press).

Classical Solutions for Gauge Fields Interacting with Higgs Mesons

(D. A. Geffen).

I am reexamining the classical solutions to the equations for Non-Abelian and Abelian gauge fields interacting with Higg's mesons, in particular the t'Hooft monopole [Nucl. Phys. B79, 276 (1974)], and the Julia-Zee Dyon [Phys. Rev. D11, 2227 (1975), and Prasad-Sommerfield, Phys. Rev. Lett. 35, 760 (1975)] for the non-abelian theories and Nielson-Olesen solutions in the Abelian theory.

(a) Non-Abelian case.

The difference between the solutions in the Abelian and non-Abelian theories is dramatic. In the former case, the magnetic field (a vortex line in a superconductor for example) falls off exponentially, characteristic of the mass generated by the spontaneous symmetry breaking due to the Higgs field. In the Non-Abelian case, the generalization of the Nielsen-Olesen solution by t'Hooft, an "electromagnetic" field is obtained which falls off like $1/r^2$ as $r \rightarrow \infty$ and includes a "magnetic field", " $\vec{B}$ " = $-\frac{\hat{r}}{er^2}$, which is identified as due to a point magnetic monopole of strength $1/e$. There is a problem with such an identification however. Although the monopole is a point magnetic charge, its energy is finite and easy to estimate numerically! This is an extraordinary result which requires a better understanding. The identification of the "electromagnetic" field by t'Hooft is made by constructing a gauge invariant anti-symmetric tensor $F_{\mu\nu}$ which reduces to a conventional looking electromagnetic field tensor when a gauge transformation is made. (The Higg's field, $\phi^a = X_a F(r)$, $a = 1,2,3$, is rotated by a gauge transformation to $\phi^{a'} = (0,0,rF)$ which is the conventional choice for the axis of spontaneous symmetry breaking in the $SU(2) \times U(1)$ theory). In this gauge, a vector potential A_μ is generated ($A_\mu = W_\mu^{a'}$, $W_\mu^{a'}$ the iso-rotated vector

potential triplet) such that $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$. The disturbing fact is that $\vec{A}$ depends only on the gauge transformation and not on the original fields W_μ^a . Another way of seeing this is to start with the Wu-Yang-t'Hooft ansatz

$$\begin{aligned} W_i^a &= \epsilon_{aij} x_j W(r) \\ W_0^a &= 0 \\ \phi^a &= x_a F(r) \end{aligned}$$

then
$$F_{\mu\nu} \equiv \frac{1}{\sqrt{\phi^a \phi^a}} \phi^b G_{\mu\nu}^b - \frac{1}{e[\phi^a \phi^a]^{3/2}} \epsilon_{abc} \phi^a (D_\mu \phi^b)(D_\nu \phi^c)$$

reduces identically to

$$F_{ij} = \frac{[\text{sign } F(r)] \epsilon_{ijk} x_k}{e r^3}, \quad F_{0\mu} = 0$$

for arbitrary $W(r)$ and $F(r)$.

If we identify F_{ij} as the magnetic field ($F_{ij} = \partial_i A_j - \partial_j A_i$) in the gauge where $\phi^a = (0, 0, rF(r))$ then the magnetic field energy is infinite due to the point magnetic charge at $r = 0$. The field energy, however, depends quadratically on W and F and is scaled to the mass m , where $\lim_{r \rightarrow \infty} \phi^a = m x_a / r$, the vacuum solution for ϕ^a . Since solutions to the non-linear field equations for W and F exist which are regular at $r = 0$ and yield finite total field energy the physical interpretation of the field F_{ij} is puzzling. It is clear, however, that the point (magnetic) charge singularity is an artifact of the

particular gauge chosen and not an indication of the existence of a particle. There is, nevertheless, present a finite, long-ranged solution to the field equations whose physical interpretation needs further exploring. Work is in progress along these lines of investigation.

(b) Abelian case.

This case is of particular interest because of the work of Raby and Ukawa (Phys. Rev. D18, 1154 (1978)) who apply the solutions of Nielsen and Olesen [Nucl. Phys. B61, 45 (1973)] to the (1+1) dimensional Abelian gauge theories where the vortices become instantons along the imaginary time axis. Raby and Ukawa make use of the limit when the Higg's boson mass, m_s , is much larger than the vector boson mass, m_w , (type II superconductor) to construct an approximate solution to the field equations in order to evaluate the action. Their results were important for giving an understanding of the role of a dilute instanton gas in spontaneous gauge symmetry breaking. The approximate solution they use, however, does not take into account a non-linear, inhomogeneous term in the equation for the Higg's field. This term, quadratic in the vector field, is a source term for the Higg's field and dominates for large r in the large Higg's mass limit they consider. Consequently, the variation in the Higg's field extends well beyond the region characterized by $1/m_s$. A preliminary analysis indicates that the action is reduced by this effect and may alter their results. Using approximation techniques, I find that the Higg's field extends much further than $1/m_s$ but does reach its vacuum value when $r \sim 1/m_w$. The Raby-Ukawa results will be reexamined as part of an ongoing program to examine the classical solutions to gauge field theories.

Strange Quark Mixing In Isosinglet Bosons (D. A. Geffen and W. Wilson).

Strange quarks, s , are much heavier than up and down quarks, u and d .

Consequently the lightest $I = 0$ mesons should be predominantly $u\bar{u}$, $d\bar{d}$ pairs with a heavier counterpart consisting of $s\bar{s}$. The lighter $I = 0$ mesons should have a near degenerate $I = 1$ partner. Pure $u\bar{u}$ or $s\bar{s}$ states are not possible because the pairs can make transitions into one another by annihilating into gluons. There are two well-known examples of such states. The vector mesons, (1^{--}) , ω , ρ are $I = 0, 1$ degenerate partners with ϕ the $s\bar{s}$ heavier counterpart. ϕ clearly contains only small amounts of $u\bar{u} + d\bar{d}$ since its decay into pions is strongly inhibited. The 2^{++} group of f_0 , A_2 and the heavier f_0' ($\Gamma(f_0' \rightarrow 2\pi) \ll \Gamma(f_0' \rightarrow K\bar{K})$) is another good example. The question becomes interesting when we note that the pseudoscalars, π , η , η' do not fit into this pattern. This indicates that some new dynamics are at play. Are there any other examples of low lying mesons that fit the pattern of the pseudoscalars?

We have been examining the 1^{++} system: $D(1285)$, $A_1(1200)$, and $E(1420)$. While the near degeneracy of the D and A_1 fits the picture seen for 1^{--} and 2^{++} , the claim by experimenters that the heavier 1^{++} E frequently decays into $S\pi$, which are composed of pure light quark states, seems to break the pattern. We have been examining these results and find that serious doubts must be raised about the evidence for this decay mode of the E . Our doubts are reinforced by the results of one experiment claiming the existence of a 0^{-+} state degenerate with the D and most likely one degenerate with the E as well. We believe therefore, that the pattern is not broken by the axial vector system.

The pseudoscalars are $q\bar{q}$ 1S_0 states, while the above 1^{--} , 2^{++} , and 1^{++} are all triplet states. It is therefore of interest to examine the 3P_1 , 3D_2 , etc. states to see which pattern they follow. The $I = 1$ $B(1235)$ is a well-established 3P_1 state (1^{+-}) with the decay $B \rightarrow \omega\pi$ the only decay mode observed. An experiment identifying an $I = 0, 1^{+-}$, the H , has just been reported

with mass 1190 ± 60 MeV and width 280 MeV, $\Gamma_H \leq 380$ MeV.

The near mass degeneracy of the B and H strongly supports the pattern observed for the triplet $q\bar{q}$ states. We further note that with $H \cong \cos\alpha \left(\frac{u\bar{u}+d\bar{d}}{\sqrt{2}} \right) + \sin\alpha \ s\bar{s}$, and with the plausible assumption that $H \rightarrow \rho\pi$ dominates the H decay just as does $B \rightarrow \pi\omega$, we can find a range for $\sin\alpha$ by using the new data for Γ_H and the well known value for Γ_B . We find $0 \leq \sin\alpha \leq 0.25$, giving strong support that the H meson is predominantly composed of light quarks and thereby follows the pattern observed for the triplet spin $q\bar{q}$ states. The pseudoscalar mesons remain unique in their quark structure. The detection of the $I = 0 \ 2^{-+}$ states is now of great interest since these are the first Regge recurrences of the γ and γ' mesons.

Neutrino Oscillations (D. A. Geffen and Hiroshi Suura).

There recently has been renewed interest in the possibility that the ν_e , ν_μ , ν_τ , neutrinos have small but finite masses and that the neutrino mass eigenstates will be mixtures of the above states. Neutrinos produced in a particle decay will, therefore, be mixtures of these mass eigenstates. When the masses of two neutrinos, call them ν_1 and ν_2 for convenience, are both very small (the order of electron volts) they will propagate in overlapping wave-packets over great distances and interact with a target hadron at the same time. If both ν_1 and ν_2 are mixtures of ν_μ and ν_e , for example, both neutrinos will undergo the same reaction, i.e.,

Under certain conditions, the incoming neutrino state will be a coherent mixture of ν_1 and ν_2 . In the plane wave approximation the relative phase of the mixture will oscillate in time or, equivalently, by path length, with phase $\Phi = (\omega_1 - \omega_2) t$, ω_j the energy of ν_j . We have been examining in detail the conditions for producing such a coherent state, $\psi_1 |\nu_1\rangle + \psi_2 |\nu_2\rangle$, and the conditions for ν_1 and ν_2 to produce an identical final state so that the cross section for reactions such as (1) will contain an oscillating interference term. In the discussion below we have restricted ourselves to the case of two mixing neutrinos.

It is clear that if the source of the neutrino state, such as a pion, (i.e., $\pi^+ \rightarrow \nu_j + \mu^+$) has a sufficiently large uncertainty in energy and momentum, then the two neutrinos ν_1 , ν_2 will be in a coherent wave packet state -- namely the pion state can decay into either a ν_1 or ν_2 and the same muon state. Similarly, if this coherent mixture of ν_1 and ν_2

wave packets as well as the wave packet of the target state have momenta spreads $\Delta p \gg |\omega_1 - \omega_2|$ ($\hbar = c = 1$) then it is obvious that ν_1 and ν_2 can produce the same final state with a definite phase relation, $(\omega_1 - \omega_2)t$, so that the reaction cross section contains an interference term proportional to $\cos(\omega_1 - \omega_2)R$, R the mean distance between neutrino source and target.

Writing the cosine as $\cos(2\pi R/L)$, we define the coherence length L , [Bilenky and Pontecorvo, Physics Reports 41, 225 (1978).], $L = 2\pi / |\omega_1 - \omega_2|$, $L \approx 4\pi R / |m_1^2 - m_2^2|$ for $k \gg m_j$.

It is quite trivial to note as well, that if the source or target is spread out over a size $D \gg L$, then the interference term must average out to zero.

The case of particular interest, therefore, is when $D \sim L$ so that interference is present but is reduced by the spatial spread of target and/or beam.

We have examined the model where we describe the neutrinos as a coherent superposition of gaussian wave packet states colliding with a target that is also taken to be in a gaussian wave packet state. The two neutrino packets overlap exactly in phase at $t=0$. When the centers of their packets are a distance R from the center of the target packet. We assume a two body reaction with the target and calculate the transition probability using T-matrix theory. The crucial point is not to neglect the variation of the energy denominators with particle energies over the energy-moment spread of the packets. We then obtain an interference term with a phase proportional to both the central values of ν_1 and ν_2 's energy and momentum difference. The latter appears since, in general, the average momenta of ν_1 and ν_2 can be different. We find that the interference term is damped by a product of three gaussians. Two factors replace the energy-momentum conserving delta functions and fix the distribution of total momentum and energy of the final

state. If the average momenta of ψ_1 and ψ_2 are not equal, however, we are left with a net damping factor after integrating over final state momenta.

The remaining gaussian factor depends, roughly, on the ratios of the sizes of the particle wave packets to the coherence length L .

When the lifetime of the source is comparable or shorter than 10^{-8} sec, the pion lifetime, we must take it explicitly into account. Calculation of the damping factor in the interference becomes much more difficult. Further calculations along these lines will be carried out if the experimental situation appears favorable. If so, we might then hope to learn some interesting things about the localization of particle states in matter.

Relativistic wave equation and mass spectrum of Gluonium (Hiroshi Suura).

I studied the spectrum of two-gluon systems in terms of a gauge invariant wave function

$$\psi^{\pm}(1,2) = (\bar{\Psi}_r, \frac{1}{2} \vec{F}_{\pm}^T(1) U(1,2) \vec{F}_{\pm}(2) \Psi). \quad (1)$$

Here $\vec{F}_{\pm}$ is a column matrix with eight field-strength components

$\vec{F}_{\pm}^a = \vec{E}^a \pm i \vec{B}^a$ ($a = 1, 2 \dots 8$), and $U(1,2)$ is an octet string operator $U(1,2) = \exp [i g \int_2^1 T^a \vec{A}^a(x) \cdot d\vec{x}]$, where T^a is an octet representation of the color spin. Thus, the wave equation (1) represents a system of two-

gluons connected by a string. The system is very similar to a light quark-antiquark system except for different spin and color-spin structures. An

equal-time wave equation for $\psi^{\pm}$ can be derived in a way similar to $q\bar{q}$ systems¹ and reads as $[M - V(r)] \psi^{\pm}(1,2) = \pm (\nabla_1 \times + \nabla_2 \times) \psi^{\pm}(1,2)$.

M is the mass eigenvalue and the equation is an obvious generalization of the

Maxwell equation $i \dot{\vec{F}}_{\pm} = \pm \vec{\nabla} \times \vec{F}_{\pm}$ for two-field points. $V(r)$

is a potential consisting of a long-range linear confinement potential $k^2 r$

and a short-range Coulomb potential. It is shown that $V(r)$ is 9/4 times the

corresponding $q\bar{q}$ potential. The factor comes from the difference in the color-

spin. If I use the charmonium potential parameters as reference and put

$$V(r) = \frac{9}{4} (0.2 r - 0.5/r)$$

in GeV units, then the equation gives $M = 2.45$ GeV for the $J = 0$ ground state

of both scalar and pseudo-scalar gluoniums. If I use the potential parameters

appropriate for light quarkonia, then the Coulomb potential becomes too strong

to obtain any reasonable value of M . It is suggested that the Coulomb

potential is actually suppressed to give only a logarithmic singularity. Because of the uncertainty in the short distance interaction I can only conclude that the ground state lies below 2 GeV if the string tension is taken to be $9/4 \times 0.1 \text{ (GeV)}^2$. (Published: Phys. Rev. Letters 44, 1319 (1980)).

1. H. Suura, Phys. Rev. D20, 1412 (1979).

Dynamical structures of the pion (Emmanuel Haqq and Hiroshi Suura).

Previously, one of us (H.S.) derived an equal-time relativistic wave equation of Breit type for a confined $q\bar{q}$ system starting from the QCD Hamiltonian.¹ The linear potential appears as the fourth component of a vector, and he showed how the normalizability condition led to a unique boundary condition, which in turn gave bag-type solutions. Subsequently, he and Geffen solved the wave equation for light $q\bar{q}$ systems² and obtained an almost perfect fit of the light quarkonia spectrum. The key was a very strong Coulomb potential ($g^2/4\pi \sim 2.3$) which pulled down the pion mass to the observed value. We reanalyzed the derivation of the wave equation and found that the correct wave equation is of Salpeter type as far as the Coulomb interaction is concerned, but of Breit type with respect to the linear potential. This new hybrid Breit-Salpeter equation has two remarkable features. First, the Coulomb singularity is suppressed to a logarithmic one, leading to a weaker binding. Secondly, the particular component of the wave function that defines the decay constant is finite at the origin, even though the wave function as a whole is singular at the origin. Thus, we have been able to obtain a reasonable value for the decay constant f_π and the rms radius. On the other hand the pion mass stays above 500 MeV for reasonable parameters. The hyperfine interaction can lower the mass, but it has not been determined yet whether

the magnetic interactions are also suppressed or not.

1. H. Suura, Phys. Rev. D20, 1412 (1979).
2. D. A. Geffen and H. Suura, Phys. Rev. D16, 3305 (1977).

Convergence of Reflectionless Approximations to Confining Potentials

(Jonathan L. Rosner, with Waikwok Kwong, Jonathan F. Schonfeld, C. Quigg, and H. B. Thacker).

Work described in last year's progress report was brought to a conclusion and has been accepted for publication [Ann. Phys. (N.Y.) 127 (1980).¹] The construction of reflectionless potentials supporting a prescribed spectrum of Schrödinger bound states was discussed and related to the inverse problem for confining potentials. A simple formula was derived for the Jost solution in a one-dimensional reflectionless potential with N bound states. This led to compact expressions for the potential and the bound-state wavefunctions in terms of the bound-state energies. For symmetric potentials, N -fold product formulas were obtained for bound-state wavefunctions and their slopes at the origin. Corresponding quantities in a confining potential were expressed in terms of infinite products. Comparison of the finite-product and infinite-product expressions allowed a demonstration of the convergence of the reflectionless results to the confining potential results as $N \rightarrow \infty$. Several sum rules satisfied by the reflectionless potential at the origin were applied to numerical studies of convergence.

Degeneracy in One-Dimensional Quantum Mechanics (Jonathan L. Rosner, with Waikwok Kwong, Jonathan F. Schonfeld, C. Quigg, and H. B. Thacker).

The degeneracy of quantum mechanical energy levels was discussed from the perspective of the inverse scattering problem. The coalescence of energy levels was illustrated for a class of potentials which are completely characterized by their bound-state spectra.²

Review of heavy quarks and new particles (Jonathan L. Rosner).

Some aspects of particles containing c, b, and heavier quarks were reviewed. It was shown how these particles can provide tests of the short-range nature of the strong interactions, and how elementary quantum mechanics can reveal the color and charge of the constituents and the number of states below flavor threshold. Some new results were presented on the inverse problem for confining potentials. Properties of flavored hadrons (masses and lifetimes), of gluonic bound states, and of particles containing very heavy objects were also discussed. This work was presented at the annual meeting of the Division of Particles and Fields, American Physical Society, Montreal, Quebec, Oct. 25-27, 1979,³ and at the Conference on Color, Flavor, and Unification, Nov. 30, Dec. 1, 1979, Irvine, Calif. (unpublished).

Semiclassical results on normalization of bound state wave functions

(Jonathan L. Rosner, with Peter Moxhay).

Some methods were introduced and tested for approximating s-wave Schrödinger wave functions at the origin, and for examining their dependence on reduced mass of the bound system. Methods were discussed both for nonsingular and singular potentials. The approximations, previously tested for power-law potentials, were found to be exact for the Hulthén potential and excellent for a wide range of parameters in Coulomb + linear potentials $V(r) = -A/r + Br$. General theorems for mass dependences were reviewed and extended; these theorems apply to potentials with a definite sign of d^2V/dr^2 . Some specific instances in which d^2V/dr^2 does not have a unique sign, and the theorems do not hold, were also discussed. If the short-distance singularity of V is Coulombic, it was shown under certain assumptions that leptonic widths of vector mesons

should decrease with increasing quark mass no more rapidly than $m_Q^{-1/2}$.⁴

Proton lifetime (Richard Robinett with Jonathan L. Rosner).

Preliminary study of Grand Unified Theories (GUT's) and their applications, especially the SU(5) model and proton decay papers, was undertaken. From the effective SU(5) Lagrangian the allowed tree graphs and resulting decay modes for proton decay were derived. The value of $|\psi(0)|^2$ relevant for proton decay was estimated by various methods (electromagnetic and strong hyperfine splittings, electromagnetic form factors). The $SO(10) \rightarrow SO(6) \times SO(4) = SU(4) \times SU(2)$ symmetry breaking scheme was investigated and the renormalization group was used to study the mass scales of the various breakdown of symmetry. Careful study of the rare $K^0 \rightarrow e^+ \mu^-$ decay in the scheme was used to establish lower bounds on the mass of the vector boson whose mass is generated by the $SU(4) \rightarrow SU(3) \times U(1)_{B-L}$ breakdown. How to express α_e^{-1} in terms of the inverse coupling constants of the various subgroups was studied in this scheme and in more generality, i.e., how to express the charge operator in terms of the subgroup operators. Study of the mass scales of the U(1) splittings and their application to the magnetic monopole problem was attempted (it didn't help) and possible applications of cosmological measurements, e.g., helium abundance, to SO(10) parameters was investigated (again, it seems not to help, for helium abundance depends on weak interaction structure, not on any new structure at higher energy.) Recently I've been studying the various E_6 GUT's and their breakdown schemes, $E_6 \rightarrow SU^3(3), SO(10) \times SO(2), SU(6) \times SU(2)$.

Quark magnetic moments and E1 radiative transitions in charmonium

(Jonathan L. Rosner, with Gabriel Karl and Sydney Meshkov).

In the long-wavelength limit, $\Gamma(\psi' \rightarrow \gamma + \chi_J) = \text{const.} \times (2J+1)p_\gamma^3$

$\Gamma(\chi_J \rightarrow \gamma + \psi) = \text{const.} \times p_\gamma^3$. The corrections to these expressions of order p_γ/m_c (m_c = mass of charmed quark) were calculated. These corrections were found to be proportional to $\langle \vec{L} \cdot \vec{S} \rangle_{\chi_J} \kappa$, where $\langle \vec{L} \cdot \vec{S} \rangle_{\chi_J} = 1, -1, -2$ for $J = 2, 1, 0$, and κ is the anomalous magnetic moment of the quark. Angular distributions of photons in the decays $\psi' \rightarrow \gamma\chi$ and $\chi \rightarrow \gamma\psi$ also were predicted; small but probably measurable deviations from the pure E1 limit were found.⁵

Lectures on quark models (Jonathan L. Rosner).

A set of five lectures was delivered at a NATO Advanced Studies Institute on Techniques and Concepts of High Energy Physics, July 2-13, 1980, St. Croix, V.I., organized by T. Ferbel. These covered developments in the physics of both light and heavy quarks. A set of lecture notes was prepared, an abridged version of which will be published in the proceedings of the summer school.⁶

Inverse scattering and the Υ family (Jonathan L. Rosner, with C. Quigg).

A quarkonium potential was constructed with the help of masses and leptonic widths of the $T(1s-4S)$ levels using the inverse scattering formalism.⁷ This potential was found to agree at all interquark separations beyond 0.06f with one constructed earlier from ψ and ψ' , providing further evidence beyond that found earlier⁸ for flavor independence of the quark-antiquark interaction. Comparison with other a priori potentials suggested that tests for a short-range Coulomb interaction (as predicted by QCD) will have to rely primarily on more

precise values for $\Gamma(T \rightarrow e^+e^-)$, on measurement of the 2S-2P spacing (predicted to be about 120 MeV for a short-range Coulomb-like interaction or in the inverse scattering formalism but about 150 MeV for an effective power-law potential), and on the discovery of heavier quarks. This work was presented at parallel session C7, 20th Int. Conf. on High Energy Physics, Madison, Wisc., July 17-23, 1980.

Magnetic moments of quarks in baryons and mesons (Jonathan L. Rosner).

Recent measurements of hyperon magnetic moments and of magnetic dipole transitions in meson radiative decays were discussed in the context of the nonrelativistic quark model. It was shown that this model works surprisingly well. Various "fine-tuning" effects were discussed, including configuration mixing in the baryons, quark anomalous moments, and SU(3) breaking. This work was presented at parallel session C8, 20th Int. Conf. on High Energy Physics, Madison, Wisconsin, July 17-23, 1980.⁹

References

1. Jonathan F. Schonfeld, Waikwok Kwong, Jonathan L. Rosner, C. Quigg, and H. B. Thacker, Fermilab-Pub-79/77 THY, November, 1979, to be published in Ann. Phys. (N.Y.) 127 (1980).
2. Waikwok Kwong, Jonathan L. Rosner, Jonathan F. Schonfeld, C. Quigg and H. B. Thacker, Fermilab-Pub-80/14-THY, January, 1980, to be published in Am. J. Phys., 1980.
3. Jonathan L. Rosner, in Particles and Fields-1980 (APS/DPF Montreal), edited by B. Margolis and D. G. Stairs, AIP Conf. Proc. No. 59, Particles and Fields subseries No. 19, Amer. Inst. of Phys., New York, 1980, p. 325.

4. Peter Moxhay and Jonathan L. Rosner, J. Math. Phys. 21, 1688 (1980).
5. Gabriel Karl, Sydney Meshkov, and Jonathan L. Rosner, Phys. Rev. Lett. 45, 215 (1980).
6. Jonathan L. Rosner, "Quark Models", Univ. of Minn. report, 1980, abridged version to be published in Proc. of the NATO Advanced Study Institute on Techniques and Concepts in High Energy Physics, July 2-13, 1980, St. Croix, V.I., ed. T. Ferbel, Plenum Press, New York, 1981.
7. C. Quigg and Jonathan L. Rosner, "Inverse scattering and the T family", Univ. of Minnesota report, 1980, to appear in Proc. 20th Int. Conf. on High Energy Physics, Madison, Wisc., July 17-23, 1980, parallel session C7 (Amer. Inst. of Physics, New York, 1980).
8. C. Quigg, H. B. Thacker, and Jonathan L. Rosner, Phys. Rev. D21, 234 (1980).
9. Jonathan L. Rosner, "Magnetic moments of quarks in baryons and mesons", Univ. of Minnesota report, 1980, to appear in Proc. 20th Int. Conf. on High Energy Physics, Madison, Wisc., July 17-23, 1980, parallel session C8 (American Inst. of Phys., New York, 1980).

Ta-Pei Cheng

My recent research concerns several theoretical and phenomenological issues related to our attempts to construct grand unified gauge theories of strong, weak and electromagnetic interactions. Ling-Fong Li (of Carnegie-Mellon University) and I have proposed a possible resolution of the "gauge symmetry hierarchy problem". Basically we raised the possibility that the problem may reside with ordinary perturbative methods in dealing with such different mass scales. We suggest that "improved perturbation method" of renormalization group equations should be helpful in seeing why symmetry hierarchy can develop for some gauge groups. The result is being published in the Proceedings of the 1980 Quanzhou Conference on Theoretical Particle Physics. Other problems we have investigated this past year include: (1) the "kinship hypothesis" and its connection to lepton-quark mass relations; (2) neutrino masses; (3) lepton number nonconserving reactions. A paper with a catalog of all simple $SU(2) \times U(1)$ models with massive neutrinos has been written up. We are in the process of completing another project on $\mu \rightarrow e\gamma$.

I have also given a series of twelve seminars on "Theoretical Aspects of Proton Decay", with an audience of theorists and experimentalists.

3. Experimental High Energy Physics

(Professors Hans Courant, Kenneth Heller, Marvin Marshak, Earl Peterson and Keith Ruddick; Research Associate Michael Shupe; Research Assistants J. Bartelt, G. Blazey, B. Collick, S. Heilig, S. Heppelman, T. Joyce, Z. Malloi, B. Mossberg, J. Povlis and D. Wahl; Undergraduate Research Assistants T. Copie, D. Feyma, M. Hirsch, H. Hogenkamp, B. Nease, J. Osen; Miners H. Bristol, W. Heikkinen)

The experimental high energy group at Minnesota has historically concentrated on strong interaction physics. In the past year, this emphasis has continued, but the scope of the program has been broadened to include several crucial tests of the currently most important theories of elementary particle interactions. The program now includes both a range of particle energies and several different types of experiments. The Fermilab (both the charged-neutral spectrometer in the M1 line and the hyperon experiments in the M2 line) and Brookhaven efforts are conventional accelerator experiments, which have a high probability of yielding significant results. The Soudan baryon decay experiment is a speculative effort, which nonetheless has a reasonable prospect to make a truly significant contribution to particle physics and astrophysics. The now-concluded Argonne ZGS program has yielded considerable information about the importance of spin in particle interactions; these spin-dependent experiments are likely to be continued at higher energies in the near future.

Table 1: Summary of Experiments Discussed in this Report

Argonne E-365	Large Angle Elastic Scattering	Published
Argonne E-407	Measurement of Depolarization in the Process $p + p \rightarrow p + X$	Published
Argonne E-418	Asymmetries in Large Angle Nucleon Nucleon Scattering	Published
Argonne E-442	Deuteron Spin Alignment in High Energy Elastic Proton-Deuteron Backscattering	Published
Argonne E-443	Measurement of dp Elastic Scattering at Small Momentum Transfers with Beams of Aligned Deuterons at Momenta Equivalent to 0.6-1.0 GeV Protons	Published
Brookhaven E-717	Polarized Lambda Production at 24 GeV/c	Published
Brookhaven E-744	Production and Polarization	Analysis
Brookhaven E-755	Study of Exclusive Processes at Large Transverse Momenta	Construction
Fermilab E-272	Electromagnetic Couplings of Vector Mesons	Published and Analysis
Fermilab E-361	Lambda Beta Decay	Analysis
Fermilab E-441	400 GeV Lambda Production from H	Published
Fermilab E-495	400 GeV Ξ^0 Production from Be	Published
Fermilab E-555	Lambda Polarization at High P_T	Construction
Fermilab E-619	Σ^0/Λ^0 Transition Moment by Primakoff Effect	Construction
Fermilab E-620	Charged Hyperon Magnetic Moments	Analysis
Fermilab E-629	Direct Photon Production	Construction
Soudan E-1	Dense Detector for Proton Decay	Tuning

a. Electromagnetic Couplings of Vector Mesons-Fermilab E272

The qualitative successes of the quark model have been one of the most important themes in particle physics during the last decade. However, when this model has been used to make quantitative predictions, the results have not been so encouraging. In particular, one outstanding problem area has been the radiative decay of vector to pseudoscalar mesons. These decays can be viewed as M1 transitions from a 3S_1 to a 1S_0 state of the quark-antiquark system. A straight-forward calculation yields a ratio of about 9 between the widths $\Gamma_{\omega\pi\gamma}$ and $\Gamma_{\rho\pi\gamma}$, compared to the previously determined experimental ratio of 25 ± 7 . Fermilab E-272 was designed to make an improved measurement of the width $\Gamma_{\rho\pi\gamma}$ in order to make a definitive statement about this discrepancy. The experiment was also intended to study radiative decays of the $K^*(890)$, in order to investigate the possible flavor-dependence of quark magnetic moments, and to obtain high statistics data on the coherent production of the several A, B and other high mass mesons.

Because of backgrounds, the electromagnetic couplings of vector mesons are best measured through the coherent Primakoff process, in which an incident meson scatters from a virtual photon of the nuclear Coulomb field to produce an $I = 1, J = 1$ meson (such as ρ , K^* , B). For example, background from the common decay $\rho^\pm \rightarrow \pi^\pm \pi^0 \rightarrow \pi^\pm \gamma \gamma$ prevents detection of the rare $\rho^\pm \rightarrow \pi^\pm \gamma$ mode. $\Gamma_{\rho\pi\gamma}$ is then best measured by the Primakoff process $\pi + Z \rightarrow \rho + Z$. This experiment requires a high resolution charged neutral spectrometer and the highest possible energy beam. As the incident energy is increased, the Coulomb cross-section is enhanced relative to the strong-interaction background, thus significantly

reducing systematic errors. Such a spectrometer using drift chambers and a liquid argon neutrals detector was built by the Minnesota-Rochester-Fermilab collaboration for E-272. Data collection runs were made in 1978 and 1979-80. The earlier set of data has now been completely analyzed and the results published [Phys. Rev. Lett. 44, 706 (1980)]. Analysis of the latter data set, which emphasized kaon-induced triggers is now in progress.

E-272 has determined the width $\Gamma_{\rho\pi\gamma}$ to be 67 ± 7 keV, in contrast to a previous lower energy Primakoff measurement of 35 ± 10 keV. Figure 3.1 shows a plan view schematic of the experimental apparatus and a ρ mass plot from some reconstructed events. Figure 3.2 shows the cross-sections obtained for various nuclear targets along with fits to the data considering both Coulomb and strong interaction effects. The absolute normalization for the cross-section data is obtained by comparison to kaon decays in the spectrometer. The systematic accuracy is much improved over the earlier result, due to the almost order of magnitude increase in the incident momentum. In the case of the lead target, the coherent strong interaction contribution to the observed cross-section at small t was less than 5 percent. As a result, the Fermilab data do not contain as strong assumptions in the coupling constant calculation as the earlier Brookhaven AGS result.

The 1978 data have also been analyzed to determine $\Gamma_{K^*K\gamma}$ through the Primakoff processes $K^- + A \rightarrow K^- \pi^0 + A$ and $K^- + A \rightarrow K_S^0 \pi^- + A$. The width has been determined as 62 ± 14 keV [Phys. Lett. B, to be published]. This is the first reported measurement of the radiative width of the $K^*(890)^-$. The ratio of the radiative widths for the K^*0 [W.

Figure 1--Schematic plan of the E-272 apparatus (upper figure). $\pi^-\pi^0$ mass plot from a Pb target.

Figure 2--Cross-sections measured in E-272 from various nuclear targets.

Carithers, *et al.*, Phys. Rev. Lett. 35, 349 (1975)] to this measurement of the K^{*-} is 1.2 ± 0.6 , which is inconsistent with unbroken SU(3) symmetry, but in agreement with quark models in which different flavor quarks possess different magnetic moments. Figure 3.3 shows $K^{*}(890)$ mass peaks and decay angles. Measurements with much improved statistical errors on this process should be available in about six months, after completion of the analysis of the 1979-1980 data run.

Progress on the experimental study of the electromagnetic couplings of vector mesons has been advanced considerably by many discussions on this subject with members of the Minnesota theoretical group. As described in last year's report, D. Geffen and W. Wilson have developed a model which explains the E-272 results in terms of an anomalous contribution to the effective quark magnetic moment as a result of the magnetic coupling of the photon to three or more gluons [Phys. Rev. Lett. 44, 370 (1980)].

E-272 has also produced data on coherent production of high mass mesons and π -e scattering, which can be used to study the electromagnetic form factor of the pion. Figure 3.4 shows a typical three-charged-pion mass distribution, while Figure 3.5 indicates preliminary cross-sections for A_1 production. These data are currently being compared with the predictions of the Chou-Yang model for coherent production. A combination of the 1978 and 1979-80 data will provide sufficient events to permit a partial wave spin-parity analysis, which is planned for the coming year.

E-272 has been performed by a Minnesota-Rochester-Fermilab collaboration.

Figure 3--(a) $K\pi$ mass peaks from E-272 for the reaction $\pi + A \rightarrow K^* + A$.
(b) Gottfried-Jackson angle distributions for the same process.

FIG. 5

b. Direct Photon Production at Large Transverse Momentum-Fermilab

E-629

In 1980, our significant experience with the charged-neutral spectrometer has permitted us to develop a new experiment, E-629, which tests quantum chromodynamics by measuring the rate for direct photon production at large transverse momentum. There are two reasons for pursuing this measurement. First, QCD permits a direct calculation of the cross-section for direct photon production at large transverse momentum in strong interactions. An experimental measurement of this quantity thus is a test of QCD, which can result in refinement and validation of this important theoretical approach. The importance of this test is considerably increased by a simultaneous measure of π^0 production, since a ratio of direct photon to neutral pion rates eliminates some structure function dependence, which could complicate the QCD comparison. Second, if QCD is correct, then large transverse momentum direct photons should be correlated with quark and gluon jets on the away side. The type of jet expected on the away side depends on the type of particles participating in the interaction. In particular, for proton beams, quark jets are expected to dominate, while for incident pions, a majority of gluon jets should be observed. If this conjecture is correct, direct photons could provide a unique trigger for exploration of quark and gluon jet production processes.

The main aim of E-629 is to demonstrate the ability of our apparatus to distinguish direct photons from the large background of photons produced by neutral pion and other decays. The experiment will use the proportional chambers and the liquid argon neutrals detector from E-272. Validation of our technique for isolating direct photons is a necessary step towards a

Tevatron experiment, in which both the direct photons and the away side jets will be observed.

E-629, which is described in more detail in the Proposal for 1981, will be done by a Minnesota-Rochester-Fermilab-Michigan State collaboration.

c. Hyperon Experiments-Fermilab E-361, 441, 495, 555, 619, 620; Brookhaven E-717, 740

The Fermilab hyperon experiments have been the work of a Minnesota-Michigan-Rutgers-Wisconsin collaboration. E-361 (Lambda-beta decay) was completed at the end of October, 1979. Preliminary analysis of the data indicates the presence of more than 50000 lambda-beta decay events with an average Λ polarization of 10 percent. The synchrotron radiation detector devised for this experiment to identify electrons worked as expected. Data from this detector together with information from the lead glass array was used to achieve a hadron rejection from the electron sample of better than one part in 10000. This rejection ratio will lead to a $\Lambda \rightarrow p\pi^-$ contamination of the $\Lambda \rightarrow pe^- \bar{\nu}$ sample of order 10 percent. It is expected that further study of the data will yield some improvement in this rejection factor.

Fermilab E-620, the charged hyperon magnetic moment experiment, was completed at the end of January, 1980. A preliminary analysis of the data indicates a yield of greater than 1.2×10^5 Σ^+ , 1.5×10^5 Ξ^- and 5.0×10^5 Σ^- events. This analysis has also shown that the Ξ^- and Σ^+ particles produced inclusively by protons are polarized. The Ξ^- polarization is the same as the Λ^0 and the Ξ^0 polarization in sign, magnitude and kinematic

behavior. The Σ^+ polarization is also similar in magnitude and kinematic behavior but opposite in sign. The sign change of the Σ^+ polarization is consistent with the predictions of a simple quark model. The observation of significant polarization in the inclusive production of these hyperons insures that the data from this experiment will yield important results on these hyperon magnetic moments. Because of the small asymmetry parameter for the decay $\Sigma^- \rightarrow n\pi^-$, the preliminary analysis of a small fraction of the data has failed to determine whether these hyperons are also polarized.

During 1980, final analysis was completed on Fermilab experiments E-441 and E-495. E-441 studied the production of neutral lambda hyperons from a hydrogen target by 400 GeV protons at many different production angles. The major conclusions from this experiment were that polarization at high energies does not depend on the target, that is $P_{\Lambda}(\text{Be tgt}) = P_{\Lambda}(\text{H}_2 \text{tgt})$. The Λ^0 polarization does, however, depend on x and p_T . This dependence can be summarized as follows: For transverse momenta below 0.7 GeV/c, the polarization is a function of p_T alone; for transverse momenta above 0.8 GeV/c, the polarization is a function of x alone. The experiment also determined the inclusive differential cross-section for the process $p + p \rightarrow \Lambda^0 + X$, as a function of x and p_T .

Fermilab E-495 studied the production of Ξ^0 from a Be target by 400 GeV protons at several production angles. The major results from this experiment were that the Ξ^0 polarization was similar to that of the lambda, the magnetic moment of the Ξ^0 is -1.236 ± 0.014 nm and the asymmetry parameter, α_{Ξ^0} , for the decay $\Xi^0 \rightarrow \Lambda\pi$ is -0.415 ± 0.011 . These data for both μ_{Ξ^0} and α_{Ξ^0} represent the best measurements made of these quantities and are the currently accepted values. This experiment

also measured the inclusive differential cross-section, as a function of x and p_T for the reaction $p + Be \rightarrow \Xi^0 + X$.

Results from E-441 and E-495, along with preliminary results from E-620, were presented at the International Conference on High Energy Physics, Madison, 1980.

Fermilab E-555 is an approved experiment to measure inclusive lambda polarization at the largest possible transverse momentum. The Σ^0 to Λ^0 transition moment will be measured in approved Fermilab experiment 619 through analysis of the Primakoff effect. Both of these experiments are under construction and awaiting running time in the M2 beam.

The Brookhaven hyperon experiments are the results of a collaboration with Brookhaven, Michigan and Massachusetts. Brookhaven E-744, which studied Σ^0 production and polarization, completed data collection in August, 1980. Preliminary analysis indicates a yield of more than 50000 Σ^0 's from the reaction $p + Be \rightarrow \Sigma^0 + X$ at 28 GeV at a laboratory production angle of 4° . The data will be analyzed to determine the Σ^0 / Λ^0 production ratio together with P_{Σ^0} . This information will permit a determination of the direct lambda polarization from data measured in previous experiments. Simple quark models predict a polarization for the Σ^0 which has the opposite sign to Λ^0 polarization.

The analysis of Brookhaven E-717 was completed in 1980 and the results published [Physics Letters 86B, 386 (1979)]. That experiment measured the polarization of Λ^0 's produced by 28 GeV protons from H, D and Be targets. No difference was seen between hydrogen and deuterium indicating that neutron and proton targets are equally effective and have the same

sign for production of polarized lambdas. The Be target yielded a slightly lower Λ^0 polarization at these energies, which may be due to rescattering effects.

d. Spin Effects in Strong Interactions-Argonne E-407,418,442,443

The current year marked the close of Minnesota's long history of experiments at the Argonne ZGS. Results from four experiments were published during the year; two used the high energy proton beam, while the others took data on intermediate energy effects using the first polarized deuteron beam ever extracted at the Argonne ZGS. Argonne E-407 was the first measurement of the depolarization parameter in high energy pp inelastic scattering. The data, shown in Fig. 3.6, were analyzed in terms of the naturality of the exchange for production of baryon resonances in pp collisions. These data [Phys. Rev. Lett. 44, 1373 (1980)] yield the first direct evidence for pion exchange in $N^*(1232)$ production. The data also show that most of the higher resonances are produced through natural parity exchange.

The analyzing power for 2, 3 and 6 GeV/c np elastic scattering at large angles was first measured in E-418. This experiment used the polarized proton beam and a liquid deuterium target. The outgoing proton was detected in proportional chambers. The outgoing neutron was converted in a brass plate and the subsequent cascade also detected in proportional chambers. The data generally show a smooth variation between earlier forward and backward np polarization data. Unlike the pp polarization, the np data cross the axis and go negative. They also do not show the dip and peak structure observed in the pp data.

Figure 6--The depolarization parameter D for $p + p \rightarrow p + X$ as measured in experiment E-407. The curve shows a Regge prediction by Hidaka assuming π dominance for $N^*(1232)$ production.

Figure 7--Tensor observables measured at $T_d = 2$ eV in the reaction $d + p \rightarrow d + p$ in E-442 and 443. The curves represent multiple scattering model fit to the data.

Argonne E-442 and 443 measured the tensor and vector analyzing power for forward and backward dp scattering. The vector polarization of the deuteron beam was oriented normal to the incident direction both transverse to and in the scattering plane. Some of the data from these experiments is shown in Figure 3.7. These measurements were made in the context of an earlier program to determine the NN scattering amplitudes and the details of multiple scattering models at kinetic energies near 1 GeV. As can be seen from the figure, these data, when combined with earlier results on p-He scattering, yield a consistent set of NN amplitudes. This agreement between experiment and theory yields considerable confidence that multiple scattering models provide good representations of the physical mechanisms in the intermediate energy region.

e. A Dense Detector for Proton Decay-Soudan E-1

At the time of writing of this report (August 1980), assembly of the Soudan proton decay detector in the mine is mostly complete, with tuning and initial data collection still scheduled for Fall 1980. As described in the report "A Dense Detector for Proton Decay", the Soudan E-1 detector has a mass of about 30 tons of taconite concrete cast around 48 layers of 72 steel tubes per layer, with each tube 3 m in length. The 3456 tubes each have a 2.8 cm inner diameter and are set on a 4 cm lattice. In each succeeding layer, the tubes are rotated by 90 degrees, in order to provide two views of each event.

The chronological progress of this detector during the past year is as follows:

Nov.-Dec. The Minnesota-Harvard-Purdue-Wisconsin proton decay

collaboration, which proposed to use a water Cherenkov detector, terminates. The Minnesota group decides to build a Phase I dense detector in the Soudan Mine. A report describing the project is prepared. DOE approves the effort as instrumentation research.

Jan.-Feb. Engineering and prototype studies are done on the mine, the proportional tubes, the taconite concrete and the electronics for the experiment. A 5 year lease with a renewal option is negotiated with the State of Minnesota for two stopes in the Soudan Mine. The decision is made to use the 23rd level for the Soudan E-1 detector, while reserving the 25th level for a future detector.

Mar.-Apr. The Regents of the University of Minnesota approve the lease and designate the area as the Soudan Underground Research Site. Production of taconite slabs with tubes begins. Usable slabs are produced and proportional tubes are tested and plateaued. Work begins on preparing the 23rd level for the first detector by installing a foundation slab, electricity, a bulkhead to keep out cold air and a floor for the instrumentation.

May-June The method for construction of the taconite concrete slabs is perfected and slab production continues. Techniques for cleaning the tubes and stringing the 50 micron signal wires are developed. First slabs are installed in the mine and gas is connected. Amplifiers and time-over-threshold modules are prototyped. Four complete slabs are connected

to a computer and used to detect cosmic rays.

July-Aug. During this period, the construction of slabs is completed, 90 percent of the tubes are cleaned and strung and 70 percent of the detector is installed in the mine and connected to gas. Sample tubes are tested in the mine environment, yielding information about counting rates. Contracts are let for production of circuit boards and assembly of amplifier and time-over-threshold electronics. Planning for Phase II detector begins.

In general, work on Soudan E-1 has proceeded as outlined in the report "A Dense Detector for Baryon Decay". The only major surprise has been a singles rate of 10 Hz for tubes lying on the outer boundary of the detector. For tubes within the detector, the counting rate is attenuated by the detector mass at a rate consistent with that expected for gammas of energy about 1 MeV. This background rate, which does not affect the operation of the detector, may result from radon and radon daughters in the mine stope.

Further plans for the Soudan Underground Research Site are discussed in the Proposal for 1981.

f. Exclusive Processes at Large Transverse Momenta-Brookhaven E-755, Argonne E-365

For many purposes, the simple quark model, which we tested in Fermilab E-272, has been enhanced by a more complete theory of the strong interactions, known as quantum chromodynamics. This QCD model has also had many qualitative successes; it must now satisfy detailed, quantitative

tests in order to determine whether it is really a reliable picture of the strong interactions. Because of theoretical difficulties, significant tests are currently limited to the high energy, large transverse momentum domain. Brookhaven AGS E-755 is an experiment designed to measure the cross-sections for a number of exclusive reactions at large momentum transfers over the range 8-18 GeV incident momentum. This energy region is high enough to reach the largest momentum transfers presently measurable.

The reactions to be measured include

$$\pi^- p \rightarrow p\pi^-; p\rho^-; pA_2^-; \pi^+\Delta^-, K^+\Sigma^-, K^+Y^{*-}$$

As shown in Fig. 3.8, these reactions share a limited number of QCD gluon diagrams. This experiment should reveal, for example, whether the cross-sections at large momentum transfer for elastic scattering and rho production are the same. Rho production cannot proceed via the Landshoff diagram, but otherwise, it should be similar to elastic scattering. This approved experiment, a Minnesota-Brookhaven collaboration, is described in more detail in the Proposal for 1981.

Elastic scattering at large transverse momenta was also the subject of an earlier experiment by a Minnesota-Argonne-Columbia collaboration at the Argonne ZGS. In 1980, an archival article on this experiment, E-365, was published [Phys. Rev. D21, 2445 (1980)], which completed the unpolarized beam Argonne program.

Publications

Publications Previously Reported

- COO-1764-354: "Mass Dependence of Schrodinger Wavefunctions," C. N. Leung and J. Rosner, J. Math. Phys. 19, 1435 (1979).
- COO-1764-360: "Quantum mechanics with applications to quarkonium", C. Quigg and J. Rosner, Phys. Reports 56C, 167-235 (1979).
- COO-1764-361: "Resource Letter NP-1: New Particles", J. Rosner, Am. J. Phys. 48, 90 (1980).
- COO-1764-362: "Constructive evidence for flavor independence of the quark-antiquark potential", C. Quigg, H. Thacker and J. Rosner, Phys. Rev. D21, 234 (1980).
- COO-1764-369: "Inverse problem for symmetric reflectionless potentials," title changed to: "On the Convergence of Reflectionless Approximations to Confining Potentials," J. Schonfeld, W. Kwong, J. Rosner, C. Quigg and H. Thacker, to be published in Ann. Phys. (N. Y.) 127 (1980).

New

- COO-1764-377: PROGRESS REPORT
- COO-1764-378: "Magnetic Properties of the Low-Lying Hadrons", W. Wilson, and D. Geffen, submitted to Phys. Rev. Lett.
- COO-1764-379: "Neutrino masses, mixings and Oscillations in $SU(2) \times U(1)$ Models of Electroweak Interactions", T. P. Cheng. Submitted for publication.
- COO-1764-380: "Gauge Hierarchy Problem", T. P. Cheng, submitted for publication.

- COO-1764-381: "Degeneracy in one-dimensional quantum mechanics", Waikwok Kwong, J. Rosner, J. Schonfeld, C. Quigg and H. B. Thacker, Fermilab-Pub-80/14-THY, Jan. 1980, to be publ. in Am. Journ. Phys. 1980.
- COO-1764-382: "Heavy quarks and new particles", J. Rosner in Particles and Fields - 1980 (APS/DPF Montreal) ed. by B. Margolis and D. G. Stairs, AIP Conf. Proc. No. 59, Part. & Fields, subseries No. 19, AIP, N.Y. (1980), p. 325.
- COO-1764-383: "Semiclassical results on normalization of bound state wave functions", P. Moxhay, J. Rosner, Journ. Math. Phys. 21, 1688 (1980).
- COO-1764-384: "Quark Magnetic moments and E1 radiative transitions in charmonium", Gabriel Karl, S. Meshkov and J. Rosner, Phys. Rev. Lett. 45, 215 (1980).
- COO-1764-385: "Quark Models", J. Rosner, U of M report 1980, to be publ. in Proc. of the NATO Advanced Study Inst. on Techniques and Concepts in High Energy Physics, July 2-13, 1980, St. Croix, V. I., ed., T. Ferbel, Plenum Press, N.Y. 1981.
- COO-1764-386: "Inverse Scattering and the γ family", C. Quigg and J. Rosner U of M report 1980, to be published in Proc. of the 20th Intern. Conf. on High Energy Physics, Madison, Wisc., July 17-23, 1980, parallel session C7 (AIP, N.Y. (1980)).
- COO-1764-387: "Magnetic moments of quarks in baryons and mesons, J. Rosner, U of M report 1980, to be published in Proc. of the 20th Intern. Conf. on High Energy Physics, Madison, Wisconsin, July 17-23, 1980, parallel session C8 (AIP, N.Y. (1980)).
- COO-1764-388: "Depolarization Parameter in PP Inclusive Scattering at 6 GeV/c", Phys. Rev. Lett. 44, 1373 (1980) (with Courant, Kagan, et al.).

- COO-1764-389: "Measurement of Wide Angle Elastic Scattering of Pions and Protons Off Protons", Phys. Rev. D21, 2445 (1980) M. Marshak, et al.
- COO-1764-390: "Radiative Decay of the K^* (890)", submitted to Physics Letters, M. Marshak et al.
- COO-1764-391: "The Soudan Mine Experiment: A Dense Detector for Baryon Decay", Proc. of the 1st Workshop on Grand Unification, Durham, N. H., April 1980, M. Marshak, et al.
- COO-1764-392: "The Spin Dependence of the Strong Interaction: An Experimental Review", Proceedings of the 1980 Intern. Symposium on High Energy Physics with Polarized Beams and Polarized Targets, Lausanne, 1980, M. Marshak, et al.
- COO-1764-393: "Polarization in Large Angle Proton-Neutron Elastic Scattering", submitted to Phys. Rev. Lett., M. Marshak, et al.
- COO-1764-394: "A Dense Detector for Baryon Decay", unpublished, H. Courant, et al.
- COO-1764-395: "Relativistic wave equation and heavy quarkonium spectra", H. Suura, W. Wilson and B. L. Young, Phys. Rev. D21, 3204 (1980).
- COO-1764-396: "Relativistic wave equation and mass spectrum of gluonium", H. Suura, Phys. Rev. Lett. 44, 1319 (1980).
- COO-1764-397: "Polarization in Inclusive Lambda Production", in High Energy Physics with Polarized Beams and Polarized Targets; ed. by G. H. Thomas, AIP, N. Y. 1979, p. 542, K. Heller, et al.
- COO-1764-398: "The First Measurement of the Ξ^0 Magnetic Moment", Phys. Lett. 86B, 386, 1979, K. Heller, et al.
- COO-1764-399: "A Study of Inclusive Λ Polarization from Hydrogen and Other Targets at 28 GeV", Phys. Lett. 90B, 319, (1980), K. Heller, et al.

COO-1764-400: "Proton Polarization in Inclusive Production at 400 GeV/c",
K. Heller, et al.

COO-1764-401: "Precise Measurement of the Ξ^0 Hyperon Decay Parameter",
K. Heller, et al.

COO-1764-402: "Precise Measurement of the Ξ^0 Magnetic Moment", K. Heller, et al.

COO-1764-403: "Polarization of Ξ^0 's Produced by 400 GeV Protons", K. Heller, et al.

COO-1764-404: "The Polarization of Hyperons Produced by 400 GeV Protons",
in Proceedings of the 20th Intern. Conf. of High Energy
Physics, 1980, K. Heller, et al.