JAN 19 1993

Progress Report

Period: November 1, 1991 to October 31, 1992

U.S. Department of Energy Grant No. DE-FG05-91ER40622

EXPERIMENTAL STUDY OF HEAVY FLAVOR PHYSICS AND SSC RESEARCH AND DEVELOPMENT

at

The University of Mississippi

Principal Investigator

James J. Reidy, Professor of Physics

Co-Principal Investigators

Lucien M. Cremaldi, Assistant Professor of Physics

Donald J. Summers, Assistant Professor of Physics

Physics Department

University, MS 38677

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

Table of Contents

	Page
1. Introduction	1
2. Progress in the Experimental and High Energy Physics Program During the Period Noveber 1, 1991 to October 31, 1992	4
2.1 E691	4
2.2 E769	4
2.3 E791	5
2.3.1 UNIX/RISC Computing Farm	6
2.3.2 FY92 Physics Analysis Effort	10
3. New Facilities	19
4. SSC Research and Development Program	19
4.1 Introduction	19
4.2 CALOR89 Development and Simulation Studies	20
4.3 Scintillating Fiber/Plate Hadron Calorimeter Test Beam Analyses	21
4.4 Scintillating Fiber/Plate Hadron Calorimeter Studies for GEM	21
4.5 Infrastructure Development	23
5. Proposals Submitted.....	27
6. Summary	27
Appendix	29
1. Presentations	
2. E691	
3. E769	
4. E791	
5. SSC R&D	

1. Introduction

The University of Mississippi High Energy Physics (UMiss HEP) Group is partially supported by the U.S. Department of Energy grant DE-FG05-91ER40622 and the Texas National Research Laboratory Commission (TNRLC) project RCFY91 30 through the SAHEP consortium. It presently consists of three tenured or tenure-track faculty (Profs. Cremaldi, Reidy and Summers), one visiting faculty, (Prof. Weinstein) two postdoctoral associates (Drs. Krish Gounder and Brent Moore), one adjunct professor (Dr. Jeff Spalding, FNAL), one technician(temporary) and five students with assistance from the physics department's electronics technician, computer systems manager, machinist/instrument maker and administrative coordinator. One of the postdoctoral associates is funded by the TNRLC and the technician has a temporary appointment based on available funds. Two of the students, Mr. Ali Rafatian and Ms. Danying Yi, having passed their Ph.D. qualifying examinations and participated in the E791 run at FNAL, are presently analyzing data and writing dissertations. A third graduate student, Eric Aitala, should finish his M.S. in physics using E791 data next summer. Another student, David Brown, will finish his M.S. in the summer of 1993 as well. Three undergraduate students and three graduate students in computer science are making significant contributions to the E791 and SSC R&D programs. A B.S. computer science student, Kevin Hendrix, who was supported by us to assist Mr. Quinn in managing the DECstation "farm", plans to accept our offer (pending support) of the position as "farm" system manager since Mr. Quinn has left to pursue graduate studies in physics at the University of Chicago.

The HEP group is involved in three Fermilab experiments, E691, E769 and E791. This series of trend-setting heavy quark experiments has been the most productive at Fermilab in the past few years. E691/E769/E791 have set the standard for high rate data acquisition systems, innovative reconstruction of large data samples and analysis techniques. These experiments have been manned by international collaborations and have produced a number of quality Ph.D. students including two at UMiss. Drs. Summers and Cremaldi are senior collaboration members and guide many day-to-day decisions. The UMiss group has expended major resources toward the success of the experiments especially in the areas of Cerenkov detectors, DA systems and muon detection. Now the group has become a leader in the large analysis effort for E791. In support of these analysis efforts a 900 MIP "farm" was assembled at UMiss, commissioned in early summer and devoted almost exclusively to the E791 program since that time. Mr. Steve Bracker from E791 was retained as a short-duration consultant to assist in this development.

The group has an ongoing SSC R&D program dealing with (a) the adaptation and use of the HETC-based detector simulation code CALOR89, (b) the development of liquid scintillator technology for use in SSC detector calorimeters, (c) the hanging file calorimeter project (i at FNAL) and (d) the calorimetry program for GEM. In addition, Prof. Reidy serves as Principal Investigator for the SAHEP program funded by the Texas National Research Laboratory

Commission and is chair of the Executive Committee for SAHEP. This program (\$1M in each of FY91 and FY92) involves a consortium of sixteen universities and Oak Ridge National Laboratory and has as its purpose the enhancement of the infrastructure of these universities thereby enabling them to make significantly greater contributions to the SSC program. The major contributions of the UMiss HEP group in these SSC efforts lie mainly in the areas of detector development and simulation, simulation code development and fast data acquisition systems.

The UMiss HEP group has cooperative programs with Jackson State University (an HBCU) and Memphis State University. These programs allow students who are interested in high energy physics to work with our group and later transfer to the University of Mississippi as Ph.D. candidates in high energy physics. At Memphis State Dr. S. Jahan and one graduate student are working with us on radiation damage studies with scintillating fibers and plates. At Jackson State University Prof. H. Zhang works with us on SSC-related detector simulation projects.

An set of milestones has been developed by the HEP group for use in planning and evaluation. These are summarized in Table I.

Table I. Milestones

<u>EXPERIMENTS</u>	<u>SSC RESEARCH & DEVELOPMENT</u>
	1990
Analysis of FNAL E769; work on Cerenkov detector, muon detector and DA system for FNAL E791.	Join SDC as charter member; hired postdoc with DOE/HEP/SSC funds to work on CALOR89; join L-Star as co-ordinator for detector simulation.
	1991
E769; running on E791; mentor Jackson State U.(HBCU) faculty member on E791; hire K. Gounder as E791 postdoc with DOE/HEP funds; hire B. Quinn with TNRLC funds; muon wall upgrade.	Join GEM; hire postdoc and faculty member on TNRLC funds; PI on DOE/SSC funded Liquid Scint. Calorimeter R&D; participate in "hanging file" test beam prep-MOU with SSCL; begin restructuring of CALOR89 to install on Unix systems.
	1992
E769; finish E791 run-begin analysis with assembly of 900 MIP "farm" using OSU loan equipment and TNRLC funds.	Install CALOR89 on PDSF at SSCL-prepare users manual (SDC-92-246); "hanging file" run at FNAL; design & fabricate optics package for prototype

scin.fiber calorimeter - test beam run at BNL; analysis of "hanging file" and scin. fiber cal. test beam data; Dr. Reidy on sabbatical at SSCL as a lead editor for GEM Technical Design Report.

1993

E769; E791; hire K. Hendrix as farm manager and a research scientist as E791 physics analysis coordinator with DOE funds; award two Ph.D.s.; initiate participation in new experiment -collider D0 upgrade?; possibly fixed target?

Hire research scientist (TNRLC funds) to direct optics R&D for GEM hybrid hadron calorimeter; hire new faculty (TNRLC funds); complete work with industry on scin fibers and plates; begin design of hybrid cal with responsibility for optics R&D and fiber/plate testing.

1994

Continue E791 analysis; join collider experiment; L. Cremaldi on sabbatical;

Complete prototype design and fabrication of optics; use in test beam run; perform analysis of test beam data for timing studies; continue fiber/plate testing.

1995

Finish E791; work on DØ or other experiment; D. Summers on sabbatical.

Upgrade optics design and fabricate initial units; detector simulation studies; muon timing system development; test beam runs on prototype.

1996

Collider experiment; install analysis software on farm.

Test beam runs; evaluate; make modifications as needed; continue development of simulation pkg.

1997

Collider data analysis

Test initial scin modules from supplier, develop trigger software; continue simulation development; install scintillator modules in GEM detector at SSCL.

1998

Collider data analysis

Continue installation and software development; initiate installation of GEM analysis software at UMiss.

1999

GEM; new initiative?

Finish installation; take data; troubleshoot!

2. Progress in the Experimental High Energy Physics Program During the Period November 1, 1991, to October 31, 1992

The progress in the main areas of research of the UMiss HEP group will be described below. Only brief narratives will be given in most cases with more complete descriptions found in the articles contained in the Appendix. The research program encompasses primarily Fermilab experiments E691, E769, E791 and the SSC R&D program with the main emphases on E791 and the SSC R&D activities. Personnel in the UMiss HEP group authored or co-authored six refereed publications, have submitted five additional preprints for publication, authored six technical reports and gave twelve presentations at national or international meetings. Since E791 is in the analysis phase and the SSC R&D mostly in its formative stage this output is quite significant for a small group. A brief description of the activities in each area follows.

2.1 E691 (Photoproduction of Charm): University of California at Santa Barbara, Carleton University., University of Cincinnati, Centro Brasileiro de Pesquisas Fisicas(CBPF), University of Colorado, FNAL, University of Mississippi, NRC Canada, Princeton University, Universidade de Sao Paulo, University of Toronto, Yale University - Prof. Cremaldi.

E691 was a high statistics charm photoproduction experiment run at FNAL in the summer of 1985. About 25 journal articles, mostly in Physical Review Letters have been published. Most of the work on E691 physics has been completed, only one or two publications are pending. These are listed below and are included in the Appendix.

- 1- Study of the decays $D \rightarrow K\pi\pi\nu$ and $D \rightarrow K^*\pi\nu$, The Tagged Photon Spectrometer Collaboration, J.C. Anjos et al., Phys. Rev. D. Rapid Communications **45**, (1992) R1.
- 2- Experimental probes of final-state interactions in D^0 -meson decays, The Tagged Photon Spectrometer Collaboration, J.C. Anjos et al., Phys. Rev. D. Rapid Communications **46**, (1992) R2177.
- 3- Study of the decays $D^+ \rightarrow K3\pi$, The Tagged Photon Spectrometer Collaboration, J. C. Anjos et al., Physical Review D. **46**, (1992) 1941.
- 4- A Dalitz plot analysis of $K\pi\pi\pi$ Decays, The Tagged Photon Spectrometer Collaboration, J.C. Anjos et al., submitted to PRL.
- 5- Dalitz plot analysis of $\Lambda_c^+ \rightarrow p K\pi$ Decays, The Tagged Photon Spectrometer Collaboration, J.C. Anjos et al., in preparation.

2.2 E769 (Hadroproduction of Charm): Centro Brasileiro de Pesquisas Fisicas(CBPF), FNAL(Fermilab), University of Mississippi, Northeastern University, University of Toronto, Tufts University, University of Wisconsin-Madison, Yale University - Profs. Crema'di and Summers; A. Rafatian.

E769 is a high statistics charm hadroproduction experiment run at FNAL's Tagged Photon Lab during the fixed target running period of '87-88. Charm states were produced by the interactions of a mixed hadronic beam (π , K, p) of 250 GeV with targets of Be, Al, Cu, and W. Experiment 769 recorded about 400 million events on 10,000 nine-track tapes and reconstructed them on a farm of Advanced Computer Project (ACP) parallel processors and UNIX/RISC computers. The reconstruction required a full years worth of CPU time on these machines and was equivalent to about 500 VAX 11/780 years of computing. This was the first time UNIX/RISC computers had been used for such a large data set in HEP.

All of the data tapes have been reconstructed. A sub-sample of events has been extracted based on the reconstruction of multiple vertices as seen in our silicon micro-vertex detector. Our latest results will be presented at the annual meeting of DPF at Fermilab in November 1992. They are appended to this document and include Feynman-x and transverse momentum distributions for charm, as well as measurements of the atomic mass dependence of charm production and a study of beauty and charm production using semi-electronic decays. There are two Fermilab preprints of our results on x_F , p_t and A-dependence of charm production (FERMILAB-Pub-92/208-E, FERMILAB-Pub-92/279-E.) The first paper has already been accepted for publication by Physical Review Letters and the second one was submitted to the same publication in October.

Presently our major participation in the E769 analysis is through the work of Ali Rafatian. Mr. Rafatian is a graduate research assistant studying for his Ph.D. at the University of Mississippi. His dissertation research will be based on the E769 data. He has participated in the running of E791 and in the reconstruction and analysis efforts at Fermilab on E769 during the last three years. He is now at UMiss to finish his work. Over the past year Mr. Rafatian has concentrated his efforts on the study of the $D^{*+} \rightarrow D^0 \pi^+$ and $D^0 \rightarrow K^- \pi^+ \pi^- \pi^+$ decays. He has prepared a preliminary draft of a paper on the results of his study. Some of the material on E769 in the Appendix includes:

- 1- Using Multiple RISC CPUs in Parallel to Study Charm Quarks, C. Stoughton and D. J. Summers, Computers in Physics (1992) 371
- 2- Feynman-x and Transverse Momentum Dependence of D^{*+} and D^0 , D^0 -bar Production in 250 GeV π^- - Nucleon Interactions, G. A. Alves *et al.*, Physical Review Letters **69**(1992) 3147.
- 3- Atomic Mass Dependence of D^{*+} and D^0 , D^0 -bar Production in 250 GeV π^- - Nucleon Interactions, G. A. Alves *et al.*, (submitted to Phys. Rev. Lett.) FERMILAB-Pub-92/279-E.
- 4- Study of $D^{*+} \rightarrow D^0 (K^- \pi^+ \pi^- \pi^+) \pi^+$ Production, A. Rafatian, E769 Technical Note.

2.3 E791 (Hadroproduction of Charm): Centro Brasileiro de Pesquisas Fisicas(CBPF), FNAL, Illinois Institute of Technology, University of Mississippi, Princeton University, Ohio

State University, Tel-Aviv University, Tufts University, Universidade Federal do Rio de Janeiro(UFRJ), University of Wisconsin - Madison, Yale University - Profs. Cremaldi, Reidy and Summers; Dr. Gounder; Aitala, Rafatian, Rich and Yi.

E791 is a very high statistics charm hadroproduction experiment that has finished taking data at FNAL's Tagged Photon Laboratory. Charmed particles were produced through π -N interactions at $E_{\pi}=500$ GeV. An unprecedented 20 billion triggers were recorded on 25000 Exabyte tapes with a loose transverse energy trigger. We expect to fully reconstruct over 200K charm particles from this data. These data tapes are to be processed at UNIX/RISC computing farms set up at the University of Mississippi, Ohio State University, CBPF and Fermilab. Krish Gounder, Don Sumners, Lucien Cremaldi, intermittent visitor Steve Bracker and student Danying Yi are the principal UMiss researchers on this project.

2.3.1 UNIX/RISC COMPUTING FARM

Introduction

The UNIX/RISC computing farm at the University of Mississippi will be used in the reconstruction of approximately 1/3 of the E791 data set as well as for SSC detector simulations and test beam data analysis. The Dec farm consists of 40 DECstations Model 5000/200 with 25 MHz MIPS R3000 CPUs - about 900 MIPS of computing. The CPU's are connected via ETHERNET. The HEP events are distributed over the 40 nodes for parallel reconstruction by specially tailored distribution software. Many thousands of raw data tapes will be processed in the next few years on the UMiss farm. This reconstruction will generate about 3000 additional data summary tapes(DSTs) which must be distributed. A copy of each DST is sent to Fermilab from where they are distributed to our other collaborators. Presently the UMiss farm represents about 40% of the E791 computing power.

Breese Quinn, paid with SSC funds, managed this farm through the summer of 1992. (He has entered graduate school in physics at the University of Chicago). Kevin Hendrix, an undergraduate computer science student, assumed Breese's responsibilities in midsummer. Kevin is closely supervised by Dr. Gounder, who has provided great support and guidance. Dr. Gounder is in charge of the E791 farm software and supervises the day-to-day running of the facility. Mr. Hendrix graduates in December, 1992, and is willing to work fulltime as the farm system manager. However, the SSC funds for this position are exhausted. Funding this position is our topmost priority.

Farm Software

In January, just after the finish of the run at Fermilab, the E791 farms were not operational. Computing farms are relatively new structures and very little software exists for event distribution. Consequently, it was not surprising that event distribution software promised by Fermilab's computing division was not available. We invited Steve Bracker to UMiss to help with this

software development. Through an enormous effort by him and Krish Gounder at UMiss, an event distribution package for the Dec processors was developed and tested in a few weeks. This package allows HEP events to be shipped to each farm node and the reconstruction results returned to common disk area. A second process running on the disk collects output from the nodes when all events are finished and writes them to tape. Reconstruction of E791 data began in February, 1992, at UMiss. At the same time Ohio State University began running a package from Argonne National Lab on their farm. E791 had its first charm peaks in March, 1992. Gounder and Bracker are largely responsible for this quick startup and for providing a basis for the highly constructive competitiveness between groups at UMiss and Ohio State. Without Steve Bracker's talent and expertise on interprocessor communication this project could not have been accomplished in such a revolutionary short time. Mr. Bracker remains in close contact with our group and visits frequently. Many experimenters are now contacting Steve Bracker and Krish Gounder for the details of the UMiss event distribution software package. A representation of the UMiss farm distribution package is shown in Figure 1.

FIGURE 1. Flow chart summarizing the UMiss farm distribution package.

FY92 Farm Processing

A schematic of the farm configuration is shown in Figure 2. Through Dr. Gounder's excellent management and hard work, the UMiss farm has averaged processing *well over* 100 tapes per month. Much of the earlier running was used for testing purposes, exploring charm particle yields

FIGURE 2. Present UMiss farm configuration.

and to establish farm strategy. In Table II we summarize E791 data processing through September, 1992. Runs processed at UMiss have been distributed to the collaboration.

Table II. E791 Data Processing - FY92

Site	nodes/type	MIPS	tapes processed	start date
UMiss *	40 Dec	900	900	2/92
OSU	40 Dec	900	900	2/92
OSU	40 Dec	800	200	8/92
Fermilab	24 IBM	600	50	6/92
CBPF	ACPII	900	0	**

* shared with SSC work

** not available due to software problems

Our first charm signals were seen in Feb/Mar, on data processed by both the UMiss and OSU farms. These two universities have shown that large HEP computing can be managed quite easily at university based sites. There is no question that farms consisting of many thousands of MIPS can be university operated.

In addition to the UMiss and OSU farms, E791 has requested farm resources from Fermilab. The

FNAL computing division has donated some ACPII systems through CBPF-Rio to E791. Neither of these farm sites has contributed in FY92 to large scale processing of E791 data.

The UMiss group has invested a great deal, both in funds and personnel, on the farm. A large fraction of the funds have come from a TNRLC infrastructure grant that is now exhausted. No direct funds have been given for this by DOE even though at DOE's direction about 30% of the farm is on loan from OSU. We summarize the UMiss farm expenditures to date below and present a detailed list for the record in the Appendix. These numbers do not include personnel costs nor space contributions. These inclusions give a total investment in the farm of nearly \$400K.

Table III. UMiss Farm Expenditures- FY92

<u>YEAR</u>	<u>AMOUNT</u>	<u>SOURCE</u>
1991	\$ 117666	TNRLC
	82809	Loan from OSU
1992	<u>54569</u>	TNRLC
TOTAL	\$ 255044	

Tape Copying

Presently the UMiss farm can average 4 runs per month. Each run (40 input tapes) generates about 15 output tapes. These output tapes contain a copy of the raw data plus physics analysis from the farm processing, such as vertex, tracking, and particle identification information, plus momentum analysis. These data summary tapes, DSTs, are copied by a tape copy facility that we have set up at UMiss with our only available analysis workstation and 2 very precious EXABYTE tape drives. Kevin Hendrix, our present acting system manager, has written the tape copy software. We have no hardware backups to this critical copy facility and quite often the tape drives and workstation must be used for other functions, interfering with its tape copy function. Timely distribution is important for prompt physics analyses so we are requesting an additional workstation for this function.

Distribution of the UMiss Farm Software to Other HEP Sites

The UMiss farm shell is one of two parallel processing farm shells developed independently at university sites on E791. The shell has undergone two major revisions and is fully documented and easily transferable to other sites. It is unique with multiple output streams and sophisticated

online processing mechanisms. The tape-to-disk based system is very robust and may be more reliable than existing farm shells using tape-to-tape based systems. It is relatively trouble-free. A single client can crash with no effect on any other part of the system. Only the input tape drive must run continuously to keep the farm operational.

The UMiss farm software has been distributed to a number of institutions across the country. It is presently in use entirely or in parts at DØ(Fermilab) and Ohio State University. We also have had enquiries by the Sky Survey Project (a research consortium of Princeton University, University of Chicago, California Institute of Technology and Fermilab), the University of California at Berkeley, and Centro Brasileiro de Pesquisas Fisicas - Rio de Janeiro regarding the possibility of adopting this farm shell.

2.3.2 FY92 PHYSICS ANALYSIS EFFORT

World's Largest Sample of Semileptonic Decays

UMiss played a major role in setting up and maintaining the muon systems at TPL. At the insistence of L.Cremaldi and D.Summers, UMiss added a second level of muon detection to the existing system at TPL, believing that it would enhance clean reconstruction of semileptonic charm decays. Graduate student Danying Yi was in charge of the muon system during the run. It should be noted that previous experiments at TPL have used only electrons for these studies.

The muon detector worked well with high efficiency. E791 is now able to reconstruct semileptonic decays in the muon channels, based on information derived from the new wall. This will be the largest sample of semileptonic charm decays ever studied by any HEP experiment. It is disappointing that DOE was not willing to support such a small but heavily endorsed project; rather, extremely limited university resources had to be committed to it - resources that were meant to expand our equipment base. As an example of the use of the muon system Figure 3 on the next page presents the results of a preliminary study with a detected muon in the final state. E791 expects over 5000 decays in this channel alone.

Muon Reconstruction Software

The muon reconstruction software has been developed by Danying Yi and Dr. Cremaldi at UMiss. This reconstruction software was installed into the E791 reconstruction package in March 1992.

Ms. Yi did the detector alignments and time calibrations in the spring of this year. This software package points spectrometer tracks into the TPL muon systems and returns a muon likelihood value based on signals detected in the muon counters. Further muon discrimination is derived from the calorimeter energy depositions. She is presently working to refine this package. Dr. Cremaldi and Ms. Yi have used this software to extract semi-muonic charm signals and $J/\Psi \rightarrow \mu\mu$ decays (see Figure 4). It is expected that about 2000 such events will be used for mass scale and calibration purposes. Ms. Yi is studying the efficiency of the muon identification code and is

working closely with Nick Witchey and Arun Tripathi from Ohio State University.

FIGURE 3. Semileptonic Decays $D^+ \rightarrow K^- \pi^+ \mu^+ \nu_\mu$ from about ten E791 runs.

FIGURE 4. Decays of $J/\Psi \rightarrow \mu\mu$ from about two E791 runs.

Cerenkov Reconstruction Software

Two large multicelled Cerenkov counters in the Tagged Photon Spectrometer provide particle identification in separating pions, kaons and protons. Some muons and electrons are also tagged by this system. As E791 reconstruction and filter ideas mature, it is evident that these counters play a major role in the charm physics that E791 will produce. Dr. Cremaldi is responsible for the Cerenkov particle identification software. The bulk of this work has been done with Tel Aviv University. The software was made fully operational in the fall of 1991. The $\pi/K/p$ particle id is working with good efficiency. The run by run calibrations for the detectors is being done by Tel Aviv University in close collaboration with UMiss.

Production studies

Eric Aitala has been working with E791 for over two years. He is writing an M.S. thesis on the A-dependence of charm production. This research is being directed by Dr. Summers. This past summer he has worked to isolate a clean sample of $D^0 \rightarrow K\pi$ decays for his studies. Dr. Gounder

has provided invaluable assistance to him. Eric has also written a compact utility routine for fitting mass plots that many in the UMiss group have adopted.

Charm Meson Results from UMiss

The UMiss group has been involved from the very beginning with extracting charm meson decay signals from it's farm runs. In the early phases of the reconstruction these charm signals are used as benchmarks to reference how well the software development is progressing. Krish Gounder has written many monitoring and stripping jobs that extract the familiar charm signals for $D \rightarrow K\pi$, $D \rightarrow K2\pi$ and $D \rightarrow K3\pi$ decays. Based on many of these programs the UMiss group has the opportunity to expand into a great many physics areas using the E791 data set. Indeed, some of the first clean $D \rightarrow KK\pi$ and $\Lambda_c \rightarrow pK\pi$ signals (see Figure 5 on page 13) based on Dr. Gounder's software were shown by the UMiss group to the collaboration. We are presently working to improve the yields.

Charm Filter Studies from UMiss

The E791 data set is so large that it is prohibitive to fully reconstruct every event. Krish Gounder has been developing off-line charm filters to be implemented on the E791 farms. He has made a major impact on the conventional code and has shown himself to be one of the very few people in E791 willing to exploit all the physics potential of the data set and point out major weaknesses of the software.

This past summer he has shown that the E791 vertexing algorithm was not nearly as efficient as it could be and encouraged improvements. He implemented a filter on the UMiss farm which tags Λ and K_S . The output from this "vee filter" has been used to reconstruct charm decays involving Λ and K_S . Using the UMiss filter he obtained the first $\Sigma^- \rightarrow \Lambda\pi^-$ on the experiment. A sample of baryon information is presented in Figure 6 on page 14. These studies have provided the groundwork for charm baryon studies at UMiss. Further studies have involved tagging interesting physics with just particle identification, in order to optimize our efficiency for gathering good physics from the 20 billion event data sample.

Krish also has been working on single prong decays, which involves a single track detected in the silicon vertex detector combined with a downstream reconstruction vee. These decays are more difficult to detect because a multiprong secondary vertex cannot be found in the silicon detector. Some examples are the two body decays, $D^+ \rightarrow K_S\pi$, $D_S^+ \rightarrow K_S K$ and $\Lambda_c^+ \rightarrow K_S p, \Lambda\pi$. In Figure 7 a representation is shown of a signal for $D^+ \rightarrow K_S\pi$ in this channel from two runs. These decays look very promising and will be the easiest to use in Λ_c polarization studies.

(a) $D^0 \rightarrow K\pi$

(β) $D^+ \rightarrow K\pi\pi$

(c) $D_s \rightarrow KK\pi$

(d) $\Lambda_c \rightarrow pK\pi$

FIGURE 5. Charm signals extracted from UMiss farm runs.

(a) $\Lambda \rightarrow p\pi$

(b) $\Xi^- \rightarrow \Lambda\pi$

(c) $\Xi(1530) \rightarrow \Lambda\pi$

(d) $\Omega^- \rightarrow \Lambda K$

FIGURE 6. Baryon signals extracted from UMiss farm runs.

FIGURE 7. Single prong decays of $D^+ \rightarrow K_s \pi^+$ and $D_s^+ \rightarrow K_s K^+$ from two UMiss farm runs.

Charm Baryon Studies at UMiss

The E791 data set is rich in Λ and K_s decays ('vees'). These decays can occur both inside the silicon vertex detector and just downstream of it in a 2-3 meter decay region. The standard silicon vertexing and vee-finding software locates these secondary decays with good efficiency. Since a large fraction of charmed baryon decays will contain Λ and K_s components, the reconstruction vees are a must for doing good charmed baryon physics.

The first large scale filtering of decays for charmed baryon studies was produced on the UMiss farm. Krish Gounder reported results from this filter to the collaboration in September, 1992. There he discussed techniques for extracting $\Lambda \rightarrow p\pi^-$, $\Xi^- \rightarrow \Lambda\pi^-$ and $\Omega^- \rightarrow \Lambda K^-$ decays. He also showed a small number of $\Lambda_c \rightarrow \Lambda\pi\pi$ decays and $\Xi_c \rightarrow \Xi\pi\pi$ decays from two runs (Figure 8). From these studies and others we conclude that E791 will have good sensitivity to many charmed baryon decay channels. We are vigorously pursuing Λ , Σ_c , Ξ_c and Ω_c decay channels.

E791 FNAL Director's Review

In November, 1992, the E791 collaboration gave a presentation before Dr. Peoples, the Director of Fermilab, about the status of the experiment. Results of the reconstruction work done at OSU and UMiss were presented and a request was made for 1/2 of the farms currently at Fermilab (~4000 MIPS) for use in the data reduction. This would double the present E791 computing resources. Dr. Peoples stated that his "dictum" is that about 2/3 of the Fermilab computing be devoted to

(a) $\Lambda_c^+ \rightarrow \Lambda \pi \pi \pi$

(b) $\Xi_c^+ \rightarrow \Xi \pi \pi$

FIGURE 8. Suggestive charmed baryon signals extracted from UMiss farm runs .

collider experiments and 1/3 to fixed target. He did not foresee any additional computer purchases in FY93 so, in the most optimistic scenario, it appears that the most computing E791 could get from Fermilab would be about 1300 MIPS. This falls woefully short of our need. An official response from Dr. Peoples has not been received yet.

D_s decays

Danying Yi will pursue topics involving D_s decays for her dissertation research. We expect that with about 30% of the E791 data sample analyzed she will be able to do a competitive D_s lifetime measurement (1000 decays) and Dalitz plot analysis of the $D_s \rightarrow KK\pi$ system. In addition to these standard studies she will be exploring semileptonic D_s decays such as $D_s \rightarrow \phi \mu \nu$ as a second activity which keeps her in touch with the muon identification systems. E791 could have as many as 1000 fully reconstructed $D_s \rightarrow \phi \mu \nu$ decays. With about 30% of the data reconstructed even 200 - 300 $\phi \mu \nu$ decays can make a major impact on current measurements.

Recently Ms. Yi and Fermilab postdoc, Sudeshna Banerjee, have reconstructed $D_s \rightarrow \phi \pi$ at a level of about 10 events per run using a candidate driven vertex technique. This technique yields about twice the presently predicted decays. A result from one run is shown in Figure 10. If this new yield holds true a sample of 1000 $\phi \pi$ decays will be reconstructed with only 1/6 of the E791 data

sample. Ms. Yi and Sudeshna are working hard on this project.

FIGURE 9. $D_s \rightarrow \phi \pi$ decays from one E791 run.

Charmed Baryon Physics

Dr. Gounder has made much progress in understanding charm baryons that decay through Λ , Σ , Ξ and Ω secondaries. He has written some crucial code in these areas and will pursue charm baryon decays. Some estimates give $BR(\Lambda_c \rightarrow \Lambda X) = 25\%$ and $BR(\Lambda_c \rightarrow \Sigma X) = 10\%$. Branching ratio measurements have proven to be very difficult for these modes and there is much room for improvement. The resonant structure of these decays is fully unexplored at this time. In addition, we will be sensitive to $\Xi_c^+ \rightarrow \Xi X (\sim 30\%)$, $\Lambda K X (\sim 10\%)$, $\Sigma K X (\sim 15\%)$ and other doubly strange baryons.

Krish has already implemented searches for $\Lambda_c \rightarrow \Lambda \pi$, $\Lambda \pi \pi \pi$, πK_s and $\pi K \pi \pi$ decays with marked success. When 10-30% of the E791 data is reconstructed we should be sensitive to many charmed baryon decays, adding lifetime measurements and resonant structure analysis to existing data. We will specially look for $\Xi_c^+ \rightarrow \Lambda K \pi \pi$, $\Xi^- \pi \pi$, $\Xi_c^0 \rightarrow \Xi^- \pi$, $\Xi^- \pi \pi \pi$, $\Omega^- K$ and $\Omega_c^- \rightarrow \Xi^- K \pi \pi$ in the near future. We expect to isolate ~ 500 -800 decays of this sort with 30% of the E791 data sample.

Finally, it is worth noting the comment on the E791 data quality from Prof. John Cumalat. His message is inserted on the next page.

Network Communications Enhancements

The E791 analysis heavily relies on network communications. Software is updated nightly on the

To: SUMMERS@FNAL
Original_To: @USR\$ROOT29:[E791]MAIL791.DIS

From: COLOHE::J_CUMALAT "cowboy john" 31-OCT-1992
To: FNAL::APPEL
Subj: Physics Information from E769 and E791

Jeff,

I received your Fedex package yesterday afternoon. I also received information from Milind and Ron Sidwell. Thank you very much for your assistance.

I also wanted to congratulate you on the data quality from E791. The signals look remarkably clean and it appears that you will have large signals. I am rather impressed!

Based on your data I find it hard to believe that the Hyperon Charm experiment (E781??) will play a significant role in hadronic production of charm, except possibly in examining production issues connected with a strange quark in the beam projectile.

We may find it of mutual advantage to have a heavy flavor workshop between our two groups next summer. This is something that Mike Sokolof was pushing for last summer. We are further along in our analysis, but it looks like you will have from 2-3 times more data and your signal to noise could well be much better than ours. We are probably using similar parallel techniques that might be worth discussing.

Best Regards,
John Cumalat

farms as well as other sites and sent to the entire collaboration. In addition, we routinely rely on network file sending and retrieval throughout the E791 collaboration.. Thus, good network communications between UMiss, Fermilab and other institutions is critical.. In response to the HEP activity and a heavily used Supercomputing Center, the university has installed a T1 line between Oxford and Jackson, MS, making communications to our campus and Dec farm much more reliable and fast. We would estimate that our networking has been up over 98% of the time in FY92.

3. New Facilities

The HEP group is presently moving to facilities in the new \$10M National Center for Physical Acoustics(NCPA) at the University. A spacious room has been renovated for the installation of our Dec computing farm. Two large laboratories, offices and a dedicated HEP network system is included. The HEP group retains their former space for specialized tasks. This gives the group about 5000ft². Both physics department and University funds have supported this move. The farm will operate under independently controlled environmental conditions and be placed on the research building's constant backup power supply. The NCPA is serviced by a fiber optic network to the UMiss Supercomputer Center, which is connected via a T1 line to NSFnet. One of the many advantages to this new facility is the increased capability for Dec farm expansion.

4. SSC Research and Development Program

4.1 Introduction

As a member of the L-star and later the GEM and SDC collaborations the HEP group has contributed primarily to the hadron calorimetry efforts. UMiss was P.I.(with Yu. Kamyshkov of ITEP/U. Tennessee as co-P.I.) on an SSC-funded program dealing with liquid scintillators which involved two beam test periods at ITEP-Moscow this past winter. This led to our initial close collaboration with Russian scientists. Our group has also collaborated with groups in the SDC program especially on the hanging file calorimeter test. Analysis of some of the data from this test is being carried out by the UMiss group (in collaboration with ANL, FNAL and U. Tennessee) as well as some radiation damage studies (in collaboration with Memphis State University) for both GEM and SDC. Presently the UMiss group is responsible for research and development on the optics package and simulation calculations for the GEM fiber calorimeter (in collaboration with Boston University, Jackson State University and U. Tennessee). Analysis of the latest beam test data from the fiber calorimeter is being carried out by UCSD, Boston University, U. Tennessee and UMiss.

The Umiss group is involved primarily with the GEM project in SSC research and development.

Professor Reidy is spending his sabbatical leave during 1992-93 at the SSC Laboratory as a lead editor on the GEM Technical Design Report and pursuing studies devoted to scintillation hadron calorimetry. To provide Dr. Reidy with adequate resources a request was made by him to the DOE/HEP for \$10K in a lab service account at the SSCL. In addition, the SSCL has requested salary funds from DOE/HEP for Prof. Reidy while he is at the SSCL. Neither request has been acted on as of 12/1/92.

The funds for this FY92 SSC R&D program did not arrive at the university until late October, 1992. This has severely hampered this work and has negatively impacted the level of progress. In spite of this a considerable effort in detector simulation and test beam studies was undertaken. The UMiss group played a pivotal role in the design and fabrication of prototype modules which were used in the test beam and in the analysis of the data.

Research Personnel

This research will be carried out primarily by personnel in the University of Mississippi High Energy Physics group. These include (with FTE equivalent spent on this project) Professor James J. Reidy (0.8), Assistant Professor Lucien Cremaldi (0.25), Assistant Professor D. Summers (0.2), Dr. Brent Moore (1.0), Mike Booke - machinist (0.2), S. Harper - programmer (0.2), and three graduate students (half-time). A new postdoctoral associate has been requested from the TNRLC to spend full-time on this project. In addition, Associate Professor S. Jahan of Memphis State University (0.3) and Visiting Assistant Professor H. Zhang (0.3) will collaborate with us. Close collaboration will be maintained with the ORNL group as well as other members of the SSCintCal collaboration.

4.2 CALOR89 Development and Simulation Studies

With the adoption of the hybrid calorimeter by GEM the activities of the UMiss group must be increased if we are to play a leading role in this program. Three areas in which we currently provide the lead role are:

- the design and fabrication of the optics package for the fiber hadron calorimeter,
- the simulation calculations for the GEM calorimeter using CALOR89 and
- low dose rate (approximately the rate at the face of the hadron calorimeter at $\eta=2.5$ and $L=10^{34}$) radiation damage studies on fibers in a copper matrix (the GEM geometry).

Dr. Moore has assumed an increasingly fundamental role in the simulation of the GEM calorimeter with evaluation of proposed geometries a top priority. For maximum efficiency it is most attractive for him to interface closely with the group of visiting scientists and staff at ORNL who are devoted to the GEM R&D program. In a TNRLC proposal we have requested support for two Russian physicists at ORNL to facilitate this interaction. By appointing two Russian scientists with

expertise in simulations, Dr. Dimitri Onoprienko and Dr. Konstatin Schmakov, as Visiting Research Scientists on our faculty and stationing them at ORNL we get the benefit of close interaction with the simulation effort. Dr. Moore has worked with Dr. Schmakov on previous simulation problems and Dr. Schmakov attended our workshop last spring. With these scientists spending at least quarter time on our campus they will benefit from interactions with our faculty and students as well as providing overlap (and ownership) for us with the larger simulation program. This interaction will be especially valuable for us in the design of the optics package since these scientists have extensive experience in light transport in fiber studies. Their expertise in GEANT will complement ours in CALOR thereby providing a powerful set of tools with which to do simulation studies. Our activities in simulation studies are well described in the following material. A partial list of publications of interest include

- CALOR89: Calorimetry Analysis and Benchmarking, Proc. of the First International Conference on Calorimetry in High Energy Physics, 1990, pg. 497
- CALOR89 Calorimeter Simulations, Benchmarking and Design Calculations, Proc. of the Symposium on Detector Research and Development for the SSC, 1990, pg. 608
- User's Guide to CALOR89-U3G, UMS/HEP/91/016
- User's Guide to CALOR89-U3G on PDSF(SSCL), SDC-92-246
- Preliminary Beam Test Results of the GEM Copper/Scintillating Fiber Hadron Calorimeter Prototype, GEM TN-92-210
- Punchthrough Calculations for Neutrons Using CALOR89, SSCL-584

Presentations of interest include

- Status of CALOR89, Rocky Mountain Consortium Workshop, Boulder, CO, July, 1992
- Calorimetry Simulation Using CALOR89 on Unix Computers, SESAPS, Durham, NC, 11/91
- CALOR Simulations of the Hanging File Calorimeter, SDC Calorimeter Meeting, 10/91

4.3 Scintillating Fiber/Plate Hadron Calorimeter Test Beam Analyses

Work is continuing on the analysis of the data from the test beam run at BNL last summer. We should complete this within the next six months. A second test beam run has been planned and funds for it requested from the TNRLC so we can complete this series of studies and prepare formal publications of the results. A preliminary evaluation of the test beam data can be found in

- Preliminary Beam Test Results of the GEM Copper/Scintillating Fiber Hadron Calorimeter Prototype, R. Carey, J. Miller, L. Sulak, B. Worstell, H. Paar, B. Moore, J. Reidy and R. Kroeger, GEM TN-92-210 (October 7, 1992).

4.4 Scintillating Fiber/Plate Hadron Calorimeter Studies for GEM

Research Objectives

In October the GEM Collaboration picked the hybrid configuration as the calorimeter for the GEM Detector. Inherent in this design is the use of the scintillating fiber technology in the hadron calorimeter. This choice followed an extensive R&D program of a pre-prototype fiber hadron calorimeter carried out by the SSCintCal collaboration. During the past year this pre-prototype was assembled and tested at Brookhaven National Laboratory. Over ten GEM technical notes have been published describing the work in this R&D program. This work contributed to GEM selecting fiber technology for the hybrid hadron calorimeter.

During the past year the UMiss High HEP group provided the leadership role in simulation calculations using CALOR89 as well as designing and fabricating the optical package for the preprototype towers. The group was one of the principals in the test beam study and has been one of the three groups involved in the data analysis for the pre-prototype tower test beam data. The primary objective of our FY93 study will be to design and optimize a fully hadronic shower containing set of scintillating fiber prototype towers embodying the GEM SSC calorimeter design developed for the TDR. It will include the integration of the design of these towers with that of the entire hadronic calorimeter system.

Although Monte Carlo simulations have shown that for jets, the undetected energy in the dead material between the EM cryostat and hadronic section is negligible compared to the energy of the neutrinos there remains a concern by some that this could lead to degradation of the performance of the calorimeter. For this reason we are working in collaboration with ORNL personnel to initiate a study of a "massless gap" for this region. It might consist of a layer of scintillating tiles with WLS readout suspended from the front face of the hadron calorimeter. The fibers could be threaded through the fiber/copper sections to PM tubes. Its signal can be used to correct for the energy loss of electrons and hadrons in the dead region. SDC has foreseen a similar device to correct for energy loss in their coil. The "massless gap" will also provide an additional longitudinal sample and will be useful for π/e rejection and muon tracking inside the calorimeter. During the past year the University of Mississippi has studied the optical coupling between the wavelength shifting fibers and plastic scintillating plates in anticipation of the need for such a device. The results for one configuration are shown in Figures 10 and 11. We are a member of the hanging file calorimeter collaboration which has its roots in SDC and have participated in the data analysis from this test beam program. In the hanging file $1 \times 1 \times 0.005 \text{ m}^3$ scintillator plates were used. Our studies on these scintillating plates will be incorporated into our massless gap program. The preliminary simulations appear quite encouraging. The energy deposition in this plate is combined with the energy detected in the electromagnetic and hadronic sections. With proper weighting, significant enhancement of the resolution can be obtained. Figure 12 shows the results of a simulation for a 100 GeV pion using a 25λ liquid krypton/copper electromagnetic section followed by the "massless gap" scintillator and then the copper/fiber hadron section. The top curve shows a linearly weighted sum of the electromagnetic plus hadronic section outputs. The center curve shows a linearly weighted sum including the "massless gap". The bottom curve shows the

Figure 10a: Relative count rate vs. position

Figure 10b: Tile/fiber groove configuration

(a)

Figure 11

(b)

Figure 12

significant improvement which one obtains if one uses a functional form for the weight of the "massless gap". These are preliminary simulations but we think that the results indicate that this technique should be given serious consideration. For this reason we are continuing the simulation effort to better understand the weighting and the scintillator studies to aid in designing a scintillating plate "massless gap" arrangement for the beam test studies. The optical studies should be finished by 9/93; the simulation studies depend on the calorimeter specifics and should provide some definitive directions two to three months after the Baseline design is chosen.

4 .5 Infrastructure Development

As part of the TNRLC-funded infrastructure program with SAHEP the UMiss group submitted an institutional plan for the period 1990-1999. Some excerpts from this plan are presented here to give DOE an idea of the extent that the group is attempting to integrate funds from the TNRLC into its program.

The Institutional Plan for the University of Mississippi High Energy Physics Group

Personnel and Support

The HEP group consists of four faculty (Assistant Professors Lucien Cremaldi and Don Summers, Professor James Reidy and Visiting Assistant Professor John Weinstein), two postdoctoral associates (Krish Gounder and Brent Moore), six graduate students (two writing their dissertations; three of the students are female) and about three undergraduate students. Prof. Weinstein and Dr. Moore position are funded by the TNRLC through the infrastructure program. The physics department machinist/instrument maker is available at no charge at approximately 0.5 FTE and the electronic technician at 0.5 FTE. The computer system manager, Sandy Harper, is shared between the high energy physics group and the acoustics group. Dr. Cremaldi spends about 1/4 time and Dr. Moore spends full time on SSC-related research; Dr. Reidy spends about half-time (except for this year when it is full-time as he is at the SSCL on sabbatical) on the SSC program.

The group was formed in 1989 and initially placed in a 2500 sq. ft. renovated underground facility which had housed a 3 MeV Dynamatron accelerator. In December 1992, the group will move to space in a new \$10M National Center for Physical Acoustics facility on the university campus. The old space which includes a high bay area will be retained by the group for research and development activities. In addition, a \$2.1M wing is being added to the physics building and will house the HEP group beginning in 1995. Using funds from the TNRLC to leverage approximately \$80,000 from DoE in equipment the UMiss HEP group has established a 42 CPU DECstation 5000/200 900 MIP computer "farm". In addition, the group has a VAXstation 3100, a VAXstation 4000, a microVAX II, a Silicon Graphics Personal Iris workstation, a Quadra 950 and approximately 10 other Macintosh computers. The computing system is connected to the

university mainframe via fiber optics. A T-1 line connects the mainframe with NFSnet. Machine shop facilities include 3 computer controlled milling machines and 2 computer controlled lathes. The UMiss HEP group receives funding from the DOE, TNRLC and the University of Mississippi. The TNRLC support for the period 4/91-3/93 and proposed support for 4/93-3/94 along with matching and leveraged support is presented in Table V on the next page. The 1992 TNRLC support includes \$10K for scintillating fiber research and \$90K for infrastructure support. Current and future support from all sources including faculty research time and university assumption of the TNRLC faculty position in 1998 is presented in Table VI. The university contribution for the past two years has been about \$200K per year. This includes 50% research time for the faculty, support for two graduate students and two undergraduates, overhead reduction for DOE, overhead waiver for TNRLC and direct cash contribution of \$30K per year as well as secretarial and shop support. In addition, about \$25K has been spent to provide suitable space for the high energy computer "farm" and other equipment. These contributions do not include the new research space which has been assigned to the HEP group. In Table VII is shown the current and proposed distribution of TNRLC Infrastructure funds.

Ten Year Research Plan: 1990-1999

We would like to expand our involvement in both basic physics research and SSC R&D. To maintain our intellectual vigor, for the benefit of our junior faculty and as an avenue for our students to obtain quality PhD's, we must initiate an involvement in a new basic physics experiment within the next year or so while we carry on analysis of the E791 data. This will most likely be a collider experiment. In addition, with the GEM detector proceeding from the TDR to construction phase we must expand our involvement in the calorimeter if we are to maintain a significant role in the GEM program. Funding which can only be obtained through the SSC/TNRLC program is critical to maintain our group effort. In order to proceed at a nondestructive pace we see that, at a minimum, we need an additional post-doc and one research scientist as well as a full time computer system manager. To provide much needed balance we would also welcome the addition at least two theorists. The TNRLC funds are currently supporting one post-doc and one faculty member. The University has agreed to assume support for the faculty position after five years of TNRLC support. Given the long range plan at the University it is unlikely that they will agree to support a further position in HEP. However, this could change dramatically if additional support from DOE/TNRLC were to provide funds for additional technical support and research scientists for a number of years. The University is presently initiating a review of the physics department and external support for the HEP program will play a major role in the response of the review team to the University about HEP needs.

To carry out our responsibilities in the experimental fiber calorimeter program which are mainly in the optics design and testing, radiation damage and scintillator evaluation areas we need a full-time **research scientist**. This person would oversee the UMiss GEM calorimeter effort. He/she would be expected to interface with the other collaborative groups as well as the simulation studies

TABLE V- TNRLC-BASED DISTRIBUTION

Category	4/91-3/92			4/92-3/93			4/93-3/94		
	TNRLC	Univ.	Lever-	TNRLC	Univ.	Lever-	TNRLC	Univ.	Lever-
	Match	Match	aged	Match	Match	aged	Match	Match	aged
PERSONNEL(FTE)									
Faculty	1.0		0.8	1.0	0.3	1.0	1.0		1.0
Perm. Res. Sci.							0.75		
Postdocs	1.0			1.0			1.0		
Grad. Students	0.3		1.0		0.5	2.0	0.5	0.5	2.0
Engineers									
Technicians			0.8			0.8			0.8
Undergrads			1.0			1.0			1.0
Other Personnel			0.2			0.25			0.25
TRAVEL(\$)	\$10K	\$ 3K		\$10K	\$ 5K		\$ 8K		
EQUIPMENT(\$)								\$ 6K	
SPACE(ft ²)			500			2500			4000
OTHER(list)			\$ 41K			\$ 45K			\$110K

TABLE VI - FUNDS FROM ALL SOURCES

SOURCE	1992	1993	1994	1995	1996	1997	1998	1999
DOE-NSF/HEP	\$ 150K	\$ 200K	\$ 260K	\$ 300K	\$ 350K	\$ 395K	\$ 450K	\$ 500K
DOE/SSCL	145K	100K	120K	160K	180K	180K	190K	230K
TNRLC-Res.	10K	110K	130K	150K	190K	190K	150K	0
TNRLC-Infra.	90K	135K	200K	200K	200K	125K	60K	0
University	161K	171K	195K	210K	240K	260K	360K	385K
TOTAL	\$ 550K	\$ 716K	\$ 905K	\$ 1020K	\$ 1160K	\$ 1100K	\$ 1210K	\$ 1115K

TABLE VII - TNRLC INFRASTRUCTURE FUNDS

CATEGORY	1992	1993	1994	1995	1996	1997	1998	1999
Personnel	\$ 80K	\$ 127K	\$ 143K	\$ 154K	\$ 165K	\$ 120K	\$ 55K	\$0
Equipment			47K	36K	25K			0
Travel	10K	8K	10K	10K	10K	5K	5K	0
Other								0
TOTAL	\$ 90K	\$ 135K	\$ 200K	\$ 200K	\$ 200K	\$ 125K	\$ 60K	0

being carried out at UMiss. This person could have a joint appointment with ORNL thereby strengthening the interactions with that lab but financial support would have to come from UMiss. Initially, support would come from TNRLC but this should be coupled with a commitment from DOE to provide increasingly greater support as time goes on. It is expected that at least two graduate students would be involved in this work at any given time. A very talented individual has already been identified who wishes to apply for this position. A **third postdoctoral associate** is needed to work half-and-half on the basic physics and SSC programs. This person would work closely with the research scientist on the SSC projects and with the faculty on the basic physics program. We have also identified a qualified individual for this position. Prof. Weinstein's position will most likely be picked up by the university so the present TNRLC funds will be used to hire an additional **faculty member** in experimental high energy physics. This person will be expected to spend at least quarter time on the SSC program. It is possible that this individual will chose to contribute to areas other than calorimetry and to a new basic physics area. We presently try to scrape up funds from several sources to pay a student as **system manager for the computer "farm"**. This is obviously not very satisfactory since the complexity of the "farm", the high level of responsibility assigned our group in the E791 data analysis and the great need for the "farm" in calorimeter simulation calculations requires greater continuity and expertise than one usually finds with students. We have nearly \$400,000 invested in the "farm" and a full-time system manager is needed to oversee and protect this investment. This person should be shared between DOE and TNRLC. To carry out the radiation damage and optics studies as well as provide us with some basic research equipment (other than computers) we request **modest equipment support**. Our present non- computer equipment inventory amounts to less than \$60K.

Finally, the HEP group has a commitment to assist faculty and students at Jackson State University(an HBCU) in participating in a physics research program. Besides obvious benefits to

their faculty we see this as an opportunity to attract minority students into physics research. However, to do this requires a great deal of time even though two faculty members have a background in high energy physics. The research scientist and new faculty member could provide the additional manpower for this. The Universities have expressed a willingness to provide some matching funds for such a project.

Efforts are also underway to form a close collaboration with Vanderbilt, U. Tennessee and ORNL for SSC research.

5. Proposals Submitted

During FY92 the following proposals were submitted to external agencies.

Title	Date	Agency	Amount
Heavy Quark Trigger Processor	12/9/91	DOE	250000
SAHEP Initiative for SSC Research and Education (Supplement)	3/26/92	TNRLC	\$ 26800
Southern Association for High Energy Physics Initiative for SSC Research and Education	4/16/92	TNRLC	90000*
Study on a New Scintillating Plate-WLS Fiber Hadron Calorimeter Design	3/31/92	TNRLC	73838
Personnel to Assist in the Development of Simulation and Optics Package for GEM Calorimetry	10/29/92	TNRLC	100995**
Studies Devoted to GEM Scintillating Fiber Calorimetry for GEM	10/29/92	TNRLC	165000**
* Funded			
** Pending			

6. Summary

The University of Mississippi High Energy Physics group is active in three Fermilab experiments and the SSC R&D program. During FY92 most of our effort have been devoted to running on E791 where we had extensive on-line responsibilities dealing with the Cerenkov counters, PWCs, low voltage monitoring system, beam counters, a "muon wall" of scintillation counters and part of

the data acquisition system. Since the end of the run in February, 1992, the emphasis has shifted toward the assembly and checkout of a 900 MIP DECstation 5000/200 computing farm which will be used for the analysis of E791. That checkout is now complete and production has begun. Preliminary physics results have been obtained with emphasis on charm semileptonic decays and charmed baryon decays.

The UMiss group is involved primarily with the GEM project in SSC research and development. Professor Reidy is spending his sabbatical leave during 1992-93 at the SSC Laboratory as a lead editor on the GEM Technical Design Report and pursuing studies devoted to scintillation hadron calorimetry. Our group has also collaborated with groups in the SDC program especially on the hanging file calorimeter test. Analysis of some of the data from this test is being carried out by the UMiss group (in collaboration with ANL, FNAL and U. Tennessee) as well as some radiation damage studies (in collaboration with Memphis State University) for both GEM and SDC. Presently the UMiss group is responsible for research and development on the optics package and simulation calculations for the GEM fiber calorimeter (in collaboration with Boston University, Jackson State University and U. Tennessee). Analysis of the latest beam test data from the fiber calorimeter is being carried out by UCSD, Boston University, U. Tennessee and UMiss.

APPENDIX

UNIVERSITY OF MISSISSIPPI
PUBLICATIONS, TALKS, AND TECHNICAL NOTES

1. Breese Quinn, Finding Charm Quarks with Parallel RISC Computers and 8mm Tapes, Meeting of the APS Southeastern Section, Durham, North Carolina (11-13 November 1991); Bulletin of the American Physical Society 36 (1991) 2748.
2. Brent Moore, Calorimetry Simulation Using CALOR89 on UNIX Computers, Meeting of the APS Southeastern Section, Durham, North Carolina (11-13 November 1991); Bulletin of the American Physical Society 36 (1991) 2749.
3. James J. Reidy, SAHEP and the TNRLC - Partners in a Vision, Meeting of the APS Southeastern Section, Durham, North Carolina (11-13 November 1991); Bulletin of the American Physical Society 36 (1991) 2749.
4. D. J. Summers, High Speed Data Acquisition at the Fermilab E791 Charm Quark Factory, Meeting of the APS Southeastern Section, Durham, North Carolina (11-13 November 1991); Bulletin of the American Physical Society 36 (1991) 2749.
5. John Weinstein, Low Mass Pseudoscalar Pseudoscalar Dynamics, Meeting of the APS Southeastern Section, Durham, North Carolina (11-13 November 1991); Bulletin of the American Physical Society 36 (1991) 2745.
6. Don Summers, High Speed Data Acquisition at the Fermilab E791 Charm Quark Factory, Physics Seminar, University of Chicago (19 August 1991).
7. Don Summers, High Speed Data Acquisition at Fermilab E791, CERN Computing and Networking Division Seminar, Genève, Switzerland (12 March 1992).
8. J.C. Anjos et al. (E691), Study of the Decays $D^+ \rightarrow K\pi\pi e\nu$ and $D \rightarrow K^*\pi e\nu$, Physical Review D45 (1992) R2177.
9. J.C. Anjos et al. (E691), Experimental Probes of Final-State Interactions in D^0 - Meson Decays, Physical Review D46 (1992) R46.
10. J.C. Anjos et al. (E691), Study of the Decays $D^+ \rightarrow K 3\pi$, Physical Review D46 (1992) 1941.
11. J.C. Anjos et al. (E691), Study of the Doubly Cabibbo-Suppressed Decay $D^+ \rightarrow \phi K^+$ and the Singly Cabibbo-Suppressed Decay $D_S^+ \rightarrow \phi K^+$, Physical Review Letters 69 (1992) 2892.
12. J.C. Anjos et al. (E691), A Dalitz Plot Analysis of $D \rightarrow K\pi\pi$ Decays, (in preparation, draft attached).

13. C. Stoughton and D. J. Summers (E769), Using Multiple RISC CPUs in Parallel to Study Charm Quarks, *Computers in Physics* 6 (1992) 371.
14. G.A. Alves et al. (E769), Feynman- x and Transverse Momentum Dependence of D^+ and D^0 , $\bar{D}^0$ Production in 250 GeV π^- - Nucleon Interactions, *Physical Review Letters* 69 (1992) 3147.
15. G.A. Alves et al. (E769), Atomic Mass Dependence of D^+ and D^0 , $\bar{D}^0$ Production in 250 GeV π^- - Nucleon Interactions, (submitted to *Physical Review Letters*); FERMILAB-Pub-92/279-E.
16. Ali Rafatian, Study of $D^{*+} \rightarrow D^0(K^-\pi^+\pi^-\pi^+)\pi^+$ Production in 250 GeV π^- - N Interactions, E769 Technical Note.
17. D.J. Summers et al., Charm Physics at Fermilab E791, Electroweak Interactions and Unified Theories, XXVIIth Rencontre de Moriond, Les Arcs, France (15-22 March 1992); UMS/HEP/92-020.
18. S. Amato, J.R.T. de Mello Neto, J. de Miranda, C. James, D.J. Summers, and S.B. Bracker, The E791 Parallel Architecture Data Acquisition System, *Nuclear Instruments and Methods A* (accepted for publication); UMS/HEP/92-019.
19. Lucien Cremaldi, E691 Dalitz Plot Analysis of $\Lambda_c^+ \rightarrow pK^-\pi^+$ Decays, Proceedings of the E791 Charm Physics Workshop, University of California, Santa Cruz (27-31 July 1992) 142.
20. D.J. Summers, Charm Strange Baryons and the 1.5 Prong, Proceedings of the E791 Charm Physics Workshop, University of California, Santa Cruz (27-31 July 1992) 147.
21. L.M. Cremaldi et al., Fermilab E791, XXVIth International Conference on High Energy Physics, Dallas, Texas (6-12 August 1992); UMS/HEP/92-021.
22. B. Moore, A. Van Smith, and J. Reidy, Punchthrough Calculations for Neutrons Using CALOR89, SSCL-584 (September 1992).
23. R. Carey, J. Miller, L. Sulak, B. Worstell, H. Paar, B. Moore, J. Reidy, and R. Kroeger, Preliminary Beam Test Results of the GEM Copper/Scintillating Fiber Hadron Calorimeter Prototype, GEM TN-92-210 (7 October 1992).
24. Brent R. Moore, User's Guide to CALOR89-U3G on PDSF, SDC-92-246 (14 April 1992).
25. John Weinstein, Remarks on Low Mass Pseudoscaler-Pseudoscaler Dynamics Including the S-Wave $I = 1$ KK System, (submitted to *Physical Review D*); UMS/HEP/92-018.

2 abstracts, reprints,
preprints removed;
conf papers cycled
separately.

**DATE
FILMED**

02/19/93