9
1-13-93 JS (2)

PREPARED FOR THE U.S. DEPARTMENT OF ENERGY,
UNDER CONTRACT DE-AC02-76-CHO-3073

PPPL-2873
UC-421

PPPL-2873

K β SPECTRA OF HELIUMLIKE IRON FROM TFTR PLASMAS

BY

A.J. SMITH, M. BITTER, K.W. HILL, ET AL.

DECEMBER, 1992

PPPL

PRINCETON
PLASMA PHYSICS
LABORATORY

PRINCETON UNIVERSITY, PRINCETON, NEW JERSEY

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

NOTICE

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial produce, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

NOTICE

This report has been reproduced from the best available copy.
Available in paper copy and microfiche.

Number of pages in this report: 25

DOE and DOE contractors can obtain copies of this report from:

Office of Scientific and Technical Information
P.O. Box 62
Oak Ridge, TN 37831;
(615) 576-8401.

This report is publicly available from the:

National Technical Information Service
Department of Commerce
5285 Port Royal Road
Springfield, Virginia 22161
(703) 487-4650

K β Spectra of Heliumlike Iron from TFTR Plasmas

A. J. Smith

Lock Haven University, Lock Haven, Pa 17745

M. Bitter, H. Hsuan, K. W. Hill, S. von Goeler, and J. Timberlake

*Plasma Physics Laboratory, Princeton University, P.O. Box 451, Princeton New Jersey
08543*

P. Beiersdorfer and A. Osterheld

Lawrence Livermore National Laboratory, Livermore, CA 94550

MASTER

Abstract

$K\beta$ spectra of heliumlike iron, Fe XXV, have been observed from TFTR plasmas with a high-resolution crystal spectrometer. The wavelength range of the Fe $K\beta$ spectrum partially overlaps the spectrum of heliumlike nickel (Ni XXVII), which is used on TFTR and JET for ion temperature measurements. The experimental arrangement made it possible to observe the Fe XXV $K\beta$ lines and their satellite transitions of the type $1s2l'3l'' \rightarrow 1s^22l'$, as well as the entire satellite spectrum of the Ni XXVII $K\alpha$ line simultaneously. In order to identify the features of the $K\beta$ spectra and to study their possible interference with the Ni XXVII spectrum, the intensity of the $K\beta$ spectrum was enhanced by injection of iron into the plasma. Accurate wavelengths and intensities have been obtained and compared with different theoretical calculations.

*Supported by DOE Contract No. DE-AC02-76-CHO-3073

I. Introduction

High resolution satellite spectra of high-Z atoms are important for the diagnostics of tokamak plasmas and solar flares, and they are also of interest in testing atomic theories which must include relativistic and quantum electrodynamics effects. In the large tokamaks such as the Tokamak Fusion Test Reactor (TFTR), the Joint European Torus (JET), the large Japanese tokamak JT-60 U, and Tore Supra, where large volume ($>40\text{m}^3$) plasmas with peak electron temperatures $\sim 5\text{ keV}$ are maintained for several seconds, hydrogenlike and heliumlike ions of high-Z atoms such as titanium, chromium, iron, and nickel, are readily produced. Line radiation from these highly stripped ions is used for electron and ion temperature diagnostics as well as for determination of charge-state distributions and ion transport coefficients. In particular, the resonance line of heliumlike nickel¹⁻⁴ is employed on TFTR, JET and Tore Supra for Doppler broadening and Doppler shift measurements in order to determine the central ion temperature as well as toroidal and poloidal plasma velocities.

In the case of TFTR and Tore Supra which use stainless steel ($\sim 70\text{ at. \% Fe}$, $\sim 20\text{ at. \% Cr}$, $\sim 7\text{ at. \% Ni}$) walls and Inconel ($\sim 70\text{ at. \% Ni}$, $\sim 15\text{ at. \% Cr}$, $\sim 7\text{ at. \% Fe}$, $\sim 2.5\text{ at. \% Ti}$) bellows protective cover plates, iron and nickel are simultaneously present as wall-derived impurities. Thus the possibility exists that the FeXXV $K\beta$ spectrum and the NiXXVII $K\alpha$ spectrum, which fall in the wavelength range from 1.57 to $1.62\text{ \AA}$, interfere with each other. It is therefore of considerable interest to determine whether a distortion of the Ni $K\alpha$ spectrum by the Fe $K\beta$ spectrum impairs the ion temperature measurements. In this paper, we investigate the $K\beta$ spectrum of heliumlike iron. Such a study is very important since experimental data on $K\beta$ spectra are scarce.

Under normal TFTR operating conditions prior to 1987, when the present data were collected, the nickel $K\alpha$ radiation was much more intense than the iron $K\beta$ radiation.⁵ Iron was therefore injected into the plasma by a laser blow-off technique in order to enhance the iron $K\beta$ emission relative to the nickel $K\alpha$ emission. Thus it was possible to extract the Fe $K\beta$ spectrum from the obtained composite spectrum and to clearly identify most of the theoretically predicted features.

The experimental wavelengths of the $K\beta$ lines and the associated satellite features were determined relative to the positions of nickel $K\alpha$ lines with an accuracy of $\sim 0.1\text{ m\AA}$.

An important result of our experiments is that the $K\beta$ features are found to be well separated from the nickel $K\alpha$ resonance line, so that there is no interference with the central ion temperature measurement. As an additional benefit, observation of the time history of the $K\beta$ radiation following the iron injection makes it possible to study the confinement of metal impurity ions in TFTR plasmas.

The paper is organized as follows: Section II presents theoretical data on the iron $K\beta$ satellite spectrum obtained from the Hebrew University-Lawrence Livermore Atomic Code (HULLAC) package. The experimental results and data analysis are described in Section III. Conclusions are given in Section IV.

II. Theory

The spectrum of interest includes the $K\beta$ lines of heliumlike iron, corresponding to the transitions $1s^2\ ^1S_0 - 1s3p\ ^1P_1$ and $1s^2\ ^1S_0 - 1s3p\ ^3P_1$, and the associated lithiumlike and berylliumlike satellites due to the transitions: $1s^22l - 1s2l'3l''$ and $1s^22s2l' - 1s2s2l'3l''$. These satellites can be formed (1) in the process of dielectronic recombination of heliumlike or lithiumlike ions or (2) by collisional inner-shell excitation of lithiumlike and berylliumlike ions from their ground states. In the first process a free electron is captured by a heliumlike or lithiumlike ion into the $n = 3$ level :

$$1s^2 + e^- \rightarrow 1s2l'3l'', \quad (1)$$

or

$$1s^22s + e^- \rightarrow 1s2s2l'3l''. \quad (2)$$

The resulting autoionizing states given in Eq. (1) can decay by two types of radiative decay schemes :

$$1s2l'3l'' \rightarrow 1s^23l, \quad (3)$$

or

$$1s2l'3l'' \rightarrow 1s^22l. \quad (4)$$

The first type provides satellites to the heliumlike $K\alpha$ resonance line. These, so-called unresolved $n=3$ satellites, lie very close to the $K\alpha$ resonance line and have been investigated rather well.⁶ Of interest for the present paper are the transitions of the second type which give rise to $n = 2$ satellites of the $K\beta$ lines. Because the plasma also contains lithiumlike ions, we expect to find satellite transitions from the dielectronic recombination of lithiumlike ions by radiative decay of the excited states given in Eq. (2):

$$1s^2 2s + e^- \rightarrow 1s2s2l'3l'' \rightarrow 1s^2 2s2l'. \quad (5)$$

The relevant transitions due to the second process of direct electron-impact excitation of inner-shell electrons from lithiumlike and berylliumlike charge states involve only the ground states of these ions:

$$1s^2 2s + e^- \rightarrow 1s2s3l + e^- \quad (6)$$

and

$$1s^2 2s^2 + e^- \rightarrow 1s2s^2 3l + e^-, \quad (7)$$

since the so-called zero-density limit applies to the low electron densities of TFTR plasmas.

In the following, we discuss the line emissivities for ions from different charge states. The emissivities of the heliumlike $K\beta$ lines can be represented by

$$\epsilon_{H\epsilon} = N_e N_{H\epsilon} C \quad (8)$$

where N_e and $N_{H\epsilon}$ are the electron and heliumlike ion densities and C is an effective excitation rate coefficient which takes into account direct excitation and radiative cascades. These rate coefficients were obtained with the HULLAC set of atomic codes, which calculate wave functions, energy levels, and radiative transition rates by the relativistic, multi-configuration parametric potential method with full configuration interaction.⁷ The collisional excitation rates were calculated in the distorted-wave approximation; an efficient technique, which performs an angular factorization of both the direct and exchange contributions to excitation cross sections and interpolates the necessary radial integrals as a function of threshold energy⁸ was used. The excitation

rate coefficients obtained from these calculations are given in Table I as a function of the electron temperature.

The emissivities of the so-called collisional satellites lines which result from inner-shell electron impact excitation of the lithiumlike and berylliumlike charge states are given by

$$\epsilon_{cs} = N_e N_{Li(Be)} C \frac{A_r^f}{\sum_f A_r^f + \sum_i A_a^{si}} \quad (9)$$

where $N_{Li(Be)}$ is the density of the lithiumlike or berylliumlike charge state, A_r^f is the radiative transition probability for the transition in question, and A_a^{si} is the autoionization probability to some final state i . The excitation rate coefficients C for the lithiumlike and berylliumlike ions were obtained by the same methods as were the rate coefficients for the heliumlike ions, while the autoionization rates were obtained from a factorization-interpolation technique using a single consistent potential for the bound and continuum orbitals.⁹ The excitation rate coefficients and transition probabilities for Fe XXIV and Fe XXIII are also listed in Table I.

The emissivities of the lithiumlike dielectronic satellites of the $K\beta$ lines can be represented by

$$\epsilon_d = N_e N_{He} F_1^*(T_e) F_2^*(s), \quad (10)$$

where the factor

$$F_1^*(T_e) = \frac{1}{2} \left[\frac{2\pi\hbar}{mkT_e} \right]^{3/2} \exp(-E_e/kT_e) \quad (11)$$

depends only on the electron temperature, and

$$F_2(s) = \frac{s_e}{s_i} \frac{A_r^f A_a^{si}}{\sum_i A_a^{si} + \sum_f A_r^f} \quad (12)$$

is the line strength. Here g_s and g_i are the statistical weights of the autoionizing states and the ground states, respectively, and E_s is the energy of the satellite level above that of the heliumlike ground state. The emissivities of the berylliumlike dielectronic satellite lines are given by an expression similar to (4) except that N_{He} is replaced by N_{Li} . The intensity factors $F_2(s)$ for the strong lithiumlike and berylliumlike dielectronic satellites calculated with the HULLAC package are listed in Tables II and III, respectively.

III. Experimental Results.

The spectra were obtained from five nearly identical ohmically heated TFTR discharges with the following parameters: The plasma current and the toroidal magnetic field were $I_p = 1.4$ MA and $B_T = 3.2$ T, and the plasma major and minor radii were $R = 2.48$ m and $a = 0.80$ m. These discharges were in steady-state conditions during the period from 1.5 to 4.5 s and had a peak electron temperature of 2.7 keV and a peak electron density of $4.3 \times 10^{13} \text{ cm}^{-3}$. Figure 1 shows the time evolution of the electron temperature T_e and electron density N_e for the TFTR discharge number 28931, which is typical of the discharges used in this experiment. The electron temperature and density data were obtained from electron cyclotron measurements and laser interferometry. Iron was injected into the plasma at 2.7 s by a laser blow-off technique. The injection led to a substantial increase (30 %) of the total radiated power, but did not affect the central electron temperature and electron density, see Fig. 1.

The spectral data were recorded with the TFTR horizontal and vertical curved crystal spectrometers.^{10,11} The horizontal spectrometer was used to measure the composite spectra of the NiXXVII $K\alpha$ and the FeXXV $K\beta$ radiation. This spectrometer permits observation of spectral features in the wavelength range 1.57 to 1.62 Å, with a spectral resolving power of $\lambda/\Delta\lambda = 7500$. The vertical spectrometer was used for simultaneous observation of the FeXXV $K\alpha$ spectrum near 1.85 Å. Both spectrometers recorded 128 spectra from a discharge with a time resolution of 10 ms per spectrum, starting at 2.5 s.

Figure 2 shows the time history of the FeXXV $K\alpha$ and $K\beta$ line intensities. These line intensities reach a maximum at 2.770 s, i.e. 70 ms after the injection of iron. The intensities then decay exponentially with an e-folding time of 300 ms, which is a measure of the particle confinement time. A more detailed understanding of these data requires

plasma modeling calculations, e. g. with the Multi-Ion Species Transport (MIST) code,¹² which takes into consideration the processes of ionization and recombination as well as ion transport due to diffusion and convection.

Figure 3 presents composite satellite spectra of the iron $K\beta$ and nickel $K\alpha$ lines from different time periods: (a) the period from 2.5 to 2.7 s before the injection of iron, and (b) the period from 2.75 to 2.88 s after injection, when the radiation from the injected iron is greater than 50% of its maximum value (see Fig. 2). The satellite spectrum of the NiXXVII $K\alpha$ line has been described earlier by Hsuan *et al.*¹ and by Bitter *et al.*⁴ The dominant features of the FeXXV $K\beta$ spectrum, the $1s^2\ ^1S_0 - 1s3p\ ^1P_1$ and $1s^2\ ^1S_0 - 1s3p\ ^3P_1$ transitions, are on the short wavelength side of the NiXXVII $K\alpha$ line between channels 140 and 160. It is evident from Figs. 3 (a) and (b) that the iron $K\beta$ spectrum is very weak compared to the nickel spectrum under normal operating conditions in TFTR, but that it is enhanced by an order of magnitude following the iron injection. The second spectrum, Fig. 3(b), also shows some $K\beta$ satellite lines in the neighborhood of the nickel $K\alpha$ line *w* and in the wavelength range of the nickel features *x*, *s* and *t*, see Gabriel's notation.¹³ These features are not apparent in Fig. 3(a). The nickel resonance line *w*, the dielectronic satellite line *j* and the combined lines *q* and *y*, as well as *l*, β and *z* (see Gabriel's notation¹³) are not distorted by the iron lines, even during strong iron injection.

Since the electron temperature T_e and the electron density N_e were not affected by the iron injection, the nickel spectrum of Fig. 3(a) can be normalized and subtracted from the composite nickel-iron spectrum of Fig. 3(b). The experimental iron $K\beta$ spectrum obtained in this way is shown in Fig. 4(a). The main features of the observed spectrum include the $K\beta_1$ and $K\beta_2$ lines (labeled 1 and 2) and the more prominent satellite transitions 3 through 9 which are identified in Table 4. In order to determine wavelengths of the observed spectrum accurately, the central channel numbers for the prominent peaks have been obtained by a least squares fit of Voigt profiles to the experimental data. These central channel numbers have then been converted to wavelengths using the equation

$$\lambda = 2d \sin(\theta_0 + \Delta\theta), \quad (13)$$

where $2d = 2.028\ \text{\AA}$ is the 2d-spacing of the quartz crystal (cut: 2243) and $\Delta\theta\ (\text{deg}) = 9.6880 \times 10^{-3}(N - N_0)$ is the dispersion of the instrument. N_0 and θ_0 are the channel number and the Bragg angle for the nickel line *w*. Since the spectrometer has no

absolute wavelength calibration, the theoretical wavelength value of 1.5884 Å for the nickel w line from Safronova's calculations¹⁴ is used as a reference value, i. e. we assigned $\lambda_0^{theor} = 1.5884 = 2d \sin(\theta_0)$ to channel number N0. The experimental wavelengths of the prominent features are shown in Table 4 together with the theoretical wavelengths of the present calculations, see Tables II and III, and those from the calculations by Safronova,¹⁴ Chen¹⁵ and Nilsen.¹⁶ Safronova uses the 1/Z expansion perturbation technique, with relativistic corrections taken into account within the framework of the Breit operator. Chen¹⁵ computes the relevant atomic data relativistically in intermediate coupling using configuration interaction in the multi-configuration Dirac-Fock model (MCDF); a generalized Breit interaction and QED corrections were also taken into account. The third set of calculations by Nilsen uses MCDF for bound states and distorted-wave Dirac continuum states for electrons.

The different sets of theoretical wavelengths listed in Table 4 are in good general agreement with the experimental wavelengths, given that these wavelengths can be shifted by a constant amount due to the fact that the spectrometer is not absolutely calibrated. The wavelengths of Safronova's calculations seem to be more consistent with the observed data; the wavelengths from our calculations tend to be smaller than the observed values by about 0.3 mÅ, while those from Chen and Nilsen tend to be 0.2 and 0.7 mÅ, respectively, larger than the observed data.

Figure 4(b) shows a synthetic FeXXV Kβ spectrum constructed from the present theoretical data. The additional parameters, i. e. the electron and ion temperatures, T_e and T_i , as well as the relative densities, N_{He} , N_{Li} and N_{Be} , of the heliumlike, lithiumlike, and berylliumlike charge states of iron, have been derived from the iron Kα spectrum which was observed with the vertical TFTR spectrometer. The synthetic spectrum shown in Fig. 4(b) was obtained using $T_e = 2.7$ keV, $T_i = 3.0$ keV, and with N_{He} , N_{Li} and N_{Be} in the ratio of 1.0 : 0.55 : 0.11.

IV Conclusions.

The Fe XXV Kβ spectrum has been obtained from TFTR discharges and found to be in good agreement with a synthetic spectrum constructed from new theoretical predictions. Since the iron Kβ spectrum was partially blended with the Ni XXVII Kα spectrum, it was necessary to enhance the intensity of the Kβ radiation by the injection of iron into

the plasma. A total of nine features in the Fe XXV $K\beta$ spectrum have been identified. The relative intensities depend on the electron temperature and charge state balance, which suggests their future use for plasma diagnostics.

Experimental wavelengths of the FeXXV $K\beta$ spectrum have been determined relative to the Ni XXVII $K\alpha$ resonance line. The satellite lines of the iron $K\beta$ spectrum do not interfere with the Ni XXVII $K\alpha$ resonance line. This is important for both TFTR and Tore Supra plasmas where iron and nickel are simultaneously present as impurities and where the Ni XXVII $K\alpha$ line is used for ion temperature measurements.

Acknowledgments. We deeply appreciate the fruitful discussions with Professor U. I. Safronova and would like to thank her for making her most recent wavelength calculations for the FeXXV $K\beta$ spectrum available to us. We also gratefully acknowledge the continuing support of D. Meade, R. J. Hawryluk and K. M. Young, the technical assistance of J. Gorman, R. Such and the TFTR operating crew, and software support by N. Schechtman. This work was supported by the U.S. Department of Energy, Contract No. DE-AC02-76-CHO-3073 and Contract No. W-7405-Eng-48.

References

- (1) H. Hsuan, M. Bitter, K. W. Hill, S. von Goeler, B. Grek, D. Johnson, L. C. Johnson, S. Sesnic, C. P. Bhalla, K. B. Karim, F. Bely-Dubau and P. Faucher, Phys. Rev. A 35, 4280(1987).
- (2) F. Bombarda, R. Giannella, E. Källne, G. J. Tallents, F. Bely-Dubau, P. Faucher, M. Cornille, J. Dubau and A. H. Gabriel, Phys. Rev. A 37, 504(1988); and H. Hsuan, M. Bitter, J. E. Rice, K. W. Hill, L. Johnson, S. L. Liew, S. D. Scott, and S. von Goeler, Rev. Sci. Instrum. 59, 2127(1988).
- (3) F. Bombarda, R. Giannella, E. Källne, G. J. Tallents, J. Quant. Spectrosc. Radiat. Transf. 41, 323(1989).
- (4) M. Bitter, H. Hsuan, V. Decaux, B. Grek, K. W. Hill, R. Hulse, L. A. Kruegel, D. Johnson, S. von Goeler and M. Zarnstoff, Phys. Rev. A 44, 1796(1991).
- (5) In 1987 the Inconel bellows cover plates were covered with graphite tiles, so that the concentration of nickel was very much reduced and iron became the dominant metal impurity in TFTR discharges. Line radiation from nickel could then no longer be used for ion temperature measurements and the spectrometer was realigned for the observation of the FeXXV $K\alpha$ line.
- (6) F. Bely-Dubau, A. H. Gabriel and S. Volonte, Mon. Not. R. Astr. Soc. 186, 405(1979); and M. Bitter, S. von Goeler, K. W. Hill, R. Horton, D. Johnson, W. Roney, N. Sauthoff, E. Silver and W. Stodiek, Phys. Rev. Lett. 47, 921(1981).
- (7) M. Klapisch, Comput. Phys. Commun. 2, 239(1971); and M. Klapisch, J. L. Schwob, B. S. Fraenkel, and J. Oreg, J. Opt. Soc. Am. 61, 148(1977).
- (8) A. Bar-Shalom, M. Klapisch, and J. Oreg, Phys. Rev. A 38, 1773(1988).
- (9) J. Oreg, A. Bar-Shalom, W. H. Goldstein, and M. Klapisch, Phys. Rev. A 44, 1750(1991).

- (10) M. Bitter, K. W. Hill, S. Cohen, S. von Goeler, H. Hsuan, L. C. Johnson, R. Raftopoulos, M. Reale, N. Schechtman, F. Spinos, J. Timberlake, N. Young and K. M. Young, Rev. Sci. Instrum. 57, 2145(1986).
- (11) K. W. Hill, M. Bitter, M. Tavernier, M. Diesso, S. von Goeler, G. Johnson, L. C. Johnson, N.. Sauthoff, N. Schechtman, S. Sesnic, F. Tenney and K. M. Young, Rev. Sci. Instrum. 56, 1165(1985).
- (12) R. A. Hulse, Nucl. Technol. Fusion 3, 259 (1983).
- (13) A. H. Gabriel, Mon. Not. R. Astr. Soc. 160, 79(1972).
- (14) U. I. Safronova, personal communication, 1992.
- (15) M. H. Chen, Atomic Data and Nuclear Data Tables 34, 301(1986).
- (16) J. Nilsen, Atomic Data and Nuclear Data Tables 37,191(1987)

Table I: Excitation rate coefficients, for direct electron impact excitation from the ground states of (a) FeXXV, FeXXI, and (b) FeXXIII at electron temperatures between 1.2 and 4.0 keV. Also listed are the autoionization probabilities A_a^s , the radiative transition probabilities A_r , and the branching ratios defined as $\beta_r = \frac{A_r^{sf}}{\sum_f A_r^{sf} + \sum_i A_a^s}$. The numbers in square brackets are powers of ten.

Table II: Intensity factors for the lithiumlike dielectronic satellite lines of the iron $K\beta$ lines; only lines with line factors larger than 1% of the strongest satellite are listed. Numbers in square brackets signify powers of 10.

Table III: Intensity factors for the berylliumlike satellites of the iron $K\beta$ lines; only lines with $F_2(s) \geq 2.0 \times 10^{12} s^{-1}$ are included. Numbers in square brackets signify powers of 10.

Table IV: Experimental wavelengths and theoretical predictions for the lines in the iron $K\beta$ spectrum; peaks are numbered as in Fig 6.

Table Ia

Charge State	Excited State	$\lambda_{T_{\text{keV}}^{\text{eor}}}$ (s ⁻¹)	A_a (s ⁻¹)	A_r (s ⁻¹)	β_r	Excitation Rate Coefficients for FeXXV and FeXXIV at different Electron Temperatures				
						1.5 keV	2.0 keV	2.5 keV	3.0 keV	3.5 keV
FeXXV	[1s3p _{3/2}] ₁	1.5732	0	1.3985[14]	0.940	5.974[-15]	2.077[-14]	4.370[-14]	7.161[-14]	1.018[-13]
	[1s3p _{1/2}] ₁	1.5751	0	1.0757[13]	0.546	1.807[-15]	5.635[-15]	1.080[-14]	1.633[-14]	2.163[-14]
	[1s2s3p _{3/2}] _{3/2}	1.5853	5.8017[12]	4.8657[13]	0.608	1.842[-15]	6.159[-15]	1.258[-14]	2.015[-14]	2.813[-14]
	[1s2s3p _{1/2}] _{1/2}	1.5864	5.0193[11]	6.1784[13]	0.813	1.060[-15]	3.596[-15]	7.436[-15]	1.293[-14]	1.693[-14]
	[1s2p _{1/2} 3s] _{3/2}	1.5878	1.0177[12]	1.0134[13]	0.252	3.078[-16]	1.054[-15]	2.194[-15]	3.571[-15]	5.051[-15]
FeXXIV	[1s2p _{1/2} 3s] _{1/2}	1.5886	2.9939[11]	4.1506[12]	0.309	6.324[-17]	2.128[-16]	4.373[-16]	7.039[-16]	9.866[-16]
	[1s2s3p _{3/2}] _{3/2}	1.5912	1.3235[12]	6.9929[13]	0.813	1.178[-15]	3.979[-15]	8.202[-15]	1.324[-14]	1.860[-14]
	[1s2s3p _{1/2}] _{1/2}	1.5913	1.7645[12]	6.3381[13]	0.738	2.302[-15]	7.728[-15]	1.586[-14]	2.549[-14]	3.572[-14]
	[1s2s3p _{1/2}] _{3/2}	1.5924	6.5669[12]	1.6307[13]	0.786	1.376[-15]	4.310[-15]	8.337[-15]	1.274[-14]	1.706[-14]
	[1s2s3p _{1/2}] _{1/2}	1.5926	1.2161[11]	3.4637[13]	0.776	5.423[-16]	1.631[-15]	3.035[-15]	4.469[-15]	5.779[-15]

Table Ib

Charge State	Excited State	$\lambda_{T_{\text{keV}}^{\text{eor}}}$ (Å)	A_a (s ⁻¹)	A_r (s ⁻¹)	β_r	Excitation Rate Coefficients for FeXXIII at different Electron Temperatures				
						1.2 keV	1.6 keV	2.0 keV	3.0 keV	4.0 keV
FeXXIII	[1s2s ² 3p _{3/2}] ₁	1.6053	6.9341[11]	1.2430[14]	0.852	1.67441[-15]	7.801[-15]	1.956[-14]	6.635[-14]	1.220[-13]

Table II

Transition	λ_{Theor} (Å)	g_s	A_a (s^{-1})	β_r	$F_2(s)$ (s^{-1})
$[1s2s3p_{3/2}]_{3/2} - 1s^22s$	1.5853	4	5.80[12]	0.608	1.41[13]
$[1s2p_{3/2}3p_{3/2}]_{3/2} - 1s^22p_{3/2}$	1.5900	4	1.66[13]	0.100	6.64[12]
$[1s2p_{3/2}3p_{3/2}]_{5/2} - 1s^22p_{3/2}$	1.5907	6	3.30[13]	0.048	9.50[12]
$[1s2p_{3/2}3p_{1/2}]_{3/2} - 1s^22p_{3/2}$	1.5912	4	3.96[13]	0.050	7.92[12]
$[1s2s3p_{3/2}]_{1/2} - 1s^22s$	1.5912	2	1.32[12]	0.813	2.15[12]
$[1s2s3p_{3/2}]_{3/2} - 1s^22s$	1.5913	4	1.76[12]	0.738	5.20[12]
$[1s2p_{3/2}3p_{1/2}]_{3/2} - 1s^22p_{1/2}$	1.5922	4	3.38[12]	0.157	2.12[12]
$[1s2s3p_{1/2}]_{3/2} - 1s^22s$	1.5924	4	6.70[11]	0.786	2.10[12]
$[1s2p_{1/2}3p_{3/2}]_{3/2} - 1s^22p_{3/2}$	1.5931	6	3.12[13]	0.663	1.24[14]
$[1s2p_{1/2}3p_{3/2}]_{3/2} - 1s^22p_{1/2}$	1.5933	4	9.02[12]	0.523	1.88[13]
$[1s2p_{3/2}3p_{3/2}]_{3/2} - 1s^22p_{3/2}$	1.5946	4	4.78[12]	0.419	8.01[12]
$[1s2p_{3/2}3p_{1/2}]_{5/2} - 1s^22p_{3/2}$	1.5949	6	6.49[12]	0.305	1.19[13]
$[1s2p_{3/2}3p_{1/2}]_{3/2} - 1s^22p_{3/2}$	1.5954	4	3.38[12]	0.362	4.89[12]
$[1s2s3d_{3/2}]_{3/2} - 1s^22p_{1/2}$	1.5980	4	4.50[12]	0.230	4.14[12]
$[1s2s3d_{5/2}]_{5/2} - 1s^22p_{3/2}$	1.6013	6	3.76[12]	0.122	2.75[12]
$[1s2s3s_{1/2}]_{1/2} - 1s^22p_{1/2}$	1.6034	2	4.33[13]	0.052	4.50[12]
$[1s2s3s_{1/2}]_{1/2} - 1s^22p_{3/2}$	1.6067	2	4.33[13]	0.054	4.68[12]

Table III

Transitions	λ_{Theor} (Å)	g_s	$A_e^{s \rightarrow 0}$ (s^{-1})	β_r	$F_2(s)$ (s^{-1})
$[1s2s2p_{3/2}3p_{3/2}]_3 - [1s^22s2p_{3/2}]_2$	1.6023	7	1.16[13]	0.112	4.55[12]
$[1s2s2p_{3/2}3p_{3/2}]_2 - [1s^22s2p_{3/2}]_2$	1.6027	5	2.11[13]	0.054	2.85[12]
$[1s2s2p_{1/2}3p_{3/2}]_2 - [1s^22s2p_{1/2}]_1$	1.6047	5	1.09[13]	0.107	2.92[12]
$[1s2s2p_{3/2}3p_{3/2}]_3 - [1s^22s2p_{3/2}]_2$	1.6052	7	1.84[13]	0.081	5.22[12]
$[1s2s2p_{3/2}3p_{3/2}]_0 - [1s^22s2p_{3/2}]_1$	1.6057	1	1.43[13]	0.367	2.62[12]
$[1s2s2p_{3/2}3p_{3/2}]_2 - [1s^22s2p_{3/2}]_1$	1.6071	5	1.56[13]	0.281	1.10[13]
$[1s2s2p_{3/2}3p_{3/2}]_2 - [1s^22s2p_{3/2}]_1$	1.6082	5	2.99[13]	0.326	2.44[13]
$[1s2s2p_{3/2}3p_{3/2}]_2 - [1s^22s2p_{1/2}]_1$	1.6095	5	1.81[13]	0.153	6.92[12]
$[1s2s2p_{3/2}3p_{3/2}]_1 - [1s^22s2p_{3/2}]_2$	1.6101	3	8.76[12]	0.503	6.61[12]
$[1s2s2p_{1/2}3p_{3/2}]_1 - [1s^22s2p_{1/2}]_0$	1.6102	3	2.53[13]	0.148	5.62[12]
$[1s2s2p_{3/2}3p_{1/2}]_2 - [1s^22s2p_{1/2}]_1$	1.6106	5	3.38[12]	0.271	2.29[12]
$[1s2s2p_{1/2}3p_{3/2}]_3 - [1s^22s2p_{3/2}]_2$	1.6108	7	3.08[13]	0.614	6.62[13]
$[1s2s2p_{1/2}3p_{3/2}]_1 - [1s^22s2p_{1/2}]_1$	1.6110	3	2.53[13]	0.425	1.61[13]
$[1s2s2p_{1/2}3p_{3/2}]_2 - [1s^22s2p_{1/2}]_1$	1.6115	5	5.72[12]	0.361	5.16[12]
$[1s2s^23d_{5/2}]_3 - [1s^22s2p_{1/2}]_1$	1.6116	5	4.22[12]	0.445	4.69[12]
$[1s2s2p_{3/2}3p_{3/2}]_2 - [1s^22s2p_{3/2}]_2$	1.6119	5	1.81[13]	0.459	2.08[13]
$[1s2s2p_{1/2}3p_{3/2}]_1 - [1s^22s2p_{1/2}]_0$	1.6120	3	4.04[12]	0.602	3.65[12]
$[1s2s2p_{3/2}3p_{1/2}]_3 - [1s^22s2p_{3/2}]_2$	1.6125	7	5.83[12]	0.442	9.02[12]
$[1s2s2p_{3/2}3p_{1/2}]_2 - [1s^22s2p_{3/2}]_2$	1.6130	5	3.38[12]	0.311	2.63[12]
$[1s2s2p_{3/2}3p_{1/2}]_2 - [1s^22s2p_{3/2}]_1$	1.6130	5	2.73[13]	0.039	2.66[12]

Table IV

Peak	Transitions	Channel Number	λ_{Expt} (Å)	λ_{Theor} (Å)	λ_a (Å)	λ_b (Å)	λ_c (Å)
1	1s3p ¹ P ₁ -1s2 ¹ S ₀	138.07	1.5734	1.5732	1.5732	1.57396	
2	1s3p ³ P ₁ -1s2 ¹ S ₀	146.02	1.5751	1.5751	1.5751	1.57584	
3	1s2s[¹ S]3p ² P _{3/2} -1s2s ² S _{1/2}	194.32	1.5854	1.5853	1.5855	1.58593	1.5858
4	1s2s[¹ S]3p ² P _{1/2} -1s2s ² S _{1/2}	200.0	1.5867	1.5864	1.5866		1.5869
w*	1s2p ¹ P ₁ -1s ² ¹ S ₀	208.14	1.5884		1.5884		
5	1s2s[³ S]3p ² P _{1/2} -1s2s ² S _{3/2}	222.16	1.5914	1.5912	1.5915	1.59190	1.5917
	1s2s[³ S]3p ² P _{1/2} -1s2s ² S _{1/2}			1.5913	1.5916	1.5919	1.5918
6	1s2p[³ P]3p ² D _{5/2} -1s2p ² P _{3/2}	230.68	1.5932	1.5931	1.5934	1.59387	1.5935
7	1s2p[³ P]3p ² P _{5/2} -1s2p ² P _{3/2}	239.11	1.5950	1.5949	1.5951	1.59567	1.5954
z*	1s2s ³ S ₁ -1s ² ¹ S ₀	279.80	1.6036		1.6037		
8	1s2s ² 3p ¹ P ₁ -1s2s ² 1S ₀	289.42	1.6056	1.6053	1.6055		
9	1s2s2p3p ³ P ₁ -1s2s2p ³ P ₂	315.02	1.6109	1.6108			

*w and z lines of NiXXVII

 λ_{Theor} from present calculations λ_a from Ref. 14 λ_b from Ref. 16 λ_c from Ref. 15

Figure Captions

Figure 1: Electron temperature and electron density as functions of time for TFTR shot 28931. The data were obtained from measurements of the electron cyclotron emission and laser interferometry. Iron was injected at 2.7 s by a laser ablation technique.

Figure 2: Time history of (a) the Fe $K\beta$ and (b) the Fe $K\alpha$ line intensities.

Figure 3: Satellite spectra of Ni XXVII $K\alpha$ and Fe XXV $K\beta$ (a) before and (b) after iron injection. The data have been accumulated from five nearly identical TFTR ohmic shots during the periods 2.5 - 2.7 s for Fig. 3 (a) and 2.75 - 2.88 s for Fig. 3 (b). The features of the Ni XXVII $K\alpha$ satellite spectrum have been identified using Gabriel's notation.¹³ The Fe XXV $K\beta$ features are significantly enhanced in the spectrum (b) which was recorded after iron injection.

Figure 4: (a) Experimental iron $K\beta$ spectrum obtained by subtracting a normalized spectrum of Fig. 3 (a) from the spectrum shown in Fig. 3 (b). (b) Synthetic iron $K\beta$ spectra constructed from present theoretical data for the measured electron and ion temperatures of $T_e = 2.7$ keV and $T_i = 2.35$ keV. The shaded area represents the satellite contributions from collisional inner-shell excitation. The solid line represents the entire spectrum including contributions from dielectronic recombination.

Figure 1

Figure 2

Figure 3

PPPL#92X0232

Figure 4

Figure 4

EXTERNAL DISTRIBUTION IN ADDITION TO UC-420

Dr. F. Paoloni, Univ. of Wollongong, AUSTRALIA
 Prof. M.H. Brennan, Univ. of Sydney, AUSTRALIA
 Plasma Research Lab., Australian Nat. Univ., AUSTRALIA
 Prof. I.R. Jones, Flinders Univ, AUSTRALIA
 Prof. F. Cap, Inst. for Theoretical Physics, AUSTRIA
 Prof. M. Heindler, Institut für Theoretische Physik, AUSTRIA
 Prof. M. Goossens, Astronomisch Instituut, BELGIUM
 Ecole Royale Militaire, Lab. de Phy. Plasmas, BELGIUM
 Commission-European, DG. XII-Fusion Prog., BELGIUM
 Prof. R. Boudiqué, Rijksuniversiteit Gent, BELGIUM
 Dr. P.H. Sakanaka, Instituto Fisica, BRAZIL
 Instituto Nacional De Pesquisas Especiais-INPE, BRAZIL
 Documents Office, Atomic Energy of Canada Ltd., CANADA
 Dr. M.P. Bachynski, MPB Technologies, Inc., CANADA
 Dr. H.M. Skarsgard, Univ. of Saskatchewan, CANADA
 Prof. J. Teichmann, Univ. of Montreal, CANADA
 Prof. S.R. Sreenivasan, Univ. of Calgary, CANADA
 Prof. T.W. Johnston, INRS-Energie, CANADA
 Dr. R. Bolton, Centre canadien de fusion magnétique, CANADA
 Dr. C.R. James,, Univ. of Alberta, CANADA
 Dr. P. Lukáč, Komenského Univerzita, CZECHO-SLOVAKIA
 The Librarian, Culham Laboratory, ENGLAND
 Library, R61, Rutherford Appleton Laboratory, ENGLAND
 Mrs. S.A. Hutchinson, JET Library, ENGLAND
 Dr. S.C. Sharma, Univ. of South Pacific, FIJI ISLANDS
 P. Mähönen, Univ. of Helsinki, FINLAND
 Prof. M.N. Bussac, Ecole Polytechnique,, FRANCE
 C. Mouttet, Lab. de Physique des Milieux Ionisés, FRANCE
 J. Radet, CEN/CADARACHE - Bat 506, FRANCE
 Prof. E. Economou, Univ. of Crete, GREECE
 Ms. C. Rinni, Univ. of Ioannina, GREECE
 Dr. T. Mui, Academy Bibliographic Ser., HONG KONG
 Preprint Library, Hungarian Academy of Sci., HUNGARY
 Dr. B. DasGupta, Saha Inst. of Nuclear Physics, INDIA
 Dr. P. Kaw, Inst. for Plasma Research, INDIA
 Dr. P. Rosenau, Israel Inst. of Technology, ISRAEL
 Librarian, International Center for Theo Physics, ITALY
 Miss C. De Palo, Associazione EURATOM-ENEA, ITALY
 Dr. G. Grosso, Istituto di Fisica del Plasma, ITALY
 Prof. G. Rostangni, Istituto Gas Ionizzati Del Cnr, ITALY
 Dr. H. Yamato, Toshiba Res. & Devel. Center, JAPAN
 Prof. I. Kawakami, Hiroshima Univ., JAPAN
 Prof. K. Nishikawa, Hiroshima Univ., JAPAN
 Director, Japan Atomic Energy Research Inst., JAPAN
 Prof. S. Itoh, Kyushu Univ., JAPAN
 Research Info. Ctr., National Inst. for Fusion Science, JAPAN
 Prof. S. Tanaka, Kyoto Univ., JAPAN
 Library, Kyoto Univ., JAPAN
 Prof. N. Inoue, Univ. of Tokyo, JAPAN
 Secretary, Plasma Section, Electrotechnical Lab., JAPAN
 S. Mori, Technical Advisor, JAERI, JAPAN
 Dr. O. Mitarai, Kumamoto Inst. of Technology, JAPAN
 J. Hyeon-Sook, Korea Atomic Energy Research Inst, KOREA
 D.I. Choi, The Korea Adv. Inst. of Sci. & Tech., KOREA
 Prof. B.S. Liley, Univ. of Waikato, NEW ZEALAND
 Inst of Physics, Chinese Acad Sci PEOPLE'S REP. OF CHINA
 Library, Inst. of Plasma Physics, PEOPLE'S REP. OF CHINA
 Tsinghua Univ. Library, PEOPLE'S REPUBLIC OF CHINA
 Z. Li, S.W. Inst. Physics, PEOPLE'S REPUBLIC OF CHINA
 Prof. J.A.C. Cabral, Instituto Superior Tecnico, PORTUGAL
 Dr. O. Petrus, AL I CUZA Univ., ROMANIA
 Dr. J. de Villiers, Fusion Studies, AEC, S. AFRICA
 Prof. M.A. Hellberg, Univ. of Natal, S. AFRICA
 Prof. D.E. Kim, Pohang Inst. of Sci. & Tech., SO. KOREA
 Prof. C.I.E.M.A.T, Fusion Division Library, SPAIN
 Dr. L. Stenflo, Univ. of UMEA, SWEDEN
 Library, Royal Inst. of Technology, SWEDEN
 Prof. H. Wilhelmson, Chalmers Univ. of Tech., SWEDEN
 Centre Phys. Des Plasmas, Ecole Polytech, SWITZERLAND
 Bibliotheek, Inst. Voor Plasma-Fysica, THE NETHERLANDS
 Asst. Prof. Dr. S. Cakir, Middle East Tech. Univ., TURKEY
 Dr. V.A. Glukhikh, Sci. Res. Inst. Electrophys. Apparatus, USSR
 Dr. D.D. Ryutov, Siberian Branch of Academy of Sci., USSR
 Dr. G.A. Eliseev, I.V. Kurchatov Inst., USSR
 Librarian, The Ukr.SSR Academy of Sciences, USSR
 Dr. L.M. Kovrizhnykh, Inst. of General Physics, USSR
 Kernforschungsanlage GmbH, Zentralbibliothek, W. GERMANY
 Bibliothek, Inst. Für Plasmaforschung, W. GERMANY
 Prof. K. Schindler, Ruhr-Universität Bochum, W. GERMANY
 Dr. F. Wagner, (ASDEX), Max-Planck-Institut, W. GERMANY
 Librarian, Max-Planck-Institut, W. GERMANY
 Prof. R.K. Janev, Inst. of Physics, YUGOSLAVIA