NOTICE

**CERTAIN DATA
CONTAINED IN THIS
DOCUMENT MAY BE
DIFFICULT TO READ
IN MICROFICHE
PRODUCTS.**

DOE/ER/53174^{T1}XTHEORETICAL STUDIES ON PLASMA
HEATING AND CONFINEMENT

DOE/ER/53174---T1

DE93 007617

~~Final~~
Progress Report

for Period May 4, 1984 - May 13, 1988

R.N. Sudan

Cornell University
Ithaca, New York 14853

January 1993

Prepared for

THE U.S. DEPARTMENT OF ENERGY
AGREEMENT NO. DE-FG02-84ER53174

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED ^{lp}

THEORETICAL STUDIES ON PLASMA HEATING AND CONFINEMENT

DOE Grant NO: DE-FG02-84ER53174

Period: May 4, 1984 - May 13, 1988

Abstract

Three principal topics are covered in this final report:

1. Stabilization of low frequency modes of an axisymmetric compact torus plasma confinement system, such as, spheromaks and FRC's, by a population of large orbit axis encircling energetic ions. Employing an extension of the 'energy principle' which utilizes a Vlasov description for the energetic ion component, it has been demonstrated that short wavelength MHD type modes are stabilized while the long wavelength tilt and precessional modes are marginally stable. The deformation of the equilibrium configuration by the energetic ions results in the stabilization of the tilt mode for spheromaks.
2. Formation of Ion Rings and their coalescence with spheromaks. A two dimensional electromagnetic PIC codes [CIDER] has been developed for the study of ion ring formation and its propagation, deformation and slowing down in a cold plasma. It has been shown that a ring moving at a speed less than the Alfvén velocity can merge with a stationary spheromak.
3. Anomalous transport from drift waves in a Tokamak. The Direct Interaction Approximation is used to obtain incremental transport coefficients for particles and heat for drift waves in a Tokamak. It is shown that the transport matrix does not obey Onsager's principle.

I. SUMMARY OF WORK ACCOMPLISHED (1984 - 1988)

A. Stabilization of Low Frequency Modes by Large Orbit Ions

1. **Stabilization of the Spheromak Tilt Instability** by *C. Litwin, R.N. Sudan and A.D. Turnbull*. A stability criterion for the tilt mode of a spheromak-ion ring hybrid configuration has been developed for the case where the ring current is small compared to the spheromak azimuthal current. It is shown that the stability is related to the distortion of the spheromak separatrix.
2. **Stability of Compact-Torus Ion-Ring Hybrid Systems to Magnetohydrodynamic Tilt and Shift Modes** by *A.D. Turnbull and R.N. Sudan*. The low-frequency stability of a compact-torus ion-ring hybrid configuration, in which part of the toroidal current is carried by a ring of high-energy ions, is studied. In particular, the magnetohydrodynamic (MHD) tilt and radial shift instabilities in spheromak-like hybrid systems are considered. The ion ring has a stabilizing effect on the tilt mode, through its modification of the magnetic geometry. For the shift mode, the stabilization is dynamic for moderate elongation, it is found that a high-energy ion component substantially enhances stability against the MHD shift mode.
3. **Stability of Field-Reversed Ion Rings** by *Alex Friedman, R.N. Sudan and Jacques Denavit*. Studies of the low-frequency stability of field-reversed ion rings in a dense uniform plasma background have been carried out using a linearized 3-D hybrid simulation model. For a moderately thick ring with aspect ratio on the order of 4:1, the azimuthal mode number $l = 1$ radial mode (magnetohydrodynamic precession) is observed to be stable, because of the favorable field gradient arising from image currents in the outer wall. The $l = 1$ axial (tilt) mode is unstable for the parameters chosen, as are the $l = 2$ radial and axial (kink) modes, and the $l = 3$ radial mode. All other modes are observed to be either stable or very weakly unstable; numerical difficulties arising from the local instability of individual orbits in the equilibrium field make it difficult to examine modes with small growth rates. The structures of the various unstable modes range from near rigid to quite complicated in displacement pattern. Some preliminary evidence for the existence of betatron resonance-driven instabilities is noted.
4. **Effect of a Strong-Current Ion Ring on Spheromak Stability** by *C. Litwin and R.N. Sudan*. The stability of a spheromak with an energetic ion ring, carrying a current comparable to the plasma current, to the tilt mode is considered. For small departures from sphericity a perturbative approach is applied to an appropriate energy

principle in order to calculate the lowest nontrivial kinetic contribution of the ion ring. An analytic stability criterion is obtained. It is seen that the prolate configuration becomes more stable while the oblate one is less stable than in the absence of the ring. The prolomak becomes stable when the ring kinetic energy exceeds the magnetic energy within the separatrix.

5. **Kinetic Stabilization of an Axisymmetric Mirror by an Energetic Ion Component** by *J.F. Krull, C.E. Seyler, and R.N. Sudan*. We examine the stability of a periodic, axisymmetric mirror with an energetic ion component to finite azimuthal mode number interchange and ballooning modes. Using the beam-plasma model, we arrive at a quadratic form of the dispersion functional, involving phase space autocorrelation functions over energetic orbits. The eigenfunction is represented in terms of a complete set of global basis functions, providing a necessary and sufficient condition for stability. Preliminary numerical results are obtained for beam ions described by a rigid rotor distribution function and by a simple Maxwellian. For the Maxwellian, the energetic ions form a "bump on tail" and optimization of the temperature can reduce MHD growth rates by as much as 85% for fixed density.

B. Formation of Ion Rings and Merging with Spheromaks

1. **Particle Simulations of Merging Ion Rings in Resistive Plasmas** by *P.M. Lyster and R.N. Sudan*. A numerical study, employing the particle code RINGA [Plasma Phys. 23, 521 (1981)], of two ion rings that merge with each other due to their mutual magnetic attraction is presented. It is shown that an ion ring with closed magnetic field lines (field reversed) can be formed by merging two weaker non-field-reversed rings. A rough calculation shows that there is an optimal plasma conductivity $\sigma \approx c^2/4\pi a R \Omega \zeta^{1/2}$ (a and R are the respective axial width and major radius of the ring, Ω is the ion cyclotron frequency in the external magnetic field, and ζ is the field reversal factor of the weaker ring) for which most of the energy of mutual magnetic attraction between the rings is dissipated during the process of merging.
2. **The Dynamics of Ion Rings in Highly Conductive Plasmas** by *P.M. Lyster and R.N. Sudan*. A numerical study employing a hybrid computer code (particle ions/fluid electrons) is made of the dynamics of translating ion rings in a highly conductive plasma. The translational kinetic energy of such rings is lost through the generation of Alfvén waves. When the initial axial velocity of the ring is greater than the local Alfvén speed in the plasma, a typical slowing down length is $z_A = (a_r/4\zeta)^{-4} v_{r0}/(v_\theta^2 v_A^2)$, where a_r is the ring radial width, $\zeta = |\delta B/B_0|$, δB is the ring

self-magnetic-field strength on axis, B_o is the strength of the axial magnetic field, $\bar{v}_{x0}$ is the initial translational velocity of the ring, v_θ is the average toroidal ring particle velocity, $v_A = B_o/\sqrt{4\pi\rho}$ is the Alfvén velocity, and ρ is the plasma mass density. Also discussed is the merging of an ion ring with a spheromak.

C. Drift Wave Turbulence and Anomalous Transport

1. **Anomalous Electron Heat Transport Driven by Low-Frequency Electromagnetic Turbulence** by *A.A. Thoul, P.L. Similon, and R.N. Sudan*. We consider the anomalous electron and heat transport in a tokamak plasma. The electrons are described by the nonlinear drift-kinetic equation. We analyze transport through the averaged response function in the presence of drift-Alfvén wave turbulence. In contrast to recent findings by Terry, Diamond, and Hahn, we conclude that magnetic fluctuations lead to a substantial transport of both parallel and perpendicular energies. The latter, previously neglected, is found to be significant and of the order of the test-particle diffusion.
2. **Study of Turbulent Transport Due to Drift-Alfvén Fluctuations** by *A.A. Thoul, P.L. Similon and R.N. Sudan*. Turbulent electron heat transport is a major loss mechanism in tokamaks. The effect of self-consistent magnetic fluttering has been recently analysed using clump theory. An alternate approach has also been presented, which includes perpendicular energy transport, and which leads to a revision of the previous conclusions. It used the renormalized response function to determine the incremental fluxes due to perturbation of the mean density and temperature profiles. The method is illustrated with a simple model with turbulent collision operator similar to the Balescu-Lenard operator. It is shown explicitly how drag and diffusion terms appear in this formalism. In the context of drift Alfvén turbulence, the perturbations of thermodynamic fluxes across magnetic surfaces are evaluated for various sources, namely, in response to external sources of a) density, b) parallel temperature, c) perpendicular temperature, and d) parallel current, the fluxes of a) particles, b) electron parallel energy, c) electron perpendicular energy, and d) parallel current are computed. This transport matrix is analysed, and it is found that Onsager symmetry relations are violated for turbulent transport.
3. **Almost two-dimensional strong turbulence in a magnetized plasma** by *M.N. Rosenbluth and R.N. Sudan*. From the direct interaction approximation (DIA) equations for strong turbulence it is shown that low-frequency fluctuations constituting two-dimensional strong turbulence in a magnetized plasma scatter off each other to de-

Research held by the International Atomic Energy Agency in London, 12 - 19 September 1984., Vol. 3, Vienna, 1985.

R.N. Sudan and P.M. Lyster, "Injection and Trapping of Energetic Ions in Compact Torus Configurations," *Comments Plasma Phys. Controlled Fusion*, 9 (1) 23, 1984.

1984

C. Litwin, R.N. Sudan, and A.D. Turnbull, "Stabilization of the Spheromak Tilt Instability," *Phys. Fluids* 27 (12), 1984.

III. PhD STUDENTS SUPPORTED BY THIS GRANT

Peter M. Lyster, "Particle Simulations of Ion Rings for Magnetic Fusion," PhD completed January 1987.

Jonathan F. Krall, "Stability of an Axisymmetric Mirror with an Energetic Ion Component," PhD completed January 1987.

Anne A. Thoul, "Anomalous Perturbative Transport in Tokamaks Due to Drift Waves Turbulence," PhD completed May 1992.

IV. OTHER PERSONNEL SUPPORTED BY THIS GRANT

Senior Research Associate: C. Litwin

Post-Docs: C.G. Schultz
J.M. Albert

END

DATE
FILMED

3 / 23 / 93