ANNUAL REPORT ON FUSION PLASMA THEORY GRANT

TASK III: Auxiliary Radiofrequency Heating of Tokamaks Covering the period November 16, 1991 to November 15, 1992

J.E. Scharer, Principal Investigator; N.T. Lam, Scientist;
M. Bettenhausen and K. Sund, Graduate Students

Abstract

The research performed under this grant during the past year has been concentrated on the following several key tokamak ICRF (Ion Cyclotron Range of Frequencies) coupling, heating and current drive issues: Efficient coupling during the L- to H- mode transition by analysis and computer simulation of ICRF antennas; analysis of ICRF cavity-backed coil antenna coupling to plasma edge profiles including fast and ion Bernstein wave coupling for heating and current drive; benchmarking the codes to compare with current JET, D-IIID and ASDEX experimental results and predictions for advanced tokamaks such as BPX and SSAT (Steady-State Advanced Tokamak); ICRF full-wave field solutions, power conservation, heating analyses and minority ion current drive; and the effects of fusion alpha particle or ion tail populations on the ICRF absorption. Research progress, publications, and conference and workshop presentations are summarized in this report.

I. Introduction

This annual report on Task III-Auxiliary Radiofrequency Heating in Tokamaks of the Fusion Plasma Theory DOE Grant Number DE-FG02-86ER53218 at the University of Wisconsin covers the research carried out during the current grant period (November 16, 1991 to November 15, 1992). This is the first year of a third 3 year grant which was issued under the same grant number. These grants evolved

MASTER

out of the previous DOE Contract Number DE-AC02-80ER53104 that supported research in this area at the University of Wisconsin-Madison from July, 15, 1980 through December 31, 1985.

The Task III research progress described in this report emphasizes ICRF (Ion Cyclotron Range of Frequencies) antenna coupling, plasma heating and current drive and is relevant to tokamak plasma confinement issues. These areas of research are relevant to theoretical interpretations of JET and TFTR experimental observations, Burning Plasma Experiment (BPX) and Toroidal Plasma Experiment (TPX/Steady State Advanced Tokamak) physics analysis and advanced tokamak design concepts (funded by Princeton University), and future radiofrequency roles in tokamaks including ITER. We are currently collaborating with Professor Miklos Porkolab and the Princeton TPX/SSAT Physics group to define ICRF roles including appropriate antennas to provide heating and current drive. Our group has maintained a continuous collaboration with Dr. Jacquinot and the JET ICRF group since the summer of '83 and annual visits in both directions and regular phone and FAX contacts have served to stimulate ICRF research activity and the exchange of ideas.

Our research program emphasizes the efficient ICRF antenna power coupling to plasmas during the L- to H-mode transition. It also addresses an improved understanding of interactions between plasma edge conditions and ICRF wave coupling which are important to future fusion experiments such as TPX and ITER. The role of ICRF in producing near steady-state operation of future tokamaks via lower hybrid/fast wave current drive synergistic effects with high efficiencies is an important issue for future tokamaks. The role of minority ion current drive which can be accomplished by ICRF at high densities is an important confinement issue which can provide high performance tokamak reactors in the future. Finally, the influence of ICRF absorption by alpha particles on the fusion breakeven, alpha-particle excited TAE mode influence on confinement, ash removal and ignition heating scenarios are important issues. Our research program, within the limitation of budget constraints, is focussed on these physics issues which will provide an attractive fusion power source during the next century.

The research progress during the past year is summarized in the following sections: II. ICRF Antenna Modelling and Wave Coupling to the Plasma; III. Wave Absorption with Alpha Particles and Current Drive Synergistic Effects; and IV. Fast Ion Interaction with ICRF and Minority Ion Current Drive. The progress report concludes with lists of the publications, conferences, reports and workshops in section V and the personnel and their degree of involvement in Appendix A.

II. ICRF Antenna Modelling and Wave Coupling to the Plasma [1,2,V.A.1,V.A.2,V.B.1,V.C.1,V.C.2,V.C.3,V.D.1,V.D.2,V.E.1]

A paper [1] has been published detailing our calculations of antenna loading for BPX using two computer codes: ANTIMP and FELICE. ANTIMP is a code that we have developed which includes antenna coil cavity effects in evaluating coupling to L- and H-mode plasma edge profiles. We have modified Brambilla's FELICE code considerably to include multiple coaxial feeders and multi-element arrays. We have also restructured the code to more readily implement future development of the code. Since much of the attention of the RF community has now shifted towards current drive for SSAT, we have modified the FELICE code to evaluate the loading resistance of a multi-element antenna array.

We have calculated the loading resistances of various antenna structures for the SSAT base line parameters (major radius = 2.25 m, minor radius = 0.5 m, central B = 3.35 T, central density = $8 \times 10^{13} / \text{cm}^3$). Calculations have been carried out for 4 and 8 strap antennas. For an inter-coil phasing of $\pi/2$ (suitable for current drive), we obtain a loading of about 40 ohms/coil for a 4 strap module without the influence of coaxial feeders to the antenna. For an 8-coil module with $\pi/2$ phase between adjacent coils, the radiation resistance/coil drops to 26 ohms. When the effects of two coaxial feeders/coil are included, this is reduced to 20 ohms/coil. This is adequate to handle 2 MW/coil or 15 MW current drive physics scenarios. Further, the power spectrum of the antenna (including plasma effects) peaks in the proper parallel wave number region ($v_{\phi z} = 1 - 3 v_{the}$) for efficient electron current drive. These preliminary results suggest that a fast wave current drive system is an attractive RF generator power at $f = 80$ MHz. We are currently evaluating the use of 40 MHz and 37.5 MHz frequency scenarios.

Brambilla [2] used an approximate method for calculating the effect of angles between the orientation of the Faraday shield, the antenna current straps and the equilibrium magnetic field on ICRF coupling. Brambilla's method may provide a reasonable estimate of the effect on the radiation resistance but likely provides poor results for the coupled power spectrum. This is because his method assumes that the Fourier decomposition is actually performed in the toroidal direction so there is an angle between the direction of the Fourier decomposition and the magnetic field due the poloidal magnetic field component (which is the reason the angles are of interest). To account for this angle, a coordinate rotation should be used prior to the finite element solution of the differential equations for the wave electric fields in the plasma. Appropriate changes to the theoretical model and the computer code are in progress.

As noted above, the coupling code has been modified to calculate the radiation resistance and coupled power spectrum for the multiple-coil antenna arrays which are of particular interest for ICRF current drive. These modifications do not account for the effects of the radial feeders which deliver the current to the straps. Addition of the feeder effects for multiple-coil antenna arrays with various feeder configurations, to the model and the computer code, is nearing completion.

Analytic work has begun on developing the model to support the addition of a 3-D recessed cavity and/or septa to the vacuum model currently used in the ICRF coupling code. These two antenna features are believed to have important effects on the coupled power spectrum and, therefore, on ICRF current drive applications and are not treated simultaneously by existing models.

III. Wave Absorption with Alpha Particles and Current Drive Synergistic Effects [3,4,V.A.3,V.B.2,V.C.3,V.D.2]

Using XWAVE (a code which implements our derivation of power conservation) [3,4], we have pointed out the possible effects of the presence of a two-ion hybrid resonance on the absorption of fast waves for a (D)T plasma with a small concentration of α particles. Specifically, we have considered an intermediate BPX plasma with the following parameters: $n_e = 1.9 \times 10^{14} \text{ cm}^{-3}$, 1:2 D:T ratio, $T_D = T_T = 10 \text{ keV}$, $n_\alpha = 0.5\% n_e$. With the heating frequency set at the fundamental D resonance on axis, a two-ion (D-T) hybrid resonance exists 30 cm towards the high field from the center. We find that the presence of this hybrid resonance can give rise to a significant enhancement of the α absorption and a large reflection of the fast wave. For the above BPX parameters, the α particles absorb up to 23% of incident fast wave power, as compared with 20% for the D particles which absorb near the center. The reflection coefficient of the fast wave also amounts to 47%. It should be pointed out that our formalism is based on a gyroradius expansion and could overestimate the enhancement in the local α particle absorbed particle density. However, this effect should be of significance for future D-T operation including that of ITER.

Recently, a synergistic effect between lower hybrid and fast wave heating has been observed on JET [5]. Under certain conditions (monopole phasing, density below a critical density, etc.), the fast electrons created by lower hybrid waves can absorb additional power from the fast wave. In this way, significant increase of the current drive efficiency has been obtained. To explain this effect, the JET team claims that 10% of the ICRF power is absorbed synergistically when a 1% fast electron (200 keV) population is present. Using XWAVE, we have simulated the presence of such a hot electron population for a (H)-D JET-like plasma. For low parallel wave numbers (corresponding to monopole phasing) and fundamental hydrogen heating, we found that the the minority ions (H) absorb up to 60% of the fast wave power, while only up to 2% is absorbed by the fast electrons. It should be pointed out that XWAVE assumes an uniform density profile near the tokamak center, and Jacquinet has suggested that the spatial inhomogeneity in the fast electrons density profile may play a role in the synergistic effects. Further, the present version of XWAVE assumes isotropic Maxwellian temperature profiles for all species. At the present time, it is fair to say that the explanation of current drive enhancement by synergy between lower hybrid and fast waves remains a challenging theoretical problem. We are continuing to examine this effect in JET and will examine its prospects and scaling for future machines such as SSAT and ITER.

IV. ICRF Analysis for Large Gyroradius and Minority Ion Current Drive Effects [3,V.A.3,V.B.2,V.C.3]

We are currently evaluating ICRF heating scenarios in cases where small concentrations of alpha particles or tail ion species are present utilizing absorption equivalent Maxwellians and the XWAVE code. This is particularly important for fundamental and second harmonic deuterium and heating scenarios. We are also developing a formalism that includes the effects of parallel and perpendicular magnetic field gradients to more rigorously analyze the fast particle - ICRF interaction.

The works of Koch and Van Eester [6] and Sauter and Vaclavik [7A] have argued that the alpha particle slowing down distribution absorption of ICRF over the transverse spacial Doppler width can be equivalently modeled by Maxwellians in the range of 300-900 keV. We are currently evaluating the effects of alpha particle absorption competing with the fundamental deuterium resonance absorption for advanced tokamak parameters. We will compare our finite gyroradius absorption results [3] with a large $k_{\perp}\rho$ result in which ∇B effects ignored by the previous authors are included to evaluate the effects of alpha particle absorption for future burn and ignition tokamak ICRF heating scenarios. We are also working on an extension of Smithe's work [7B] to include a compact representation of the wave fields in order to evaluate the interaction with large gyroradius particles including parallel and perpendicular magnetic field gradients.

A proper analysis and code development to examine fast ion tail heating and alpha particle absorption by ICRF is an important future research area for the fusion program. The stabilization of sawtooth processes arising from the $m=1$ kink mode and transition to the monster sawtooth regime by fast ions produced by ICRF heating is an important stabilization mechanism for future fusion machines. The ability of ICRF to control the heating profile as well as the concentration and temperature of the fast ions leading to monster sawtooth periods of very long duration is of major importance for good confinement in future machines. Recent works by Porcelli, et al [8] have outlined the enhanced confinement window. Improved understanding and the development of wave heating codes which examine the interaction with the fast particle distributions is being examined by our group.

ICRF heating and modification of the alpha particle distribution could play an important role in stabilization of toroidal Alfvén eigenmodes (TAE) which are of concern for burning and ignition fusion machines such as ITER. Work by Chang et al. [9] has examined the role of ICRF heating in preferentially providing enhanced radial transport to remove lower energy alpha particle ash for a fusion burning tokamak device. The versatility of ICRF heating and fast particle interaction needs to be examined in more detail to yield the benefits that it can provide for a successful fusion program towards burning, demonstration, and commercial fusion reactors producing electric power.

We have been working to develop a self-consistent wave equation for fast waves and ion-Bernstein waves, valid near resonance and mode-conversion layers, that is not limited by a small gyroradius approximation, and that incorporates the effects of perpendicular and parallel magnetic field gradients, as well as allowing for non-Maxwellian equilibrium distribution functions. Fast ions whose gyroradii are comparable to or larger than the local perpendicular wavelength may arise either from heating or from fusion reactions (alphas). Additionally, in hot plasmas the wavelength of the ion-Bernstein mode becomes comparable even to thermal ion gyroradii as it propagates away from the mode conversion layer. To fully investigate the physics of such situations requires a formulation of the wave equation that does not depend on a small gyroradius expansion.

Attempts at such a formulation lead naturally to an integral equation approach. One such approach which we investigated was that of Sauter et al. [7A], who developed an equation allowing short wavelengths and gradients in density and temperature. However, we found that contrary to their claims, they did not incorporate the perpendicular magnetic field gradient properly. It is difficult to extend their work to correctly include such gradients.

We are currently pursuing an integral equation approach similar to Smithe's [7B] but including the generalizations mentioned above. Smithe's work included parallel gradient effects but was limited to small gyroradii and to Maxwellian distribution functions. He developed a constitutive relation for the current density at a particular location in terms of an integral over the Fourier representation of the wave electric field. When the Fourier representation is discretized the integral equation becomes a matrix equation for the values of E on a set of points across the solution interval. Inversion of the matrix yields the solution. Here the number of grid points is determined by the necessity to resolve the shortest wavelengths rather than the longer scale over which the wavelength changes. We are investigating alternative representations of the wave field along the lines of wavelet transformations which may lead to more compact matrices and possible extension to two dimensions.

Fast ion distributions are, by nature of their origin, non-Maxwellian. Consequently any investigation of their effects on wave propagation and absorption must allow for such distributions. The wave solutions can then be used by bounce-averaged Fokker-Planck codes to generate self-consistent distributions. We are working to develop the constitutive relation between J and E using expansions of the distribution appropriate for kinetic codes. Among many applications, this will be useful for studies of the effects of fusion alphas, and studies of minority ion current drive [5].

REFERENCES

- [1] N. T. Lam, J. Scharer and M. Bettenhausen, "Effects of the L- to H-mode Transition on a Recessed Coil Antenna in BPX", Nucl. Fusion, **31**, 2141 - 2146 (1991).
- [2] M. Brambilla, "Theory of Bernstein Wave Coupling with Loop Antennas," Nuclear Fusion, vol. 28, no. 4, pp. 549-563, (1988).
- [3] R. Sund and J. Scharer, "Full Wave Field Solutions and Power Conservation for the Ion Cyclotron Range of Frequencies (ICRF) in Nonuniform Plasmas," Physics of Fluids B3, 1326 (1991).
- [4] J. E. Scharer, N. T. Lam and M. Bettenhausen, "ICRF H-mode Coupling and Heating in BPX and JET", 18th European Conference on Controlled Fusion and Plasma Physics, **15c**, Part III, pp. 381-384 (1991).
- [5] J. Jacquinet, Private communication and presented at Boulder ICRF workshop, 1992.
- [6] R. Kock and D. Van Eester, "Ion cyclotron Heating Scenarios for NET Revisited," Nuclear Fusion, **29**, 59, (1989).
- [7A] O. Sauter and C. Vaclavik, "Non-local Effects of Alpha Particles on Second-Harmonic Heating," poster at Sherwood Theory Conference, Seattle, WA, (1991).
- [7B] D. Smithe et al., Nuclear Fusion **27**, 1319 (1987).
D. Smithe et al., Phys. Rev. Lett. **29**, 801 (1987).
- [8] Porcelli et al., "Review of Fast Ion Stabilization of Monster Sawteeth," Invited paper to be published, Berlin EPS, (1991).
- [9] C.S. Chang, "Control of Energetic Ion Confinement by Ion Cyclotron Range of Frequency Waves," Phys. Fluids B **3**, 259, (1991).

V. List of Publications and Papers During the Present Grant Period (1991-92).

A. Published Conference Papers

1. M. Porkolab,...., N. T. Lam, J. Scharer, et al., "Non-inductive Current Drive Scenarios and Bootstrap Currents in the Steady State Advanced Tokamak", Brussels RF Heating Conference (1992).
2. N. T. Lam, J. Scharer and M. Bettenhausen, "Effects of the L- to H-Mode Transition in BPX on a recessed coil antenna", AIP Conf. Proc. **224**, 209-212 Charleston (1991).
3. J. Scharer, N. T. Lam, R. Sund and M. Bettenhausen, "ICRF Heating in BPX", AIP Conf. Proc. **224**, 227-230, Charleston (1991).

B. Conference Abstracts and Posters

1. M. Bettenhausen, J. Scharer and N. T. Lam, "A study of the coupled power spectrum for ICRF on JET and BPX", BAPS 36, Tampa (1991).)
2. J. Scharer, R. Sund, M. Bettenhausen and N. Lam, "ICRF heating in BPX and JET", BAPS 36, Tampa (1991).

C. Workshop Papers

1. J. Scharer, "ICRF Heating and Current Drive Studies for Advanced Tokamaks", Fifth Boulder Workshop on ICRF, Boulder (April, 1992).
2. J. Scharer, N. T. Lam, M. Bettenhausen and R. Sund, "ICRF Heating and Current Drive Studies for Advanced Tokamaks", GA SSAT workshop, San Diego (1992)
3. J. Scharer, N.T. Lam, M. Bettenhausen and R. Sund, "ICRF Antenna Coupling and Wave Absorption by Fast Particles", Princeton ICRF Physics Workshop, August, 1992.

D. Reviewed Journal Papers

1. N. T. Lam, J. Scharer and M. Bettenhausen, "Effects of the L- to H-mode transition on a recessed coil antenna in BPX", Nucl. Fusion, **31**, 2141-2146 (1991).
2. D. B. Batchelor, ...J. E. Scharer, et al., "Burning Plasma Experiment Special Issue, VI. ICRF Heating", Fusion Technology, **21**, 1214-1242 (1992).

E. Reports

1. N.T. Lam, J.E. Scharer and M. Bettenhausen, "Effects of the L- to H-Mode Transition in BPX on a Recessed Coil Antenna (June 1991).
2. J.E. Scharer, O.C. Eldridge, S.F. Chang, Y.S. Zhang, M. Bettenhausen and N.T. Lam, "Electron Cyclotron Wave Propagation, Absorption, and Backscatter Measurements in a Laboratory Plasma." Submitted to IEEE Plasma Sciences, 1992.

Appendix A Grant Modalities

During the current grant period (11/16/91 - 11/15/92 - \$70K) the participating personnel and their approximate degree of involvement has been as follows:

- A. John Scharer, Principal Investigator (15% for 12 months)
- B. Dr. N. Lam, Scientist (10% for 12 months)
- C. M. Bettenhausen, graduate student (50% for 12 months) Including DEC workstation operations management for General Fusion Theory and Computational Research.
- D. R. Sund, graduate student (33% for 12 months)
- E. Y. Mouzouris, undergraduate student assisting in fusion computational runs for independent studies course credit.

END

**DATE
FILMED**

1 / 28 / 93