~~UCSD 010000~~
Progress Report for Grant No. DOE DE-FG03-90ER40546
under the Outstanding Junior Investigator Program

DOE/ER/40546--T1

DOE/ER/40546--T1

DE93 000648

STUDIES IN THEORETICAL PARTICLE PHYSICS

University of California at San Diego

Received by OST
OCT 0 9 1992

Principal Investigator: David B. Kaplan

August 1, 1992

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

TABLE OF CONTENTS

1. Overview	2
2. Chiral Fermions on the Lattice	3
3. Weak Scale Baryogenesis	5
4. Other Topics	8
5. Recent Publications	9
6. Recent Conference Talks	10

[1] OVERVIEW

This research progress report is being submitted to the Department of Energy for renewal of grant no. DOE DE-FG03-90ER40546 under the Outstanding Junior Investigator Program. The grant supports myself (David B. Kaplan) and a graduate student (Martin Schmaltz) in carrying out fundamental research in theoretical particle physics.

This report focuses on research on three distinct areas of particle physics:

1. Chiral Fermions on the Lattice. We are exploring a recent idea I proposed for putting chiral fermions on the lattice. In particular, we are analyzing $1 + 1$ dimensional chiral gauge theories, planning to include dynamical fermions in numerical simulations, and to begin exploration of $3 + 1$ dimensional theories.

2. Weak Scale Baryogenesis. A continuation of work on baryogenesis in the early universe from weak interactions. In particular, an investigation of baryogenesis occurring during the weak phase transition through anomalous baryon violating processes in the standard model of weak interactions.

3. Other Topics. This past year I have also been involved in several other areas of research: one has been an analysis of parity violating nuclear forces, the other an attempt to render the electric dipole moment of the neutron immune from quantum gravity corrections.

A discussion of this work is found in the next three sections, followed by a list of publications produced since my proposal for renewing this grant last year. Footnoted publications in boldface are from the last twelve months.

[2] Chiral Fermions on the Lattice

A. Background

The inability to find a nonperturbative regulator for chiral gauge theories has been a notorious problem for the last twelve years or so. The problem arises from the fact that naive fermions on the lattice always appear to be “doubled”, so that the continuum limit of the theory always consists of vector-like fermion representations. The doubled fermions are just the modes at the corners of the Brillouin zone. In a theory of Dirac fermions one can add a “Wilson term” to push the modes at the Brillouin boundary off to the cutoff, leaving a single Dirac fermion in the continuum limit of the theory. However, there has previously existed no way of devising a lattice regulated theory that had a continuum limit with a chiral representation of fermions.

A strong argument for why such a lattice chiral gauge theory shouldn't exist is that not all such theories can have continuum limits, due to anomalies. On the lattice all classical symmetries of the action are symmetries of the quantum theory, and so if a chiral current is to have an anomalous divergence in the continuum, it should be explicitly broken on the lattice. This is how the $U(1)_A$ anomaly is recovered in lattice QCD with Wilson fermions: the Wilson term explicitly breaks the symmetry on the lattice, and this lattice divergence becomes the anomalous divergence in the continuum limit. However, if one wants to simulate a chiral gauge theory, one cannot break the chiral symmetries on the lattice without breaking gauge invariance, so one cannot apparently reproduce the correct anomaly structure in the continuum.

B. Recent and Proposed Research

Recently I proposed a way to evade the anomaly problem and recover a chiral spectrum in the continuum limit*. The method makes use of the fact that when a Dirac fermion in $2n + 1$ dimensions is coupled to a space dependent mass term with a domain wall structure (i.e, $m(x) = m_0$ for $x > 0$, $m(x) = -m_0$ for $x <$

---

* D.B. Kaplan, “A Method For Simulating Chiral Fermions on the Lattice”, UCSD-PTH-92-16.

0) then there exists a state bound to the $2n$ dimensional defect—a “zero mode”—which is chiral with respect to the $2n$ dimensional γ_5 matrix. The effective low energy theory on the domain wall is then a theory of a chiral fermion propagating in $2n$ dimensions. On the lattice, this theory is afflicted with doubler zero modes which are chirally paired, but a Wilson term which eliminates these doublers may be incorporated in the $2n + 1$ dimensional theory, since the fermion field is Dirac.

Anomalies are easily understood in this theory: the effective low energy theory is not *just* a theory of the zero-modes—the heavy scattering states which are not bound to the domain wall do not entirely decouple, but give rise to a $2n + 1$ dimensional Chern-Simons term. This term exactly accounts for the anomaly. One finds that the anomalous divergence of the $2n$ dimensional currents on the domain wall is due to a current contribution flowing onto or off of the domain wall from the heavy fermions, via the Chern-Simons term, as first discussed by Callan and Harvey. As the coefficient of the Chern-Simons term is equal to the $2n$ dimensional anomaly, it vanishes in the case where the zero-mode representation is anomaly-free (even if chiral). Thus only in anomaly free chiral theories do the heavy fermion modes entirely decouple, leaving us with an effective $2n$ dimensional theory. The effect of the Chern-Simons term. This picture has been confirmed numerically*.

To date my proposal for chiral lattice fermions has only been analyzed for background gauge fields. It is necessary for a complete simulation of $2n$ dimensional chiral gauge theories to have dynamical gauge bosons that only propagate in $2n$ dimensions. To this end I suggested in my paper that the gauge action should have unequal β coefficients for the plaquettes parallel and perpendicular to the domain wall. In particular, the plaquettes perpendicular to the domain wall should have small or vanishing β . In order to test whether this works as hoped, I have applied for supercomputer time at the San Diego Supercomputer Center, and with Schmaltz' and Jansen's help, plan to investigate the $1 + 1$ dimensional Schwinger model formulated as a $2 + 1$ dimensional theory with a domain wall. Known analytic and numerical results will provide good benchmarks to see if the model works as expected. This is the project I expect to take up most of my time in the coming year.

---

* K. Jansen, “Chiral Fermions and Anomalies on a Finite Lattice”, UCSD-PTH-92-18.

[3] Weak Scale Baryogenesis

The photon to baryon ratio observed in the universe today is about 10^{-9} . What is remarkable is that there is *any* asymmetry between baryons and anti-baryons, since we see none on the microscopic level. In the late sixties Sakharov pointed out that baryogenesis would require (1) Baryon number violation; (2) C and CP violation; (3) Nonthermal processes. In the mid-seventies it was realized that the CP violating late decay of particles from Grand Unified Theories (GUTS) could lead to a baryon asymmetry. This is the "standard" picture. One of the great drawbacks of the standard baryogenesis scenario is that it is not testable—all the interesting physics occurs at such a high temperature that the only relic around today is the baryon asymmetry itself.

In the mid-seventies t'Hooft realized that baryon number was violated in the $SU(2) \times U(1)$ theory of weak interactions, now a greatly tested and successful theory. This so-called "anomalous" baryon violation was merely a theoretical curiosity though, since any such process was hugely suppressed at zero temperature. However, starting with speculations by Dimopoulos and Susskind in the late 1970's, and by Kuzmin, Rubakov and Shaposhnikov in the mid- 1980's, it has since been generally agreed upon that these baryon violating processes might have occurred in thermal equilibrium when the temperature of the universe was greater than the $SU(2) \times U(1)$ symmetry breaking scale (about 250 GeV).

Baryogenesis requires more than just baryon violation. The ingredients that are needed, aside from anomalous baryon violation, are CP violation and a departure from thermal equilibrium. The latter is possible, as Kuzmin *et al.* pointed out, if the weak phase transition is first order. If such is the case (and many viable models appear to undergo a first order weak phase transition), then there will always be some supercooling and departure from equilibrium.

As for CP violation, it is easy to show that Kobayashi-Maskawa (KM) CP violation is always too small to account for the observed baryon asymmetry, due to suppression from small Yukawa couplings. (Basically, the reason is that a dimensionless measure of CP violation in the KM model yields a number of order 10^{-20} , which is much smaller than the observed baryon to entropy ratio of 10^{-10} .)

As a result, a successful theory of weak scale baryogenesis necessarily involves an extension of the standard model to incorporate extra CP violation.

This explains why weak scale baryogenesis is such an interesting topic to pursue: from the observed baryon asymmetry one is led to constraints on the symmetry breaking sector of the weak interactions (in order to obtain a first order phase transition) as well as predictions for new particles and new CP violation at the weak scale—all with experimental consequences.

Having all of the necessary ingredients in an acceptable theory does not necessarily lead to baryogenesis and it is important to construct and analyze explicit models. I have been involved in this endeavour with coauthors A. E. Nelson (UCSD) and A. G. Cohen (BU) for the last two years. In 1990 we developed the first explicit and phenomenologically acceptable model for weak scale baryogenesis based on an extension of the standard model involving right handed neutrinos and a CP violating neutrino mass matrix*. During the weak phase transition the phase boundary sweeps through space. In this model, CP violation in the neutrino mass matrix leads to a net lepton number flux being reflected in front of the moving phase boundary, even though there is no net lepton number impinging on the domain wall. This excess lepton number is then converted into baryon number in the unbroken phase by anomalous weak processes. The phase boundary then passes over the baryons, which saves them from decay since anomalous baryon violating processes are highly suppressed in the broken phase. As a consequence of this model, we find that in order to get a sufficiently large baryon asymmetry, the tau neutrino mass is constrained to lie in the experimentally accessible range

$$5 \text{ MeV} \lesssim m_{\nu_\tau} \lesssim 35 \text{ MeV} .$$

More recently we have been investigating models in which the requisite CP violation is in the Higgs sector**. By making use of the mechanism of “spontaneous baryogenesis” developed in an earlier work by Cohen and myself*** we showed how

* A.G. Cohen, D.B. Kaplan and A.E. Nelson, *Phys. Lett.* B245 (1990) 561; *Nucl. Phys.* B349 (1991) 727.

** A.G. Cohen, D.B. Kaplan and A.E. Nelson, *Phys. Lett.* B263 (1991) 86; A.E. Nelson, D.B. Kaplan and A.G. Cohen, *Nucl. Phys.* B373 (1992) 453.

*** A. G. Cohen and D. B. Kaplan, *Phys. Lett.* B199 (1987) 251; *Nucl. Phys.* B308 (1988) 913.

a first order weak phase transition with Higgs CP violation and broad domain wall structure can temporarily generate a chemical potential for axial top quark number (only the top quark is involved, due to its large Yukawa coupling). Creating an asymmetry in left handed top quark number biases the anomalous $SU(2)$ events in the unbroken phase and is converted into a baryon number asymmetry.

The effect we find is much larger than those proposed by others.

Our most recent work on the subject has been to analyze the case of thin domain walls. Then we find that the mechanism first discussed in the neutrino model works as well in the two higgs model with CP violation in the top quark Yukawa coupling. We find that in this regime the baryon asymmetry can be orders of magnitude larger than estimated by other parties*.

We have also recently considered the issue of Debye screening of gauge charges during the reflection process. Others have suggested that this will dilute the effect we have found, but we have recently found that not to be the case**.

In a separate piece of research I showed that electroweak baryogenesis could easily explain why the dark matter density is 10–100 times the baryon density. The reason is that baryons are produced via instantons, which require the concomitant production of other fermions. As a result it is natural to cogenerate dark matter along with baryon number. One finds that the dark matter to baryon energy density ratio may be understood as the ratio of the weak to strong physics scales...basically the W to proton mass ratio***.

In the coming year I expect to work on a review of the subject, as well as perhaps an analysis of supersymmetric models.

---

* A.E. Nelson, D.B. Kaplan and A.G. Cohen, *Nucl. Phys.* B373 (1992) 453.

** A.G. Cohen, D.B. Kaplan and A.E. Nelson, "Debye Screening and Baryogenesis During the Electroweak Phase Transition", UCSD-PTH-92-19.

*** D.B. Kaplan, *Phys. Rev. Lett.* 68 (1992) 741.

[4] Other Topics

1. STRANGE MATRIX ELEMENTS AND PARITY VIOLATING PION-NUCLEON COUPLINGS

This past year M. Savage and I wrote a paper that explored the chiral structure of possible pion nucleon parity violating interactions*. We found a number of multipion vertices previously ignored, and concluded that in the $I = 1$ channel a large effect could arise from nonzero strange matrix elements in nucleons. This claim was bolstered by an analysis I performed of nonleptonic hyperon decays which, when assuming $SU(3)$ symmetry, yields strong evidence for large strange matrix elements in the nucleon**. This subject is of topical interest as there are numerous experiments presently looking for parity violation in nuclear physics.

2. CP VIOLATION AND GRAVITY

K. Choi, A.E. Nelson and I recently wrote a paper discussing an issue or much discussion lately about whether there is any way to understand why “Planck slop” (operators induced by quantum gravity) doesn’t lead to a large electric dipole moment of the neutron, by infecting the strong interactions with CP violation from the weak sector. We suggested that this could be avoided if CP was a gauge symmetry – i.e, if CP was a discrete subgroup of a local continuous symmetry that had been spontaneously broken. We concluded that this was possible in theories with dimensional compactification, and catalogued the constraints such theories must satisfy***.

---

* D.B. Kaplan and M. Savage, “An Analysis of Parity Violating Pion Nucleon Couplings”, UCSD-PTH-92-04.

** D.B. Kaplan, *Phys. Lett.* B275 (1992) 137.

*** K. Choi, D.B. Kaplan and A.E. Nelson, “Is CP a Gauge Symmetry?”, UCSD-PTH-92-11.

[5] RECENT PUBLICATIONS[†]

David B. Kaplan

- [1] Why There Is Something Rather Than Nothing: Matter from Weak Interactions (with A.G. Cohen and A.E. Nelson), Nucl. Phys. B373 (1992) 453.
- [2] A Single Explanation For Both the Baryon and Dark Matter Densities, Phys. Rev. Lett. 68 (1992) 741.
- [3] Strong Evidence for a New Strange Matrix Element, Phys. Lett. B275 (1992) 137.
- [4] An Analysis of Parity Violating Pion-Nucleon Interactions, UCSD/PTH-92-04.
- [5] Is CP a Gauge Symmetry? (with K. Choi and A. E. Nelson), UCSD/PTH-92-11.
- [6] A Method for Simulating Chiral Fermions on the Lattice, UCSD/PTH-92-16.
- [7] Debye Screening and Baryogenesis at the Electroweak Phase Transition, UCSD/PTH-92-19.

[†] Research papers from the last twelve months and not included in last year's grant renewal proposal.

[6] RECENT CONFERENCE TALKS[†]

David B. Kaplan

- [1] A New Mechanism for Generating Fermion Masses With Implications for Flavor, *Proceedings of the 15th Johns Hopkins Workshop on Current Problems in Particle Theory (1991)*, ed. G. Domokos and S. Kovesi-Domokos, World Scientific.
- [2] Dark Matter from the Electroweak Anomaly, UCSD PTH/91-32, to appear in *Proceedings of the International Workshop on Electroweak Symmetry Breaking (Hiroshima, 1991)*, World Scientific.

[†] Conference talks not included in last year's grant renewal proposal.

END

DATE
FILMED

11 / 23 / 92