41348 J. Q.
PPPL

PRINCETON
PLASMA PHYSICS
LABORATORY

PPPL-TH92-2
UC-420,427

PPPL-TH92-2

**PRINCETON UNIVERSITY
PLASMA PHYSICS LABORATORY**

**THEORY DIVISION
QUARTERLY REPORT***

April 1-June 30, 1992

*This work was supported by U.S. Department of Energy Contract No. DE-AC02-76-CHO-3073.

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

NOTICE

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial produce, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

NOTICE

This report has been reproduced directly from the best available copy.
Number of pages in this report: 15

Available to DOE and DOE contractors from the:
Office of Scientific and Technical Information
P.O. Box 62, Oak Ridge, TN 37831;
Prices available from (615) 576-8401.

Available to the public from the:
National Technical Information Service
U.S. Department of Commerce
5285 Port Royal Road, Springfield, Virginia 22161
(703) 487-4650

PPPL THEORY NEWS is a quarterly bulletin presenting condensed versions of selected research reports recently printed and distributed by the PPPL Theory Division at PPPL. Lists of printed reports and publications, external presentations at professional meetings, and recent visitors to the group are also included. Further information may be obtained by contacting the individual authors. Copies of PPPL Reports may be requested from the R & P Office at PPPL. Our address is:

Princeton Plasma Physics Laboratory
James Forrestal Campus
P.O. Box 451
Princeton, NJ 08543

Phone: (609) 243-2656
Fax: (609) 243-2662

Roscoe B. White, *Editor*
Terry Greenberg, *Copy Production & Layout*

Contents

1	Long-Wavelength Microinstabilities in Toroidal Plasmas	3
2	Alpha Particle Destabilization of TAE modes in TFTR DT Experiments	8
3	Theory Publication List	12
4	Theory Visitor List	13

1. Long-Wavelength Microinstabilities in Toroidal Plasmas

W.M. Tang and G. Rewoldt

Realistic kinetic toroidal eigenmode calculations have been carried out to support a proper assessment of the influence of long-wavelength microturbulence on transport in tokamak plasmas. This is a very important and relevant issue in that scattering diagnostics have generally indicated that the largest fluctuation amplitudes occur at the longest measurable wavelengths. The recent implementation of new diagnostics (beam-emission spectroscopy and reflectometry) capable of measuring much longer wavelength fluctuations in major devices such as TFTR has greatly heightened interest in the development of appropriate theoretical models of such phenomena. Moreover, results from empirical confinement scaling studies have indicated that Bohm-like trends characteristic of long-wavelength modes could be supported. Additional motivation for these eigenmode studies is provided by the need to produce realistic self-consistent benchmarks for toroidal gyrokinetic and gyrofluid codes currently under very active development.

Fully two-dimensional (r, θ) mode structures for long-wavelength trapped-ion and toroidal ion-temperature-gradient (ITG) modes have been calculated using a comprehensive toroidal finite-element code with the unique capability for evaluating large-scale kinetic behavior extending over many rational surfaces. Improvements over previously used numerical methods [1] were necessary in order to tractably evaluate eigenvalues for realistic tokamak parameters. An important new result from the present studies is that in contrast to the long-accepted picture predicted by one-dimensional radial eigenmode calculations [2] and supported by earlier idealized (very weak magnetic shear) two-dimensional calculations [1], trapped-ion eigenmodes are not strongly localized between rational surfaces. This is due to the fact that toroidal microinstabilities generally exhibit a strong ballooning character which can be associated with poloidal localization effects caused both by magnetic trapping as well as unfavorable curvature. The resultant coupling of the poloidal (m) harmonics produces radial structures which are relatively insensitive to ion Landau damping at the rational surfaces. This trend can be seen in the plot of a representative 2-D trapped-ion mode eigenfunction in Fig. 1. Note that the magnitude of $\tilde{\Phi}$ peaks around $\theta = 0$ (B-field minimum) and exhibits no significant sensitivity in the radial direction to the location of the rational surfaces there. This is more clearly displayed in Fig. 2 which shows the radial structure of the eigenfunction at $\theta = 0$ and also at $\theta = \pi$ (B-field maximum). Since ballooning effects are minimal at $\theta = \pi$, the sensitivity to the rational surfaces, as expected, is evident here. Of course, the amplitude of the eigenfunction is clearly much smaller than the dominant component localized near the B-field minimum.

The results in Figs. 1 and 2 illustrate an important general trend found in the present studies. Specifically, even at very long wavelengths (*i.e.*, $k_{\theta} \rho_i \ll 1$, $n < 10$), the mode structure of toroidal microinstabilities is dominantly influenced by ballooning features and not by the radial fine-structure effects associated with ion Landau damping dynamics near the rational surfaces. This suggests that a one-dimensional ballooning representation analysis might capture the essential characteristics of these eigenmodes. Calculations were accordingly carried out using a comprehensive kinetic ballooning code [3] normally used to

Fig. 1. Fully 2-D (r, θ) trapped-ion eigenmode structure ($n=5$) illustrates large scale kinetic behavior extending over many rational surfaces for generic TFTR-like parameters (i.e., $R = 248$ cm, $a = 85$ cm, $B = 5$ T, $n(0) = 6 \times 10^{13}$ cm $^{-3}$, $T_e(0) = T_i(0) = 5$ keV, and parabolic-type q -profile with $q_0 = 1$ and $q_a = 3.2$).

Fig. 2. Comparison of the small amplitude radial structure at $\theta = \pi$ (no ballooning) with the large amplitude radial structure at $\theta = 0$ (maximum ballooning) for the $n = 5$ trapped-ion eigenmode of Fig. 1.

evaluate moderate to high toroidal mode number (n) instabilities. The linear spectral properties (growth rate and real frequency vs. toroidal mode number) for a representative TFTR plasma were obtained and are displayed in Fig. 3. One-dimensional kinetic ballooning code results are represented by circles while the fully 2-D results are denoted by squares. Good agreement (less than 50% difference) is found for $n = 3$ through $n = 10$; i. e., the 1-D analysis provides reasonable estimates of the linear eigenvalues down to very long wavelengths. Below $n = 3$, these drift-type instabilities (ω scaling roughly with n) become sufficiently weak so that the ballooning character no longer dominates. Hence, the fully 2D-type calculations become necessary here. However, at the higher n -values, where the 2D-computations become increasingly expensive, the present results demonstrate that the 1D-code can be applied with some confidence. This is computationally analogous to earlier MHD studies which showed that 1-D ballooning codes could produce results consistent with those from the 2-D PEST-code at moderate values of n [4]. Results of the type illustrated in Fig. 3 demonstrate that, employed together, our 2-D and 1-D codes provide a unique capability for calculating the linear properties of toroidal microinstabilities over the entire spectral range of interest. It is interesting to note here that results [5] from very recent beam-emission spectroscopy measurements of long wavelength fluctuation properties in TFTR plasmas have indicated

Fig. 3. Representative linear spectrum (growth rate and real frequency vs. toroidal mode number) for realistic TFTR L-mode discharge parameters.

a possible peaking of the spectrum near $k_{\theta}\rho_i = 0.1$ – a trend qualitatively similar to that displayed in Fig. 3.

In order to gain better insight into the nonlinear behavior of electrostatic microinstabilities in tokamak plasmas, a fully toroidal 3-D gyrokinetic particle code has recently been developed to study the properties of ion temperature gradient (ITG) driven modes [6]. Comparisons between results from the linear phase of these simulations and our realistic toroidal eigenmode calculations have provided valuable self-consistent benchmarks for the particle code. In particular, the simulations indicate that unstable modes of moderate (m, n) with ballooning-type structures are dominant in the linear stage. Reasonable agreement is found with respect to the magnitude of the eigenvalues (within about 25%) as well as with the characteristic features of the eigenfunctions. Regarding the latter, the mode structures exhibit significant ballooning in poloidal angle due to unfavorable curvature around the B-field

minimum and a radial localization around the temperature gradient maximum with little sensitivity to ion Landau damping at the rational surfaces. Future benchmarking studies of this type in support of further development of the toroidal gyrokinetic code will be carried out and will involve, for example, the implementation of important effects associated with trapped-electron dynamics and collisional dissipation.

References

- [1] R. Marchand, W.M. Tang, and G. Rewoldt, *Phys. Fluids* **23**, 1164 (1980).
- [2] N.T. Gladd and D.W. Ross, *Phys. Fluids* **16**, 1706 (1973).
- [3] G. Rewoldt, W.M. Tang, and R.J. Hastie, *Phys. Fluids* **30**, 807 (1987).
- [4] R.L. Dewar, J. Manickam, R.C. Grimm, and M.S. Chance, *Nucl. Fusion* **21**, 493 (1981).
- [5] R.J. Fonck, N. Bretz, G. Cosby, R. Durst, E. Mazzucato, R. Nazikian, S. Paul, S. Scott, W.M. Tang, and M. Zarnstorff, *Proceedings of 19th European Physical Society Conference on Controlled Fusion and Plasma Heating*, invited paper (1992).
- [6] S.E. Parker and W.W. Lee, invited paper 1B2, *International Sherwood Fusion Theory Conference*, Santa Fe, NM (1992).

2. Alpha Particle Destabilization of TAE modes in TFTR DT Experiments

G. Y. Fu and C. Z. Cheng

The toroidicity-induced shear Alfvén eigenmode (TAE) [1] has received increasing attention recently. It has been shown theoretically [2–4] and experimentally [5,6] that the TAE mode can be destabilized by energetic/alpha particles in tokamaks. The excitation of the TAE modes may degrade the confinement of fusion alpha particles in an ignited tokamak. Because the TAE mode is a global toroidal shear Alfvén eigenmode, direct comparison between theory and experimental measurement is possible.

The TAE modes have discrete frequencies located inside the toroidicity-induced continuum gap. If one neglects the toroidicity, then the shear Alfvén spectrum can be described in its cylindrical form $\omega^2 = (n - m/q)^2 v_A^2 / R^2$ where n/m is the toroidal/poloidal mode number. Note that the Alfvén spectrum is labeled by each pair of (n, m) and is a continuous function of minor radius r since both the safety factor q and the Alfvén speed v_A depend on r . In particular, the spectrum is degenerate at $q = q_g(r = r_g) = (2m + 1)/2n$ where the continuum spectrum with the poloidal mode number m crosses with that of $m + 1$. This degeneracy is broken by the toroidicity through the mode coupling between the mode m and the mode $m + 1$, and a continuum gap is created at $q = q_g$. The TAE mode has the following distinct characteristics: 1) In the ideal MHD limit, it is a marginally stable mode with a finite real frequency given approximately by $\omega = v_A/2qR$; 2) The mode is localized near the gap location at $r = r_g$ (between two rational magnetic surfaces) with a radial width on the order of r_g^2/R .

The TAE mode can be destabilized by fusion alpha particles through wave-particle resonant interaction. The TAE mode becomes unstable by tapping the free energy associated with the pressure gradient of the alpha particles. This mechanism can be easily understood by considering the linear power transfer from the particles to the TAE mode, which is given by

$$P_\alpha = \pi \int d^3x \int d^3v q_e^2 \left(\frac{\omega_*}{\omega} F + T \frac{\partial F}{\partial E} \right) (v_\parallel E_\parallel + v_d E_\perp)^2 \delta(\omega - k_\parallel v_\parallel) \quad (1)$$

where q_e is the charge, ω_* is the diamagnetic frequency, F is the particle distribution function, T is the particle temperature, v_d is the particle magnetic drift velocity, $E_\parallel$ and $E_\perp$ are the parallel and perpendicular components of the electric field of the TAE mode respectively, and $k_\parallel$ is the parallel wave vector. Physically, the $\frac{\partial F}{\partial E}$ term gives the usual Landau damping associated with the velocity space gradient (for monotonic energy distribution assumed here). On the other hand, the ω_* term is destabilizing and is associated with the pressure inhomogeneity. The balance of these two terms determines whether the particles give or lose their energy to the wave. For fusion alpha particles, it is strongly destabilizing since $\omega_*/\omega \gg 1$ due to the steep alpha pressure profile.

In this work, the stability of the TAE mode for the TFTR DT experiments is studied with the NOVA-K global stability code. A quadratic dispersion relation is used to calculate perturbatively the destabilizing effects of fusion alpha particles, as well as the stabilizing effects of thermal

particles. In addition, the collisional damping due to trapped electrons has been added into the NOVA-K code. These kinetic effects may be treated perturbatively since they are smaller than the ideal MHD terms by a factor of $\beta_j \ll 1$ for each species j . In calculating the trapped electron's collisional effects, we solve exactly the bounce-averaged drift kinetic equation for electrons, including the pitch angle scattering collisional operator. Thus, no assumption on the ordering of ν_e/ω is made, where ν_e is the electron's collisional frequency.

To study TAE instabilities for TFTR DT experiments, we will consider the super shot operational scenario. The TFTR parameters for the DT experiments are taken to be: the major radius $R = 245\text{cm}$, the minor radius $a = 80\text{cm}$, the toroidal magnetic field $B = 5\text{T}$, the alpha particle pressure scale length $L_\alpha = 0.3a$. The core plasma parameters are scaled according to the TFTR super shot beam power scaling with steady state plasma: $\langle \beta \rangle = 0.13 + 0.025P_b$, $T_e(0) = 5.7 + 0.11P_b$, $T_i(0) = 7.2 + 0.75P_b$, $n_e(0) = 2.2 + 0.16P_b$, where P_b is the neutral beam heating power in unit of MW. Furthermore, we choose plasma profiles as, $P = P(0)(1 - \Psi)^2$, $n_e = n_e(0)(1 - 0.8\Psi)$, $q(0) = 1.1$, $q(1) = 5.5$, $q'(0) = 1.0$, $q'(1) = 14.0$, where Ψ is the normalized poloidal flux and the prime indicates the derivative with respect to Ψ . These profiles are characteristic of TFTR super shots. Figure 1 shows the critical volume averaged beta of alpha particles $\langle \beta_\alpha \rangle$ for TAE instabilities versus the core plasma β . The projected range of $\langle \beta_\alpha \rangle$ in steady state DT operation is shown by the shaded area. We see that the $n = 1$ TAE mode is stable except for a small range of core plasma beta value around $\beta = 0.4\%$ where the mode is weakly unstable. On the other hand, the $n = 2$ TAE mode is unstable for $\beta < 0.6\%$. Note that the critical alpha beta for TAE instabilities decreases rapidly as plasma β decreases. This is because ion Landau damping is the dominating damping mechanism. The electron Landau damping rate and the collisional damping rate are both very small, on the order of 10^{-2} of the ion Landau damping rate. Since the ion Landau damping is roughly proportional to $\beta_i^{-3/2} \exp(1/9\beta_i)$, and β_i decreases as beam power decreases, the instability scenario for the $n = 1$ TAE mode is proposed by turning off the beam power. As the beam power is turned off, the alpha particle beta decreases much slower than the core plasma beta and can remain above the threshold for the $n = 1$ instability for several hundreds of ms. Figure 2 shows a typical time dependent trajectory of β_α and v_α/v_A obtained from a TRANSP simulation of the TFTR DT experiment based on a super shot run (#55848). The arrow indicates the direction of time. We observe that during the increasing phase of β_α (v_α/v_A increases as density increases), the $n = 1$ TAE mode is stable. However, after the beam power is turned off, the alpha particle beta remains relatively constant due to long slowing-down time as v_α/v_A decreases and the TAE mode becomes unstable.

Several approximations are made in obtaining the results described above. We have neglected the finite gyro-radius effects and the finite drift orbit width effects of the alpha particles. We have also neglected the parallel electric field related to the finite ion Larmor radius. These finite orbit effects are expected to be important for high- n TAE modes ($n > 1$ for TFTR) and to increase the stability threshold for these short wavelength modes.

In our recent work [7] on the alpha particle destabilization of high- n TAE modes, we have taken into account the finite orbit width effects of alpha particles and the parallel electric field

Figure 1: Critical alpha particle β , $\langle \beta_\alpha \rangle_{crit}$, as a function of the plasma β .

Figure 2: Critical alpha particle β , $\langle \beta_\alpha \rangle_{crit}$, as a function of v_α/v_A .

in the calculation of the collisional damping due to the trapped electrons. With orbit width effects, the alpha destabilizing contribution has a maximum as a function of the mode number corresponding to $k_{\theta} \rho_{\alpha}$ on the order of unit. Work is in progress to incorporate these effects into the NOVA-K code.

References

- [1] C. Z. Cheng and M. S. Chance, Phys. Fluids **29**, 3695 (1986); C. Z. Cheng, L. Chen, and M. S. Chance, Ann. Phys. (N. Y.) **161**, 21 (1984).
- [2] G. Y. Fu and J. W. Van Dam, Phys. Fluids B**1**, 1949 (1989).
- [3] C. Z. Cheng, G. Y. Fu, and J. W. Van Dam, in Theory of Fusion Plasmas 1988, ed. J. Vaclavik, F. Troyon, and E. Sindoni (Societa Italiana di Fisica/Editrice Compositori, Bologna, 1989), p. 259.
- [4] C. Z. Cheng, Phys. Fluids, B **3**, 2463 (1991).
- [5] K. L. Wong et al., Phys. Rev. Lett. **66**, 1874 (1991).
- [6] W. W. Heidbrink, E. J. Strait, E. Doyle, and R. Snide, Nucl. Fusion **31**, 1635 (1991).
- [7] G. Y. Fu and C. Z. Cheng, Report PPPL-2852, 1992.

3. Theory Publication List 4/92 - 6/92

Fisch, N.J., *Radiation from Mildly Relativistic Electrons in Tokamak Plasmas*, in Nonlinear and Relativistic Effects in Plasmas, edited by V. Stefan (AIP, NY, 1992), 103-110.

Fisch, N.J. and J-M. Rax, *Current-Drive by Lower Hybrid Waves in the Presence of Energetic Alpha-Particles*, Nucl. Fusion **32**, 549-556 (April, 1992) and PPPL 2786.

Fisch, N.J., W. Tighe, E. Valeo, and S. Suckewer, *Two-laser Approach to Shorter Wavelength X-ray Lasing*, in Nonlinear and Relativistic Effects in Plasmas, edited by V. Stefan (AIP, NY, 1992), 314-324.

Fu, G.Y. and C.Z. Cheng, *Excitation of High- n Toroidicity-induced Shear Alfvén Eigenmodes by Energetic Particles and Fusion Alpha Particles in Tokamaks*, Phys. of Fluids B and PPPL Report 2852.

Hahm, T.S., *Trapped Particle Dynamics in Toroidally Rotating Plasmas*, Phys. of Fluids B, PPPL Report 2843.

Perkins, F.W. et al., *Nondimensional Transport Scaling in TFTR: Is Tokamak Transport Bohm or GyroBohm?* Phys. of Fluids and Tech. Report J-386.

Pon, N., *Bootstrap Dependence on Plasma Profile Parameters*, Nucl. Fusion and PPPL Report 2854.

Stotler, D.P., D.P. Coster, B.J. Braams, B. A.B. Ehrhardt, C.F.F. Karney, M. Petravic, *Scrape-Off Layer Modeling Using Coupled Plasma and Neutral Transport Codes*, in Proceedings of 10th Plasma Surface Interaction Conference, Monterey, CA, March 30 - April 3, 1992, to appear in J. of Nuclear Materials: PPPL Report 2842.

Valeo, E. and S. Cowley, *A Microsphere-Based Short-Wavelength Recombination X-Ray Laser*, Phys. Rev. A and PPPL Report 2835.

Valeo, E. J. and S.C. Cowley, K. Krushelnick, W. Tighe and S. Suckewer, *A Microsphere-Based X-Ray Laser* in Proceedings of Third International Colloquium on X-Ray Lasers, Schliersee, Germany, May 18-22, 1992.

Voss, K.E. and N.J. Fisch, *Inverse Problem for Bremsstrahlung Radiation*, Phys. Fluids B4, 762-763 (March, 1992), and PPPL Report 2793.

4. Theory Visitor List 4/92 - 6/92

Bondeson, Anders, Centre de Recherches en Physique des Plasmas, Ecole Polytechnique Federale de Lausanne, Lausanne, Switzerland.

Eastman, Timothy, National Science Foundation, Washington, DC, April.

Ichiguchi, Kasuji, National Institute for Fusion Science, Nagoya, Japan, April.

Laurence, Peter, University of Rome, Rome, Italy, April - May.

Merkel, Peter, IPP, Garching, Germany, April - June.

Ottaviani, Maurizio, ENEA Frascati Energy Centre, Frascati, Rome, Italy.

Salas, Angel, CIEMAT, Madrid Spain, April, June - August.

Schneider, Ralf, IPP, Garching, Germany, March - April.

Shoucri, Magdi, Hydro-Quebec, Varennes, Quebec, May.

Takamura, Hisanori, National Institute for Fusion Science, Nagoya, Japan, April.

Zonca, Fulvio, ENEA-Centro Ricerche Energia, Frascati, Italy, April - May.

EXTERNAL DISTRIBUTION IN ADDITION TO UC-420

Dr. F. Paoloni, Univ. of Wollongong, AUSTRALIA
 Prof. M.H. Brennan, Univ. of Sydney, AUSTRALIA
 Plasma Research Lab., Australian Nat. Univ., AUSTRALIA
 Prof. I.R. Jones, Flinders Univ, AUSTRALIA
 Prof. F. Cap, Inst. for Theoretical Physics, AUSTRIA
 Prof. M. Heindler, Institut für Theoretische Physik, AUSTRIA
 Prof. M. Goossens, Astronomisch Instituut, BELGIUM
 Ecole Royale Militaire, Lab. de Phy. Plasmas, BELGIUM
 Commission-Europeen, DG. XII-Fusion Prog., BELGIUM
 Prof. R. Bouciqué, Rijksuniversiteit Gent, BELGIUM
 Dr. P.H. Sakanaka, Instituto Fisica, BRAZIL
 Instituto Nacional De Pesquisas Espaciais-INPE, BRAZIL
 Documents Office, Atomic Energy of Canada Ltd., CANADA
 Dr. M.P. Bachynski, MPB Technologies, Inc., CANADA
 Dr. H.M. Skarsgard, Univ. of Saskatchewan, CANADA
 Prof. J. Teichmann, Univ. of Montreal, CANADA
 Prof. S.R. Sreenivasan, Univ. of Calgary, CANADA
 Prof. T.W. Johnston, INRS-Energie, CANADA
 Dr. R. Bolton, Centre canadien de fusion magnétique, CANADA
 Dr. C.R. James, Univ. of Alberta, CANADA
 Dr. P. Lukács, Komenského Univerzita, CZECHO-SLOVAKIA
 The Librarian, Culham Laboratory, ENGLAND
 Library, R61, Rutherford Appleton Laboratory, ENGLAND
 Mrs. S.A. Hutchinson, JET Library, ENGLAND
 Dr. S.C. Sharma, Univ. of South Pacific, FIJI ISLANDS
 P. Mähönen, Univ. of Helsinki, FINLAND
 Prof. M.N. Bussac, Ecole Polytechnique, FRANCE
 C. Mouttet, Lab. de Physique des Milieux Ionisés, FRANCE
 J. Radet, CEN/CADARACHE - Bat 506, FRANCE
 Prof. E. Economou, Univ. of Crete, GREECE
 Ms. C. Rinni, Univ. of Ioannina, GREECE
 Dr. T. Muai, Academy Bibliographic Ser., HONG KONG
 Preprint Library, Hungarian Academy of Sci., HUNGARY
 Dr. B. DasGupta, Saha Inst. of Nuclear Physics, INDIA
 Dr. P. Kaw, Inst. for Plasma Research, INDIA
 Dr. P. Rosenau, Israel Inst. of Technology, ISRAEL
 Librarian, International Center for Theo Physics, ITALY
 Miss C. De Palo, Associazione EURATOM-ENEA, ITALY
 Dr. G. Grosso, Istituto di Fisica del Plasma, ITALY
 Prof. G. Rostangni, Istituto Gas Ionizzati Del Cnr, ITALY
 Dr. H. Yamato, Toshiba Res & Devel Center, JAPAN
 Prof. I. Kawakami, Hiroshima Univ., JAPAN
 Prof. K. Nishikawa, Hiroshima Univ., JAPAN
 Director, Japan Atomic Energy Research Inst., JAPAN
 Prof. S. Itoh, Kyushu Univ., JAPAN
 Research Info. Ctr., National Inst. for Fusion Science, JAPAN
 Prof. S. Tanaka, Kyoto Univ., JAPAN
 Library, Kyoto Univ., JAPAN
 Prof. N. Inoue, Univ. of Tokyo, JAPAN
 Secretary, Plasma Section, Electrotechnical Lab., JAPAN
 S. Mori, Technical Advisor, JAERI, JAPAN
 Dr. O. Mitarai, Kumamoto Inst. of Technology, JAPAN
 J. Hyeon-Sook, Korea Atomic Energy Research Inst., KOREA
 D.I. Choi, The Korea Adv. Inst. of Sci. & Tech., KOREA
 Prof. B.S. Liley, Univ. of Waikato, NEW ZEALAND
 Inst of Physics, Chinese Acad Sci PEOPLE'S REP. OF CHINA
 Library, Inst. of Plasma Physics, PEOPLE'S REP. OF CHINA
 Tsinghua Univ. Library, PEOPLE'S REPUBLIC OF CHINA
 Z. Li, S.W. Inst Physics, PEOPLE'S REPUBLIC OF CHINA
 Prof. J.A.C. Cabral, Instituto Superior Tecnico, PORTUGAL
 Dr. O. Petrus, AL I CUZA Univ., ROMANIA
 Dr. J. de Villiers, Fusion Studies, AEC, S. AFRICA
 Prof. M.A. Hellberg, Univ. of Natal, S. AFRICA
 Prof. D.E. Kim, Pohang Inst. of Sci. & Tech., SO. KOREA
 Prof. C.I.E.M.A.T, Fusion Division Library, SPAIN
 Dr. L. Stenflo, Univ. of UMEA, SWEDEN
 Library, Royal Inst. of Technology, SWEDEN
 Prof. H. Wilhelmson, Chalmers Univ. of Tech., SWEDEN
 Centre Phys. Des Plasmas, Ecole Polytech, SWITZERLAND
 Bibliotheek, Inst. Voor Plasma-Fysica, THE NETHERLANDS
 Asst. Prof. Dr. S. Cakir, Middle East Tech. Univ., TURKEY
 Dr. V.A. Glukhikh, Sci. Res. Inst. Electrophys. Apparatus, USSR
 Dr. D.D. Ryutov, Siberian Branch of Academy of Sci., USSR
 Dr. G.A. Eliseev, I.V. Kurchatov Inst., USSR
 Librarian, The Ukr.SSR Academy of Sciences, USSR
 Dr. L.M. Kovrizhnykh, Inst. of General Physics, USSR
 Kernforschungsanlage GmbH, Zentralbibliothek, W. GERMANY
 Bibliothek, Inst. Für Plasmaforschung, W. GERMANY
 Prof. K. Schindler, Ruhr-Universität Bochum, W. GERMANY
 Dr. F. Wagner, (ASDEX), Max-Planck-Institut, W. GERMANY
 Librarian, Max-Planck-Institut, W. GERMANY
 Prof. R.K. Janev, Inst. of Physics, YUGOSLAVIA

END

**DATE
FILMED
02/19/93**