MASTER

MASS SPECTROMETRIC STUDIES OF HYDRAZINE
PHOTOOXIDATION BY ILLUMINATED CHLOROPLASTS

DISCLAIMER

This book was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

Richard Radmer and Otto Ollinger
Martin Marietta Laboratories
1450 South Rolling Road
Baltimore, Maryland 21227

Running Title: Hydrazine Photooxidation by Chloroplasts

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

ESB

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency Thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

SUMMARY

Mass spectrometric techniques were used to directly monitor the products evolved during the course of hydrazine (NH_2NH_2) photooxidation by chloroplasts exposed to short saturating flashes or continuous high light. We found that:

- (1) Molecular N_2 was the sole volatile product of hydrazine photooxidation. Isotopic studies showed that the N-N bond remained intact during the $\text{NH}_2\text{NH}_2 \rightarrow \text{N}_2$ transformation. Under conditions in which spurious side reactions were minimized (see item 3 below), the N_2 yield was equal to the O_2 yield during H_2O photooxidation.
- (2) In the presence of 3-(3,4-dichlorophenyl)-1,1-dimethylurea, N_2 was evolved, but only on the first flash, suggesting that N_2 was formed by the combination of single electron oxidation products of hydrazine.
- (3) In addition to its production by Photosystem II, N_2 can also be generated by a series of secondary reactions mediated by superoxide. This "extra" N_2 evolution can be eliminated by the addition of superoxide dismutase.

Our results indicate that hydrazine can be used as a reliable probe of Photosystem II provided that a) N_2 evolution (rather than O_2 uptake) is monitored, and b) precautions are taken to minimize spurious side reactions. Under conditions in which the participation of superoxide is minimized, N_2 evolution accurately reflects the photooxidation of hydrazine by Photosystem II.

ABBREVIATIONS

PSII, photosystem II; PSI, photosystem I; MS, mass spectrometer; DCMU, 3-(3,4-dichlorophenyl)-1,1-dimethylurea; Fd, ferredoxin; MV, methylviologen; SOD, superoxide dismutase.

Key Words: hydrazine, photosynthesis, photosystem II, mass spectrometer

INTRODUCTION

Hydrazine (NH_2NH_2) can serve as an electron donor to PSII and thus can be used to study reactions near the site of O_2 evolution (1,2). At low concentrations, NH_2NH_2 , like hydroxylamine (NH_2OH), can effect a delay in the normal O_2 flash yield pattern without significantly decreasing O_2 evolution capacity (3). Thus this compound is one of a rather small group of electron donors that can be oxidized by PSII and not destroy the O_2 -evolving system (4).

Early hydrazine photooxidation studies (1,2) showed that the rate of hydrazine-supported O_2 uptake could significantly exceed the rate of PSII turnover. The excess O_2 uptake was attributed to the interaction of hydrazine or one of its oxidation products with dissolved oxygen (1).

In a previous report (5), we described the results of experiments in which we used mass spectrometric techniques to directly monitor the products generated during the photooxidation of hydroxylamine by isolated chloroplasts. We found that molecular N_2 , formed by the combination of two one-electron oxidation products of NH_2OH , was the only stable reaction product.

In this communication, we describe experiments in which we studied the processes involved during the photooxidation of hydrazine by PSII. We found that, as in the case of hydroxylamine, molecular N_2 was the only significant stable reaction product. However, unlike hydroxylamine, hydrazine can also be oxidized via a series of secondary reactions, which also produce N_2 . Our data were consistent with a reaction scheme in which the interaction of NH_2NH_2 and O_2^- serve to generate the quantities of N_2

above and beyond that attributable to the direct oxidation of NH_2NH_2 by PSII. In agreement with this scheme, we found that the stoichiometry between PSII-generated oxidant and N_2 could be varied widely by altering the concentration of internally generated O_2^- .

MATERIALS AND METHODS

The mass spectrometric apparatus and techniques for directly monitoring the gas exchange elicited by short saturating flashes (5,10) and continuous light (6,7) were described earlier.

Tris-extracted chloroplasts were used for all experiments. Chloroplasts were isolated (8) from greenhouse or market spinach in STN medium (0.4 M sucrose, 0.05 M Tris-HCl, 0.01 M NaCl, pH 7.4). These chloroplasts were extracted by diluting a 1-2 ml aliquot (~ 4 mg chl/ml) to 100 ml with a high-Tris solution (0.4 M sucrose, 0.01 M NaCl, 0.75 M Tris-HCl, pH 8.1) and stirring this suspension for 20 minutes in a cold room under dim light (9). The chloroplasts were then centrifuged, washed, and resuspended in STN medium. The extraction efficiency, based on the extent of inhibition of the Hill reaction, was greater than 95%.

[¹⁵N] hydrazine sulfate (95 atom % ¹⁵N) was obtained from Prochem, London, U.K.

RESULTS AND DISCUSSION

Products of hydrazine photooxidation

Figure 1 shows the relative flash yields -- at m/e values corresponding to likely oxidation products -- when chloroplasts were subjected to a train of saturating flashes (3-s spacing) in the presence of 1 mM NH_2NH_2 . Note that molecular N_2 was the only volatile stable oxidation product formed in significant amounts; there was no apparent evolution of nitrogen oxides. Similar results were obtained in continuous light: only m/e values corresponding to molecular N_2 increased when chloroplasts were subjected to strong continuous light in the presence of 10 mM NH_2NH_2 and saturating amounts of ferredoxin and NADP.

We specifically searched for the production of NH_3 [a likely product of NH_2NH_2 oxidation (11)] by running a similar series of experiments both at pH 7.4 and at pH's 9.0 and 9.5, near the pK_a of the NH_3 - NH_4^+ couple. In these experiments, we used a dry ice-acetone trap to remove most of the H_2O , thus greatly facilitating the sensitivity of detection at m/e = 17 (NH_3). We found no evidence evolution of NH_3 evolution during the NH_2NH_2 photooxidation process; any putative NH_3 production would be less than 1% of the observed N_2 production. The absence of significant NH_3 production under these conditions also seems to rule out the production of azide (N_3^-), a non-volatile compound not detectable with our experimental technique, since azide production is accompanied by NH_3 production during the NH_2NH_2 oxidation reaction (11).

Figure 2 shows the results obtained when we ran the same experiment in the presence of ^{15}N -labeled substrate. As shown in the top traces,

the major product appeared at $m/e = 30$, the spectral position of $^{15}\text{N}_2$, when the reaction was run with 95 atom % ^{15}N hydrazine. A small amount of $^{14,15}\text{N}_2$ at $m/e = 29$, consistent with the isotopic composition of the substrate, was also evolved.

The bottom traces of Fig. 2 show the results obtained when the same experiment was run in the presence of equimolar amounts of labeled and unlabeled substrate. These data suggest that little, if any, N interchange, manifest as isotopic scrambling, occurs during the production of molecular dinitrogen from hydrazine. If the nitrogen isotopes were to be completely randomized during the series of reactions leading to N_2 , one would predict that the $^{15}\text{N}_2: ^{14,15}\text{N}_2: ^{14}\text{N}_2$ ratio would be 0.23:0.50:0.27. In reality, quite the opposite was observed; the data of Fig. 2* indicate that less than 3% of the N atoms interchanged, suggesting that the N-N bond remains intact during the conversion of NH_2NH_2 to N_2 .

* We routinely observe that the 29/28 ratio of the N_2 evolved during oxidation of unlabeled NH_2NH_2 or NH_2OH is about a factor of two higher than one would predict from normal isotopic abundance. This problem can be corrected, albeit with a substantial loss of time response and signal-to-noise ratio, by interposing a dry ice-acetone trap between the inlet and the MS ionizer. This spurious signal probably reflects the protonation of N_2 within the MS; i.e., $\text{N}_2 + \text{H}^+ \rightarrow \text{N}_2\text{H}^+$ analogous to the well-known reaction

When ^{15}N -labeled NH_2NH_2 is photooxidized, a comparable H_2O -dependent signal is observed at $m/e = 31$ ($^{15}\text{N}_2\text{H}^+$).

The fact that N_2 was evolved on the first flash in all of these experiments indicates that, like NH_2OH , the oxidation of NH_2NH_2 involves the production of a one-electron photooxidation product. Subsequent (non-photochemical) interactions lead to the evolution of molecular N_2 .

Effect of DCMU

Table I is a compilation of the initial fluorescence yields observed after a saturating flash with different donors (H_2O , NH_2OH , and NH_2NH_2) in the presence and absence of DCMU. These results show that, in the presence of DCMU, NH_2NH_2 , like NH_2OH (see also Ref. 12), inhibits the fluorescence decay following a saturating flash (compared to the noninhibited case). This, in turn, suggests that NH_2NH_2 , like NH_2OH , inhibits the back reaction between $P680^+$ (the oxidized PSII donor) and Q^- (the reduced primary acceptor) and that the reduction of $P680^+$ by NH_2NH_2 is irreversible, in contrast to the reduction of $P680^+$ by the O_2 system (13).

Figure 3 shows a comparison of the N_2 evolution pattern elicited by one flash and four flashes in the presence of DCMU (added in complete darkness). A comparison of the two traces indicates that N_2 was evolved, but only on the first flash. A similar comparative experiment with NH_2OH yielded virtually identical results (data not shown), corroborating an earlier report (5). This indicates that, like NH_2OH , the oxidation of NH_2NH_2 is strictly a PSII reaction under these conditions.

The irreversibility of the NH_2NH_2 -DCMU system, like that of the NH_2OH -DCMU system, can thus be attributed to the ability of these donor

systems to immediately discharge oxidizing equivalents as N_2 . This contrasts with the normal DCMU-inhibited system, in which the oxidizing equivalents are retained (primarily as S_2) and can subsequently back react with Q^- .

Stoichiometry and the Involvement of O_2

Table II summarizes the results of a series of experiments in which the flash yields of N_2 evolution and O_2 uptake were determined as a function of acceptor and O_2 tension using NH_2NH_2 and NH_2OH as donors. Several salient points can be gleaned from these data:

- 1) At low O_2 tensions, the N_2 yield with NH_2NH_2 was about one-half that observed with NH_2OH . Since with NH_2OH the yield is $1/2$ N_2 per trap per flash (5), the data suggest that $1/4$ N_2 per trap per flash was evolved under low O_2 conditions.
- 2) At high O_2 tensions, the NH_2NH_2 N_2 flash yields were approximately double those observed at the low conditions, whereas the NH_2OH N_2 yields were unaffected by O_2 tension. This suggests that N_2 is formed by either the reaction of molecular O_2 with an NH_2NH_2 oxidation product or NH_2NH_2 with an O_2 reduction product.
- 3) O_2 uptake is strongly correlated with the production of N_2 from NH_2NH_2 (but not from NH_2OH). The (approximately) two-fold difference in N_2 evolution between low- O_2 and

high- O_2 conditions is associated with large differences in O_2 uptake in all three cases.

In sum, the data of Table II point to the participation of O_2 (or its reduction products) in some of the reactions leading to the production of N_2 from NH_2NH_2 . These secondary reactions probably underlie the unusual stoichiometry of these data (see also Ref. 1).

Hydrazine Oxidation Under High-Light Conditions

Figures 4 and 5 show the results obtained when N_2 evolution and O_2 uptake were monitored during hydrazine photooxidation under various conditions in continuous high light. These data, obtained using either Fd-NADP or MV as the terminal electron acceptor, show the rates as a function of NH_2NH_2 concentration under high O_2 conditions (> 1 mM, Figs. 4A and 5A) and as a function of O_2 tension (Figs. 4B and 5B). Experiments were done both in the presence and absence of saturating amounts of SOD.

One of the most striking aspects of these data is the large effect of SOD. In all cases (except zero O_2) the addition of SOD resulted in a large decrease in N_2 evolution and O_2 uptake. SOD also dramatically affected the shape of the substrate saturation curve. As shown in Figs. 4A and 5A, the rates of N_2 evolution and O_2 uptake saturated at ca. 30 mM NH_2NH_2 with both the Fd-NADP and MV acceptor systems in the presence of SOD; in the absence of added SOD there was no apparent substrate saturation up to 100 mM. These data suggest that superoxide can play a large role in N_2 evolution as well as O_2 uptake during hydrazine photooxidation. The addition of SOD seems to control side reactions so that N_2 evolution accurately reflects PSII activity.

The results obtained in the presence of SOD can be satisfactorily explained by the following set of reactions:

Equations 1-3 illustrate the (possibly idealized) production of molecular N_2 via the one-electron oxidation product of NH_2NH_2 . These reactions probably predominate under conditions of low NH_2NH_2 , low O_2 , and low light (Figs. 1 and 2, low O_2 point in Fig. 4B). The participation of diimine ($\text{NH} = \text{NH}$) in this reaction sequence is consistent with earlier reports (11). Diimine can be produced by the oxidation of NH_2NH_2 ; in the absence of other unsaturated species (e.g., olefins) it disproportionates.

Equations 4 and 5 illustrate the generation of superoxide (O_2^-) by PSI (see e.g. Ref. 14) and its breakdown by dismutation. The latter reaction can be greatly accelerated by the addition of SOD. The production of O_2^- by PSI is strongly dependent upon the terminal electron acceptor. In the presence of MV, O_2^- is efficiently generated at very high rates (equal to those set by limitations in electron transport) at rather modest O_2 tensions (15). It also formed, albeit at somewhat lower rates, in the presence of Fd (14). This reaction probably underlies

the somewhat anomalous results shown in Fig. 4B, which suggest that at O_2 tensions not much removed from zero the reduction of O_2 competes very favorably with that of NADP under the chosen reaction conditions.

In the presence of SOD, N_2 evolution depends on O_2 only to provide a terminal electron acceptor (via MV or Fd). The O_2 saturation kinetics of this PSI-mediated reaction are shown for the case of MV by the rapid rise of N_2 evolution in Fig. 5B (+ SOD curve). The equivalent N_2 curve for the Fd-NADP case (Fig. 4B) is masked by the apparent switch of Fd oxidation from NADP to O_2 at rather low O_2 tensions.

The PSII-mediated level of N_2 evolution with saturating NH_2NH_2 in the presence of SOD is about 100 $\mu\text{moles mg chl}^{-1} \text{ hr}^{-1}$ with Fd-NADP and about 175-200 $\mu\text{moles mg chl}^{-1} \text{ hr}^{-1}$ with MV. These differences probably reflect disparities in the efficiency of the two acceptor systems under the chosen assay conditions. In the presence of SOD, the N_2 evolution/ O_2 uptake stoichiometry was near unity in all cases in accordance with the reaction scheme

In the absence of SOD, the observed reactions are much more difficult to explain. Equations 6 and 7 illustrate two possibly relevant

secondary reactions by which N_2 production (and O_2 uptake) could be enhanced (see Refs. 16 and 17).

Equation 6 describes the attack of O_2 by the hydrazyl (oxidized hydrazine) radical, a reaction which competes with disproportion (Eqn. 2). Equation 7 describes the attack of NH_2NH_2 by O_2^- . Since this reaction competes with the reaction of Eqn. 5 for O_2^- , it can be effectively eliminated by the addition of large amounts of SOD. Note that if these reactions were to both proceed rapidly compared to the chain termination reactions (Eqns. 2 and 5), they would form the basis of an autocatalytic chain reaction sequence leading to the production of large amounts of N_2 per PSII turnover.

The data of Figs. 4 and 5, in particular the effect of SOD, indicate that the oxidation of NH_2NH_2 by O_2^- makes a substantial contribution to N_2 evolution (and O_2 uptake) in the absence of added SOD. The observed N_2 evolution rates with Fd-NADP are probably consistent with a scheme in which equal amounts of N_2 are generated by PSII (Eqns. 1-3) and PSI (via Eqns. 4, 7, and 1-3). A similar mechanism may also predominate with MV. The lack of any significant increase of N_2 evolution with increasing O_2 concentration suggests that the attack of O_2 by the hydrazyl radical (Eqn. 6) does not make a substantial contribution to N_2 evolution in these experiments.

We have no ready explanation for the anomalously high rate of O_2 uptake observed in the absence of SOD, particularly at high NH_2NH_2 and O_2 concentrations (Fig. 5A). These observations may reflect the participation of O_2^- in additional side (dark) reactions that result in the oxidation of NH_2NH_2 and the production of non-volatile products rather than molecular N_2 (11).

ACKNOWLEDGEMENTS

We thank A. Dudeck for technical assistance and J. Golbeck and B. Velthuys for critical readings of the manuscript. This work was supported by the Department of Energy, Contract DE-AC02-76ER03326, and the National Science Foundation, Grant PCM77-20526.

REFERENCES

- 1 Mantai, K. E. and Hind, G., (1971) *Plant Physiol.* 48, 5-8
- 2 Heath, R.L. (1971) *Biochim. Biophys. Acta* 245, 160-164
- 3 Diner, B. and Joliot, P. (1977) in *Encyclopedia of Plant Physiology*, Vol. 5, (Trebst, A. and Avron, M., ed.), pp. 187-205, Springer-Verlag, Berlin
- 4 Velthuys, B. (1980) *Ann. Rev. Plant Physiol.* 31, 545-567
- 5 Radmer, R. (1979) *Biochim. Biophys. Acta* 546, 418-425
- 6 Radmer, R. and Kok, B. (1976) *Plant Physiol.* 58, 336-340
- 7 Radmer, R. and Ollinger, O. (1980) in *Methods in Enzymology*, Vol. 69, Part C, pp. 547-560, Academic Press, New York
- 8 Schwartz, M. (1966) *Biochim. Biophys. Acta* 112, 204-212
- 9 Yamashita, I. and Butler, W. (1969) *Plant Physiol.* 44, 435-445
- 10 Radmer, R. and Ollinger, O. (1980) *FEBS Lett.* 110, 57-61
- 11 Raphaelian, L. (1966) in *Encyclopedia of Chemical Technology* (Kirk, R. and Othmer, D., eds.), pp. 164-196, Interscience Pub., New York
- 12 Bennoun, P. (1970) *Biochim. Biophys. Acta* 216, 357-363
- 13 Radmer, R. and Kok, B. (1975) *Ann. Rev. Biochem.* 44, 409-433
- 14 Allen, J.F. (1977) in *Superoxide and Superoxide Dismutases* (Michelson, A.M., McCord, J.M., and Fridovich, I., eds), pp. 417-436, Acad. Press, New York
- 15 Farrington, J.A., Ebert, M., Land, E.J. and Fletcher, K. (1973) *Biochim. Biophys. Acta* 314, 372-381

- 16 Jain, S.K. and Hochstein, P. (1979) Biochim. Biophys. Acta 586,
128-136
- 17 Misra, H.P. and Fridovich, I. (1976) Biochemistry 15, 681-687
- 18 Doschek, W. and Kok, B. (1972) Biophys. J. 12, 832-838

Table I. Initial Fluorescence Yield After a Saturating Flash with Different Donors in the Presence and Absence of DCMU

	<u>Fluorescence</u>
No added donor (H ₂ O oxidation)	0
Flash + 0.1 s dark	0.07
DCMU (10 ⁻⁵ M)	0.13
Flash + 2 s dark (+DCMU)	0.69
Flash + 10 s dark (+DCMU)	0.27
NH ₂ OH (2mM)	0
Flash + 0.1 s dark	0.06
DCMU (10 ⁻⁵ M)	0.41
Flash + 10 s dark (+DCMU)	0.77
Flash + 5 min dark (+DCMU)	0.50
NH ₂ NH ₂ (5mM)	0
Flash + 0.1 s dark	0.08
DCMU (10 ⁻⁵ M)	0.40
Flash + 10 s dark (+DCMU)	0.74
Flash + 5 min dark (+DCMU)	0.50

The tabulated fluorescence was calculated from the expression $(F_i - F_{i \text{ min}}) / (F_m - F_{i \text{ min}})$ where F_i = initial fluorescence level, $F_{i \text{ min}}$ = initial fluorescence level in dark adapted samples in the absence of DCMU, and F_m = maximum fluorescence level attained after prolonged illumination (> 10 s). The F_i 's in the three cases were identical; F_m 's varied by about 10%. Procedures were similar to those of Doschek and Kok (18). A 4 min dark period preceded each run.

The reaction mixture for each experiment contained 50 mM Tricine (pH 7.4), 5 mM MgCl₂, and chloroplasts (6 µg chl/ml) in addition to the reagents given in the table. The tabulated values for each of the three cases were obtained in the order shown using the same sample of unextracted, O₂-evolving, chloroplasts. Similar results with NH₂OH and NH₂NH₂ were obtained with Tris-extracted chloroplasts.

Table II. N₂ and O₂ Flash Yields as a Function of Acceptor and O₂ Tension

[O ₂]	Acceptor	<u>Hydrazine</u>		<u>Hydroxylamine</u>	
		N ₂ Evolution	O ₂ Uptake	N ₂ Evolution	O ₂ Uptake
Low	None	23	14	48	4
High	None	46	52	40	11
Low	NADP	21	10	42	1
High	NADP	33	36	40	2
Low	MV	18	11	42	5
High	MV	40	59	47	35

Tabulated values refer to relative peak height of the first flash after 5 min dark; the N₂ and O₂ flash yields are not directly comparable. Subsequent flash yields with NH₂OH did not change significantly (see ref. 5); those with NH₂NH₂ decreased 10-20% by the tenth flash.

O₂ levels: high, 32 to 47% (430 to 630 μM); low, 1.8 to 2.0% (24 to 27 μM)

Acceptor concentrations: 0.4 mM NADP plus 20 μg ferredoxin plus 1 mM MgCl₂;

Methylviologen (MV): 10⁻⁴ M.

FIGURE LEGENDS

Fig. 1. Relative flash yields at different m/e values observed when 10 μ l Tris - extracted chloroplasts (5.8 mg chl/ml) were subjected to a series of flashes (arrows) in the presence of 1 mM NH_2NH_2 . Numbers given after the m/e values refer to different attenuations; e.g. $m/e = 28$ was attenuated tenfold compared to $m/e = 30$. The different traces were sequentially obtained in various orders after a dark time of 5 min. See text and Ref. 5 for other details.

Fig. 2. Relative flash yields at different m/e values when 10 μ l Tris - extracted chloroplasts (6.1 mg chl/ml) were subjected to a series of flashes in the presence of 1 mM $^{15}\text{NH}_2^{15}\text{NH}_2$ (top) or an equivalent mixture (500 μ M each) of labeled and unlabeled NH_2NH_2 (bottom).

Fig. 3. Relative yields of N_2 elicited by one flash and four flashes (3 sec spacing) in the presence of NH_2NH_2 (1 mM) and DCMU (10^{-4} M).

Fig. 4. Rates of N_2 evolution and O_2 uptake in continuous high light as a function of NH_2NH_2 concentration and O_2 concentration with Fd-NADP as the electron acceptor. The experiments of Fig. 4A were all done at high O_2 (> 1 mM); those of 4B were all done using 30 mM NH_2NH_2 . In addition to substrate and O_2 the

reaction medium contained 50 mM Tricine (pH 7.4), 30 mM methylamine, 1 mM MgCl_2 , and 500 units catalase. Acceptor system: 40 μg Fd/ml, 2 mM NADP. Chloroplast concentration was equivalent to 50 μg chl/ml. Saturating amounts of SOD (1200 units/ml, determined using MV, see legend of Fig. 5) were added as noted in figure.

Fig. 5. Rates of N_2 evolution and O_2 uptake in continuous high light as a function of NH_2NH_2 concentration and O_2 concentration with MV (10^{-4} M) as acceptor. See legend of Fig. 4 for details. The dotted line in Fig. 5B illustrates an alternate interpretation of the data. The existence and significance of this apparent maximum was not studied in detail.

Fig 1

↑
EVOLUTION

m/e = 30

m/e = 29

m/e = 28

TIME →

Fig 2

Fig 3

A

B

A

B

Fig. 1