REPORT
to
DoE/ARM

DOE/ER/61062--T1

DE93 002638

by

C. Gautier, Paul Ricchiazzi, Pete Peterson and D. Lavallée
UC Santa Barbara

R. Frouin and D. Lubin
UC San Diego

S. Lovejoy (McGill U., Montreal, Canada)

and

D. Schertzer (Meteo. Nat., Paris, France)

May 6, 1992

Our group has been very active over the last year, analyzing a number of data sets to characterize multifractal cloud properties and assess the effects of clouds on surface radiation properties (spectral and broadband). The data sets analyzed include:

- AVHRR observations of clouds over the ocean
- SPOT observations of clouds over the ocean
- SSM/I observations of clouds over the ocean
- pyranometer data with all-sky photographs
- pyrgeometer data all-sky photographs
- spectral surface irradiance all-sky photographs

In addition, a number of radiative transfer computations have been performed to help in the interpretation of these observations or provide theoretical guidance for their analysis.

Finally a number of radiative transfer models have been acquired and tested to prepare for the interpretation of ARM/CART data.

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

402
MASTER

Multifractal Data Analysis and dynamic cloud modeling

McGill/Paris groups:

Radiation budget studies and remote sensing techniques depend heavily on clouds inhomogeneity, which turns out to be quite extreme over a wide range of scales. In the framework of multifractals, the relationship between cloud inhomogeneity and the internal scattered radiation field can be evaluated at all scales by establishing relationships between their respective orders of singularities and their associated codimensions. Recent work in this direction can be divided into empirical and theoretical categories.

a) Theoretical and numerical simulation

1) Exact radiative transfer results can be obtained when homogeneity is present only along the vertical: i) 1-D dimensional case or the so called "two stream approximations" where the conserved flux and total radiance are sufficient for a complete description ii) multifractal plane parallel atmosphere where a "K integral" is necessary to complete the description . When inhomogeneity is extreme along the horizontal as well as along the vertical (in 2-D cases) we have obtained some asymptotic results and more generally numerical results simplified radiative transfer models, which nevertheless are exact for the above cases i) and ii).

2) In the different cases studied, we find (with numerics and some theoretical arguments) that the radiation field shares a fundamental intermittency property with the concentration field, since the Levy index α_r of multifractality ($0 \leq \alpha_r \leq 2$: $\alpha_r = 0$ corresponds to the monofractal β -model, $\alpha_r = 2$ corresponds to the maximum of multifractality) is apparently preserved by the radiative transfer ($\alpha_r = \alpha_R$, R for "radiation"). However, the radiation field results from a non-linear integration of the concentration field and has distinguishing "dressed" properties such as divergence of moments.

3) We have theoretically considered the problem of multiple scattering in optically thick multifractal clouds, developing asymptotic expansions for single, double and n photon scattering statistics where the large parameter is the most probable optical thickness at the large scale. These calculations show how to quantify radiative "channeling" through optically thin regions.

b) Empirical results:

We performed (perhaps the first) systematic analysis of the scaling of energy spectra of satellite radiances over five wavelength channels and spanning the range of scales 160m to 4000km (the entire mesoscale). The study mostly involved 15 consecutive scenes of AVHRR data (1.1 km resolution, 512X512 pixels) taken over the same location at same local time in February 1986. This data was chosen because it was expected to provide a very sensitive indicator of the mesoscale break in the scaling predicted by the standard model of atmospheric dynamics (the "mesoscale gap"). Over the entire range, with surprisingly little scene to scene variation, the (isotropic) energy spectrum ($E(k)$) was found to follow the scaling form $E(k) \sim k^{-\beta}$ where k is a wavenumber, and β is the spectral exponent. This type of behavior is exactly as predicted by the unified scaling model of the atmosphere as the outcome of anisotropic nonlinear cascade dynamics. It is hard to see how these results can be reconciled with the standard model. These results indicate that the pertinence of using multifractal models of clouds to perform radiative transfer calculations over very wide ranges of scale. This is discussed in more details in Lovejoy et al., 1992.

Santa Barbara/McGill Groups

A study of the scaling properties of satellite data at different channels has been performed. For this purpose, we analyzed all 5 channels of the AVHRR sensor obtained from the NOAA-9 satellite. The universal parameters were estimated for 15 individual scenes and for all the scenes considered as whole (statistical average performed over 15 different samples). For the later, the values are listed in Table 1 below. Preliminary results have been presented at Fall AGU Meeting (Lavalley et al., 1991a) and at European Geophysical Society Meeting (Lavalley et al., 1992a, and Tessier et al., 1992a). Implications of these results will be discussed at AGU Spring Meeting (Tessier et al., 1992b).

Table 1

	α	C_1	H
Channel 1	1.15	0.1	0.41
Channel 2	1.10	0.09	0.41
Channel 3	1.11	0.07	0.31
Channel 4	1.45	0.1	0.52
Channel 5	1.32	0.1	0.50

The Lévy index α and the parameter C_1 have been evaluated using the double trace moments (DTM), the parameter H is deduced from the slope β of the energy spectra curve and the values of the parameters α and C_1 .

Furthermore, scaling properties of cloud liquid water derived from the SSM/I sensor aboard the DMSP-F8 have also been studied for three different samples. Cloud liquid water has been retrieved by a linear combination of several microwave measurements. These results will be presented at the Spring AGU (Lavalée et al., 1992b).

Another issue investigated is the anisotropic properties of the scaling. This has been addressed using the AVHRR data discussed above. However, in this case only channel 1 and 4 have been used until now, and the analysis have been performed for each of the 15 individual samples. For this purpose, the difference gray level has been estimated for different distances (distance between the two pixels used to estimate the difference) and different angles (angles between the orientation of the two pixels used to estimate the difference). The scaling properties can be studied by looking at the "straightness" of the curves of the statistical moments of the difference gray level as a function of distance on a log-log graph. The slope of these curves gives the values of scaling exponents. When the values of the scaling exponents is the same for different angles there is isotropic scaling; otherwise the scaling is anisotropic. Preliminary results indicate anisotropic scaling.

Effect of Broken Clouds on Surface Solar Irradiance

San Diego Group - R. Frouin

Solar irradiance data collected during the FIFE experiment over Kona Prairie have been analyzed in view of interpreting the observations in terms of broken clouds effects. Pyranometer measurements in the presence of cumulus clouds have revealed, when the sun was not obscured, substantially higher values than those expected on a clear day. Fig. 1 contrasts the situations on August 4 and August 8, 1989 at site 16 (39° 04'N and 96° 30'W). On August 8, cumulus clouds started to develop around 11:30 local time, were fully developed by early afternoon, and disappeared around 16:00 local time, which manifested in short periods of alternate high and low solar irradiance values. The high values largely exceeded those observed at the same time on August 4, a clear sky day, by as much as 130 Wm^{-2} between 13:00 and 14:00 local time.

In order to quantitatively assess the cloud effects we have run a 3-D radiative transfer model based on Monte Carlo model simulations. This model, developed by one of our colleague (Dr. F-M Breon), has been run with cumulus cloud field properties (cloud fractional coverage, optical thickness, aspect ratio and average cloud separation) characterized from all-sky camera and lidar observations. This cloud field has been assumed to move with the wind field at 4 ms^{-1} in the southward direction. The results from the computations are presented in Fig. 2. They clearly show the same tendencies as the observations : values higher than clear conditions at the edges of and in-between clouds and lower in the middle of clouds. The regularity in the changes of the insulation field comes from the regularity of the cloud field chosen.

Effects of clouds on spectral and broadband surface irradiance

Santa Barbara/San Diego Groups

As a result of work sponsored by the NSF, we have obtained surface radiation data from Palmer Station, Antarctica, located just north of the Antarctic Circle at 64 46' S, 64 04' W. We are in the process of analyzing this data to establish a correlation between stratospheric ozone abundance (e.g., from Total Ozone Mapping Spectrometer aboard Nimbus 7) and the surface

irradiance of PAR, UVA, and UVB, as part of our NSF project. However, an additional goal which encompasses our ARM objectives, is to use this information together with sophisticated radiative transfer methods and satellite measurements to more fully understand how clouds modulate the earth's radiation budget. We also want to investigate the scaling properties of these data sets.

The data consists of radiometric measurements in several different spectral bands for a ~60 day sequence during the austral spring. This time sequence represents a maximal variation of conditions, ranging from high surface albedo and low sun angle in the early spring to high solar angle and low albedo in the late spring. An accurate record of sky conditions is available from an all-sky video camera and 35mm photographs. Table 2 lists some of the spectral bands represented in the data set and a brief explanation of each. Measurements were taken every few minutes in daylight hours for the fixed bandwidth sensors.

Except for the TUVB, Eppley PSP and PIR all the fixed band sensors were contained within a Biospherical Instruments GTR-100 radiometer. A unique feature of this radiometer are the sensors covering the 580 - 680 nm and 730 - 1000 nm bands. These sensors were designed to closely match the spectral response of the AVHRR channel 1 and channel 2 sensors aboard the NOAA polar orbiting satellites. We expect these tailor made sensors will help us integrate the satellite and ground data into our radiative transfer models without introducing extra assumptions about the spectral shape of the radiation.

Clearly, the success of our analysis depends on an accurate cross-calibration of the ground and satellite AVHRR detectors. Biospherical Instruments provided us with a pre-deployment calibration of the AVHRR channel 1 sensor. However, because they lacked the necessary optical equipment appropriate for the near IR they were unable to do the same for the AVHRR channel 2 sensor. We were able to obtain some help from Santa Barbara Research Center, a local company specializing in satellite optical systems. Through an act of good will they provided us the use of their facilities and some help from their technical staff. From this effort we have determined post-deployment calibration constants for the channel 1 and 2 sensors.

Table 2

Fixed Bandwidth Detectors

936 nm	Narrow band sensor centered on the water vapor absorption band.
410 nm	Narrow band sensor placed at a wavelength maximizing atmospheric transmission.
560-680 nm	Wide band sensor with spectral response closely matching the satellite-based AVHRR channel 1 sensor.
730-1000 nm	Wide band sensor with spectral response closely matching the satellite based AVHRR channel 2 sensor.
280-2800 nm	Total solar irradiance (Eppley PSP).
5-50 μ m	Total IR sensor (Eppley PIR).
280-400 nm	Total UV radiometer.
400-700 nm	Total photosynthetically active radiation (PAR).

Scanning Spectrometers

5-20 μ m	Fourier transform IR spectrometer (FTIR)
280-700 nm	UV Spectrometer

Figure 3 shows data from day 251 for AVHRR channel 1 and the 936 nm channel. The solid lines are measurements from the GTR while the dotted lines indicate solar irradiance at the top of the atmosphere. The much greater attenuation of the 936 nm reading vs AVHRR channel 1 is a consequence of additional absorption due to atmospheric water vapor in the 936 nm band. We expect to use a comparison of these two channels to estimate the total column water vapor.

The conditions over Palmer Station on this day were overcast in the morning with partial clearing by noon. An interesting feature of the AVHRR channel 1 panel is that the surface irradiance achieves a value of about 1.4 times the extraterrestrial irradiance at local noon. This probably indicates detection of direct solar illumination plus some additional radiation scattered off of cloud edges. This example illustrates the importance of devising data evaluation algorithms which do not reject data points because they violate "common sense" notions of what the radiation values should be.

Also, after comparing data from this radiometer with one week's worth of AVHRR data over Antarctica, we have reason to believe that this radiometer can be used to provide a better radiometric calibration for the satellite sensors, under clear or completely overcast sky conditions. This is done in three steps: 1) measuring atmospheric transmission with the radiometer, 2) using a delta-Eddington (or any suitable) radiative transfer model to estimate cloud optical depth as a function of atmospheric transmission, and 3) using the same radiative transfer model to estimate planetary albedo (as would be measured by AVHRR) as a function of cloud optical depth. We have not yet analyzed enough satellite data to say for certain, but our results so far suggest that this method works if the solar zenith angle is less than 70 degrees. Our corrections derived from this method agree well with those derived by NASA/GISS from images of the Libyan desert. If the solar zenith angle is larger than 70 degrees, cloud bidirectional and edge effects confuse the issue. The advantage of using this AVHRR-specific radiometer instead of a total shortwave pyranometer (such as the common Eppley PSP) is that the uncertainty due to water vapor is eliminated. We repeated the above procedure using an Eppley PSP as the surface radiometer, and the results were not as promising. For partly cloudy skies, this radiometer (in conjunction with an all-sky camera) allows us to directly relate surface irradiance to satellite-measured planetary albedo.

Preliminary results from the analysis of these data will be presented at the 1992 Spring AGU meeting (Ricchiazzi et al., 1992).

Antarctic FTIR Spectroradiometer Data Reduction

Using some helpful suggestions from Dr. H. Revercomb (U. Wisconsin), an improved calibration algorithm was developed for the CalSpace FTIR Spectroradiometer measurements made at Palmer Station, Antarctica during August-November, 1991. This revised algorithm includes emission from the air column between the blackbody calibration source and the Bomem MB-100 Michelson interferometer, and includes an empirical correction for frost buildup on the blackbody when operating below the dew point. As of 4/15/91 a set of 360 Antarctic mid-IR emission spectra has been processed using the new algorithm. An average of four spectra were measured per day over a 90-day period at Palmer Station, and this data set represents a springtime longwave climatology for the region. The data set contains many clear-sky spectra which can be compared to FASCOD3P, but consists mostly of cloudy-sky spectra, as the Southern Ocean is one of the cloudiest regions on earth. Data analysis is therefore proceeding along the lines of computing longwave cloud forcing.

Figure 3 shows results from the FTIR for the days Sept. 7 through Sept. 10. Included in each plot are black body curves for 170, 220 and 270 K. We will use data of this type to gauge the temperature of the cloud bottoms. The FTIR radiance between 800 and 1200 cm^{-1} is particularly useful for this purpose because, except for an ozone absorption band at 1040 cm^{-1} , there is only some relatively weak H_2O absorption. We are able to infer the brightness temperature of the cloud bottom because the H_2O absorption in this band is only significant in a cloud layer. Combined with similar satellite measurements and an assumed atmospheric temperature profile we will determine the altitude of the top and bottom of the cloud layer (assuming there is only one cloud layer).

Figure 4 illustrates the procedure. Assuming a surface temperature near 273 K, Sept. 10 can be identified as a day with low clouds. That is, the cloud bottom temperature is very close to the surface temperature, implying low level clouds. By the same reasoning Sept. 7 and 8 must have had high clouds because the cloud bottom temperature is much less. The Sept. 9 plot

illustrates an IR spectral profile for clear conditions. Here spectral features due to CO₂ (650 cm⁻¹) and O₃ (1040 cm⁻¹) are clearly evident.

Preliminary results from this data analysis will be presented (Lubin, 1992) at the Spring AGU Meeting.

FIGURE CAPTIONS

- Fig. 1: Surface Solar Irradiance Observations Aug. 4 (clear) and Aug. 8 (cloudy).
- Fig. 2: Computed Surface Solar Irradiance for Aug. 8 using Monte Carlo Simulations.
- Fig. 3: Surface Solar Irradiance, Sept. 8. Dotted line indicates solar irradiance at the top of the atmosphere.
- Fig. 4: FTIR observations, Sept. 7 - Sept. 10. Blackbody curves are shown for $T=170, 200$ and 270 K.

REFERENCES

- Lavallée, D. Schertzer and S. Lovejoy, 1992a: On the quantification of multifractal properties in Geophysics. *Annales Geophysicae, supplément II, 10*, p. C337.
- Lavallée, D., C. Gautier, D. Jourdan, B. Anderson, S. Lovejoy, Y. Tessier, C. Hooge, D. Schertzer and F. Schmitt, 1992b: The quantification of the universal multifractal properties in Geophysics. *EOS, supplément April 7*, p. 58.
- Lavallée, D., Y. Tessier, S. Lovejoy and D. Schertzer, D. Jourdan, C. Gautier and B. Anderson, 1991a: Specific and nonspecific multifractal analysis techniques and satellite cloud radiance from 200m to 4000 km. *EOS, supplément october 29*, p.91.
- Lovejoy, S., D. Schertzer, P. Silas, Y. Tessier and D. Lavallée, 1992. The Unified scaling model of atmospheric dynamics and systematic analysis of scale invariance in cloud radiance. Prepare for *Annales Geophysicae*, 11/3/92
- Lubin, D., 1992: Measurements of atmospheric infrared emission by an FTIR spectroradiometer on the Antarctic peninsula. *EOS (supplement), April 7*, p. 70.
- Ricchiazzi, P., D. Lubin, C. Gautier, C. R. Booth: A Ground-Truthing Radiometer System for AVHRR Climate Studies. *EOS (supplement), April 7*, p. 71.
- Tessier, Y, P. Silas, F. Francis, S. Lovejoy, D. Lavallée, D. Jourdan, C. Gautier, B. Anderson and D. Schertzer, 1992c: Double trace moment and spectral analysis of cloud radiance over the range 160m to 4000km. *EOS (supplement), April 7*, p. 58.
- Tessier, Y, S. Lovejoy and D. Schertzer, 1992a: Universal multifractal exponents of in situ and remotely sensed rain and cloud fields. *Annales Geophysicae, supplément II, 10*, p. C340.

AUGUST 4 VS AUGUST 8, 1969 - SITE 16

PYRANOMETER MSMTS -- PARABOLA SITE

OBSERVATION

MODELLED

FIG 4

END

**DATE
FILMED**

12 / 22 / 92