Biological Conversion of Synthesis Gas Culture Development

Topical Report

**K.T. Klasson
R. Basu
E.R. Johnson
E.C. Clausen
J.L. Gaddy**

March 1992

Work Performed Under Contract No.: DE-FG21-90MC27225

For
U.S. Department of Energy
Office of Fossil Energy
Morgantown Energy Technology Center
Morgantown, West Virginia

By
University of Arkansas
Department of Chemical Engineering
Fayetteville, Arkansas

MASTER

262
DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

This report has been reproduced directly from the best available copy.

Available to DOE and DOE contractors from the Office of Scientific and Technical Information, P.O. Box 62, Oak Ridge, TN 37831; prices available from (615)576-8401, FTS 626-8401.

Available to the public from the National Technical Information Service, U.S. Department of Commerce, 5285 Port Royal Rd., Springfield, VA 22161.

Biological Conversion of Synthesis Gas Culture Development

Topical Report

**K.T. Klasson
R. Basu
E.R. Johnson
E.C. Clausen
J.L. Gaddy**

Work Performed Under Contract No.: DE-FG21-90MC27225

**For
U.S. Department of Energy
Office of Fossil Energy
Morgantown Energy Technology Center
P.O. Box 880
Morgantown, West Virginia 26507-0880**

**By
University of Arkansas
Department of Chemical Engineering
Fayetteville, Arkansas 72701**

March 1992

ACKNOWLEDGMENT

Financial support for this work has been provided by the Morgantown Energy Technology Center, U. S. Department of Energy under grant number DE-FG21-90MC27225.

TABLE OF CONTENTS

	<u>Page</u>
ACKNOWLEDGMENT	i
1.0 INTRODUCTION.....	1
1.1 Biological Water Gas Shift.....	1
1.2 Sulfur Gas Utilization.....	3
1.3 Purpose	5
2.0 EXPERIMENTAL SCREENING PROCEDURES.....	6
3.0 RESULTS AND DISCUSSION.....	11
3.1 Water Gas Shift Studies.....	11
3.1.1 H ₂ Production from Acetate.....	15
3.1.1.1 Bioreactor and Operating Conditions.....	16
3.1.1.2 Analytical Techniques.....	17
3.1.1.3 Results.....	17
3.2 H ₂ S Removal Studies.....	21
3.3 COS Degradation.....	28
3.3.1 COS Degradation by Selected CO-Utilizing Bacteria.....	28
3.3.1.1 CO Conversion by Bacterial Cultures.....	30
3.3.1.2 Materials and Methods.....	31
3.3.1.3 Results and Discussion.....	32
3.3.2 Indirect COS Utilization by Chlorobia.....	43
4.0 CONCLUSIONS.....	44
5.0 LITERATURE CITED.....	45

LIST OF TABLES

	<u>Page</u>
1.1 Bacteria Capable of Carrying Out the Water Gas Shift Reaction.....	2
1.2 Microorganisms Capable of Reducing H ₂ S to Elemental Sulfur.....	4
2.1 Composition of Solutions.....	7
3.1 Initial Results of Water Gas Shift Reaction Studies	12
3.2 Modified Media Compositions.....	14
3.3 Results of Water Gas Shift Reactions Studies Using a Modified Medium.....	14
3.4 Medium Composition for H ₂ Production from Acetate.....	16
3.5 Initial Screening of Growth Conditions of H ₂ S Reducers.....	29

LIST OF FIGURES

	<u>Page</u>
2.1 Cell Concentration Calibration Curve as a Function of Optical Density at 580 nm for <i>C. thiosulfatophilum</i>	9
2.2 Cell Concentration Calibration Curve (Based Upon Chlorophyll Content) as a Function of Optical Density at 670 nm for <i>C. thiosulfatophilum</i>	10
3.1 <i>R. rubrum</i> Grown in the Presence of Glutamic Acid.....	18
3.2 <i>R. rubrum</i> Grown in the Presence of Glutamic and Acetic Acids	18
3.3 <i>R. rubrum</i> Grown in the Presence of NH ₄ Cl.....	19
3.4 <i>R. rubrum</i> Grown in the Presence of NH ₄ Cl and Acetic Acid.....	19
3.5 <i>R. rubrum</i> Grown in the Presence of Urea.....	20
3.6 <i>R. rubrum</i> Grown in the Presence of Urea and Acetic Acid.....	20
3.7 <i>R. palustris</i> Grown in the Presence of Glutamic Acid.....	22
3.8 <i>R. palustris</i> Grown in the Presence of Glutamic and Acetic Acids.....	22
3.9 <i>R. palustris</i> Grown in the Presence of NH ₄ Cl.....	23
3.10 <i>R. palustris</i> Grown in the Presence of NH ₄ Cl and Acetic Acid.....	23
3.11 <i>R. palustris</i> Grown in the Presence of Urea.....	24
3.12 <i>R. palustris</i> Grown in the Presence of Urea and Acetic Acid.....	24
3.13 <i>R. gelatinosus</i> Grown in the Presence of Glutamic Acid.....	25
3.14 <i>R. gelatinosus</i> Grown in the Presence of Glutamic and Acetic Acids	25
3.15 <i>R. gelatinosus</i> Grown in the Presence of NH ₄ Cl.....	26
3.16 <i>R. gelatinosus</i> Grown in the Presence of NH ₄ Cl and Acetic Acid.....	26
3.17 <i>R. gelatinosus</i> Grown in the Presence of Urea.....	27
3.18 <i>R. gelatinosus</i> Grown in the Presence of Urea and Acetic Acid.....	27

	<u>Page</u>
3.19 CO Uptake by <i>P. productus</i> in the Presence of COS.....	33
3.20 CO Uptake by <i>E. limosum</i> in the Presence of COS.....	34
3.21 CO Uptake by <i>C. ljungdahlii</i> in the Presence of COS.....	35
3.22 CO Uptake by <i>R. rubrum</i> in the Presence of COS.....	36
3.23 COS Degradation by <i>P. productus</i> in Comparison to Medium Alone.....	37
3.24 COS Degradation by <i>E. limosum</i> in Comparison to Medium Alone	38
3.25 COS Degradation by <i>C. ljungdahlii</i> in Comparison to Medium Alone.....	39
3.26 COS Degradation by <i>R. rubrum</i> in Comparison to Medium Alone.....	40
3.27 Initial COS Utilization Rate as a Function of Initial CO Utilization Rate for <i>P. productus</i> , <i>E. limosum</i> and <i>R. rubrum</i>	42

1.0 INTRODUCTION

1.1 Biological Water Gas Shift

Since very little has been published concerning the biological conversion of CO and water to H₂ and CO₂ according to the reaction of Equation (1.1), the initial screening process was based mostly upon previous work conducted in the University of Arkansas laboratories.

In the past, two photosynthetic purple bacteria, *Rhodospirillum rubrum* and *Rhodopseudomonas gelatinosa* have been found to be capable of carrying out the water gas shift reaction (1). *R. rubrum* carries out the reaction both in the light and in the absence of light, but requires light for growth. *R. gelatinosa*, on the other hand, carries out the reaction in the dark and utilizes CO as a carbon source for growth. Since both of these organisms were not originally identified as H₂ producers in the literature, other bacteria with similar physiologies may also have H₂ producing capabilities. Thus, bacteria with the following characteristics were sought as possible water gas shift biocatalysts:

1. be a purple non-sulfur photosynthetic bacterium in the *Rhodospirillaceae* family;
2. be capable of anaerobic growth in the light;
3. have an optimal temperature of 25-40°C; and
4. show documented growth on simple substrates such as CO₂ or acetate.

A list of bacteria satisfying the above criteria is shown in Table 1.1 (2 3). Based upon this list, three organisms (one from each genus) were selected for further study in the laboratory. The choice of organisms was based upon the ability of the organisms to use simple acids as substrates and

Table 1.1

Bacteria Capable of Carrying Out the Biological
Water Gas Shift Reaction

ATCC #	Name	Optimum Temperature (°C)
<u>Rhodospirillum</u>		
25903	<u>rubrum</u>	30-35
25093	<u>tenuis</u> *	30
14031	<u>molischianum</u>	30
27871	<u>photometricum</u>	---
15798	<u>fulvum</u>	30
<u>Rhodopseudomonas</u>		
17001	<u>palustris</u>	30-37
19567	<u>viridis</u>	25-30
25092	<u>acidophilia</u>	25-30
17011	<u>gelatinosa</u> **	30
11166	<u>capsulata</u>	25-30
17023	<u>sphaeroides</u>	25-30
<u>Rhodomicrobium</u>		
17100	<u>vannielii</u>	30

* - Also named - Rhodocyclus tenuis

** - Also named - Rhodocyclus gelatinosus

optimal growth conditions. Together with *R. rubrum* and *R. gelatinosa* (see Table 1.1), a representative collection of bacteria from the Rhodospirillaceae family was thus obtained. Each of the bacterial cultures was obtained from the American Type Culture Collection (Rockville, MD).

1.2 Sulfur Gas Utilization

A number of microbial processes for the oxidation of H_2S to SO_4^{2-} or elemental sulfur have been reported in the literature. The sulfur bacteria comprise the phototropic and the chemolithotrophic bacteria, and the growth conditions may be anaerobic to aerobic. Based upon literature information (2,4), the list presented in Table 1.2 was compiled with the following considerations in mind:

1. the bacterium must be capable of anaerobic growth and sulfide reduction;
2. the bacterium must show elemental sulfur accumulation outside the cell;
3. the bacterium must have an optimal temperature of 25-40°C; and
4. the bacterium must have shown documented growth on simple substrates such as CO_2 or acetate.

Of the five choices shown in Table 1.2, only 2 were selected for further study at present: *Chlorobium thiosulfatophilum* and *C. phaeobacteroides*. *C. thiosulfatophilum* was obtained from ATCC (Rockville, MD) and *C. phaeobacteroides* was kindly supplied by Dr. T. Bergstein, a visiting scientist at Washington University, St. Louis, MO. *Ectothiorhodospira abdelmalekii* was found not to be suitable for sulfur gas conversion due to its alkaline growth conditions at pH 8.5 and high salinity (14-16%) requirements (5). *Thiobacillus denitrificans* has also been initially rejected, not being

phototrophic and mainly producing sulfate as the end product. *Chloroflexus aurantiacus* has not been selected for immediate study, but future plans may include studies of this organism. *C. aurantiacus* is capable of reducing S^{2-} to elemental sulfur and does not produce substantial amounts of sulfate. However, only 50% of the initial sulfur present in the medium was accounted for as elemental sulfur in one study (6).

Table 1.2
Microorganisms Capable of Reducing H_2S to Elemental Sulfur

ATCC #	Name
35917	<u>Ectothiorhodospira abdelmalekii</u>
	<u>Chlorobium</u>
17092	<u>thiosulfatophilum</u>
N/A	<u>phaeobacteroides</u>
29362	<u>Chloroflexus aurantiacus</u>
23642	<u>Thiobacillus denitificans</u>

In addition to the reduction of H_2S , the removal of carbonyl sulfide (COS), also present in the synthesis gas, is also desired. COS spontaneously undergoes a reaction with water to produce H_2S and CO_2 (7) as described by the equation:

Thus, the removal of COS may be seen as a removal of H₂S produced according to Equation (1.2). It has also been proposed that CO-utilizing microorganisms may utilize COS due to their molecular similarities. Three cultures, *Peptostreptococcus productus* (ATCC 35244), *Rhodospirillum rubrum* (ATCC 25903) and *Eubacterium limosum* (ATCC 8486) are presently being considered as possible COS-utilizing organisms. Both *P. productus* and *E. limosum* are anaerobic organisms capable of consuming CO and producing acetate. Acetate may also be used as the required carbon source for organisms carrying out the water gas shift reaction. *R. rubrum* is the obvious first choice since it is one of the most promising microorganisms for the water gas shift reaction studies. Other cultures may include selected H₂S-reducing cultures, since H₂S is a product in the reaction between COS and water (Equation 1.2).

1.3 Purpose

The overall objective of this task is to develop a system of bacterial cultures that is capable of simultaneously producing H₂ and removing sulfur compounds from synthesis gas. Cultures are being screened and compared for their ability to:

- a. convert CO and water to CO₂ and H₂ by the water gas shift reaction (H₂ production);
- b. remove H₂S from synthesis gas while producing elemental sulfur as the preferred by-product; and
- c. remove COS from synthesis gas by producing H₂S, elemental sulfur or other sulfur compound as a by-product.

2.0 EXPERIMENTAL SCREENING PROCEDURES

All experiments have been carried out using a basal medium containing per 100 mL: Pfennig mineral solution, 5 mL; Pfennig trace metal solution, 0.1 mL; B-vitamin solution, 0.5 mL; yeast extract (Difco), 0.05 or 0.1 g; and NaHCO_3 , 0.25 or 0.4 g (see Table 2.1 for solution compositions). All ingredients (except for NaHCO_3) were mixed with distilled water, heated and boiled for 2 min under anaerobic conditions. After cooling and the addition of NaHCO_3 , the medium was dispensed into serum bottles (100 or 150 mL nominal volume). The bottles were capped with butyl rubber stoppers and aluminum crimp seals. Anaerobic conditions were maintained throughout the preparation by working under an atmosphere of N_2/CO_2 (75%/25%) or H_2/CO_2 (75%/25%). Half-filled bottles were steam-sterilized at 15 psig for 20 min.

If strict anaerobicity was necessary, 2 mL of reducing agent solution (2.5% $\text{Na}_2\text{S}\cdot 9\text{H}_2\text{O}$ or cysteine-HCl $\cdot\text{H}_2\text{O}$) were added per 100 mL of medium before inoculation. In addition to above medium, this basal medium was supplemented with specific carbon substrates or growth factors as the various cultures required. A 5% inoculum was typically employed in these batch screening experiments and the pH was maintained close to 7.0.

In monitoring the progress of the experiments, gas phase analyses were performed using gas chromatography, liquid phase sulfide concentrations were monitored with a sulfide probe, cell density was measured by optical density on a spectrophotometer and pH was measured by employing a Corning pH meter. Liquid and gas samples, 1.5 mL and 0.4 mL, respectively, were withdrawn from the reactors during the fermentation and analyzed for cell concentration and gas phase composition. The dry cell weight concentration for Rhodospirillaceae were obtained by measuring light scatter at 540 nm in a

Table 2.1

Composition of Solutions

<u>Pfennig Mineral Solution (5)</u>	<u>g/L</u>
KH ₂ PO ₄	10.0
MgCl ₂ ·6H ₂ O	6.6
NaCl	8.0
NH ₄ Cl	8.0
CaCl ₂ ·2H ₂ O	1.0
 <u>Pfennig Trace Metal Solution (5.6)</u>	 <u>g/L</u>
ZnSO ₄ ·7H ₂ O	0.1
MnCl ₂ ·4H ₂ O	0.03
H ₃ BO ₃	0.3
CoCl ₂ ·6H ₂ O	0.2
CuCl ₂ ·H ₂ O	0.01
NiCl ₂ ·6H ₂ O	0.02
Na ₂ MoO ₄ ·2H ₂ O	0.03
FeCl ₂ ·4H ₂ O	1.5
Na ₂ SeO ₃	0.01
 <u>B-Vitamin Solution (6)</u>	 <u>mg/L</u>
Biotin	20
Folic acid	20
Pyridoxal HCl	10
Lipoic acid	60
Riboflavin	50
Thiamine HCl	50
Ca-D-pantothenate	50
Cyanocobalamin	50
P-aminobenzoic acid	50
Nicotinic acid	50

Spectronic 21 (Milton Roy Co., Rochester, NY) and converting it to cell density using a calibration curve ($DCW(mg/l) = 395.4 \cdot ABS_{540}$).

The presence of elemental sulfur crystals makes measurement of cell concentration by optical density difficult. Two methods have thus been used for measuring the cell concentration of *C. thiosulfatophilum*: direct o.d. measurement at 580 nm and o.d. measurement of the pigment chlorophyll at 670 nm. For the measurement of pigment, 1 mL of solution was first centrifuged at 15,000 rpm for 4 min. The supernatant water was then discarded. Methanol (1 mL) was then added to dissolve the chlorophyll present in the cells, followed by centrifugation for an additional 2 min. Methanol (1.5 mL) was then added to the new supernatant and the absorbance measured at 670 nm.

A calibration curve for cell concentration in mg/L as a function of optical density at 580 nm is shown in Figure 2.1 and a calibration for cell concentration based upon chlorophyll content in mg/L as a function of optical density at 670 nm is shown in Figure 2.2. As is noted, both calibrations yielded straight lines. These calibration curves were constructed at a late stage in the fermentation when all of the elemental sulfur had been converted to sulfate. Thus, no interference of sulfur on the direct cell concentration measurements at 580 nm was observed.

Gas phase CO, CO₂ and H₂ analyses were performed on a gas chromatograph (Perkin-Elmer Sigma 300, Norwalk, CT) using an 1.8 m stainless steel column packed with Carbosphere, 60/80 mesh. The oven temperature was held constant at 135°C while the injector and thermal conductivity detector temperatures were 175°C. Helium was used as the carrier gas at a flow rate of 40 mL/min.

Figure 2.1 Cell Concentration Calibration Curve as a Function of Optical Density at 580 nm for *C. thiosulfatophilum*.

Figure 2.2 Cell Concentration Calibration Curve (Based upon Chlorophyll Content) as a Function of Optical Density at 670 nm for *C. thiosulfatophilum*.

The gas phase H_2S and COs concentrations were measured by gas chromatography using a Hewlett-Packard (Kennett Square, PA) HP 5890 Series II gas chromatograph and HP 7673 integrator equipped with a thermal conductivity detector. The column was a 1/8" x 6' teflon column packed with Chromosorb 107, 80/100 mesh (Altech Assoc., Inc., Deerfield, IL). The oven temperature was 80°C, and the detector and injector temperatures were kept at 175°C. the carrier gas was helium at a flow rate of 30 mL/min. Liquid phase sulfur concentrations ($H_2S(l)$, HS^- , S^{2-}) were measured using a Corning Silver/Sulfide Electrode (Corning Glass Works, Medfield, MA) and an Orion Specific Ion meter (Orion Research, Inc., Cambridge, MA).

3.0 RESULTS AND DISCUSSION

3.1 Water Gas Shift Studies

The initial water gas shift work concentrated on growing lyophilized cells in simple basal medium with the addition of acetate and biocarbonate under a H_2/CO_2 atmosphere (75%/25%) by successive transfer. After good growth had repeatedly been obtained in this medium, small amounts of CO were added to the gas phase at a partial pressure of approximately 0.16 atm. If hydrogen was produced in these initial studies, a transfer was made to identical medium and CO was again added after growth was observed.

The results of these initial experiments with the four Rhodospirillaceae are shown in Table 3.1. As is noted, all of the cultures except *R. gelatinosa* were identified as H_2 -producers under the conditions employed. All of the organisms showed good growth on acetate, with H_2 production the most prominent so far in cultures reduced with cysteine-HCl. So far the most promising microorganisms are *R. palustris*, together with previously studied *R. rubrum*. *R. rubrum* has previously been shown to grow and completely convert all CO

Table 3.1

Initial Results of Water Gas Shift Reaction Studies

<u>Culture</u>	<u>Reducing Agent</u>	<u>Growth</u>	<u>H₂ Production</u>	<u>Microscope Observation</u>
<u>R. tenuis</u>	Na ₂ S	++	ns	Thin rod-shaped, form large dense mats and clusters
	Cysteine-HCl	++	+	
	None	++	ns	
<u>R. palustris</u>	Na ₂ S	+++	ns	Thin crooked rod shaped, form small star-like clusters
	Cysteine-HCl	+++	+++	
	None	+++	+++	
<u>R. gelatinosa</u>	Na ₂ S	++	ns	Thin rod shaped, non-descriptive
	Cysteine-HCl	++	--	
	None	++	ns	
<u>R. vannielii</u>	Na ₂ S	+	ns	Lemon shaped, flagellated, form chains and aggregates
	Cysteine-HCl	++	++	
	None	++	ns	

ns - not studied

- - negative test (no H₂ production)

+ - positive results; ++ more positive; +++ highly positive results

present in the gas phase to hydrogen at initial CO partial pressures up to 2 atm (10).

Continued studies, however, where gas phase compositions and stoichiometric relationships were closely followed, showed that even though substantial amounts of H₂ were produced, CO was not consumed. These experiments were conducted in triplicate.

Based upon documented research conducted with *R. rubrum* (11), it is likely that the observed H₂ production was based on a nitrogen system which liberates H₂ from organic substrates (such as acetate) under ammonia or N₂ limitations. This theory is in agreement with the results of the present study, where H₂ production was initiated after substantial growth had been observed. It is likely that cell growth was accompanied by a limitation of ammonia and with the subsequent onset of H₂ production.

More recent experiments were directed toward utilizing a modified medium (see Table 3.2). In this medium, the yeast extract content was reduced and NH₄Cl was added to inhibit the formation of H₂ from acetate. The results from these experiments showed that no H₂ production occurred in the modified medium (see Table 3.3). Good growth was observed for *R. palustris* and slight growth was observed for *R. vaniellii* and *R. tenue*. The weak growth observed with the two latter strains is possibly related to the reduction of yeast extract in the new medium.

acclimating the cultures to H₂ production/CO consumption. In addition, a small effort has been focused on the apparent production of H₂ from organic substrates such as acetate. Acetate may be seen as a liquid intermediate which may be produced from CO by cultures such as *Peptostreptococcus productus* (4), according to Equation (3.1):

Table 3.2
Modified Media Compositions

	Original Medium per 100 mL	Modified Medium per 100 mL
Pfennig Minerals Solution	5 mL	5 mL
Pfennig Trace Metal Solution	0.1 mL	0.1 mL
B-vitamins Solution	0.5 mL	0.5 mL
Yeast Extract	0.1 g	0.005 g
NaHCO ₃	0.25 g	0.25 g
NH ₄ Cl	--	0.27 g

For solutions composition and medium preparation, see reference (2)

Table 3.3
Results of Water Gas Shift Reaction Studies
Using a Modified Medium

<u>Culture</u>	<u>Growth</u>	<u>H₂ Production</u>
<i>R. tenuis</i>	++	--
<i>R. palustris</i>	+++	--
<i>R. vaniellii</i>	+	--

- negative test (no H₂ production)
+ positive results, ++ more positive; +++ highly positive results

The produced acetate could then be converted to H₂ by several members of the Rhodospirillaceae family.

3.1.1 H₂ Production from Acetate

Several members of the Rhodospirillaceae family have been investigated for possible production of H₂ from acetate. The acetate may be seen as a liquid intermediate which may be produced by e.g. *P. productus*. The production of H₂ from a variety of organic substrates has previously been shown in the literature (13-15). The supply of nitrogen and chemical configuration were shown to strongly affect H₂ production in these cases. Since it is generally considered that the enzyme nitrogenase is responsible for the release of H₂ in photosynthetic bacteria (16-18), experiments were conducted with various nitrogen sources according to the following test pattern:

Cultures: *Rhodospirillum rubrum* (ATCC 25903)
Rhodopseudomonas palustris (ATCC 17001)
Rhodocyclus gelatinosus
 (*Rhodopseudomonas gelatinosa*)(ATCC 17001)

Nitrogen Source: Glutamic acid, 0.106% (w/v)
 NH₄Cl, 0.038%
 Urea, 0.021% (base nitrogen content the same)

Carbon Source: None other than CO₂
 Sodium acetate 0.576%

Each of the above organisms were evaluated for H₂ production using a certain nitrogen source with or without acetate addition. For additional medium constituents and preparation see Table 3.4 and reference (19).

Table 3.4

Medium Composition, for H₂ Production
from Acetate per 100 mL

Minerals Solution	5 mL
Pfennig Trace Metals Solution	0.1 mL
B-vitamins Solution	0.5 mL
Yeast Extract	0.02g
NaHCO ₃	0.25g
Nitrogen Source	as indicated in text
Sodium Acetate	as indicated in text

The minerals solution contained 10.0 g/L KH₂PO₄, 6.6 g/L MgCl₂·6H₂O, 8.0 g/L NaCl, and 1.0 g/L CaCl₂ 2H₂O. For other solutions see reference (10)

3.1.1.1 Bioreactor and Operating Conditions

Batch fermentations were carried out in 150 mL glass serum bottles (Wheaton Glass Co., Millville, NJ). The bottles were sealed using butyl rubber septum stoppers and aluminum crimp seals (Bellco Glass Co., Vineland, NJ). Before sterilization, 75 mL of medium were added to each bottle under a He/CO₂ (75%/25%) atmosphere. The bottles were autoclaved at 15 psig for 20 minutes.

Once sterile, the bottles were inoculated with 5 mL of seed culture. The seed cultures had been previously growing in conditions similar to the experiment before they were used. Argon (20 mL) was then added to each bottle as an inert tracer gas with a gas-tight syringe. This procedure allowed calculation of the amounts of H₂ through the fermentation.

The experiments were carried out in a modified incubator shaker (Model G25, New Brunswick Scientific Co., Edison, NJ) at 150 rpm and 30°C. Tungsten light, necessary for growth, was supplied in excess in the shaker.

3.1.1.2 Analytical Techniques

Liquid and gas samples, 2.5 mL and 0.4 mL, respectively, were withdrawn from the reactors during the fermentation and analyzed for cell concentration and gas phase composition. Cell growth was obtained by measuring light scatter at 540 nm in a Spectronic 21 (Milton Roy Co., Rochester, NY).

Gas analyses were performed on a gas chromatograph (Perkin-Elmer Sigma 300, Norwalk, CT) using an 1.8 m stainless steel column packed with Carbosphere, 60/80 mesh. The oven temperature was held constant at 135°C while the injector and thermal conductivity detector temperatures were 175°C. Helium was used as the carrier gas at a flow rate of 40 mL/min. The acetate concentration was analyzed with a Varian (Walnut Creek, CA) model GC-3400 gas chromatograph and a model DS-604 data station. The GC was equipped with a 3 mm x 1.8 m, 100/200 mesh, Poropak QS (Alltech, Deerfield, IL) column and a flame ionization detector. The oven temperature was maintained at 190°C, the detector and injector temperatures were both 240°C, and the carrier gas was helium at a flow rate of 40 mL/min.

3.1.1.3 Results

The results of the acetate fermentations conducted with *R. rubrum* are displayed in Figures 3.1-3.6. As noted in the figures, the growth (OD₅₄₀) increased sharply and then leveled off towards the end of the fermentation. The highest optical density reached was 1.2 in the medium supplemented with NH₄Cl and sodium acetate (or acetic acid) (see Figure 3.4). The only H₂ production with this culture was observed in the medium supplemented with glutamic and acetic acids (Figure 3.2). The amount produced was 0.4 mmol after 200 hrs, but decreased to 0.2 mmol after 500 hrs. Apparently the produced H₂ was being consumed during the latter part of the fermentation.

Figure 3.1 *R. rubrum* grown in the presence of glutamic acid.

Figure 3.2 *R. rubrum* grown in the presence of glutamic and acetic acids.

Figure 3.3 *R. rubrum* grown in the presence of NH_4Cl .

Figure 3.4 *R. rubrum* grown in the presence of NH_4Cl and acetic acid.

Figure 3.5 *R. rubrum* grown in the presence of urea.

Figure 3.6 *R. rubrum* grown in the presence of urea and acetic acid.

In Figures 3.7-3.12 the results for the fermentations using *R. palustris* are displayed. In general, the medium supplemented with acetate supported the best growth. In addition, the culture in the medium with only glutamic acid grew quite well. The highest optical densities ranged from 1.2-1.4. The production of H₂ was observed in several cases (see Figures 3.7, 3.8, 3.10, 3.12) with the most promising results for the experiment with glutamic and acetic acids (Figure 3.8). In this experiment the culture produced 1.4 mmol of H₂ in 200 hrs and showed no signs of H₂ disappearance. As is noted in Figure 3.8, growth (OD), H₂ production and acetate consumption almost ceased after 200 hrs of fermentation.

The experiments conducted with *R. gelatinosus* showed similar patterns for growth and H₂ production as with the other two purple bacteria. The results are shown in Figures 3.13-3.18. As with *R. palustris*, the best growth with *R. gelatinosus* was obtained in medium supplemented with glutamic and/or acetic acids. The typical growth obtained was an OD of 1.3-1.7. The highest H₂ production, 0.9 mmol, occurred in the medium with glutamic and acetic acids after 150 hrs (see Figure 3.14). This production level remained constant throughout the rest of the fermentation.

To summarize, all of the cultures produced H₂ most efficiently in the medium with glutamic and acetic acids. The largest H₂ production, 1.4 mmol, was observed using *R. palustris*. H₂ production rates for this case are presented later in this report.

3.2 H₂S Removal Studies

The two most likely bacterial candidates capable of converting H₂S to elemental sulfur have been investigated for use in a co-culture with *R. rubrum*. These two bacteria, *Chlorobium thiosulfatophilum* and *Chlorobium*

Figure 3.7 *R. palustris* grown in the presence of glutamic acid.

Figure 3.8 *R. palustris* grown in the presence of glutamic and acetic acids.

Figure 3.9 *R. palustris* grown in the presence of NH_4Cl .

Figure 3.10 *R. palustris* grown in the presence of NH_4Cl and acetic acid.

Figure 3.11 *R. palustris* grown in the presence of urea.

Figure 3.12 *R. palustris* grown in the presence of urea and acetic acid.

Figure 3.13 *R. gelatinosus* grown in the presence of glutamic acid.

Figure 3.14 *R. gelatinosus* grown in the presence of glutamic and acetic acids.

Figure 3.15 *R. gelatinosus* grown in the presence of NH_4Cl .

Figure 3.16 *R. gelatinosus* grown in the presence of NH_4Cl and acetic acid.

Figure 3.17 *R. gelatinosus* grown in the presence of urea.

Figure 3.18 *R. gelatinosus* grown in the presence of urea and acetic acids.

phaeobacteroides, are both members of the green sulfur photosynthetic bacteria family. Both bacteria were subjected to parametric studies covering growth and H₂S uptake in order to select the best bacterium for further study.

The conditions studied and overall results of these studies are shown in Table 3.5. As is noted in the table, both bacteria are very similar in behavior for the conditions studied. Both grew well in basal medium with or without yeast extract present. Hydrogen consumption was noted for both bacteria during the latter part of the fermentation as the H₂S (added as Na₂S) was depleted from the medium. Only *C. thiosulfatophilum* has been tested so far for the utilization of acetate (a substrate for *R. rubrum*) or acetate plus NH₄Cl. *C. thiosulfatophilum* was not inhibited by acetate and, in fact, grew on acetate as a carbon source. The bacterium also grew on basal medium supplemented with NH₄Cl and acetate.

Later studies with the two bacteria have shown that *C. phaeobacteroides* often fails to grow on basal medium, while *C. thiosulfatophilum* always grows in basal medium. Also, *C. phaeobacteroides* will thus be chosen as the bacterium for H₂S conversion to elemental sulfur.

3.3 COS Degradation

3.3.1 COS Degradation by Selected CO-Utilizing Bacteria

Synthesis gas, a major product of coal gasification, typically contains 1-2 sulfur by volume, with 95-99% or more of the sulfur present as H₂S (20,21). Typical levels of carbonyl sulfide (COS) in coal-derived synthesis gas range from 0.03-0.07% by volume (22), although COS concentrations have been observed in the 5-10% range (20,21).

Table 3.5

Initial Screening of Growth Conditions of H₂S Reducers

<u>Condition Studied</u>	<u>C. thiosulfatophilum</u>	<u>C. phaeobacteroides</u>
Is yeast extract needed for growth?	No	No
Does it grow in basal medium	Yes	Yes
Is acetate used?	Yes	--
Does acetate inhibit growth?	No	--
Does it use H ₂ as reducing agent?	Yes	Yes
Does it grow in basal medium supplemented with NH ₄ Cl and acetate?	Yes	--

A dash indicates that the condition has not yet been studied.

Although COS is present in only small quantities, it poses serious problems to equipment and the environment. COS is corrosive to both iron and steel (23), and is a precursor to the formation of sulfur oxide derivatives, which are highly regulated environmental pollutants. COS also poses significant problems to downstream catalysts as a catalyst poison. Examples of catalysts for downstream processing of synthesis gas to methane include nickel and potassium-based catalysts, as well as mixtures of iron/chromium oxides and zinc/copper oxides (24,25). Poisons for these catalysts include chlorine and sulfur (26,27).

It has been recently shown in the University of Arkansas laboratories that some co-utilizing bacteria are capable of degrading COS (28). The results of these initial tests with various co-utilizing bacteria comparing their ability to degrade COS in the presence of CO are shown in this report. Optimum conditions for each culture were employed as noted in the literature or determined in the laboratory. The respective rates at which COS was degraded by each culture were compared to the rate of chemical reaction of COS with water:

The gas phase composition was held constant in comparing the different cultures.

3.3.1.1 CO-Conversion by Bacterial Cultures

Four CO-utilizing bacteria were chosen to be tested for their ability to degrade COS:

1. *Peptostreptococcus productus*
2. *Eubacterium limosum*
3. *Clostridium ljungdahlii*
4. *Rhodospirillum rubrum*

Of the above, *C. ljungdahlii*, *P. productus*, and *E. limosum* convert CO to acetate by the equation

P. productus was isolated from anaerobic sewage-digester sludge and has been reported to have a doubling time of 1.5 h at 37°C and pH 7 (12). *E. limosum* exhibited a doubling time of 7 h under the same conditions (29). *P. productus* has been shown to be able to tolerate H₂S and COS concentrations as high as 20% by volume (30).

C. ljungdahlii was isolated from chicken waste at the University of Arkansas (31) and, in addition to Reaction 2, carries out the following reaction (32):

The CO conversion to ethanol is favored at low pH and under nutrient limitation.

R. rubrum, as mentioned previously, is a photosynthetic bacterium that is capable of converting CO and H₂O to HCO₃⁻, H⁺, and H₂ by the following reaction (33):

R. rubrum does not utilize CO for growth, but is able to utilize a wide variety of carbon sources, including acetate, malate, and yeast extract. Tungsten light is required for growth, but is not required for CO utilization (34). *R. rubrum* was shown to be inhibited by COS concentrations above 6% (30).

3.3.1.2 Materials and Methods

Peptostreptococcus productus, Strain U-1, was supplied by M. P. Bryant, University of Illinois, Department of Dairy Science. *Eubacterium limosum*,

Strain 8486, and *Rhodospirillum rubrum*, Strain 25903, were obtained from the American Type Culture Collection, Rockville, MD. *Clostridium ljungdahlii*, Strain PETC, was isolated from chicken waste in the University of Arkansas laboratories.

The medium for the bacteria was the same as the original medium of Table 3.2 with the addition of 0.1 mL of 0.1% resazurin. The medium pH was 7.0 for *P. productus*, *E. limosum*, and *R. rubrum*, and 5.0 for *C. ljungdahlii*. The experimental techniques employed in these batch comparison studies were as described previously (19,28).

3.3.1.3 Results and Discussion

The experiments were designed so that CO and COS served as cosubstrates for the organisms. The utilization of CO in the presence of COS by the four bacteria is shown in Figures 3.19-3.22. The disappearance of CO in the medium alone is also shown for comparison purposes. As is noted in the figures, the rate of CO consumption increased with inoculum size. Furthermore, all organisms yielded essentially complete conversion for all inoculum sizes in 25 h or less. CO was not degraded in the medium alone.

The uptake of COS in the presence of CO by the bacteria is shown in Figures 3.23-3.26. *P. productus*, *E. limosum*, and *R. rubrum* showed an ability to degrade COS faster than the reaction of COS with water in the medium. *C. ljungdahlii*, on the other hand, was not shown to utilize or degrade COS any faster than medium alone. It should be pointed out that the COS profiles with medium alone were run at identical pH levels and temperatures as the CO-COS degradation experiments. Thus, for experiments with *P. productus* and *E. limosum*, a medium at pH 7 and 37°C was used, and for experiments with *C. ljungdahlii* a medium at pH 5 and 37°C was used. A medium at pH 7 and 30°C was

Figure 3.19 CO uptake by *P. productus* in the presence of COS.

Figure 3.20 CO uptake by *E. limosum* in the presence of COS.

Figure 3.21 CO uptake by *C. ljungdahliae* in the presence of COS.

Figure 3.22 CO uptake by *R. rubrum* in the presence of COS.

Figure 3.23 COS degradation by *P. productus* in comparison to the medium alone.

Figure 3.24 COS degradation by *E. limosum* in comparison to the medium alone.

Figure 3.25 COS degradation by *C. ljungdahlii* in comparison to the medium alone.

Figure 3.26 COS degradation by *R. rubrum* in comparison to the medium alone.

used for experiments with *R. rubrum*.

In analyzing the results presented in Figures 3.19-3.26 it was noted from the slopes of the curves that the rate of COS degradation for *P. productus* and *E. limosum* tended to decrease when the concentrations of both CO and COS were very low. It is possible that the rate of COS degradation could exhibit a corresponding decrease. If the difference in initial rates of COS degradation with cells and in medium alone is plotted as a function of the difference in CO uptake rates by the cells and in medium alone for *P. productus*, *E. limosum*, and *R. rubrum*, a nearly linear trend is observed (Figure 3.27). At least-squares fit forced through the origin yields a slope of 0.055 mol COS/mol CO. This ratio of initial rates may be compared to the initial amounts of gases present, which were approximately 0.06 mol COS/mol CO in all experiments. This would indicate similar kinetic utilization modes for both CO and COS, with COS possibly being "used" in place of CO. However, since the initial ratios (mol COS/mol COS) were kept constant in all experiments and isotope labeling was not employed, this conclusion is speculative.

The results of this study support the hypothesis that several organisms that are capable of utilizing CO are also capable of degrading CO-like compounds, such as COS. Three organisms, *Peptostreptococcus productus*, *Rhodospirillum rubrum*, and *Eubacterium limosum*, were shown to metabolize COS at rates much higher than that of the reaction of COS with H₂O. *Clostridium ljungdahlii*, on the other hand, was not found to have an ability to degrade COS. H₂S was detected as a major gas-phase product, but stoichiometric product yields have not yet been determined.

Figure 3.27 Initial COS-utilization rate as a function of initial CO-utilization rate for *P. productus*, *E. limosum* and *R. rubrum*.

COS degradation by *C. ljungdahlii* (see Figure 3.25) may have been limited by the low operating pH (pH 5). In comparing the rates of COS degradation by the medium alone, it seen that the rate (slope of curve) at pH 5 is much lower than the rate at pH 7. Perhaps *C. ljungdahlii* would have shown the ability to degrade COS if an initial pH of 7.0 were utilized.

In any case, it has been demonstrated that COS can be degraded by some CO-utilizing organisms faster than in medium alone. Particularly effective was *R. rubrum*, which completely degraded the added COS in <20 h for cell concentrations above 55 mg/L. Possible mechanisms for degradation include, but are not limited to, dual substrate utilization of both COS and CO. Studies to elucidate the reaction mechanism and microbial kinetics are in progress.

It was observed during the COS degradation studies that H₂S was a major product formed in the gas phase for each of the three bacteria capable of degrading COS (data not shown). No other gas-phase products were formed, and liquid-phase products were not detected or measured. Product yields in verifying process stoichiometry are under study.

3.3.2 Indirect COS Utilization by Chlorobia

COS undergoes a chemical reaction with water to produce H₂S and CO₂ by the equation:

Thompson *et al.* (7) studied the reaction and showed that the kinetics of the reaction could be described by a first order irreversible rate expression over a temperature range of 15-40°C. They showed that the rate constant varied with temperature by the equation:

$$k = 1.06 \times 10^{12} e^{-22710/RT} \quad (3.5)$$

where k is the first order rate constant, sec^{-1} ;

R is the ideal gas constant, $1.987 \text{ cal/gmole}^\circ\text{C}$; and

T is the absolute temperature, $^\circ\text{K}$.

At 37°C , the rate constant is thus calculated as 0.37 h^{-1} .

C. thiosulfatophilum can utilize COS indirectly, by utilizing the H_2S produced by the chemical reaction in Equation (3.2). If this is indeed a satisfactory process, it will eliminate the need for an additional organism (such as *Rhodospirillum rubrum* or *Peptostreptococcus productus*) for COS degradation.

4.0 CONCLUSIONS

Based upon the results of this culture screening study, *R. rubrum* is recommended for biocatalysis of the water gas shift reaction and *C. thiosulfatophilum* is recommended for H_2S conversion to elemental sulfur. Both bacteria require tungsten light for growth and can be co-cultured together if H_2S conversion is not complete (required concentration of at least 1 ppm), thereby presenting H_2 uptake by *C. thiosulfatophilum*.

COS degradation may be accomplished by utilizing various CO-utilizing bacteria or by indirectly converting COS to elemental sulfur after the COS first undergoes reaction to H_2S in water. The second alternative is probably preferred due to the low expected concentration of COS relative to H_2S .

5.0 LITERATURE CITED

1. M. P. Dashkevich and R. L. Uffen, "Identification of a Carbon Monoxide-Metabolizing Bacterium as a Strain of Rhodopseudomonas gelatinosa (Molisch) van Niel," *Int. J. Sys. Bacteriol.*, 29(2), 145 (1979).
2. American Type Culture Collection, "Catalog of Bacteria and Phages," 17th ed., American Type Culture Collection, Rockville, MD (1989).
3. "Bergey's Manual of Determinative Bacteriology," 8th ed., Eds., R. E. Buchanan and N. E. Gibbons, The Williams & Williams Company, pp. 26 (1974).
4. "Bergey's Manual of Determinative Bacteriology," 8th ed., Eds., R. E. Buchanan and N. E. Gibbons, The Williams & Williams Company, pp. 35, 51, 456 (1974).
5. J. F. Imhoff and H. G. Trüper, "Ectothiorhodospira abdelmalekii sp. nov., a New Halophilic and Alkaliphilic Phototropic Bacterium," *Zbl. Bakt. Hyg., I. Abt. Orig. C2*, 228 (1981).
6. M. T. Madigan and T. D. Brock, "Photosynthetic Sulfide Oxidation by Chloroflexus aurantiacus, a Filamentous, Photosynthetic, Gliding Bacterium," *J. Bacteriol.*, 122, 782 (1975).
7. H. W. Thompson, C. F. Kearton and S. A. Lamb, "The Kinetics of the Reaction Between Carbonyl Sulfide and Water," *J. Chem. Soc.*, 1033 (1935).
8. M. J. McInerney, M. P. Bryant and N. Pfennig, "Anaerobic Bacterium that Degrades Fatty Acids in Syntrophic Association with Methanogens," *Arch. Microbiol.*, 122, 129 (1979).
9. B. R. S. Genther, M. P. Davis and M. P. Bryant, "Features of Rumen and Sewage Sludge Strains of Eubacterium limosum, a Methanol- and H₂-CO₂-Utilizing Species," *Appl. Environ. Microbiol.*, 42, 12 (1981).
10. K. T. Klasson et al., "Advanced Studies of the Biological Conversion of Coal Synthesis Gas to Methane," Final Report for Contract No. DE-AC21-86MC23281 for Morgantown Energy Technology Center, p. 136, November 1989.
11. P. M. Vignais, A. Colbeau, J. C. Willison, and Y. Jouanneau, *Advances in Microbial Physiology*, 26, 155 (1985).
12. W. H. Lorowitz and M. P. Bryant, *Appl. Environ. Microbiol.*, 47, 961 (1984).
13. H. Gest and M. D. Kamen, *J. Bacteriol.*, 58, 239 (1949).

14. H. Gest and M. D. Kamen, *Science*, 109, 558 (1949).
15. J. M. Siegel and M. D. Kamen, *J. Bacteriol.*, 61, 215 (1951).
16. P. Hillmer and H. Gest, *J. Bacteriol.*, 129, 732 (1977).
17. K. Watanabe, J. S. Kim, K. Ho, L. Buranakarl, T. Kampee, and H. Takahashi, *Agric. Biol. Chem.*, 45, 217 (1981).
18. B. A. MacIer, R. A. Felroy, and J. A. Bassham, *J. Bacteriol.*, 138, 446 (1979).
19. M. D. Ackerson, E. C. Clausen, and J. L. Gaddy, "Biological Conversion of Synthesis Gas," Quarterly Report No. 1-1, U. S. Department of Energy, Grant No. DE-FG21-90MC27225 (September-December 1990).
20. A. G. Eickmeyer and H. A. Gangriwala, *Energy Progress*, 1, 9-12, (1981).
21. L. S. Galstaun and R. F. Geosits, *Energy Prog.* 5, 156-160, (1985).
22. A. L. Kohl and F. C. Riesenfeld, *Gas Purification*, 4th ed., Gulf Publishing, Houston, TX (1985).
23. M. Caillet and A. Galerie, *Ceram.* 30, 171-194 (1980).
24. C. A. Euker, Jr., and R. D. Wessenhott, *Energy Prog.* 1, 12-16 (1981).
25. L. J. Kaplan, *Chem. Eng.*, 89(6), 64-66 (1982).
26. J. T. Richardson, *Hydro. Proc.* 52(1), 91-95 (1973).
27. H. Jockel and B. E. Tribskorn, *Hydro. Proc.* 52(1), 93-98 (1973).
28. K. D. Smith, K. T. Klasson, M. D. Ackerson, E. C. Clausen, and J. L. Gaddy, *Appl. Biochem. Biotechnol.* 28/29, 787-796 (1991).
29. B. R. S. Genthner and M. P. Bryant, *Appl. Environ. Microbiol.* 43, 70-74 (1983).
30. J. L. Vega, K. T. Klasson, D. E. Kimmel, E. C. Clausen, and J. L. Gaddy, *Appl. Biochem. Biotechnol.* 24/25, 329-340 (1990).
31. S. Barik, S. Prieto, S. B. Harrison, E. C. Clausen, and J. L. Gaddy, *Appl. Biochem. Biotechnol.* 17/18, 363-378 (1988).

32. J. L. Vega, S. Prieto, B. B. Elmore, E. C. Clausen, and J. L. Gaddy,
Appl. Biochem. Biotechnol. 20/21, 781-797 (1989).
33. L. M. Brown, MS thesis, University of Arkansas, Fayetteville, AR (1987).
34. K. T. Klasson, J. P. Cowger, C. W. Ko, J. L. Vega, E. C. Clausen, and J.
L. Gaddy, *Appl. Biochem Biotechnol.* 24/25, 317-328 (1990).

END

**DATE
FILMED**

9 / 14 / 92