CARBON DIOXIDE RESEARCH PROGRAM

Dr. Park S. Nobel

Open-top chambers at University of California at Riverside

(A)

(B)

(C)

Gas exchange measurements in open-top chambers at UCR (A), in walk-in growth rooms at UCLA (B), and in Conviron growth chambers at UCLA (C)

MASTER

CO₂ Exchange, Environmental Productivity Indices, and
Productivity of Opuntia ficus-indica under Current and Elevated
CO₂ Concentrations

Principal Investigator - Dr. Park S. Nobel

	Page
Project 1 PHYSIOLOGY - Dr. Muyi Cui	2
Project 2 PARTITIONING - Dr. Yiqi Luo	5
Project 3 ROOTS - Dr. Patsy Miller	10
Project 4 MORPHOLOGY - Dr. Yiqi Lou	15
Project 5 ABSORPTANCE - Dr. Muyi Cui	18
Project 6 NUTRIENTS AND SECONDARY COMPOUNDS - Dr. Patsy Miller	21
Project 7 PRODUCTIVITY - Dr. Park Nobel	23

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

Project 1 PHYSIOLOGY - Gas Exchange of Opuntia ficus-indica
Exposed to Elevated CO₂

Objective: Determine the short and long term responses of photosynthesis and stomatal conductance to elevated CO₂ for mother cladodes and newly formed daughter cladodes. Specifically, CO₂ and H₂O flux rates, g^s, and diel patterns for developing daughter cladodes of the same size (or age) will be monitored.

Location: The research will be conducted in three open-top chambers, L (ambient CO₂), M (+150 ppm CO₂), and H (+350 ppm CO₂), located at the University of California Riverside research garden.

Plant material: Five replicates of two cohorts will be used from 12 planted cladodes in each chambers. Plants assigned for the measurements are B5, B6, B7, C5, C6, C7, J5, J6, J7, K5, K6, and K7 (Figure 1).

Measurements: Gas exchange will be measured using a LiCor 6200 portable system. Total number of measurements per hour: 1 cohorts x 5 replicates x 3 chambers = 15. Total number of days required to finish one round of measurement for 2 cohorts = 2 days.

Schedule: First phase - measurements on planted cladodes. Gas exchange measurements will be conducted on the same planted cladodes (total of 5 replicates) for 24 hours beginning the day after the cladodes are planted in wet soil

(day 0) and will be measured 0, 14, 30, 45, 60, 90, and 120 days after planting.

Second phase - measurements on daughter cladodes. J_{CO_2} and J_{H_2O} will be measured continuously for 24 hours 0, 3, 7, 14, 30, 45, 60, 90, and 120 days after newly initiated Spring 1992 daughter cladodes reach a measurable size. Five replicate daughter cladodes will be measured in the L, M, and H chambers.

Third phase - measurements on two cohorts. Gas exchange measurements will start when Spring 1993 daughter cladodes are appearing on the Spring 1992 cladodes (in spring, 1993). The measurements will be conducted as close to 0, 3, 7, 14, 30, 60, 90, 120 days after the initiation of daughter cladodes as possible. Five replicate sets of Spring 1992 cladodes and their Spring 1993 daughter cladodes will be measured in the L, M, and H, chambers.

Fourth phase - follow-up study. If, after the first year of study, great differences are observed for plants exposed to different levels of CO_2 , gas exchange will also be conducted during the second year on plants subjected to other environmental stresses, such as shading, water stress and high temperature.

Results expected: 1) Diel J_{CO_2} and total carbon fixation vs days after exposure to elevated CO_2 for mother cladodes
2) WUE and g^s vs days after exposure for mother cladodes

- 3) diel J_{CO_2} and total carbon fixation vs age of newly appeared daughter cladodes
- 4) WUE and g^s vs days after exposure for newly appeared daughter cladodes

Project 2 PARTITIONING - Effects of elevated CO₂ on carbon
partitioning of Opuntia ficus-indica

Objectives: 1) Study how CO₂ concentration affects root/shoot ratio and carbon partitioning of the CAM plants; 2) investigate relevant physiology such as net CO₂ exchange and nonstructural carbohydrate.

Growth room experiment

Location: Two existing walk in growth rooms at the Laboratory of Biomedical and Environmental Sciences, UCLA (23-172 A and 23-172 B), will be used. One room will be maintained at ambient levels of carbon dioxide and one at 350 ppm above ambient concentrations. The floor space in each room is 3.25 m x 2.85 m = 9.26 m².

Plant material: A total of 88 similar sized cladodes from 1 ascension of O. ficus-indica from Mexico (either 1282, 1296, or 1300, which are the ascensions that will be planted in the center m² of the open top chambers) will be collected from the UCR plantation. Eight cladodes (4 replicates x 2 sampling times) will be planted in small pots with a area of ~ 160 cm², 28 cladodes in medium pots with an area of ~440 cm² (4 replicated x 7 sampling times), and 8 cladodes in large pots with an area of ~1610 cm² (4 replicates x 2 sampling times) for a total of 44 cladodes in each walk in growth room, one room with ambient CO₂ and one with CO₂ at +350 ppm above ambient.

Space requirement per room: If the pot sizes are as described above, 3.5 m^2 is needed in each growth room. The three sizes of pots will not be arranged randomly, but will be arranged to maximize the space available in the growth room.

Pre-planting Measurements: Before planting the length, width, thickness at 4 locations around edge of cladode, and fresh weight will be measured and the surface area traced to establish the initial condition of the mother cladode.

Growth medium: Plants will be planted in a mixture of 50% soil from the University of California at Riverside research garden and 50% sandbox sand and will be watered every other day with 1/10 strength Hoagland's solution.

Measurements: 1) Dry mass of roots and chlorenchyma and water storage parenchyma tissues of shoots in the medium size pots will be measured at day 7, 14, 30 or 45, 60, 75, and 90.

2) Plants in the small and large size pots will be harvested at day 30/45 and 90.

3) Net gas exchange (NCE) and total non-structural carbohydrates (TNC) will be measured on day 30/45 and 90 on cladodes growing in all three sized pots under elevated and ambient levels of CO_2 (4 replicates x 3 pot sizes x 2 CO_2 levels x 2 sampling dates = 48 samples).

Expected results: 1) Time course of root/shoot ratio in the medium size pots in two CO_2 environments, 2) root/shoot

ratio, NEC, and TNC for three pot sizes at two harvest times under two CO₂ conditions.

Follow-up experiment: Root/shoot ratios and carbon partitioning will be measured with two carbon dioxide levels, two light levels and three drought periods in the UCLA walk in growth rooms.

Field experiment

Location: This study will be conducted in the L (ambient CO₂), M (+150 ppm CO₂), and H (+350 ppm CO₂) open top chambers at the University of California Riverside research gardens.

Plant material: Cladodes were selected from the edge columns and rows in each chamber, columns A and L and rows 1 and 8 (Table 1). Four cladodes from the L, M, and H open top chambers will be labeled at each sampling period. Cladodes A2, B1, K1, and L2 will be sampled in each chamber 14 days after the cladodes are planted in wet soil; cladodes A3, A4, L3 and L4 sampled on day 30; cladodes G1, H1, I1, and J1 sampled on day 60; and cladodes C1, D1, E1, and F1 sampled on day 90 (Table 1).

Schedule: Labeling with ¹⁴CO₂ will be done at midnight on day 12, 28, 58, and 88, after planting.

Measurements: 1) Samples of cladode tissue will be collected with a cork borer 6 times one day after labeling and 3 times two days after labeling.

2) Carbon dioxide exchange (4 replicates x 1 cohort x 3 chambers = 12 samples per hr) and TNC may be measured on day 30/45 and 90.

Root/shoot ratio: After all other measurements are completed, above and below-ground tissue of each cladode will be dried and weighted to calculate root/shoot ratios. After a cladode is removed from a open topped chamber to determine root biomass and root/shoot ratios, a replacement cladode from the plantation at UCR will be planted in its place.

Expected results: Time course of root/shoot ratio under three CO₂ conditions. Carbon partitioning patterns at four times and NCE and TNC at two times under three CO₂ concentrations.

Table 1. Location of cladodes to be sampled for ^{14}C and root biomass 7, 14, 21, 30, 45, 60, 75, and 90 days after planting in the L (ambient carbon dioxide), M (+150 ppm carbon dioxide), and H (+350 ppm carbon dioxide) open top chambers at the University of California Riverside research garden.

Day	Chamber	Cladode	Measurements
7	L	A5, A6, L5, L6	R/S ratio
	M	A5, A6, L5, L6	Root morphology
	H	A5, A6, L5, L6	
14	L	A3, A4, L3, L4	^{14}C translocation
	M	A3, A4, L3, L4	R/S ratio, Root morphology
	H	A3, A4, L3, L4	
21	L	A7, B8, L7, K8	R/S ratio
	M	A7, B8, L7, K8	Root morphology
	H	A7, B8, L7, K8	
30	L	A2, B1, K1, L2	^{14}C translocation
	M	A2, B1, K1, L2	NCE, TNC
	H	A2, B1, K1, L2	R/S ratio, Root morphology
45	L	C8, D8, E8, F8	R/S ratio
	M	C8, D8, E8, F8	Root morphology
	H	C8, D8, E8, F8	
60	L	G1, H1, I1, J1	^{14}C translocation
	M	G1, H1, I1, J1	R/S ratio
	H	G1, H1, I1, J1	Root morphology
75	L	G8, H8, I8, J8	R/S ratio
	M	G8, H8, I8, J8	Root morphology
	H	G8, H8, I8, J8	
90	L	C1, D1, E1, F1	^{14}C translocation
	M	C1, D1, E1, F1	NCE, TNC
	H	C1, D1, E1, F1	R/S ratio, Root morphology

Project 3 ROOTS - Root initiation and development with elevated CO₂, shading, and root pruning

Objectives: 1) Understand effects of elevated carbon dioxide on establishment of O. ficus-indica by quantifying root initiation on newly planted cladodes, determining costs of root construction, and documenting changes in morphology and anatomy; and 2) to measure the effects of reducing photosynthetic photon flux density (PPFD) and root pruning on root production of O. ficus-indica growing with three levels of carbon dioxide.

Growth Room Experiment

Project 3 will utilize the same location, plant material, and pots described for Project 2 - PARTITIONING

Sampling: This project will focus on cladodes growing in the medium sized pots. Four randomly chosen cladodes will be harvested 7, 14, 30, 45, 60, 75, and 90 days after planting. Remaining plants will be consolidated after each harvest to reallocate space in the growth room for additional experiments.

Measurements: 1) At each harvest the number of roots originating from the base and from the areoles will be counted to determine if elevated carbon dioxide affects location of root initiation.

2) The branching pattern of the root mass in each pot will be described.

- 3) Roots will be divided into 3 classes, i.e., main roots, second order, and third order roots. As the root mass increases, only random sub-samples of the root mass will be separated into size classes.
- 4) While the root mass is of a manageable size, the length and area of all roots will be measured. As roots become more abundant, sub-sampling will be used by measuring randomly chosen sections that add up to 10 cm of root length; 5 replicated 10 cm lengths of each class of roots will be measured.
- 5) Sub-samples and the total root mass will be dried and weighed.
- 6) At day 30/45 and 90, nitrogen concentration, ash, and caloric content will be measured on four replicates of the three root size classes from cladodes growing in the medium size pots in each carbon dioxide treatment (4 replicates x 3 root sizes x 2 CO₂ levels x 3 pot sizes x 2 sampling periods = 144 samples).
- 7) The distance from the root tip to where suberization and lignification occurs will be measured on a sub-sample of root tissue using staining techniques.
- 8) The number of periderm layers at various distances from the root tip will be measured on sub-samples of root tissue to determine where cortex cells are shed.

Field Experiments

Root Initiation

Location: This study will be conducted in the A (- chamber ambient CO₂), L (ambient CO₂), M (+150 ppm CO₂), H (+350 ppm CO₂) open top chambers and the F (- chamber +350 ppm CO₂) area at the University of California Riverside research garden.

Plant material: Cladodes were selected from the edge columns and rows in each chamber, columns A and L and rows 1 and 8 (Figure 1).

Sampling schedule: Cladodes will be sampled for root biomass 7, 14, 21, 30, 45, 60, 75, and 90 days after planting (Table 1). The complete root system of each cladode will be traced out and excavated by hand (4 replicates x 5 chambers x 8 periods = 160).

Measurements: Parameters measured on the recovered roots will be the same as those described previously for day 30/45 for roots recovered from cladodes growing the medium sized pots in the walk in growth room at UCLA.

PPFD Reduction and Root Pruning

Location: The research will be carried out in the A, L, M, H, and F areas/chambers at UCR.

Plant material: In each chamber cladodes A3, A4, A5, A6, L3, L4, L5, and L6 will be used for the light reduction experiment; cladodes A2, A7, B1, B8, K1, K8, L2 and L7 will be used for

the root pruning experiment. Cladodes G1, H1, I1, and J1 will be the controls for both experiments.

Manipulations: Light reduction - After cladodes from locations A5, A6, L5 and L6 are harvested replacement cladodes will be planted on day 7 and shaded to reduce PPFD by 50%. Sixty days after planting these cladodes will be harvested, on day 67 (4 replicates x 5 chambers = 20 samples). After cladodes from locations A3, A4, L3, and L4 are harvested on day 14, replacement cladodes will be planted on day 14 and shaded to reduce PPFD by 90%. Sixty days after planting these cladodes will be harvested, on day 75, (4 replicates x 5 chambers = 20 samples). Cladodes G1, H1, I1, and J1 harvested on day 60 will be the controls for the two shading treatments.

Root reduction - After cladodes from locations A7, B8, L7, and K8 are harvested on day 21, replacement cladodes will be planted on day 21. For the following 60 days, root growth will be restricted to 50% of that measured at day 7, 14, 21, 30, 45, and 60. Sixty days after planting, on day 81, these cladodes will be harvested (4 replicates x 5 chambers = 20 samples). After cladodes from locations A2, B1, K1, and L2 are harvested on day 30, replacement cladodes will be planted on day 30. For the following 60 days, root growth will be restricted to 10% of that measured at day 7, 14, 21, 30, 45, and 60. Sixty days after planting, on day 90, these cladodes will be harvested (4 replicates x 5 chambers = 20

samples). Cladodes G1, H1, I1 J1 harvested on day 60 will be the controls for the two root reduction treatments.

Measurements: Parameters measured on the recovered roots will be the same as those described previously for day 30/45 for roots recovered from cladodes growing the medium sized pots in the walk in growth room at UCLA.

Project 4 MORPHOLOGY - Kinematics of cladode expansion

Preliminary experiment in growth rooms

Objective: learn how to measure movement trajectory of areoles, stained dots, and trajectory of cell and stomatal cells from areoles and to measure cell growth velocity; practice mathematical manipulation.

Plant Material: The cladodes use in growth room experiment of Project 2 - PARTITIONING will also be used for this project.

Measurements: (1) measure movement of stained dots and areoles, plot movement trajectory; (2) count cell number, measure cell length, cell growth velocity; (3) stomatal cell differentiation and flow with cladode expansion.

Experiment in the growth rooms

Objective: quantitative description of both the displacement and the expansion of cladode tissue elements, including lateral meristem activity, cell density gradient, cell length, stomata flow.

Location: two existing walk in growth rooms at the Laboratory of Biomedical and Environmental Sciences, UCLA (23-172A and 23-172B), will be used. One room will be maintained at ambient level of carbon dioxide and one at 350 ppm above ambient concentration.

Schedule: This experiment will begin after the growth room experiments for Projects 2 and 3 are completed.

Plant material: Forty-eight similar sized cladodes from 1 ascension of O. ficus-indica from Mexico (either 1228, 1296, or 1300) will be used (2 CO₂ levels X 2 light levels X 3 drought periods X 4 replicates = 48 plants). Cladodes will be planted in the medium size pots used in Project 2, in a mixture of 50% soil from the University of California at Riverside research garden and 50% sandbox sand and will be watered every other day with 1/10 strength Hoagland's solution for control.

Treatments: 1) Light levels will be full growth room light and 50% reduction in growth room light. 2) The three drought periods will be normal watering, withhold watering so water potential drops below -0.5 MPa for 10 days, and withhold watering so water potential drops below -0.5 MPa for 20 days. 3) Two CO₂ levels, ambient and +350 ppm CO₂.

Measurements and Analysis Methods: trace aerole movement to get movement trajectory, trace cell movement from areoles and then describe the trajectory with the fluid flow equation such as:

$$\frac{Da}{Dt} = \frac{\partial a}{\partial t} + v \cdot \nabla a \quad (1)$$

where

a is a growth field variable, t is time, and v is growth velocity (rate of displacement from a material origin in the plant).

Equation means the material rate of change (cell specific) equals

the rate of change at the position instantaneously occupied by the tissue element, plus the rate of change due to displacement to a new position.

Measure cell length (or cell density), in the case of one-dimensional growth, then

$$\text{Rate of cell division} = \frac{1}{l} \left[\frac{\partial u}{\partial t} - u \frac{\partial l}{\partial x} \right]$$

where l is cell length, u is growth rate, x is position.

Expected results: Trajectory of areole movement, strain rates, local production rates, cell division rates, and cell elongation rates at two CO_2 , water, and light conditions.

Project 5 ABSORPTANCE - Light absorption and reflection in
Opuntia ficus-indica exposed to elevated CO₂

Objectives: To determine light absorptance and reflectance for whole cladodes and tissue sections across the chlorenchyma in O. ficus-indica. The relationship between tissue structure and its functions in light absorption and CO₂ fixation will be investigated.

Location: This project will be carried out in the A (ambient CO₂ - chamber), L (ambient CO₂ + chamber), M (+150 ppm CO₂), and H (350 ppm CO₂) chambers/locations at the UCR research gradens.

Plant material: Planted cladodes D7, E7, F7, G7, H7 and I7 in each location will be reserved for this project. Ten replicate samples from Spring 1992 cladodes will be obtained on each sampling day.

Schedule: Experiments will start when a sufficient number of Spring 1992 cladodes reach a width of 5 cm, or two weeks after daughter cladode initiation. Ten replicated samples (3.12 cm²) will be obtained from cladodes of similar size in each chamber/location on days 14, 30, 45 and after 2, 3, 5, 7, and 10 months. Five samples will be used for light measurements and five for chlorophyll content and acidity.

Measurements: 1) The thickness of the cuticle, cpidermis, chlorenchyma, and water storage parenchyma will be measured using a stereo microscope and a bright field microscope.

Publishable photos will be taken using a scanning electron microscope.

2) Reflectance will be measured using a LI-1800 integrating sphere with a LI-1820 photo sensor at 450, 550, 680, and 860 nm and scanned from 300 nm to 1000 nm.

3) Tissue absorptance and transmittance through the chlorenchyma will be determined on the rest of the tissue using the radiospectrometer by repeated removing 1.0 mm sections from the bottom of the tissue. After each 1.0 mm section is removed absorptance and transmittance will be remeasured until all mesophyll tissues are removed. Forward transmitting, forward scattering, and backward scattering light across the intact tissues will also be scanned with a fiber optic probe at 450, 550, 680, and 860 nm.

4) Transmittance and reflectance by cuticle and epidermis layers, which have been obtained by removing the mesophyll tissue with a razor blade, will be determined at 450, 550, 680, and 860 nm.

5) Chlorophyll contents and acidity will be measured on 1.0 mm thick peridermal sections of the chlorenchyma tissue.

Expected results: 1) Absorptance, transmittance, and reflectance vs carbon dioxide

2) Thickness of the cuticle, CHL, and WSP

3) Chlorophyll and chloroplast distribution across the CHL

4) Acid contents across the CHL vs CO₂

5) Light distribution across CHL

6) Combining the effects of CO₂ and light distribution of
assess the adaptive features of O.ficus-indica.

Project 6 NUTRIENTS AND SECONDARY COMPOUNDS - Effects of Elevated Carbon Dioxide on macro- and micro-nutrient concentrations and precursors of secondary quantitative compounds in Opuntia ficus-indica

Objective: Determine the effect of elevated carbon dioxide on macro- and micro-nutrient concentrations and on secondary compounds in O. ficus-indica grown under various levels of carbon dioxide.

Location: This study will be carried out in the A (ambient CO₂ - chamber) L (ambient CO₂ +chamber), M (+150 ppm CO₂), and H (+350 ppm CO₂) chambers/locations at the UCR research gardens.

Plant material: This project will utilize Spring 1992 cladodes from planted cladodes C2, D2, E2, F2, G2, H2, I2, and J2. Measurements will be made when the daughter cladodes are about 10 cm in length. Cladode tissue will be separated into cuticle, chlornychema, and water storage paranychema.

Measurements: 1) nitrogen concentration (5 replicated x 5 treatments = 25 samples) using micro Kjeldahl digestion.
2) macro- and micro-nutrient concentrations (5 replicates x 5 treatments = 25 samples) using DC arc optical emission spectrometry, 50 mg of tissue
3) caloric content (5 replicates, 3-4 internal replicated may be required due to variability of the small samples, x 5

treatments = 25 samples, but 75 - 100 measurements) using a Gentry bomb calorimeter, 1 gram of tissue

4) mucilage content using methods described in Goldstein and Nobel 1991, 4 grams of tissue (13 mm cork borer)

5) isoprenoids/terpenoids/essential oils (for methods see pp 107-116 of Harborne - Gas liquid chromatography and thin layer chromatography)

7) phenolic compounds (for methods see Ragan and Jensen 1977)

8) tanin (for methods see page 84-88 of Harborne)

Project 7 Productivity - The productivity of Opuntia ficus-indica growing will elevated concentrations of carbon dioxide

Objective: To measure the productivity of O. ficus-indica growing under three concentration os carbon dioxide.

Location: This project will be carried out in the A (- chamber ambient CO₂), L (ambient CO₂), M (+150 ppm CO₂), H (+350 ppm CO₂), and F (- chamber +350 ppm CO₂) at the University of California Riverside experimental garden.

Plant material: The 24 planted cladodes in the center m² in each chamber/area will be reserved for productivity measurements, i.e., rows 3, 4, 5, and 6 of columns D, E, F, G, H, and I.

Measurements: 1) The length of each daughter cladode growing on the 24 planted cladodes in the center m² of each chamber/area will be measured monthly (24 cladodes x 5 locations = 120).

2) After stem area index (SAI) reaches 4, cladodes will be harvested monthly or every two months as needed to maintain a SAI of 4.

3) Harvested material will be dried and weighted.

4) The center m² of planted cladodes will be harvested after two years exposure to elevated carbon dioxide.

References on elevated carbon dioxide and CAM plants

- Black, C. C., Jr. 1986. Effects of CO₂ concentrations on photosynthesis and respiration of C₄ and CAM plants. In H. Z. Enoch and B. A. Kimball (eds.). Carbon dioxide enrichment of greenhouse crops, Vol. 2: Physiology, Yield, and Economics, CRC Press, Boca Raton, FL pp. 29-40.
- Garcia de Cortazar, V. and P. S. Nobel. 1990. Worldwide environmental productivity indices and yield predictions for a CAM plant, Opuntia ficus-indica, including effects of doubled CO₂ levels. Agri. For. Meteor. 49:261-279.
- Holtum, J. A. M., M. H. O'Leary and C. B. Osmond. 1983. Effect of varying CO₂ partial pressure on photosynthesis and on carbon isotope composition of carbon-4 of malate from the Crassulacean acid metabolism plant Kalanchoe daigremontiana Hamet et Perr. Plant Physiol. 71:602-609.
- Huerta, A. J. and I. P. Ting. 1988. Effects of various levels of CO₂ on the induction of Crassulacean acid metabolism in Portulacaria afra (L.) Jacq. Plant Physiol. 71: 602-609.
- Idso, S. B., B. A. Kimball, M. G. Anderson, and S. R. Szarek. 1986. Growth response of a succulent plant, Agave vilmoriniana, to elevated CO₂. Plant Physiol. 80: 796-797.
- Nobel, P. S. 1991. Environmental productivity indices and productivity for Opuntia ficus-indica under current and elevated atmospheric CO₂ levels. Plant, Cell Envir. 14:637-646.
- Nobel, P. S. and V. Garcia de Cortazar. 1991. Growth and predicted productivity of Opuntia ficus-indica for current and elevated carbon dioxide. Agron. J. 83: 224-230.
- Nobel, P. S. and T. L. Hartsock. 1986. Short-term and long-term responses of crassulacean acid metabolism plants to elevated CO₂. Plant Physiol. 82:604-606.
- Nobel, P. S. and J. A. Palta. 1989. Soil O₂ and CO₂ effects on root respiration of cacti. Plant and Soil 120:263-271.
- Szarek, S. R., P. A. Holthe, and I. P. Ting. 1987. Minor physiological response to elevated CO₂ by the CAM plant Agave vilmoriniana. Plant Physiol. 83:938-940.

Measuremnts at UCR

Day	Measurement	Time	People
Day 0	gas exchange - 144	2 days	M/P
Day 7	R UCR - 20 excavations		P
	R UCR replant 20 cladodes/shade 50%		P
Day 12	14C labeling		Y
Day 14	Gas exchange - 144	2 days	M/P
	S UCR 14C collection 3 x 4 = 12		Y
	R UCR - 20 excavations		P
	R UCR replant 20 cladodes/shade 90%		P
Day 16	S UCR 14C collection 3 x 4 = 12		Y
Day 21	R UCR - 20 excavations		P
	R UCR - replant 20 cladodes/root cut 50%		P
Day 28	14C labeling		Y
Day 30	Gas exchange - 144	2 days	M/P
	Gas exchange on Yiqi 14C cladodes		M/Y
	R UCR - 20 excavations		P
	R UCR recut roots 50% - 20		P
	R UCR replant 20 cladodes/root cut 90%		P
	S UCR 14C collection 3 x 4 = 12		Y
Day 32	14C UCR collection 3 x 4 = 12		Y
Day 45	Gas exchange	2 days	M/P
	R UCR - 20 excavations		P
	R UCR recut roots 50% - 20		P
	R UCR recut roots 90%		P
Day 58	14C labeling		Y

Day 60	Gas exchange	2 days	M/P
	S UCR 14C collection 3 x 4 = 12		Y
	R UCR - 20 excavations		P
	R UCR recut roots 50% - 20		P
	R UCR recut roots 90% -20		P
Day 62	14C collection 3 x 4 = 12		Y
Day 67	R UCR - 20 excavations/shade		P
Day 75	R UCR - 20 excavations		P
	R UCR - 20 excavations/shade		P
	R UCR cut roots 50% - 20		P
	R UCR cut roots 90% - 20		P
Day 88	14C labeling		Y
Day 81	R UCR harvest cut root 50% - 20		P
Day 90	Gas exchange	2 days	M/P
	Gas exchange Yiqi 14C cladodes UCR		M/Y
	S UCR 14C collection 3 x 4 = 12		Y
	R UCR - 20 excavations		P
	R UCR Harvest cut roots 90% - 20		P
Day 92	14C collection 3 x 4 = 12		Y

Measurements at UCLA

Day	Measurement	Time	People
Day 0	Cladodes planted in walk in rooms		Y/M/P
Day 7	S UCLA, cladode expansion - 8		Y
	R UCLA - 8 pots		P
Day 14	S UCLA, cladode expansion - 8		Y
	R UCLA - 8 pots		P
Day 30	Gas exchange/TNC - UCLA		M/Y
	S UCLA, cladode expansion - 24		Y
	R UCLA - 8		P
	R UCLA - N, ash, cal - 72		P
Day 45	R UCLA - 8		P
Day 60	S UCLA, cladode expansion - 8		Y
	R UCLA - 8		P
Day 75	S UCLA, cladode expansion - 8		Y
	R UCLA - 8		P
Day 90	S UCLA, cladode expansion - 24		Y
	R UCLA - 8		P
	R UCLA - N, ash, cal - 72		P
	Gas exchange/TNC UCLA - 24		Y/M

N

28

W

1
2
3
4
5
6
7
8

E

A B C D E F G H I J K L

F FACE

S

H +350

M +150

L ambient + chamber

A ambient - chamber

END

DATE
FILMED

2/17/93