FINAL PROJECT REPORT

Department of Energy Grant **DE-FG05-88ER60628**, "Utilization, cycling and vertical transport of particulate organic matter in the coastal marine environment", 11/15/87 - 5/14/92, M. R. Landry (Principal Investigator).

SUMMARY OF COMPLETED PROJECT

This project was funded as part of the California Basin Study (CaBS), a DOE-funded regional program investigating production, cycling, transport, and fate of organic matter, chemical tracers, and pollutants in the Southern California Bight. The study area, adjacent to Los Angeles, was of programmatic interest due to its heavy concentration of energy-related activities, including offshore oil drilling and natural seeps, shipping, nuclear power facilities, and industrial and municipal ocean waste disposal. It was also of scientific interest because the wide continental margin in the region, pot-marked with natural sediment traps in the form of deep basins with restricted inputs and outputs, was ideal for integrating water-column and benthic studies and tracing the fates of *in situ* production and introduced pollutants.

Our role in the CaBS Program was to investigate the flux of particulate matter through the water column, emphasizing the relationship between macrozooplankton feeding and particle flux. Experimental studies of grazing-flux interactions were conducted on cruises in October 1985, May and October 1986, April and October 1987, February 1988, and January and July 1990. Experiments involved short-term, drift of moored arrays of sediment traps, ^{14}C incubations (to determine the depth distribution of particle flux and primary production rates), water column sampling (to measure the depth distributions of temperature, salinity, and particulate carbon, nitrogen, biogenic silica, and chlorophyll concentrations), and zooplankton collections (to determine population abundances, gut fluorescences, and evacuation rates of dominant taxa). The common thread in this approach was the use of phytoplankton pigments - chlorophyll *a* and phaeopigment degradation products - as tracers of phytoplankton standing stocks, macrozooplankton grazing rates, and grazing-related particle flux. Grazing rates were determined from the mean content of phytoplankton pigment in the guts, population densities in the euphotic zone, and average rates of gut clearance. Grazing-related particle flux was determined from the quantity of phaeopigments captured in sediment traps below the euphotic zone. In addition to pigments, particulate mass, carbon, nitrogen, and biogenic silica fluxes were measured from the more than 400 sediment traps deployed in the area from 1985-1990.

MASTER
DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

The main results of our investigations are summarized below. Details can be found in appended papers and manuscripts:

SEDIMENT TRAP PERFORMANCE (Landry, Peterson Andrews, in press): When comparisons were made at the same time and with the same analytical techniques, we found little significant or consistent difference in rate estimates between short-term drifting traps and long-term moored traps, between preserved and unpreserved traps in short-term experiments, between different preservatives (Formalin and mercury) in long-term experiments, between different designs of small cylindrical traps (Lorenzen closing traps vs VERTEX multitrap), and between deep-moored, cylindrical traps and large conical traps. Problems were found in comparing flux estimates from large conical traps and small cylindrical multitrap in the upper water column. Flux estimate from the two trap designs agreed within 5% for all measured parameters (pigments, carbon, nitrogen, silica, and mass) in the protected basin where currents were slow ($\sim 0.5 \text{ km d}^{-1}$). However, the large cones gave estimates 6 times less than the multitrap when compared in the upper water column where the flow environment was stronger. Historical flux estimates in the Southern California Bight were largely based on deployments of large conical traps and showed little variation in flux with depth, or frequently a significant increase in flux from 100 m to the bottom.

FLUX RATES (Landry, Peterson and Andrews, in press): Seasonal patterns in carbon and nitrogen flux were not evident in our data, but strong seasonality, with spring maxima and summer minima, were observed for export of phaeopigments and biogenic silica from the euphotic zone. In assessing mesoscale variability in the Santa Monica Basin, all of the seasonal features of particle flux were reproduced on replicate moored arrays separated by 18 km. Flux averages at comparable depths for the two stations agreed within 1%. Time-averaged rates of particulate flux for long-term deployments in 1990 were 121, 18.8, 1.5, 67 and 633 $\text{mg m}^{-2} \text{ d}^{-1}$ at 110 to 135m for carbon, nitrogen, phaeopigment, biogenic silica and mass, respectively. Flux estimates at the basin floor (835-860 m) were 50, 6.5, 0.64, 42 and 575 $\text{mg m}^{-2} \text{ d}^{-1}$, for the same parameters. The former estimates are constrained by and in good agreement with independent assessments of new production from nitrate in the euphotic zone (e.g., Jackson et al., 1989; Small et al., 1989). The latter agree with rates previously inferred from the sedimentary record using ^{210}Pb as a tracer. In addition, the difference in carbon estimates in the water column between the euphotic zone and the basin floor is consistent with carbon requirements for bacterial growth and metabolism at intermediate depths.

GUT PIGMENTS (Landry, Peterson and Fagerness, MS): Larger zooplankton taxa were more abundant during winter-spring cruises than summer-autumn cruises. Gut pigments of larger, migratory taxa (*Calanus pacificus*, *Metridia lucens*, *Pleuromamma* spp., large euphausiids and ostracods) and often some of the smaller, non-migratory forms (*Clausocalanus* spp., *Acartia* spp., and appendicularians) indicated diel periodicity in feeding intensity with highest gut pigments generally at night or in the early morning. Feeding periodicity was weakest when chlorophyll concentration and mean gut pigment content (ng chl equ. ind⁻¹) was highest in the spring. The night:day ratio of community gut pigment (CGP) varied from 1.02 to 1.95 for the winter-spring period and 1.77 to 3.39 for the autumn period. Migrating taxa explained most of the day-night difference in the winter-spring but relatively little of the difference in the autumn. Small species and developmental stages (< 1.5 mm body length) dominated daytime (95.1-99.8%) and nighttime (63.6-96.2%) CGP during the autumn and were important, though relatively less so, in the winter-spring cruises (58.2-76.9% - day; 37.7-53.7% - night). Copepods accounted for the majority of CGP for May 1986 and April 1987 cruises. Salps (*Thalia* sp.) dominated in February 1988 and were generally more important in winter-spring than autumn, as were euphausiids. Pteropods made a significant contribution to CGP in October 1985. Appendicularians (*Oikopleura* sp.) were major grazers on all cruises, but their contribution to CGP surpassed that of copepods on the three autumn cruises.

GRAZING RATES VS EXPORT FLUXES (Landry, Lorenzen and Peterson, MS): Grazing rate estimates for all mesozooplankton populations, but particularly copepods and salps, were higher during spring-winter cruises than in the autumn. Except for February 1988 when salps were abundant, copepods and appendicularians dominated, accounting for at least 80% of the estimated grazing impact on phytoplankton by the mesozooplankton community. Community grazing estimates varied seasonally from 1.7 to 7.3 mg Chl *a* m⁻² d⁻¹ and 6 to 15% Chl standing stock d⁻¹. All winter-spring estimates were substantially higher than those for the autumn cruises in terms of chlorophyll ingested, but grazing as percentage of chlorophyll standing stock cleared per day showed no consistent seasonal signal. On average, grazing accounted for a loss of 11.7% of chlorophyll standing stock per day, with a range of 6 to 18% d⁻¹.

Pigment flux from sediment traps below the euphotic zone were highest during the winter-spring cruises (1.3 to 2.1 mg Ph m⁻² d⁻¹) compared to autumn (0.4 to 1.0 mg Ph m⁻² d⁻¹). The percentage of chlorophyll standing stock lost to phaeopigment flux varied from 1.7 to 5.1% d⁻¹ with no seasonal trend. On average, only 26.5% (3.4% std. dev.) of estimated mesozooplankton grazing appeared in the traps as phaeopigment flux. Grazing estimates account

for the utilization of 29 to 55% (mean = 42%) of primary production on winter -spring cruises and 16 to 24% (mean = 19%) of primary production on summer-autumn cruises.

We concluded from these results that about 75% of the mesozooplankton grazing in the Southern California Bight was functionally indistinguishable from "microzooplankton" in that the resulting fecal debris was recycled within the euphotic zone. Although they met the formal size definition of "mesozooplankton," the important net-collected grazers in the Bight were quite small (i.e., small species and developmental stages of copepods ≤ 1 mm, mean size of appendicularians ≤ 0.5 mm, and small *Thalia* sp. ≤ 2 mm). Our results are consistent with predicted effects of body size on fecal pellet sinking rate and decomposition. Most of the material that enters sediment traps appears to do so as slowly settling, reworked detritus or marine snow aggregates.

PERSONNEL

The following individuals received partial salary support from DOE Grant DE-FG05-88ER60628:

M. R. Landry, Professor, Department of Oceanography, University of Hawaii
 C. C. Andrews, Research Associate, Department of Oceanography, University of Hawaii
 K. E. Selph, Graduate Student, Department of Oceanography, University of Hawaii
 W. K. Peterson, Oceanographer III, School of Oceanography, University of Washington
 C. Park, Visiting Scientist, Chungnam University, South Korea

PUBLISHED PAPERS AND SUBMITTED MANUSCRIPTS:

The following papers and manuscripts acknowledge full or partial support from DOE Grant DE-FG05-88ER60628:

Jackson, G. A., F. Azam, A. F. Carlucci, R. W. Eppley, B. Finney, D. S. Gorsline, B. Hickey, C.-A. Huh, R. A. Jahnke, I. R. Kalpan, M. R. Landry, L. F. Small, M. I. Venkatesan, P. M. Williams, and K. M. Wong. 1989. Elemental cycling and fluxes off Southern California. *EOS* 70: 146-149.

Small, L. F., M. R. Landry, R. W. Eppley, F. Azam, and A. F. Carlucci. 1989. Role of plankton in the carbon and nitrogen budgets of Santa Monica Basin, California. *Mar. Ecol. Prog. Ser.* 56:57-74.

- Hassett, R. P. and M. R. Landry. 1990. Seasonal changes in feeding rate, digestive enzyme activity, and assimilation efficiency of the copepod *Calanus pacificus* in Puget Sound, Washington. *Mar. Ecol. Prog. Ser.* 62: 203-210.
- Hassett, R. P. and M. R. Landry. 1990. Effects of diet and starvation on digestive enzyme activity and feeding behavior of *Calanus pacificus*. *J. Plank. Res.* 12: 991-1010.
- Landry, M.R., V.L. Fagerness and W.K. Peterson. 1991. Ontogenetic patterns in the distribution of *Pseudocalanus* spp. during upwelling off the coast of Washington, USA. Proc. 4th Intern. Conf. Copepoda *J. Plank. Soc. Japan*, Spec. Vol. 1991: 451-466.
- Landry, M.R., W.K. Peterson and C.C. Andrews. In press. Particulate flux in the water column overlying Santa Monica Basin. *Prog. Oceanogr.*
- Park, C. and M.R. Landry. Subm. Egg production by the subtropical copepod *Undinula vulgaris*. *Mar. Biol.*
- Huh, C.-A., T.-L. Ku, S. Luo, M.R. Landry, and P.M. Williams. Subm. Fluxes of thorium isotopes in the Santa Monica Basin. *Earth Planet. Sci. Lett.*
- Landry, M.R., W.K. Peterson and V.L. Fagerness. MS. Zooplankton grazing in the Southern California Bight. I. Population abundances and gut pigment contents of mesozooplankton. To be submitted to *Mar. Ecol. Prog. Ser.*
- Landry, M.R., C.J. Lorenzen and W.K. Peterson. MS. Zooplankton grazing in the Southern California Bight. II. Mesozooplankton grazing impact and pigment flux. To be submitted to *Mar. Ecol. Prog. Ser.*

The following are publications supported by previous DOE research grants but published during the award period for Grant DE-FG05-88ER60628:

- Hassett, R. P. and M. R. Landry. 1988. Short-term changes in feeding and digestion by the copepod *Calanus pacificus*. *Mar. Biol.* 99: 63-78.
- Landry, M. R. and V. L. Fagerness. 1988. Behavioral and morphological influences on predatory interactions among marine copepods. *Bull. Mar. Sci.* 43: 509-529.
- Herman, A. J., B. M. Hickey, M. R. Landry, and D.F. Winter. 1989. Coastal upwelling dynamics. Pages 211-254 in M.R. Landry and B.M. Hickey (eds.), *Coastal Oceanography of Washington and Oregon*. Elsevier.
- Hickey, B. M. and M. R. Landry. 1989. Process interactions and prospectus for future research on the Washington/Oregon shelf. Pages 595-600 in M.R. Landry and B.M. Hickey (eds.), *Coastal Oceanography of Washington and Oregon*. Elsevier.
- Landry, M.R. and B.M. Hickey (eds.). 1989. *Coastal Oceanography of Washington and Oregon*. Elsevier.
- Landry, M. R. and C. J. Lorenzen. 1989. Abundance, distribution, and grazing impact of zooplankton on the Washington shelf. Pages 175-210 in M.R. Landry and B.M. Hickey (eds.), *Coastal Oceanography of Washington and Oregon*. Elsevier.
- Landry, M. R., J. Postel, J. Newman, and W. K. Peterson. 1989. Broad-scale distributional patterns of hydrographic variables on the Washington/Oregon shelf. Pages 1-40 in M.R. Landry and B.M. Hickey (eds.), *Coastal Oceanography of Washington and Oregon*.

APPENDICES

- Jackson, G. A., F. Azam, A. F. Carlucci, R. W. Eppley, B. Finney, D. S. Gorsline, B. Hickey, C.-A. Huh, R. A. Jahnke, I. R. Kalpan, M. R. Landry, L. F. Small, M. I. Venkatesan, P. M. Williams, and K. M. Wong. 1989. Elemental cycling and fluxes off Southern California. *EOS* 70: 146-149.
- Small, L. F., M. R. Landry, R. W. Eppley, F. Azam, and A. F. Carlucci. 1989. Role of plankton in the carbon and nitrogen budgets of Santa Monica Basin, California. *Mar. Ecol. Prog. Ser.* 56:57-74.
- Hassett, R. P. and M. R. Landry. 1990. Seasonal changes in feeding rate, digestive enzyme activity, and assimilation efficiency of the copepod *Calanus pacificus* in Puget Sound, Washington. *Mar. Ecol. Prog. Ser.* 62: 203-210.
- Hassett, R. P. and M. R. Landry. 1990. Effects of diet and starvation on digestive enzyme activity and feeding behavior of *Calanus pacificus*. *J. Plank. Res.* 12: 991-1010.
- Landry, M.R., V.L. Fagerness and W.K. Peterson. 1991. Ontogenetic patterns in the distribution of *Pseudocalanus* spp. during upwelling off the coast of Washington, USA. Proc. 4th Intern. Conf. Copepoda *J. Plank. Soc. Japan*, Spec. Vol. 1991: 451-466.
- Landry, M.R., W.K. Peterson and C.C. Andrews. In press. Particulate flux in the water column overlying Santa Monica Basin. *Prog. Oceanogr.*
- Park, C. and M.R. Landry. Subm. Egg production by the subtropical copepod *Undinula vulgaris*. *Mar. Biol.*
- Huh, C.-A., T.-L. Ku, S. Luo, M.R. Landry, and P.M. Williams. Subm. Fluxes of thorium isotopes in the Santa Monica Basin. *Earth Planet. Sci. Lett.*

Copies of the following two manuscripts will be sent when they are submitted for publication:

- Landry, M.R., W.K. Peterson and V.L. Fagerness. MS. Zooplankton grazing in the Southern California Bight. I. Population abundances and gut pigment contents of mesozooplankton. To be submitted to *Mar. Ecol. Prog. Ser.*
- Landry, M.R., C.J. Lorenzen and W.K. Peterson. MS. Zooplankton grazing in the Southern California Bight. II. Mesozooplankton grazing impact and pigment flux. To be submitted to *Mar. Ecol. Prog. Ser.*

*reprints + preprints
removed. obs*

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

END

**DATE
FILMED**

12 / 8 / 92