DOE/ER-53266-42

Technical Progress Report**PLASMA CONFINEMENT THEORY AND TRANSPORT SIMULATION***David W. Ross*

Fusion Research Center, The University of Texas at Austin, Austin, Texas 78712

May 1, 1992 - April 30, 1993

U. S. Department of Energy, Office of Energy Research, Grant DE-FG05-88ER53266

SUMMARY OF PROGRESS

The objectives continue to be: (1) to advance the transport studies of tokamaks, including development and maintenance of the Magnetic Fusion Energy Database, and (2) to provide theoretical interpretation, modeling and equilibrium and stability studies for TEXT-Upgrade. Recent publications and reports [1-12], and conference presentations [13-25] of the Fusion Research Center theory group are listed on attached pages.

A. Magnetic Fusion Energy Database

The Magnetic Fusion Energy Database (MFEDB) continues to expand and evolve in order to serve the needs of the fusion community [13-15]. Our principal effort this year has been to strengthen connections with major laboratories to develop collaborations and greater use of the database. We received positive responses from several of these, notably ASDEX, JT-60, and T-10. To demonstrate the feasibility of this collaborative work, we have 1) initiated a pilot-project transport scaling study, 2) prepared to participate in the effort of the H-mode working group (J. DeBoo, chair) to develop a profile database of well-diagnosed shots, and 3) proposed to provide data-handling for the various edge physics groups. We organized a discussion on these and other collaborations during the APS meeting in November, 1992. Dr. Miner will attend the next meeting of the H-mode group at the JET laboratory in April, 1993. We are also intensifying our contact with the ITER central and U. S. home teams.

Preliminary work for these projects is nearly complete and includes defining the additional tables and generic plasma parameters necessary for cross-machine comparisons. Needed statistical functions are being defined within the *Mathematica*® interface [12], described below and in the 1991-92 report. We are working closely with Dr. Tsui and Dr. Wootton of to establish an edge physics section of MFEDB, beginning with TEXT data. Dr. Yagi, from JAERI, visited in January, 1993, and succeeded in transferring a large number of JT-60 shots to our repository. He and Dr. Shirai also became familiarized with using the database.

Two new network interfaces have been developed. The first is an RPC service which allows programs that have Internet access to retrieve data directly from the MFEDB host computer. A simple RPC client is available as a template for those who would like to incorporate it in their code. A *Mathematica*® package [12] has been developed using the RPC service to allow direct access to MFEDB from within a *Mathematica* session. The package includes a custom set of functions and also accepts standard SQL queries. The data is returned as *Mathematica* lists that can be manipulated by any *Mathematica* function. Users can incorpo-

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

rate MFEDB data directly into their *Mathematica* calculations.

In order to enter data received in arbitrary formats, a new code has been written which automatically parses the input file and stores the data in the database. In direct response from participants in the Numerical Tokamak Project, new data types, including density and temperature fluctuation data, will be added to the database as they become available. The fusion community is currently investigating standard data formats and data access software to provide a mechanism for comparing codes and allowing codes to communicate. The MFEDB will comply with these standards allowing users to retrieve data from the database in the "standard" file format, as well as through the *Mathematica* interface.

B. Support for TEXT-Upgrade and Diagnostics

1. A set of magnetic probes will be used for plasma shape reconstruction in TEXT-Upgrade. We have done a study to determine the minimum number of coils necessary to achieve acceptable magnetic reconstruction [16], using the equilibrium and fitting code EFIT.¹ The optimum placement of the probes was investigated, taking into account the variety of plasma shapes to be produced and the effect of measurement errors. Iron core effects, which are accounted for here,² have also been studied in a collaboration with JET [7].

¹L. L. Lao, H. St. John, R. D. Stambaugh, et al., Nucl. Fusion **25**, 1611 (1985).

²E. R. Solano, G. H. Nielson, and L. L. Lao, Nucl. Fusion **30**, 1107 (1990).

2. The apparatus that we have designed and assembled for the measurement of the magnetic-field profile on TEXT-U,³ utilizing successive charge-exchange collisions, is awaiting operation of the tokamak.

³P. M. Valanju, R. D. Bengtson, and A. Ouroua, Rev. Sci. Instrum. **61**, 2935 (1990).

3. Pending the operation of TEXT-U, our concluding work on the use of the Heavy Ion Beam Probe (HIBP) to measure local density fluctuations has been accepted for publication [9]. The HIBP signal is sensitive to modulation due to density fluctuations along the entire beam trajectory. We present a model of the HIBP experiment on the TEXT tokamak that includes profile information for equilibrium and fluctuating parameters, allows for differences in the radial and poloidal characteristics of the fluctuations, and uses realistic beam trajectories. We show how experimental measurements can be analyzed for modulation without extensive model calculations. Calculations show that for low- to medium-density plasmas path effects are likely to be present, but do not dominate the measured signals.

4. Time and space correlation properties of the thermal Electron Cyclotron radiation have been analyzed [17]. A comparison with the analogous properties of the ECE fluctuations induced by various non-equilibrium plasma phenomena was performed and the plasma-diagnostic potential of the EC radiation fluctuations was investigated.

5. Extensive measurements of impurity rotation in TEXT were reported for Ohmic plasmas [18]. Comparison with simple neoclassical theory suggests that rotation can be predicted.⁴ We have extended the comparisons to a more complete neoclassical theory which predicts impurity rotation explicitly.⁵ Comparisons of rotation velocities of different impurities (C⁺⁴, O⁺⁴, He⁺¹) were made. With this work as background we have attempted to understand perturbations of rotation by EML, ECH and biased electrode. We have also collaborated in a

general review of rotation in Ohmic plasmas [10].

⁴R. D. Hazeltine, Phys. Fluids 17 (5), 1974.

⁵Y. B. Kim, P. H. Diamond, and R. J. Groebner, Phys. Fluids B 3 (8), 1991.

C. Transport Studies

1. Edge physics and L-H transition

In collaboration with TEXT and the IFS, we are pursuing a broad range of edge physics and H-mode studies, many of which are related to the experiments described above.

a. We have developed a kinetic theory of the neutral distribution function, including charge-exchange (CX) and ionization, in the long CX mean-free-path (mfp) limit. This theory is used to calculate the contribution of CX to the plasma viscosity and toroidal acceleration [19]. We analytically calculate the exact toroidal acceleration in a tokamak plasma which has been biased by a radial probe, as in TEXT experiments. The toroidal acceleration is found to be affected by: (i) viscosity, (ii) space-charge, (iii) current in metallic conductors, and (iv) changes in total enclosed charge. The contribution of CX viscosity in the long mfp limit is found to be the dominant feature in determining toroidal acceleration.

b. Traditionally, neoclassical transport calculations ignore poloidal variation of the poloidal magnetic field. Near an X-point of a diverted plasma, the poloidal field is small, causing guiding centers to linger at that poloidal position. Indeed, the poloidal transit frequency approaches zero near the separatrix, so that v^* becomes locally large for any plasma temperature. We have presented a model flux function with an X-point [20], and with it solved the drift kinetic equation to calculate the first order perturbation to the distribution function. X-point effects in the plateau regime were then calculated inside the separatrix.

c. Recent DIII-D and TEXT experiments show a sudden increase in plasma poloidal rotation as a signature of L to H transition. We have emphasized that shear in the radial electric field near the plasma edge indicates a non-neutral space charge distribution [21]. For such space charge a natural source is the "halo" effect: the relatively large ion (poloidal) gyroradius precludes perfect tracking between n_i and n_e . We build a simple model, including the halo, to find the relation between plasma poloidal flow and radial electric field. The resulting integro-differential equation shows that a Maxwell-Boltzmann electron distribution (no flow) does not allow space charge. Only in the presence of a sheared velocity can space charge develop.

2. Oscillating gas puff

Transport simulations of modulated gas-feed experiments in TEXT [4,6] have been continued [22], using scalings corresponding to trapped-particle mode theory, but with diagonal and off-diagonal transport coefficients. Because of nonlinear functional dependences on plasma parameters, the amplitude and phase of the perturbed density are quite different from those of a simple diffusion-convection (D-V) model and tend to be insensitive to variations in numerical factors in the transport coefficients. The amplitudes and phases in the simulation results seem to disagree with the data in contradictory ways, particularly at higher density. Work continues with a variety of models to attempt to resolve the discrepancies and fit the data.

3. Other transport studies

a. Transport phenomena were investigated arising due to an up-down asymmetry of ripple and helical perturbations of the magnetic field in tokamaks and stellarators [23]. Taking into account this asymmetry is important for devices with vertically displaced magnetic axis or with single-null divertor and, in particular, for one version of the ITER project. The asymmetry manifests itself in the different magnitude of the ripple or helical perturbation in the up and down parts of the magnetic surface and leads to certain peculiarities of the toroidally trapped, locally trapped, and transitional orbits. The most pronounced effect is the net radial drift of the banana particles which is absent in conventional neoclassical theory. Transport associated with the up-down asymmetry is estimated.

b. We developed an approach to turbulent diffusion in tokamaks where the toroidal drift is considered as a perturbation to the particle motion in the electromagnetic fluctuations [24]. The analysis of the simplest single scale case,⁶ when the width of the neoclassical trajectory, D , was equal to the characteristic perpendicular scale of electromagnetic oscillations, L , is extended to an arbitrary ratio of D/L . The diffusion mechanism can substantially differ from that of the single scale case. In particular, small neoclassical displacement of the electrons, $D \ll L$, gives rise to the diffusion exceeding the Hamiltonian diffusion. Due to the dependence of the trajectory on the particle energy this process finally results in spatial diffusion. On the other hand, for $D \gg L$, two types of trajectories are formed: open, along which particles can move in the direction of the toroidal drift, and closed, with particles trapped near extrema of the fluctuating potential. The transport process in this case is defined by the fine structure of the transition layer between these two types of orbit. It was shown that the neoclassical diffusion coefficient in a turbulent plasma is weakly dependent on the value of D/L and generally determined by the scale length of the fluctuations, L .

⁶P. N. Yushmanov, Comments Plasma Phys. and Contr. Fusion 14 (1992) 313.

c. We assisted in the investigation of the electrostatic wake of a superthermal test electron in a magnetized plasma [12]. The potential is determined for a test electron with $v_{\parallel} \gg v_{Te}$ in a uniform magnetized plasma ($\omega_{ce} \gg \omega_{pe}$). In the frame of the test electron, part of the spatially oscillatory potential has spherical symmetry to the rear of the electron and is zero ahead of the electron. A second part, which makes the potential continuous at the plane containing the electron, is oscillatory radially but decreases almost monotonically in the axial direction.

D. Alfvén Waves

The code presented previously to study kinetic effects in antenna-driven toroidal Alfvén waves⁷ has been rewritten to make use of improved numerical methods and to include coupling to more than three poloidal harmonics [8,24]. Toroidal Alfvén Eigenmodes and Global Alfvén Eigenmodes are identified as resonances in the antenna impedance. Mode structure, energy deposition profiles, and damping rates are investigated as functions of the kinetic (effective-resistivity) model. Results have been compared with analytical calculations,⁸ and, in collaboration with General Atomics [8], will be compared with non-ideal MHD calculations.

⁷W. Q. Li, S. M. Mahajan, D. W. Ross, and M. Chu, Bull. Am. Phys. Soc. 36, (1991).

⁸R. R. Mett and S. M. Mahajan, Phys. Fluids B 4, 2885 (1992).

REPORTS AND PUBLICATIONS 1992-93

Names underlined are those supported by the contract.

publication status update:

1. R. V. Bravenec, K. W. Gentle, B. Richards, D. W. Ross, D. C. Sing, A. J. Wootton, D. L. Brower, N. C. Luhmann Jr., W. A. Peebles, C. X. Yu, T. P. Crowley, J. W. Heard, R. L. Hickok, P. M. Schoch, and X. Z. Yang, *Core Turbulence and Transport Studies on the TEXT Tokamak*, Phys. Fluids B 4, 2127 (1992).
2. D. W. Ross, M. L. Sloan, A. J. Wootton, P. M. Schoch, J. W. Heard, T. P. Crowley, R. L. Hickok, and V. Simcic, *Effect of Beam Attenuation Modulation on Fluctuation Measurements by Heavy Ion Beam Probe*, Rev. Sci. Instruments 63, 2232 (1992).
3. S. M. Mahajan, P. M. Valanju, and W. L. Rowan, *Closed, Form Invariant Solutions of Convective-Diffusive Systems with Applications to Impurity Transport*, Phys. Fluids B 4, 2495 (1992).
4. B. Richards, M. E. Austin, R. V. Bravenec, D. L. Brower, K. W. Gentle, R. L. Hickok, N. C. Luhmann, A. Ouroua, W. A. Peebles, P. E. Phillips, D. W. Ross, W. L. Rowan, P. M. Schoch, P. M. Valanju, J. C. Wiley, A. J. Wootton, X. Z. Yang, and C. X. Yu, *Tests of Plasma Transport Models via Perturbation Techniques*, Nucl. Fusion 32, 567 (1992).
5. P. M. Valanju, M. D. Calvin, R. D. Hazeltine, and E. R. Solano, *The Effect of Charge Exchange on Plasma Flows*, Phys. Fluids B 4, 2675 (1992).

additional reports and publications

6. K. W. Gentle, B. Richards, M. E. Austin, R. V. Bravenec, D. L. Brower, R. F. Gandy, W. L. Li, P. E. Phillips, D. W. Ross, W. L. Rowan, P. M. Schoch, P. M. Valanju, and A. J. Wootton, *Equilibrium and Perturbed Fluxes and Turbulence Levels in a Tokamak: Implications for Models*, Phys. Rev. Lett. 68, 2444 (1992).
7. D. P. O'Brien, L. L. Lao, E. R. Solano, M. Garribba, T. S. Taylor, J. G. Cordey, and J. J. Ellis, *Equilibrium Analysis of Iron Core Tokamaks using a Full Domain Method*, Nucl. Fusion 32, 1351 (1992).
8. M. S. Chu, A. D. Turnbull, J. M. Greene, L. L. Lao, M. S. Chance, H. L. Berk, B. N. Breizman, W. Q. Li, D. M. Lindberg, S. M. Mahajan, R. R. Mett, D. W. Ross, J. W. Van Dam, J. C. Wiley, H. Ye, J. Candy, and M. N. Rosenbluth, *Theory of Toroidicity-induced Alfvén Eigenmodes in a Finite-beta Arbitrary Cross-section Tokamak and Analysis of Nonideal Effects, Stability Thresholds, and Nonlinear Behavior*, in *Plasma Physics and Controlled Nuclear Fusion Research 1992* (Proc. 14th Int. Conf. Wuerzburg, IAEA, Vienna, 1992), paper IAEA-CN-56/D-2-2, to be published.
9. J. W. Heard, T. P. Crowley, D. W. Ross, P. M. Schoch, R. L. Hickok Jr., and B. Z. Zhang, *Path Integral Effects in Heavy Ion Beam Probe Density Measurements: A comparison of Simulation Results and Experimental Data*, to be published in Rev. Sci. Instrum. (1992).

10. W. L. Rowan, A. G. Meigs, E. R. Solano, P. M. Valanju, M. D. Calvin, and R. D. Hazeltine, *Rotation in Ohmically Heated Tokamaks: Experiment and Theory*, to be published in Phys. Fluids (1993).
11. A. A. Ware and J. C. Wiley, *The Electrostatic Wake of a Superthermal Test Electron in a Magnetized Plasma*, The University of Texas Report IFSR #561, 1992, revision in progress.
12. J. C. Wiley, W. H. Miner Jr., and D. W. Ross, *MDI: Mathematica Database Interface for the MFEDB*, The University of Texas, Report FRCR #411 1992.

American Physical Society, Division of Plasma Physics, November, 1992:

13. W. H. Miner Jr., R. T. McCann, and J. C. Wiley, *New Data, New Data Types and Future Directions of the MFE Database*, Bull. Am. Phys. Soc. **37**, 1503 (1992).
14. J. C. Wiley, W. H. Miner Jr., and D. W. Ross, *RPC and Mathematica Interfaces for the MFE Database*, Bull. Am. Phys. Soc. **37**, 1543 (1992).
15. R. T. McCann, W. H. Miner Jr., and J. C. Wiley, *Using LEX and YACC to Build a Parser for the MFEDB*, Bull. Am. Phys. Soc. **37**, 1393 (1992).
16. W. B. Xu, E. R. Solano, and A. J. Wootton, *Optimal Magnetic Probe Placement in TEXT-Upgrade*, Bull. Am. Phys. Soc. **37**, 1385 (1992).
17. G. Cima and P. M. Valanju, *Correlation Properties of ECE Fluctuations*, Bull. Am. Phys. Soc. **37**, 1389 (1992).
18. A. G. Meigs, W. L. Rowan, D. Crockett, K. W. Gentle, E. Solano, R. L. Hickok, P. M. Schoch, X. Z. Yang, and B. Z. Zhang, *Rotation Experiments in TEXT*, Bull. Am. Phys. Soc. **37**, 1385 (1992).
19. M. D. Calvin, P. M. Valanju, R. D. Hazeltine, and E. R. Solano, *Charge-Exchange Viscosity and Toroidal Acceleration*, Bull. Am. Phys. Soc. **37**, 1432 (1992).
20. E. R. Solano and R. D. Hazeltine, *X-Point Effect on Neoclassical Kinetic Theory*, Bull. Am. Phys. Soc. **37**, 1601 (1992).
21. H. Xiao, P. M. Valanju, and R. D. Hazeltine, *Gyrosheath Near the Tokamak Edge*, Bull. Am. Phys. Soc. **37**, 1433 (1992).
22. D. W. Ross, B. Richards, and J. C. Wiley, *Comparison of Transport Models in Simulation of Modulated Gas Feed Experiments*, Bull. Am. Phys. Soc. **37**, 1389 (1992).
23. P. N. Yushmanov, J. Hernandez, J. R. Cary, and S. G. Shasharina, *Transport in Tokamak with Up-down Asymmetry*, Bull. Am. Phys. Soc. **37**, 1366 (1992).
24. P. N. Yushmanov and A. I. Smolyakov, *Neoclassical Transport in a Turbulent Plasma for Arbitrary Ratio of Regular and Fluctuation Orbit Sizes*, Bull. Am. Phys. Soc. **37**, 1475 (1992).
25. W. O. Li, D. W. Ross, J. C. Wiley, S. M. Mahajan, and R. R. Mett, *Numerical Studies of Toroidal Alfvén Waves*, Bull. Am. Phys. Soc. **37**, 1605 (1992).

END

**DATE
FILMED**

4 / 19 / 93