DOE/ER/60503--352

DE92 019916

ANNUAL REPORT

1991-1992

UNITED STATES DEPARTMENT OF ENERGY

DE-FG05-87ER60503

"A PHYSICO-CHEMICAL STUDY OF SOME
AREAS OF FUNDAMENTAL SIGNIFICANCE TO BIOPHYSICS"

S. P. McGlynn
Principal Investigator
Boyd Professor

Devendra Kumar
Co-Principal Investigator
Associate Professor-Research

April 30, 1992

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

TABLE OF CONTENTS

PAGE

INTRODUCTION	4
--------------------	---

ITEM 1. PUBLICATION ACTIVITY (1991-1992),
BIBLIOGRAPHY OF TITLES PUBLISHED,
SYMPOSIA/SEMINARS/REVIEWS, OTHER ACTIVITIES
AND PRESENT SCIENTIFIC PERSONNEL

A. Publication Activity 1991-1992	6
B. Bibliography of Titles	8
C. Symposia Attended	12
D. Other Activities	
(i) Dedicated Synchrotron Beam Lines for Radiation Signature	15
(ii) Laboratory Guests	20
(iii) Recognition and Awards	20
E. Present Scientific Personnel	21

ITEM 2. SCIENTIFIC REPORT

A. Radiation Signatures	23
B. Photoelectron Spectroscopy of Biologically Active Molecules	38
C. Laser Optogalvanic Effect	47
D. Magnetic Circular Dichroism	65
E. Photochemistry of Halogenated Molecules	77
F. Density Effects on High-n Rydbergs	81

INTRODUCTION

A comprehensive list of publications for the year 1991-92, which includes manuscripts published and/or in process of publication and/or in preparation for submission, is given in Section 1.A, Publication Activity 1991-1992.

The seventeen (17) titles (Nos. 1-17) of Item 1.A for the 1991-1992 year constitute the gist of this annual report. These titles represent original work completed and published, in press or submitted during the current reporting period.

The Scientific Report of Item 2 provides a summary of the research in progress in the Molecular Spectroscopy Group.

Since many items listed in previous years were published some years later and since we like to be orderly in our approach to Annual Publication Lists, we also provide in Section 1.B, Bibliography of Titles, information on the fate of all unpublished items from previous annual lists. This section also provides new titles pertinent to the present reporting period (1991-1992).

ITEM 1. PUBLICATION ACTIVITY 1991-1992,
BIBLIOGRAPHY OF TITLES PUBLISHED,
SYMPOSIA/SEMINARS/REVIEWS,
OTHER ACTIVITIES
AND PRESENT SCIENTIFIC PERSONNEL

A. PUBLICATION ACTIVITY 1991-1992

DOE Grant No. DE-FG05-87ER60503

1. L. Klasinc, Lj. Pasa-Tolic, H. Spiegl, J. V. Knop, and McGlynn, "Positional and Conformational Dependence of Long-Range Interaction in 3,16- and 3,17-Androstanediones," J. Math. Chem. 8, 179 (1991).
-- -Paper, published
2. L. Nyulaszi, T. Vezpremi, J. Reffy, G. Zsombok, T. Cvitas, B. Kovac, L. Klasinc, and S. P. McGlynn, "Photoelectron Spectra of Halogenofurans," Int. J. Quant. Chem: Chem. Symp. 25, 479 (1992).
-- -Paper, published
3. Lj. Pasa-Tolic, L. Klasinc, H. Spiegl, J. V. Knop and S. P. McGlynn, "Ab Initio Calculations on 5 α -Androstane," Int. J. Quant. Chem. 41, 815 (1992)
-- -Paper, published
4. S. P. McGlynn, K. Wittel and L. Klasinc, "The Orbital Concept as a Foundation for Photoelectron Spectroscopy," p. 63-98 of Theoretical Models of Chemical Bonding, Part 3 of Molecular Spectroscopy, Electronic Structure and Intramolecular Interactions, ed. Z. B. Maksic, Springer Verlag, Berlin, 1991.
-- -Paper, published
5. R. Armstrong, "A Study of the Optogalvanic Effect in Water and Argon," a report presented to LaSER (Louisiana Stimulus for Excellence in Research), January, 1992.
-- -Report, submitted
6. D. Kumar and S. P. McGlynn, "Analysis of Radiofrequency Discharges in Plasma," patent accepted, 1992.
-- -Patent, granted
7. S. P. McGlynn and M. N. Varma, "Radiation Signatures", Proceedings of the Council of the European Community Workshop on Biophysical Modelling of Radiation Effects, Padova, Italy, September 1991, Adam Hilger (IOP), Bristol, England (in press).
-- -Paper, in press
8. S. P. McGlynn, K. Rupnik, M. N. Varma and L. Klasinc, "Radiation Signatures for Biology," Int. J. Quant. Chem., Quant. Biol. Symp.
-- -Paper, in preparation

9. M. Eckert-Maksic, Z. B. Maksic, M. Hodoscek, and K. Rupnik, "Intra- and Extramolecular Electrostatic Potentials in Vitamin C", J. Mol. Struct. (THEOCHEM).
-- -Paper, in press
10. S. P. McGlynn and D. Kumar, "The Laser Optogalvanic (LOG) Effect in Radiofrequency Discharges," Proceedings of the International Society for Optical Engineering Meeting on *Lasers, Sensors and Spectroscopy*, Los Angeles, California, January 1992.
-- -Paper, in press
11. A. Alebic-Juretic, T. Cvitas and L. Klasinc, "Ozone Destruction on Powders," Ber. Bunsenges. Phys. Chem.
-- -Paper, in press
12. T. Pasinszki, T. Veszpremi, M. Feher, B. Kovac, L. Klasinc and S. P. McGlynn, "The Photoelectron Spectra of Methyl Pseudohalides," Int. J. Quantum Chem. Quantum Chem. Symp.
-- -Paper, in press
13. K. Rupnik, and S. P. McGlynn, "Molecular Lesion Spectra as Radiation Signatures", Radiation Research.
-- -Paper, submitted
14. U. Asaf, R. Reininger, K. Rupnik and S. P. McGlynn, "Pressure Shift and Electron Scattering Length in Atomic and Molecular Gases," J. Phys. Chem.
-- -Paper, submitted
15. U. Asaf, S. Felps, K. Rupnik, and S. P. McGlynn, "Electron Scattering in Dense Hydrogen-Argon Gas Mixtures: A Pressure Shift Study," Phys. Rev. A.
-- -Paper, submitted
16. M. Surgi, D. Giamalva and L. Klasinc, "Correlation Between Ozonolysis Rate and π -Ionization Potential of 10,10'-Dimethyl-9,9'-Biacridylidene." Bull. Slov. Chem. Soc.
-- -Paper, submitted
17. S. P. McGlynn, K. Rupnik and M. N. Varma, "Molecular Lesion Spectra are Radiation Signatures," 1991.
-- -Patent, disclosed

B. BIBLIOGRAPHY OF TITLES

The bibliography of nineteen (19) titles constitutes the bulk of this Report. These are separated into three (3) categories. The first of these consists of one (1) title and provides information pertaining to the 1989-1990 Annual Report; the second of these consists of four (4) titles and provides information pertaining to the 1990-1991 Annual Report; the third of these consists of fourteen (14) new titles and provides reprints or preprints pertinent to the present reporting period (1991-1992).

The serial numbers assigned to the titles are in numerical order and represent the number of original manuscripts/reports/dissertations published since the institution of the original DOE grant several years ago.

1. **Completion of the 1989-1990 Annual Report**

323. S. P. McGlynn, K. Wittel and L. Klasinc, "The Orbital Concept as a Foundation for Photoelectron Spectroscopy," in Theoretical Models of Chemical Bonding, Part 3 of Molecular Spectroscopy, Electronic Structure and Intramolecular Interactions, ed. Z. B. Maksic, Springer Verlag, Berlin, 1991, p. p. 63-98.

(7 Xerox copies of the Chapter appended to this report).
The 1989-1990 Annual Report is now complete.

2. **Completion of the 1990-1991 Annual Report**

335. L.Nyulaszi, T. Vezpremi, J. Reffy, G. Zsombok, T. Cvitas, B. Kovac, L. Klasinc, and S. P. McGlynn, "Photoelectron Spectra of Halogenofurans," Int. J. Quant. Chem: Quant. Chem. Symp. 25, 479 (1991)
(7 reprints appended to this report).

336. L. Klasinc, Lj. Pasa-Tolic, H. Spiegl, J. V. Knop, and S. P. McGlynn, "Positional and Conformational Dependence of Long-Range Interaction in 3,16 - and 3,17 - Androstanediones," J. Math. Chem. 8, 179 (1991).
(7 reprints appended to this report).

337. Lj. Pasa-Tolic, L. Klasinc, H. Spiegl, J.V. Knop, and S. P. McGlynn, "Ab initio Calculations on 5 α - Androstanes," Int. J. Quant. Chem.: Quant. Biol. Symp. 41, 815 (1992)
(7 reprints appended to this report).

338. D. Kumar and S. P. McGlynn, "Analysis of Radiofrequency Discharges in Plasma," Patent Application (accepted), Patent number to be issued.
(7 copies appended to the 1990-1991 Annual Report).

Apart from the issuance of Patent Number for item #338, the 1990-1991 Report is complete.

3. **Bibliography for the 1991-1992 Annual Report**

340. S. P. McGlynn and M. N. Varma, "Radiation Signatures", Proceedings of the Council of the European Community Workshop on Biophysical Modelling of Radiation Effects, Padova, Italy, September 1991, Adam Hilger (IOP), Bristol, England (in press).
(7 preprints appended to this report).

341. S. P. McGlynn, K. Rupnik, M. N. Varma and L. Klasinc, "Radiation Signatures for Biology," *Int. J. Quant. Chem., Quant. Biol. Symp.*, in preparation for submission.
(7 preprints appended to this report).
342. M. Eckert-Maksic, Z. B. Maksic, M. Hodoscek, and K. Rupnik, "Intra- and Extramolecular Electrostatic Potentials in Vitamin C" *J. Mol. Struct. THEOCHEM*, (in press).
(7 preprints appended to this report).
343. S. P. McGlynn and D. Kumar, "The Laser Optogalvanic (LOG) Effect in Radiofrequency Discharges," *Proceedings of the International Society for Optical Engineering Meeting on Lasers, Sensors and Spectroscopy*, Los Angeles, California, January 1992 (in press).
(7 preprints appended to this report).
344. A. Alebic-Juretic, T. Cvitas and L. Klasinc, "Ozone Destruction on Powders," *Ber. Bunsenges. Phys. Chem.* (in press).
(7 preprints appended to this report).
345. T. Pasinszki, T. Veszpremi, M. Feher, B. Kovac, L. Klasinc and S. P. McGlynn, "The Photoelectron Spectra of Methyl Pseudohalides," *Int. J. Quantum Chem. Quantum Chem. Symp.* (in press).
(7 preprints appended to this report).
346. K. Rupnik, and S. P. McGlynn, "Molecular Lesion Spectra as Radiation Signatures", *Radiation Research* (submitted).
(7 preprints appended to this report).
347. U. Asaf, R. Reininger, K. Rupnik and S. P. McGlynn, "Pressure Shift and Electron Scattering Length in Atomic and Molecular Gases," *J. Phys. Chem.* (submitted).
(7 preprints appended to this report).
348. U. Asaf, S. Felps, K. Rupnik, and S. P. McGlynn, "Electron Scattering in Dense Hydrogen-Argon Gas Mixtures: A Pressure Shift Study," *Phys. Rev. A*. (submitted).
(7 preprints appended to this report).
349. M. Surgi, D. Giamalva and L. Klasinc, "Correlation Between Ozonolysis Rate and π -Ionization Potential of 10,10'-Dimethyl,-9,9'-Biacridylidene." *Bull. Slov. Chem. Soc.* (submitted).
(7 preprints appended to this report).
350. S. P. McGlynn, K. Rupnik and M. N. Varma, "Molecular Lesion Spectra are Radiation Signatures," *Patent Application* (pending).
(7 copies of patent disclosure form appended to this report).

351. R. Armstrong, "A Study of the Optogalvanic Effect in Water and Argon," a report presented to LaSER (Louisiana Stimulus for Excellence in Research), January 1992.
(7 copies appended to this report)
352. Annual Report for 1991-1992. A Bibliography of Titles, Report of Work in Progress and Work Completed (Items 340-353) for the period of 1991-1992 of Research Grant Number DE-FG05-87ER 60503.
(7 copies submitted to The United States Department of Energy, Office of Health and Environmental Research on April 30, 1992).
353. Comprehensive 3-Year Report 1989-1992. A Bibliography of Titles Submitted in 1989-1992. A Brief Resume of Research Activities and Principal Results Obtained During 1989-1992, with Comments on the Research Performed under Contract No. DE-FG05-87ER 60503.
(7 copies submitted to The United States Department of Energy, Office of Health and Environmental Research, on April 30, 1992).

C. SYMPOSIA/SEMINARS/REVIEWS

1. L. Klasinc, Lj. Pasa-Tolic, S. P. McGlynn and T. P. Zivkovic ac, "Low-Rank Perturbation Approach Applied to Steroids, " IV. International Conference on Mathematical and Computational Chemistry, in Bled, Slovenia, June 4th, 1991.
- - -Invited Lecture
2. S. P. McGlynn, "Spin Uncoupling In Rydberg States," IV. International Conference on Mathematical and Computational Chemistry, Bled, Slovenia, June 5th, 1991.
- - -Plenary Lecture
3. S. P. McGlynn, G. Findley and A. Findley, "The Geometry of Genetics", IV. International Conference on Mathematical and Computational Chemistry, Bled, Slovenia, June 6th, 1991.
- - -Plenary Lecture
4. S. P. McGlynn, G. Findley and A. Findley, "The Geometry of Genetics", to the faculties of Biochemistry and Biophysics, The University of Split, Split, Croatia, June 11th
- - -Invited Lecture
5. S. P. McGlynn, "Scattering of Rydberg Electrons", given to the faculty of Arts & Sciences at The University of Split, Split, Croatia, June 13th, 1991.
- - -Invited Lecture
6. S. P. McGlynn, "Technology Transfer: University to Industry," given at the University of Ljubljana, Ljubljana, Slovenia and the Ministry of Science, Research and Technology of the Republic of Slovenia on June, 14th, 1991.
- - -Invited Lecture
7. L. Klasinc, "Quantum Chemical Calculations on Steroids," given at the International Conference on Mathematics, Chemistry and Computers, Dubrovnik, Croatia, June 24-29, 1991.
- - -Invited Lecture
8. S. P. McGlynn, "Molecular Lesion Spectra: Radiation Signature and Biological Gateway" Workshop on Biophysical Modelling and Radiation Effects. Organized by the U. S. Department of Energy and The Radiation Protection Research Program of the Commission of European Communities, Padua, Italy, September 1-5, 1991.,
- - -Presentation

9. S. P. McGlynn visited the new synchrotron research center in Trieste, Italy on September 6th and 7th, 1991. This center, funded by the U. N. and Italy, hosted a discussion concerned with a joint U. S. /Eastern European/Italian research program on radiation signatures. Professor McGlynn was one of the discussants.

- - -Discussant

10. S. P. McGlynn reviewed the research program of the Radiation and Solid State Laboratory of New York University, October 17th, 18th and 19th, 1991; on behalf of the U. S. Department of Energy. His co-reviewers were Harel Weinstein (Mt. Sinai Medical Center) and A. W. Castleman (Penn State).

- - -Reviewer

11. D. Kumar and S. P. McGlynn, "Dependence of Optogalvanic Signal on Laser Excitation Region in a Low Power Neon RF Discharge," 44th Annual Gaseous Electronic Conference, Albuquerque, New Mexico, October 22-25, 1991.

- - -Poster

12. S. P. McGlynn, K. Rupnik, M. N. Varma and L. Klasinc, "Radiation Signatures," Fourth Winter Conference of the Inter-American Photochemical Society at Clearwater Beach, Florida on January 1-5, 1992.

- - -Chairman

13. S. P. McGlynn, K. Rupnik, M. N. Varma and L. Klasinc, "Radiation Signatures for Biology," Sanibel Symposia on *Atomic, Molecular and Condensed Matter Theory*, St. Augustine, Florida; March 14-21, 1992.

- - -Poster

14. T. Pasinszki, T. Veszpremi, M. Feher, B. Kovac, L. Klasinc and S. P. McGlynn, "The Photoelectron Spectra of Methyl Pseudohalides," Sanibel Symposia on *Atomic, Molecular and Condensed Matter Theory*, St. Augustine, Florida; March 14-21, 1992.

- - -Poster

D. OTHER ACTIVITIES

(i) DEDICATED SYNCHROTRON BEAM LINES FOR RADIATION
SIGNATURES

A primary concern of biological modelling is the relationship between radiation and cancer. This concern has lasted more than fifty years and has not produced much success. Indeed, Dr. K. H. Chadwick (Commission of European Communities) reported recently (Chadwick, 1991)* that the subjects of radiation safety and risk evaluation had effectively become a fully political domain: the scientific disagreement on issues and conclusions within the European Community had become so pervasive that the EC officialdom, being forced to enact standards, had to do so in the absence of much credible scientific input. He followed this statement with a plea to the European Radiation Science community to reclaim authority in its chosen area. Dr. M. N. Varma (U. S. Department of Energy), who followed Dr. Chadwick, stated categorically (Varma, 1991)* that present approaches and concepts were not adequate, that they had in fact failed, and that a new paradigm was necessary in order to revive the field and give it direction.

Following the Padua meeting, at which the comments by Chadwick and Varma were made, Dr.'s McGlynn and Leo Klasinc went to the Synchrotron Trieste (Trieste, Italy) where they participated in discussions of plans to fund and construct a beam line dedicated to studies of biological radiation damage caused by photons in the range $\lambda > 4\text{\AA}$. These plans have been formalized in a preliminary way and a copy of them is appended. A preliminary list of beam-line users is also appended.

*"Workshop on Biophysical Modelling and Radiation Effects, "Organized by the U. S. Department of Energy and the Radiation Protection Research Program of the Commission of European Communities, Padua, Italy, September 1-5, 1991.

Radiation Signatures Bending Magnet Beamline

Each bending magnet on the ELETTRA storage ring is intended to be the source of 3 beamlines, with each beamline taking 12 mrad of the horizontal beam divergence. The beam from the bending magnet is split immediately after the light port, before any of the major front end components. Each beamline will therefore have to pay the full costs for a front end and a beamline control system. The major components of the beamline as a whole are as follows: Front End, Beamline Control System, Beam Transport System, Monochromator and Experimental Station. These sections will be discussed individually.

(1) **FRONT END:** The front end basically consists of all the components between the light port and the shielding wall. These are mainly safety features, designed to protect both people and the storage ring. The distance from the bending magnet source to the light port is approximately 2 m, and the distance from the light port to the shielding wall (the "Front End") is approximately 6 m. The shielding wall is about 1.5 m thick. All components before the light port are considered to be part of the machine and are therefore paid for by Sincrotrone Trieste. The main components in the front end are a safety magnet, safety shutter, fast valve, beam stopper and residual gas analyser. In addition there are a number of gauges, pumps, valves, bellows, connecting pipes, crosses and flange adaptors. The cost of a front end is likely to be 200 ML.

(2) **BEAMLINE CONTROL SYSTEM:** The beamline control system hardware is a VME crate and controls all of the beamline up to the experimental station. If the experimental station has its own control system (e.g. a FC or similar) this can be interfaced to the VME crate, to give control of the monochromator and beamline vacuum system. The beamline control system controls all of the front end safety features, and is also interfaced to the machine control system in order to monitor various beam parameters. The software for the beamline control system is being paid for by

Sincrotrone Trieste, but the hardware, i.e. the VME crate, has to be paid for from the beamline budget. This is likely to cost 50 ML per beamline.

(3) **BEAM TRANSPORT SYSTEM:** The beam transport system includes all the components between the shielding wall and the entrance slit of the monochromator, and all components between the exit slit of the monochromator and the experimental station. In reality this means just two major components: a prefocusing mirror chamber before the monochromator and a higher order suppressor after the monochromator. Since neither of these two components require high precision optics or alignment, the total cost of the beam transport system (including vacuum equipment) should be approximately 200 ML.

(4) **MONOCHROMATOR:** The monochromator for the radiation studies beamline has to be fairly special, since it is required to cover the energy range of approximately 20-2000 eV. A number of gratings will be required, although there is no need for a very high resolving power. A very approximate guess for the cost of this monochromator, including optics, mechanics and vacuum system, is approximately 500 ML.

(5) **EXPERIMENTAL STATION:** The experimental station is the area where the specimens are placed to be irradiated. It also includes the experimental "hutch" around the experimental area and any facilities needed at the beamline itself. Costing of an experimental station is very difficult until we know what facilities will be required. Since the project does not involve vacuum, complex mechanical systems or intricate detectors, a reasonable estimate could be 250 ML.

The total cost for this beamline will be approximately 1.2 BL, about 1 million USD.

This cost estimate assumes that all the complex laboratory facilities that will be required for the biological projects will be provided free of charge either from PolyBios, ICGB or the Sincrotrone Trieste BSL (Biological Support Lab).

15-20 m

Higher-Order
Suppressor

Monochromator
Chamber

Exit
Slit

Experimental
Station

Shielding
Wall

Prefocusing
Mirror
Chamber

Beam
Stopper

Fast
Valve

Safety
Shutter

Safety
Magnet

BENDING
MAGNET

PRELIMINARY LIST OF BEAMLINE USERS

Professor S. P. McGlynn	LSU, Boyd Professor of Chemistry.
Professor L. Klasinc	LSU and University of Zagreb, Croatia.
Dr. K. Rupnik	LSU.
Professor J. Battista	LSU, Microbiology.
Professor K. Pisk	Zagreb, General Director Ruder Boskovic Institute, Croatia.
Dr. D. Morris	Sincrotrone Trieste (ST), Physicist.
Professor Falaschi	Director International Center for Genetic Engineering and Biology, (ICGEB) Trieste, Italy .
Professor Rindi	Health Physicist at Sincrotrone Trieste
Dr. G. Tromba	Health Physicist at Sincrotrone Trieste
Professor Rosei	Director, Sincrotrone Trieste
Dr. M. Folkard	Radiation Biologist The Gray Laboratory, England.
Dr. G. Foster	Medical Physicist King's College, London, England
Professor Bettazoni	Molecular Biologist; ICGEB
Poly Bios Laboratory	a Biological Polymer Laboratory on the same site as Sincrotrone Trieste but independent of it.
Royal Holloway & Bedford New College London, England	3 Biologists with immediate interest in water window soft X-rays: Dr.'s Tom Ford, Tony Stead and Robin Cotton.

(ii) LABORATORY GUESTS

1. Russell Armstrong, Louisiana Stimulus for Excellence in Research (LaSER) Young Scholar Program, Summer 1991.
2. Professor John Murrell, School of Chemistry & Molecular Sciences, University of Sussex, Falmer, Brighton, United Kingdom, November 11-14, 1991.
3. Professor Danko Bosanac, Department of Physical Chemistry, Ruder Boskovic Institute, Zagreb, Croatia, December 12-14, 1991
4. Professor Tohru Azumi, Department of Chemistry, Tohoku University, Aoba-Ku, Japan, December 24-28, 1991.
5. Dr. Uri Asaf, Racah Institute of Physics, The Hebrew University of Jerusalem, Jerusalem, Israel, February 29 - March 14, 1992.
6. Dr. Ante Graovac, Ruder Boskovic Institute, Zagreb, Croatia, March 27-28, 1992.
7. Professor J. V. Knop, Director, Computing Center, Heinrich-Heine University, Dusseldorf, Germany, April 23-24, 1992.

(iii) RECOGNITION AND AWARDS

1. Dr. K. Rupnik was selected United States National Representative for the International Society for Theoretical Chemical Physics.
2. Dr. S. P. McGlynn is a founding member of The International Society for Molecular Electronics and Biocomputing (ISMEBC) which was established in September, 1991.
3. Dr. L. Klasinc is a founding member of The International Society for Molecular Electronics and Biocomputing (ISMEBC) which was established in September, 1991.

E. PRESENT PERSONNEL

Tanya Ducre

Student Worker

T. Eaton

Visiting Assistant Professor
McNeese State University
Lake Charles, Louisiana

Margaret Edwards

Secretary

W. S. Felps

Senior Research Associate

L. Klasinc

Visiting Professor
Ruder Boskovic Institute, University of Zagreb
Zagreb, Croatia

D. Kumar

Associate Professor-Research

S. P. McGlynn

Boyd Professor of Chemistry

R. C. Mohanty

Visiting Professor
Southern University
Baton Rouge, Louisiana

K. Rupnik

Research Associate

J. D. Scott

Senior Staff Scientist
Center for Advanced Microstructures
and Devices
Louisiana State University

ITEM 2. SCIENTIFIC REPORT

A. RADIATION SIGNATURES

The Radiation Signature Paradigm

A major goal of biophysical modelling is to relate physical events such as ionization, excitation, etc. to the production of radiation carcinogenesis. All types of radiations are of concern since each one is potentially carcinogenic. This concern has lasted more than fifty years and has had very limited success. It is now quite clear that new paradigms are necessary and these must include the biological, chemical and physical aspects of radiation-matter interactions.

A description of the physical events is provided by track or ion distribution structures. The track structure is a function of radiation quality, and it can be considered to be the "physical signature" of the radiation. Unfortunately, the uniqueness characteristics of these signatures are dissipated in biological systems in $\sim 10^{-10}$ s. Nonetheless, it is our contention that this physical disturbance of the biological system eventuates later, at $\sim 10^0$ s or greater, in molecular lesion spectra, which we dub the "lesion signature" or "radiation signature" and which also characterize the causal radiation.^{1,2}

The best way to understand the radiation signature paradigm is by inspection of Fig. 1, which represents a spatio-temporal path from a normal, just insulted cell to a fully transformed carcinogenic cell. Much radiation physics work has been concentrated at the extreme lower left of this pathway: track structures at the 10^{-10} s level and ion distribution at the 10^{-6} s level. Unfortunately, the observed tracks are too far removed, in a spatio-temporal sense, from clinical manifestations of cancer. Clinical manifestations of cancer may take more than 20 years to develop, which places any possible radiation signature based on malignancy or mutation type at 10^8 to 10^9 s. Thus it becomes necessary to follow the radiation track through a whole host of chemical, biochemical, biological repair and emergent cellular happenings before its manifestation as cancer. The radiation signature paradigm is aimed at finding an intermediate point on the spatio-temporal path. We suggest that the most appropriate place to seek such a "signature" is at the junction of the chemical and biological regimes. Such a point is shown in the middle of Fig. 1. If such a signature exists at this point and provides identification capability for the causal radiation, it should also serve as a gateway point for the bioprocessing which ultimately ends in malignancy. The correlation of the two ends of the spatio-temporal path might then be significantly facilitated by relating both ends to the midway point.

It is our contention that such a signature is provided by the spectrum (or histogram) of DNA molecular lesions. These signatures have the advantages of (1) high fidelity, (2) persistence times some ten orders of magnitude greater than track structures, and (3) a space-time cellular position which is approximately half way between initial insult and ultimate biological dysfunction.

Molecular lesion spectra contain information about the mechanism of radiation damage at the molecular level. That is, the types of lesions produced by a given radiation surely reflect the variety of different physico-chemical-biological processes that take place after radiation insult. In fact, some lesions may be a result of some particular radiation: for example, double strand breaks and multiply locally damaged sites that are mainly (but not solely) associated with high density radiations.

In this work, in order to clarify terminology in radiation signature research we introduce certain definitions:

(i) Molecular Lesion: A molecular lesion in DNA is defined differently in different fields. For example, definitions at the organismal, cellular and sub-cellular levels vary widely. We shall adopt an operational definition, and take a molecular lesion in DNA to be any damage, whether constitutional or conformational, that can be measured reproducibly by different investigators. It is implied, then, that a good assay method exists for the lesion. It is also implied, for now anyway, that irradiations and measurements be done in the vicinity of 0°C since, otherwise, differential repair rates might vitiate reproducibility.

(ii) Sub-lesion: We suppose sub-lesions to be post-radiative damage usually consisting of loci at which energy has been deposited or ionizations have occurred. We suppose them to lie in the time domain in which physics has terminated and chemistry (but not biology) has initiated. The important point is that they are unstable and that processing of them may produce more than one type of molecular lesion. That is, branching may take place.

(iii) Molecular Lesion Spectra: We define a molecular lesion spectrum (MLS) as a plot of lesion frequency (or probability), usually in bar graph form, versus lesion type (categorized, for example, by indices 1,2,3,...). The sum of bars is normalized to 100. The set of lesions should be complete; that is, every lesion, regardless of how low its frequency, should be represented. Reality, however, intrudes: we have neither detected all possible lesions produced by a particular radiation nor learned to perform reliable assay for the limited set that has been detected. Thus, we must make do with a constrained definition which replaces "complete set" with "maximal detected set that can be reliably assayed".

(iv) Radiation Signature: The term radiation signature implies the existence of a set of post-irradiation properties of DNA that can be used to identify the radiation which caused them. The search for such a property set is an urgent desideratum of the radiation safety, insurance and environmental fields. It is our contention that molecular lesion spectra provide such a signature. Any individual possesses a variety of signatures: the "name" signature is used to validate a contract; the bank card or microchip to obtain money from an automatic teller; the metallized parking card to gain entry to an automobile lot; or a voice recognition system to open a pension door. In any event, signature and purpose must be related. The purpose of this work is to define a radiation signature in DNA which not only serves as an identifier of the radiation which produced the signature but also functions as an appropriate gateway to the cellular processings which ultimately result in cell carcinogenesis. We contend that molecular lesion spectra serve as such a nexus.

(v) The Spatio-Temporal Pathway: There is a path (perhaps many paths) from normal to carcinogenic cell. The point of reference for a signature based on ion distributions is $\sim 10^{-7}$ s, at the lower left of the curve. The point of reference for a signature based on carcinogenesis (i. e., on medicine) is years. These two points are too far removed in both organismal (or cellular) space and in time to permit other than an allusive correlation of one with the other. It is our contention that an intermediate point provides signature for one end (i. e., radiation) and gateway to the other (i. e., cancer), and permits a real two-fold correlation which, in a single end-to-end correlation, might forever remain suppositional. We contend that molecular lesion spectra fit those requirements.

PRESENT RESEARCH DIRECTIONS

In this work, which is part of a long-term radiation signature study, the focus is on molecular lesion spectra as radiation identifiers. The objects of the research are: (1) to demonstrate that MLS can serve as radiation signatures for various radiations and to study the "sensitivity" and precision of these signatures; (2) to expand the data bases for such research; and (3) to develop tools for feature recognition and extraction.

We have shown that these goals are realistic; however, any attempt to formulate relations between signatures and carcinogenesis must remain futuristic.

Indeed, we intend, at least for now, to keep the accent solely on molecular lesion spectra as signatures of a causative radiation.

The experimental data bases for the great majority of radiations are very fragmented, so much so that a systematic study of radiation signatures is largely inhibited. The situation would improve enormously if more and better data, such as those of the Peaks³, Setlows⁴ and Ward⁵, were available. However, the advent of new experimental techniques, such as the polymerase chain reaction⁶ or laser technology^{7,8} promise much in unraveling the difference between high and low intensity radiation effects, in generating new data on the intensity-dependent photochemistry of DNA, in elucidating the effects of biphotonic processes⁹, etc. In sum, the systematic elucidation of lesion types and frequencies, with proper regard paid to dosimetry (e.g., energy absorbed, cell concentration of insult, etc.) and biology (e.g., *in vitro*, *in vivo*, cell cycle stage, etc.), is a massive task that must be engaged in order to put the concept of spectrum as signature on a firm footing.

The concept of a signature (i.e., feature recognition) implies the existence of a set of object characteristics that can discriminate between different objects and/or classes of objects. A signature should possess such characteristics as repeatability (i.e., confidence) and ease of recognition (i.e., ease of discrimination from threshold "noise"). However, since the characteristics of a signature need not always be explicitly circumscribed by rules, some element of adaptability in the models used for feature extraction, particularly when the experimental data are collected under conditions that are exceedingly difficult to standardize, appears to be mandatory. For that reason, we have also introduced methods for the extraction of radiation signatures from diverse data sets: that is, an adaptive lesion model, which originates in parallel distributed processing (PDP) methods and which is related to adaptive nonlinear (neural) networks (ANN) and/or genetic algorithms (GA). It might be argued that the use of feature (or pattern) recognition techniques in the present instance is redundant in the cases where the MLS are visually different. One answer is that visual differentiation is a qualitative sorting, and that one should make the sorting quantitative. To that end, each of the MLS is viewed as a "vector" in a space spanned by the MLS (or patterns), and the distinctiveness of patterns is quantified as the degree of orthogonality of the representative vectors. A second, and more relevant answer is that feature recognition techniques become necessary when one deals with data bases that are less than optimal¹⁰, when one queries the minimum number of lesions adequate for a signature¹⁰, or when one pursues any presumptive connection of a causal radiation and clinical outcome.

Our most complete study concerns the three electromagnetic radiations presented below.

THE SPECTRUM OF MOLECULAR LESIONS

Somewhat complete assays exist for only three different electromagnetic radiations, namely

Radiation	Energy	Source of data
VIS (limit of UV-A)	405 nm	Peak and Peak, (1987) ³
UV-C	254 nm	Setlow and Setlow, (1972) ⁴
γ -RAY	¹³⁷ Cs (~0.6 MeV)	Ward, (1988) ⁵
	⁶⁰ Co (~1.1 MeV)	

The molecular lesion spectra (or histograms) for these three radiations have been constructed and are given in Figure 2 as bar graphs of lesion frequency versus lesion type. The bar sum in any one graph is arbitrarily normalized to 100 .

The molecular lesions assayed in the three data sets are

Index	Name	Abbreviation
1	cytidine hydrate	ch
2	thymine dimer	d
3	pyrimidine adduct	pa
4	DNA-protein cross-link	dpc
5	DNA-DNA cross-link	ddc
6	single strand break	ssb
7	double strand break	dsb
8	locally multiply damaged sites	lmds

Indeed, these eight lesions are those for which assay data are most generally available. The source authors^{3,4,5} and Cadet and Vigny¹¹ should be consulted for definitions of the various kinds of lesions.

The three data sets refer to the DNA of various cells and of different biological entities. All data refer to the number of lesions produced by an F37

(D37) radiation fluence. Intercomparison of the three data sets, as is done in Fig. 2, must be accompanied with certain reservations:

- - Peak and Peak³ studied DNA lesions in mammalian cells at 405 nm. The F37 fluence was equivalent to $\sim 4 \text{ MJ/m}^2$. The cell line was a cultured P3 teratocarcinoma strain.
- - The Setlow and Setlow⁴ data for 254nm refer to vegetative mammalian cells at room temperature, again at an F37 fluence which, in this case, corresponded to 10 J/m^2 .
- - The Ward⁵ data also refer to mammalian cells at an F37 dose. However, in this instance of ionizing radiation, the kill efficiency was much higher than for either 405 or 254nm. That is, in a relative sense, the γ -rays produced low levels of DNA damage. The high mortality level is apparently attributable to the ability of γ -rays to produce damage in which more than one moiety in a localized region of the DNA suffers damage, the so-called "locally multiply-damaged sites" ((lmds))⁵. It is important to point out that the (lmds) category, index 8, encompasses a large variety of lesions, many of them categorizable under indices 1 through 7.

The defects then, are: (i), the supposition that F37 is an appropriate standard for radiation dosage; (ii), the use of different cell lines under different experimental conditions, both physical and biological; (iii), the lack of fluence considerations, and a disregard for the manner in which lesion interactions relate to fluence and fluence rate on the one hand and mortality and lesion production on the other.

Intercomparison of the MLS of Fig. 2 is revealing. The 405nm distribution is quite similar to the γ -ray distributions if the (lmds) lesion category, category 8, of the latter is disregarded; and both the 405nm and γ -ray distributions are markedly different from the 254nm spectrum. According to radiation quality considerations, on the other hand, it is the 405 and 254nm distributions which should exhibit similarity, and both should be quite different from the γ -ray distribution. It is clear, then, that radiation quality does not provide an apt classification of DNA damage at the molecular level, and that the nature and loci of the energy deposition events is the significant descriptor. In specific, the 254nm distribution reflects direct absorption by, and deposition in the DNA bases, events which are totally impossible at 405nm and statistically improbable at $\sim 1 \text{ MeV}$; the similarity of the 405nm and γ -ray spectra, the (lmds) category of the latter excepted, is probably a consequence of energy deposition in the aqueous environment, directly for γ -rays and through the intermediacy of some third-party-chromophore for 405nm.

The construction of a portfolio of trustworthy radiation signatures is inseparable from the existence of reliable, quantitative data sets of molecular lesions. The imposition of the reliability consideration probably implies that repair processes be inoperable. That is, irradiation must be made at temperature near 0°C in order to inhibit repair. Enzymatic repair rates are functions of lesion type and, hence, the relative concentrations of lesions types may change with post-irradiation time. Should that be the case, the fidelity of radiation signatures would be diminished.

Apart from a subcategorization of existing lesion sets (e.g., the dimer set), the signature fidelity would also be enhanced by the discovery and assay of new lesion sets. In view of the importance of the OH· radical and other oxygen-hydrogen entities, it is surprising, for example, that nitrogenous species are not implicated in the indirect production of molecular lesions. It is known that irradiation of protein produces generous amounts of NH₂·, a radical isoelectronic with and no less reactive than OH·, yet lesions produced by NH₂· seem not to occur or, at least, not to have been detected^{12,13}.

At this time, there is no unique, generally acceptable standard for biological dose. The data used in this work are based on a biological normalization related to the "efficiency of killing", namely the dose (D37) or fluence (F37) which permits a survivor fraction of 37%. This type of normalization presents at least two difficulties. Firstly, it appears that the ratio of lesion production to cell mortality may be a function of the radiation. For example, if 405nm is, as it appears to be, inefficient in killing, a D37 dose at 405nm implies measurement of the lesion production at very high fluence. That is (D37/F37) for the γ-ray region may well be more than 10⁶ times larger than (D37/F37) for the 405nm region. This question, in turn, generates the second difficulty: if the quadratic component in the linear-quadratic dependence of effect on dose¹⁴ is dominant, or merely important, the distribution of lesion intensity in the lesion spectra will vary with dose (and fluence), and the concept of a signature will run into difficulty. Thus, it is imperative that the effects of fluence on lesion distribution be investigated in detail. If it is found that lesion distributions are dose dependent, this will require an extension of the concept of a signature promulgated here to one which acquires multidimensionality in the sense that, for electromagnetic radiation of wavelength λ , the signature $S = S(F)$ will have to be replaced by $S = S(\lambda, F)$. In terms of the adaptive PDP approach advocated here, such a change poses no problem. In fact, it should produce better signature discrimination characteristics. However, it surely would increase the

experimental workload, and reasons of economy alone could cause the signature concept, even if fully validated in a multidimensional sense, to lose utility.

Two final points remain. It is now known¹⁵ that the ability of UV to damage a given DNA base is determined by base sequence in the vicinity of the photoproduct, and by the flexibility of DNA at the photoproduct site. This raises the unwholesome spectre of a lesion spectrum for which $S = S(\lambda, F, d)$, where d represents a complete or partial enumeration of DNA sequence data. The second point concerns the rate of fluence delivery to the sample. The γ -ray data set used here contains an (lmds) lesion type which is thought to categorize highly-ionizing radiations. Now, if the rate of a fixed fluence delivery can be increased, say by a factor of 10^6 in a high power laser beam, to the point where multiphoton events become significant or even dominate, there exists the possibility that even 405nm spectra will contain an (lmds), index 8, lesion type. Such an occurrence would do much to destroy the existing orthogonalities of the three MLS patterns of Fig. 2.

ADAPTIVE NEURAL NETWORKS FOR RADIATION SIGNATURES

The MLS of Fig. 2 appear to be excellent radiation signatures. This conclusion, however, is a subjective, visual judgement and, in order to quantify it, one must resort to feature (or pattern) recognition approaches which provide some measure of pattern distinctiveness.

Many neural network algorithms and architectures, all of them originating in adaptive parallel distributive processing (PDP)¹⁶, have particular advantage in problems involving mappings, topologies and feature classifications. We have chosen to use a counterpropagating network¹⁷, the features being cast as vectors such that the measure of feature distinctiveness becomes a matter of vector projections and orthogonalizations. In the present instance, if each of the 3 patterns (i.e., the 3 MLS) constitutes an orthogonal basis vector for a 3 vector-space of patterns, the patterns can be considered to be unique radiation identifiers. On the other hand, as the degree of non-orthogonality of the 3 patterns grows, so also does their lack of uniqueness as radiation signatures.

The NeuralWare version¹⁸ of the adaptive, counterpropagating network is used here. This is an open, well-described system which is both upgradeable and expandable, and which has been tested in feature recognition operations. The workings of this network are shown in Fig. 3. The network consists of processing elements (PE), represented by squares in Fig. 3, linked by adjustable weight

connectors, shown as lines in Fig. 3. The basic operation consists of the network learning how best to discriminate the different features. During this learning (or training) period, the magnitudes of the weight connectors alter, or adapt, in order to produce the best weighted classification. Information passes up from the lowest (input) layer through intermediate (hidden) layers to the output layer by means of summations and linear or non-linear transfer functions. In a vector sense, rotation of the weight vectors is made to occur around a unit sphere until they are at the center of those input vectors that create a maximum output 19,20,21. Further processing selects one PE (with closest weight vector and highest output) as a "winner", in which case it is given a value of "1"; non-selection, on the other hand, is assigned a value of "0"; and the final results are coded in a standard vector form (0, ..., 1, ..., 0), etc.

Three molecular lesion spectra, each consisting of as many as eight lesion types, constituted the input. Each PE represents a particular lesion type and is the window through which the experimental occurrence frequency is introduced. The three outputs are shown in Fig. 3 and are (1,0,0), (0,1,0), and (0,0,1). Fidelity (i.e., orthogonality of the vectors) is practically complete (i.e., >99%). That is, the MLS, in a truly quantitative sense, are faithful radiation signatures.

There are a variety of network architectures available. The counterpropagating network used here happens to be particularly advantageous for handling data sets that are either restricted in the number of lesions for which frequency data are available or of mixed quality because of lesion assay methods. Other network architectures, such as the backpropagating network, are also under study. The results of this study,¹⁰ indicate that the radiation signature approach can indeed identify the causative radiation and also provide valuable information on the role that specific lesion types play in the radiation signature. As a result, we expect to be able to limit molecular lesion spectra to a "minimum set" of lesions and to define those lesions which are the crucial identifiers for a given radiation.

REFERENCES

1. S. P. MCGLYNN AND K. RUPNIK, How to model molecular lesions and Adaptive lesion model. *The 24th DOE Contractors' Meeting on Radiation Signatures: Facts or Fancy?*, Columbia University, New York, April 15-17th (1991).
2. S.P. MCGLYNN AND M. VARMA, Radiation signatures. In *Proceedings of the CEC Workshop on Biophysical Modelling of Radiation Effects*, Padua, Italy, September 2-5, Adam Hilger (IOP) Bristol, England, 1992 (in press).
3. J. G. PEAK AND M. J. PEAK, Comparison of initial yields of DNA-to -protein crosslinks and single-strand breaks induced in cultured human cells by far- and near-ultraviolet light, blue light and X-rays. *Mutat. Res.* **246**, 187-191 (1991).
4. R. B. SETLOW AND J. K. SETLOW, Effects of radiation on polynucleotides. *Ann. Rev. Biophys. Bioeng.* **1**, 293-346 (1972).
5. J. F. WARD, DNA damage produced by ionizing radiation in mammalian cells: Identities, mechanisms of formation, and repairability. *Progress in Nucleic Acid Research and Molecular Biology*. **35** (W.E. Cohen and K. Moldave, Eds.), pp. 95-125. Academic Press, New York, 1988.
6. M. J. MCPHERSON, G. R. TAYLOR, AND P. QUIRKE, PCR: A practical approach. Oxford University Press, New York, 1991.
7. I. E. KOICHEVAR, Cytotoxicity and mutagenicity of excimer laser radiation. *Lasers in Surgery & Medicine* **9**, 440-445 (1989).
8. I. E. KOICHEVAR AND L. A. BUCKLEY, Photochemistry of DNA using 193nm excimer laser radiation. *Photochem. Photobiol.* **51**, 527-532 (1990).
9. T.W. MASNYK AND K. W. MINTON, Formation of single and double strand breaks in DNA ultraviolet irradiated at high intensity. *Photochem. Photobiol.* **54**, 99-107 (1991).
10. K. RUPNIK, M. VARMA, S.P.MCGLYNN, AND L. KLASINC, in preparation for submission 1992.

11. J. CADET AND P. VIGNY, The photochemistry of nucleic acids. *Bioorganic Photochemistry* . Vol. 1, (H. Morrison, Ed.) , pp.1 -272 . J. Wiley & Sons, New York, 1990.
12. H. YOSHIDA , W. E. BLOCH, K. B. JACOBSEN AND J. E. TURNER, Measurement of free ammonia produced by X -irradiation of glycylglycine in aqueous solution. *Radiat. Res.* **121**, 257-261(1990) .
13. A. W. CASTELMAN, Clusters of Atoms and Molecules. *Private communication*, Department of Chemistry, Pennsylvania State University, University Park, PA 16802 .
14. K. H. CHADWICK AND H. P. LEENHOUTS , The molecular theory of radiation. Springer- Verlag, New York, 1981.
15. M. M. BECKER AND Z. WANG , Origin of UV damage in DNA . *J. Mol. Biol.* **210**, 429-438 (1989).
16. J. L. MCCLELLAND AND D. RUMELHARD , Parallel Distributed Processing. MIT Press, Cambridge, MA ,1988.
17. R. HECHT-NIELSEN, Neurocomputing. Addison-Wesley, Reading, PA , 1990.
18. NeuralWorks, *NeuralWare, Inc.*, Penn Center West, Building IV, Suite 227, Pittsburgh, PA 15276, 1991.
19. T. KOHONEN, Self-organization and associative memory. Springer, New York, 1988.
20. S. GROSSBERG, Structures of mind and brain. Reidel, Boston, MA, 1982.
21. B. WIDROW AND M. HOFF , Adaptive switching circuits. *IRE WESCON Convention Record, Part 4*, pp 96-104, August 23-26, 1960.

FIGURE CAPTIONS

FIGURE 1

The post-radiative spatio-temporal development of a cell from "normal" to "abnormal". The radiative insult occurs at the lower left. Clinical effects occur at the extreme upper right. The molecular lesion spectra introduced in this work are located about midway along the curve. In the same way as MLS are mapped onto the radiations, there exists the possibility that clinical effects might be mapped onto MLS.

FIGURE 2

Examples of molecular lesion spectra (lesion type versus lesion frequency) for 405nm, 254nm and γ -ray radiations. The numbers on the abscissae correspond to the lesion indices in the text.

FIGURE 3

Test of the trained counterpropagating network for 254nm, 405nm, and γ -ray radiations. This network is discussed in the text. Input occurs in the bottom layer (8 elements or lesions), normative and selection (competition) processes occur in the middle layers, and output occurs in the top layer. A normalized histogram of all variable weights in the network and the RMS error of the output layer during the process are shown at the top.

COMPUTATIONAL SPACE

FIGURE 1

B. PHOTOELECTRON SPECTROSCOPY OF BIOLOGICALLY ACTIVE MOLECULES

Carbonyl and Thiocarbonyl Lone Pair Ionizations:

Thionyl analogues of molecules containing carbonyl groups may be used to replace their biologically-active oxygen counterparts, and, in the process, to produce important new effects. From an electronic point of view, the main difference between the two sets of compounds is vested in the lower electronegativity of sulphur, which causes thio-compounds to exhibit a lower ionization potential (usually, a sulphur lone-pair ionization). Indeed, the high electron density on sulphur in the highest-energy, occupied MO of these compounds makes them preferable targets for electrophiles and oxidation agents in both chemical systems and living organisms. It is the importance of this latter property that makes study of their electronic structure imperative. Photoelectron (PE) spectroscopy, although restricted to the gas phase, is central to such studies.

We have measured the HeI PE spectra of some thioderivatives of oxamide, $\text{H}_2\text{N}.\text{CO}.\text{CO}.\text{NH}_2$: $\text{N}^{\text{O}},\text{N}^{\text{O}}$ -dimethyl-monothiooxamide (1), its analogue $\text{N}^{\text{S}},\text{N}^{\text{S}}$ -dimethylmonothiooxamide (2), dithiooxamide (3), N,N' -dimethyldithiooxamide (4), and $\text{N},\text{N},\text{N}',\text{N}'$ -tetramethyldithiooxamide (5). The results of CNDO/2 calculations, the PE spectra of the related compounds acetamide, thioacetamide, thiourea and their methyl-derivatives, and HeI/HeII intensity considerations are used to assign the spectra. HeI/HeII spectra for 5 are shown in figure 1.

Our previous PE studies of oxamides were concerned with the assignment of ionization events¹, the effects of N-alkylation on ionization energies^{2,3} and the energy separation of n^+/n^- ionization events⁴. These studies led to additivity rules for N-alkyl shifts, and the establishment of such rules as empirical guides for the interpretation of PE spectra^{1,3,5}. Even though oxamide is both planar and s-trans in the crystal and even though the N-alkyl oxamides are generally s-trans, there is no reason to expect that these compounds,

depending on their environment, will not exhibit some twisting around the carbonyl-carbonyl bond. Isaksson and Liljefors⁸ used PE spectroscopy to study the conformation of oxamides and, were able to demonstrate that the s-cis and s-trans conformations exhibited quite different PE spectra. In an extension of this research, Henriksen et al.⁹ studied some cyclic and open chain mono and dithiooxamides. They reported a relatively large splitting of both the ns^+/ns^- orbital pair (0.7eV) and the π_{SCN^-}/π_{SCN^+} orbital pair (~0.5eV) in the open chain N-alkyl-derivatives. These splittings, we believe, should be small because of the non-planar s-trans conformations and the essential, single C-C bond character of these compounds.

Henriksen et al.⁹ noted that the PES of N^O,N^O -dimethylthiooxamide and its N^S,N^S -analogue are replicated rather well by a simple addition of the spectra of thioformamide to dimethylformamide and dimethylthioformamide to formamide, respectively. Our analysis, which concurs in their conclusion, makes use of acetamide, thioacetamide and the N-dimethyl derivatives of formamide and thioformamide to achieve the same result. However, the Henriksen argument, namely that the additive property is a facet of the large energy differences of the component spectra does not hold. As an example, the π_{NCS} and π_{NCO} bands of N^O,N^O -dimethylthiooxamide are also found to be simple additions of those for dimethylformamide and thioacetamide, and they show no splitting even though they possess essentially the same energy in both component molecules. Thus, we disagree with Henriksen et al.,⁹ in the sense that we believe "additivity" to be a consequence of both the non-planar conformation and the low p- electron density on the carbon atoms of dimethylthiooxamide.

This interpretation also suffices to understand an additive behavior of the dithiooxamides (which, we suggest, also occurs). In specific, the splitting in dithiooxamide, in which we have assigned $ns^+,\pi_{SCN^-},ns^-,\pi_{SCN^+}$ bands, is

much smaller than in thiourea, although the former compound imposes no prohibitions on the existence of a planar conformation. The relatively high ng^+/ng^- splitting that has been observed actually speaks in favour of a nearly planar conformation. The HeII spectrum of dithiooxamide indicates substantial loss of intensity in the 8.48, 9.34 and 12.55 eV bands, which we assign as ng^+ , ng^- and σ_{CS} , respectively. A gain of intensity is observed in the 9.08, 9.5 (sh) 12.05 and 13.30 eV bands which, consequently, we assign to ionizations of π -type.

The primary evidence for two nearly-degenerate π_{NCS} levels in N-alkyl-dithiooxamides is provided by the HeII and HeI spectra of tetramethyldithiooxamide. The HeII spectrum indicates substantial loss of intensity in the first two ionization bands, as well as in the 12eV band which we attribute to one of the σ_{CS} ionizations. It is suggested, then, that this molecule is unlike dithiooxamide in that it is highly non-planar. The same, by comparison, holds for N,N'-dimethyldithiooxamide. The observation that π_2 lies above the two σ_{CS} ionizations is in agreement with the spectra of the other two compounds: this ionization band clearly lies higher than the two σ_{CS} events in the dimethyl derivative, and occludes one of them in the tetramethyl derivative. The lowest p-ionization event in these compounds is expected to lie around 14eV.

In conclusion, the present results indicate a weak ng^+/ng^- interaction and a nearly additive behavior of the two composite parts, at least insofar as the low energy π -ionizations of monothio- and dithiooxamide derivatives are concerned.

Fig.1. Expanded low ionization energy region of the HeI and HeII PES of 5.

REFERENCES

1. J. L. Meeks, H. J. Maria, P. Brint and S. P. McGlynn, *Chem. Rev.*, 75, 603 (1975).
2. J. L. Meeks, J. F. Arnett, D. B. Larson and S. P. McGlynn, *J. Am. Chem. Soc.*, 97, 3905 (1975).
3. J. L. Meeks and S. P. McGlynn, *J. Am. Chem. Soc.*, 97, 5079 (1975).
4. D. Dougherty, P. Brint and S. P. McGlynn, *J. Am. Chem. Soc.*, 100, 5597 (1978).
5. L. Klasinc and S. P. McGlynn, "The photoelectron spectroscopy of double-bonded CC,CN,NN and CO groups" in "The Chemistry of Double-bonded Functional Groups", S. Patai, ed., John Wiley & Sons, 1989.
6. E. M. Ayerst and J. R. C. Duke, *Acta Cryst.* Z, 588 (1954).
7. D. B. Larson and S. P. McGlynn, *J. Mol. Spectry.*, 47, 469 (1973).
8. R. Isaksson and T. Liljefors, *JCS Perkin* 2, 1815 (1980).
9. L. Henriksen, R. Isaksson, T. Liljefors and J. Sandstroem, *Acta Chem. Scand., Ser. B*, B35, 489 (1981).

Steroids:

We have had concern for some time for the electronic structure of steroids.¹ Theory and experiment suggest that ribbon-like orbitals of these compounds exist and that these orbitals can mediate the transmission of electronic effects over as many as ten saturated single CC bonds. As a result, we have been particularly interested in the long-range interactions of substituents located at different positions on the A and D rings of a steroid molecule, and their dependence on the presence of interpolated, localized double bonds. The experimental results, consisting of studies on fast intramolecular ET processes in steroid solutions² as well as PES, FTMS and ¹³C NMR investigations, and the theoretical results, consisting of semiempirical MNDO calculations, indicate the existence of an efficient long-range coupling between substituents located at the 3- and 16- positions of the steroidal framework.

Here, we wish to present ab initio SCF (STO-3G) results for 5 α -androstande, to compare these with experiment and with MNDO computations, to discuss molecular geometry and electronic structure, and to ascertain whether such concepts as "ribbon-like MO's", "T- and L-type MO's" and "electronic transmission effects", which originated in the MNDO algorithm, retain their validity in STO-3G.

Geometry:

Unfortunately, x-ray crystallographic data are not available for 5 α -androstande, and calculation and experiment can not be compared. MNDO calculations, were found to exhibit much larger average deviations of bond lengths than did STO-3G (especially for C-H bond lengths greater than 1.1Å). The bond angle values for C-C-C- angles were found to be generally smaller than the corresponding STO-3G values, and the H-atoms were generally "more" equatorial or "more" axial. The MNDO geometry was found to be more planar (i.e., the angle between the least squares fitted molecular planes

C2,C3,C5,C7,C8,C10 and C9,C11,C13,C14,C16 being 1.9° , while the angle between the C10-C19 and C13-C18 vectors is 10.5° ; Fig.2).

Figure 2. Superposition of *ab initio* STO-3G (front) and MNDO (behind) optimized geometries for the C-skeleton of 5a-androstane (adjusted to match at C7).

On the basis of the above results, one concludes that the simple STO-3G basis, which approximates each AO by a single contracted function designed to mimic an STO, gives quite good results for the 5 α -androstande geometry.

Electronic Structure:

According to its photoelectron (PE) spectrum, 5 α -androstande has two characteristic low energy ionizations of 9.21 eV and 9.49 eV which, assuming Koopmans' theorem, correspond to the release of electrons from the two highest occupied MO's of the molecule. These energies, both absolutely and relatively, are in excellent accord with the STO-3G results.

Qualitative arguments indicate that these two topmost orbitals contain C2p AO's aligned somewhat in the molecular plane. Indeed, MNDO calculation yields a HOMO with constituent C2p AO's aligned along the short molecular axis (T-type MO) and a SHOMO in which the C2p AO's are aligned along the long molecular axis (L-type MO). It is believed that such MO's, best described as a mixture of σ_{CC} and π_{CH_2} orbitals, and colloquially termed "ribbon orbitals," are essential for the transmission of electronic effects in large saturated polycyclic hydrocarbon molecules.

According to MNDO calculations of the androstanones, the T-type MO interacts with 3-substituents, while the L-type MO interacts with 17-substituents, the same exclusivity being retained for 3,17-disubstitution. The 16-substituent, however, interacts with both types of MO and, as a consequence, so also do dual substituents, one each in the 3 and 16 positions. This behavior leads one to assume that, in 3,16- disubstitution, long range electronic interaction should be heavily promoted. In the STO-3G calculation, the order of the two orbitals is reversed relative to the MNDO situation: that is, the HOMO is now L-type and the SHOMO is T-type. The electron distribution of these L-and T-type MO's is somewhat lumped on the inner B and C ring carbons according to MNDO

results, but is more evenly distributed over all four rings in the STO-3G calculation. In particular, the electron density at C3 of the L-type HOMO and C16 of the T-type SHOMO is considerably higher in STO-3G than in MNDO. Thus, the STO-3G calculations suggest ribbon-like MO's that are better suited for the transmission of electronic effects. They also suggest that such transmissions will be particularly effective in the case of 3,16-disubstitution. The energy inversion of L-and T-type MO's that occurs in the two computations is not really discriminatory because of the near accidental degeneracy of these ribbon-like MO's (Δ MNDO = 0.117eV and Δ STO-3G = -0.154eV). The salient point is that the STO-3G computation both validates and enhances the concept of ribbon-like MO's for both the SHOMO and HOMO of 5 α -androstane and, by extension, for steroids generally.

REFERENCES

1. N. S. Bhacca and L. Klasinc, Z. Naturforsch., 40A, 706 (1985);
 L. Klasinc, B. Ruscic, N. S. Bhacca and S. P. McGlynn, Int. J. Quantum Chem. QBS, 12, 161 (1986);
 T. Cvitas, B. Kovac, Lj. Pasa-Tolic, B. Ruscic, L. Klasinc, J. V. Knop, N. S. Bhacca and S. P. McGlynn, Pure Appl. Chem., 61, 2139 (1989);
 L. Klasinc, Lj. Pasa-Tolic, B. Kovac, and S. P. McGlynn, Int. J. Quantum Chem. Symp., 16, 331 (1989);
 Lj. Pasa-Tolic, B. Kovac, L. Klasinc, J. V. Knop, C. Kubli-Garfias and S. P. McGlynn, Croat. Chem. Acta, 62, 37 (1989);
 L. Klasinc, Lj. Pasa-Tolic, H. Spiegl, J. V. Knop and S. P. McGlynn, Computers Chem., 4, 287 (1990);
 L. Klasinc, Lj. Pasa-Tolic, H. Spiegl, J. V. Knop and S. P. McGlynn, J. Math. Chem., 8, 179 (1991) in press.
2. See G. L. Closs and J. R. Miller, Science, 240, 3673 (1988) and references cited therein.

C. LASER OPTOGALVANIC EFFECT

1. INTRODUCTION

The laser optogalvanic (LOG) effect is the change of electrical impedance of a plasma produced by resonant absorption of laser radiation by a plasma moiety¹. Plasma is a complex mixture of different species (neutrals, ions, etc.) in various states of excitation. Under steady-state conditions (e.g., in dc discharges), there is a dynamic equilibrium between these species and a well-defined impedance is offered to the flow of current in the plasma. Laser excitation of a species in the discharge results in a net deposition of energy in that species and, consequently, the dynamic equilibrium is disturbed. This perturbation results in a change in the impedance of the discharge and a LOG signal is generated. For radiofrequency (rf) discharges operating at a frequency f , a modulation of the dynamic equilibrium occurs at a frequency f (or $2f$, depending on the process and/or the life-time of the state relative to $1/f$). Laser induced perturbations in the steady-state equilibrium (for dc discharges) and in the modulated equilibrium (for rf discharges) can be picked-up by suitable circuitry and the LOG signal can be extracted.

The LOG effect is an extremely sensitive and powerful spectroscopic/detection tool with wide-ranging applications. It has important applications in (i) the spectroscopy of highly excited (i.e., Rydberg) states of atoms and molecules, (ii), the study of various dynamic processes in flames and plasmas, and (iii), in pollution monitoring and control as an extremely sensitive detection tool. Both dc and rf discharges may be used to generate the plasma for LOG studies. While dc discharges are simple to generate, they are unfortunately characterized by a non-uniform plasma (discrete discharge regions) and by the risk of electrode contamination by reactive gases/species present in the discharge. The rf discharge, on the other hand, has the advantage of spatial uniformity (except for a sheath in the vicinity of the electrodes), a

"contactless" nature (i.e., no metal electrodes inside the discharge tube), and background noise levels that are not critically dependent on either the gas pressure or gas type.²

Despite its importance, the mechanism of generation of LOG signals is not completely understood. This lack of understanding, which has severely impeded exploitation of the effect, is primarily due to the complexity of the plasma and a lack of knowledge about various rates/cross-sections, etc. The temporal profiles of LOG signals, which contain considerable mechanistic information, are crucial to the understanding of the various mechanisms in the generation of these signals. Unfortunately, the temporal profiles of LOG signals, are very complex and very sensitive to discharge conditions (e.g., discharge configuration, electrical input power, rf source characteristics, laser intensity, region of the plasma excited by the laser, etc.), and, presumably for those reasons, models for the LOG effect are either wrong or incomplete.

2. MODELS

The electrical impedance of a plasma depends on charge carrier density and mobility. Consequently, the two phenomenological models that have been proposed for the generation of LOG signal are: (i), direct or collision-assisted ionization rate changes (IRC) of the excited plasma atoms/molecules produced by laser excitation, which causes an increase of electron-ion pair density and, hence, increased conductivity;¹ and (ii), an electron temperature coupling model in which the increased electron kinetic energy, caused by laser energy deposition, causes an increase of discharge conductivity by enhancing electron mobility or collisional ionization rates.^{3,4} Mechanism (ii) was invoked in order to explain the generation of LOG signals by low energy (e.g., vibrational) optical excitation to levels far below ionization threshold. Mechanism (ii), though successful in the case of unchopped IR irradiation

of molecular discharges,⁴ fails to explain the following important features of LOG signals⁵⁻⁷:

- - -No correlation exists between LOG signal intensity and the energy gap from terminal laser-excited level to ionization threshold.
- - -No correlation exists between LOG signal intensities of various entities and their respective ionization potentials.

These correlative deficiencies were overcome when the optoacoustic (OA) effect was invoked as an intermediate mechanism in the production of LOG signals from such low-energy photons. Production of LOG signals via an OA effect has been known for over a decade⁸ and several simultaneous OA and LOG studies have been carried out.^{6,9} However, an important aspect of the mechanism by which the OA wave generates LOG signal was not correctly understood. Heating and cooling, which accompany the OA wave, were incorrectly supposed to generate the LOG signal via mechanism (ii) listed above.

New Results

Our recent¹⁰ work on LOG effects in rf discharges in iodine has shown that LOG signals generated via the optoacoustic (OA) effect need not involve thermal effects (mobility changes) or ionization rate changes in the final step of such signal production. A simple perturbation of the distribution of positive ions (or other charged species) by the pressure wave which accompanies an OA effect can generate an OA-mediated (OAM) LOG signal directly. As an example, the pressure wave induced displacement of the positive ion sheath (I^+) in the vicinity of the electrodes in a low power rf discharge in iodine vapor generates a LOG signal whose polarity is reversed with the reversal of the direction of propagation of the pressure wave. It is clear that the pressure wave generates these LOG signals by an actual physical movement of the positive ions in the sheath region (i.e., the "sensitive" region) of the discharge. The "sensitive" region, in which the OA wave generates a LOG signal,

may be confined to the vicinity of the electrodes (as in the case of low power rf discharge in iodine) or, it may be spread out over the discharge region (e.g., in a low power rf discharge in neon).

Thus, the LOG signal can be generated by two distinct mechanisms. These are (a), ionization rate changes (IRC); and (b), displacement of charged species in the "sensitive" region of the discharge. Mechanism (b) is usually OA-mediated (OAM component). Mechanism (ii), as listed previously, if it is at all operative, should have the same temporal profile as the OAM component except that the polarity of a LOG signal induced by a mechanism (ii) will not be inverted when the direction of propagation of the OA wave is reversed. We conclude that, under our experimental conditions, mechanism (ii) is either inoperative or negligibly small. Consequently, the temporal profile of a LOG signal consists, in general, of two physically distinct components: one attributable to ionization rate changes, the other to an optoacoustic effect.

We have developed a method which, under suitable conditions, identifies and disentangles these two distinct components in the temporal profile of a LOG signal even when they exhibit extensive overlap.^{11,12} This method is based on a technique which treats the plasma as an integral rf circuit component. The underlying reason for the success of this approach is the fact that the IRC and the OA wave do not perturb the electrical resonance characteristics of the rf plasma in identical ways.

3. EXPERIMENTAL SET-UP AND PROCEDURE

A flash-lamp pumped dye laser (Chromatix CMX-4, $\sim 1\mu\text{s}$ pulse width, resolution $\sim 0.1\text{cm}^{-1}$ with intracavity etalon and $\sim 3\text{cm}^{-1}$ without, beam diameter $< 0.5\text{mm}$) is used for plasma excitation. Most rf discharge studies have used chopped CW lasers and lock-in amplifiers, a configuration which is not suited for mechanism studies because the LOG response must be averaged over the chopper on-off period. The rf discharge/pulsed laser conjunction is configured such that the laser pulse

width, $\tau \simeq 1\mu\text{s}$ is much larger than the rf period, $\Gamma \simeq 33\text{ns}$. This configuration is readily achieved.^{10,11} Since the laser pulse samples a number of rf oscillations to produce a time-averaged signal, laser triggering need not be synchronous with the rf cycle. A resolution approaching that of the laser pulse width is readily achievable. This configuration is of higher sensitivity than the $\tau \ll \Gamma$ configuration and is better suited to extraction of the slow LOG signal components that concern this work. The $\tau \ll \Gamma$ configuration requires synchronization, and is particularly apt for field-effect studies and the resolution of high-speed transients.

A schematic representation of the apparatus is shown in Fig. 1. A discussion of it is available elsewhere.^{10,11} A number of points are worth noting here. First, laser excitation is transverse, not axial; this excitation geometry provides better linearity of LOG signal with laser intensity, it is often more sensitive, and it permits the probing of different plasma regions. Second, the resonantly-tuned pick-up coil is coupled to the rf discharge by virtue of extension of this discharge beyond the electrodes; laser excitation produces a change in the power transferred to the pick-coil and this change, which is the LOG signal, is recovered by demodulation in the signal recovery section. Third, the tuning and de-tuning of the driver/detector circuits to and away from resonance constitutes a significant tactic for separating the ionization and optoacoustic components of the LOG signal; these tuning/de-tuning techniques are discussed elsewhere.^{11,12}

The rf resonance profile of the driver/detector circuit is obtained by scanning the rf oscillator frequency f at constant amplitude and recording the dc voltage in the demodulator of the signal recovery section, all in the absence of laser excitation. Such a profile, pertinent to the apparatus of Fig. 1, is shown in Fig. 2.

The tactic referred to in the previous paragraph consists then of recording LOG signals at f_0 , the resonance maximum, at $f_0 - \Delta f$ and $f_0 + \Delta f$, where $\Delta f < 0.5\text{ MHz}$. Changes in the profile of the LOG signal may also be monitored as the rf frequency is

scanned across the resonance peak. The effects observed are vested in, and derive from the following statements about the nature of plasma and plasma perturbations:

- - -Plasma at equilibrium has a certain impedance.
- - -Since the plasma is coupled to the driving circuit, it controls the electrical resonance frequency of the combined driver/plasma circuit.
- - -Laser induced changes in the equilibrium ionization rate (IRC component) alter the plasma impedance (effective inductance or capacitance) because of an alteration of the electron-ion pair number density. This transient change in plasma impedance produces a transient shift of the electrical resonance frequency. An increase of ionization shifts the resonance response curve of Fig.2 to the left whereas a decrease produces a shift to the right.
- - -An optoacoustic effect causes no change in the electron-ion pair density and, hence, in the impedance (i.e., effective inductance or capacitance, the resonance determinative elements) of the plasma. Therefore, the resonance response curve of Fig. 2 will not be displaced to either left or right. The OAM LOG signal generation, is attributable, in this view, to a simple change of amplitude of the response curve.

In view of the above, the nature of the experiment is quite transparent. The solid curve of Fig. 2 represents the response curve of the steady state plasma. When the plasma is perturbed by laser excitation such that an increase of ionization occurs, the LC circuit characteristics are altered and the response curve, now shown dotted, shifts slightly to the left. If the rf frequency was initially at point P, laser excitation will now produce a transient positive LOG signal because the dotted curve is more responsive at P than the solid curve. Under similar conditions, but with the rf frequency initially at R, a transient negative LOG signal will be produced. And, if the rf frequency is initially at Q, even though the laser produces an increase of ionization, no signal will result. These changes are indicated by arrows in Fig. 2. They represent the expected behavior of the IRC component:^{11,12} null at Q, positive at P and negative at R.

The OAM component is produced not by an alteration of the rate of ionization but largely by changes in the distribution of positive charges and possibly in the mobility of electrons. Such changes, being largely redistributive in character, will not produce any left-right shift of the response curve but may change its amplitude. Thus, in contrast to the IRC component, which is nulled, the OAM component will be maximal at Q , and will decrease according to the response curve profile as the rf frequency varies on either side of Q .

Experiment, shown in Fig. 3, is in full accord with the above expectations. Since the OAM signal magnitude follows the response curve whereas the IRC signal magnitude follows the derivative of the response curve, the pure OAM signal may be observed at Q , where the derivative is zero (Fig. 3A). The pure IRC signal may only be recovered subtractively. Such "pure" IRC signals are shown in Fig. 4. It is important to note that our treatment is truly perturbative, in the sense of a small change of steady state conditions, and that a relatively tight coupling must exist between the plasma and the driving circuit in order for the plasma perturbations produced by the laser to affect RLC circuit behavior.

4. PLASMA PROCESSES

Plasma is a very complex state of matter about which our ignorance is extensive. Since the goal of this work is to initiate a discussion of LOG signals in terms of energy level diagrams for the plasma constituents, it is clear that a specification of those plasma processes critical to the LOG effect must first be elaborated. An attempt at such a specification now follows:

Direct Photoionization: An instantaneous process that should produce a "fast" IRC signal whose rise time is the convolution of laser pulse and detection electronics rise times (and, therefore, coincident in the present instance with the laser pulse rise time of $\sim 1\mu\text{s}$) but whose decay time will be largely controlled by the slow process of plasma re-equilibration. Plasma re-equilibration times are functions of the particular plasma,

the characteristics of the laser excited levels of the plasma constituents and the impedance of the electrical power source which controls the stimulus for re-equilibration.

Collisional Photoionization: An instantaneous process (relative, that is, to the 1 μ s pulse used in this work) because, at the pressure range ~1mTorr to few Torr prevalent in plasma, the collision frequencies are 1GHz or larger.

Metastable-Metastable Penning Ionization: A process which is usually symbolized

where M^* is the metastable excited state of the ground state plasma entity M ; this process dominates at low rf powers ($\leq$ few Watts) and moderate pressures ($\geq$ few Torr).^{13,14} Any perturbation of the M^* population will be followed by a slow collisionally-dominated re-equilibration process which may extend into the 10¹-10² μ s range. When dominant, as in a 5 Torr neon rf discharge powered at 1W to be discussed later, depletion of the metastable level by non-ionizing laser excitation will produce a decrease of the ionization rate^{13,14} and, hence, a negative IRC signal component. By the same token, given that collisions are the dominant re-equilibration mode, the decay characteristics of the level produced by laser excitation may also dictate an OAM component, which of course produces a positive OAM signal, and may control the trailing parts of both the OAM and IRC components of the LOG response.

Radiation Trapping: A process of sequential emission-reabsorption which is particularly important for high-extinction ground state absorption events in atoms (where, unlike molecules, there is no Stokes loss). For example, in an rf discharge in Neon at 5 Torr, the 1s₂ and 1s₄ states (Paschen notation) are so strongly trapped, that their effective lifetimes increase by a factor of $\sim 10^3$ (from ns to μ s).¹⁵ Since the 1s₂

(1s4) --->ground state photons lie in the VUV they can readily photoionize the M* or other excited levels and give rise to a delayed IRC component of the LOG signal.

Acoustic Effects: That part of the laser excitation energy which by non-radiative processes, whether collisionally induced or not, is converted into an acoustic shock wave that disturbs the equilibrium distribution of positive ions. Such a disturbance represents a "flow" of current and will give rise to a LOG signal. This acoustically-mediated (OAM) component has been misinterpreted in many works. Indeed, the temporal LOG profiles, recorded by many workers in both rf and dc discharges, are entirely OAM in nature.¹¹

Mobility Changes: A thermally-induced change of electron mobility that has been invoked, for example, to explain the change of plasma discharge current that occurs at onset of CO₂ lasing.³ These electron mobility effects are attributed not to electron kinetic energy changes but rather to collision rate changes dictated by thermally-induced density changes of atoms, molecules and ions in the plasma. If true, no change of the equilibrium ionization rate should occur and no IRC component should be produced. An OAM component could result but, if it did, it should not exhibit any directional dependence such as is observed in iodine plasmas.¹⁰ Thus, we can conclude that this component is certainly not particularly significant in the latter parts of the LOG signal.

5. REPRESENTATIVE SYSTEMS

The six effects of the previous section are not easy to disentangle. In general, then, the inventiveness of the experimenter is the dominant determinant in devising situations which either isolates one of the six effects or at least limits the number of such effects that go into the production of a given LOG signal. This is now demonstrated in a few situations:

Isolation of an OAM Component: The levels connected by laser excitation should not be significantly involved in the maintenance of the discharge, and the excited level should not be easily ionizable. A system best fitting these requirements¹⁶ is provided by a molecule which possesses low-lying excited states and for which the constituent atoms and atomic ions control the discharge characteristics. The I₂/I system fits these requirements and, in accord with expectation, pulsed laser excitation of the B $\leftarrow$ X transition of I₂ in an iodine plasma yields a LOG signal consisting solely of an OAM component. The term diagram appropriate to this situation is shown in Fig.5.

The use of a LOG detector as a sensitive, low-pressure "microphone" has been reported by us earlier.^{17,18} The OAM component was picked-up from an rf discharge in I₂/He mixture. The non-radiative energy detectivity was shown to be $\sim 10^{-9}$ J per pulse. Signal averaging is expected to improve this detectivity limit by a factor of 10^2 - 10^3 . This "microphone" has a wide dynamic range (no diaphragms to be displaced) and is contactless in nature (suitable for high temperature and corrosive environments).

Isolation of an IRC Component: Of the levels connected by laser excitation, the upper should be either in the ionization continuum or so close to threshold that collisional ionization is guaranteed whereas the lower level, for maximum signal, should not be M*. Examples of such situations, for pre-threshold ionization, are available in Xenon, Cesium and Iodine discharges. An appropriate term diagram is given in Fig. 5 (I atoms).

Isolation of a Trap Component: Of the levels connected by laser excitation, the upper (intermediate or I) state must be tightly connected to the trap level, T, whereas the lower state must be extensively populated (e.g., the metastable or M* state) in order to yield a relatively high density of T states. An example is available in the low-power

(0.2W) rf discharge in 5 Torr neon.¹⁹ The sequence of events which follows laser excitation are:

- - -Depletion of M^* population, causing a decrease of the ionization rate and onset of a negative IRC signal coincident with laser pulse.
- - -Non-radiative loss from state I which generates an OAM component more or less coincident with but possibly slightly delayed (say 1 to 2 μ s) relative to the laser pulse.
- - -Radiative trapping, following $I \rightarrow T$ processes, which gradually increases the ionization rate as $[T]$ increases. Thus, an IRC trap peak should appear at 5 -8 μ s following excitation. A delayed OAM component may also be launched.
- - -Since M^* / M^* collisions determine re-equilibration rates, long tails extending possibly to 100 μ s are expected in both the OAM and IRC components.

A representative term scheme is shown in Fig.6. The temporal profiles are given in Fig. 3 and 4.

6. CONCLUSIONS

The LOG effect is sensitive and versatile. It has enormous analytic potential, whether independent from or adjunct to ICP technologies. Its potential as a diagnostic for plasma processes, particularly as these become crucial to microelectronic fabrication techniques, is promising. As shown, plasma is also an excellent acoustic wave detector (i.e., microphone). And, finally, it is hoped this work demonstrates that appropriate experimentation can provide a detailed resolution of the processes important to plasma stabilization and maintenance.

REFERENCES

1. For a review of the optogalvanic effect and its various applications see (a) B. Barbieri, N. Boverini, and A. Sasso, "Optogalvanic Spectroscopy," *Rev. Mod. Phys.* 62, 603 (1990);
(b) C. R. Webster and C. T. Rettner, "Laser Optogalvanic Spectroscopy of Molecules," *Laser Focus* 19, 41 (1983);
(c) J. E. M. Goldsmith and J. E. Lawler, "Optogalvanic Spectroscopy," *Contemp. Phys.* 22, 235 (1981);
(d) "Optogalvanic Spectroscopy and its Applications," *J. Phys. Colloq.* C7, (1983).

2. R. E. Muenchausen, R. D. May, and G. W. Hills, *Opt. Commun.* 48, 317 (1984).
3. S. Mofatt and A. L. S. Smith, *J. Phys. D: Appl. Phys.* 17, 59 (1984).
4. C. R. Webster and R. T. Menzies, *J. Chem. Phys.* 78, 2121 (1983).
5. R. A. Keller and E. F. Zalewski, *Appl. Opt.* 19, 3301 (1980).
6. E. Armondo, M. G. DiVito, K. Ernst and M. Inguscio, *Opt. Letters*, 9, 530, (1984).
7. R. A. Keller, B. E. Warner, E. F. Zalewski, P. Dyer, R. Engleman, Jr., and B. A. Palmer, *J. Phys. Colloq.* C7, 23 (1983).
8. C. T. Rettner, C. R. Webster, and R. N. Zare, *J. Phys. Chem.* 85, 1105 (1981).
9. (a) C. Hameau, E. Arimondo, J. Wascot, and P. Glorieux, *Opt. Commun.*, 53, 375 (1985).
 (b) A. Sasso and M. G. DiVito, *Opt. Commun.* 66, 270 (1988).
10. D. Kumar, P. L. Clancy, and S. P. McGlynn, *J. Chem. Phys.* 90, 4008 (1989).
11. D. Kumar and S. P. McGlynn, *J. Chem. Phys.* 93, 3899 (1990).
12. D. Kumar and S. P. McGlynn, *Chem. Phys. Lett.* 176, 536 (1991),
13. G. A. Bickel and K. K. Innes, *Appl. Opt.* 24, 3620 (1985).
14. A. Sasso, M. Ciocca, and E. Arimondo, *J. Opt. Soc. Am.* B5, 1484 (1988).
15. Y. Uetani and T. Fujimoto, *Opt. Commun.* 49, 258 (1984).
16. C. Demuyne and J. L. Destombes, *IEEE J. Quantum Electron.* 17, 575 (1981).
17. D. Kumar, L. Klasinc, P. L. Clancy, and S. P. McGlynn, *IEEE Trans. Ultrasonics, Ferroelectrics, and Frequency Control* UFEC-33, 513 (1986).
18. D. Kumar, L. Klasinc, P. L. Clancy, R. V. Nauman, and S. P. McGlynn, *Can. J. Phys.* 64, 1107 (1986).
19. A. Ben-Amar, G. Erez, and R. Shuker, *J. Appl. Phys.* 54, 3688 (1983).

OPTOGALVANIC SET-UP WITH RF DISCHARGE

Figure 1.

Fig 2. Response curve (solid) in the absence of laser excitation and (dotted) when the plasma is laser irradiated.

Figure 3: TEMPORAL PROFILES, $V(f)$, of the LOG signal at three different rf frequencies:
 A. Top trace, $V(f_0)$; B. Middle trace, $V(f_0 - \Delta f)$; C. Bottom Trace, $V(f_0 + \Delta f)$; $f_0 = 29.24\text{MHz}$
 (resonance); $\Delta f = 0.2\text{MHz}$.

Figure 4: Temporal profiles of the IRC LOG component obtained by difference methods;
Upper trace, $V(f_o - \Delta f) - V(f_o)$; Lower trace, $V(f_o + \Delta f) - V(f_o)$.

Figure 5. Energy level diagram for I_2/I system.

Figure 6. Energy level diagram illustrating trap levels and generation of delayed IRC and OAM LOG effects attributable to trapping.

D. MAGNETIC CIRCULAR DICHROISM (MCD)

1. INTRODUCTION

In this work, which is part of a long-term MCD research effort in this Laboratory,¹ the focus is on an electronic MCD study² of HI. MCD is a spectroscopic method that probes the earliest, electronic events in photophysics/photochemistry. MCD spectroscopy, the difference of absorptivity of right and left circularly polarized light, refers to a situation in which a static magnetic field is applied parallel to the direction of the polarized beam. MCD spectroscopy in the region $2000 > \lambda > 1400 \text{ \AA}$ of small polyatomic molecules is the focus of our investigations in this laboratory. In Circular Dichroism (CD) spectroscopy one measures the difference in absorption of left and right circularly polarized light caused by the torques exerted on the molecular constituents by the electromagnetic field. Consequently, the measurements are dependent on the molecular structure as well as on the changes that occur during interactions with the field and/or with the molecular environment. In MCD measurements, on the other hand, the two forces, circularly polarized light and (static) magnetic field, are coupled.

Circular dichroism spectroscopy has been applied to a great variety of structures ranging from small chiral molecules to larger structures such as eukaryotic cells in solutions. MCD spectra are not so readily available. Indeed, MCD spectra are very scarce even for small molecules and are virtually non-existent for the Vacuum Ultraviolet (VUV) region. This, despite the fact that all molecular systems must exhibit electronic MCD spectra. Because of its general presence, MCD has a considerable potential and could become the spectroscopy of choice for the study of molecular DNA damage, in particular local DNA molecular lesions.

In order for such to come to pass, some of the experimental and interpretive difficulties now facing MCD techniques and analysis must be removed. Several methods have been proposed for the interpretation of electronic MCD and the birefringence of small molecular systems. Most earlier models related directly to Zeeman effects (line shifts) in atoms. A review of work prior to 1966 has been given by Buckingham and Stephens.³ A more recent discussion has been given by Michl and Thulstrup,⁴ and is oriented toward the interpretation of larger molecule MCD. This latter approach uses Mueller-Stokes matrices. A similar but more general interpretation of the polarized light parameters is also available within the framework of statistical optics⁵ using polarization and coherency matrices. Nonlinear magnetic field effects⁶ obtained using more coherent sources of radiation have also been the subject of much recent investigations and interpretative efforts.

Initial attempts to interpret MCD spectra were all concerned with systems of high symmetry in solutions, systems in which the linewidths were large relative to atomic Zeeman splits. Consequently, only electronic terms were considered, the interpretation of vibrational and rotational spectral structures being either impossible or irrelevant. The earliest modellings of magnetic effects on small molecules include the work of Hause on the IR spectrum of the NO molecule⁷ and of Eberhardt on the electronic spectrum of the ICl molecule.⁸

2. EXPERIMENTAL

Experimental MCD work in this Laboratory has focussed on the VUV region and on particular groups of molecules and atoms: simple gaseous molecular systems⁹ such as CH₃I which exhibit both Rydberg and valence electronic spectral structures.

The VUV absorption and MCD spectra of the gases HI molecule in the region 250 to 146nm (region of the first Rydberg transitions and beyond) have been studied at various pressures below 1 Torr (Fig's. 1-4). This molecule is of particular interest since it is a simple diatomic that exhibits resolved rotational and vibrational electronic structures above the CaF₂ cutoff of 140nm. Progress in MCD measurements beyond this limit requires optics that are not presently available. An expansion of MCD spectroscopy below this limit was one of the subjects discussed at a recent Workshop¹⁰ on MCD. Some possibilities related to the use of synchrotron radiation in MCD studies beyond this limit were addressed at that Workshop.

Our present research efforts deal with the systematic study of the electronic spectra of HI. The analysis of the VUV absorption and MCD spectra of HI reveals some new results concerning the electronic structure of HI. First, quantitative information related to the large spin-orbit parameters ($>5000\text{ cm}^{-1}$) of the HI molecule are now available. Indeed, the observed MCD spectra encompass the two most intense absorption bands which directly reflect spin-orbit splitting. However, a small shift of $\sim 20\text{ cm}^{-1}$ is observed at the center of the 62320 cm^{-1} line. The precise interpretation of the relative positions of line centers will demand a quantitative interpretation of high resolution rotational and vibrational electronic spectra. Secondly, unexpectedly strong MCD signals occur for all absorption bands (first Rydberg bands from ~ 54600 to 68600 cm^{-1}), some of which appear to be in complete discord with some existing state assignments as s, d or p. Thirdly, many of the observed MCD structures do not meet either of the two specifications for A (derivative) and B (absorption) signal types, signatures which frequently are used for the analysis of low-resolution MCD spectra. Thus, a model which incorporates explicit rotational and vibrational structures is much in need.

3. RESULTS

Magnetic field strength effects are observed in the MCD spectrum in the region around 56740 cm^{-1} : the intensity of the MCD lines is linear with field, but no peak to peak shifts are observed in the A structures (Fig. 5). An analysis of the observed spectra as a combination of A structures, an approach similar to that used in earlier studies of opto-magnetic activity, indicates that an interpretation can be found in terms of a field induced "mixing" of electronic states. By fitting the observed MCD spectra with two derivative functions extracted from the MCD spectra in the energy region around 56740 cm^{-1} , the separation (or splitting) between the pairs of "doublets" for the first four MCD bands (between 55830 - 62330 cm^{-1}) is estimated to be in the range 20 to 90 cm^{-1} . However, we do not know the nature of the states that are admixed by the field.

The magnetic momentum ratio obtained from the MCD spectra for the first (around 55830 cm^{-1}) and second (around 56790 cm^{-1}) bands is: $m_1/m_2 \sim 2.25$, and the ratio between the first and fourth (around 67330 cm^{-1}) bands is $m_1/m_4 \sim 2$. The magnetic moment obtained from the MCD spectrum in the region of the third band (around 60840 cm^{-1}) is difficult to estimate. However, the results indicate that, even with present experimental MCD limitations, the technique is capable of generating certain fundamental insights into the electronic processes in molecular systems.

The MCD results also call attention to the coupling mechanisms of electronic and nuclear motions in the rovibronic levels of Rydberg states. Thus we have begun an analysis based on a previously - developed spin-uncoupling model. This model was recently applied to a study of the VUV spectra of CH_3I .¹¹ It was shown that uncoupling of the spin angular momentum of the optical electron from the molecular axis was essentially

complete in some s- Rydberg states. We believe that the same model, can account for the main MCD characteristics found in the HI spectra.

We have also initiated a computational analysis of the observed spectra using methods such as Fourier transform and convolution and deconvolution procedures. The results of the calculations indicate that the observed roelectronic envelope of the first electronic transition can be correlated directly to the observed MCD spectra. If successful, this study will provide very specific information about low-n Rydberg electrons in magnetic fields.

References

1. J. D. Scott, W. S. Felps and S. P. McGlynn, Nuc. Instrum. and Meth. 152, 231 (1978).
2. K. Rupnik, W. S. Felps, and S. P. McGlynn, OSA Annual Meeting Technical Digest 1990, Vol. 15, Optical Society of America, Washington, D. C., 1990,p.144.
3. A. D. Buckingham and P. J. Stephens, Ann. Rev. Phys. Chem. 17, 399 (1966).
4. J. Michl and E. W. Thulstrup, "Spectroscopy with Polarized Light", VCH, New York, 1986.
5. J. W. Goodman, "Statistical Optics", Wiley, New York, 1985.
6. Y. R. Shen, "The Principles of Nonlinear Optics", Wiley, New York, 1984.
7. J. L. Aubel, and C. D. Hause, J. Chem. Phys., 44, 2659, (1966).
8. A. F. Stalder, and W. H. Eberhardt, J. Chem. Phys. 47, 1445 (1967).
9. S. P. McGlynn, J. D. Scott and W. S. Felps, J de Phys. (Paris) 43,C2, 305 (1982).
10. Workshop on Magnetic Circular Dichroism, CAMD / LSU, December 17, 1990, (unpublished)

11. J. A. Dagata, M. A. Scott, W. S. Felps and S. P. McGlynn, in
UNDERSTANDING MOLECULAR PROPERTIES, J. Avery,
A. E. Hansen, and J. P. Dahl, eds. (D. Reidel, Dordrecht, 1987) p. 277.

Figure Captions

- Fig.1. VUV (high and low resolution) absorption spectra (upper) derivative of absorption (middle) and MCD spectra (lower) of HI in the region of the first 5p-6s Rydberg band at $\sim 55830 \text{ cm}^{-1}$.
- Fig.2. Absorption at various pressures (upper), derivative of absorption (middle) and MCD spectra (lower) of HI in the region of the second 5p-6s Rydberg band at $\sim 56790 \text{ cm}^{-1}$.
- Fig.3. High-resolution absorption (upper) and MCD spectra (lower) of HI molecules in the region of the third 5p-6s Rydberg band at $\sim 60890 \text{ cm}^{-1}$.
- Fig.4. Absorption (upper) and MCD spectra (lower) of HI molecules in the region of the fourth 5p-6s Rydberg band at $\sim 62330 \text{ cm}^{-1}$.
- Fig.5. MCD spectra of HI molecules in the region of the second 5p-6s Rydberg band: magnetic field strengths are: 1, 2 and 4 T.

55.6 55.8 56.0
 $\nu \longrightarrow$ (10^3 cm^{-1})

Fig. 1.

Fig. 2.

Fig. 3.

Fig. 4.

Fig. 5.

E. PHOTOCHEMISTRY OF HALOGENATED MOLECULES

An examination of the VUV absorption spectra of halogenated molecules shows that, in general, only the spectra of the normal-alkyl iodides^{1,2} and bromides^{3,4} exhibit "regular" structure at shorter wavelength. These absorption features are readily fitted by means of the Rydberg equation to yield ionization energies that agree well with those obtained from Photoelectron Spectroscopy (PES). An example of such regular structure exhibited in the spectrum of CH_3I is shown in Figure 1. The classes of halogenated molecules that do not exhibit this regular Rydberg structure are extensive: alkyl chlorides;^{5,6} other singly or multiply substituted alkyl iodides,² bromides³ and chlorides;⁶ any alkyl halide which also contains one or more different halogen atoms;^{2,7} vinyl iodide;² iodoacetylene;⁸ and the cyanogen halides.⁹ The absence of regular structure, as illustrated by ICN, is shown in Figure 1.

We reasoned in the case of the cyanogen halides⁹ that the presence of intravalence transitions could account for the absence of "regular" Rydberg structure. A meaningful extension of such an analysis to halogenated molecules larger than a triatomic was found to be difficult because of the increased complexity of the molecular orbital structure.

Cyanogen iodide was found to be extremely photosensitive: if the ICN sample was not flowed rapidly, molecular iodine peaks appeared strongly both in the absorption and magnetic circular dichroism (MCD) spectrum of ICN. Evidence for the presence of a cyanogen photoproduct was not found, probably because of the inherent weakness and broadness of its absorption features.¹⁰ BrCN and ClCN are evidently much less photosensitive: absorption by their photoproducts (Br_2 ; Cl_2 ; $(\text{CN})_2$), was not detected even though their extinction coefficients are similar to those of the parent molecules. A similar situation regarding the non-observance of photoproducts exist for the alkyl iodides,

bromides and chlorides, compounds we have investigated using both absorbance and MCD.

We are presently investigating the possibility that the stainless steel wall of the absorption cell may serve as a trap for halogen photoproducts and prevent their observation. All our previous absorption studies have utilized either 33mm or 100mm pathlength stainless steel cells with either MgF_2 or LiF_2 windows. We now plan to use either a home made 100mm Teflon cell or a variable short path length stainless steel cell which was purchased from McPherson, the manufacturer of the VUV monochromator. The pathlength of the cell is determined by the thickness of the spacer that is used between two cell windows and has a lower limit of 0.0015 inch or $\sim 0.04\text{mm}$. Use of the teflon cell should limit and change the nature of any wall reactions. The short path length cell will allow investigations at sample densities that are ~ 2600 times greater than previously studied and, thus, also reduce the importance of wall reactions of the photoproducts.

Another aim of this research is an understanding of the effects of added perturber gases on photochemical processes. Oxygen or NO is reported¹¹ to increase the quantum yield of photodissociation of methyl iodide at 2537Å from 0.003 for pure CH_3I to unity when either is present in excess relative to CH_3I . We have found, however, that the photo-dissociation of CH_3I remains negligible when contaminated with air at sample ratios of 1 part CH_3I to 1000 parts air for VUV radiation from 2000-1150Å. Our finding might be interpreted in terms of wall recombinations. Consequently, our planned experiments with the teflon cell and the short path length cell will address this question as well.

REFERENCES:

1. R. A. Boschi and D. R. Salahub, Mol. Phys. 24, 289 (1972).
2. R. A. Boschi and D. R. Salahub, Mol. Phys. 24, 753 (1972).
3. G. C. Crosley and B. R. Russell, J. Chem. Phys. 62, 848 (1975).
4. M. A. Baig, J. P. Connerade and J. Hormes, J. Phys. B 15, L5 (1982).
5. D. T. Truck, D. R. Soloman, and D. A. Armstrong, J. Mol. Spectrosc. 78, 31 (1979).
6. J. W. Raymonda, L. O. Edwards, and B. R. Russell, J. Amer. Chem. Soc. 96, 1708 (1974).
7. A. Penner and A. Amirav, J. Chem. Phys. 93, 8576 (1990).
8. D. R. Salahub and R. A. Boschi, Chem. Phys. Lett. 16, 320 (1972).
9. W. S. Felps, K. Rupnik and S. P. McGlynn, J. Phys. Chem. 95, 639 (1991).
10. R. E. Conners, J. L. Roebber and K. Weiss, J. Chem. Phys. 60, 5011 (1974).
11. J. R. Maier and J. P. Simons, Adv. Photochem. 2, 137 (1964).

Figure 1 Electronic absorption spectra of CH₃I and I-CN in the spectral region of the lowest energy ionization potential, 2e and 2 π , respectively. The dashed line in each spectrum is the base line.

F. DENSITY EFFECTS ON HIGH-N RYDBERGS

The spectroscopic study of perturber number density effects, both atomic and molecular, has emerged as a very efficient approach to study the mechanisms of momentum and energy transfer and deposition in density dependent, highly-energetic electronic processes. These mechanisms are considered to be important precursors of the more complicated intermolecular electronic processes in photochemistry /photophysics, as well as in the radiation chemistry and physics of larger biological entities in the in vivo environment. The focus of the work was on the electron scattering and the solvatochromic shifts of molecular Rydbergs, and it suggests that spectroscopic shifts of the Rydberg states may be used to generate scattering cross sections for nonpolar perturbers.

Highly excited Rydberg states are extremely sensitive to the surrounding medium because of their "large orbital" nature. Following the experiments of Amaldi and Segré¹, in which gaseous perturbers modulated the excited states of Na and K atoms, Fermi presented a simple model² of the observed energy shifts. According to Fermi, the interaction between a highly excited absorber and a perturbing medium can be decomposed into two independent, additive terms. These are the scattering term $\Delta_{sc}(\rho)$, (Eq.1.) describing the interaction between the optical electron and perturber, and a polarization term $\Delta_p(\rho)$, (Eq. 2) describing the interaction between the cationic core and perturber. The sum of these two terms (Eq. 3) is the total energy shift of the high-n Rydberg state $\Delta_t(\rho)$:

$$\Delta_{sc}(\rho) = (2\pi \hbar^2 / m) A \rho \quad (1)$$

$$\Delta_p(\rho) = -10\alpha e^2 \rho^{4/3} \quad (2)$$

$$\Delta_t(\rho) = \Delta_{sc}(\rho) + \Delta_p(\rho) \quad (3)$$

where m and e are the electronic mass and charge of the electron; A is the electron scattering length of the atom or molecule; ρ is the number density of the perturber gas; and α is the macroscopic polarizability of the perturber medium. According to experiment, the density dependence of the total energy shift is always linear³; therefore, at large perturber densities, the $\rho^{4/3}$ dependence (Eq.2) fails to describe the experimental observation. It was found³ that a modification of the polarization term proposed by Alekseev and Sobel'man⁴ within the impact approximation to line-broadening theory could be applied to the experimental results with success:

$$\Delta_p(\rho) = -10.78(\alpha e^2/2)^{2/3} (\pi v)^{1/3} \rho \quad (4)$$

where v is the relative thermal velocity of the colliding absorber-perturber pairs. Recent experiments on the CH_3I /nitrogen⁵ system provided a very sensitive test of the Fermi theory in general and the polarization term in particular. The observation of a zero shift over the entire experimental density range supports the evidence about the role of polarizability, here given in Eq.(4). According to equation (3), the small positive electron scattering length⁵ of nitrogen produces a "blue" energy shift which is cancelled by the "red" polarization energy shift. This mutual cancellation occurs as a result of both the linear density dependence of the two shift terms and the equality of the two shift rates. Indeed, this zero shift can be reproduced by calculation using $-\Delta_p(\rho) = \Delta_{sc}(\rho)$, an equality which yields $A = 1.9 \times 10^{-9} \text{cm}$.

We have compared three sets of scattering length data: one from measurements of density effects on high- n Rydberg states of CH_3I , a second based on TOF (time of flight) data, and a third based on the drift of electron swarms. The latter two data sets yield A only after application of MERT extrapolations to zero energy. O'Malley derived⁶ the MERT (modified effective range theory) expressions for the phase shifts of the partial waves of a scattering system

consisting of an electron and neutral polarizable atom or molecule. The expressions can be used to extrapolate data to low and experimentally inaccessible energies. The primary adjustable parameter in the MERT expression is the electron scattering length. A comparison of scattering lengths from pressure shift data for the Rydberg states $n=10-16$ (0.13-0.05eV) with those from the MERT expansions yields generally good agreement.

It is now clear that all models for electron scattering length, A , of atomic and molecular systems contain the polarizability α . Consequently, ~~we~~ have investigated the role of atomic and molecular polarizabilities in such number density effects. For example, following the idea of atomic effective-range theory, which applies to neutral electron scatterers, we find that a simple linear correlation brings five rare gas atoms into a common frame. This correlation is purely empirical. It was found⁷ that a linear correlation exists between the electron scattering length, A , of the atomic perturbers, as measured by pressure shift method, and the polarizabilities for He, Ne, Ar, Kr, and Xe gases. The correlation obtained with the data from swarm experiments confirms that finding. This correlation, however, is considerably different in intent and complexity from the modified effective-range models developed for rare gases. By using one single parameter, the polarizability, it provides scattering lengths at the low-energy limit. The nature of the scattering potential, (whether repulsive or attractive) and the sign (blue or red shift) are also contained in the correlative behavior, since it predicts both size and sign of electron scattering length.

In our recent work the accent was on the role of molecular polarizability. There is now agreement that the empirical relation between polarizability and the scattering length for the rare gases is linear. Questions remain, however, about the molecular polarizability. We have observed that the anisotropic, so called non-spherical, polarizability of some molecules (H_2 and N_2 , for example) correlates very well with their electronic scattering lengths. This empirical

finding leads to questions about the possible role which the two types of response functions (atomic and molecular polarizabilities) play in the elucidation of various mechanisms in density dependent electronic processes.

On Fig. 1., we show a correlation between electronic scattering lengths and average atomic and molecular polarizabilities for both atoms and molecules. The data are from this study. Two separate correlations exist: (1) correlation between α , atomic polarizability for He, Ne, Ar, Kr, Xe and (2), a correlation between molecular (H_2 , N_2 , C_2H_6 , C_3H_8) polarizabilities and corresponding electronic scattering lengths. The two lines have essentially identical intercepts of 1.23, and slopes that are in the approximate ratio 2:1.

This result indicates that the "more anisotropic" molecular systems "see" electrons differently than "more isotropic" systems. It is also interesting that CH_4 , which is as "more isotropic" molecule, lies close to the atomic line whereas the CO_2 molecule, with a well known quadrupolar characteristic, does not align with the other molecules.

The correlation of electron scattering lengths and polarizabilities indicates that Rydberg electron scattering by atomic and molecular perturbers should be described in terms of a spatio-temporal response functions of perturbers. This work is in progress.

EXPERIMENTAL

Perturber density effects on alkali metal absorbers may only be studied at elevated temperatures. Rydberg spectra of molecular absorbers, however, can be studied at room temperature or even lower, being dependent only on the freezing point of the absorber/perturber system. Thus, maintenance of the sample cell at room temperature permits studies of density effects well below vibrational thermal-excitation thresholds. Further advantages of molecular absorbers are the relative insensitivity of band widths of the absorber over a large

experimental perturber density range (up to $5\text{-}10 \times 10^{20} \text{ cm}^{-3}$) and an asymptoticity of energy at principal quantum numbers as low as $n=9$ or 10 . A detailed description of the experimental techniques of pressure shift measurements for high- n Rydbergs of CH_3I in different perturber gases is given elsewhere^{3,5}.

Density effects on CH_3I were studied at room temperature in the photoabsorption and photoionization spectra in transparent and non-transparent perturbers, respectively. The shifts of discrete nd ($^2E_{3/2}$) and nd' ($^2E_{1/2}$) transitions were measured^{3,8,9} at a moderate resolution of $5\text{-}6 \text{ meV}$. The CH_3I absorber pressure lay in the range $0.1\text{-}1 \text{ mbar}$ whereas perturber pressures, $0.5\text{-}50 \text{ bar}$, were much larger. Photoionization spectra were recorded using a synchrotron light source. In the autoionization range ($E_n > I_v$), the photocurrent spectra appear as inverted phototransmission spectra. It is found, as expected, that the current maxima coincide exactly with the places of transmission minima for the nd' ($^2E_{1/2}$) transitions. The photoionization spectrum of CH_3I in the presence of propane perturber is shown in Fig. 2. The red shift of the nd' series members is readily seen by comparison with a pure methyl iodide reference which was recorded simultaneously.

Photocurrent steps have been observed recently¹⁰ in the photoionization spectra of both pure and perturbed CH_3I below the first ionization potential. These subthreshold photoionization spectra have been interpreted and have been used for the evaluation of density effects.¹¹ These spectra constitute one of the few known instances of vibrational subthreshold autoionization. The photocurrent steps are identified as nd Rydberg transitions originating in excited vibrational levels of the electronic ground state and which, subsequently, autoionize into the vibrational ground state of the CH_3I^+ ion.

COMPARISON OF A WITH TOF AND SWARM SCATTERING LENGTHS

The total energy shift of the high- n Rydberg states becomes asymptotic at $n=9$ or 10. The average asymptotic energy shifts of CH_3I are plotted as a function of perturber number in Fig. 3. The density dependencies are consistently linear. The typical experimental slope error is approximately $0.1\text{-}0.3 \times 10^{-23} \text{ eV cm}^3$. Table 1 presents the total, scattering and polarization shift rates: Δ_t/ρ ; Δ_{sc}/ρ and Δ_p/ρ respectively. These values are in good agreement with early¹² and recent¹³ pressure shift data, the latter of which used "doppler-free" laser techniques for Rydbergs of alkali metal absorbers. The scattering shift is obtained by subtracting the polarization shift from the total shift. The macroscopic polarizabilities and relative thermal velocities required for calculating the polarization shift (Eq.4) are also shown in Table 1.

The density dependence of the scattering shift Δ_{sc} is specially important to considerations of the evolution of band structure in insulators. The $\Delta_{sc}(\rho) \approx V_0(\rho)$ equivalence, where V_0 is the electron energy at the bottom of the conduction band, is the crucial connector. Evidence from photoelectric injection experiments in both rare and molecular gases^{14,15} suggests that "band-like" behavior occurs well below the critical density in these systems.

The upper density limit of the pressure shift experiment reported here depends on the mass of the scatterer and may extend as high as $5 - 10 \times 10^{20} \text{ cm}^{-3}$. In this density range, the accuracy of the photoelectric method^{16,17} is low (30 - 50 meV) because of photocathode surface effects, whereas the accuracy of Δ_{sc} from pressure shifts is an order of magnitude higher.

The scattering length governs the size and sign of the potential barrier ($\Delta_{sc} \approx V_0$) experienced by the optical electron. Electrons of large Rydberg orbitals are quasifree. Close to zero energy, the scattering of these electrons is governed by the same cross section that is "active" in momentum transfer scattering (σ_m) or in TOF experiments (σ_T), and the three types of cross sections may not be

distinguished¹⁸. For electrons at the zero energy limit, the cross sections are determined by the scattering length¹⁸:

$$\sigma_{T,m}(k \rightarrow 0) = 4\pi A^2 \quad (5)$$

where k is the wave number of the electron, $k^2 = (2m/\hbar^2)E$. The dependence of the total or momentum transfer cross section on electron energy is determined from electron attenuation and drift measurements, respectively. The cross section data are then extrapolated to zero energy using a four-parameter MERT expansion⁶, the scattering length being the primary fitting parameter. Scattering lengths, as obtained from high- n CH_3I Rydbergs and from electron scattering cross sections of both TOF and swarm origins are shown in Table 2. The TOF (A_{tof}) and swarm (A_{sw}) scattering lengths compare well with those (A) measured by the pressure shift method. The typical error of A_{tof} and A_{sw} values is 1-2%. Consequently, we find that the scattering length obtained from density effects constitutes a "zero energy" scattering length despite the finite electron energy of $n=9-16$ asymptotic Rydberg transitions (0.05-0.13eV on average, respectively). O'Malley⁶ had earlier proposed a simplified extrapolation for the zero energy scattering length (A_0). For true "zero energy" one should only consider Rydberg states $n \rightarrow \infty$, in which case $k = (na_0)^{-1} \rightarrow 0$. Assuming that $\tan \eta_0/k$ is proportional to the energy shift, η_0 being the phase shift, the "extrapolated" value of A_0 may be obtained⁶ as

$$A_0 = A - (\pi/3a_0)\alpha k \quad (6)$$

or, more simply, as

$$\pm A_0 = \pm A - \alpha\pi/3n \quad (7)$$

Our attempts to extrapolate to near-zero energy by subtraction of the linear k term from the experimental values of A did not give satisfactory results: The A_0 values exhibited a large deviation from A_{tof} and A_{sw} "zero energy" values. For example, by substituting the polarizabilities 16.73, 27.29 and $17.54a_0^3$ and using $n=13$ (0.08eV), we find $A_0 = -4.39a_0, -8.16a_0, -3.93a_0$ for Kr, Xe and methane,

respectively. These "zero energy" values differ by 50% from the scattering lengths A obtained at $n=9-16$ as well as the MERT values (A_{sw} , A_{tof}) of Table 2. More than one value of A has been obtained from MERT analyses of swarm and TOF data for the rare gases (see Table 2). Almost all pressure shift scattering lengths lie within 10% of the swarm or TOF scattering lengths. Indeed, the swarm and TOF A values may differ more in themselves than in their deviance from the pressure shift values. A comparison of results follows:

- - - A is lower by ~10% than either $A_{tof}^{19,20}$ or A_{sw}^{21} for helium; and A for neon is also lower by ~10% than the A_{sw} value cited by Crompton and O'Malley²². The scattering length for argon obtained by pressure shift studies is very close to the average of $A_{tof}^{20,23,24}$ and A_{sw}^{25} .

- - - The scattering length of Kr may be compared with the results of Weyhreter et al.²³ and Buckman and Lohmann²⁶; A_{tof} obtained by the latter authors agrees to within 10% with our result^{3,27}. Buckman and Lohmann²⁶ also analyzed the momentum transfer cross sections for Kr given by Frost and Phelps²⁸. For Xe²⁹, comparison is again made with A_{tof} from Weyhreter et al.²³ and an earlier analysis by O'Malley⁶ of the σ_m results published by Pack, Phelps and coworkers.³⁰

- - - Only isotropic long-range potentials are considered in the theory of electron scattering by rare gas atoms.^{6,25} The theoretical description of the extrapolation for simple molecules, however, necessitates also a concern for anisotropic potentials. Chang³¹ presented such a theory for molecules and found that, for hydrogen, true molecular effects contribute only about 1% to the total cross section. His theory agrees well with TOF cross sections for hydrogen. The pressure shift value of A for hydrogen ($0.93 a_0$), however, is lower by 30% than the scattering lengths derived by Chang from the momentum transfer cross sections of Crompton et al.³² and the TOF data of Ferch et al.³³, both of which cite $1.27 a_0$.

- - Chang also applied his theory³¹ to the swarm data of Engelhardt et al. for nitrogen³⁴, and obtained $A_{sw} = 0.44a_0$. Because of the large contribution of anisotropic molecular potentials, $4\pi A^2$ does not yield a zero energy cross section (see Eq.5). Indeed, to obtain a correct $\sigma_T(0)$, a factor³¹ of ≈ 1.58 must be inserted into Eq. 5. This factor³¹ originates in an especially high ratio of Q/A for nitrogen, where Q is the quadrupole momentum. The scattering length of nitrogen as obtained from pressure shifts is then in good agreement with Chang's value. Thus, the case of the nitrogen perturber demonstrates that the parameter obtained from pressure shifts is always the scattering length and not the square root of the cross section (Eq.5.), two values which are equivalent in the Fermi model, for atomic scatterers only.

- - Methane is used widely in conjunction with inert gases in track detectors, and its transport coefficients have special importance. The application of MERT to such a highly symmetrical molecule as methane has not been treated theoretically. The scattering length of methane, as obtained from pressure shifts ($-2.61a_0$), is intermediate to two recent A values. Ferch et al.³⁵ employed a four parameter MERT^{6,25} fit for the total scattering cross section data getting $A = -2.475a_0$. In a refined analysis based on drift velocity and the longitudinal and transverse diffusion coefficients of electrons, Schmidt³⁶ obtained $A = -2.755a_0$ using a six parameter MERT.

- - Gee and Freeman³⁷ evaluated momentum transfer cross sections for methane and ethane at very low electron energies. For methane, for which the cross section was followed down to 0.001eV , the values of σ_m were found to be far higher than any existing data might suggest. No MERT analysis of these σ_m data has been made. However, the values of σ_m at $\sim 0.01\text{eV}$ compare well with the cross sections for both methane and ethane obtained from pressure shift studies. For ethane and propane, the TOF cross sections do not extend to low enough electron energies³⁸ to merit comparison with the shift data.

--CO₂ has important applications in laser and plasma physics. Accurate computed scattering cross sections have become available³⁹ for CO₂ at very low energies. These computed values exhibit a rapid increase with decreasing electron energy, possibly due to a nearly bound $l=0$ state in the e-CO₂ potential; the computations also agree well with the TOF measurements of Ferch et al.⁴⁰ A zero energy extrapolation, however, is not available. Nonetheless, when Eq.5 is arbitrarily applied at 0.07 eV (which is the lowest energy for which measurements are available⁴⁰), one finds $A=-4.3a_0$, in good agreement with the pressure shift value of $-4.23 a_0$ ⁴¹. The attractive character of the scattering potential, which is clear from the sign of the pressure shift, inhibits occurrence of a normal Ramsauer minimum in the σ_T vs electron energy plot⁴⁰, and a very broad minimum extending from 1.0 eV to a prominent resonance at 3.8eV is observed instead.

REFERENCES

1. E. Amaldi and E. Segré, *Nuovo Cimento* 11, 145, 1934.
2. E. Fermi, *Nuovo Cimento* 11, 157, 1934.
3. A. M. Köhler, R. Reininger, V. Saile and G. L. Findley, *Phys. Rev. A* 35, 79 (1987); R. Reininger, E. Morikawa and V. Saile, *Chem. Phys. Lett.* 159, 276, 1989.
4. V. A. Alekseev and I. I. Sobel'man, *Sov. Phys. JETP* 22, 882 (1966).
5. U. Asaf, J. Meyer, R. Reininger and I. T. Steinberger, *J. Chem. Phys.*, in press.
6. T. F. O'Malley, *Phys. Rev.* 130, 1020, 1963.
7. K. Rupnik, U. Asaf and S. P. McGlynn, *J. Chem. Phys.* 92, 2303, 1990.
8. U. Asaf, W. S. Felps, K. Rupnik, S. P. McGlynn and G. Ascarelli, *J. Chem. Phys.* 91, 5170 1989; U. Asaf, W. S. Felps and S. P. McGlynn, *Phys. Rev. A* 40, 5458, 1989.
9. J. Meyer, U. Asaf and I. T. Steinberger, *J. Chem. Phys.* 94, 1820, (1991).
10. J. Meyer, U. Asaf and R. Reininger, submitted to *Phys. Rev. A*.
11. I. T. Steinberger, U. Asaf, G. Ascarelli, G. Reisfeld and M. Reshotko, *Phys. Rev. A* 42, 3135, (1990).
12. S. Y. Ch'en and M. Takeo, *Rev. Mod. Phys.* 29, 20, 1957;
D. K. L. Tan and S. Y. Ch'en, *Phys. Rev. A* 2, 1124, 1970.
13. K. H. Weber and K. Niemax, *Z. Phys. A* 307, 13, 1982.
14. R. Reininger, U. Asaf and I. T. Steinberger, *Chem. Phys. Lett.* 90, 287, 1982.

15. U. Asaf, I. T. Steinberger and R. Reininger, Chem. Phys. Lett. 100, 363 (1983).
16. R. Reininger, U. Asaf and I. T. Steinberger, Phys. Rev. B 28, 4426, 1983.
17. U. Asaf and I. T. Steinberger, Chem. Phys. Lett. 128, 91, 1986.
18. H. S. W. Massey, E. H. S. Burhop and H. B. Gilbody, Electronic and Ionic Impact Phenomena, Vol. 1, Oxford, 1969.
19. J. Ferch, W. Raith and K. Schröder, J. Phys. B: At. Mol. Phys. 13, 1481, 1980.
20. S. J. Buckman and B. Lohmann, J. Phys. B: At. Mol. Phys. 19, 2547, 1986.
21. R. W. Crompton, Elford and Robertson, Aust. J. Phys. 23, 667, 1970.
22. T. F. O'Malley and R. W. Crompton, J. Phys. B: At. Mol. Phys. 13, 3451, 1980.
23. M. Weyhreter, B. Barczik, A. Mann and F. Linder, Z Phys. D. 7, 333 (1988).
24. J. Ferch, B. Granitza, B. Masche and W. Raith, J., Phys. B: At. Mol. Phys.
25. G. N. Haddad and T. F. O'Malley, Aust. J. Phys. 35, 35, 1982.
26. S. J. Buckman and B. Lohmann, J. Phys. B: At. Mol. Phys. 20, 5807, 1987.
27. U. Asaf, M. Sielberg, J. Meyer, R. Reininger, G. Reisfeld and I. T. Steinberger, submitted to Chem. Phys.
28. L. S. Frost and A. V. Phelps, Phys. Rev. A 1538, 1964.
29. I. T. Steinberger, U. Asaf, G. Ascarelli, G. Reisfeld and M. Reshotko, Phys. Rev. A42, 3135, 1990.
30. J. L. Pack and A. V. Phelps, Phys. Rev. 121, 798, 1961; J. L. Pack, R. E. Voshall and A. V. Phelps, Phys. Rev. 127, 2084, 1962.

31. E. S. Chang, J. Phys. B: At. Mol. Phys, 14, 893, 1981.
32. R. W. Crompton, D. K. Gibson and A. I. McIntosh, Aust. J. Phys. 22, 715, 1969.
33. J. Ferch, W. Raith and K. Schröder, J. Phys. B: At. Mol. Phys., 13, 1481, 1980.
34. A. G. Engelhardt, A. V. Phelps and C. G. Risk, Phys. Rev., 35A, 1566, 1964.
35. J. Ferch, C. Masche and W. Raith, J. Phys. B: At. Mol. Phys. 14, L97, 1981.
36. B. Schmidt, J. Phys. B: At. Mol. Phys. 24, 4809, 1991.
37. N. Gee and G. R. Freeman, Phys. Rev., A20, 1152, 1979;
N. Gee and G. R. Freeman, Phys. Rev., A22, 301, 1980.
38. O. Sueoka and S. Mori, J. Phys. B: At. Mol. Phys. 19, 4035 (1986).
39. M. A. Morrison, N. F. Lane and L. S. Collins, Phys. Rev. A15, 2186, 1977.
40. J. Ferch, B Granitza and W. Raith, J. Phys. B: At. Mol. Phys. 18, L445, 1985.
41. U. Asaf, I. T. Steinberger, J. Meyer and R. Reininger, J. Chem. Phys. 95, 4070, 1991.

TABLE 1.

The averaged total, scattering and polarization shift rates (i.e., the slopes of energy shift vs. perturber number density plots) for the high- n Rydbergs of CH_3I ($n=10-16$) are cited in units of $10^{-23} \text{ eV cm}^{-3}$. Average polarizabilities (Landolt-Börnstein, Zahlenwerte und Funktionen, I Bd., Springer, Berlin, 1951) and the relative thermal velocities (as calculated from the Boltzmann distribution at 300°K) are cited in units of 10^{-24} cm^3 and 10^4 cm/s , respectively. References for A are given in brackets.

gas	Δ_t/ρ	Δ_{sc}/ρ	Δ_p/ρ	α	v	[ref]
He	2.45	2.73	-0.29	0.20	11.4	[3]
Ne	0.09	0.45	-0.35	0.396	6.0	[3]
Ar	-4.86	-4.04	-0.82	1.65	4.5	[3,8,14]
Kr	-8.6	-7.6	-1.01	2.48	3.5	[3,27]
Xe	-16.8	-15.5	-1.29	4.04	3.0	[29]
H ₂	1.62	2.35	-0.73	0.81	17.9	[8]
N ₂	0*	0.88	-0.88	1.76	5.2	[5]
CH ₄	-7.86	-6.63	-1.23	2.60	6.4	[9]
C ₂ H ₆	-10.05	-8.42	-1.63	4.47	5.1	[9]
C ₃ H ₈	-12.93	-10.99	-1.94	6.29	4.4	[9]
CO ₂	-11.8	-10.71	-1.09	2.65	4.3	[41]

* error $\sim 0.15 \cdot 10^{-23} \text{ eV cm}^3$

TABLE 2

Electron scattering lengths (in units of Bohr radius, a_0) from pressure shift (A), from MERT analysis based on cross sections of TOF (A_{tof}) and from swarm data (A_{sw}). References for A are as given Table 1. References for A_{sw} and A_{tof} are given in brackets.

gas	(A)	(A_{tof})	[ref]	(A_{sw})	[ref]
He	1.08	1.95 1.16	[19] [20]	1.19	[21]
Ne	0.17	--	--	0.214	[22]
Ar	-1.63 -1.55	-1.593 -1.492 -1.449	[20] [23] [24]	-1.492	[25]
Kr	-3.02	-3.478 -3.19	[23] [26]	-3.32	[26,28]
Xe	-6.12	-6.527	[23]	-6.0	[6,30]
H ₂	0.93	1.27	[31,32]	1.27	[31,33]
N ₂	0.35	--	--	0.44	[31,34]
CH ₄	-2.61	-2.475	[33]	-2.755	[36]
C ₂ H ₆	-3.33	--	--	--	
C ₃ H ₈	-4.31	--	--	--	
CO ₂	-4.23	--	--	--	

FIGURE CAPTIONS

- FIG. 1: Scattering lengths of rare gases and molecules, denoted: square from Table 1 and 2. are plotted vs. polarizability. The symbol: diamond represents the results of linear correlations (solid lines), for rare gases (slope = -0.265) and for molecules (slope = -0.144).
- FIG. 2: a: Phototransmission of pure CH_3I (0.1 mbar). Rydberg states of the nd' ($^2E_{1/2}$) series are marked at the absorption maxima.
b: Autoionization current spectrum of CH_3I perturbed with dense propane gas for which $\rho = 1.29 \times 10^{20} \text{cm}^{-3}$. The series members are marked at the maxima.
- FIG. 3: Total energy shift averaged over the $n=10-16$ Rydberg transitions of CH_3I for 11 perturber gases. References are given in Table 1.

FIGURE 1

FIGURE 2

FIGURE 3

END

**DATE
FILMED**

10/02/92