Coal Liquefaction Process Streams Characterization and Evaluation

The Use of Selective Chemical Reactions and Carbon NMR Spectroscopy in the Speciation and Quantification of Oxygen Functional Groups in Coal Resids

Topical Report

Prepared by:

University of Chicago
Department of Chemistry
5735 South Ellis Avenue
Chicago, IL 60637

L. M. Stock
C. Cheng

Prepared for:

CONSOL Inc.
Research & Development
4000 Brownsville Road
Library, PA 15129

F. P. Burke
R. A. Winschel
S. D. Brandes

November 1992

Under Contract to:

United States Department of Energy
Contract No. DE-AC22-89PC89883

U.S. DOE Patent Clearance was provided by
Chicago Operations Office on November 5, 1992.

PROJECT ASSESSMENT

Introduction

Under subcontract from CONSOL Inc. (U.S. DOE Contract No. DE-AC22-89PC89883), the University of Chicago speciated and quantified the oxygen functional groups in coal-derived resids. Hydroxyl functional groups in the resids were selectively derivatized by phase-transfer methylation. In order to characterize and quantify the ether groups in the resids, single-electron transfer reactions were used to cleave certain ether groups to produce phenols, which then were derivatized by phase-transfer methylation. The methylated products of reaction were evaluated by ^{13}C -NMR. The full report authored by the University of Chicago researchers is presented here. The following assessment briefly highlights the major findings of the project, and evaluates the potential of the method for application to coal derived materials. These results will be incorporated by CONSOL into a general overview of the application of novel analytical techniques to coal derived materials at the conclusion of this contract.

Summary

This study demonstrated the feasibility of using phase-transfer methylation reactions for the derivatization of hydroxyl functionalities and single-electron transfer reactions to cleave ether groups in distillation resid materials derived from direct coal liquefaction. ^{13}C -NMR was used to analyze the reaction products. It was demonstrated that the techniques can be used to speciate and quantify about half of the oxygen in resids. The remaining half of the oxygen is presumed to be present as unreactive ethers, probably dibenzofurans. Hydroxyl concentrations were determined on five resid samples and ether cleavage reactions were conducted on three of those five. Further development of this general analytical strategy as a process development tool is justified based on these results.

Program Description

This report describes the work performed at the University of Chicago under a subcontract from CONSOL Inc., Research and Development. CONSOL's prime contract to the U.S. Department of Energy (Contract

No. DE-AC22-89PC89883, "Coal Liquefaction Process Streams Characterization and Evaluation") established a program for the analysis of direct coal liquefaction derived materials. The program involves a number of participating organizations whose analytical expertise is being applied to these materials. This Participants Program has two main objectives. The broad objective is to improve our understanding of fundamental coal liquefaction chemistry to facilitate process improvement and new process development. The specific approach to achieving this objective is to provide a bridge between direct coal liquefaction process development and analytical chemistry by demonstrating the application of various advanced analytical methods to coal liquefaction materials. The methodologies (or techniques) of interest are those which are novel in their application for the support of coal liquefaction and those which have not been fully demonstrated in this application. CONSOL is providing well-documented samples from different direct coal liquefaction production facilities to the program participants. The participants are required to interpret their analytical data in context to the processing conditions under which the samples were generated. The methodology employed is then evaluated for its usefulness in analyzing direct coal liquefaction derived materials.

Participant's Methodology

The University of Chicago speciated and quantified the hydroxyl and ether functional groups in the pyridine-soluble portion of 850°F⁺ distillation resids. This was accomplished by: 1) derivatization of the hydroxyl groups in five pyridine-soluble resids by phase-transfer methylation, and 2) ether cleavage reactions of three of the five resids with Lewis acids and single-electron transfer reagents followed by methylation, and 3) quantitation of the reaction products by ¹³C-NMR spectroscopy. The pyridine-soluble resids are solids at room temperature. The original process samples, from which the pyridine-soluble resids were generated, were produced at the Wilsonville pilot plant. The samples are composites of samples taken over long periods of single runs at Wilsonville. Two major processing parameters were varied among the Wilsonville runs: feed coal and reactor configuration (thermal/catalytic vs catalytic/catalytic). Samples were taken from two locations: between the first- and second-stage reactors, and after the second-stage reactor. These

samples are expected to represent different extents of coal liquefaction. The experimental procedures for phase-transfer methylation, Lewis acid and single-electron transfer ether cleavage reactions, and ^{13}C -NMR spectroscopy are described on pages 4 through 6 of the attached report.

Participant's Major Findings

The following principal observations for the derivatization of the hydroxyl groups in the pyridine-soluble portion of coal liquefaction resids, and the ether cleavage and subsequent methylation of the product phenols, were reported by the University of Chicago. An expanded discussion can be found in the attached report, pages 7 through 14

Derivatization of the hydroxyl groups in the pyridine-soluble resids was accomplished by reaction with a phase-transfer base (tetrabutylammonium hydroxide) and subsequent methylation of the product oxyanions with $^{13}\text{CH}_3\text{I}$. The number and kind of hydroxyl groups present in the samples were determined by ^{13}C -NMR. Three types of hydroxyl groups were found to be present: hindered phenols, unhindered phenols, and carboxylic acids. It was concluded that the carboxylic acids are not native to the samples, but arise in the preparation and handling of the pyridine-soluble resids and by reaction with atmospheric CO_2 during the derivatization procedure. The ratio of hindered to unhindered phenols was about 1:2. The quantitative analysis of total hydroxyl concentration agreed reasonably well with phenolic -OH concentration values obtained by the Fourier transform infrared (FTIR) method used by CONSOL.

Ether cleavage was accomplished in three ways: 1) with boron tribromide (BBr_3), a Lewis acid, 2) potassium-naphthalene in tetrahydrofuran, and 3) with lithium in dioxane (the last two are different single-electron transfer reagents). The BBr_3 reaction proved unsuccessful; this Lewis acid apparently catalyzes degradative condensation reactions in the resid as well as the desired ether cleavage reactions. The apparent condensation reactions resulted in a reduction in the number of derivatizable hydroxyl groups in the Lewis acid treated sample (relative to the initial pyridine-soluble resid sample). The first of the two single-electron transfer reagents (potassium-naphthalene) increased the number of derivatizable hydroxyl groups in two of three pyridine-soluble resids

reacted, indicating that some ethers were cleaved. Use of the second single-electron transfer reagent (lithium in dioxane) was unsuccessful in increasing the number of derivatizable hydroxyl groups; it is presumed, therefore, that no ethers were cleaved.

It was proposed by the University of Chicago on the basis of their findings and the assumption that the quantity of total oxygen in the resids (which is determined by difference) is correct, that approximately 40% of the total oxygen (determined by difference) in the pyridine-soluble resids is phenolic. Another 0 to 10% was found to be cleavable ethers. The remaining 50% of the oxygen, it was speculated, exists in the resids as non-cleavable (via single-electron transfer reaction) ethers, such as dibenzofurans or dinaphthofurans, or possibly, dialkyl ethers.

CONSOL Evaluation

The phase transfer methylation technique was employed successfully to derivatize the available hydroxyl species in the pyridine-soluble resid samples. The total quantity of phenolic -OH determined by this method agrees reasonably well with the value obtained by FTIR, with the exception of two samples in which solubility in chloroform (the NMR solvent) was poor. In addition to measuring the total hydroxyl content, the derivatization/NMR method provides classification of the hydroxyl groups into hindered, nonhindered and carboxylic acid classes. Indications are that the carboxylic acids are an artifact of sample preparation and handling.

Although not optimized for coal-derived resid samples, the use of single-electron transfer reagents for selective cleavage of ether groups in the pyridine-soluble resids was demonstrated. The use of two different reagents, one which was effective and the other which was not, illustrates the need for a more detailed study with this type of reagents, 1) to find the optimum reagent system for use with coal-derived materials for the cleavage and subsequent analysis of the ethers, and 2) to better understand the fundamental chemistry which occurs when coal-derived materials are reacted with these reagents.

Each phase-transfer methylation experiment requires approximately five days per sample (two days for a duplicate reaction, two days for work-up, and one day for NMR data acquisition). Ether cleavage reactions require an additional five days for sample preparation, reaction and work-up. The reagents, which are commercially available, cost approximately \$20.00 for each phase transfer methylation reaction and about \$50.00 for each single-electron transfer reaction followed by phase transfer methylation. The ^{13}C -NMR spectrometer used in this study costs approximately \$350,000, and the ^1H -NMR spectrometer costs approximately \$150,000.

Further Development

The use of potassium-naphthalene as a single-electron transfer reagent to induce ether cleavage met with success in this limited study. Many different alkali metal, electron shuttling agents and solvent combinations could be applied to the study of fossil fuels. The optimum combination must be determined empirically. It is recommended that a study be undertaken to determine the most effective combination for ether cleavage in coal-derived materials.

The Lewis acid (BBr_3) used in this study proved to be ineffective because of its promotion of degradative condensation reactions as well as ether cleavage. It is recommended that additional studies be made with BBr_3 and other Lewis acids to determine if milder reaction conditions could promote the ether cleavage reactions over the condensation reactions.

Because the determination of the relative proportion of oxygen functional groups ultimately lies in relying on the value obtained for the total oxygen in the samples, it is recommended that total oxygen content be determined in a more rigorous manner than by difference.

The original resids used in this work were obtained by pyridine extraction of distillation resids. The ^{13}C -NMR analysis of the reaction products was conducted in chloroform solution. Although the methylated products are expected to be more soluble than the original resids, chloroform may not completely solubilize the entire product. Therefore, it is recommended that pyridine or some other good coal-liquid solvent be employed as the NMR solvent in future work.

Participant's Statement of Work

Oxygen is the the principal heteroatom in most coal liquids. Though it often is inferred to be important in coal liquefaction chemistry, its chemical form in coal liquids is ill-defined. The derivatization of hydroxyl oxygen (phenolic, alcoholic, and carboxylic) with ^{13}C labeled methyl iodide, followed by ^{13}C -NMR results, in the case of petroleum-derived resids, in clearly distinguishable and quantifiable signals in the NMR. Selective cleavage of etheric bonds followed by derivatization with ^{13}C methyl groups and identification of the derivatized species by ^{13}C -NMR has been demonstrated with model compounds. These techniques can be used to quantitatively speciate oxygen; however, they have not been demonstrated with process-derived coal liquids, or investigated for their ability to answer questions pertaining to the chemistry of coal conversion. As such, they fit well within the scope of the participants program.

The derivatization of the hydroxyl oxygen, and cleavage and derivatization of the etheric oxygen, followed by subsequent characterization by ^{13}C -NMR of a sample set of six samples will allow a demonstration of the value of these techniques for the examination of the oxygen compounds in coal liquefaction-derived resids. Samples will be derivatized with ^{13}C labeled reagents and analyzed via ^{13}C -NMR to yield a total hydroxyl oxygen value for each of the six samples. Three of the samples (as indicated on the attached list) will be selectively cleaved at ether linkages utilizing two different methods. In the one method, the ethers will be cleaved by forming anion radicals with an electron transfer reagent. This reagent will cleave heterocyclic and non-heterocyclic ethers. The cleavage products will be marked with labeled reagents. NMR spectra of the starting material, alkylated starting material, and the alkylated product of the ether-cleaved material will allow for quantitation of the oxygen-containing species in the pyridine-soluble resid samples. The second method entails cleavage of the ether groups in the starting pyridine-soluble resid sample with a boron trihalide reagent to be chosen by the research group at the University of Chicago. This reagent will cleave non-heterocyclic ethers but not heterocyclic ethers. The treated sample then will be alkylated with a ^{13}C labeled reagent and analyzed by ^{13}C NMR. Comparison of the NMR spectra of the starting material, the

alkylated starting material and the alkylated cleaved materials produced by both methods will allow for quantitative speciation of the oxygen-containing compounds in the pyridine-soluble resid samples.

Six samples will be supplied by CONSOL to the University of Chicago. These samples have been selected so that the utility of the method described above for the measurement of the forms of oxygen can be assessed in application with pyridine-soluble coal liquefaction resids. The samples will be supplied to the University of Chicago with the following information, as available: elemental analyses, phenolic -OH concentration by FTIR, calorific value, hydrogen classes by $^1\text{H-NMR}$, and the full history of the sample (plant, process conditions, age, and storage conditions). The six samples are the pyridine-soluble portion of the 850°F distillation residual materials. Sample size will be at least 10 g. The six pyridine-soluble resid samples will be brittle pitch-like materials that will be supplied as approximately minus 60 mesh powder.

Final Report

**THE USE OF SELECTIVE CHEMICAL REACTIONS AND CARBON NMR
SPECTROSCOPY IN THE SPECIATION AND QUANTIFICATION OF OXYGEN
FUNCTIONAL GROUPS IN COAL RESIDS**

Submitted to:

**CONSOL, INC.
Research and Development
4000 Brownsville Road
Liberty, PA 15129**

by:

**Leon M. Stock - Principal Investigator
Carlos Cheng - Research Associate**

**University of Chicago
Department of Chemistry
Chicago, Illinois 60637**

August, 1992

Coal Liquefaction Process Streams Characterization and Evaluation

The Use of Selective Chemical Reactions and Carbon NMR Spectroscopy in the Speciation and Quantification of Oxygen Functional Groups in Coal Resids

Topical Report

Prepared by:

University of Chicago
Department of Chemistry
5735 South Ellis Avenue
Chicago, IL 60637

L. M. Stock
C. Cheng

Prepared for:

CONSOL Inc.
Research & Development
4000 Brownsville Road
Library, PA 15129

F. P. Burke
R. A. Winschel
S. D. Brandes

November 1992

Under Contract to:

United States Department of Energy
Contract No. DE-AC22-89PC89883

U.S. DOE Patent Clearance was provided by
Chicago Operations Office on November 5, 1992.

PROJECT ASSESSMENT

Introduction

Under subcontract from CONSOL Inc. (U.S. DOE Contract No. DE-AC22-89PC89883), the University of Chicago speciated and quantified the oxygen functional groups in coal-derived resids. Hydroxyl functional groups in the resids were selectively derivatized by phase-transfer methylation. In order to characterize and quantify the ether groups in the resids, single-electron transfer reactions were used to cleave certain ether groups to produce phenols, which then were derivatized by phase-transfer methylation. The methylated products of reaction were evaluated by ^{13}C -NMR. The full report authored by the University of Chicago researchers is presented here. The following assessment briefly highlights the major findings of the project, and evaluates the potential of the method for application to coal derived materials. These results will be incorporated by CONSOL into a general overview of the application of novel analytical techniques to coal derived materials at the conclusion of this contract.

Summary

This study demonstrated the feasibility of using phase-transfer methylation reactions for the derivatization of hydroxyl functionalities and single-electron transfer reactions to cleave ether groups in distillation resid materials derived from direct coal liquefaction. ^{13}C -NMR was used to analyze the reaction products. It was demonstrated that the techniques can be used to speciate and quantify about half of the oxygen in resids. The remaining half of the oxygen is presumed to be present as unreactive ethers, probably dibenzofurans. Hydroxyl concentrations were determined on five resid samples and ether cleavage reactions were conducted on three of those five. Further development of this general analytical strategy as a process development tool is justified based on these results.

Program Description

This report describes the work performed at the University of Chicago under a subcontract from CONSOL Inc., Research and Development. CONSOL's prime contract to the U.S. Department of Energy (Contract

No. DE-AC22-89PC89883, "Coal Liquefaction Process Streams Characterization and Evaluation") established a program for the analysis of direct coal liquefaction derived materials. The program involves a number of participating organizations whose analytical expertise is being applied to these materials. This Participants Program has two main objectives. The broad objective is to improve our understanding of fundamental coal liquefaction chemistry to facilitate process improvement and new process development. The specific approach to achieving this objective is to provide a bridge between direct coal liquefaction process development and analytical chemistry by demonstrating the application of various advanced analytical methods to coal liquefaction materials. The methodologies (or techniques) of interest are those which are novel in their application for the support of coal liquefaction and those which have not been fully demonstrated in this application. CONSOL is providing well-documented samples from different direct coal liquefaction production facilities to the program participants. The participants are required to interpret their analytical data in context to the processing conditions under which the samples were generated. The methodology employed is then evaluated for its usefulness in analyzing direct coal liquefaction derived materials.

Participant's Methodology

The University of Chicago speciated and quantified the hydroxyl and ether functional groups in the pyridine-soluble portion of 850°F⁺ distillation resids. This was accomplished by: 1) derivatization of the hydroxyl groups in five pyridine-soluble resids by phase-transfer methylation, and 2) ether cleavage reactions of three of the five resids with Lewis acids and single-electron transfer reagents followed by methylation, and 3) quantitation of the reaction products by ¹³C-NMR spectroscopy. The pyridine-soluble resids are solids at room temperature. The original process samples, from which the pyridine-soluble resids were generated, were produced at the Wilsonville pilot plant. The samples are composites of samples taken over long periods of single runs at Wilsonville. Two major processing parameters were varied among the Wilsonville runs: feed coal and reactor configuration (thermal/catalytic vs catalytic/catalytic). Samples were taken from two locations: between the first- and second-stage reactors, and after the second-stage reactor. These

samples are expected to represent different extents of coal liquefaction. The experimental procedures for phase-transfer methylation, Lewis acid and single-electron transfer ether cleavage reactions, and ^{13}C -NMR spectroscopy are described on pages 4 through 6 of the attached report.

Participant's Major Findings

The following principal observations for the derivatization of the hydroxyl groups in the pyridine-soluble portion of coal liquefaction resids, and the ether cleavage and subsequent methylation of the product phenols, were reported by the University of Chicago. An expanded discussion can be found in the attached report, pages 7 through 14

Derivatization of the hydroxyl groups in the pyridine-soluble resids was accomplished by reaction with a phase-transfer base (tetrabutylammonium hydroxide) and subsequent methylation of the product oxyanions with $^{13}\text{CH}_3\text{I}$. The number and kind of hydroxyl groups present in the samples were determined by ^{13}C -NMR. Three types of hydroxyl groups were found to be present: hindered phenols, unhindered phenols, and carboxylic acids. It was concluded that the carboxylic acids are not native to the samples, but arise in the preparation and handling of the pyridine-soluble resids and by reaction with atmospheric CO_2 during the derivatization procedure. The ratio of hindered to unhindered phenols was about 1:2. The quantitative analysis of total hydroxyl concentration agreed reasonably well with phenolic -OH concentration values obtained by the Fourier transform infrared (FTIR) method used by CONSOL.

Ether cleavage was accomplished in three ways: 1) with boron tribromide (BBr_3), a Lewis acid, 2) potassium-naphthalene in tetrahydrofuran, and 3) with lithium in dioxane (the last two are different single-electron transfer reagents). The BBr_3 reaction proved unsuccessful; this Lewis acid apparently catalyzes degradative condensation reactions in the resid as well as the desired ether cleavage reactions. The apparent condensation reactions resulted in a reduction in the number of derivatizable hydroxyl groups in the Lewis acid treated sample (relative to the initial pyridine-soluble resid sample). The first of the two single-electron transfer reagents (potassium-naphthalene) increased the number of derivatizable hydroxyl groups in two of three pyridine-soluble resids

reacted, indicating that some ethers were cleaved. Use of the second single-electron transfer reagent (lithium in dioxane) was unsuccessful in increasing the number of derivatizable hydroxyl groups; it is presumed, therefore, that no ethers were cleaved.

It was proposed by the University of Chicago on the basis of their findings and the assumption that the quantity of total oxygen in the resids (which is determined by difference) is correct, that approximately 40% of the total oxygen (determined by difference) in the pyridine-soluble resids is phenolic. Another 0 to 10% was found to be cleavable ethers. The remaining 50% of the oxygen, it was speculated, exists in the resids as non-cleavable (via single-electron transfer reaction) ethers, such as dibenzofurans or dinaphthofurans, or possibly, dialkyl ethers.

CONSOL Evaluation

The phase transfer methylation technique was employed successfully to derivatize the available hydroxyl species in the pyridine-soluble resid samples. The total quantity of phenolic -OH determined by this method agrees reasonably well with the value obtained by FTIR, with the exception of two samples in which solubility in chloroform (the NMR solvent) was poor. In addition to measuring the total hydroxyl content, the derivatization/NMR method provides classification of the hydroxyl groups into hindered, nonhindered and carboxylic acid classes. Indications are that the carboxylic acids are an artifact of sample preparation and handling.

Although not optimized for coal-derived resid samples, the use of single-electron transfer reagents for selective cleavage of ether groups in the pyridine-soluble resids was demonstrated. The use of two different reagents, one which was effective and the other which was not, illustrates the need for a more detailed study with this type of reagents, 1) to find the optimum reagent system for use with coal-derived materials for the cleavage and subsequent analysis of the ethers, and 2) to better understand the fundamental chemistry which occurs when coal-derived materials are reacted with these reagents.

Each phase-transfer methylation experiment requires approximately five days per sample (two days for a duplicate reaction, two days for work-up, and one day for NMR data acquisition). Ether cleavage reactions require an additional five days for sample preparation, reaction and work-up. The reagents, which are commercially available, cost approximately \$20.00 for each phase transfer methylation reaction and about \$50.00 for each single-electron transfer reaction followed by phase transfer methylation. The ^{13}C -NMR spectrometer used in this study costs approximately \$350,000, and the ^1H -NMR spectrometer costs approximately \$150,000.

Further Development

The use of potassium-naphthalene as a single-electron transfer reagent to induce ether cleavage met with success in this limited study. Many different alkali metal, electron shuttling agents and solvent combinations could be applied to the study of fossil fuels. The optimum combination must be determined empirically. It is recommended that a study be undertaken to determine the most effective combination for ether cleavage in coal-derived materials.

The Lewis acid (BBr_3) used in this study proved to be ineffective because of its promotion of degradative condensation reactions as well as ether cleavage. It is recommended that additional studies be made with BBr_3 and other Lewis acids to determine if milder reaction conditions could promote the ether cleavage reactions over the condensation reactions.

Because the determination of the relative proportion of oxygen functional groups ultimately lies in relying on the value obtained for the total oxygen in the samples, it is recommended that total oxygen content be determined in a more rigorous manner than by difference.

The original resids used in this work were obtained by pyridine extraction of distillation resids. The ^{13}C -NMR analysis of the reaction products was conducted in chloroform solution. Although the methylated products are expected to be more soluble than the original resids, chloroform may not completely solubilize the entire product. Therefore, it is recommended that pyridine or some other good coal-liquid solvent be employed as the NMR solvent in future work.

Participant's Statement of Work

Oxygen is the the principal heteroatom in most coal liquids. Though it often is inferred to be important in coal liquefaction chemistry, its chemical form in coal liquids is ill-defined. The derivatization of hydroxyl oxygen (phenolic, alcoholic, and carboxylic) with ^{13}C labeled methyl iodide, followed by ^{13}C -NMR results, in the case of petroleum-derived resids, in clearly distinguishable and quantifiable signals in the NMR. Selective cleavage of etheric bonds followed by derivatization with ^{13}C methyl groups and identification of the derivatized species by ^{13}C -NMR has been demonstrated with model compounds. These techniques can be used to quantitatively speciate oxygen; however, they have not been demonstrated with process-derived coal liquids, or investigated for their ability to answer questions pertaining to the chemistry of coal conversion. As such, they fit well within the scope of the participants program.

The derivatization of the hydroxyl oxygen, and cleavage and derivatization of the etheric oxygen, followed by subsequent characterization by ^{13}C -NMR of a sample set of six samples will allow a demonstration of the value of these techniques for the examination of the oxygen compounds in coal liquefaction-derived resids. Samples will be derivatized with ^{13}C labeled reagents and analyzed via ^{13}C -NMR to yield a total hydroxyl oxygen value for each of the six samples. Three of the samples (as indicated on the attached list) will be selectively cleaved at ether linkages utilizing two different methods. In the one method, the ethers will be cleaved by forming anion radicals with an electron transfer reagent. This reagent will cleave heterocyclic and non-heterocyclic ethers. The cleavage products will be marked with labeled reagents. NMR spectra of the starting material, alkylated starting material, and the alkylated product of the ether-cleaved material will allow for quantitation of the oxygen-containing species in the pyridine-soluble resid samples. The second method entails cleavage of the ether groups in the starting pyridine-soluble resid sample with a boron trihalide reagent to be chosen by the research group at the University of Chicago. This reagent will cleave non-heterocyclic ethers but not heterocyclic ethers. The treated sample then will be alkylated with a ^{13}C labeled reagent and analyzed by ^{13}C NMR. Comparison of the NMR spectra of the starting material, the

alkylated starting material and the alkylated cleaved materials produced by both methods will allow for quantitative speciation of the oxygen-containing compounds in the pyridine-soluble resid samples.

Six samples will be supplied by CONSOL to the University of Chicago. These samples have been selected so that the utility of the method described above for the measurement of the forms of oxygen can be assessed in application with pyridine-soluble coal liquefaction resids. The samples will be supplied to the University of Chicago with the following information, as available: elemental analyses, phenolic -OH concentration by FTIR, calorific value, hydrogen classes by $^1\text{H-NMR}$, and the full history of the sample (plant, process conditions, age, and storage conditions). The six samples are the pyridine-soluble portion of the 850°F distillation residual materials. Sample size will be at least 10 g. The six pyridine-soluble resid samples will be brittle pitch-like materials that will be supplied as approximately minus 60 mesh powder.

Final Report

**THE USE OF SELECTIVE CHEMICAL REACTIONS AND CARBON NMR
SPECTROSCOPY IN THE SPECIATION AND QUANTIFICATION OF OXYGEN
FUNCTIONAL GROUPS IN COAL RESIDS**

Submitted to:

**CONSOL, INC.
Research and Development
4000 Brownsville Road
Liberty, PA 15129**

by:

**Leon M. Stock - Principal Investigator
Carlos Cheng - Research Associate**

**University of Chicago
Department of Chemistry
Chicago, Illinois 60637**

August, 1992

TABLE OF CONTENTS

	Page
LIST OF TABLES	ii
LIST OF FIGURES	iii
I. EXECUTIVE SUMMARY	1
II. EXPERIMENTAL PART	4
III. DISCUSSION	6
IV. RECOMMENDATIONS	14
V. REFERENCES	16
VI. TABLES AND FIGURES	16
VII. APPENDICES	26

LIST OF TABLES

Tables	Page
I. Results of the Phase-Transfer Methylation of CONSOL Samples	16
II. Comparison of Phenolic OH quantities as determined by carbon NMR spectroscopy and IR solution spectroscopy	17
III. Results of the Phase-Transfer Methylation of CONSOL Samples after Reaction with Boron Tribromide	17
IV. Results of the Phase-Transfer Methylation of CONSOL Samples After Reaction with Potassium and Naphthalene	17
V. Comparison of Phenolic Hydroxyl Quantities Before and After Treatment with Boron Tribromide	18
VI. Comparison of Phenolic Hydroxyl Quantities Before and After Single Electron Transfer with Potassium and Naphthalene	18
VII. Comparison of Phenolic Hydroxyl Quantities Before and After Single Electron Transfer with Lithium	19

LIST OF FIGURES

Figures	Page
I. Typical carbon-13 NMR spectrum of resid that has been methylated once with $^{13}\text{CH}_3\text{I}$	19
II. Typical ^2H NMR Spectrum of a resid (CONSOL 04) methylated twice with CD_3I	20
III. IR absorbance spectrum of nonmethylated resid 01 (8 mg/mL in THF) ratioed against a THF blank.	21
IV. IR absorbance spectrum of methylated sample 01 (8.5 mg/mL in THF) ratioed against a THF background.	22
V. The carbon-13 NMR spectrum of a resid sample (CONSOL 03) methylated once with $^{13}\text{CH}_3\text{I}$ in the presence of carbon dioxide	23
VI. The carbon-13 NMR spectrum of sample 01 methylated twice in the presence of 0.05 M concentration of Cr (III) acetylacetonate.	24
VII. The carbon-13 NMR spectrum of sample 01 in the presence of 0.1M concentration of Cr(III) acetylacetonate.	25

The Use of Selective Chemical Reactions and Carbon NMR Spectroscopy In the Speciation and Quantification of Oxygen Functional Groups In Coal Resids

I. Executive Summary

The broad objectives of this program are to identify and evaluate promising analytical techniques that will be useful in coal liquefaction process development and to demonstrate their direct application to coal liquefaction resids. For our phase of the work, there were two subtasks: first, to characterize and quantify the hydroxylic groups that exist in a selected group of resids that were derived from the liquefaction of coal and, second, to study new chemical methods of cleaving ether linkages in these resids and to characterize and quantify these ether groups.

The first task was accomplished by reacting the resids with a phase-transfer base, tetrabutylammonium hydroxide (E1), and subsequently methylating the oxyanions formed with $^{13}\text{CH}_3\text{I}$ (E2). The number and types of hydroxylic groups were determined by

qualitative and quantitative analyses of the carbon NMR spectra of the methylated products.

The carbon NMR spectra of the methylated samples indicated that the hydroxylic groups found in the resids could be subdivided into hindered phenols, unhindered phenols and carboxylic acids. For reasons that are explained thoroughly in the discussion, it is our conclusion that the carboxylic acids are not really native to the resids. Rather, these compounds were incorporated into the samples before they were received in Chicago through oxidative processes during handling and storage. Further, we showed that the concentration of these compounds are increased in side reactions between carbanions and adventitious carbon dioxide. Accordingly, these substances are not significant to the main

thrust of this study. The phenolic content of the six resids that we studied ranged from about 0.5 meq/g to slightly more than 1 meq/g of the resid; approximately two-thirds of the phenols are of the unhindered type and about a third are of the hindered type.

The quantities of phenolic hydroxyl groups determined from the carbon NMR spectroscopy of the methylated samples agree quite well with IR spectroscopic data for 4 of the 6 resid samples. The remaining discrepancies may be attributed to three sources. First, there are some problems in obtaining homogeneous solutions of the samples. Second, whereas the IR values were obtained from the THF solutions of the resids (1), our samples were the pyridine soluble portions of the resids. Third, the IR data require the assumption of an average extinction coefficient, while NMR results are based on direct observations. Time limitations of the research contract did not allow us to study this point.

The second task was accomplished by applying three chemical methods of effecting ether cleavage reactions to the resids, and subsequently phase-transfer methylating the treated samples with $^{13}\text{CH}_3\text{I}$ to determine the extent of cleavage.

In the first method, resid samples were treated with boron tribromide, a Lewis acid which is commonly used to cleave alkyl aryl ethers (especially methyl aryl ethers) and dialkyl ethers (2). The samples were subsequently methylated with $^{13}\text{CH}_3\text{I}$, and the extent of methylation was determined by carbon NMR spectroscopy. Treatment of the resid samples with the Lewis acid gave partly insoluble products. This was accompanied by a substantial decrease in the concentrations of phenolic hydroxyl groups, together with a change in the relative proportion of hindered and unhindered phenolic hydroxyl groups. Clearly, boron tribromide catalyzes degradative condensation reactions and ether cleavage reactions. The degradative condensation reactions apparently occur more rapidly than any ether cleavage reactions. Thus, this method does not appear to be generally suitable for investigations of ether functional groups in resids.

The second method consisted of reacting the resid samples with potassium-naphthalene, a single electron transfer reagent, in dry THF. The samples were

subsequently phase-transfer methylated with $^{13}\text{CH}_3\text{I}$ and their carbon NMR were obtained and analyzed quantitatively. This second method noticeably increased the phenolic hydroxylic groups of two of the three samples and the ratio of hindered to unhindered

CONCLUSION

Hindered Phenols (meq/g) 0.17	SET ether cleavage	Hindered Phenols (meq/g) 0.28
Unhindered phenols (meq/g) 0.37		Unhindered Phenols 0.42

phenolic hydroxylic groups by a modest amount. In contrast to the reactions with the Lewis acid, this single electron transfer reagent preserved the chemical integrity of the samples. From these results, it is our conclusion that single electron transfer still offers the best possibilities of characterizing and quantifying the etheric groups in resids.

The third method was carried out by refluxing a particular resid sample with lithium in dry dioxane, another single electron transfer reaction. The reaction did not increase the quantity of phenolic groups and the relative proportion of hindered to unhindered phenolic groups remained practically unchanged.

The Illinois No. 6 Coal (Burning Star) from which three of the resid samples were derived has around 10% oxygen (determined by difference), about 6 meq/g. Several lines of evidence suggest that about half of these oxygen atoms occur in phenolic compounds (3). The other half are thought to be mainly ethers (3). The resids that were derived from this coal contain much less oxygen than the original coal, about 2.4 meq/g. The new results and the IR data indicate that approximately 0.5 to 1 meq/g occur in phenolic hydroxyl groups, a large decrease from the raw coal. The ether cleavage reactions indicate that about 0.2 meq/g of the remaining 1.4 to 1.9 meq/g oxygen occur in ethers that are susceptible to single electron transfer reagents. Uncertainties about the meaning of the by-difference analytical data for oxygen render any further discussion highly speculative. But if we adopt that view that the by-difference oxygen content is accurate, about half of the atoms in the resid have been speciated with confidence. And assuming the proportion of

the different kinds of organic oxygen compounds are similar in the coals and the resids, it can be inferred that the remaining oxygen atoms must be in the form of single electron transfer reagent-resistant ethers. Since it has been well established (3) that alkyl aryl ethers and benzyl aryl ethers are readily cleaved by potassium/naphthalene, we conclude that 1.2 to 1.7 meq/g of oxygen are present in the form of refractory ethers presumably of the dibenzofuran/dinaphthofuran or the dialkyl type.

II. Experimental:

A. Materials and Equipment:

Sample No.	Coal Feed	Samples	
		Source/Conditions	Comments
01	Illinois No. 6	Run 250	Interstage
02	Illinois No. 6	Run 257	Recycle
03	Illinois No. 6	Run 257	Interstage
04	Pittsburgh	Run 259	Recycle
05	Pittsburgh	Run 259	Interstage
06	Wyodak and Anderson	Run 258	Interstage

Resid samples were obtained from CONSOL, Inc. These resid samples are pyridine-soluble fractions of distillation residues obtained from the liquefaction of various coals (characteristics of samples are enumerated above). Analytical information on these samples can be found in the Appendix. Tetrabutylammonium hydroxide (40% by weight aqueous solution), lithium ribbon (0.75 mm thick, 19 mm wide, 99.9%) and boron tribromide (1.0 M in CH_2Cl_2) were purchased from Aldrich; $^{13}\text{CH}_3\text{I}$ and CD_3I were purchased from Cambridge Isotope Laboratories. THF was initially dried and refluxed over lithium aluminum hydride and dried further over potassium benzophenone ketyl. Dioxane was dried directly over sodium benzophenone ketyl.

Carbon-13 NMR spectra were recorded on the General Electric QE-300 Mhz Spectrometer and ^2H NMR spectra were recorded on the Varian XL-400 Spectrometer.

The carbon NMR spectra were obtained using CDCl_3 solutions while deuterium NMR spectra were obtained using methylene chloride solutions.

B. Phase-Transfer Methylation:

The resid sample (0.5 g) was stirred in THF (15 - 20 mL) under nitrogen. After stirring for 30 minutes, 40% aqueous tetrabutylammonium hydroxide solution (3 mL ~4.5 mmol) was added. The solution was allowed to stir for four hours, after which the flask was cooled in an ice/water bath, and $^{13}\text{CH}_3\text{I}$ was added via a 1.0 mL syringe (0.4 mL ~6.4 mmol). The nitrogen inlet was then replaced with a glass stopper. After 48 hours of stirring, the solution was quenched with 1.0 N HCl (until neutral pH). The solvent was removed by rotary evaporation, and water (50 mL) was added to the resid. The resid/water mixture was sonicated and then filtered and washed with at least one liter of water. The mixture was finally washed with 50/50 water/methanol solution (100 mL), dried on the filter, and dried under vacuum for several hours.

The dried sample was dissolved once more in THF (15 - 20 mL), stirred and subjected to exactly the same procedure described above to ensure complete methylation.

C. Carbon-13 NMR Spectroscopy:

N-butyl ether (30 - 60 mg) was weighed directly into a preweighed NMR tube. Next, the dried methylated resid (100 mg) was weighed into the tube. A dark-purple solution of Cr(III) acetylacetonate in CDCl_3 (70 mg in 1.5 mL CDCl_3) was prepared and added finally to the mixture in the NMR tube.

The NMR signals in appropriately broadened spectra were quantified by cutting out each of the peaks and weighing them on the microgram balance. For every spectrum, at least 3 photocopies were cut out to determine the number of phenolic hydroxylic groups corresponding to the signals.

D. Reaction with Boron Tribromide

In a 100 mL round bottom flask, the resid sample (1 g) was dissolved and stirred in methylene chloride (40 mL). The flask was equipped with a condenser. Boron tribromide

(16 mL of 1.0 M solution in methylene chloride ~ 16 mmol) was quickly added to the flask by means of a syringe. The solution was then refluxed for 24 hours. The solution was cooled and quenched with water, then neutralized with 0.1 N NaOH. The solvent was removed by rotary evaporation, and water (100 mL) was added to the resid. The mixture was then filtered and washed further with water (1 L) and dried under vacuum.

The dried boron tribromide-treated resid was then methylated as in B.

E. Single Electron Transfer with Potassium and Naphthalene

Potassium (2 g ~ 51 mmol) was added to dry THF (80 mL). Naphthalene (0.6 g) was then added and the solution was stirred until it turned a dark-blue color. The resid (2 g) was finally added, and stirring continued (under nitrogen) for 100 hours. At the end of 100 hours, the excess potassium was collected on a filter. The solvent was removed by rotary evaporation, and the mixture was washed repeatedly with acetonitrile (3 X 25 mL) to remove naphthalene. Water (100 mL) was added to the resid mixture. The solution was neutralized with aqueous HCl. The mixture was sonicated, filtered and washed with excess water (1 L). After drying on the filter, the resid was dried under vacuum for several hours, and subjected to methylation as in B.

F. Single Electron Transfer with Lithium Ribbon

Pieces of lithium (0.2 g ~ 29 mmol) were cut and weighed under Ar and added to a flask filled with dry dioxane (30 mL) that was equipped with a reflux condenser and an inlet for Ar. Resid (2 g) was subsequently added to the mixture. The mixture was refluxed for 24 hours. After refluxing, excess lithium was collected on a filter. The solvent was removed by rotary evaporation. Water (100 mL) was added to the mixture, and the solution was neutralized with aqueous HCl. The mixture was sonicated, filtered and washed with excess water (1 L). After drying on the filter, the resid was dried under vacuum for several hours, and subjected to methylation as in B.

III. Discussion:

The characterization and quantification of hydroxylic species in the CONSOL samples were carried out by reacting the resids with tetrabutylammonium hydroxide to form the oxyanions (E1) and subsequently methylating the oxyanions with $^{13}\text{CH}_3\text{I}$ (E2). The

carbon NMR spectra showed the presence of resonances between 50 and 60 ppm which are indicative of successful methylation (Figure I). In addition, the carbon spectra also showed a signal at 29 ppm. This signal is due to methylene ^{13}C native to the resid samples. When methylation was carried out with CD_3I , the ^2H NMR of the methylated sample showed the presence of a broad peak between 3 and 4 ppm (Figure II), which is further evidence of successful methylation.

In the carbon spectra, the resonances of the O-methylation products typically appear in three distinct regions (2)(Figure I). The first is centered at about 61 ppm, which is attributed to the methylation of hindered phenols. The second is centered at 55 ppm; this prominent peak is attributed to the methylation of simple unhindered phenols. The third signal is centered at 51 ppm; this is due to the methylation of aliphatic and aromatic carboxylic acids.

It is pertinent to note that the NMR assignments are based on a detailed study in which chemical and spectroscopic methods were used to assign resonances in coal (3, 4). The assignments are summarized in Scheme 1.

Scheme 1

Region 1 (Region centered at 61 ppm)

Methyl Vinyl Ethers

Primary Alkyl Methyl Ethers, RCH_2OCH_3

Primary Benzyl Methyl Ethers, $\text{ArCH}_2\text{OCH}_3$

Hindered Aryl Methyl Ethers

Region 2 (Region Centered at 55 ppm)

Secondary Alkyl Methyl Ethers, R_2CHOCH_3

Secondary Benzyl Methyl Ethers, $ArCH(R)OCH_3$

Unhindered Aryl Methyl Ethers, $PhenylOCH_3$, $NaphthylOCH_3$

Region 3 (Region Centered at 51 ppm)

Tertiary Alkyl Methyl Ethers, R_3COCH_3

Methyl Alkyl and Aryl Esters, $ArCOOCH_3$ and $RCOOCH_3$

The data tabulated in Tables I, III and IV were obtained from samples that were methylated twice (See Experimental Section) with $^{13}C_3H_7I$ to ensure complete methylation. This precaution was taken notwithstanding our IR observations (Figures III and IV) of CONSOL Sample 01 which suggested that a single treatment with $^{13}C_3H_7I$ completely methylated the resid.

The quantity of carboxylic hydroxyl groups in the samples appeared to be excessive. Tables I and IV show that treatment with potassium and naphthalene followed by methylation leads to an unexpected increase in the abundance of carboxylic groups. This suggests that the carboxylic signal may be largely an artifact; the phase-transfer base tetrabutylammonium hydroxide has a tendency to absorb carbon dioxide from the atmosphere, and entrapment by carbanions (E3) (3) may lead to the increase in the

resonance centered at 51 ppm. The plausibility of this line of reasoning was confirmed when CONSOL Sample 03 was methylated once in the presence of carbon dioxide. The carbon spectrum of this particular sample exhibited a greatly increased resonance at 51 ppm (compare Figures I and V). In addition, the CONSOL Coal Company, which prepared the resid samples, has suggested that much of the carbonyl oxygen may have been acquired during sample preparation and storage. We therefore thought it best to disregard the signals at 51 ppm.

Under ideal conditions, the areas of the carbon resonances are directly proportional to the number of carbon nuclei which gives rise to the signals. Unfortunately, this is not generally true due to several factors (5). First, carbon nuclei even within the same molecule can differ in their spin-lattice relaxation times, T_1 , by several orders of magnitude. And in cases where proton-carbon decoupling is necessary, the Nuclear Overhauser Enhancement (NOE) limits the quantitative usefulness of carbon NMR spectra. In taking NMR spectra, we have therefore utilized conditions recommended to circumvent the above problems, namely, the use of relatively long relaxation times (2 seconds), a flip angle of 90° , the use of gated decoupling (meaning proton-carbon decoupling takes place only during acquisition) and the use of a paramagnetic compound, Cr (III) acetylacetonate, which provides an effective relaxation mechanism to shorten T_1 values and quench NOE. Preliminary experiments showed that under the above conditions and with the addition of 0.03 to 0.05M concentrations Cr (III) acetylacetonate, relative concentrations of pure compounds anisole and methyl caproate could be obtained quantitatively.

However, when the carbon NMR spectrum of the CONSOL resids were obtained in the presence of 0.05 M Cr (III) acetylacetonate and n-butyl ether as internal standard, the

signals of the standard were very narrow compared to the resonances of the resid. In addition, the four carbon atoms due to n-butyl ether did not integrate equally as required (Figure VI). As the concentration of the chromium compound was increased, the peaks of n-butyl ether noticeably widened until at about 0.1M Cr (III), the peaks did not change further. At this point, the four signals of the standard gave equal integrations (Figure VII). Therefore 0.1 M Cr (III) acetylacetonate was used in all quantitative carbon NMR analyses.

Table II compares the quantity of phenolic hydroxyl groups that were determined by carbon NMR spectroscopy with the values that were obtained from IR solution spectroscopy (carried out by the CONSOL Coal Company) (1). Reproducibility for the IR values is ± 0.09 (1). For the carbon NMR data, duplicate experiments of CONSOL 02 and 03 establishes a comparable precision of ± 0.1 . The agreement between the two sets of values is quite good for samples 01 to 04, where the accuracy is $\leq \pm 0.1$ and much less so for the other two samples (accuracy approximately ± 0.4). There are several plausible reasons for this discrepancy. First, it should be pointed out that whereas the samples subjected to phase-transfer methylation and carbon NMR spectroscopy were the pyridine-soluble fractions of coal liquefaction resids, the IR values were obtained from THF solutions of whole resids. Second, we also observed that, with the exception of sample 05, all the other samples were quite soluble in chloroform, the solvent used to obtain the carbon NMR spectra. The chloroform solution of sample 05 required sonication for one hour before it became homogeneous enough for NMR spectroscopy. Third, whereas the IR data required the assumption of an average extinction coefficient, NMR results are based on direct observations.

It should also be mentioned that the resids were only partly soluble in THF. In taking the IR spectra of sample 01, we observed that whereas the unreacted sample did not form a homogeneous solution in THF, the methylated sample 01 dissolved in THF with ease. Hydrogen bonds between the hydroxylic groups constitute some of the attractive forces that

bind the nonmethylated resid clusters together. Methylation disrupts these hydrogen bonds, therefore rendering the resid more soluble.

Three chemical methods of cleaving ethers were utilized on the CONSOL resids. In the first method, the resids were refluxed with boron tribromide, which has been reported to cleave alkyl aryl ethers and even dialkyl ethers (2, 6). In the second method, the resids were treated with potassium and naphthalene, a single electron-transfer reagent which has been reported to cleave diaryl ethers and alkyl aryl ethers (7). In the third method, one resid sample was refluxed with lithium metal, which was reported to cleave dibenzofuran-type compounds (8).

The products of the boron tribromide-treated reactions were subsequently subjected to phase-transfer methylation with $^{13}\text{CH}_3\text{I}$. If boron tribromide had succeeded in effecting ether cleavage, the increase in methylation would be detected and measured by carbon NMR spectroscopy. We observed that treatment with the Lewis acid led to the formation of products that were less soluble in chloroform. A comparison of the quantity of phenolic hydroxyl groups of CONSOL samples 03, 05 and 06 before and after treatment with boron tribromide (refer to Table V) shows that, for samples 03 and 05, the quantity of hydroxyl groups dropped steeply and less so for sample 06.

The cleavage of ethers by boron tribromide have been carried out under reaction conditions of varying severity (2, 6, 9). We chose to reflux the resids with the Lewis Acid in methylene chloride as recommended by Bhafl and Kulkarni (9). Apparently, this leads to predominant degradative condensation, as suggested by Table V. However, an examination of the ratio of hindered to unhindered phenolic hydroxyl groups in Table V shows a substantial increase in samples 03 and 05 when these resids were reacted with boron tribromide, which strongly suggests that ether cleavage had taken place concurrently with the condensation reactions. We therefore suggest that milder reaction conditions could be found which should promote ether cleavage reactions at the expense of degradative condensation reactions.

In the second method, samples 03, 05 and 06 were subjected to a single electron transfer reaction with potassium and naphthalene in dry THF (E4, E5), and subsequently phase-transfer methylated with $^{13}\text{CH}_3\text{I}$ (E6). In contrast to the reactions with the Lewis

acid, no change was observed in the solubility of the resid samples before and after single electron transfer. An examination of Table VI indicate that sample 06 did not undergo ether cleavage. Both the quantity of hindered and unhindered phenolic hydroxyl groups remained the same within experimental errors. For sample 03, the quantity of hindered phenolic hydroxyl groups went up by more than 20% while that of the unhindered phenolic hydroxyl groups remained the same within experimental errors, leading to a slight but significant increase in the ratio of hindered to unhindered phenolic hydroxyl groups. For sample 05, both types of phenolic hydroxyl groups increased, with the hindered phenolic hydroxyl groups increasing by more than 50%, while the increase in the unhindered phenolic hydroxyl groups did not exceed the limits of experimental errors, leading to a significant increase in the ratio of hindered to unhindered phenolic hydroxyl groups. The data for these samples suggest that a modest amount of ether cleavage was effected by potassium and naphthalene.

In the third method, sample 02 was subjected to another single electron transfer reaction with lithium as the electron source in the absence of any electron shuttling agent, and with the use of dry dioxane as solvent. Keumi and associates (8) have reported the successful cleavage of dibenzofurans using lithium as an electron source and since heterocyclic aromatic carbon-oxygen bonds are the most resistant towards cleavage, we had hoped that this reaction would dramatically increase the quantity of phenolic groups. After refluxing the resid and lithium in dioxane for 24 hours, the resid was once again phase-transfer methylated with $^{13}\text{CH}_3\text{I}$. The NMR analysis of the methylated resid showed no significant change in the quantity of both hindered and unhindered phenolic hydroxylic groups (Refer to Table VII).

Elemental analyses of the parent coals of the resid samples indicate that these coals possess oxygen contents ranging from 10 - 14 % by weight. More particularly, Illinois No. 6 coal (Burning Star), from which three of the samples were derived possess around 10% oxygen, which corresponds to 6 meq/g. In contrast, the elemental analyses of the derived resid samples indicate that these resids contain about 40% of the oxygen found in the coals, around 2.4 meq/g. Our analytical results and IR data (Refer to table II) indicate that about 0.5 to 1 meq/g of the 2.4 meq/g occur in phenolic hydroxyl groups. The ether cleavage reactions indicate that about 0.2 meq/g of the remaining 1.4 to 1.9 meq/g oxygen occur in ethers that are susceptible to single electron transfer cleavage reactions. This means that less than half of the oxygen atoms in the resids have been speciated with confidence. What about the other half? What kind of oxygen compounds comprise the other half?

Several lines of evidence suggest that half of the oxygen atoms in Illinois No. 6 coal consists of phenolic compounds, and the other half consists mainly of ether compounds (3). If we assume this proportion to hold in the resids, then it can be inferred that the unspciated half must consist of ethers. Since it has been well established (3) that alkyl aryl ethers and aryl benzyl ethers are readily cleaved by potassium/naphthalene, it can be

inferred that 1.2 to 1.7 meq/g of oxygen are present in the form of refractory ethers presumably of the dibenzofuran/dinaphthofuran type, but the dialkyl type cannot be excluded on the basis of our observations. But this conclusion holds true only to the extent that elemental analyses of the resids are accurate. It should be pointed out that the oxygen content is determined by difference, and therefore subject to the sum of the errors of all the other elements. Further, these resids have a relatively low oxygen content (3 - 4% oxygen) so that an error of 1 to 2% would lead to a very different proportion of ethers. Therefore, our conclusions about the amount of ethers in these resids are highly conjectural.

IV. Recommendations:

The phase-transfer methylation of coal liquefaction resids and the carbon NMR spectroscopic analysis of the methylated resids is an effective, precise method of quantifying and speciating the hydroxylic groups in the resids. The following steps are recommended to ensure that the areas of carbon resonances are directly proportional to the number of carbon nuclei giving rise to the resonances. First, the use of gated decoupling. Second, the use of a flip angle of 90° and a T_1 of a few seconds. Third, and most importantly, use at least 0.1M of Cr (III) acetylacetonate as a relaxing agent.

The discrepancy in the quantity of phenolic groups for sample 05 as determined by IR and carbon NMR is most probably due to the difference in solubility of the sample in THF and chloroform. The use of another NMR solvent in which sample 05 is more soluble may close the gap between the two values.

The use of chemical methods to cleave ethers in the resids met with little success. The use of boron tribromide, a Lewis acid, led to predominant degradative condensation instead of the desired ether cleavage. This compound has been reported to cleave dialkyl ethers (2), which are almost impossible to cleave by any other method. Any continued investigation of the use of Lewis acids to cleave ethers should focus on the use of milder reaction conditions to promote ether cleavage and to minimize degradative condensation reactions.

Compared to Lewis acids, single electron transfer methods are much less destructive to resids. Studies on ether cleavage reactions and the characterization of ether compounds in resids should therefore focus on single electron transfer reagents. Our preliminary studies suggest that lithium by itself is practically useless and that potassium/naphthalene leads to minimal amount of ether cleavage within resids. The literature contains extensive information (9) on single electron transfer-induced ether cleavage reactions using lithium, sodium and potassium with varying electron shuttling agents and solvents. It is very difficult to predict which combination of alkali metal, electron shuttling agent and solvent will best promote ether cleavage reactions in fossil fuels. The effectivity of each particular combination can only be determined empirically.

Our assumption that the other oxygen compounds within the resids exist predominantly in the form of ethers is based entirely on elemental analyses of oxygen (determined by difference), and on the known distribution and proportion of oxygen compounds in the coals from which these were derived. Since the % oxygen is determined by difference, it is of course subject to the sum of the errors of all the other elements. And since these resids contain merely a fraction (3 - 4% oxygen by weight for resids compared to 10 - 14% oxygen for coals) of the oxygen of the parent coals, an error of 1 or 2% (which is reasonable) would suggest a much smaller amount of ethers or even the absence of ethers. The very modest increase in the quantities of both hindered and unhindered phenols resulting from the treatment of the resids with potassium/naphthalene, which has been shown to be very effective for the cleavage of alkyl aryl ethers and benzyl aryl ethers (2) in fossil fuels definitely suggest that there is very little of these types of compounds in these resids. We can therefore infer that the ethers, if they exist, must be mostly dialkyl and/or benzofuran/dibenzofuran type ethers.

It is therefore highly important to obtain more reliable estimates of oxygen content, preferably direct estimates. Only then would we have a reliable estimate of the relative quantities and proportion of ether groups in resids.

V. References

- 1). G. A. Robbins, R. A. Winschel, F. P. Burke, Prepr. Am. Chem. Fuel Soc., Div. of Fuel Chem., **30** (4), 155 - 163 (1986).
- 2). F. L. Benton, T. E. Dillon, J. Am. Chem. Soc., **64**, 1128 (1942).
- 3). L. M. Stock, R. S. Willis, Journal of Organic Chemistry, **50**, 3566 (1985).
- 4). L. M. Stock, New Trends in Coal Science, Y. Yurum (ed.), Kruwer Academic Publishers, 287 - 303 (1988).
- 5). J. N. Shoolery, Progress in NMR Spectroscopy, Vol. II, 79 - 93 (1977).
- 6). J. F. W. McOmie, M. L. Watts, Chem. Ind. (London), 1658 (1963).
- 7). D. R. Moore, J. Org. Chem., **26**, 3596 (1961).
- 8). T. Keumi, C. Murata, Y. Sasaki, H. Kitajima, Synthesis, 634 (1980).
- 9). M. Vivekananda Bhatt, S. U. Kulkarni, Synthesis, 249 - 282 (1983).

VI. Tables and Figures:

Table I. Results of the Phase-Transfer Methylation of CONSOL Samples^a

Sample No.	[OH] in meq/g resid			Total Phenolic [OH] meq/g	Ratio of Hindered to Unhindered Phenolic [OH]
	Hindered Phenolic [OH]	Unhindered Phenolic [OH]	Carboxylic [O ₂ C]		
01	0.31	0.77	0.46	1.08	0.40
02	0.22	0.37	0.45	0.59	0.59
03	0.25	0.52	0.36	0.77	0.48
04	0.25	0.52	0.33	0.77	0.48
05	0.17	0.37	0.24	0.54	0.46
06	0.25	0.52	0.36	0.77	0.48

^a Sample 02 and 03 are the average of two determinations. The Carbon NMR spectra of the samples are included in the appendix.

Table II. Comparison of Phenolic OH quantities as determined by carbon NMR spectroscopy and IR solution spectroscopy

Sample No.	IR ^a	Phenolic [OH] in meq/g resid	¹³ C NMR
01	1.07		1.08
02	0.50		0.59
03	0.68		0.77
04	0.69		0.77
05	0.92		0.54
06	1.09		0.77

^aThe IR values were determined using IR solution spectroscopy in THF. Values were determined by CONSOL Coal Company. For experimental details, refer to Reference 1.

Table III. Results of the Phase-Transfer Methylation of CONSOL Samples after Reaction with Boron Tribromide^a

Sample No.	[OH] in meq/g resid			Total Phenolic [OH]	Ratio of Hindered to Unhindered Phenolic [OH]
	Hindered Phenolic [OH]	Unhindered Phenolic [OH]	Carboxylic [OH]		
03	0.15	0.18	0.27	0.33	0.83
05	0.11	0.18	0.07	0.29	0.61
06	0.23	0.46	0.28	0.69	0.50

^aThe Carbon NMR spectra of the samples are included in the Appendix.

Table IV. Results of the Phase-Transfer Methylation of CONSOL Samples After Reaction with Potassium and Naphthalene^a

Sample No.	[OH] in meq/g resid			Total Phenolic [OH]	Ratio of Hindered to Unhindered Phenolic [OH]
	Hindered Phenolic [OH]	Unhindered Phenolic [OH]	Carboxylic [OH]		
03	0.32	0.48	0.79	0.80	0.67

Sample No.	[OH] in meq/g resid			Total Phenolic [OH]	Ratio of Hindered to Unhindered Phenolic [OH]
	Hindered Phenolic [OH]	Unhindered Phenolic [OH]	Carboxylic [OH]		
05	0.28	0.42	0.52	0.71	0.67
06	0.25	0.49	0.43	0.74	0.51

*The Carbon NMR spectra of the samples are included in the Appendix.

Table V. Comparison of Phenolic Hydroxyl Quantities Before and After Treatment with Boron Tribromide

Sample No.	Before Treatment with BBr ₃				After Treatment with BBr ₃			
	OH in meq/g resid		Total Phenolic OH	Ratio	OH in meq/g resid		Total Phenolic OH	Ratio
	Hindered Phenolic OH	Unhindered Phenolic OH			Hindered Phenolic OH	Unhindered Phenolic OH		
03	0.25	0.52	0.77	0.48	0.15	0.18	0.33	0.83
05	0.17	0.37	0.54	0.46	0.11	0.18	0.29	0.61
06	0.25	0.52	0.77	0.48	0.23	0.46	0.69	0.50

Table VI. Comparison of Phenolic Hydroxyl Quantities Before and After Single Electron Transfer with Potassium and Naphthalene

Sample No.	Before Treatment with K/Naphthalene				After Treatment with K/Naphthalene			
	OH in meq/g resid		Total Phenolic OH	Ratio	OH in meq/g resid		Total Phenolic OH	Ratio
	Hindered Phenolic OH	Unhindered Phenolic OH			Hindered Phenolic OH	Unhindered Phenolic OH		
03	0.25	0.52	0.77	0.48	0.32	0.48	0.80	0.67
05	0.17	0.37	0.54	0.46	0.28	0.42	0.71	0.67
06	0.25	0.52	0.77	0.48	0.25	0.49	0.74	0.51

Table VII. Comparison of Phenolic Hydroxyl Quantities Before and After Single Electron Transfer with Lithium

Sample No.	Before Treatment with Lithium				After Treatment with Lithium			
	OH in meq/g resid		Total Phenolic OH	Ratio	OH in meq/g resid		Total Phenolic OH	Ratio
	Hindered Phenolic OH	Unhindered Phenolic OH			Hindered Phenolic OH	Unhindered Phenolic OH		
02	0.22	0.37	0.59	0.59	0.23	0.40	0.63	0.58

Figure I

Typical carbon-13 NMR spectrum of resid (CONSOL 03) that has been methylated once with $^{13}\text{CH}_3\text{I}$. The resonances at 14, 19, 32 and 70 ppm are due to the 4 carbons of n-butyl ether.

Figure II
Typical ^2H NMR Spectrum of a resid (CONSOL 04) methylated twice with CD_3I .
The resonance at 5.3 ppm is due to the solvent, methylene chloride, of natural deuterium abundance and resonance at 6 ppm is due to $\text{CDCl}_2\text{CDCl}_2$.

Figure III
IR absorbance spectrum of non-methylated resid 01 (8 mg/mL in THF) ratioed against a THF blank. The phenolic O-H stretch is found at 3300 cm^{-1}

Figure IV
IR absorbance spectrum of methylated sample 01 (8.5 mg/mL in THF) ratioed against a THF background. The phenolic O-H stretch at 3300 cm^{-1} is absent.

Figure V

The carbon-13 NMR spectrum of a resid sample (CONSOL 03) methylated once with $^{13}\text{CH}_3\text{I}$ in the presence of carbon dioxide. Note the increased intensity of the resonance at 50 - 51 ppm due to methylation of carboxylic acids (compare with Figure I). The signals at 12, 18, 30 and 69 ppm are due to the 4 carbons of n-butyl ether. The smaller signals at 12, 18, 23 and 58 ppm are due to the carbons of tetrabutylammonium hydroxide.

Figure VI
The carbon-13 NMR spectrum of sample 01 methylated twice in the presence of 0.05 M concentration of Cr (III) acetylacetonate. Note the narrow carbon resonances of n-butyl ether and their unequal areas.

Figure VII

The carbon-13 NMR spectrum of sample 01 in the presence of 0.1M concentration of Cr(III) acetylacetonate. The carbon resonances of n-butyl ether have become wider. The areas integrate equally.

VII. Appendices

A. Raw Data and Calculations

Table VIII. Raw Data for Table I

Sample No.	Wt. n-butyl ether (g)	Wt. Resid (g)	Wt. Signal 70 ppm (g)	Wt Signal 61 ppm (g)	Wt. Signal 55p ppm (g)	Wt. Signal 51 ppm
01	0.0286	0.115	0.00858	0.06454	0.16066	0.09721
02A	0.0303	0.0993	0.01270	0.05378	0.09415	0.12223
02B	0.0452	0.0815	0.02653	0.06569	0.10719	0.12201
03A	0.0316	0.0943	0.01868	0.09243	0.19308	0.12458
03B	0.0367	0.1142	0.0250	0.11046	0.21786	0.13957
04	0.0357	0.0756	0.02453	0.07720	0.16390	0.10274
05	0.0238	0.0939	0.02363	0.09742	0.20899	0.13873
06	0.0253	0.0936	0.01469	0.08150	0.17154	0.11748

Sample Calculation:

Sample 01, Table I

$$\frac{0.0286 \text{ g n-butyl ether}}{130.23 \text{ g/mole}} \times 2 \times 1000 \times 0.0108 = 0.0047 \text{ meq } ^{13}\text{C for } \alpha\text{C n-butyl ether}$$

where 0.0108 is factor for ^{13}C natural abundance

$$0.00858 \text{ g 70ppm} = 0.0047 \text{ meq } \alpha\text{C } ^{13}\text{C}$$

$$\text{quantity of hindered phenolic OH} = \frac{0.06454}{0.00858} \times \frac{0.0047}{0.115} = 0.31 \text{ meq/g resid}$$

$$\text{quantity of unhindered phenolic OH} = \frac{0.16066}{0.00858} \times \frac{0.0047}{0.115} = 0.77 \text{ meq/g}$$

$$\text{quantity of carboxylic OH} = \frac{0.09721}{0.00858} \times \frac{0.0047}{0.115} = 0.46 \text{ meq/g}$$

Table IX. Raw Data for Table III

Sample No.	Wt. n-butyl ether (g)	Wt. Resid	Wt. Signal 70 ppm (g)	Wt Signal 61 ppm (g)	Wt. Signal 55p ppm (g)	Wt. Signal 51 ppm
03	0.0214	0.0469	0.03584	0.07016	0.08464	0.13030
05	0.0364	0.0912	0.03054	0.04982	0.08470	0.03414
06	0.0404	0.0898	0.03479	0.10589	0.21346	0.13228

Table X. Raw Data for Table IV

Sample No.	Wt. n-butyl ether (g)	Wt. Resid	Wt. Signal 70 ppm (g)	Wt Signal 61 ppm (g)	Wt. Signal 55p ppm (g)	Wt. Signal 51 ppm
03	0.0273	0.0907	0.01253	0.08220	0.12219	0.20000
05	0.0390	0.0850	0.02183	0.08167	0.11880	0.14714
06	0.0286	0.0897	0.02408	0.11466	0.22390	0.19774

B. NMR Spectra for Tables I, III and IV

Table I, 01

Table I, 02A

Table I, 02B

Table I, 03A

Table I, 03B

Table I, 04

Table I, 05

Table I, 06

Table III, 03

Table III, 05

Table III, 06

Table IV, 03

Table IV, 05

Table IV, 06

C. Analytical Data for Resid Samples

ANALYSIS OF PYRIDINE-SOLUBLE PORTION
OF THE DISTILLATION RESID

Ref. No. 1

Plant: Wilsonville
Run Number: 250
Sample Designator: R-1236
Sampling Point: Interstage
Feed Coal: Illinois No.6
Burning Star #2

Ultimate (Wt%, as Det.)

Carbon:	88.18
Hydrogen:	6.04
Nitrogen:	1.60
Sulfur:	0.49
Oxygen (diff.)	3.69

Proton Distribution (%)

Condensed Aromatics	27.8
Uncondensed Aromatics	6.0
Cyclic Alpha	17.7
Alkyl Alpha	9.4
Cyclic Beta	14.6
Alkyl Beta	13.8
Gamma	10.7

Phenolic -OH Concentration
(THF-soluble basis, meq/g): 1.07

Phenolic Peak Location (cm⁻¹): 3292

Pyridine Soluble (% of Resid): 91.09

ANALYSIS OF PYRIDINE-SOLUBLE PORTION
OF THE DISTILLATION RESID

Ref. No. 2

Plant: Wilsonville
Run Number: 257
Sample Designator: V-131B
Sampling Point: Recycle
Feed Coal: Illinois No.6
Burning Star #2

Ultimate (Wt%, as Det.)

Carbon:	87.52
Hydrogen:	7.38
Nitrogen:	0.90
Sulfur:	0.15
Oxygen (diff.):	4.05

Proton Distribution (%)

Condensed Aromatics	16.1
Uncondensed Aromatics	4.8
Cyclic Alpha	15.6
Alkyl Alpha	9.2
Cyclic Beta	19.3
Alkyl Beta	21.6
Gamma	13.3

Phenolic -OH Concentration
(THF-soluble basis, meq/g): 0.50

Phenolic Peak Location (cm^{-1}): 3302

Pyridine Soluble (% of Resid): 86.49

ANALYSIS OF PYRIDINE-SOLUBLE PORTION
OF THE DISTILLATION RESID

Ref. No. 3

Plant: Wilsonville
Run Number: 257
Sample Designator: R-1235
Sampling Point: Interstage
Feed Coal: Illinois No.6
Burning Star #2

Ultimate (Wt%, as Det.)

Carbon:	87.72
Hydrogen:	7.32
Nitrogen:	0.97
Sulfur:	0.28
Oxygen (diff.):	3.71

Proton Distribution (%)

Condensed Aromatics	17.3
Uncondensed Aromatics	6.3
Cyclic Alpha	15.8
Alkyl Alpha	8.7
Cyclic Beta	18.6
Alkyl Beta	19.8
Gamma	13.3

Phenolic -OH Concentration
(THF-soluble basis, meq/g): 0.68

Phenolic Peak Location (cm⁻¹): 3300

Pyridine Soluble (% of Resid): 80.21

ANALYSIS OF PYRIDINE-SOLUBLE PORTION
OF THE DISTILLATION RESID

Ref. No. 4

Plant: Wilsonville
Run Number: 259
Sample Designator: V-131B
Sampling Point: Recycle
Feed Coal: Pittsburgh Seam
Ireland Mine

Ultimate (Wt%, as Det.)

Carbon:	89.83
Hydrogen:	6.38
Nitrogen:	1.19
Sulfur:	0.38
Oxygen (diff.)	2.22

Proton Distribution (%)

Condensed Aromatics	23.5
Uncondensed Aromatics	6.0
Cyclic Alpha	17.8
Alkyl Alpha	9.2
Cyclic Beta	14.6
Alkyl Beta	16.8
Gamma	12.1

Phenolic -OH Concentration
(THF-soluble basis, meq/g): 0.69

Phenolic Peak Location (cm^{-1}): 3296

Pyridine Soluble (% of Resid): 84.51

ANALYSIS OF PYRIDINE-SOLUBLE PORTION
OF THE DISTILLATION RESID

Ref. No. 5

Plant: Wilsonville
Run Number: 259
Sample Designator: R-1235
Sampling Point: Interstage
Feed Coal: Pittsburgh Seam
Ireland Mine

Ultimate (Wt%, as Det.)

Carbon:	89.27
Hydrogen:	6.07
Nitrogen:	1.31
Sulfur:	0.62
Oxygen (diff.):	2.73

Proton Distribution (%)

Condensed Aromatics	26.2
Uncondensed Aromatics	8.3
Cyclic Alpha	17.1
Alkyl Alpha	8.9
Cyclic Beta	14.0
Alkyl Beta	15.0
Gamma	10.5

Phenolic -OH Concentration
(THF-soluble basis, meq/g): 0.92

Phenolic Peak Location (cm⁻¹): 3295

Pyridine Soluble (% of Resid): 83.63

ANALYSIS OF PYRIDINE-SOLUBLE PORTION
OF THE DISTILLATION RESID

Ref. No. 6

Plant: Wilsonville
Run Number: 258
Sample Designator: R-1235
Sampling Point: Interstage
Feed Coal: Wyodak and Anderson Seam
Black Thunder Mine

Ultimate (Wt%, as Det.)

Carbon:	89.19
Hydrogen:	5.59
Nitrogen:	1.57
Sulfur:	0.06
Oxygen (diff.)	3.59

Proton Distribution (%)

Condensed Aromatics	29.2
Uncondensed Aromatics	7.3
Cyclic Alpha	17.4
Alkyl Alpha	9.0
Cyclic Beta	12.1
Alkyl Beta	14.2
Gamma	10.8

Phenolic -OH Concentration
(THF-soluble basis, meq/g): 1.09

Phenolic Peak Location (cm^{-1}): 3294

Pyridine Soluble (% of Resid): 71.45

END

DATE
FILMED

4 / 3 / 93