PREDICTIONS OF A NON-GAUSSIAN MODEL FOR LARGE SCALE STRUCTURE

Z. H. Fan & J. M. Bardeen

*Department of Physics, FM-15
The University of Washington
Seattle, Washington 98195*

June 26, 1992

RECEIVED
JUN 29 1992
COTI

PREPARED FOR THE U.S. DEPARTMENT OF ENERGY

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Department of Energy, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the product or process disclosed, or represents that its use would not infringe privately-owned rights. By acceptance of this article, the publisher and/or recipient acknowledges the U.S. Government's right to retain a nonexclusive, royalty-free license in and to any copyright covering this paper.

MASTER

ABSTRACT

A modified CDM model for the origin of structure in the universe, based on an inflation model with two interacting scalar fields, is analyzed to make predictions for the statistical properties of the density and velocity fields and the microwave background anisotropy. The initial gauge-invariant potential ζ which is defined as $\zeta = \delta\rho/(\rho + p) + 3\varphi$, where φ is the curvature perturbation amplitude and p is the pressure, is the sum of a Gaussian field ϕ_1 and the square of a Gaussian field ϕ_2 . We assume a Harrison-Zel'dovich scale-invariant power spectrum for ϕ_1 , and a log-normal 'peak' power spectrum for ϕ_2 . The location and the width of the peak are described by parameters k_c and a , respectively. Comparing our theoretical calculations for two dimensional density angular correlation function to the results of the APM survey, we set certain parameter regime for k_c , a and the amplitude ratio of Gaussian and non-Gaussian fields. The model is motivated to some extent by inflation models with two interacting scalar fields, but is mainly interesting as an example of a model whose statistical properties change with scale. On small scales, it is almost identical to a standard scale-invariant Gaussian CDM model. On scales near the location of the peak of the non-Gaussian field, the distributions have long tails in high positive values of the density and velocity fields. Thus, it is easier to get large-scale streaming velocities than the standard CDM model. We calculate the quadrupole amplitude of fluctuations of the cosmic microwave background radiation and the rms variation of the temperature field smoothed with a 10° FWHM Gaussian, and find a reasonable agreement with the new COBE results.

Biasing should be important in our model. We discuss one of the possible biasing schemes very briefly, and further discussions on this will be in our future work.

Subject headings: large scale structure -- statistics -- cosmic background radiation
— cosmology — galaxies: clustering

1. INTRODUCTION

The formation of large scale structure is an important subject in cosmology. Among the various models attempting to explain large scale structures, the cold dark matter (CDM) model is one of the most successful models. It has three basic assumptions:

(1). The initial perturbations arise from quantum fluctuations during inflation and are statistically Gaussian random fields with a scale-invariant power spectrum (Bardeen, Steinhardt, & Turner 1983).

(2). The perturbations evolve in a universe with approximately 90% of the non-relativistic matter in the form of weakly interacting particles with negligible velocity dispersion.

(3). The cosmological density parameter for the present universe is dominated by the non-relativistic matter and is $\Omega = 1$, as expected from inflation.

For a Gaussian field, all statistical properties can be described by two-point correlation function $\xi(\vec{x})$, or the power spectrum which as defined by Salopek *et al.* (1989) is

$$P(k) = \frac{k^3}{2\pi^2} \int d^3x \exp(-i\vec{k} \cdot \vec{x}) \xi(\vec{x}), \quad (1.1)$$

the power per unit logarithmic interval in k .

The initial perturbation field is best expressed in terms of the gauge-invariant potential ζ

$$\zeta = \delta\rho/(\rho + p) + 3\varphi$$

where p is the pressure, and φ is the curvature perturbation amplitude. (Bardeen *et al.* 1986 hereafter BBKS). Scale-invariance means that the primordial ζ , evaluated when all astrophysically relevant scales have wavelength larger than the Hubble radius, has a power spectrum with roughly the same amplitude on each logarithmic interval in wave number k . $P(k) \approx \text{constant}$.

The standard CDM model works quite well at small scales if the Hubble constant $H_0 \approx 50 \text{ km sec}^{-1} \text{ Mpc}^{-1}$ (Davis *et al.* 1985), but seems less successful on scales larger than $10 \text{ Mpc } h^{-1}$ (where h is the Hubble constant in units of

100 km sec⁻¹ Mpc⁻¹). (White *et al.* 1987; Efstathiou, Sutherland, & Maddox 1990). Two recent statistically well-founded pieces of evidence are the galaxy-galaxy angular correlations from the APM survey, and the QDOT redshift survey (Maddox *et al.* 1990, Saunders *et al.* 1991, Efstathiou *et al.* 1990). This means that we need some extra power on those scales comparing to the prediction of the standard CDM model.

There are models which have completely different mechanisms from CDM to explain the large scale structure (Hogan 1991; Hill, Schramm, & Fry 1989; Ostriker & Cowie 1981; Turok 1989). Still in the framework of CDM, there are some ways to modify the standard model. Efstathiou *et al.* (1990) consider a non-zero cosmological constant model. Bond *et al.* (1991) consider the effect of an unstable 17keV massive neutrino, and propose a mechanism which increases the neutrino species' parameter θ . The basic idea in these two models is to get larger characteristic scale $\lambda_{eq} \approx 2\pi \times 3.7\theta^{1/2}/(\Omega h) \text{ (Mpc}^{-1}\text{)}$, which is the comoving horizon scale at matter and radiation equality (BBKS). The other way is to drop out the assumption of a scale-invariant power spectrum (Salopek *et al.* 1989, Kofman *et al.* 1989), but still assume a Gaussian field. There are more parameters to be tuned in this kind of models to fit the observational results. There is another way which gives up the Gaussian assumption. Peebles (1983) is the first to point out that it is possible to get some non-Gaussian fluctuations by non-linear interactions, and they may have some interesting effects on large scale structures. A lot of work has been done on generating non-Gaussian fields (Wise 1988; Salopek 1991; Kofman *et al.* 1989; Ostriker & Cowie 1981; Turok 1989; La & Steinhardt 1989). Recently Matarrese *et al.* (1990, 1991a, 1991b, hereafter MMLM), and Weinberg and Cole (1992, Preprint, hereafter WC) have done simulations on various kinds of non-Gaussian models. They found that the skewness of the density perturbation field is an important parameter to classify different kinds of non-Gaussian models. The general conclusion is that skew-positive models produce more high density clumpy structures than Gaussian models which have zero skewness, while skew-negative models give more void structures. (MMLM; WC). MMLM favor skew-negative models arguing they have bubble-like structures which agree with the observations (Geller & Huchra 1989). But the quantitative

calculations show that skew-negative models give too many voids comparing to the observations while skew-positive models agree with observations on topological properties better (WC). WC also pointed out the disadvantages of skew-positive models, e.g., they need very strong biasing to create empty voids. In MMLM papers, they consider a scale-invariant power spectrum, and in WC, a power law spectrum is considered.

In this paper, we present a model which is basically a skew-positive model. Our model thus shares some general properties of the latter. For example, there are more high density clumpy structures, and the formation of quasars and galaxies occurs earlier than that predicted by the standard CDM. But we have two important differences from MMLM and WC. One is that in our model, we assume the initial perturbation field is a combination of a Gaussian and a non-Gaussian field rather than a purely non-Gaussian field. Motivated by a particular class of double inflation models (Salopek, Bond, & Bardeen 1989, Bardeen 1992), the gauge-invariant potential ζ is assumed to be a combination of a Gaussian and the square of a Gaussian field

$$\zeta = \phi_1 + \phi_2^2 \quad (1.2)$$

where ϕ_1 and ϕ_2 are two independent Gaussian fields. The other difference is that we use a non-scale invariant power spectrum for ϕ_2 , which is

$$P_{\phi_2}(k) = B^{1/2} \exp\{-2a[\ln(k/k_e)]^2\} \quad (1.3)$$

where a , and k_e are parameters which represent the width and the location of the peak of the power spectrum, and $B^{1/2}$ is the amplitude of the non-Gaussian field. For ϕ_1 , we assume the Harrison-Zel'dovich spectrum: $P_{\phi_1}(k) = A$, where A is a constant.

The statistical properties change with scale: On small scales, Gaussian statistics are dominant, but near characteristic scale related to k_e the non-Gaussian contributions are important.

We use the same normalization condition as WC to normalize the total amplitude of density perturbation field to 1 when averaged over a sphere of 8 Mpc^{-1} . (WC; Davis & Peebles 1983). Then the ratio of the Gaussian and the non-Gaussian fields A/B is another adjustable parameter.

This paper mainly concentrates on using the linear theory to compare our model with the standard CDM by discussing statistical properties and two-point correlation functions which are directly related to the power spectrum. We assume $h = 0.5$ and $\Omega = 1$, as conventional for CDM, though direct observations of h are increasingly convergent toward a value of 0.8.

The paper is organized as following: In Sec.2, we discuss the distribution function of the total density field, and show how it changes with different smoothing scales, different characteristic scales as characterized by k_c , and different ratios A/B . In Sec.3, by comparing the theoretical calculations with the results of the APM (the Automatic Plate Measuring, Maddox *et al.* 1990) survey, we set an acceptable parameter regime for a , k_c , and A/B . In Sec.4, the properties of velocity fields are discussed. In Sec.5, we consider the fluctuations on the cosmic microwave background radiation. The quadrupole term and the rms variation smoothed with $\theta_{FWHM} = 10^\circ$ Gaussian are calculated. They are compared to the new Cosmic Background Explorer (COBE) results, and we find that our model agree with the COBE results reasonably. In Sec.6, Several realizations are presented to compare the visual impressions with the standard CDM model. In Sec.7, we make some discussions and comments on our work.

2. THE DISTRIBUTION FUNCTION OF THE DENSITY FIELD

In this section, we consider the distribution function of the density perturbation field which is a combination of a Gaussian and a non-Gaussian field.

The power spectrum of linear density perturbations is related to that of the initial gauge-invariant potential ζ through a transfer function $T(k)$

$$P_{\delta\rho/\rho}(k) = T^2(k)P_{\zeta}(k) \quad (2.1)$$

For CDM models, detailed calculations give (Bond & Efstathiou 1984)

$$T(k) \propto k^2 / \{1 + [ak + (bk)^{3/2} + (ck)^2]^\nu\}^{1/\nu} \quad (2.2)$$

where $a = 6.4(\Omega h^2)^{-1}$ Mpc, $b = 3.0(\Omega h^2)^{-1}$ Mpc, $c = 1.7(\Omega h^2)^{-1}$ Mpc and $\nu = 1.13$. For k much smaller than $k_{eq} = 2\pi/\lambda_{eq}$, $T(k) \propto k^2$. Transforming to the spatial coordinates gives $\delta\rho/\rho \propto -\nabla^2\zeta$. We use this approximation to discuss the qualitative statistical properties of the density perturbation fields, but for the normalization based on $\langle (\delta\rho/\rho)^2 \rangle$ averaged over a sphere of $8 \text{ Mpc} h^{-1}$, the full transfer function (2.2) is used.

With $\zeta = o_1 + (o_2)^2$, we have

$$\delta\rho/\rho \propto -\nabla^2 o_1 - \nabla^2 o_2^2 \quad (2.3)$$

where o_1 and o_2 are two independent Gaussian fields. Their power spectra are given in the last section. a , k_e and A/B are parameters which are adjustable in our model. We normalize $\delta\rho/\rho$ to 1 when averaged over a sphere with a radius $8 \text{ Mpc} h^{-1}$. There is no any bias in this paper.

Write $\delta\rho/\rho$ as

$$\delta\rho/\rho = \sigma_{0G}\nu_G + 2\sigma_{02}\sigma_{22}y \quad (2.4)$$

where ν_G is the dimensionless Gaussian part, and y is the dimensionless non-Gaussian part. σ_{0G} , σ_{02} and σ_{22} are defined as

$$\sigma_{0G}^2 = \int d\ln k P_G(k) \quad (2.5)$$

$$\sigma_{02}^2 = \int d\ln k P_{\delta_2}(k) \quad (2.6)$$

$$\sigma_{\gamma_2}^2 = \int d \ln k \quad k^4 P_{\phi_2}(k) \quad (2.7)$$

where P_G stands for the power spectrum of the Gaussian density perturbations.

We need to consider the joint distribution function for ν_G and y . Since they are independent of each other, the joint distribution is:

$$P(\nu_G, y) d\nu_G dy = P(\nu_G) P(y) d\nu_G dy \quad (2.8)$$

where $P(\nu_G)$ is simply a Gaussian distribution

$$P(\nu_G) = \frac{1}{(2\pi)^{1/2}} \exp(-\nu_G^2/2), \quad (2.9)$$

and $P(y)$ is obtained from the joint distribution function of ϕ_2 , $\nabla \phi_2$, and $\nabla^2 \phi_2$

$$P(y) = \frac{4\pi}{(2\pi)^{5/2}(1-\gamma_2^2)^{1/2}} \int dy_0 dv v^2 / |y_0| \exp \left[-\frac{y_0^2}{2(1-\gamma_2^2)} - \frac{(y + \gamma_2 v^2/3)^2}{2y_0^2(1-\gamma_2^2)} + \frac{\gamma_2(y + \gamma_2 v^2/3)}{1-\gamma_2^2} - (1/2)v^2 \right] \quad (2.10)$$

where $\gamma_2 = \sigma_{12}^2/(\sigma_{02}\sigma_{22})$, and $\sigma_{12}^2 = \int d \ln k \quad k^2 P_{\phi_2}(k)$.

Let ν be the total dimensionless density perturbation, i.e., $\nu = \sigma_0^{-1} \delta\rho/\rho$, where σ_0^2 is $< (\delta\rho/\rho)^2 >$. Then

$$\nu_G = (\sigma_0 \nu - 2\sigma_{02}\sigma_{22}y)/\sigma_{0G} \quad (2.11)$$

Replace ν_G in eq.(2.9) by eq.(2.11), and integrate over y to get the distribution function for ν ,

$$P(\nu) d\nu = \left\{ \int \exp \left[-\frac{(\sigma_0 \nu - 2\sigma_{02}\sigma_{22}y)^2}{2\sigma_{0G}^2} \right] P(y) dy \right\} d\nu \quad (2.12)$$

Define $c_1 = \sigma_0/\sigma_{0G}$ and $c_2 = 2\sigma_{02}\sigma_{22}/\sigma_{0G}$. When $c_2 \rightarrow 0$, i.e., the contribution of the non-Gaussian part goes to zero, (2.12) becomes a Gaussian distribution. On the other hand when $c_1 \rightarrow \infty$, (2.12) tends to have the form of (2.10) with $y = (\delta\rho/\rho)/(2\sigma_{02}\sigma_{22})$.

The behavior of the distribution (2.12) depends on a , k_c , and A/B . If we consider a smoothed field, it also depends on the smoothing scale R_f . We use a Gaussian smoothing function.

In Fig.1, we plot the purely non-Gaussian distribution $P(y)$ with different γ_2 . They are skew-positive, and have higher probabilities at high σ -level perturbations than that of the Gaussian distribution. The smaller the γ_2 is, the sharper peaked at zero the distribution is, and it gets broader when γ_2 goes to 1.

Fig.2 shows the behavior of $P(\nu)$ with different k_z , A/B , and R_f . When the smoothing scale R_f is small, the Gaussian field is dominant, and the distribution is nearly symmetric. For $R_f = 10 \text{ Mpc h}^{-1}$, and $R_f = 20 \text{ Mpc h}^{-1}$, the distribution function is no longer symmetric, and has a long tail at high positive values, which is the general property of a skew-positive model related to the earlier formation of quasars and galaxies than that of the standard CDM. The larger the A/B is, the more Gaussian the distribution tends to be. Note that for $R_f = 2 \text{ Mpc h}^{-1}$, our approximation (2.3) is not valid. But the tendency described above is true.

Comparing with observations, the standard CDM model works quite well on small scales $r < 10 \text{ Mpc h}^{-1}$, but on scales larger than 10 Mpc h^{-1} , more power than predicted by the standard CDM model is needed to explain the strong correlations on those scales. (Davis & Peebles 1983; Maddox *et al* 1990) One of the advantages of our model is that the statistical properties of perturbation fields depend on scale, and on small scales, Gaussian fields are dominant, thus keeps the advantages of Gaussian fields (WC). By choosing parameters a , k_z and A/B properly, we can get large correlations on scales larger than 10 Mpc h^{-1} . In next section, we calculate the angular correlation function of density perturbations, and compare it with the APM survey to get an acceptable parameter regime.

3. THE LIMITATIONS ON PARAMETERS

Using the APM technique, Maddox *et al* (1990) got more reliable results than those of the Lick survey (Groth & Peebles 1977) on the two dimensional correlation function $w(\theta)$ shown in Fig.3. The dotted line is the result of the standard CDM model with $\Omega = 1$ and $h = 0.5$. It is clearly seen that there is insufficient power on large scales.

The angular correlation function is related to the spatial two-point correlation function by an integral equation (Peebles 1980). Here we use $\Omega = 1$, which corresponds to $q_0 = 0.5$. The same selection function as in Maddox *et al* (1990) is used in our calculations.

In our model, the power spectrum for the total density field is

$$P(k) = T^2(k)[P_{\phi_1}(k) + P_{\phi_2}(k)] \quad (3.1)$$

where $T(k)$ is the transfer function given by (2.2).

$P_{\phi_1}(k) = A$ is used. $P_{\phi_2}(k)$ is related to the power spectrum of ϕ_2 by the following equation:

$$P_{\phi_2}(k) = \frac{k^3}{2\pi^2} \int \frac{d^3\vec{k}_1}{(2\pi)^3} |\phi_2(\vec{k}_1)|^2 |\phi_2(\vec{k} - \vec{k}_1)|^2 \quad (3.2)$$

We calculate $w(\theta)$ for different a , k_c , and A/B . It is known that from quantum fluctuation theories (Salopek *et al.* 1989), we can not get a very sharp peak in the power spectrum for ϕ_2 . Thus we restrict a to be less than 1. Calculations show that:

(1). For $k_{eq}/k_c < 5$, the curve of $w(\theta)$ is always too steep to match the APM results, which means that the characteristic scale related to k_c is too small to account for the extra power on large scales needed by the observational results.

(2). For $k_{eq}/k_c > 14$, the curve is always too flat, i.e., the characteristic scale is too large.

(3). In the range $5 \leq k_{eq}/k_c \leq 14$, we can find proper a and A/B to fit the APM results. In Table.1, we list the proper range for a , k_c , and A/B . For

example, when $k_{eq}/k_c = 10$, $a = 0.5$, then A/B should be in the range $0.0165 \leq A/B \leq 0.0302$. The two dimensional correlation functions $w(\theta)$ for $k_{eq}/k_c = 10$, $a = 0.5$ are shown in Fig.4 along with the results calculated using the standard CDM model.

Bardeen, Bond & Efstathiou (1987) discussed what kind of a power spectrum of the peculiar velocity field is needed to match the observations. They concluded that it is better to have a spectrum with a wider peak shifted to a larger scale than the predictions of the standard CDM model. In Fig.5, we plot the power spectra of the total velocity field of our model with a group of parameters and that of the standard CDM model. We do get a wider peak which shifts to a larger scale. Calculations on 3-dim. rms velocity for different smoothing scales are listed in Table.2. We see, the rms velocities of our model are larger than those predicted by the standard CDM model. On small scales they seem to be too large. But here we do not consider any biasing. For a skew-positive model like ours, the biasing should be stronger than that of the standard CDM model (WC 1991) because the density distribution is less homogenous, and the probabilities for large positive density perturbations are higher than those of the Gaussian case. The strong biasing will reduce the rms peculiar velocities. At the same time it will also reduce the quadrupole perturbation a_2 of the microwave background radiation, and make a_2 be too small. Decreasing a , i.e., flattening the peak of the power spectrum will increase a_2 , but the power on small scales will be increased as well since our power spectrum of non-Gaussian field is logarithmically symmetric respect to k_c . One possible way to get rid of this problem is to make the peak asymmetric, and put more power on large scales. An appropriately designed effective scalar field potential in inflation can accomplish this.

4. STATISTICAL ANALYSIS ON THE PECULIAR VELOCITY FIELD

In this section, we discuss the correlations and the distribution function of the peculiar velocity field using the linear theory, and compare them with those of the standard CDM model.

First, we consider the correlation of 3-dim. velocity $\vec{v}(\vec{x})$:

$$\xi_{vb}(r) = \langle \vec{v}(\vec{x}) \cdot \vec{v}(\vec{x} - \vec{r}) \rangle \quad (4.1)$$

This quantity gives us information on the bulk velocity field. A positive $\xi_{vb}(r)$ means that on average particles separated by r move coherently toward the same direction.

Using the linear theory, $\xi_{vb}(r)$ is related to the power spectrum of the initial perturbation field by

$$\xi_{vb}(r) = \int d \ln k \quad T^2(k)/k^2 P_\zeta(k) \sin(kr)/(kr) \quad (4.2)$$

where $P_\zeta(k)$ stands for the initial power spectrum of the gauge-invariant potential ζ . $T(k)$ is the transfer function of (2.2).

We calculate $\xi_{vb}(r)$ for several groups of parameters as shown in Fig.6. On comoving scales less than the characteristic scale, the correlations are larger than those of the standard CDM model. This is consistent with the physical picture. In our model, there are more dense regions around the characteristic scale, and on scales less than that, more pairs of particles move toward the same dense clump, thus we get larger ξ_{vb} . For points whose separation is larger than the typical scale of clumps, they have higher probabilities to move in the way either falling to the same clump from opposite directions or falling to different clumps. Both of them tend to reduce ξ_{vb} .

Now we consider the correlation of $\vec{v}(\vec{x}) \cdot \hat{r}$:

$$\xi_{vi}(r) = - \langle (\vec{v}(\vec{x}) \cdot \hat{r}) (\vec{v}(\vec{x} - \vec{r}) \cdot \hat{r}) \rangle \quad (4.3)$$

$\xi_{vi}(r)$ represents the property of the infall velocity. The minus sign in (4.3) means that we define the correlation to be positive if two points move toward each other.

$\xi_{vi}(r)$ is related to the power spectrum by:

$$\xi_{vi}(r) = \int d \ln k \quad T^2(k)/k^2 P_\zeta(k) \left[\frac{\sin(kr)}{(kr)} + \frac{2 \cos(kr)}{(kr)^2} - \frac{2 \sin(kr)}{(kr)^3} \right] \quad (4.4)$$

Fig.7 shows $\xi_{vi}(r)$. On certain scales, our model gives larger correlations, and the coherent scale is larger than that of the standard CDM model. This is again consistent with that our model is skew-positive and has more clumpy structures.

By analysing the correlations of the streaming velocity and the infall velocity, it shows that our model can have larger coherent motion and the coherent scales than those predicted by the standard CDM model.

Furthermore, we calculate the rms velocity of $\vec{v}(\vec{x}, R)$, the average velocity over certain scales R ,

$$\vec{v}(\vec{x}, R) \equiv \frac{1}{V} \int_{|\vec{r}| \leq R} d^3 \vec{r} \quad \vec{v}(\vec{x} + \vec{r}) \quad (4.5)$$

where V is the volume of the sphere with a radius R .

The rms value reflects how strong the motion is coherent on scales less than R . And the rms of $\vec{v}(\vec{x}, R)$ can also be calculated from the power spectrum:

$$\langle \vec{v}(\vec{x}, R) \cdot \vec{v}(\vec{x}, R) \rangle = \int d \ln k \quad T^2(k)/k^2 P_\zeta(k) W_1^2(kR) \quad (4.6)$$

where

$$W_1(kR) = \frac{3[-kR \cos(kR) + \sin(kR)]}{(kR)^3}.$$

Fig.8 shows the behaviors of the rms values. We see that for the same scale which is less than the characteristic scale, our model has larger rms velocity than that of the standard CDM model. In another word, we can get larger regions in which the motion is coherent. Some observations (Dressler 1989) show that there is a very large coherent flow, $\sim 600 \text{ kms}^{-1}$, extending at least 30 Mpc h^{-1} . It is very hard to get such large coherent motion over such large scales in the standard CDM model. While in our model, we can have such large coherent motions over $20 - 30 \text{ Mpc h}^{-1}$. This is again because we have dense structures over large scales in our model, which are more like ‘Great Attractors’.

Another quantity we calculate is the rms of $v_r(\vec{x}, R)$. Here $v_r(\vec{x}, R)$ is defined as

$$v_r(\vec{x}, R) \equiv \frac{1}{V} \int_{|\vec{r}| \leq R} d^3 \vec{r} \quad \vec{v}(\vec{x} + \vec{r}) \cdot \hat{r} \quad (4.7)$$

The rms of $v_r(\vec{x}, R)$ is related to the power spectrum by

$$\langle v_r^2(\vec{x}, R) \rangle = \int d \ln k \quad T^2(k)/k^2 P_\zeta(k) W_2^2(kR) \quad (4.8)$$

and

$$W_2(kR) = \frac{3[kR \sin(kR) - 4 \sin^2(kR/2)]}{(kR)^3}.$$

This rms value shows that how strong the gravitation is and what a large region it can affect. This can be seen from the Virial theorem: $v \sim (GM/R)^{1/2}$, where M is the mass in the sphere. Since we have more large and dense structures, we expect to get larger rms of v_r .

The results are plotted in Fig.9. we do see there are larger values than those of the standard CDM model on scales less than the characteristic scale.

The analysis above is mainly related to the power spectrum. To understand the non-Gaussian perperties further, we need to calculate the distribution function of the velocity field.

Write the total velocity as

$$\vec{v}_{total} = \vec{v}_G + \vec{v}_{NG} \quad (4.9)$$

where $\vec{v}_G$ stands for the Gaussian part of the velocity field, and $\vec{v}_{NG}$ is the non-Gaussian part.

Using the linear theory, we have

$$\vec{v}_G = \nabla \phi_1.$$

$$\vec{v}_{NG} = \nabla \phi_2^2.$$

Let us first consider the distribution of the purely non-Gaussian part $\vec{v}_{NG}$. Since ϕ_2 is a Gaussian field, $\nabla_i \phi_2$ ($i = 1, 2, 3$) is also a Gaussian field. Because $\langle \phi_2 \nabla_i \phi_2 \rangle = 0$, the joint probability of ϕ_2 and $\nabla_i \phi_2$ is simply:

$$P(\phi_2, \eta_1, \eta_2, \eta_3) d\phi_2 d\eta_1 d\eta_2 d\eta_3 = \left(\frac{1}{2\pi}\right)^2 \frac{1}{\sigma_{\eta_2}} \left(\frac{3^{1/2}}{\sigma_{\eta_2}}\right)^3 \exp\left(-\frac{\phi_2^2}{2\sigma_{\phi_2}^2}\right) \exp\left[-\frac{3(\eta_1^2 + \eta_2^2 + \eta_3^2)}{2\sigma_{\eta_2}^2}\right] d\phi_2 d\eta_1 d\eta_2 d\eta_3 \quad (4.10)$$

With $\nu_2 = \phi_2/\sigma_{02}$ and $\eta^2 = \eta_1^2 + \eta_2^2 + \eta_3^2$, (4.10) turns out to be

$$P(\nu_2, \eta) d\nu_2 d\eta = \frac{1}{(2\pi)^{1/2}} \left(\frac{54}{\pi}\right)^{1/2} \frac{1}{\sigma_{12}^3} \exp(-\nu_2^2/2) \exp[-3\eta^2/(2\sigma_{12}^2)] \eta^2 d\nu_2 d\eta \quad (4.11)$$

where σ_{02} and σ_{12} are defined as before.

Use $\vec{v}_{NG} = 2\phi_2 \nabla \phi_2$, we can find the distribution function for v_{NG} . It is

$$P(v_{NG}) dv_{NG} = \frac{1}{(2\pi)^{1/2}} \left(\frac{54}{\pi}\right)^{1/2} \frac{1}{\sigma_{12}^3} \frac{v_{NG}^2}{8\sigma_{02}^2 \sigma_{12}^2 \nu_2^2} \exp(-\nu_2^2/2) \exp[-3v_{NG}^2/(8\sigma_{02}^2 \sigma_{12}^2 \nu_2^2)] |\nu_2^{-3}| d\nu_2 dv_{NG} \quad (4.12)$$

With $x = v_{NG}/(2\sigma_{02}\sigma_{12})$, the distribution function for x is

$$P(x) dx = \frac{1}{(2\pi)^{1/2}} \left(\frac{54}{\pi}\right)^{1/2} x^2 \int_{-\infty}^{\infty} \exp(-\nu_2^2/2) \exp[-3x^2/(2\nu_2^2)] |\nu_2^{-3}| d\nu_2 dx \quad (4.13)$$

We see there is no parameter in it, and the distribution for x is unique and independent of a and k_r . This is different from the density field which distribution depends on the parameter α_2 .

In Fig.9, we show the behaviors of $P(x)$ along with the distribution function for the Gaussian velocity field. We see they are quite different. The non-Gaussian distribution has non-zero probability at $x = 0$, and a long tail at high x values.

Now we consider the distribution for the total velocity. Since $\vec{v}_G$ and $\vec{v}_{NG}$ are independent of each other, the joint distribution is the product of the distribution functions for v_{Gi} and for v_{NGi} ($i = 1, 2, 3$):

$$\begin{aligned} P(v_{G1}, v_{G2}, v_{G3}, v_{NG1}, v_{NG2}, v_{NG3}) dv_{G1} dv_{G2} dv_{G3} dv_{NG1} dv_{NG2} dv_{NG3} \\ = \frac{1}{(2\pi)^{3/2}} \frac{27}{8\sigma_{1G}^3 \sigma_{12}^3 \sigma_{12}^3} \exp\left[-\frac{3(v_{G1}^2 + v_{G2}^2 + v_{G3}^2)}{2\sigma_{1G}^2}\right] \\ \int_{-\infty}^{\infty} \exp(-\nu_2^2/2) \exp\left[-\frac{3(v_{NG1}^2 + v_{NG2}^2 + v_{NG3}^2)}{8\sigma_{02}^2 \sigma_{12}^2 \nu_2^2}\right] |\nu_2^{-3}| d\nu_2 \\ dv_{G1} dv_{G2} dv_{G3} dv_{NG1} dv_{NG2} dv_{NG3} \end{aligned} \quad (4.14)$$

where σ_{1G} is the rms of v_G , $\sigma_{1G} = \langle v_G^2 \rangle^{1/2}$.

Use $v_i = v_{Gi} + v_{NGi}$ ($i = 1, 2, 3$) as variables, we get

$$\begin{aligned} P(v_1, v_2, v_3, v_{NG1}, v_{NG2}, v_{NG3}) dv_1 dv_2 dv_3 dv_{NG1} dv_{NG2} dv_{NG3} \\ = \frac{1}{(2\pi)^{3/2}} \frac{27}{8\sigma_{1G}^3 \sigma_{12}^3 \sigma_{12}^3} \exp\left[-\frac{3(v^2 + v_{NG}^2 - 2\vec{v} \cdot \vec{v}_{NG})}{2\sigma_{1G}^2}\right] \\ \int_{-\infty}^{\infty} \exp(-\nu_2^2/2) \exp\left[-\frac{3(v_{NG1}^2 + v_{NG2}^2 + v_{NG3}^2)}{8\sigma_{02}^2 \sigma_{12}^2 \nu_2^2}\right] |\nu_2^{-3}| d\nu_2 \\ dv_1 dv_2 dv_3 dv_{NG1} dv_{NG2} dv_{NG3} \end{aligned} \quad (4.15)$$

Define $b_1 = \sigma_{02}\sigma_{12}/\sigma_{1G}$ and $b_2 = \sigma_{11}/\sigma_{1G}$, where σ_{11} is defined as $\langle v^2 \rangle^{1/2}$. Integrate over all other variables except $v'_1 = v_1/\sigma_{11}$, we get (hereafter we still use v for v')

$$p(v)dv = \frac{54}{(2\pi)^{3/2}} \frac{b_2^2}{(6b_1)} v \exp(-3b_2^2 v^2/2) f(v) dv \quad (4.16)$$

where

$$f(v) = \int_{-\infty}^{\infty} d\nu \int_0^{\infty} dx \quad x |\nu_2^{-3}| \exp(-\nu_2^2/2) \exp[-3x^2/(2\nu_2^2)] \exp(-6b_1^2 x^2) [\exp(6b_1 b_2 x v) - \exp(-6b_1 b_2 x v)] \quad (4.17)$$

The distribution $P(v)$ depends on b_1 and b_2 which are functions of a , k_c and A/B . As $b_1 \rightarrow 0$, i.e., the contribution from the non-Gaussian field goes to zero, the distribution (4.17) tends to the distribution for a Gaussian velocity field which is

$$P(v)dv = \left(\frac{54}{\pi}\right)^{1/2} v^2 \exp(-3v^2/2) dv \quad (4.18).$$

When $b_1 \rightarrow \infty$, i.e., the Gaussian part goes to zero, (4.17) goes to (4.13).

Fig.10 shows $P(v)$ for several groups of parameters. We see that for the smoothing scale equal to 2 Mpc h^{-1} , the distribution is very near the distribution for a purely Gaussian velocity field, as it is for larger smoothing scales when A/B is larger. When the smoothing scale is near k_c^{-1} and A/B is smaller, the distribution has a long tail in high velocities. Thus as the density field, the statistical properties of the velocity field changes with scales.

Thus there are two effects make it easier to get high velocities on certain scales than using the standard CDM model. One is from our particular choice of the power spectrum for the non-Gaussian field, and this gives higher rms velocities. The other effect is purely statistical. We get a long tail in high velocities which means that we have higher probabilities to get high velocities than the standard CDM model.

One thing needs to be mentioned. Our analysis in this paper is for the mass field, and we do not take into account any biasing effect. Since our model, like all other skew-positive models, has larger probabilities for high positive density perturbations, the biasing effect should be important. In the last section, we put forward a possible biasing scheme for our model, more detailed work will be done in our future work.

5. FLUCTUATIONS OF COSMIC MICROWAVE BACKGROUND RADIATION

Fluctuations of the cosmic microwave background radiation (CMBR) provide us the most restrictive constraints to theoretical models. Recent reports from the COBE show that the rms quadrupole amplitude of temperature fluctuations is $13 \pm 4 \mu\text{K}$, and the rms sky variation, smoothed with a 10° FWHM Gaussian, is $31 \pm 7 \mu\text{K}$.

In this section, we calculate the rms quadrupole amplitude and the rms variation of the temperature field using our combination model, and it shows that our model can agree with the COBE results quite well.

The multipole amplitude is defined as (Bond & Efstathiou 1987)

$$\Delta T_l^2 = \frac{1}{4\pi} \sum_{m=-l}^{m=+l} |a_{lm}|^2 \quad (5.1)$$

and $a_l^2 \equiv \langle |a_{lm}|^2 \rangle$ is related to the power spectrum of the gauge-invariant initial potential ζ as (Peebles 1982)

$$a_l^2 = \pi H^4 \int_0^\infty \frac{dk}{k} P_\zeta(k) j_l^2(k\tau_0) \quad (5.2)$$

where $\tau \equiv \int dt/a$ is the proper time, and the subscript zero stands for the present value.

The two point correlation function of the smoothed temperature field is (Wilson & Silk 1981)

$$C(\theta, \theta_s) = \int_0^\infty C(\phi) R(\phi, \theta, \theta_s) \phi d\phi \quad (5.3)$$

where $\theta_s = 0.425\theta_{FWHM}$ is the Gaussian smoothing angle,

$$R(\phi, \theta, \theta_s) = \frac{1}{2\theta_s^2} \exp\left(-\frac{\theta^2 + \phi^2}{4\theta_s^2}\right) I_0\left(\frac{\theta\phi}{2\theta_s^2}\right),$$

and $C(\phi)$ is the unsmoothed two point correlation function. After subtracting the mean and dipole term, it is (Górski 1991)

$$C(\phi) = \frac{1}{4} H^4 \int_0^\infty \frac{dk}{k} P_\zeta(k) [j_0^2(k\tau_0\phi) - j_0^2(k\tau_0) - 3j_1^2(k\tau_0)(1 - \frac{\phi^2}{2})]. \quad (5.4)$$

The rms variation $\sigma(\theta_s)$ is defined as (Bond & Efstathiou 1987)

$$\sigma(\theta_s) \equiv [C(0, \theta_s)]^{\frac{1}{2}} \quad (5.5)$$

Integrating over ϕ , we get the rms variation is

$$\sigma^2(\theta_s) = \frac{1}{4} H^4 \int_0^\infty \frac{dk}{k} P_\zeta(k) [F(1; \frac{3}{2}; -(k\tau_0\theta_s)^2) - j_0^2(k\tau_0) - 3j_1^2(k\tau_0)(1-2\theta_s^2)]. \quad (5.6)$$

where $F(\alpha; \gamma; z)$ is the confluent hypergeometric function, and

$$F(1; \frac{3}{2}; -(k\tau_0\theta_s)^2) = \int_0^1 \exp[(k\tau_0\theta_s)^2(t^2 - 1)] dt.$$

The results are listed in Table.3. For the parameters we use, the main contributions to $(\Delta T/T)_2$ and $\sigma(\theta_{FWHM} = 10^\circ)$ come from the Gaussian field. For $(\Delta T/T)_2^2$, the non-Gaussian contribution is only $\sim 1.7\%$ of the Gaussian contribution. For $\sigma^2(\theta_{FWHM} = 10^\circ)$, it is $\sim 14.7\%$.

It is known that the quadrupole fluctuations are the results of very large scale perturbations, and large θ_{FWHM} correspond to large scales, too. We use a 'peak' power spectrum for the non-Gaussian field, thus it is important only on certain scales which depend on k_c and a . For the k_c and a we choose here, they are smaller than those scales which contribute to $(\Delta T/T)_2$ and $\sigma(\theta_{FWHM} = 10^\circ)$. Reduce a and k_c , the contributions will be larger. The contribution to σ from non-Gaussian field depends on θ_{FWHM} . Fig.12 shows the contributions to σ^2 from the Gaussian field and from the non-Gaussian field relative to θ_{FWHM} . When $\theta_{FWHM} > 3^\circ$, the Gaussian part is larger. While for $\theta_{FWHM} < 3^\circ$, the non-Gaussian part is dominant.

Compare with the COBE results on the quadrupole term and the rms variation with $\theta_{FWHM} = 10^\circ$ (Smoot *et al.* 1992), we find that our model can fit both the angular correlation function of the density perturbation field and the CMBR fluctuations reasonably well.

6. MORPHOLOGY

In this section, we use the adhesion model (Weinberg & Gunn 1990; Shandarin & Zel'dovich 1989; Shandarin 1988; Kofman, Pogosyan & Shandarin 1990) to construct several realizations of the velocity fields both for the standard CDM model and for our model. Because of the limitation in the dynamic range for doing realizations, we use $a = 0.5$, and $k_z = k_{z,1}/5$ for the non-Gaussian field.

Fig. 13(a) and 13(b) are slices of the velocity fields for the standard CDM model and our combinational model using the linear theory. It can be seen that in our model, there are some large clumps, while in the standard CDM model, it is more random and homogeneous on such large scales. The large coherent structures in our model make the correlations on large scales larger than those in the standard CDM model. Fig.14(a) and 14(b) correspond to Fig.13(a) and 13(b) respectively, but some non-linear effects are taken into account by using the adhesion model with the 'finite ν ' method (Weinberg *et al.* 1990; Shandarin 1988). We see that the patterns are almost the same as linear realizations (Fig.11), but some small scale structures are smoothed out. (Weinberg *et al* 1990)

We have two different ways of analysing statistical properties including non-linear evolution effects in the framework of the adhesion model (Shandarin 1988). One is the simulation method with finite ν (Nusser & Dekel 1990; Weinberg *et al* 1990), the other is the 'skeleton method' in the limit of ν going to zero (Kofman *et al* 1990). Future work will apply these adhesion model techniques more extensively.

7. DISCUSSION

We discuss a model in which the initial perturbation field ζ is a combination of a Gaussian and a square of a Gaussian field with some particular power spectra. There are three parameters in this model being adjusted: a , k_c and A/B . By comparing with the APM survey, we get certain acceptable ranges for them. Then we compare our model with the standard CDM model.

Our model is basically a skew-positive model, and there are more large scale clumpy structures and fewer voids than the standard CDM model (see Fig.13(b) and 14(b)) if we assume that galaxy traces mass. This seems to be inconsistent with observations. Thus some biasing mechanisms are needed in our model. Since it is a skew-positive, we expect the existence of a strong biasing. WC pointed out that with certain biasing, the skew-positive models can agree with the observations best on topological properties whereas both Gaussian and skew-negative models have too many voids. For our model, we use 'peak' power spectrum for the non-Gaussian field, and the characteristic scale is much larger than the typical galaxy scale. Thus one possible biasing scheme is still to follow the basic idea of 'peak' biasing (BBKS 1986; Kaiser 1988), but assume that the non-Gaussian field only changes the large scale 'background' field, and has no contributions to the number of peaks with galaxy scales. High density regions over certain large scales enhances the number of peaks of the Gaussian part, which are higher than a certain threshold (BBKS 1986). We will consider this farther in subsequent papers.

Our theoretical motivation is based on a specific double inflation model, but our model is an interesting example of certain kind of models whose statistical properties change with scale. On small scales, it is basically Gaussian dominant, thus the advantages of Gaussian models on small scale still exist. By adjusting parameters properly, we can fit both the APM two-dimensional correlations and the COBE results quite well. Analysis of the peculiar velocity fields shows that on scales less than the characteristic scale related to k_c , the rms bulk velocities as well as the correlations of the bulk velocity are larger than those of the standard CDM model. Further more the non-Gaussian field have a long tail at high positive values

for the velocity distribution. These two effects make it easier to get large bulk velocities than the standard CDM model. On scales larger than the characteristic scale, the rms bulk velocities become less than those predicted by the standard CDM. This is because we have dense and large clumps in our model, and on scales less than the typical scale of clumps, matter mainly moves toward the same clump, and gives larger bulk velocities. While on scales larger than the typical scale, matter moves mainly falling either into the same clump from opposite sides or into different clumps. This makes the bulk velocities smaller. Our model seems to be more attractive if ‘Great Attractors’ really exist in the universe. We also analyse the average infall velocity over a sphere with certain radius R . The peak of the rms velocity distribution with R is higher and broader than the standard CDM model, and the peak shifts to larger length scales depending on the characteristic scale we choose for the non-Gaussian field. This again shows that our model is more consistent with a ‘Great Attractor’ model than standard CDM.

The calculations on the fluctuations of the cosmic microwave background radiation show that with $\Omega = 1$, and $h = 0.5$, our model can agree with the results of both the APM survey and the COBE.

Couchman *et al.* (1992) made some simulations with the standard CDM model trying to trace galaxies when they cluster, and they claimed that there is a significant bias in the velocities of galaxies relative to the dark matter. Thus they suggest that by considering the anti-bias, the standard CDM model might be consistent with the observations of large bulk peculiar velocity. We do not consider this effect. What we do here is to use the same normalization condition ($\sigma_8 = 1$) for both the standard CDM model and for our model to compare them and look for the differences between these two models.

The ability to fit observational data on a range of scales cannot be considered a success of the model, in that the model has plenty of adjustable parameters. However it is interesting to explore some of the ways non-standard statistics can affect comparison with observations.

While this work is mainly based on linear perturbation theory, we plan to use the adhesion model to gain insight into the non-linear evolution of large scale

structure in these models.

Acknowledgements

This research was supported in part by the Department of Energy under Grant No. DE-FG06 91ER40614.

REFERENCES

- Bardeen, J. M. 1988, in The Second Guo Shou-jing Summer School on Particle Physics and Cosmology, ed. A. Zee (Gordon and Breach, London, 1988)
- Bardeen, J. M., Bond, J. R., Kaiser, N., & Szalay A. S. 1986. ApJ, 304, 15 (BBKS)
- Bardeen, J. M., Bond, J. R., & Efstathiou, G. 1987, ApJ, 321, 28
- Bardeen, J. M., Steinhardt, P. J., & Turner, M. S. 1983, Phys. Rev. D., 28, 679
- Bardeen, J. M. 1992, in preparation
- Bond, J. R. 1991, Phys. Lett. B., 265, 245
- Bond, J. R., & Efstathiou, G. 1984, ApJ, 285, L45
- Bond, J. R., & Efstathiou, G. 1987, MNRAS, 226, 655
- Couchman, H. M. P., & Carlberg, R. G. 1992, ApJ, 389, 453
- Davis, M., & Peebles, P. J. E. 1983, ApJ, 267, 465
- Davis, M., Efstathiou, G., Frenk, C. S., & White, S. D. M. 1985, ApJ, 292, 371
- Dressler, A. 1989, in Large Scale Structures and Peculiar Motions in The Universe, ed. David W. Latham and L. A. Nicolaci da Costa. (Astronomical Society of the Pacific), p19
- Efstathiou, G., Sutherland, W. J., & Maddox, S. J. 1990, Nature, 348, 705
- Efstathiou, G., Kaiser, N., Saunders, W., Lawrence, A., Rowan-Robinson, M., Ellis, R. S., & Frenk, C. S. 1990, MNRAS, 247, 10p
- Geller, M. J., & Huchra, J. P. 1989, Science, 246, 897
- Górski, K. M. 1991, ApJ, 370, L5

- Groth, E. J., & Peebles, P. J. E. 1977. *ApJ*, 217, 385
- Hogan, C. 1991, *ApJ*, 369, 300
- Hill, C. T., Schramm, D. N., & Fry, J. N. 1989. *Comments Nucl. Part. Phys.*, 19, 25
- Kaiser, N. 1988. in *Evolution of Large Scale Structures in the Universe*, IAU Symposium 130, ed. J. Audouze & A. Szalay (Dordrecht: Reidel), p43
- Kofman, L., Blumenthal, G. R., Hodges, H., & Primack, J. R. 1989, in *Large Scale Structures and Peculiar Motions in the Universe*, ed. David W. Latham and L. A. Nicolaci da Costa, (Astronomical Society of the Pacific), P339
- Kofman, L., Pogosyan, D., & Shandarin, S. 1990. *MNRAS*, 242, 200
- La, D., & Steinhardt, P. J. 1989. *Phys. Lett.*, B220, 375
- Maddox, S. J., Efstathiou, G., Sutherland, W. J., & Loveday, J. 1990. *MNRAS*, 242, 43p
- Matarrese, S., Lucchin, F., Messina, A., & Moscardini, L. 1991b. *MNRAS*, 253, 35 (MMLM)
- Messina, A., Moscardini, L., Lucchin, F., & Matarrese, S. 1990. *MNRAS*, 245, 244 (MMLM)
- Moscardini, L., Matarrese, S., Lucchin, F., & Messina, A. 1991a. *MNRAS*, 248, 424 (MMLM)
- Nusser, A., & Dekel, A. 1990. *ApJ*, 362, 14
- Ostriker, J. P., & Cowie, L. L. 1981. *ApJ*, 245, L127
- Peebles, P. J. E. 1980. *The Large Scale Structure of the Universe*. (Princeton: Princeton University Press).
- Peebles, P. J. E. 1982. *ApJ*, 263, L1
- Peebles, P. J. E. 1983. *ApJ*, 274, 1
- Salopek, D. S., Bond, J. R., & Bardeen, J. M. 1989. *Phys. Rev.*, D40, 6
- Salopek, D. S. 1991. Preprint. Fermilab-pub-91/161-A

- Saunders, W., Frenk, C. S., Rowan-Robinson, M., Efstathiou, G., Lawrence, A., Kaiser, N., Ellis, R. S., Crawford, J., Xia, X. Y., & Parry, I. 1991, *Nature*, 349, 32
- Shanarin, S. F. 1989, in *Large-Scale Structures and Peculiar Motions in the Universe*, ed. D. W. Latham and L. A. Nicolaci da Costa. (Astronomical Society of the Pacific), p189
- Shandarin, S. F., & Zel'dovich, Ya. B. 1989, *Rev. Mod. Phys.*, 61, 185
- Smoot, G. F., Bennett, C. L., Kogut, A., Wright, E. L., Aymon, J., Boggess, N. W., Cheng, E. S., De Amici, G., Gulkis, S., Hauser, M. G., Hinshaw, G., Lineweaver, C., Loewenstein, K., Jackson, P. D., Janssen, M., Kaita, E., Kelsall, T., Keegstra, P., Lubin, P., Mather, J., Meyer, S. S., Moseley, S. H., Murdock, T., Rokke, L., Silverberg, R. F., Tenorio, L., Weiss, R., & Wilkinson, D. 1992, preprint
- Turok, N. 1989, *Phys. Rev. Lett.*, 63, 262
- Weinberg, D. H., & Cole, S. 1991, preprint.
- Weinberg, D. H., & Gunn, J. E. 1990, *MNRAS*, 247, 260
- White, S. D. M., Frenk, C. S., Davis, M., & Efstathiou, G. 1987, *ApJ*, 313, 505
- Wise, M. B. 1988, in *The Early Universe*. (NATO ASI series. Series C. Mathematical and Physical Science; No. 219)
- Wilson, M. L., & Silk, J. 1981, *ApJ*, 243, 14

FIGURE CAPTIONS

Figure 1: Density distribution functions of the purely non-Gaussian fields. The horizontal axis is $y = \delta\rho/\rho/(2\sigma_{02}\sigma_{22})$. The solid line is for $\gamma = 0.1$, the dotted line is for $\gamma = 0.3$, the dash-dot-dot-dot line is for $\gamma = 0.5$, and the long dashed line is for $\gamma = 0.8$.

Figure 2: Density distribution functions of the total density fields. The horizontal axes are $\nu = \delta\rho/\rho / <(\delta\rho/\rho)^2>^{1/2}$. (a) $a = 0.5$, $k_{eq}/k_c = 10.$, and $BP = 50\%$. (Here BP is defined the same as in the Table.1.) The solid line is for $R_f = 20 \text{ Mpc h}^{-1}$, the dash-dot-dot-dot line is for $R_f = 10 \text{ Mpc h}^{-1}$, and the dotted line is for $R_f = 2 \text{ Mpc h}^{-1}$. (b) $a = 0.5$, $k_{eq}/k_c = 10.$, and $R_f = 10 \text{ Mpc h}^{-1}$. The solid line is for the purely non-Gaussian distribution, the dash-dot-dot-dot line is for $BP = 50\%$, the dotted line is for $BP = 20\%$, the long dashed line is for Gaussian distribution. (c) $R_f = 10 \text{ Mpc h}^{-1}$, $BP = 50\%$. The solid line is for $a = 0.5$, $k_{eq}/k_c = 5.$, and the dotted line is for $a = 0.5$, $k_{eq}/k_c = 10.$.

Figure 3: The dots show estimates of the angular correlation function $w(\theta)$ for galaxies in the APM galaxy survey. The dotted line is the simulation results of standard CDM with $\Omega = 1$ and $h = 0.5$. The thin solid and dashed lines are the results of the linear theory for $\Omega = 0.2$ scale-invariant CDM model with $h = 1$ and 0.75 , respectively. The thick solid line is the N -body simulation results for $\Omega = 0.2$ and $h = 0.9$. (Reproduced from Efstathiou *et al.* (1990) with permission)

Figure 4: The angular correlation function $w(\theta)$. The solid line is for the standard CDM model with $\Omega = 1$ and $h = 0.5$. The dashed line is for $a = 0.5$, $k_{eq}/k_c = 10.$, and $BP = 13\%$. The dotted line is for $a = 0.5$, $k_{eq}/k_c = 10.$, and $BP = 15\%$. The dash-dot-dot-dot line is for $a = 0.5$, $k_{eq}/k_c = 10.$, and $BP = 20\%$. All of the later three are consistant with the APM data.

Figure 5: Power spectrum of velocity fields in units of $(100 \text{ km s}^{-1})^2$ without biasing. The solid line is for $a = 0.5$, $k_{eq}/k_c = 10.$ and $BP = 20\%$. The dotted line is for the standard CDM model with $\Omega = 1$, and $h = 0.5$.

Figure 6: The correlation function $\xi_{11}(R)$ of 3-dim. velocity field in units of $(100 \text{ km s}^{-1})^2$. The solid line is for $a = 0.5$, $k_{eq}/k_c = 10.$, and $BP = 50\%$. The

dotted line is for $a = 0.5$, $k_{eq}/k_c = 10.$ and $BP = 20\%$. The long dashed line is for the standard CDM model with $\Omega = 1$, and $h = 0.5$.

Figure 7: The correlation function $\xi_{11}(R)$ of $\vec{v}(\vec{r}) \cdot \hat{r}$ in units of $(100 \text{ km s}^{-1})^2$. $a = 0.5$, $k_{eq}/k_c = 10$. The solid line is for $BP = 50\%$. The dotted line is for $BP = 20\%$. The long dashed line is for the standard CDM model with $\Omega = 1$, and $h = 0.5$.

Figure 8: The rms of $\vec{v}(\vec{r}, R) = 1/V \int_{|\vec{r}'| \leq R} d^3\vec{r}' \vec{v}(\vec{r} + \vec{r}')$ in units of 100 km s^{-1} . $a = 0.5$, $k_{eq}/k_c = 10$. The solid line is for $BP = 50\%$. The dotted line is for $BP = 20\%$. The long dashed line is for the standard CDM model with $\Omega = 1$, and $h = 0.5$.

Figure 9: The rms of $v_r(\vec{r}, R) = 1/V \int_{|\vec{r}'| \leq R} d^3\vec{r}' \vec{v}(\vec{r} + \vec{r}') \cdot \hat{r}$ in units of 100 km s^{-1} . $a = 0.5$, $k_{eq}/k_c = 10.$ The solid line is for $a = 0.5$, $k_{eq}/k_c = 10.$ and $BP = 50\%$. The dotted line is for $a = 0.5$, $k_{eq}/k_c = 10.$ and $BP = 20\%$. The dash-dot-dot-dot line is for $a = 0.5$, $k_{eq}/k_c = 5$, and $BP = 50\%$. The long dashed line is for the standard CDM model with $\Omega = 1$, and $h = 0.5$.

Figure 10: The velocity distribution function. The horizontal axis is $\nu = v / \langle v^2 \rangle^{1/2}$. The solid line is for the purely non-Gaussian velocity distribution. The long dashed line is for the velocity distribution of a Gaussian field

Figure 11: The distribution functions of total velocity fields. The horizontal axes are $\nu = v / \langle v^2 \rangle^{1/2}$. In (a), (b) and (c), the solid line is for the purely non-Gaussian distribution, and the long dashed line is for the velocity distribution of a Gaussian field. (a) $a = 0.5$, $k_{eq}/k_c = 10.$ and $BP = 50\%$. The dash-dot-dot-dot line is for $R_f = 20 \text{ Mpc h}^{-1}$. The dotted line is for $R_f = 10 \text{ Mpc h}^{-1}$. The dashed line is for $R_f = 2 \text{ Mpc h}^{-1}$. (b) $a = 0.5$, $k_{eq}/k_c = 10$, and $R_f = 10 \text{ Mpc h}^{-1}$. The dotted line is for $BP = 50\%$. The dashed line is for $BP = 20\%$. (c) $R_f = 10 \text{ Mpc h}^{-1}$, and $BP = 50\%$. The dashed line is for $a = 0.5$, $k_{eq}/k_c = 10.$ The dotted line is for $a = 0.5$, $k_{eq}/k_c = 5.$

Figure 12: The contributions to $\sigma^2(\theta_{FWHM})$. The dashed line is the Gaussian contribution, and the solid line is the non-Gaussian contribution. $a = 0.5$, $k_{eq}/k_c = 10$, and $BP = 20\%$. The horizontal axis is $\theta_{FWHM} (\text{arcmin})$.

Figure 13: Vector plots of one slice of initial velocity fields. The areas are $128 \times 128(\text{Mpch}^{-1})^2$. (a) The standard CDM model with $\Omega = 1$, and $h = 0.5$. (b) The combination model with $a = 0.5$, $k_{eq}/k_c = 5.$, and $BP = 25\%$.

Figure 14: Vector plots of one slice of present velocity fields by using the adhesion model. The areas are $128 \times 128(\text{Mpch}^{-1})^2$. (a) The standard CDM model with $\Omega = 1$, and $h = 0.5$. (b) The combination model with $a = 0.5$, $k_{eq}/k_c = 5.$, and $BP = 25\%$.

TABLE 1: Parameter Regime

k_{eq}/k_c	a	$BP(\%)$
5 – 8	> 0.5	< 20
9	0.5	17.7 – 29.3
10	0.5	13.8 – 22.9
11	0.5	12.4 – 20.5
12	0.5	11.2 – 18.5
13	0.5	10.2 – 16.9
13	0.4	13.9 – 23.1
14	0.4	12.8 – 21.3

Note.— The third column BP is the non-Gaussian contribution to $\langle (\delta\rho/\rho)^2 \rangle$ on 8 Mpc h^{-1} as a percentage, chosen to fit the APM angular correlation function $w(\theta)$.

TABLE 2: The 3-dim. rms peculiar velocity

$a = 0.5, k_{eq}/k_c = 10, BP = 15\%$				
	$\sigma_v(3.2)$	$\sigma_v(15)$	$\sigma_v(25)$	$\sigma_v(40)$
Standard CDM	759	408	294	207
Combination Model	777	473	357	253

Note.— The numbers in the parentheses stand for the smoothing scales in units of Mpc h^{-1} , and the rms velocity σ_v is in units of kms^{-1} .

TABLE 3: Temperature fluctuations

$h = 0.5, \Omega = 1, a = 0.5, k_{\epsilon q}/k_c = 10, BP = 20\%$		
	$(\Delta T/T)_2$	$\sigma(\theta_{FWHM} = 10^\circ)$
Standard CDM	5.2×10^{-6}	1.06×10^{-5}
Combination Model	4.7×10^{-6}	1.01×10^{-5}

Fig. 1

Fig. 2(a)

Fig. 2(b)

Fig. 2(c)

Fig.3

Fig.4

Fig.5

Fig.10

Fig.11(a)

Fig.11(b)

Fig.11(c)

Fig.12

Fig.13(a)

Fig.13(b)

Fig.14(a)

Fig.14(b)

END

DATE
FILMED
6/9/93